

45

Señores
DIVISION DE NUEVAS CREACIONES
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
Santafé de Bogotá, D.C.

SUPERINDUSTRIA Y COMERCIO
Radicacion : 00015142 00000001 Folios: 53
Fecha (AMD): 2000-05-05 12:31:16
Tramite : 002 PATENTES D 1 REGISTRO/ 329 COMPLEMEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Re.: Código 02-01-09
Expediente No. 00-15.142
Solicitud de Patente
SEPRACOR, INC.

En relación con el Expediente de la referencia y con el fin de dar cumplimiento a la preceptuado por el artículo 4o. literal d.) numeral 3 de la Convención de París adjunto a este escrito sírvanse encontrar la copia certificada de la solicitud de patente de la cual se reivindica prioridad, la cual se encuentra debidamente traducida en sus apartes pertinentes.

De Ustedes Atentamente,

GERMAN CASTILLO GRAU
T.P. No. 2978 del C.S.J.



CONSULADO DE COLOMBIA
WASHINGTON, D.C.



46

No. 757

EL CONSULADO DE COLOMBIA

CERTIFICA:

Que es auténtica la firma de PATRICK O. HATCHETT.
quien el día 28 DE MARZO DEL 2000
desempeñaba las funciones de OFICIAL DE AUTENTICACIONES DEPARTAMENTO DE ESTADO.
la cual aparece al pie de los documentos anexos relacionados con APLICACION PARA PATENTE.

Dado en Washington, D.C. a los VEINTINUEVE (29) días del mes de
MARZO. del año dos mil (2000)

Pagado Impuesto de Timbre 7=

Pagado Derecho Consular 15=

Total 22=

JAQUELINE ESPITIA ARIAS
Vicecónsul

(El Consulado no asume ninguna
responsabilidad por el contenido de los
documentos anexos)

MINISTERIO DE RELACIONES
EXTERIORES

256129
JADELINE ESPITA
CUYA FIRMA APARECE EN EL
DOCUMENTO DESEMPEÑA
LAS FUNCIONES INDICADAS.

NO SE ASUME
RESPONSABILIDAD DEL TEXTO


2000 MAY -2 A 11:41

Malpica
JEFE DE LEGALIZACIONES
SANTAFE DE BOGOTA D.C.

United States of America

0032620-3

3 10
47



DEPARTMENT OF STATE

To all to whom these presents shall come, Greetings:



I Certify That the document hereunto annexed is under the Seal of the Patent and Trademark Office, Department of Commerce, United States of America, and that such Seal is entitled to full faith and credit.*

In testimony whereof, I, Madeleine K. Albright, Secretary of State, have hereunto caused the seal of the Department of State to be affixed and my name subscribed by the Assistant Authentication Officer, of the said Department, at the city of Washington, in the District of Columbia, this twenty-eighth day of March, 2000.

Madeleine K. Albright

Secretary of State


By *Paul J. Haney*

Assistant Authentication Officer,
Department of State

Issued pursuant to CHXIV, State of
Sept. 15, 1789, 1 Stat. 68-69; 22
USC 2657; 22 USC 2651a; 5 USC
301; 28 USE 1733 et. seq.; 8 USC
1443(f); RULE 44 Federal Rules of
Civil Procedure.

**For the contents of the annexed document, the Department assumes no responsibility*
This certificate is not valid if it is removed or altered in any way whatsoever

PA 220624



Columbia

48

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

March 14, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/122,394

FILING DATE: March 02, 1999



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

P. SWAIN
Certifying Officer

COVER SHEET FOR PROVISIONAL APPLICATION FOR PATENT

Commissioner for Patents
PROVISIONAL PATENT APPLICATION
Washington, DC 20231

Request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(b)(2).

JCS41 U.S. PTO
60/122394
03/02/99



550
49

Docket Number	4821-344-888	Type a plus sign (+) inside this box -	+
---------------	--------------	---	---

INVENTOR(s) APPLICANT(s)			
LAST NAME	FIRST NAME	MIDDLE INITIAL	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Rubin Barberich	Paul Timothy	D. J.	Sudbury, MA Concord, MA

TITLE OF THE INVENTION (280 characters max)

METHODS AND COMPOSITIONS USING (+) NORCISAPRIDE IN COMBINATION WITH PROTON PUMP INHIBITORS OR H₂ RECEPTOR ANTAGONISTS

CORRESPONDENCE ADDRESS:

PENNIE & EDMONDS LLP
1667 K Street, N.W.
Washington, D.C. 20006
(202) 496-4400

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification	Number of Pages	35	☐ Small Entity Statement
☐ Drawing(s)	Number of Sheets	0	☐ Other (specify)

METHOD OF PAYMENT (check one)

☐ A check or money order is enclosed to cover the Provisional filing fees.	ESTIMATED PROVISIONAL FILING FEE AMOUNT
☒ The Commissioner is hereby authorized to charge the required filing fee to Deposit Account Number 16-1150.	☒ \$150 ☐ \$75

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No. ☐ Yes, the name of the U.S. Government agency and the Government contract number are: _____

Respectfully submitted,

Signature Stanton T. Lawrence, III (42,234)
STANTON T. LAWRENCE, III
PENNIE & EDMONDS LLP

REGISTRATION NO. 25,736 (if appropriate)

Date _____

☐ Additional inventors are being named on separately numbered sheets attached hereto. Total number of cover sheet pages. 1

PROVISIONAL APPLICATION FILING ONLY

50

**METHODS AND COMPOSITIONS
USING (+) NORCISAPRIDE IN COMBINATION WITH
PROTON PUMP INHIBITORS OR H₂ RECEPTOR ANTAGONISTS**

1. FIELD OF THE INVENTION

The invention relates to methods and compositions for the prevention, treatment, or management of gastrointestinal disorders or symptoms thereof, by administering one or more agent(s) or compound(s) that simultaneously or sequentially act on a 5-HT₃ receptor, a 5-HT₄ receptor, and either an H₂ receptor or a proton pump.

2. BACKGROUND OF THE INVENTION

Gastrointestinal disorders are common disorders that affect the gastrointestinal tract, *i.e.*, the stomach and intestines. Various gastrointestinal disorders exist, including: gastro-esophageal reflux disease, emesis, gastrointestinal motility dysfunction, gastrointestinal ulcers, pathological hypersecretory conditions, and gastric hyperacidity. These diseases may be treated by various non-invasive means, such as administering to a patient a therapeutic agent, such as ZANTAC® (ranitidine), TRITEC® (ranitidine), AXID® (nizatidine), TAGAMET® (cimetidine), PREVACID® (lansoprazole), PEPCID®, PEPCID AC® ACID CONTROLLER™, MYLANTA AR ACID REDUCER™ (famotidine), PRILOSEC® (omeprazole), and others. New pharmaceutical compounds and preparations are continually being developed.

U.S. Patent Nos. 4,962,115, 5,057,525, and 5,137,896 (collectively "Van Daele") disclose N-(3-hydroxy-4-piperidenyl)benzamides. These compounds are said to stimulate the motility of the gastrointestinal system. Van Deale states that the *cis* and *trans* diastereomeric racemates of these compounds may be obtained separately by conventional methods, and the *cis* and *trans* diastereomeric racemates may be further resolved into their optical isomers. One such racemate, Cisapride, is chemically named *cis*-4-amino-5-chloro-N-[1-[3-(4-fluorophenoxy)propyl]-3-methoxy-4-piperidinyl]-2-methoxybenzamide. Schapira *et al.*, *Acta Gastroenterolog. Belg.*, LIII:446-457 (1990). Cisapride is used primarily to treat gastroesophageal reflux disease ("GERD"), which is characterized as the backward flow of the stomach contents into the esophagus. Cisapride is commercially available as the racemic mixture of the *cis*(-) and *cis*(+) diastereomeric enantiomers of cisapride known as PROPULSID®.

7-8
51

Benzamide derivatives have several prominent pharmacological actions due to their effects on neuronal systems modulated by the neurotransmitter serotonin. It has been reported that a major site of production and storage of serotonin is the enterochromaffin cell of the gastrointestinal mucosa. It was also reported that serotonin provides a powerful intestinal transit and decreasing absorption time, as occurs with diarrhea. This stimulating action is also associated with nausea and vomiting.

Because of their modulation of the serotonin neuronal system in the gastrointestinal tract, some benzamide derivatives are effective antiemetic agents and are used to control vomiting during cancer chemotherapy or radiotherapy. Costall *et al.*, *Neuropharmacology*, 26:1321-1326 (1987). This action is the result of an ability to block serotonin at specific sites, particularly Type-3 5-hydroxytryptamine (5-HT₃) receptors. Clarke *et al.*, *Trends in Pharmacological Sciences*, 10:385-386 (1989). Chemotherapy and radiation therapy can induce nausea and vomiting by damaging enterochromaffin cells in the gastrointestinal tract. As a result, the neurotransmitter serotonin is released and stimulates both afferent vagal nerve fibers (thus initiating the vomiting reflex) and serotonin receptors in the chemoreceptor trigger zone of the area postrema region of the brain. The anatomical site for this action of the benzamide derivatives, and whether such action is central (CNS), peripheral, or a combination thereof, remains unresolved. Barnes *et al.*, *J. Pharm. Pharmacol.*, 40:586-588 (1988).

A second prominent action of certain benzamide derivatives is in augmenting gastrointestinal smooth muscle activity from the esophagus to the proximal small bowel, thus accelerating esophageal and small intestinal transit, as well as facilitating gastric emptying and increasing lower esophageal sphincter tone. Decktor *et al.*, *Eur. J. Pharmacol.*, 147: 313-316 (1988). Although the benzamide derivatives are not cholinergic receptor agonists *per se*, the aforementioned smooth muscle effects may be blocked by muscarinic receptor blocking agents such as atropine or inhibitors of neuronal transmissions, such as the tetrodotoxin type that block sodium channels. Fernandez and Massingham, *Life Sci.*, 36: 1-14 (1985). Similar blocking activity has been reported for the contractile effects of serotonin in the small intestine. Craig and Clarke, *Brit. J. Pharmacol.*, 96: 247P (1989). It is believed that the primary smooth muscle effects of some benzamide derivatives are the result of an agonist action upon a class of serotonin receptors referred to as 5-HT₄ receptors, which are located on interneurons in the myenteric plexus of the gut

52
wall. Clarke *et al.*, *Trends in Pharmacological Sciences*, 10: 385-386 (1989) and Dumuis *et al.*, *N. S. Arch. Pharmacol.*, 340: 403-410 (1989). Activation of these receptors subsequently enhances the release of acetylcholine from parasympathetic nerve terminals located near surrounding smooth muscle fibers. It is the combination of acetylcholine with its receptors on smooth muscle membranes which is the actual trigger for muscle contraction.

It has been reported that cisapride enters the central nervous system and binds to 5-HT₄ receptors. This indicates that cisapride may have centrally-mediated effects. Cisapride is a potent ligand at 5-HT₄ receptors, which are located in several areas of the central nervous system. Dumuis *et al.*, *N.S. Arch. Pharmacol.*, 340: 403-410 (1989). Modulation of serotonergic systems may have a variety of behavioral effects.

The co-administration of racemic cisapride with other therapeutic agents causes inhibitory problems with the metabolism of cisapride by the liver. For example, ketoconazole has a pronounced effect on cisapride kinetics resulting from the inhibition of the metabolic elimination of cisapride and leads to an 8-fold increase of the steady-state plasma levels. *Physician's Desk Reference*®, Medical Economics Co., Inc., p. 1308-1309, 52nd Edition (1998). Interaction of racemic cisapride and other therapeutic agents can also potentiate cardiovascular side effects, such as cardiotoxicity. This potentiation occurs when other drugs present in the patient's system interfere with the metabolism of cisapride, thereby causing a build up of racemic cisapride in the body.

These interactions are a significant drawback to the use of racemic cisapride; in particular, because racemic cisapride is often used before, with, or immediately after another therapeutic agent. In addition, administration of racemic cisapride to a human has been found to cause adverse effects such as cardiac arrhythmia, including ventricular tachycardia, ventricular fibrillation, Q_T prolongation, and torsades de pointes, central nervous system ("CNS") effects, increased systolic pressure, interactions with other drugs, diarrhea, abdominal cramping, and cardiac depression.

Racemic cisapride in humans is metabolized mainly by oxidative N-dealkylation of the piperidine nitrogen or by aromatic hydroxylation occurring on either the 4-fluorophenoxy or benzamide rings. Meuldermans *et al.*, *Drug Metab. Dispos.*, 16(3):410-419 (1988); and Meuldermans *et al.*, *Drug Metab. Dispos.*, 16(3):403-409 (1988).

9-54
53

Norcisapride, chemically named 4-amino-5-chloro-N-(3-methoxy-4-piperidinyl)-2-methoxybenzamide, is an active metabolite of cisapride.

5 Recently, investigators have reported that the optically pure (+) stereoisomer of the cisapride metabolite norcisapride exhibits many useful characteristics, but without certain side effects of racemic cisapride. Specifically, U.S. Patent No. 5,739,151 discloses a method of eliciting an antiemetic effect and treating other conditions using optically pure (+) norcisapride.

10 Other agents or compounds which have been studied for the treatment of gastrointestinal diseases include proton pump inhibitors and H_2 receptor antagonists. Proton pump inhibitors treat gastrointestinal diseases by inhibiting H^+K^+ ATPase and thereby regulating acidity in gastric juices. H_2 receptor antagonists inhibit histamine binding with H_2 receptors to regulate gastric acid secretion. Goodman & Gilman, *The Pharmacological Basis of Therapeutics*, 9th Edition, pp. 901-915 (1996).

15 The concentration gradients of gastric acids across cell membranes vary widely. Perhaps the largest gradient in the body occurs across the plasma membrane of the parietal cells of the stomach lining, which secrete hydrochloric acid into gastric juice. Since the concentration of hydrochloric acid in gastric juice may be as high as 0.1 M and the concentration of H^+ in the cells is about 10^{-7} M, parietal cells can secrete H^+ ions against a
20 concentration gradient of about 1 million to 1. A membrane-bound enzyme called H^+K^+ ATPase facilitates active transport of H^+ across membranes against concentration gradients in exchange for K^+ to form gastric hydrochloric acid. For each molecule of cytosolic ATP hydrolyzed to ADP and phosphate, 2 H^+ ions are transported across the plasma membrane from the cytosol to the stomach. Goodman & Gilman, *The
25 Pharmacological Basis of Therapeutics*, 9th Edition, pp. 901-915 (1996).

Proton pump inhibitors suppress gastric acid secretion, the final step of acid production, by specific inhibition of the H^+K^+ ATPase enzyme system at the secretory surface of gastric parietal cells. Proton pump inhibitors include benzimidazole compounds, for example, omeprazole (PRILOSEC®), lansoprazole (PREVACID®), and pantoprazole.
30 These proton pump inhibitors contain a sulfinyl group situated between substituted benzimidazole and pyridine rings. At neutral pH, omeprazole, lansoprazole, and pantoprazole are chemically stable, lipid soluble, weak bases that are devoid of inhibitory activity. These uncharged weak bases reach parietal cells from the blood and diffuse into

10-58
54

the secretory canaliculi, where the drugs become protonated and thereby trapped. The protonated species rearranges to form a sulfenic acid and a sulfenamide, the latter species capable of interacting with sulfhydryl groups of H^+-K^+ ATPase. Full inhibition occurs with two molecules of inhibitor per molecule of enzyme. The specificity of the effects of proton pump inhibitors is believed to derive from: a) the selective distribution of H^+-K^+ ATPase; b) the requirement for acidic conditions to catalyze generation of the reactive inhibitor; and c) the trapping of the protonated drug and the cationic sulfenamide within the acidic canaliculi and adjacent to the target enzyme. Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Edition, pp. 901-915 (1996).

H_2 receptor antagonists competitively inhibit the interaction of histamine with H_2 receptors. They are highly selective and have little or no effect on H_1 receptors. Although H_2 receptors are present in numerous tissues, including vascular and bronchial smooth muscle, H_2 receptor antagonists interfere remarkably little with physiological functions other than gastric acid secretion. H_2 receptor antagonists include nizatidine (AXID®), ranitidine (ZANTAC® and TRITEC®), famotidine (PEPCID AC®), and cimetidine (TAGAMET®). Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Edition, pp. 901-915 (1996).

H_2 receptor antagonists inhibit gastric acid secretion elicited by histamine, other H_2 agonists, gastrin, and, to a lesser extent, muscarinic agonists. H_2 receptor antagonists also inhibit basal and nocturnal acid secretion, and this effect contributes in a major way to their clinical efficacy.

Although therapeutic agents are available, there remains a need for a more effective, broad-spectrum treatment or therapy for gastrointestinal disorders. For example, it is desirable to have safe and effective methods and compositions for preventing, treating, and managing gastrointestinal disorders without adverse effects or adverse drug-drug interactions.

3. SUMMARY OF THE INVENTION

The invention encompasses the prevention, treatment, or management of gastrointestinal disorders by biological action at three different sites: action at $5-HT_3$ receptors, $5-HT_4$ receptors, and either H_2 receptors or proton pumps. The use of triple action is considered to provide an unexpectedly superior therapeutic profile. The use of this

456
55

triple action therapy can be less toxic and/or more potent for preventing, treating, and managing gastrointestinal disorders than the use of agents that act on any of the three sites alone.

5 In one embodiment, three therapeutic agents or compounds are used, one providing action at 5-HT₃ receptors, another providing action at 5-HT₄ receptors, and the third being either a proton pump inhibitor or an H₂ receptor antagonist. For example, in a preferred embodiment, cisapride, ondansetron, and either a proton pump inhibitor or an H₂ receptor antagonist are used to prevent, treat, or manage gastrointestinal disorders. In a
10 more preferred embodiment, either optically pure (+) cisapride or optically pure (-) cisapride, or a pharmaceutically acceptable salt thereof, optically pure R(+) ondansetron, or a pharmaceutically acceptable salt thereof, and either a proton pump inhibitor or an H₂ receptor antagonist are used.

15 In another embodiment, the invention encompasses the synergistic use of two or more agents or compounds to provide triple site action on 5-HT₃ receptors, 5-HT₄ receptors, and either H₂ receptors or proton pumps. In a preferred embodiment, two agents or compounds are used, one providing dual action at 5-HT₃ receptors and 5-HT₄ receptors, and another providing action either at H₂ receptors or proton pumps. For example, in this preferred embodiment, optically pure (+) norcisapride, or a pharmaceutically acceptable salt
20 thereof, which has both 5-HT₃ and 5-HT₄ activity, and either a proton pump inhibitor or H₂ receptor antagonist are used to prevent, treat, or manage gastrointestinal disorders.

This invention also encompasses the use of two separate pharmaceutical compositions adapted for the prevention, treatment, or management of a patient suffering from gastrointestinal disorders or symptoms thereof, one which comprises a therapeutically
25 effective amount of optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, and another which a therapeutically effective amount of either a proton pump inhibitor or H₂ receptor antagonist.

Alternatively, the invention also encompasses pharmaceutical compositions comprising optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, in
30 combination with a either a proton pump inhibitor or an H₂ receptor antagonist. Such single unit dosage forms comprise from about 0.5 mg to about 500 mg of optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, and from about 1 mg to about

6042224-020249

12-51
56
200 mg of a proton pump inhibitor or from about 1 mg to about 2400 mg an H₂ receptor antagonist, in a suitable carrier.

5 The pharmaceutical compositions and methods of the invention, particularly those which comprise a therapeutically effective amount of optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, can be used to prevent or alleviate symptoms of gastrointestinal disorders, while reducing or avoiding adverse effects associated with administration of conventional treatments such as conventional 5-HT₃ receptor antagonists, 5-HT₄ receptor agonists or antagonists, H₂ receptor antagonists, and proton pump inhibitors. 10 Moreover, the compositions and methods of the present invention encompass the treatment, prevention or management of gastrointestinal disorders while reducing or avoiding adverse drug-drug interactions which are known to occur with the use of existing commercial agents, such as racemic cisapride.

15 4. DETAILED DESCRIPTION OF THE INVENTION

The invention encompasses the prevention, treatment, or management of gastrointestinal disorders by biological action at three different sites, action on 5-HT₃ receptors, 5-HT₄ receptors, and either H₂ receptors or proton pumps (e.g., inhibition of H⁺-K⁺-ATPase enzyme system). The use of triple-site action is considered to provide an 20 unexpectedly superior therapeutic profile to conventional treatment of gastrointestinal disorders. The use of this triple action therapy can be safer and/or more effective for preventing, treating, and managing gastrointestinal disorders than the use of agents that act on any of the three sites alone.

More specifically, the invention also encompasses the use of at least three 25 agents or compounds to prevent, treat, or manage gastrointestinal disorders, or symptoms thereof, one providing action at 5-HT₃ receptors, another providing action at 5-HT₄ receptors, and a third providing action at either proton pump systems or H₂ receptors. Preferably, the three compounds are cisapride, ondansetron, and either a proton pump inhibitor or an H₂ receptor antagonist. More preferably, the three compounds are either 30 optically pure (+) cisapride or optically pure (-) cisapride, or a pharmaceutically acceptable salt thereof, optically pure R(+) ondansetron or a pharmaceutically acceptable salt thereof, and either a proton pump inhibitor or H₂ receptor antagonist.

13 58
57

5 The invention also encompasses the synergistic use of at least two agents or compounds to provide triple action on 5-HT₃ receptors, 5-HT₄ receptors, and either H₂ receptors or the proton pump mechanism in the prevention, treatment, or management of gastrointestinal disorders. Preferably, two agents or compounds are used, one providing dual action at 5-HT₃ receptors and 5-HT₄ receptors, and another providing action at either H₂ receptors or proton pump systems. More preferably, the two agents or compounds are optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, and either an H₂ receptor antagonist or a proton pump inhibitor.

10 Gastrointestinal disorders that can be treated by the compositions and methods of the invention include, but are not limited to, disorders of the upper and lower gastrointestinal system, gastro-esophageal reflux disease ("GERD"), emesis, gastrointestinal motility dysfunction, gastrointestinal ulcers, pathological hypersecretory conditions, and gastric hyperacidity. The gastrointestinal disorders additionally include, but are not limited to, dyspepsia, gastroparesis, constipation, post-operative ileus, intestinal pseudo obstruction, 15 gastric ulcers, duodenal ulcers, heartburn, acid indigestion, erosive esophagitis, sour stomach, upset stomach and Zollinger-Ellison Syndrome.

In a most preferred embodiment, the invention encompasses the use of the optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, and at least one of a proton pump inhibitor or an H₂ receptor antagonist, in preventing, treating, or 20 managing gastrointestinal disorders, or symptoms thereof. It should be noted that racemic norcisapride, or a pharmaceutically acceptable salt thereof, may be used as an alternative to optically pure (+) norcisapride in the methods and compositions of the invention, although optically pure (+) norcisapride is preferred. Similarly, optically pure stereoisomers or active metabolites of the proton pump inhibitors or the H₂ antagonists may also be use as 25 alternatives where appropriate, as well as pharmaceutically acceptable salts thereof. Specific examples by way of illustration only are set forth herein.

Without being limited by theory, it is believed that utilization of an agent or compound having dual activity as a 5-HT₃ receptor antagonist and a 5-HT₄ receptor agonist 30 along with either a proton pump inhibitor or an H₂ receptor antagonist provides triple-site action, which surprisingly results in clearer dose-related definitions of efficacy, diminished adverse effects, a superior therapy due to synergistic activity, and accordingly, an improved therapeutic index. For example, optically pure (+) norcisapride has dual activity as a 5-HT₃

14
58

receptor antagonist and a 5-HT₄ receptor agonist, and triple-site action is achieved when used with either a proton pump inhibitor or an H₂ receptor antagonist. It is, therefore, more desirable to use the compositions and methods of the invention than to use a 5-HT₃ receptor antagonist, a 5-HT₄ receptor agonist, an agent or compound having dual activity as a 5-HT₃ receptor antagonist and a 5-HT₄ receptor agonist, a proton pump inhibitor, or an H₂ receptor antagonist alone.

The use of the optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, with either a proton pump inhibitor or an H₂ receptor antagonist, in preventing, treating, or managing gastrointestinal disorders, in accordance with the present invention is consider to reduce or avoid adverse effects associated with existing commercial treatment for gastrointestinal disorders, such as racemic cisapride. Further, this embodiment is considered to reduce or avoid adverse drug-drug interactions associated with racemic cisapride.

The invention also encompasses methods of preventing, treating, or managing a condition caused by a dysfunction of 5-HT₃ receptors, 5-HT₄ receptors, or proton pumps or H₂ receptors by administering a therapeutically effective amount of optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, with a therapeutically effective amount of either a proton pump inhibitor or an H₂ receptor antagonist.

The invention encompasses methods of preventing, treating, or managing GERD, emesis, gastrointestinal motility dysfunction, gastrointestinal ulcers, pathological hypersecretory conditions, or gastric hyperacidity which comprise administering to a patient a therapeutically effective amount of optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, with either a proton pump inhibitor or an H₂ receptor antagonist. In specific embodiments, the invention also encompasses the use of these agents in combination for preventing, treating, or managing erosive esophagitis, dyspepsia, gastroparesis, constipation, post-operative ileus, intestinal pseudo obstruction, gastric ulcers, duodenal ulcers, heartburn, acid indigestion, erosive esophagitis, sour stomach, upset stomach, and Zollinger-Ellison Syndrome.

The invention, which encompasses the use of triple action therapy, may optionally further include the use of one or more additional therapeutic agents known to treat gastrointestinal disorders. Examples of such additional therapeutic agents include, but

are not limited to, hydroxyzine (ANTARAX®); diphenhydramine (BENADRYL PARENTAREL®); prochlorperazine (COMPAZINE®); dronabinol (MARINOL®); promethazine (PHENERGAN®); meclizine (ANTIVERT®); trimethobenzamide (TIGAN®); thiethylperazine (TORECAN®); perphenazine (TRILAFON®); sucralfate (CARAFATE®); and the like, and where applicable, optically pure stereoisomers or active metabolites thereof.

The administration of two or more therapeutic agents used in accordance with the methods of the invention may be concurrent, sequential, or both, *i.e.*, optically pure (+) norcisapride, or a pharmaceutically acceptable salt thereof, and either a proton pump inhibitor or an H₂ receptor antagonist, and optionally an additional therapeutic agent, may be administered as a combination, concurrently but separately, or by sequential administration.

The methods and compositions of this invention are considered to provide the benefit of reducing or avoiding adverse effects associated with prior methods and compositions used in the treatment of gastrointestinal disorders. *See, e.g., Physician's Desk Reference®*, Medical Economics Co., Inc., 52nd Edition (1998 and 1999).

The terms "adverse effects" and "adverse side effects," as used herein, include, but are not limited to, cardiac arrhythmia, cardiac conduction disturbances, appetite stimulation, weight gain, sedation, gastrointestinal distress, headache, dry mouth, constipation, diarrhea, and drug-drug interactions. The term "cardiac arrhythmia" includes, but is not limited to, ventricular tachyarrhythmia, torsades de pointes, Q_T prolongation, and ventricular fibrillation.

The term "gastrointestinal disorder," as used herein, includes, but is not limited to, gastrointestinal motility dysfunction, GERD, emesis, gastrointestinal ulcers, pathological hypersecretory conditions, gastric hyperacidity, erosive esophagitis, dyspepsia, gastroparesis, constipation, post-operative ileus, intestinal pseudo obstruction, gastric ulcers, duodenal ulcers, heartburn, acid indigestion, erosive esophagitis, sour stomach, upset stomach, and Zollinger-Ellison Syndrome.

The term "patient," as used herein, refers to a mammal, particularly a human.

The term "racemic," as used herein, is defined as a mixture of the (+) and (-) enantiomers of a compound wherein the (+) and (-) enantiomers are present in approximately a 1:1 ratio.

16-61
60

The phrase "optically pure," as used herein, means that the composition contains greater than about 90% of the desired stereoisomer by weight, preferably greater than about 95% of the desired stereoisomer by weight, and more preferably greater than about 99% of the desired stereoisomer by weight, based upon the total weight of the active ingredient, *e.g.*, norcisapride when used to qualify (+) norcisapride. The term "substantially free," as used herein, means less than about 10 weight percent, preferably less than about 5 weight percent, and more preferably less than about 1 weight percent of the undesired stereoisomer, *e.g.*, (-) norcisapride, is present according to the invention.

10 The phrase "5-HT₃ receptor antagonist," as used herein, means a compound capable of binding reversibly to a Type-3 5-hydroxytryptamine receptor. 5-HT₃ receptor antagonists include, but are not limited to, granisetron (KYTRIL®), metoclopramide (REGLAN®), ondansetron (ZOFRAN®), renzapride, zacopride, tropisetron, and where applicable, optically pure stereoisomers or active metabolites thereof.

15 The phrase "5-HT₄ receptor agonist," as used herein, means a compound capable of binding reversibly to a Type-4 5-hydroxytryptamine receptor. 5-HT₄ receptor agonists include, but are not limited to, norcisapride and cisapride, and where applicable, optically pure stereoisomers or active metabolites thereof.

20 The phrase "proton pump," as used herein, refers to H⁺-K⁺ ATPase, a membrane-bound enzyme that facilitates active transport of H⁺ across membranes against a concentration gradient.

25 The phrase "proton pump inhibitor," as used herein, refers to any agent or compound that inhibits or suppresses gastric acid secretion by inhibition of the H⁺-K⁺ ATPase enzyme system at the secretory surface of gastric parietal cells. Proton pump inhibitors include, but are not limited to, prazole derivatives, such as omeprazole lansoprazole, pantoprazole, rabeprazole, and where applicable, optically pure stereoisomers such as optically pure (+) lansoprazole, optically pure (-) lansoprazole, optically pure (+) omeprazole, optically pure (-) omeprazole, optically pure (+) rabeprazole, optically pure (-) rabeprazole, optically pure (+) pantoprazole and optically pure (-) pantoprazole, or active
30 metabolites thereof. Active metabolites of proton pump inhibitors suitable for use according to the invention include, but are not limited to, hydroxy-omeprazole, hydroxy-lansoprazole, the carboxylic acid derivative of omeprazole, and desmethyl-pantoprazole, and where applicable, optically pure stereoisomers thereof. Omeprazole, lansoprazole,

12-62
61

5 pantoprazole, and rabeprazole, for example, may be prepared by syntheses known to those of ordinary skill in the art, particularly from U.S. Patent Nos. 4,544,750, 4,620,008, 4,620,008, 4,758,579, 5,045,552, 5,374,730, 5,386,032, 5,470,983, and 5,502,195, the disclosures of which are each incorporated herein by express reference thereto.

10 The phrase "H₂ receptor antagonist," as used herein, refers to any agent or compound that competitively inhibits the interaction of histamine with H₂ receptors. H₂ receptor antagonists include, but are not limited to, cimetidine, famotidine, ranitidine, nizatidine, and where applicable, optically pure stereoisomers or active metabolites thereof. Active metabolites of H₂ receptor antagonists include, but are not limited to, N2-monodesmethylnizatidine and where applicable, optically active stereoisomers thereof.

15 Nizatidine may be prepared by synthesis known to those of ordinary skill in the art, particularly from U.S. Patent Nos. 5,541,335 and 5,700,945, the disclosures of which are each incorporated herein by express reference thereto. Ranitidine may be prepared by synthesis known to those of ordinary skill in the art, particularly from U.S. Patent No. 5,118,813, the disclosure of which is each incorporated herein by express reference thereto. Cimetidine may be prepared by synthesis known to those of ordinary skill in the art, particularly from U.S. Patent Nos. 4,413,129, 4,855,439, 4,886,910, 4,886,912, and 5,118,813, the disclosures of which are each incorporated herein by express reference thereto. Moreover, famotidine may be prepared by synthesis known to those of ordinary skill in the art, described in, for example, Guimaraens *et al.*, *Contact Dermatitis*, 31(4):259 (1994).

20 The diseases prevented, treated, or managed in the invention are herein used consistently according to *Stedman's Medical Dictionary*, 26th Edition, Williams and Wilkins (1995).

25 The terms "gastro-esophageal reflux disease" or "GERD," as used herein, is defined as a condition characterized by the backward flow of the stomach contents into the esophagus.

30 The term "gastrointestinal ulcer," as used herein, is defined as a condition characterized by a lesion or lesions on the surface of the lining of the gastrointestinal tract, caused by superficial loss of tissue, usually accompanied by inflammation. Gastrointestinal ulcers include, but are not limited to, duodenal ulcers, and gastric ulcers.

042224 0309

18-63
62

The phrase "erosive esophagitis," as used herein, is defined as a condition characterized by inflammation of the lower esophagus from regurgitation of acid gastric contents, usually due to malfunction of the lower esophageal sphincter.

5 The term "dyspepsia," as used herein, is defined as a condition characterized by an impairment of the power or function of digestion that can arise as a symptom of primary gastrointestinal dysfunction or as a complication due to other disorders, such as appendicitis, gallbladder disturbances, or malnutrition.

10 The term "gastroparesis," as used herein, is defined as a paralysis of the stomach brought about by a motor abnormality in the stomach or as a complication of diseases, such as diabetes, progressive systemic sclerosis, anorexia nervosa, or myotonic dystrophy.

15 The term "constipation," as used herein, is defined as a condition characterized by infrequent or difficult evacuation of feces resulting from conditions, such as lack of intestinal muscle tone or intestinal spasticity.

 The phrase "post-operative ileus," as used herein, is defined as an obstruction in the intestine due to a disruption in muscle tone following surgery.

20 The terms "preventing" and "prevention," as used herein, are defined respectively as to stop or hinder and the act of stopping or hindering conditions or disorders in a patient who is at risk of suffering from such conditions or disorders, including, but not limited to, patients who suffer from stress.

 The phrase "intestinal pseudo-obstruction," as used herein, is defined as a condition characterized by constipation, colicky pain, and vomiting, but without evidence of physical obstruction.

25 The phrase "gastric hyperacidity," as used herein, is defined as a condition characterized by an abnormally high degree of acidity in the gastric juices.

30 The phrase "therapeutically effective amount," as used herein, is defined as that amount of a therapeutic agent, which, alone or in combination with other drugs, provides a therapeutic benefit in the prevention, treatment, or management of gastrointestinal disorders, but not limited to, gastrointestinal motility dysfunction, GERD, emesis, gastrointestinal ulcers, pathological hypersecretory conditions, gastric hyperacidity, or symptoms thereof. Different therapeutically effective amounts may be applicable for each disorder, as will be readily known by those of ordinary skill in the art.

152020 - 152020

Optically pure (+) norcispripide may be obtained from a racemic mixture of cispripide, the chemical synthesis of which can be performed according to the method described in European Patent Application No. 0,076,530 A2 published April 13, 1983, or U.S. Patent Nos. 4,962,115, 5,057,525 or 5,137,896, the disclosures of which are each incorporated herein by express reference thereto. *See also*, Van Daele, *et al.*, *Drug Development Res.*, 8:225-232 (1986). The metabolism of cispripide to norcispripide is described in Meuldermans, W., *et al.*, *Drug Metab. Dispos.*, 16(3):410-419 (1988) and Meuldermans, W., *et al.*, *Drug Metab. Dispos.*, 16(3):403-409 (1988). The preparation of racemic norcispripide is also known to those of ordinary skill in the art, particularly in view of EP 0,076,530 A2 and U.S. Patent No. 5,137,896 to Van Daele, the disclosures of which are each incorporated herein by express reference thereto. Optically pure stereoisomers of the chiral compounds discussed herein may also be obtained from the racemic mixture by resolution of the enantiomers using conventional means, for example, from an optically active resolving acid. The resolution of racemic compounds is also known to those of ordinary skill in the art, particularly from Jacques, J., *et al.*, *Enantiomers, Racemates and Resolutions*, Wiley-Interscience, New York (1981); Wilen, S. H., *et al.*, *Tetrahedron*, 33:2725 (1977); Eliel, E. L. *Stereochemistry of Carbon Compounds*, McGraw-Hill, NY, (1962); Wilen, S. H. *Tables of Resolving Agents and Optical Resolutions*, E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN, p. 268 (1972).

In addition to separation techniques, such as those described above, optically pure compounds may be synthesized by stereospecific synthesis using methodology well known to those of ordinary skill in the art. Chiral synthesis can result in products of high enantiomeric purity. However, in some cases, the enantiomeric purity of the product is not sufficiently high. The skilled artisan will appreciate that the separation methods described above may be used to further enhance the enantiomeric purity of the active stereoisomers which are obtained by chiral synthesis.

For example, optically pure (+) norcisapride may also be prepared from the racemic norcisapride mixture by enzymatic biocatalytic resolution. This synthesis is known to those of ordinary skill in the art, particularly from U.S. Patent Nos. 5,057,427 and 5,077,217, the disclosures of which are each incorporated herein by express reference thereto.

20-65
64

5 The magnitude of a prophylactic or therapeutic dose of the active ingredients discussed herein, e.g., (+) norcisapride, a 5-HT₃ antagonist, a 5-HT₄ agonist or antagonist, a proton pump inhibitor, or an H₂ receptor antagonist, in the acute or chronic management of diseases and disorders described herein, will vary with the nature and severity of the condition to be prevented, treated, or managed and the route of administration. For example, oral, mucosal (including rectal), parenteral (including subcutaneous, intramuscular, bolus injection, and intravenous), sublingual, transdermal, nasal, buccal, and like may be employed. Dosage forms include tablets, caplets, troches, lozenges, 10 dispersions, suspensions, suppositories, solutions, capsules, soft elastic gelatin capsules, patches, and the like. The dose, and perhaps the dose frequency, will also vary according to the age, body weight, and response of the individual patient. Suitable dosing regimens can be readily selected by those skilled in the art with due consideration of such factors.

15 In general, the total daily dosage of a 5-HT₃ receptor antagonist, a 5-HT₄ receptor agonist, or a therapeutic agent providing dual action at 5-HT₃ receptors and 5-HT₄ receptors for the conditions described herein, is from about 0.5 mg to about 500 mg, preferably from about 1 mg to about 350 mg, and more preferably from about 2 mg to about 250.

20 Suitable daily dosage ranges of proton pump inhibitors can be readily determined by those skilled in the art. In general the total daily dosage of a proton pump inhibitor for the conditions described herein, such as lansoprazole, pantoprazole, rabeprazole, omeprazole, or optically pure stereoisomers or active metabolites thereof, is from about 1 mg to about 200 mg, preferably from about 5 mg to about 150 mg, and more preferably from about 10 mg to about 100 mg.

25 In addition, suitable daily dosage ranges of H₂ receptor antagonists can be readily determined by those skilled in the art. For example, see the *Physician's Desk Reference*®, Medical Economics Co., Inc., 52nd Edition (1999) for suitable dosages presently used for known H₂ receptor antagonists. For example, ranitidine may be administered using an oral daily dose range from about 1 mg to about 800 mg, preferably 30 from about 100 mg to about 600 mg, and more preferably from about 250 mg to about 500 mg. For cimetidine, the oral daily dose range may be from about 1 mg to about 2400 mg, preferably from about 400 mg to about 1600 mg, more preferably from about 600 mg to about 1000 mg. For famotidine, the oral daily dose range may be from about 1 mg to about

2168
65

200 mg, preferably from about 10 mg to about 80 mg, more preferably from about 15 mg to about 50 mg. For nizatidine, the oral daily dose range may be from about 1 mg to about 600 mg, preferably from about 50 mg to about 500 mg, more preferably from about 250 mg to about 350 mg.

5 In managing the patient, the therapy may be initiated at a lower dose, e.g., from about 0.5 mg to about 10 mg of (+) norecisapride and from about 1 mg to about 5 mg of a proton pump inhibitor or from about 1 mg to about 5 mg of an H₂ receptor antagonist, and increased up to the recommended daily dose or higher depending on the patient's global response. It is further recommended that children, patients over 65 years, and those with
10 impaired renal or hepatic function, initially receive low doses, and that they be titrated based on individual response(s) and blood level(s). It may be necessary to use dosages of each active ingredient outside the ranges disclosed herein in some cases, as will be apparent to those of ordinary skill in the art. Furthermore, it is noted that the clinician or treating
15 physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

Pharmaceutical compositions for use in the present invention can comprise a 5-HT₃ receptor antagonist and 5-HT₄ receptors agonist, or a therapeutic agent providing dual action at 5-HT₃ receptors and 5-HT₄ receptors with either a proton pump inhibitor or
20 H₂ receptor antagonist as the active ingredients, and may additionally contain a pharmaceutically acceptable carrier, and optionally, other therapeutic ingredients.

In one embodiment, the pharmaceutical compositions comprise three therapeutic agents, a 5-HT₃ receptor antagonist and a 5-HT₄ receptor agonist with a proton pump inhibitor or H₂ receptor antagonist, and may additionally contain a pharmaceutically
25 acceptable carrier, and optionally, other therapeutic ingredients. In a preferred embodiment, the pharmaceutical compositions comprise two therapeutic agents, one providing dual action at 5-HT₃ receptors and 5-HT₄ receptors, and the other is either a proton pump inhibitor or H₂ receptor antagonist.

As used herein, the term "pharmaceutically acceptable salt" refers to a salt
30 prepared from pharmaceutically acceptable non-toxic acids including, but not limited to inorganic acids, organic acids, solvates, hydrates, or clathrates thereof. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, and phosphoric. Appropriate organic acids may be selected, for example, from aliphatic, aromatic,

carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, benzoic camphorsulfonic, citric, fumaric, gluconic, isethionic, lactic, malic, mucic, tartaric, para-toluenesulfonic, glycolic, glucuronic, maleic, furoic, glutamic, benzoic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic, stearic, sulfanilic, alginic, galacturonic, and the like. Particularly preferred acids are hydrobromic, hydrochloric, phosphoric, and sulfuric acids. In one embodiment, a 5-HT₃ receptor antagonist, a 5-HT₄ receptor agonist, or a therapeutic agent providing dual action as a 5-HT₃ receptor antagonist and a 5-HT₄ receptor agonist are administered as the free base or hydrate. For example, optically pure (+) norcisapride is administered as a free base or hydrate.

The invention also contemplates the use of active agents or compounds that are acidic, in which salts may be prepared from pharmaceutically acceptable non-toxic bases including organic, inorganic bases, solvates, hydrates, or clathrates thereof. Such inorganic bases include, but are not limited to, metallic salts of aluminum, calcium, lithium, magnesium, potassium, sodium, and zinc. Appropriate organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.

In practical use, the active agents in the pharmaceutical compositions of the invention can be combined as the active ingredients in intimate admixture with a pharmaceutically acceptable carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms and comprises a number of components depending on the form of preparation desired for administration. The compositions of the present invention include, but are not limited to, suspensions, solutions and elixirs; aerosols; or excipients, including, but not limited to, starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like. Preferably, the pharmaceutical composition is in the form of an oral preparation. Because of their ease of administration, tablets and capsules are preferred and represent the most advantageous oral dosage unit form, in which case solid pharmaceutical excipients are employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques.

Preferably, the oral pharmaceutical compositions of the present invention may be administered in single or divided doses, from one to four times a day. The oral

dosage forms may be conveniently presented in unit dosage forms and prepared by any methods well known in the art of pharmacy.

Pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete pharmaceutical unit dosage forms, such as capsules, cachets, soft elastic gelatin capsules, tablets, caplets, or aerosols sprays, each containing a predetermined amount of the active ingredients, as a powder or granules, or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil liquid emulsion. Such compositions may be prepared by any methods well known in the art of pharmacy, but all methods include the step of bringing into association one or more active ingredient(s) with the carrier. In general, the compositions are prepared by uniformly and intimately admixing the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation. Oral solid preparations are preferred over oral liquid preparations. One preferred oral solid preparation is capsules, but the most preferred oral solid preparation is tablets.

For example, a tablet may be prepared by compression or molding, optionally, with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form, such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, granulating agent, surface active or dispersing agent, or the like. Molded tablets may be made by molding, in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Preferably, each tablet, cachet, caplet, or capsule contains from about 0.5 mg to about 500 mg of optically pure (+) norcisapride, more preferably from about 1 mg to about 350 mg, in combination with from about 1 mg to about 200 mg of a proton pump inhibitor or from about 1 mg to about 2400 mg of an H_2 receptor antagonist.

Pharmaceutical compositions of the present invention suitable for oral administration may be formulated as a pharmaceutical composition in a soft elastic gelatin capsule unit dosage form by using conventional methods well known in the art. *See, e.g., Ebert, Pharm. Tech*, 1(5):44-50 (1977). Soft elastic gelatin capsules have a soft, globular gelatin shell somewhat thicker than that of hard gelatin capsules, wherein a gelatin is plasticized by the addition of plasticizing agent, *e.g.,* glycerin, sorbitol, or a similar polyol. The hardness of the capsule shell may be changed by varying the type of gelatin used and

24-89
68

the amounts of plasticizer and water. The soft gelatin shells may contain a preservative, such as methyl- and propylparabens and sorbic acid, to prevent the growth of fungi. The active ingredient may be dissolved or suspended in a liquid vehicle or carrier, such as vegetable or mineral oils, glycols, such as polyethylene glycol and propylene glycol, triglycerides, surfactants, such as polysorbates, or a combination thereof.

In addition to the common dosage forms set out above, the compounds of the present invention may also be administered by controlled release means or delivery devices that are well known to those of ordinary skill in the art, such as those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, the disclosures of which are each incorporated herein by express reference thereto. These pharmaceutical compositions can be used to provide slow or controlled-release of one or more of the active ingredients therein using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or the like, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, may be readily selected for use with the pharmaceutical compositions of the invention. Thus, single unit dosage forms suitable for oral administration, such as tablets, capsules, gelcaps, caplets, and the like, that are adapted for controlled-release are encompassed by the present invention.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations may include: 1) extended activity of the drug; 2) reduced dosage frequency; and 3) increased patient compliance.

Most controlled-release formulations are designed to initially release an amount of drug that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic effect over an extended period of time. In order to maintain this constant level of drug in the body, the

25-10
69

drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body.

5 The controlled-release of an active ingredient may be stimulated by various inducers, for example pH, temperature, enzymes, water, or other physiological conditions or compounds. The term "controlled-release component" in the context of the present invention is defined herein as a compound or compounds, including, but not limited to, polymers, polymer matrices, gels, permeable membranes, liposomes, microspheres, or the like, or a combination thereof, that facilitates the controlled-release of the active ingredient.

10 Pharmaceutical compositions of the present invention may also be formulated for parenteral administration by injection (subcutaneous, bolus injection, intramuscular, or intravenous), and may be dispensed in a unit dosage form, such as a multidose container or an ampule. Such compositions for parenteral administration may be in the form of suspensions, solutions, emulsions, or the like in aqueous or oily vehicles, and
15 in addition to the active ingredients may contain one or more formulary agents, such as dispersing agents, suspending agents, stabilizing agents, preservatives, and the like.

Another route of administration is transdermal delivery, for example, via an abdominal skin patch.

20 The invention is further defined by reference to the following examples, describing in detail the preparation of the compound and the compositions of the present invention, as well as their utility. It will be apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the purpose and interest of this invention.

25 5 EXAMPLES

5.1 EXAMPLE 1: Receptor Binding

5-HT₁ Receptor

30 Racemic norcisapride, racemic cisapride and their (+)- and (-)- stereoisomers were tested (Cerep, Celle l'Evescault, France) for binding to 5-HT₁ receptor subtypes derived from N1E-115 cells.

Following incubation with the appropriate ligands, the preparations were rapidly filtered under vacuum through GF/B glass fiber filters and washed with ice-cold

26-71
70

buffer using a Brandel or Packard cell harvester. Bound radioactivity was determined with a liquid scintillation counter (LS 6000, Beckman) using a liquid scintillation cocktail (Formula 989).

5 Specific radioligand binding to the receptor was defined as the difference between total binding and nonspecific binding determined in the presence of an excess of unlabeled ligand. Results were expressed as a percent inhibition of specific binding obtained in the presence of the compounds. IC_{50} were determined using concentrations ranging from 3×10^{-10} M to 10^{-5} M to obtain full competition curves and were calculated by
10 non-linear regression analysis. The results are shown in Tables 3 and 4 below.

Other active ingredients, such as granisetron, metoclopramide, ondansetron, renzapride, zacopride, tropisetron, and the like, can be tested using the methods described above.

5-HT₄ Receptor

15 Racemic norcisapride, racemic cisapride and their (+)- and (-)- stereoisomers were tested (Cerep, Celle l'Evescault, France) for binding to 5-HT₄ receptor subtypes derived from guinea-pig striata.

Following incubation with the appropriate ligands, the preparations were
20 rapidly filtered under vacuum through GF/B glass fiber filters and washed with ice-cold buffer using a Brandel or Packard cell harvester. Bound radioactivity was determined with a liquid scintillation counter (LS 6000, Beckman) using a liquid scintillation cocktail (Formula 989).

Specific radioligand binding to the receptor was defined as the difference
25 between total binding and nonspecific binding determined in the presence of an excess of unlabeled ligand. Results were expressed as a percent inhibition of specific binding obtained in the presence of the compounds. IC_{50} were determined using concentrations ranging from 3×10^{-10} M to 10^{-5} M to obtain full competition curves and were calculated by non-linear regression analysis. The results are shown in Tables 3 and 4 below.

30 Other active ingredients, such as cisapride and the like, can be tested using the methods described above.

27 412
71
VIRVIR

TABLE 3

IC₅₀ (nM) Values for Binding to 5-HT₃ and 5-HT₄ Sites

5

Compound	5-HT ₃	5-HT ₄	5-HT ₃ /5-HT ₄ Ratio
(±) Norcisapride	8.2	686	0.012
(+) Norcisapride	4.5	331	0.014
(-) Norcisapride	30.4	1350	0.023

10

TABLE 4

IC₅₀ (nM) Values for Binding to 5-HT₃ and 5-HT₄ Sites

15

Compound	5-HT ₃	5-HT ₄	5-HT ₃ /5-HT ₄ Ratio
(±) Cisapride	365	169	2.2
(+) Cisapride	310	340	0.9
(-) Cisapride	2790	199	14.0

Agonist activity at 5-HT₄ receptor sites may also be assessed using an assay based on the ability of active compounds to increase cyclic AMP production in mouse embryo colliculi neurones grown in tissue culture (See Dumuis *et al.*, *N. S. Arch. Pharmacol.*, 340:403-410 (1989)).

25

30

28-70
92

5.2 EXAMPLE 2: Oral Formulation

Tablets

Formula	Quantity per Tablet in mg.		
	A	B	C
Active Ingredients:	5.0	10.0	25.0
(+) Norcisapride			
Lansoprazole	5.0	15.0	30.0
Lactose BP	57.0	92.0	107.0
Starch BP	20.0	20.0	25.0
Microcrystalline Cellulose	10.0	10.0	10.0
Hydrogenated Vegetable Oil	1.5	1.5	1.5
Polyvinylpyrrolidinone	1.5	1.5	1.5
Compression Weight	100.0	150.0	200.0

The active ingredients, (+) norcisapride and lansoprazole, are sieved through a suitable sieve and blended with the lactose until a uniform blend is formed. Suitable volumes of water are added and the powders are granulated. After drying, the granules are then screened and blended with the remaining excipients. The resulting granules are then compressed into tablets of desired shape. Tablets of other strengths may be prepared by altering the ratio of active ingredient to the excipient(s) or the compression weight.

2974
73

5.3 EXAMPLE 3: Oral Formulation

Tablets

Formula	Quantity per Tablet in mg.		
	A	B	C
Active Ingredients: (+) Norcisapride	5.0	10.0	25.0
Famotidine	10.0	20.0	40.0
Lactose BP	38.5	73.5	43.5
Starch BP	30.0	30.0	60.0
Pregelatinized Maize Starch BP	15.0	15.0	30.0
Magnesium Stearate BP	1.5	1.5	1.5
Compression Weight	100.0	150.0	540.0

The active ingredients, (+) norcisapride and famotidine, are sieved through a suitable sieve and blended with lactose, starch, and pregelatinized maize starch until a uniform blend is formed. Suitable volumes of water are added and the powders are granulated. After drying, the granules are then screened and blended with the remaining excipients. The resulting granules are then compressed into tablets of desired shape. Tablets of other strengths may be prepared by altering the ratio of active ingredient to the excipient(s) or the compression weight.

39-73
79

5.4 EXAMPLE 4: Oral Formulation

Tablets

5	Formula	Quantity per Tablet in mg.		
		A	B	C
	Active Ingredient: (+) Norcisapride	5.0	10.0	25.0
	Lactose BP	48.5	43.5	78.5
	Starch BP	30.0	30.0	30.0
10	Pregelatinized Maize Starch BP	15.0	15.0	15.0
	Magnesium Stearate BP	1.5	1.5	1.5
	Compression Weight	100.0	100.0	150.0

15 The active ingredient, (+) norcisapride, is sieved through a suitable sieve and blended with lactose, starch, and pregelatinized maize starch until a uniform blend is formed. Suitable volumes of water are added and the powders are granulated. After drying, the granules are then screened and blended with the remaining excipients. The resulting granules are then compressed into tablets of desired shape. Tablets of other strengths may be prepared by altering the ratio of active ingredient to the excipient(s) or the compression weight.

20 While the present invention has been described with respect to the particular embodiments, it will be apparent to those skilled in the art that various changes and modifications may be made without departing from the spirit and scope of the invention as defined in the claims. Such modifications are also intended to fall within the scope of the appended claims.

30

31-16
75

THE CLAIMS

What is claimed is:

- 5 1. A method of treating gastrointestinal disorders in a patient which comprises administering a therapeutically effective amount of one or more agent(s) or compound(s) that simultaneously or sequentially act on 5-HT₃ receptors, 5-HT₄ receptors, and either proton pumps or H₂ receptors, or an optically pure stereoisomer or active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- 10 2. A method of treating gastrointestinal disorders in a patient which comprises administering a therapeutically effective amount of an agent or compound that antagonizes 5-HT₃ receptors and agonizes 5-HT₄ receptors, and a therapeutically effective amount of at least one of a proton pump inhibitor or an H₂ receptor antagonist, or an optically pure
15 stereoisomer or active metabolite thereof, or a pharmaceutically acceptable salt thereof.
3. A method of treating gastrointestinal disorders in a patient which comprises administering to said patient a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer, and a
20 therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
4. The method of claim 3, wherein the patient is a human.
- 25 5. The method of claim 3, wherein the gastrointestinal disorder is gastrointestinal motility dysfunction.
6. The method of claim 5, wherein the gastrointestinal motility dysfunction is
30 selected from the group consisting of dyspepsia, gastroparesis, constipation, post-operative ileus, and intestinal pseudo obstruction.

32-77-
76

7. The method of claim 3, wherein the gastrointestinal disorder is gastro-esophageal reflux disease, emesis, gastrointestinal ulcers, pathological hypersecretory condition, and gastric hyperacidity.
- 5 8. The method of claim 7, wherein the pathological hypersecretory condition is Zollinger-Ellison Syndrome.
- 10 9. The method of claim 7, wherein the gastric hyperacidity is selected from the group consisting of heartburn, acid indigestion, sour stomach, erosive esophagitis, and upset stomach.
- 15 10. The method of claim 3, wherein the amount of (+) norcisapride administered is from about 0.5 mg to about 500 mg.
- 20 11. The method of claim 10, wherein the amount of (+) norcisapride administered is from about 1 mg to about 350 mg.
- 25 12. The method of claim 3, wherein the proton pump inhibitor is administered and is selected from the group consisting of omeprazole, lansoprazole, rabeprazole, pantoprazole, hydroxy-omeprazole, desmethyl-pantoprazole, hydroxy-lansoprazole, and an optically pure stereoisomer thereof.
- 30 13. The method of claim 12, wherein the amount of proton pump inhibitor is from about 1 mg to about 200 mg.
14. The method of claim 13, wherein the amount of proton pump inhibitor is from about 5 mg to about 150 mg.
15. The method of claim 3, wherein the H₂ receptor antagonist is administered and is selected from the group consisting of cimetidine, ranitidine, famotidine, nizatidine, and an optically pure stereoisomer or an active metabolite thereof.

33-18
77

16. The method of claim 15, wherein the amount of H₂ receptor antagonist is from about 1 mg to about 2400 mg.
- 5 17. The method of claim 3, wherein at least one of (+) norcisapride and the proton pump inhibitor is administered orally.
18. The method of claim 17, wherein (+) norcisapride and the proton pump are orally administered as a tablet or a capsule.
- 10 19. The method of claim 3, wherein at least one of (+) norcisapride and the H₂ receptor antagonist is administered orally.
- 15 20. The method of claim 3, wherein the proton pump inhibitor or the H₂ receptor antagonist is administered together with (+) norcisapride parenterally, transdermally, rectally or sublingually.
- 20 21. The method of claim 3, which comprises administering either the proton pump inhibitor or H₂ receptor antagonist, and the (+) norcisapride, concurrently or sequentially.
- 25 22. A method of treating gastrointestinal motility dysfunction in a patient which comprises administering to said patient a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer and a therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.
- 30 23. A method of treating emesis in a patient which comprises administering to said patient in need of such prevention, treatment, or management a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer and a therapeutically effective amount of at least

one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

5 24. A method of treating gastro-esophageal reflux disease in a patient which
comprises administering a therapeutically effective amount of (+) norcisapride, or a
pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer and a
therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor
antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a
10 pharmaceutically acceptable salt thereof.

25. A method of treating gastro-esophageal reflux disease in a patient which
comprises administering to said patient in need of such treatment a therapeutically effective
amount of: (a) cisapride, or a pharmaceutically acceptable salt or an optically pure
15 stereoisomer thereof; (b) ondansetron, or a pharmaceutically acceptable salt or an optically
pure stereoisomer thereof; and (c) a therapeutically effective amount of at least one of a
proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an
active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20 26. The method of claim 25, wherein optically pure (+) cisapride or optically
pure (-) cisapride, or a pharmaceutically acceptable salt thereof, is administered.

27. The method of claim 25 or 26, wherein optically pure R(+) ondansetron is
administered.

25 28. The method of claim 25, wherein the proton pump inhibitor administered is
selected from the group consisting of omeprazole, lansoprazole, pantoprazole, rabeprazole,
hydroxy-omeprazole, hydroxy-lansoprazole, the carboxylic derivative of omeprazole, and
desmethyl-pantoprazole.

30 29. The method of claim 25, wherein the amount of proton pump inhibitor
administered is from about 1 mg to about 200 mg.

34-79
78

35 80
79

30. The method of claim 25, wherein the H₂ receptor antagonist administered is selected from the group consisting of cimetidine, famotidine, ranitidine, nizatidine, and N2-desmethylnizatidine.

5

31. The method of claim 25, wherein the amount of H₂ receptor antagonist administered is from about 1 mg to about 2400 mg.

32. The method of claim 25, wherein at least one of optically pure (+) cisapride, optically pure (-) cisapride, optically pure R(+) ondansetron, and the proton pump inhibitor is administered orally.

10

33. The method of claim 25, wherein at least one of cisapride, optically pure R(+) ondansetron, and the H₂ receptor antagonist is administered orally.

15

34. A method of preventing or managing gastrointestinal disorders in a patient which comprises administering to said patient in need of prevention or management a therapeutically effective amount of a 5-HT₃ antagonist, a 5-HT₄ agonist, and at least one of an H₂ receptor antagonist or a proton pump inhibitor, or an optically pure stereoisomer or active metabolite thereof, or a pharmaceutically acceptable salt thereof.

20

35. A method of preventing or managing gastrointestinal disorders in a patient which comprises administering a therapeutically effective amount of an agent or compound that acts on 5-HT₃ receptors and 5-HT₄ receptors, and a therapeutically effective amount of at least one of a proton pump or an H₂ receptor, or an optically pure stereoisomer or active metabolite thereof, or a pharmaceutically acceptable salt thereof.

25

36. A method of preventing or managing gastrointestinal disorders in a patient which comprises administering to said patient a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer, and a therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

30

36-81
80

37. A method of preventing or managing gastrointestinal motility dysfunction in a patient which comprises administering to said patient a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer and a therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

38. A method of preventing or managing emesis in a patient which comprises administering to said patient in need of such prevention, treatment, or management a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer and a therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

39. A method of preventing or managing gastro-esophageal reflux disease in a patient which comprises administering a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer and a therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

40. A method of preventing or managing gastro-esophageal reflux disease in a patient which comprises administering to said patient in need of such prevention or management a therapeutically effective amount of: (a) cisapride, or an optically pure stereoisomer or a pharmaceutically acceptable salt thereof; (b) ondansetron, or a pharmaceutically acceptable salt or an optically pure stereoisomer thereof; and (c) a therapeutically effective amount of at least one of a proton pump inhibitor, an H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

41. The method of claim 40, wherein optically pure (+) cisapride or optically pure (-) cisapride, or a pharmaceutically acceptable salt thereof, is administered.

34-82
81

42. The method of claim 40, wherein optically pure R(+) ondansetron, or a pharmaceutically acceptable salt thereof, is administered.

5 43. The method of claim 40, wherein the proton pump inhibitor administered is selected from the group consisting of omeprazole, lansoprazole, pantoprazole, rabeprazole, hydroxy-omeprazole, hydroxy-lansoprazole, the carboxylic acid derivative of omeprazole, and desmethyl-pantoprazole.

10 44. The method of claim 40, wherein the amount of proton pump inhibitor administered is from about 1 mg to about 200 mg.

15 45. The method of claim 40, wherein the H₂ receptor antagonist administered is selected from the group consisting of cimetidine, famotidine, ranitidine, nizatidine, and N2-desmethylnizatidine.

46. The method of claim 40, wherein the amount of H₂ receptor antagonist administered is from about 1 mg to about 2400 mg.

20 47. The method of claim 40, wherein at least one of cisapride, R(+) ondansetron, and the proton pump inhibitor is administered orally.

48. The method of claim 40, wherein at least one of cisapride, R(+) ondansetron, and the H₂ receptor antagonist is administered orally.

25 49. A pharmaceutical composition adapted for the treatment of a patient suffering from a gastrointestinal disorder which comprises a therapeutically effective amount of (+) norcisapride, or a pharmaceutically acceptable salt thereof, substantially free of its (-) stereoisomer; and a therapeutically effective amount of at least one of a proton
30 pump inhibitor, H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

50. The pharmaceutical composition of claim 49, wherein the proton pump inhibitor is present and is selected from the group consisting of omeprazole, pantoprazole, rabeprazole, lansoprazole, hydroxy-omeprazole, hydroxy-lansoprazole, the carboxylic acid derivative of omeprazole, and desmethyl-pantoprazole.

51. The pharmaceutical composition of claim 49, wherein the H₂ receptor antagonist is present and is selected from the group consisting of cimetidine, ranitidine, famotidine, nizatidine, and N2-desmethylnizatidine.

52. A pharmaceutical composition adapted for the treatment of a patient suffering from a gastrointestinal disorder which comprises: (a) a therapeutically effective amount of an optically pure stereoisomer of cisapride, or a pharmaceutically acceptable salt thereof; (b) a therapeutically effective amount of optically pure R(+) ondansetron, or a pharmaceutically acceptable salt thereof; and (c) a therapeutically effective amount of at least one of a proton pump inhibitor, H₂ receptor antagonist, or an optically pure stereoisomer or an active metabolite thereof, or a pharmaceutically acceptable salt thereof.

53. The pharmaceutical composition of claim 52, wherein the proton pump inhibitor is present and is selected from the group consisting of omeprazole, pantoprazole, rabeprazole, lansoprazole, hydroxy-omeprazole, hydroxy-lansoprazole, the carboxylic acid derivative of omeprazole, and desmethyl-pantoprazole.

54. The pharmaceutical composition of claim 52, wherein the H₂ receptor antagonist is present and is selected from the group consisting of cimetidine, ranitidine, famotidine, nizatidine, and N2-desmethylnizatidine.

55. The method of claim 12 wherein the proton pump inhibitor is optically pure (+) pantoprazole, optically pure (-) pantoprazole, optically pure (+) rabeprazole, optically pure (-) rabeprazole, optically pure (+) lansoprazole, optically pure (-) lansoprazole, optically pure (+) omeprazole, or optically pure (-) omeprazole.

~~39-84~~
83

56. The pharmaceutical composition of claim 50 or 53 wherein the proton pump inhibitor is (+) pantoprazole, optically pure (-) pantoprazole, optically pure (+) rabeprazole, optically pure (-) rabeprazole, optically pure (+) lansoprazole, optically pure (-) lansoprazole, optically pure (+) omeprazole, or optically pure (-) omeprazole.

10

15

20

25

30

40 85
84

**METHODS AND COMPOSITIONS
USING (+) NORCISAPRIDE IN COMBINATION WITH
PROTON PUMP INHIBITORS OR H₂ RECEPTOR ANTAGONISTS**

ABSTRACT

5

The invention relates to methods and compositions for the prevention, treatment, or management of gastrointestinal disorders or symptoms thereof, employing two or more agents or compounds to provide a triple site action on 5-HT₃ receptors, 5-HT₄ receptors, and at least one of H₂ receptors and proton pumps.

10

15

20

25

30

41 86
85

ESTADOS UNIDOS DE AMÉRICA

A TODOS A QUIENES LLEGUE EL PRESENTE:

DEPARTAMENTO DE COMERCIO DE LOS ESTADOS UNIDOS

Oficina de Marcas y Patentes

14 de marzo de 2000..

LA PRESENTE CERTIFICA QUE LA ADJUNTA ES UNA COPIA TOMADA DE LOS REGISTROS DE LA OFICINA DE MARCAS Y PATENTES DE LOS ESTADOS UNIDOS, DE LOS DOCUMENTOS DE LA SOLICITUD DE PATENTE ABAJO IDENTIFICADA Y LA CUAL CUMPLE CON LOS REQUISITOS PARA QUE LE SEA CONFERIDA UNA FECHA DE RADICACIÓN SEGÚN 35 USC 111.

NÚMERO DE SOLICITUD 60/122,394

FECHA DE RADICACIÓN: 2 de marzo de 1999

Por Autoridad del
COMISIONADO DE MARCAS Y PATENTES

(FIRMA) P. SWAIN
Oficial de Certificaciones

(SELLO DORADO DE OBLEA)

42
86
87

No. 0032620-3

Estados Unidos de América
DEPARTAMENTO DE ESTADO

A todos a Quienes Llegue el Presente, Os Saludo:

Certifico que el documento adjunto se encuentra bajo el sello de la Oficina de Marcas y Patentes del Departamento de Comercio de los Estados Unidos de América y que a dicho sello debe dársele plena fe y crédito.

En Testimonio de lo cual, Yo, Madeleine K. Albright, Secretaria de Estado, he ordenado al Oficial de Autenticaciones suscribir mi nombre y colocar el sello del Departamento de Estado en la Ciudad de Washington, Distrito de Columbia, hoy 28 de marzo de 2000.

Madeleine K. Albright - Seecretaria de Estado
Patrick O. Harchett - Asistente del Oficial de
Autenticaciones. Departamento de Estado

EN ESPAÑOL: La legalización de la firma y cargo anterior por parte del Consulado de Colombia en Washington, D.C. de fecha 29 de marzo de 2000, bajo el No. 757; y por el Jefe de Legalizaciones del Ministerio de Relaciones Exteriores, de fecha 2 de mayo de 2000, bajo el No. 256129.

43 88
89

**METODOS Y COMBINACIONES QUE UTILIZAN (+) NORCISAPRIDA EN
COMBINACION CON INHIBIDORES DE BOMBA DE PROTONES O
ANTAGONISTAS DEL RECEPTOR H₂.**

REIVINDICACIONES:

1. Un método para tratar trastornos gastrointestinales en un paciente, que consiste en administrar una cantidad terapéuticamente efectiva de uno o más agentes o compuestos que simultánea o secuencialmente actúen en receptores 5-HT₃, receptores 5-HT₄ y bien sea bombas de protones o receptores H₂, o un estereoisómero ópticamente puro o metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.
2. Un método para tratar trastornos gastrointestinales en un paciente, que consiste en administrar una cantidad terapéuticamente efectiva de un agente o compuesto que antagonice con receptores 5-HT₃ y agonice con receptores 5-HT₄, y una cantidad terapéuticamente efectiva de al menos un inhibidor de bomba de protones o un antagonista del receptor H₂, o un estereoisómero ópticamente puro o metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.
3. Un método para tratar trastornos gastrointestinales en un paciente que consiste en administrar a dicho paciente una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-), y una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, un antagonista del

receptor H₂, o un estereoisómero ópticamente puro o metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

4. El método de la reivindicación 3, en donde el paciente es un ser humano.
5. El método de la reivindicación 3, en donde el trastorno gastrointestinal es disfunción de la motilidad gastrointestinal.
6. El método de la reivindicación 5, en donde la disfunción de la motilidad gastrointestinal es seleccionada del grupo consistente de dispepsia, gastroparesis, constipación íleo posoperativo y pseudo obstrucción intestinal.
7. El método de la reivindicación 3, en donde el trastorno gastrointestinal es enfermedad de reflujo gastroesofágico, emesis, úlceras gastrointestinales, condición hipersecretora patológica e hiperacidez gástrica.
8. El método de la reivindicación 7, en donde la condición hipersecretora patológica es Síndrome de Zollinger-Ellison.
9. El método de la reivindicación 7, en donde la hiperacidez gástrica es seleccionada del grupo consistente de pirosis, indigestión ácida, estómago amargo, esofagitis erosiva y malestar estomacal.
10. El método de la reivindicación 3, en donde la cantidad de (+)norcisaprida administrado es de aproximadamente 0.5 mg a aproximadamente 500 mg.
11. El método de la reivindicación 10, en donde la cantidad de (+)norcisaprida administrada es de aproximadamente 1 mg a aproximadamente 350 mg.
12. El método de la reivindicación 3, en donde el inhibidor de bomba de protones es administrado y seleccionado del grupo consistente de omeprazol, lasoprazol,

45 89

rabeprazol, pantoprazol, hidroxi-omeprazol, desmetil-pantoprazol, hidroxi-lansoprazol y un estereoisómero ópticamente puro de lo mismo.

13. El método de la reivindicación 12, en donde la cantidad del inhibidor de bomba de protones es de aproximadamente 1 mg a aproximadamente 200 mg.

14. El método de la reivindicación 13, en donde la cantidad de inhibidor de bomba de protones es de aproximadamente 5 mg a aproximadamente 150mg.

15. El método de la reivindicación 3, en donde el antagonista del receptor H_2 es administrado y seleccionado del grupo consistente de cimetidina, ranitidina, famotidina, nizatidina, y un estereoisómero ópticamente puro o un metabolito activo de lo mismo.

16. El método de la reivindicación 15, en donde la cantidad del antagonista del receptor H_2 es de aproximadamente 1 mg a aproximadamente 2400 mg.

17. El método de la reivindicación 3, en donde al menos uno de (+)norcisaprida y el inhibidor de bomba de protones es administrado por vía oral.

18. El método de la reivindicación 17, en donde (+)norcisaprida y la bomba de protones son administrados por vía oral como tableta o cápsula.

19. El método de la reivindicación 3, en donde al menos uno de (+)norcisaprida y el antagonista del receptor H_2 es administrado por vía oral.

20. El método de la reivindicación 3, en donde el inhibidor de bomba de protones o el antagonista del receptor H_2 es administrado junto con (+)norcisaprida por vía parenteral, transdermal, rectal o sublingual.

46
90

21. El método de la reivindicación 3, que consiste en administrar bien sea el inhibidor de bomba de protones o el antagonista del receptor H₂, y el (+)norcisaprida, de manera concurrente o secuencial.

22. Un método para tratar disfunción de la motilidad gastrointestinal en un paciente que consiste en administrar a dicho paciente una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-) y una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H₂, o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

23. Un método para tratar emesis en un paciente, que consiste en administrar a dicho paciente que requiera tal prevención, tratamiento o manejo, una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-) y una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H₂, o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

24. Un método para tratar enfermedad de reflujo gastroesofágico en un paciente, que consiste en administrar a dicho paciente una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-) y una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H₂, o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

25. Un método para tratar enfermedad de reflujo gastroesofágico en un paciente, que consiste en administrar a dicho paciente que requiera tal tratamiento, una

47 92
91

cantidad terapéuticamente efectiva de: (a) cisaprida, o una sal farmacéuticamente aceptable o un estereoisómero ópticamente puro de lo mismo; (b) ondasetron, o una sal farmacéuticamente aceptable o un estereoisómero ópticamente puro de lo mismo; y (c) una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

26. El método de la reivindicación 25, en donde (+)cisaprida ópticamente puro o (-) cisaprida ópticamente puro, o una sal farmacéuticamente aceptable de lo mismo es administrado.

27. El método de la reivindicación 25 ó 26, en donde el R(+)ondasetron es administrado.

28. El método de la reivindicación 25, en donde el inhibidor de bomba de protones administrado es seleccionado del grupo consistente de omeprazol, lansoprazol, pantoprazol, rabeprazol, hidroxí-omeprazol, hidroxí-lansoprazol, el derivado carboxílico de omeprazol y desmetil-pantoprazol.

29. El método de la reivindicación 25, en donde la cantidad de inhibidor de bomba de protones administrado es de aproximadamente 1 mg a aproximadamente 200 mg.

30. El método de la reivindicación 25, en donde el antagonista del receptor H_2 administrado es seleccionado del grupo consistente de cimetidina, famotidina, ranitidina, nizatidina y N2-desmetilnizatidina.

31. El método de la reivindicación 25, en donde la cantidad del antagonista del receptor H_2 administrada es de aproximadamente 1 mg a aproximadamente 2400 mg.

48-92

32. El método de la reivindicación 25, en donde al menos uno de (+)cisaprida ópticamente puro, (-)cisaprida ópticamente puro, R(+)-ondasentron ópticamente puro y el inhibidor de bomba de protones es administrado por vía oral.

33. El método de la reivindicación 25, en donde al menos uno de cisaprida, R(+)-ondasentron ópticamente puro y el antagonista del receptor H_2 es administrado por vía oral.

34. Un método para prevenir o manejar trastornos gastrointestinales en un paciente, que consiste en administrar a dicho paciente que requiera tal prevención o manejo, una cantidad terapéuticamente efectiva de un antagonista $5-HT_3$, un agonista $5-HT_3$, y al menos uno de un antagonista del receptor H_2 o un inhibidor de bomba de protones, o un estereoisómero ópticamente puro o metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

35. Un método para prevenir o manejar trastornos gastrointestinales en un paciente, que consiste en administrar una cantidad terapéuticamente efectiva de un agente o compuesto que actúe en los receptores $5-HT_3$, y en los receptores $5-HT_3$, y una cantidad terapéuticamente efectiva de al menos una bomba de protones o un receptor H_2 , o un estereoisómero ópticamente puro o metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

36. Un método para prevenir o manejar trastornos gastrointestinales en un paciente, que consiste en administrar a dicho paciente una cantidad terapéuticamente efectiva de (+)-norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-), y una cantidad terapéuticamente efectiva de al menos un inhibidor de bomba de protones, un antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

49-94
93

37. Un método para prevenir o manejar disfunción de la motilidad gastrointestinal en un paciente, que consiste en administrar a dicho paciente una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-), y una cantidad terapéuticamente efectiva de al menos un inhibidor de bomba de protones, un antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

38. Un método para prevenir o manejar emesis en un paciente, que consiste en administrar a dicho paciente que requiera tal prevención, tratamiento o manejo, una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-), y una cantidad terapéuticamente efectiva de al menos un inhibidor de bomba de protones, un antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

39. Un método para prevenir o manejar enfermedad de reflujo gastroesofágico en un paciente, que consiste en administrar una cantidad terapéuticamente efectiva de (+)norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-), y una cantidad terapéuticamente efectiva de al menos un inhibidor de bomba de protones, un antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

40. Un método para prevenir o manejar enfermedad de reflujo gastroesofágico en un paciente, que consiste en administrar a dicho paciente que requiera tal prevención, tratamiento o manejo, una cantidad terapéuticamente efectiva de: (a) cisaprida, o una sal farmacéuticamente aceptable o un estereoisómero ópticamente puro de lo mismo; (b) ondasetron, o una sal farmacéuticamente aceptable o un estereoisómero ópticamente puro de lo mismo; y (c) una cantidad terapéuticamente

50 94

efectiva de al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

41. El método de la reivindicación 40, en donde (+)cisaprida ópticamente puro o (-) cisaprida ópticamente puro, o una sal farmacéuticamente aceptable de lo mismo es administrado.

42. El método de la reivindicación 40, en donde el R(+)ondasentron, o una sal farmacéuticamente aceptable de lo mismo es administrado.

43. El método de la reivindicación 40, en donde el inhibidor de bomba de protones administrado es seleccionado del grupo consistente de omeprazol, lansoprazol, pantoprazol, rabeprazol, hidroxi-omeprazol, hidroxi-lansoprazol, el derivado carboxílico de omeprazol y desmetil-pantoprazol.

44. El método de la reivindicación 40, en donde la cantidad de inhibidor de bomba de protones administrado es de aproximadamente 1 mg a aproximadamente 200 mg.

45. El método de la reivindicación 40, en donde el antagonista del receptor H_2 administrado es seleccionado del grupo consistente de cimetidina, famotidina, ranitidina, nizatidina y N2-desmetilnizatidina.

46. El método de la reivindicación 40, en donde la cantidad del antagonista del receptor H_2 administrada es de aproximadamente 1 mg a aproximadamente 2400 mg.

47. El método de la reivindicación 40, en donde al menos uno de (+)cisaprida, R(+)ondasentron, y el inhibidor de la bomba de protones es administrado por vía oral.

51 96
95

48. El método de la reivindicación 40, en donde al menos uno de cisaprida, R(+)-ondasentron, y el antagonista del receptor H_2 es administrado por vía oral.

49. Una composición farmacéutica adaptada para el tratamiento de un paciente que sufra de un trastorno gastrointestinal, que comprende una cantidad terapéuticamente efectiva de (+)-norcisaprida, o una sal farmacéuticamente aceptable de lo mismo, sustancialmente libre de su estereoisómero(-), y una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

50. La composición farmacéutica de la reivindicación 49, en donde está presente el inhibidor de bomba de protones y es seleccionado del grupo consistente de omeprazol, pantoprazol, rabeprazol, lansoprazol, hidroxi-omeprazol, hidroxi-lansoprazol, el derivado de ácido carboxílico de omeprazol, y desmetil-pantoprazol.

51. La composición farmacéutica de la reivindicación 49, en donde está presente el antagonista del receptor H_2 y es seleccionado del grupo consistente de cimetidina, ranitidina, ramotidina, nizatidina y N2-desmetilnizatidina.

52. Una composición farmacéutica adaptada para el tratamiento de un paciente que sufra de un trastorno gastrointestinal, que comprende; (a) una cantidad terapéuticamente efectiva de un estereoisómero ópticamente puro de cisaprida, o una sal farmacéuticamente aceptable de lo mismo; (b) una cantidad terapéuticamente efectiva de R(+)-ondasentron ópticamente puro, o una sal farmacéuticamente aceptable de lo mismo; y (c) una cantidad terapéuticamente efectiva de al menos uno de un inhibidor de bomba de protones, antagonista del receptor H_2 , o un estereoisómero ópticamente puro o un metabolito activo de lo mismo, o una sal farmacéuticamente aceptable de lo mismo.

52 ~~94~~
96

53. La composición farmacéutica de la reivindicación 52, en donde está presente el inhibidor de la bomba de protones y es seleccionado del grupo consistente de omeprazol, pantoprazol, rabeprazol, lansoprazol, hidroxi-omeprazol, hidroxi-lansoprazol, el derivado de ácido carboxílico de omeprazol y desmetil-pantoprazol.

54. La composición farmacéutica de la reivindicación 52, en donde está presente el antagonista del receptor H_2 y es seleccionado del grupo consistente de cimetidina, ranitidina, famotidina, nizatidina y N2-desmetilnizatidina.

55. El método de la reivindicación 12, en donde el inhibidor de la bomba de protones es (+)pantoprazol ópticamente puro, (-)pantoprazol ópticamente puro, (+)rabeprazol ópticamente puro, (-)rabeprazol ópticamente puro, (+)lansoprazol ópticamente puro, (-) lansoprazol ópticamente puro, (+)omeprazol ópticamente puro, o (-)omeprazol ópticamente puro.

56. La composición farmacéutica de la reivindicación 50 ó 53, en donde el inhibidor de la bomba de protones es (+)pantoprazol ópticamente puro, (-) pantoprazol ópticamente puro, (+)rabeprazol ópticamente puro, (-)rabeprazol ópticamente puro, (+)lansoprazol ópticamente puro, (-) lansoprazol ópticamente puro, (+)omeprazol ópticamente puro, o (-) omeprazol ópticamente puro.

Señores
DIVISION DE NUEVAS CREACIONES
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
Santafé de Bogotá, D.C.

SUPERINDUSTRIA Y COMERCIO
Radicacion : 00015142 00000001
Fecha (AMD): 2000-05-05 12:31:16
Tramite : 002 PATENTES D 1 REGISTRO/ 329 COMPLETEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Re.: Código 02-01-09
Expediente No. 00-15.142
Solicitud de Patente
SEPRACOR, INC.

En relación con el Expediente de la referencia y con el fin de dar cumplimiento a la preceptuado por el artículo 4o. literal d.) numeral 3 de la Convención de París adjunto a este escrito sírvanse encontrar la copia certificada de la solicitud de patente de la cual se reivindica prioridad, la cual se encuentra debidamente traducida en sus apartes pertinentes.

De Ustedes Atentamente,

GERMAN CASTILLO GRAU
T.P. No. 2978 del C.S.J.

105

RESUMEN CON EL OBJETO Y LA FINALIDAD DE LA INVENCION

La invención se refiere a método para la prevención, tratamiento y manejo de trastornos gastrointestinales y síntomas de lo mismo. Un método de la invención consiste en la administración de uno o más compuestos que actúen en 5-HT₃, 5-HT₅, y ya sea en bombas de protones o en receptores H₂. Un método específico de la invención consiste en la administración de cantidades terapéuticamente específicas de norcisaprida(+) y al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H₂, y esteroisómeros ópticamente puros

ANEXOS DE LA SOLICITUD

- Certificado de Existencia y Representación Legal cuando el solicitante sea Persona Jurídica ☐
- Los poderes debidamente otorgados, o mención de su protocolización ante la SIC ☐
- Recibo de la tasa respectiva ☐
- Recibo de la tasa por concepto de excedente de palabras en la publicación ☐
- Si se reivindica prioridad Andina: ☐
- Copia de la primera solicitud ☐
- Traducción oficial ☐
- Si se reivindica prioridad Unionista: ☐
- Copia certificada de la primera solicitud: ☐
- Nota: Debe presentarse dentro de los tres (3) meses, contados a partir de la fecha de radicación de la solicitud posterior ☐
- Traducción simple ☐
- Capítulo descriptivo ☐
- Dibujos, planos necesarios ☐
- Capítulo reivindicatorio ☐
- Tarjeta archivo propietarios según formato ☐
- Tarjeta archivo temático según formato ☐
- Extracto para publicación según formato ☐
- Arte final 12 x 12 cms Y 6x6 cms ☐
- Otros Documentos ☐

SOLICITO LA CONCESION DE LA PATENTE DE INVENCION Y DECLARO QUE LA INVENCION ES NUEVA Y DE MI PROPIEDAD

8108
106

y metabolitos activos de lo mismo. La invención también se refiere a composiciones farmacéuticas que son adaptadas para el tratamiento de trastornos gastrointestinales. Una composición farmacéutica específica de la invención comprende cantidades terapéuticas efectivas de norcisaprida(+) y al menos uno de un inhibidor de bomba de protones, un antagonista del receptor H_2 , y esteroisómeros ópticamente puros y metabolitos activos de lo mismo.

2do. EXAMEN DE FORMA PATENTE DE INVENCION

Folio 112

Radicación no 00-15142

Concepto Técnico no 270

Clasificación Internacional de Patentes A 61 K 31/165 II 01/445

Practicado el examen de acuerdo a lo establecido en el Art. 38 de la Decisión 486 se determinó que esta solicitud de Patente de Invención:

- SI ☒ NO ☐ Contiene un resumen de la invención (Art.26-e).
- SI ☒ NO ☐ Contiene el título o nombre de la invención (Art.27-d).
- SI ☒ NO ☐ Contiene la descripción de la invención (Art.26-b).
- SI ☐ NO ☐ NO ES APLICABLE ☒ Contiene los dibujos necesarios para comprender la invención (Art.26-d)
- SI ☒ NO ☐ Contiene una o más reivindicaciones. (Art.26-c).
- SI ☐ NO ☐ NO ES APLICABLE ☒ Han sido desarrollados los productos y/o procedimientos a partir de recursos genéticos o de sus productos derivados de los que cualquiera de los Países Miembros es país de origen. (Art. 26- h).
- SI ☐ NO ☐ NO ES APLICABLE ☒ Los productos y/o procedimientos han sido obtenidos o desarrollados a partir de los conocimientos tradicionales de las comunidades indígenas, afroamericanas, o locales de los países miembros, de acuerdo a lo establecido en la Decisión 391 y sus modificaciones y reglamentaciones vigentes.(Art.26-i).
- SI ☐ NO ☐ La invención se refiere a un producto relativo a un material biológico. (Art.26-j).
- SI ☒ NO ☐ La prioridad coincide con la materia reivindicada.
- SI ☒ NO ☐ **CUMPLE LOS REQUISITOS TÉCNICOS**

OBSERVACIONES Y OTROS: SI ☐ NO ☐ ver hoja(s) anexa(s)

1. Respecto a la primera observación del examinador técnico en el primer examen de forma, el solicitante señala que el significado y alcance de las reivindicaciones 49 y 52 están perfectamente definidos.
2. Se adjunta un nuevo resumen (folio 105), además del Extracto para Publicación y de la tarjeta de Archivo Temático.

EXAMINADOR TÉCNICO

202040

Bogotá D. C., Marzo 1, 2002

Folio 113

2020
Bogotá, D.C.

416
2002-04-29

Doctor(a)
GERMÁN CASTILLO GRAU
Apoderado(a)

REFERENCIA:	Radicación No.	00-15142
	Trámite	002
	Evento	001
	Actuación	416
	Oficio No.	432
	Folios	001

Teniendo en cuenta que la solicitud de privilegio de patente que se tramita bajo el expediente indicado en la referencia, reúne los requisitos de forma establecidos en los artículos 26 y 27 de la Decisión 486 de la Comisión de la comunidad andina, y en las demás disposiciones vigentes sobre la materia, se envía el extracto de la solicitud a la Oficina de comunicaciones para efecto de su publicación en la Gaceta de Propiedad Industrial, conforme lo establece el artículo 40 de la misma Decisión.

Cúmplase

Dado en Bogotá, D.C., a los **11 MAR. 2002**


ALIX CÉSPEDES DE VERGEL
Jefe de la División de Nuevas Creaciones

Bogotá, D.C., **03-11-04** El extracto de esta solicitud fue publicado en el número **515** de la Gaceta de Propiedad Industrial, de fecha **29-04-02**. El término hábil señalado por la ley expiró el **29-07-02** y SI ☒ NO ☐ se presentaron oposiciones por parte de terceros.

202040

Carrera 13 No. 27-00 Pisos 5, 7 y 10
Conmutador: 382 0840
Fax: 382 2695
E-mail: info@sic.gov.co
www.sic.gov.co
Bogotá, D.C., Colombia