

Señor Jefe

DIVISION DE NUEVAS CREACIONES

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO

Santa Fe de Bogotá, D.C.

02-01-09

Re.: Expediente No. 00 - 28.930

Solicitud de Patente

SMITHKLINE BEECHAM BIOLOGICALS s.a

SUPERINDUSTRIA Y COMERCIO
Radicacion: 00028930 00020001
Fecha (AMD): 2000-05-09 10:10:22
Trabite : 002 PATENTES D 1 REGISTRO/ 329 COMPLEMEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Folios: 18

Debidamente traducida, adjunto acompaño la solicitud de patente básica Inglesa No. 9909077.1 la cual le solicito agregar a este expediente para los efectos legales respectivos.

De Usted atentamente,

GERMAN CASTILLO GRAU

T.P.#2.978 del C.S.J.



The
Patent
Office

B 45180



INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

AmBrewer

Dated 7 April 2000

2d, 2e 2f: If there are further applicants please provide details on a separate sheet of paper

3 An address for service in the United Kingdom must be supplied

Please mark correct box.

3b: If you have appointed an agent, all correspondence concerning your application will be sent to the agent's United Kingdom address.

☐	Second applicant (if any) If you are applying as a corporate body please give: Corporate Name Country (and State of Incorporation, if appropriate)
2d	
2e	If you are applying as an individual or one of a partnership please give in full: Surname: Forenames:
2f	In all cases, please give the following details: Address: UK postcode (if applicable) Country ADP number (if known)
3	Address for service details 3a Have you appointed an agent to deal with your application? Yes ☒ No ☐ go to 3b ↓ please give details below Agent's name A W R TYRRELL Agent's address SmithKline Beecham Corporate Intellectual Property Two New Horizons Ccourt Brentford, Middlesex TW8 9EP Postcode Agent's ADP number
3b	If you have not appointed an agent please give a name and address in the United Kingdom to which all correspondence will be sent: Name Address Postcode number ADP number (if known) Daytime telephone (if available)

99
701

4 Reference number

4. Agent's or
applicant's reference FB/RH/ B45180
number (if applicable)

5 Claiming an earlier application date

5. Are you claiming that this application be treated as having been filed on the date of filing of an earlier application?

Yes ☐

No ☒ go to 6



please give details below

☐ number of earlier
application or patent
number

☐ filing date (day month year)

☐ and the Section of the Patents Act 1977 under which you are
claiming:

15(4) (Divisional) ☐ 8(3) ☐ 12(6) ☐ 37(4) ☐

6 Declaration of priority

6. If you are declaring priority from previous application(s), please give:

Country of Filing

Priority application
number
(if known)

Filing Date
(day, month, year)

Please mark correct box

Please mark correct box

6 If you are declaring priority from a
PCT Application please enter 'PCT'
as the country and enter the country
code (for example, GB) as part of
the application number

Please give the date in all number
format, for example, 31/05/90 for
31 May 1990

- any applicant is not an inventor
- there is an inventor who is not
- an applicant, or
- any applicant is a corporate body.

8 Please supply duplicates of claim(s), abstract, description and drawings).

Please mark correct box(es)

9 You or your appointed agent (see Rule 90 of the Patents Rules 1990) must sign this request.

Please sign here **á**

A completed fee sheet should preferably accompany the fee.

7 Inventorship

7. Are you (the applicant or applicants) the sole inventor or the joint inventors?

Please mark correct box

Yes ☐ No ☒ ☐ A Statement of Inventorship on Patents form 7/77 will need to be filed (see Rule 15).).

8 Checklist

8a Please fill in the number of sheets for each of the following types of document contained in this application

Continuation sheets for this Patents Form 1/77

Claim(s) 2 Description 16

Abstract Drawing(s)

8b Which of the following documents also accompanies the application?

Priority documents (please state how many)

Translation(s) of Priority documents (please state how many)

Patents Form 7/77 - Statement of Inventorship and Right to Grant

Patents Form 9/77 - Preliminary Examination Report

Patents Form 10/77 - Request for Substantive Examination

9 Request

I/We request the grant of a patent on the basis of this application.

Signed A W R Tyrrell Date: 19/04/99

A W R TYRRELL

(day month)

year)

Chartered Patent Attorney
Attorney for the Applicant

Please return the completed form, attachments and duplicates where requested, together with the prescribed fee to either;

☐ The Comptroller or
The Patent Office
Cardiff Road
Newport
Gwent
NP9 1RH

☒ The Comptroller
The Patent Office
25 Southampton Buildings
London
WC2A 1AY

100
702

Novel Compositions

This invention relates to novel vaccine formulations, methods for preparing them and their use in therapy. In particular the present invention relates to combination vaccines for administration to children and the elderly.

Streptococcus pneumoniae is a gram positive bacteria responsible for considerable morbidity and mortality, particularly in the young and aged. Expansive colonisation of the respiratory tract, and middle ear, especially in young children, is the single most common cause for hospital visits in the US. The bacteria may become invasive, infecting the lower lungs and causing pneumonia. The rate of pneumococcal pneumonia in the US for persons over 60 years of age is estimated to be 3 to 8 per 100,000. In 20% of cases this leads to bacteremia, and other manifestations such as meningitis, with a mortality rate close to 30% even with antibiotic treatment. There are 90 known serotypes of *Streptococcus pneumoniae* which are determined by the structures of the capsular polysaccharide surrounding the bacteria, and this is its major virulence factor.

A 17 - valent pneumococcal vaccine (Moniarix) is known, based on the purified polysaccharides of the pneumococcal serotypes most commonly involved in invasive disease. The method of purification of these polysaccharides was disclosed in European Patent 72513 B1. Vaccine efficacy trials with lower valent vaccines demonstrated a 70 to 90% efficacy with respect to serotypes present in the combination. Case controlled studies in the US in persons >55 years using a 14 valent vaccine demonstrated 70% efficacy (Mills, O.F., and Rhoads, G.G; J. Clin. Epidemiol. (1996); Vol.49(6) 631-636). Inclusion of additional polysaccharides (to make a 23-valent pneumococcal vaccine) were accepted on the basis of an adequate serological response, even though there was clinical efficacy data lacking (K. R. Brown In 'Combined Vaccines and Simultaneous Administration', Ed. Williams et al. New York Academy of Sciences 1995 pp 241-249).

Simultaneous immunisation with a 23-valent pneumococcal vaccine (Pnu-Immune 23 - Lederle USA) and influenza vaccine (Fluarix) has been studied (TJ Fletcher, TW Tunnicliffe, K Hammond, K Roberts, JG Ayres; (1997) Br. Med. J. 314: 1663) and no significant immunological differences were noted between simultaneous immunisation, or immunisations one month apart.

Pneumococcal polysaccharides can be rendered more immunogenic in infants by chemically coupling them to protein carriers, and clinical efficacy trials are being performed to verify this concept for efficacy in preventing infant Otitis media.

Infants primed with a 7 to 11-valent conjugate pneumococcal vaccine may be boosted with the 23-valent plain polysaccharide pneumococcal vaccine in order to increase the serotype coverage and IgG concentrations (Block, SL, JA. Hedrick, R.A. Smith, R.D. Tyler, M. Giordani, M.D. Blum, J. Sadoff, E. Keegan; Abstract G-88, 37th ICAAC, Toronto (1997); Anderson, E.L., Kennedy, D.J., Geldmacher, K.M., Donnelly, J., Mendelman, P; The Journal of Paediatrics, May 1996. Vo. 128, n°5, Part 1, 649-653).

Respiratory Syncytial virus (RSV) occurs in seasonal outbreaks, peaking during the winter in temperate climates and during the rainy season in warmer climates (DeSilva LM & Hanlon MG. Respiratory Syncytial Virus: a report of a 5-year study at a children's hospital. *J Med Virol* 1986;19:299-305). Wherever the area, RSV tends to have a regular and predictable pattern and other respiratory viral pathogens that occur in outbreaks are rarely present concurrently (Glezen WP & Denny FW. Epidemiology of acute lower respiratory disease in children. *N. Engl. J. Med.* 1973;288:498-505).

RSV infection is almost certainly underdiagnosed in adults, in part because it is considered to be an infection of children. Consequently, evidence of the virus in adults is not sought in order to explain respiratory illness. In addition, RSV is difficult to identify in nasal secretions from individuals who have some degree of partial immunity to the virus, as do the large majority of adults. Young to middle-age adults typically develop a persistent cold-like syndrome when infected with RSV. Elderly individuals may develop a prolonged

101
103

respiratory syndrome which is virtually indistinguishable from influenza, with upper respiratory symptoms which may be accompanied by lower respiratory tract involvement, including pneumonia. Institutionalised elderly populations are of particular concern, because they comprise large numbers of susceptible individuals clustered together. The spread of infection through such a population, many of whom have multiple medical problems which may predispose them to a more severe course of the disease, is difficult to control.

Furthermore, reports of recent studies evaluating the impact of RSV infection as a cause of hospitalisation in adults and in community dwelling healthy elderly further point to an important role of RSV infection in severe lower respiratory tract disease in these populations (Dowell SF, Anderson LJ, & Gary Jr HE, *et al. J. Infect Dis* 1996;174:456-462; Falsey AR, Cunningham CK, & Barker WH, *et al. J. Infect Dis* 1995;172:389-394). Dowell identified RSV as one of the four most common pathogens causing severe lower respiratory tract disease resulting in hospitalisation of adults (Dowell SF, Anderson LJ, & Gary Jr HE, *et al. J. Infect Dis* 1996;174:456-462). Falsey demonstrated that serious RSV infections in elderly persons are not limited to nursing homes or outbreak situations. Rather, RSV infection is a predictable cause of serious illness among elderly patients residing in the community. Similar to hospitalisations for influenza A, those related to RSV infections were associated with substantial morbidity, as evidenced by prolonged hospital stays, high intensive care admission rates, and high ventilatory support rates (Falsey AR, Cunningham CK, & Barker WH, *et al. J. Infect Dis* 1995;172:389-394).

These studies point to the medical and economic need for an effective vaccine which can prevent severe complications of RSV infection in infants, adults and both community dwelling healthy and institutionalised elderly.

SmithKline Beecham has undertaken the development of an RSV vaccine based on the extracellular domain of the F and G surface glycoproteins of RSV strain A. The use of an adjuvant is of primary importance for a subunit recombinant vaccine. Alum is the only adjuvant currently licensed for human use. However, Alum has a limited adjuvant effect

on the humoral response, and is known to induce mainly a TH2 cellular response (Byars NE, Nakano G, & Welch M, *et al. Vaccine* 1991;9:309-318). It has been found, however, that 3 De-O-acylated monophosphoryl lipid A (3D-MPL) in combination with Alum improves the humoral response and stimulates preferentially a TH1-type immunity. The Aluminium salt with 3D-MPL adjuvant is denoted Alum/3D-MPL.

The present invention provides a vaccine composition comprising:

- (a) one or more *Streptococcus pneumoniae* polysaccharides either conjugated to a protein or peptide, or non-conjugated; and
- (b) an RSV antigen

in combination with an adjuvant which is a preferential stimulator of TH1 cell response.

The trend towards combination vaccines has the advantage of reducing discomfort to the recipient, facilitating scheduling, and ensuring completion of regiment; but there is also the concomitant risk of reducing the vaccine's efficacy. It would be, therefore, advantageous to make vaccine combinations which meet the needs of a population, and which, in addition, do not interfere with each other. It would be of further advantage if the combination of the vaccines results in synergy with resulting improvement of one or both vaccines efficacy, or improved correlates of protections for one or both vaccines. This is achieved by the vaccine composition of the invention which is of great benefit for administration to children or the elderly who may be particularly at risk of *Streptococcus pneumoniae*, and/or RSV infection.

Optionally the vaccine composition of the invention additionally comprises one or more of a number of other antigens such as an antigen against *influenza* virus. Currently available influenza vaccines include whole inactivated virus vaccines, split particle vaccines, and subunit vaccines. Inactivated influenza vaccines, of all kinds, are usually trivalent vaccines. They contain antigens derived from two influenza A virus strains and one

102
404

influenza B strain. A standard 0.5ml dose of most of them contain 15µg of haemagglutinin component from each strain.

The procedure for determining the strains to be incorporated into an influenza vaccine is a complex process which involves collaboration between the World Health Organisation, national health authorities and vaccine manufacturers.

There are two conjugation methods generally used for producing immunogenic polysaccharide constructs: (1) direct conjugation of carbohydrate and protein; and (2) indirect conjugation of carbohydrates and protein via a bifunctional linker or spacer reagent. Generally, both direct and indirect conjugation require chemical activation of the carbohydrate moiety prior to derivatisation. See for example US 5,651,971 and Dick & Beurret, "Glycoconjugates of Bacterial Carbohydrate Antigens," Conjugate Vaccines, J.M. Cruse & R.E. Lewis (eds), Vol. 10, 48 - 114 (1989).

The *Streptococcus pneumoniae* polysaccharides, if conjugated, are conjugated to either a protein or a peptide. Polysaccharide antigens have been conjugated to a number of T helper proteins. These provide T-helper epitopes. Representative proteins include Diphtheria Toxoid, Tetanus toxoid, and protein D or its lipidated derivative lipoprotein D from *Haemophilus influenzae* B. Other suitable protein carriers include, Diphtheria Crm 197 and the major non structural protein from *influenzae*, NS1 (particularly amino acid 1-81).

The *Streptococcus pneumoniae* polysaccharide in the composition of the invention is preferably a vaccine with a number of valencies, preferably at least an 11-valent vaccine, for example a 17- or 23-valent vaccine. Most preferably the 23-valent pneumococcal polysaccharide or the 11-valent conjugate vaccine.

Suitable RSV antigens for inclusion in vaccines include an inactivated RSV virus, such as a formalin inactivated RSV virus, or antigens derived from the RSV virus, preferably human RSV envelope glycoproteins, such as the RSV F or G protein or immunogenic fragments

thereof as disclosed for example in US Patent 5149650, or a chimeric polypeptide comprising at least one immunogenic fragment from both RSV F and G proteins, advantageously an RSV FG chimeric protein as disclosed for example in US patent 5194 595 preferably expressed from CHO cells.

The RSV antigen in the composition of the invention is preferably an RSV FG chimeric protein.

Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in International Patent Application No. 92/116556.

Preferably, the size of the particles of 3D-MPL are small enough to be sterile filtered through a 0.22micron membrane (as described in European Patent number 0689454).

3D-MPL will be present in the range of 10µg - 100µg preferably 25-50µg per dose wherein the antigen will typically be present in a range 2-50µg per dose.

Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of Quillaja Saponaria Molina. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with an carrier.

QS21 is a Hplc purified non-toxic fraction of a saponin from the bark of the South American tree Quillaja Saponaria Molina and its method of its production is disclosed (as QS21) in US patent No. 5,057,540.

103
105

Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen. Thus vaccine compositions which form part of the present invention may include a combination of QS21 and cholesterol.

Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; preferably 1:5 to 5 : 1 and often substantially 1 : 1. The preferred range for optimal synergy is 2.5 : 1 to 1 : 1 3D MPL: QS21.

Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or alum.

A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and tween 80. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

In a preferred aspect aluminium hydroxide (alum) or aluminium phosphate will be added to the composition of the invention to enhance immunogenicity.

In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with 3D-MPL and alum.

EXAMPLES

1. Formulation process:

H₂O-diluted FG antigen (2µg:1/10 HD) was adsorbed for 15 min on 50 µg of Al(OH)₃. When used, 5µg of 3D-MPL were added to the preparation as a suspension of 100 nm particles and incubated for 30 min. Formulations were then buffered with 10-fold concentrated PBS pH 7.4. When used, the 23 valent pneumococcal vaccine (11.5µg:1/50 HD) was added 15 min. after the addition of the buffer solution. For groups without FG the same formulation sequence was followed so that the 3D-MPL was first adsorbed on Al(OH)₃ before adding the 23 valent pneumococcal vaccine. Fifteen minutes after the addition of the concentrated buffer or the 23 valent pneumococcal vaccine phenoxxyethanol (5mg/ml) was added to the formulations as preservative.

All incubations were carried out at room temperature with agitation. The formulations were prepared simultaneously for the 2 injections with a 7-day maturation of the finalised formulations before the first injection.

Composition of formulation constituents:

COMPONENT	BATCH NUMBER	CONCENTRATION (µg/ml)	BUFFER
FG	54/023	265	10 PO ₄ , 150NaCl pH6.8
Al (OH) ₃	96A0089	10380	H ₂ O
23-Valent	Pneumune	1150	saline
3D-MPL	109	964	H ₂ O

2. Immunization protocol:

11 groups of 10 mice were immunised by different routes (50 µl) at days 0 and 28 with various formulations (see *Table I*). Group 7 and 8 were immunised with live RSV by the intra-nasal route (60µl). Sera were obtained at days 28 (28 d Post I) and 42 (14 d Post II). On day 42, spleen and node cells were taken from 5 mice of groups 4, 5, 6, 7, 8, 9, as well as from 5 naïve Balb/c mice (group 1, not immunised).

105
107

3. Humoral response.

3.1. Anti-FG antibodies:

All humoral results were performed for 10 mice/group (individual response for the anti-FG titers and pooled sera for the isotype profile) and cellular results were presented for 5 mice/group.

Individual sera were obtained 28 days after the first Immunization and 14 days after the second immunisation and were tested for the presence of FG specific Ig antibodies and their isotype (IgG2a, IgG1) distribution.

The assay protocol was as follows: coating overnight at 4°C with 50 µl of purified FG 54/023 (1 µg/ml) per well, saturation 1h at 37°C, incubation with sera 1h30 at 37°C, incubation with anti-mouse Ig biotin 1/1500 (or IgG1, IgG2a biotin 1/1000) 1h30 at 37°C, incubation with strepta-peroxydase 1/2500 30 min at 37°C, incubation with OPDA Sigma 15 min at RT, stop with H₂SO₄ 2N.

OD were monitored at 490 nm and the titers determined by linear regression ($y=a.\log x + b$): titer = serum dilution giving 50% reduction of the maximal OD. 95% confidence limits (IC 95%) were calculated for each group.

Individual sera obtained 14d Post II were tested for the presence of neutralising antibodies using the following protocol: 50 µl of serial two-fold dilutions of sera (first dilution 1/250) were incubated for 1 hour at 37°C with 50 µl of a mixture containing 500 pfu of RSV-A/Long (Lot 14) and guinea pig complement in a 96 well plate in duplicate. 100 µl of a HEp-2 cell suspension at 10⁵ cells/ml were then added to each well and the plates were incubated for 4 days at 37°C in the presence of 5% CO₂.

The supernatants were then aspirated, and after addition of a 100 µl of a WST-1 preparation (dilution 1/12.5) the plates were further incubated for 24H at 37°C in the presence of 5% CO₂. The OD were monitored at 595 nm and the titers determined by linear regression ($y=a.\log x + b$): titer = serum dilution giving 50% reduction of the maximal OD observed for the uninfected cells.

Controls in test included a pool of randomly chosen human sera (Human pool) and Sandoglobuline (lot 069, generic human IgG produced by Sandoz).

3.2. Anti-Pneumococcal Polysaccharide IgG:

Murine IgG to pneumococcal polysaccharides types 6B, 14, 19F and 23F was measured by ELISA in a method adapted from the CDC protocol. This protocol includes the addition of soluble cell wall polysaccharide (CPS) to the sera to inhibit the measurement of CPS antibodies. CPS is a phosphoryl choline containing teichoic acid common to all pneumococci. It is present under the capsule, and antibodies to it are only weakly

protective. Since CPS is linked to the capsular polysaccharide, there is usually 0.5 to 1% CPS contaminating the purified capsular polysaccharide used to coat the ELISA plates. Thus, measurement of the CPS antibodies can confound the interpretation ELISA results with respect to the capsular polysaccharide.

The ELISA was performed with polysaccharides coated at 20, 5, 10 and 20 µg/ml in carbonate buffer for types 6B, 14, 19F and 23F respectively. Sera was pre-mixed with the equivalent of 500 µg/ml CPS in undiluted sera, and incubated for 30 minutes before addition to the ELISA plate. Murine IgG was detected with Jackson ImmunoLab goat anti-murine IgG (H+L) peroxidase at 1:2000 dilution. The titration curves were referenced to polysaccharide specific murine monoclonal antibodies of known concentration for each serotype using logistic log comparison by SoftMax Pro. The monoclonals used were HASP4, PS14/4, PS19/5 and PS23/22 for types 6B, 14, 19F and 23F respectively. Due to the limited quantity of sera available, pooled sera was tested, thus statistical analysis is not available.

4. Cellular response

Spleen and lymph node cells were isolated 14d Post II from groups 4-9 and from naïve mice (Group 1) for use as a negative control for the FG-specific cellular response analysis. Samples were analysed for both FG-specific lymphoproliferation and cytokine (IFN-γ + IL-5) secretion.

Proliferation was evaluated after a 96h incubation of 4×10^5 cells/well of 96 well plates with 200 µl of media containing 10 to 0.03 µg/ml of FG (3-fold dilutions). Upon ^3H -thymidine incorporation, the FG specific proliferation was measured following our standard protocol. Cytokine induction was evaluated after 96 h incubation of 2.5×10^6 cells per well of 24 well with 1 ml of media containing 10µg to 0.01µg of FG (10-fold dilutions). Supernatants were then harvested to determine the quantity of IFN-γ and IL-5 induced by ELISA following our standard protocol.

Results

1. GROUPS.

10 groups received two immunisations of various formulations containing either the 23-valent vaccine or FG or a combination of both formulated with Aluminium hydroxide and 3D-MPL or Aluminium hydroxide. Group 1 constitutes the control for the CMI studies. Groups 2 and 9 will allow the evaluation of the immunogenicity of the 23 Valent pneumococcal vaccine upon intraperitoneal immunization which was used in the literature for the evaluation of 23 Valent pneumococcal vaccine, and upon IM immunization, Groups 3 and 10 allow a parallel analysis to groups 2 and 9 except that the 23 Valent pneumococcal vaccine is formulated with Alum/3D-MPL. Groups 2 and 3 also constitute

106
108 X

controls for the impact of Alum/3D-MPL on the 23 Valent pneumococcal vaccine when it is not combined to FG Alum/3D-MPL. Group 4 will allow the evaluation of the immune response induced upon IM Immunization of the combination of the 23 Valent pneumococcal vaccine and FG Alum/3D-MPL. Groups 5 and 6 constitute a control for the evaluation of the impact of Alum/3D-MPL on FG when it is not combined with the 23 Valent pneumococcal vaccine. The RSV live immunisations was a control for the immune response induced upon natural RSV IN infection (Groups 7). Finally, group 9 is a control for the impact of Alum/3D-MPL alone.

TABLE 1

Groups	Antigen	Adjuvant	Route
1	none	none	
2	23 Valent Pneumune(11.5µg)	none	IM
3	23 Valent Pneumune (11.5µg)	Alum/3D-MPL	IM
4	23 Valent Pneumune (11.5µg) / FG (2µg)	Alum/3D-MPL	IM
5	FG (2µg)	Alum/3D-MPL	IM
6	FG (2µg)	Alum	IM
7	RSV Live (Lot 14: 10 ⁵ PFU)	none	IN
8	none	Alum/3D-MPL	IM
9	23 Valent Pneumune (11.5µg)	none	IP
10	23 Valent Pneumune (11.5µg)	Alum/3D-MPL	IP

2. HUMORAL RESPONSE.

2.1. Anti-FG antibodies

- The analysis of specific anti-FG antibodies at 28 d Post I and 14 d Post II shows the induction of similar antibody responses upon Immunization with FG Alum/3D-MPL / 23-Valent (Group 4) or FG Alum/3D-MPL alone (Group 5) with slightly higher titers observed for FG Alum/3D-MPL / 23-Valent (*Figure 1*). The Immunization with FG Alum (Group 6) induces expectedly lower antibody titers at both time points. Statistical analysis shows that there is indeed a significant difference between the anti-FG Ig titers induced by Groups 4 -6 at 28 d Post I and 14 d Post II.

RESULTS

- The analysis of anti-RSVA neutralizing antibodies (*Figure 2*) presents a parallel profile as the one observed for the anti-FG Ig responses.
- Based on the above results, the anti-FG / anti RSV neutralizing antibody ratio (28d PostII) was calculated and showed that the ratio of FG Alum/3D-MPL / 23-Valent, FG Alum/3D-MPL and FG Alum compares to the ratio induced by the RSV IN group (*Figure 3*).
- An in depth statistical analysis on individual anti-FG Ig, IgG1 and IgG2a isotype titers showed that the addition of the 23-valent vaccine to FG Alum/3D-MPL maintains the Ig and IgG1 responses and increases significantly the IgG2a responses (*Figure 4A*). In addition, the analysis showed that the IgG1/IgG2a ratio and the IgG1 titer of FG Alum/3D-MPL / 23 valent and FG Alum/3D-MPL were similar although significantly different from FG Alum (*Figure 4B*). The three formulations had significant differences in their IgG2a titers.

2.2. Anti-Pneumococcal Polysaccharide IgG

While mice and other rodents may produce IgM against polysaccharide immunogens, they do not normally produce IgG against polysaccharides. This is because their immune system lacks the signals to induce isotype switching to a T-independent antigen such as a polysaccharide. Thus, the measurement of anti-polysaccharide IgG in mice is a sensitive test for the induction of an improved T-independent immune response.

The concentrations of IgG induced against four of the 23 polysaccharides are presented in *Figure 5*. As we had noted in other experiments, no IgG was produced against polysaccharide types 6B and 23F. However, there was measurable IgG produced against types 14 and 19F.

- Groups that did not contain 23-Valent vaccine did show any detectable IgG to polysaccharides 14 and 19F, confirming the specificity of the measurements.
- A comparison of the route of immunisation reveals that when the 23-valent vaccine is adjuvanted with Alum/3D-MPL, the intra-muscular route appears to be better than inter-peritoneal for type 19F, whereas the reverse is true for type 14. With plain 23-Valent, there does not seem to be a significant difference.
- 23Valent + Alum/3D-MPL induces greater IgG for types 14 and 19F, in both IM and IP immunisation routes when compared to 23-Valent alone. When combined with FG,

107
409

however, there is a reduction in the IgG response. Nevertheless, the IgG response to 19F in 23-Valent+FG/Alum/3D-MPL is still greater than that induced by 23-Valent alone (compare groups 3 and 4), and the response to type 14 is improved.

3. CELLULAR RESPONSE.

- The induced FG-specific lymphoproliferation did not show any difference between FG Alum/3D-MPL +/- 23 Valent and FG Alum, both in spleen cells and in lymph node cells (*Figure 6*).
- The analysis of the production of IL-5 and IFN- γ suggests that the mixture of the 23 valent vaccine with FG Alum/3D-MPL does not hamper the production of IFN- γ but could increase somewhat the production of IL-5 (*Figure 7*). This observation is confirmed by a 2 to 8 fold difference in the IFN- γ / IL-5 ratio observed at two doses of FG (10 and 1 μ g FG/ml) used for *in vitro* restimulation. However, the FG Alum/3D-MPL / 23-valent vaccine still induces a much higher IFN- γ / IL-5 ratio than FG Alum.

CONCLUSIONS

The analysis of the induced anti-FG Ig specific and anti RSVA neutralizing antibody titers shows that the combination of FG Alum/3D-MPL with the 23 valent pneumococcal vaccine does not hamper the induction of the response observed with FG Alum/3D-MPL alone. The quality of the response measured by the anti-FG/anti-RSVA neutralizing antibody ratio also remains unchanged and close to the ratio observed upon natural infection with RSV true the IN route.

Analysis of the induced FG specific cell mediated response suggest that the addition of the 23 valent vaccine to FG Alum/3D-MPL does not affect the induction of lymphoproliferation in both spleen cells and lymph node cells. Furthermore, the induction of a Th1 type response typically observed with FG Alum/3D-MPL is not hampered by the addition of the 23 valent as measured by *in vitro* cytokine production of FG specific spleen and lymphnode cells and appears to be enhanced when analysing the FG specific isotype antibody distribution. Indeed, the production of IFN- γ , marker of a Th1 response, remains unchanged upon addition of the 23 valent vaccine to FG Alum/3D-MPL while the production of IL-5, marker of a Th2 response is only slightly increased. Interestingly, the addition of the 23 valent vaccine to FG Alum/3D-MPL significantly increases the production of IgG2a antibodies, marker of a Th1 response, while it is not affecting the IgG1 (marker of a Th2 response) nor the IgG1/IgG2a response.

Thus the combination of FG Alum/3D-MPL and 23-valent vaccine does not affect the response observed with FG Alum/3D-MPL and therefore the advantages linked to the FG Alum/3D-MPL formulation versus FG Alum i.e., induction of high primary and secondary neutralizing antibody responses and a Th1 response as measured by the presence of IgG2a antibodies and the induction by high levels of IFN- γ are maintained.

Combination of 23Valent Pneumune with FG Alum/3D-MPL results in increased IgG production in 2 of 4 polysaccharides tested, but most dramatically for type 19F. 23-Valent alone with Alum/3D-MPL gave the highest IgG concentrations.

In conclusion, there is a favourable combination of 23-Valent pneumococcal vaccine with RSV FG Alum/3D-MPL in which there is a synergistic effect for both vaccines in that both vaccines show improvements in certain immunological tests, and no inhibition in any tests.

103
710

Claims

1. A vaccine composition comprising:
 - (a) one or more *Streptococcus pneumoniae* polysaccharides either conjugated to a protein or peptide, or non-conjugated; and
 - (b) an RSV antigenin conjunction with an adjuvant which is a preferential stimulator of TH1 cell response.
2. A vaccine composition according to claim 1 in which the *Streptococcus pneumoniae* polysaccharide is conjugated preferentially to a protein from the group comprising; Protein D from Haemophilus influenzae B, a lipidated version thereof (lipoprotein D); Tetanus toxin, and Diphtheria Toxin.
3. A vaccine composition according to claim 1 or claim 2 which additionally comprises a carrier.
4. A vaccine composition according to any one of claims 1 to 3 in which the preferential stimulator of TH1-cell response is selected from the group of adjuvants comprising: 3D-MPL, 3D-MPL wherein the size of the particles of 3D-MPL is preferably about or less than 100nm, QS21, a mixture of QS21 and cholesterol, and a CpG oligonucleotide.
5. A vaccine composition according to claim 4 in which the preferential stimulator of TH1-cell response is 3D-MPL.
6. A vaccine composition according to any one of claims 1 to 5 in which the *Streptococcus pneumoniae* polysaccharide is a 23-valent pneumococcal polysaccharide vaccine.

7. A vaccine composition according to any one of claims 1 to 5 in which the *Streptococcus pneumoniae* polysaccharide is an 11-valent conjugated pneumococcal polysaccharide vaccine.
8. A vaccine composition according to any one of claims 1 to 7 in which the RSV antigen is a human RSV envelope glycoprotein.
9. A vaccine composition according to any one of claims 1 to 8 in which the RSV antigen is chimeric FG or a fragment thereof.
10. A vaccine composition according to any one of claims 1 to 9 in which an influenza virus antigen is additionally present.
11. A vaccine composition according to any one of claims 1 to 10 in which the carrier is selected from the group comprising aluminium hydroxide, aluminium phosphate and tocopherol and an oil in water emulsion.

109
912

LEGALIZACIONES

OFICINA DE PATENTES

Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

El suscrito, un funcionario debidamente autorizado conforme a los Artículos 74(1) y (4) de la Ley de Desreglamentación y Contratación de 1994 para firmar y expedir certificados en nombre del Contralor General, certifico que la copia adjunta es una copia correcta de los documentos archivados originalmente con respecto a la solicitud de patente que allí se identifica.

De acuerdo con las Reglamentaciones de Patentes de 1982 (Re-registro de Compañías), si una compañía nombrada en este certificado y en los documentos adjuntos se ha vuelto a registrar según la Ley de Compañías de 1980 con el mismo nombre con el que fue registrada inmediatamente antes del re-registro, salvo la sustitución o inclusión como la última parte del nombre de las palabras "sociedad anónima" o su equivalente en galés, las referencias al nombre de la compañía y sus documentos adjuntos serán tratadas como referencias al nombre con el cual fue re-registrada.

De acuerdo con las reglas, las palabras "sociedad anónima" (o en inglés *public limited company*) pueden ser reemplazadas por las letras p.l.c, plc. P.L.C ó PLC.

El re-registro bajo la Ley de Compañías no constituye una nueva entidad legal sino simplemente somete a la compañía a ciertas normas de ley adicionales.

(FIRMA) ANDREW BREWER

FECHA: 7 de abril de 2000

(SELLO DE LACRE)

110
713 5

OFICINA DE PATENTES

Referencia No. **RB/RH/B45180**

No. **9909077.1** 20 de abril de 1999

OFICINA DE PATENTES: Solicitud de adjudicación de una Patente.

Forma 1/77

Ley de Patentes de 1977

1. TITULO DEL INVENTO: **"COMPOSICIONES NOVEDOSAS"**

2. INFORMACIÓN DEL SOLICITANTE:

2a. Nombre: **SmithKline Beecham Biologicals S.A.**

Pais: **Bélgica**

2b N/A

2c Dirección: **Rue de l'Institut 89
B-1330 Rixensart
Bélgica**

Código Postal:

PAIS. **BELGICA**

ADP No. **05781117001**

2d **Segundo Solicitante:**

N/A

3. Ha nombrado Ud. a alguien que le represente en su solicitud? SI

Nombre del Agente: **A M R TYRRELL**

SmithKline Beecham

Corporate Intellectual Property

Two New Horizons Court,

Brentford, Middlesex

Código postal **TW8 9EP**

No. ADP

4. Número de Referencia: **FB/RH/B45180**

5. Solicita Ud. que esta solicitud sea tratada como si hubiera sido radicada en la fecha de presentación de una solicitud anterior: NO

6. No aplica

7. Es usted el único inventor o inventor conjunto? NO

8. Cuántas hojas describen esta solicitud? 16 - Reivindicaciones: 2-

Resumen: - - Dibujos - 4 -

9. Solicitud: Solicitamos la adjudicación de una patente teniendo en cuenta esta solicitud.

(FIRMA) A W R TYRRELL

Fecha: 19/04/99

Abogado de Patentes

Favor devolver la forma diligenciada y sus anexos y duplicados, junto con su valor

a:

The Comptroller ó
The Patent Office
Cardiff Road
Newport
Gwent
NP9 1RH

The Comptroller
The Patent Office
25 Southampton Buildings
London
WC2A 1AY

112
115

COMPOSICIONES NOVEDOSAS

REIVINDICACIONES:

1. Una composición de vacuna que comprende:
 - a) uno o más polisacáridos *Streptococcus pneumoniae* conjugados a una proteína o a un péptido, o no conjugados; y
 - b) un antígeno RSV,junto con un adyuvante que es un estimulador preferencial de respuesta de la célula TH1.
2. Una composición de vacuna según la reivindicación 1, en la cual el polisacárido *Streptococcus pneumoniae* es conjugado preferencialmente a una proteína del grupo compuesto por: Proteína D de Haemophilus influenzae B, una versión lipitada de la misma (lipoproteína D); toxina de Tétano, y Toxina de Difteria.
3. Una composición de vacuna según la reivindicación 1 o 2, que adicionalmente comprende un vehículo.
4. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 3, en la cual el estimulador preferencial de la respuesta de la célula TH1 es seleccionado del grupo de adyuvantes compuestos por: 3D-MPL, 3D-MPL, en donde el tamaño de partículas de 3D-MPL es preferiblemente alrededor de o menos de 100 nm, QS21, una mezcla de QS21 y colesterol, y un oligonucleótido CpG.

113
116

5. Una composición de vacuna según la reivindicación 4, en la cual el estimulador preferencial de la respuesta de célula TH1 es 3D-MPL.
6. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 5, en la cual el polisacárido *Streptococcus pneumoniae* es una vacuna polisacárida neumococo de 23 valentes.
7. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 5, en la cual el polisacárido *Streptococcus pneumoniae* es una vacuna polisacárida neumococo de 11 valentes.
8. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 7, en la cual el antígeno TSV es una glicoproteína humana RSV envolvente.
9. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 8, en la cual el antígeno RSV es un FG quimérico o un fragmento del mismo.
10. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 9, en la cual un antígeno del virus de influenza está presente adicionalmente.
11. Una composición de vacuna según cualquiera de las reivindicaciones 1 a 10, en la cual el vehículo es seleccionado del grupo compuesto por hidróxido de aluminio, fosfato de aluminio y tocoferol y una emulsión aceite en agua.