

CAVELIER

ABOGADOS

WEB SITE: www.cavelier-abogados.com
E-MAIL: caveller@colomsat.net.co

EDIFICIO SISKI
CARRERA 4 No. 72 - 35
BOGOTÁ 8, COLOMBIA

SUPERINDUSTRIA Y COMERCIO

Radicación: 00065554 00000001 Folios: 143
Fecha (AM): 2000-11-17 16:57:43
Trámite: 002 PATENTES D I REGISTRO/ 329 COMPLEMEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Señora Jefe de la
DIVISION DE NUEVAS CREACIONES
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
E. S. D.

Referencia : Expediente No. 00.065.554

Solicitante : NEUROGEN CORPORATION

Asunto : Solicitud de patente para:
"PIRROLCARBOXAMIDAS FUSIONADAS, UNA NUEVA
CLASE DE LIGANDOS DE RECEPTORES CEREBRALES
DE GABA"

Fecha de solicitud : Agosto 31 de 2000

TRAMITE 002 EVENTO 001 ACTUACION 329

1. Yo, Emilio FERRERO, obrando como apoderado de la solicitante en el asunto de la referencia, atentamente, presento con este escrito la copia certificada de la solicitud de patente Estadounidense No. 09/387,311 del 31 de agosto de 1999, junto con la respectiva traducción de su carátula y reivindicaciones de la cual se reclama prioridad en esta solicitud de patente.

2. Ruego a la Señora Jefe se sirva aceptar este escrito con sus anexos, y se continúe con el trámite de la solicitud.

Bogotá, 17 NOV. 2000
Señora Jefe,



EMILIO FERRERO
T.P.A. No. 31.929

COLO 10611-801-005-2
ASN-ID-41

NIT. 860.041.367-3



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

September 27, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 09/387,311

FILING DATE: August 31, 1999



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

L. Edelen

L. EDELEN
Certifying Officer

14A
A 182

08/31/99

1000 U.S. PTO

UTILITY PATENT APPLICATION TRANSMITTAL

Submit an original and a duplicate for fee processing
(Only for new nonprovisional applications under 37 CFR 1.53(b))

ADDRESS TO:

Assistant Commissioner for Patents
Box Patent Application
Washington, D.C. 20231

Attorney Docket No. 99,894
First Named Inventor Pamela Albaugh
Express Mail No. EL007910125US
Total Pages 105

1000 U.S. PTO

09/30/99

1000 U.S. PTO

08/31/99

APPLICATION ELEMENTS

1. ☒ Transmittal Form with Fee
2. ☒ Specification (including claims and abstract) [Total Pages 100]
3. ☐ Drawings [Total Sheets]
4. ☐ Oath or Declaration [Total Pages]
 - a. ☐ Newly executed
 - b. ☐ Copy from prior application [Note Boxes 5 and 17 below]
 - i. ☐ Deletion of inventor(s) Signed statement attached deleting inventor(s) named in the prior application
5. ☐ Incorporation by Reference: The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Microfiche Computer Program
7. ☐ Nucleotide and/or Amino Acid Sequence Submission
 - a. ☐ Computer Readable Copy
 - b. ☐ Paper Copy
 - c. ☐ Statement verifying above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers
9. ☐ Power of Attorney
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)
 - ☐ PTO-1449 Form
 - ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Return Receipt Postcard (Should be specifically itemized)
14. ☐ Small Entity Statement(s)
 - ☐ Enclosed
 - ☐ Statement filed in prior application; status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
16. ☒ Other: Certificate of Mailing by "Express Mail."

17. ☐ This is a CONTINUING APPLICATION. Please note the following:

- a. ☐ This is a ☐ Continuation ☐ Divisional ☐ Continuation-in-part of prior application Serial No. _____.
- b. ☐ Cancel in this application original claims ____ of the prior application before calculating the filing fee.
- c. ☐ Amend the specification by inserting before the first line the sentence:
This is a ☐ continuation ☐ divisional ☐ continuation-in-part of application Serial No. _____, filed _____.
- d. ☐ The prior application is assigned of record to _____.

09387311-083499

145
183

UTILITY PATENT APPLICATION TRANSMITTAL				Attorney Docket No. 99,604	
APPLICATION FEES					
BASIC FEE				\$ 780.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	42 -20=	22	x \$18.00	\$ 396.00	
Independent Claims	12 -3=	9	x \$78.00	\$ 702.00	
☐ Multiple Dependent Claims(s) if applicable				+ \$270.00	
				\$	
Total of above calculations =				\$1,858.00	
Reduction by 50% for filing by small entity =				\$()	
☐ Assignment fee if applicable				+ \$40.00	
				\$	
TOTAL =				\$1,858.00	

18. ☒ Please charge my Deposit Account No. 13-2490 in the amount of \$1,858.00.

19. ☐ A check in the amount of \$_____ is enclosed.


20. The Commissioner is hereby authorized to credit overpayments or charge any additional fees of the following types to Deposit Account No. 13-2490:

a. ☒ Fees required under 37 CFR 1.16.

b. ☒ Fees required under 37 CFR 1.17.

c. ☒ Fees required under 37 CFR 1.18.

21. ☒ The Commissioner is hereby generally authorized under 37 CFR 1.136(a)(3) to treat any future reply in this or any related application filed pursuant to 37 CFR 1.53 requiring an extension of time as incorporating a request therefor, and the Commissioner is hereby specifically authorized to charge Deposit Account No. 13-2490 for any fee that may be due in connection with such a request for an extension of time.

22. CORRESPONDENCE ADDRESS	
Name	McDonnell Boehnen Hulbert & Berghoff
Address	32 nd Floor, 300 South Wacker Drive
City, State, Zip	Chicago, Illinois 60606
23. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED	
Name	Steven J. Sarussi Reg. No. 32,784
Signature	
Date	August 31, 1999

CERTIFICATE OF MAILING BY "EXPRESS MAIL" UNDER 37 CFR § 1.16: The undersigned hereby certifies that this Transmittal Letter and the paper, as described in paragraph 1 hereinabove, are being deposited with the United States Postal Service with sufficient postage as "Express Mail Post Office to Addressee" in an envelope addressed to: Asst. Commissioner for Patents, Box Patent Application, Washington, D.C. 20231, on the 31st day of August, 1999. Express Mail No. EL007910139US.

BY: 

09387311-0831

196
134

**CERTIFICATE OF MAILING BY "EXPRESS MAIL"
(NEW PATENT APPLICATION)**

(Attorney Docket No.: 99,694)

Express Mail No. EL007910125US

Deposited: August 31, 1999

I hereby certify that the attached correspondence, identified below, is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" under 37 CFR § 1.10 on the date indicated above and is addressed to the Asst. Commissioner for Patents, Box Patent Application, Washington, DC 20231.

By: Christopher Jaffee

Patent Application of: Pamela Albaugh, Gang Lin, Kenneth Shaw and Alan Hutchison

Title: Novel Fused Pyrrolocarboxamides; A New Class of GABA Brain Receptor Ligands

- ☒ Utility Patent Application Transmittal
- ☒ Patent Application (100 pages, including cover sheet, claims, and abstract)
- ☐ Check (\$_____)
- ☒ Postcard
- ☐ Declaration and Power of Attorney (a true copy)
- ☐ Preliminary Amendment
- ☐ Drawings (sheets)
- ☐ Information Disclosure Statement
- ☐ Small Entity Form
- ☐ Other:

09387311-0831 9

199
185

APPLICATION FOR UNITED STATES LETTERS PATENT
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

5

(MBHB Case No. 99,694)

Title: FUSED PYRROLECARBOXAMIDES; A NEW CLASS OF
GABA BRAIN RECEPTOR LIGANDS

10

Inventors:

Pamela Albaugh
81 Long Hill Road
Clinton, CT 06413

15

Gang Liu
491 Pesaro Street
Agoura, CA 91301

20

Kenneth Shaw
83 Sheephill Road
Weston, CT 06883

25

Alan Hutchison
29 Kimberly Lane
Madison, CT 06443

30

Assignee: Neurogen Corporation
Branford, Connecticut
A Corporation of the State of Delaware

09387311-0
FE D - FFEZBEGD
FE

198
186

**FUSED PYRROLECARBOXAMIDES; A NEW CLASS OF
GABA BRAIN RECEPTOR LIGANDS**

BACKGROUND OF THE INVENTION

5

Field of the Invention

This invention relates to novel fused pyrrolecarboxamides which selectively bind to GABA_A receptors. This invention also relates to pharmaceutical compositions comprising such compounds. It further relates to the use of such compounds in treating anxiety, sleep and seizure disorders, and overdoses of benzodiazepine-type drugs, and enhancing alertness.

10

Description of the Related Art

15

γ -Aminobutyric acid (GABA) is regarded as one of the major inhibitory amino acid transmitters in the mammalian brain. Over 30 years have elapsed since its presence in the brain was demonstrated (Roberts & Frankel, J. Biol. Chem 187: 55-63, 1950; Udenfriend, J. Biol. Chem. 187: 65-69, 1950). Since that time, an enormous amount of effort has been devoted to implicating GABA in the etiology of seizure disorders, sleep, anxiety and cognition (Tallman and Gallager, Ann. Rev. Neuroscience 8: 21-44, 1985). Widely, although unequally, distributed through the mammalian brain, GABA is said to be a transmitter at approximately 30% of the synapses in the brain. In most regions of the brain, GABA is associated with local inhibitory neurons and only in two regions is GABA associated with longer projections. GABA mediates many of its actions through a complex of proteins localized both on cell bodies and nerve endings; these are called GABA_A receptors. Postsynaptic responses to GABA are mediated through alterations in chloride conductance that generally, although not invariably, lead to hyperpolarization of the cell. Recent investigations have indicated that the complex of proteins associated with postsynaptic GABA responses is a major site of action for a number of structurally unrelated compounds capable of modifying postsynaptic responses to GABA.

20

25

0 387311-083199
661E80-1FE78E

Depending on the mode of interaction, these compounds are capable of producing a spectrum of activities (either sedative, anxiolytic, and anticonvulsant, or wakefulness, seizures, and anxiety).

1,4-Benzodiazepines continue to be among the most widely used drugs in the world. Principal among the benzodiazepines marketed are chlordiazepoxide, diazepam, flurazepam, and triazolam. These compounds are widely used as anxiolytics, sedative-hypnotics, muscle relaxants, and anticonvulsants. A number of these compounds are extremely potent drugs; such potency indicates a site of action with a high affinity and specificity for individual receptors. Early electrophysiological studies indicated that a major action of benzodiazepines was enhancement of GABAergic inhibition. The benzodiazepines were capable of enhancing presynaptic inhibition of a monosynaptic ventral root reflex, a GABA-mediated event (Schmidt et al., 1967, Arch. Exp. Path. Pharmacol. 258: 69-82). All subsequent electrophysiological studies (reviewed in Tallman et al. 1980, Science 207: 274-81, Haefley et al., 1981, Handb. Exptl. Pharmacol. 33: 95-102) have generally confirmed this finding, and by the mid-1970s, there was a general consensus among electrophysiologists that the benzodiazepines could enhance the actions of GABA.

With the discovery of the "receptor" for the benzodiazepines and the subsequent definition of the nature of the interaction between GABA and the benzodiazepines, it appears that the behaviorally important interactions of the benzodiazepines with different neurotransmitter systems are due in a large part to the enhanced ability of GABA itself to modify these systems. Each modified system, in turn, may be associated with the expression of a behavior.

Studies on the mechanistic nature of these interactions depended on the demonstration of a high-affinity benzodiazepine binding site (receptor). Such a receptor is present in the CNS of all vertebrates phylogenetically newer than the bony fishes (Squires & Braestrup 1977, Nature 166: 732-34, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). By using tritiated diazepam, and a variety of other compounds, it has

200
188

been demonstrated that these benzodiazepine binding sites fulfill many of the criteria of pharmacological receptors; binding to these sites *in vitro* is rapid, reversible, stereospecific, and saturable. More importantly, highly significant correlations have been shown between the ability of benzodiazepines to displace diazepam from its binding site and activity in a number of animal behavioral tests predictive of benzodiazepine potency (Bracstrup & Squires 1978, Br. J. Psychiatry 133: 249-60, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). The average therapeutic doses of these drugs in man also correlate with receptor potency (Tallman et al. 1980, Science 207: 274-281).

10 In 1978, it became clear that GABA and related analogs could interact at the low affinity (1 mM) GABA binding site to enhance the binding of benzodiazepines to the clonazepam-sensitive site (Tallman et al. 1978, Nature, 274: 383-85). This enhancement was caused by an increase in the affinity of the benzodiazepine binding site due to occupancy of the GABA site. The data were interpreted to mean that both GABA and benzodiazepine sites were allosterically linked in the membrane as part of a complex of proteins. For a number of GABA analogs, the ability to enhance diazepam binding by 50% of maximum and the ability to inhibit the binding of GABA to brain membranes by 50% could be directly correlated. Enhancement of benzodiazepine binding by GABA agonists is blocked by the GABA receptor antagonist (+) bicuculline; the stereoisomer (-) bicuculline is much less active (Tallman et al., 1978, Nature, 274: 383-85).

20 Soon after the discovery of high affinity binding sites for the benzodiazepines, it was discovered that a triazolopyridazine could interact with benzodiazepine receptors in a number of regions of the brain in a manner consistent with receptor heterogeneity or negative cooperativity. In these studies, Hill coefficients significantly less than one were observed in a number of brain regions, including cortex, hippocampus, and striatum. In cerebellum, 25 triazolopyridazine interacted with benzodiazepine sites with a Hill coefficient of 1 (Squires et al., 1979, Pharma. Biochem. Behav. 10: 825-30, Klepner et al. 1979, Pharmacol. Biochem.

201
189

Behav. 11: 457-62). Thus, multiple benzodiazepine receptors were predicted in the cortex, hippocampus, striatum, but not in the cerebellum.

Based on these studies, extensive receptor autoradiographic localization studies were carried out at a light microscopic level. Although receptor heterogeneity has been demonstrated (Young & Kuhar 1980, J. Pharmacol. Exp. Ther. 212: 337-46, Young et al., 1981 J. Pharmacol Exp. ther 216: 425-430, Niehoff et al. 1982, J. Pharmacol. Exp. Ther. 221: 670-75), no simple correlation between localization of receptor subtypes and the behaviors associated with the region has emerged from the early studies. In addition, in the cerebellum, where one receptor was predicted from binding studies, autoradiography revealed heterogeneity of receptors (Niehoff et al., 1982, J. Pharmacol. Exp. Ther. 221: 670-75).

A physical basis for the differences in drug specificity for the two apparent subtypes of benzodiazepine sites has been demonstrated by Sieghart & Karobath, 1980, Nature 286: 285-87. Using gel electrophoresis in the presence of sodium dodecyl sulfate, the presence of several molecular weight receptors for the benzodiazepines has been reported. The receptors were identified by the covalent incorporation of radioactive flunitrazepam, a benzodiazepine which can covalently label all receptor types. The major labeled bands have molecular weights of 50,000 to 53,000, 55,000, and 57,000 and the triazolopyridazines inhibit labeling of the slightly higher molecular weight forms (53,000, 55,000, 57,000) (Sieghart et al. 1983, Eur. J. Pharmacol. 88: 291-99).

At that time, the possibility was raised that the multiple forms of the receptor represent "isoreceptors" or multiple allelic forms of the receptor (Tallman & Gallager 1985, Ann. Rev. Neurosci. 8, 21-44). Although common for enzymes, genetically distinct forms of receptors have not generally been described. As we begin to study receptors using specific radioactive probes and electrophoretic techniques, it is almost certain that isoreceptors will emerge as important in investigations of the etiology of psychiatric disorders in people.

202 !!
190

The GABA_A receptor subunits have been cloned from bovine and human cDNA libraries (Schoenfield et al., 1988; Duman et al., 1989). A number of distinct cDNAs were identified as subunits of the GABA_A receptor complex by cloning and expression. These are categorized into α , β , γ , δ , , and provide a molecular basis for the GABA_A receptor heterogeneity and distinctive regional pharmacology (Shivvers et al., 1980; Levitan et al., 1989). The γ subunit appears to enable drugs like benzodiazepines to modify the GABA responses (Pritchett et al., 1989). The presence of low Hill coefficients in the binding of ligands to the GABA_A receptor indicates unique profiles of subtype specific pharmacological action.

Drugs that interact at the GABA_A receptor can possess a spectrum of pharmacological activities depending on their abilities to modify the actions of GABA. For example, the beta-carbolines were first isolated based upon their ability to inhibit competitively the binding of diazepam to its binding site (Nielsen et al., 1979, Life Sci. 25: 679-86). The receptor binding assay is not totally predictive about the biological activity of such compounds; agonists, partial agonists, inverse agonists, and antagonists can inhibit binding. When the beta-carboline structure was determined, it was possible to synthesize a number of analogs and test these compounds behaviorally. It was immediately realized that the beta-carbolines could antagonize the actions of diazepam behaviorally (Tenen & Hinrich, 1980, Nature 288: 609-10). In addition to this antagonism, beta-carbolines possess intrinsic activity of their own opposite to that of the benzodiazepines; they become known as inverse agonists.

In addition, a number of other specific antagonists of the benzodiazepine receptor were developed based on their ability to inhibit the binding of benzodiazepines. The best studied of these compounds is an imidazodiazepine (Hunkeler et al., 1981, Nature 290: 514-516). This compound is a high affinity competitive inhibitor of benzodiazepine and beta-carboline binding and is capable of blocking the pharmacological actions of both these classes of compounds. By itself, it possesses little intrinsic pharmacological activity in animals and humans (Hunkeler et al., 1981, Nature 290: 514-16; Darragh et al., 1983, Eur. J. Clin. Pharmacol. 14: 569-70). When

09387311 - 8319
6 FEB

a radiolabeled form of this compound was studied (Mohler & Richards, 1981, Nature 294: 763-65), it was demonstrated that this compound would interact with the same number of sites as the benzodiazepines and beta-carbolines, and that the interactions of these compounds were purely competitive. This compound is the ligand of choice for binding to GABA_A receptors because it
5 does not possess receptor subtype specificity and measures each state of the receptor.

The study of the interactions of a wide variety of compounds similar to the above has led to the categorizing of these compounds. Presently, those compounds possessing activity similar to the benzodiazepines are called agonists. Compounds possessing activity opposite to benzodiazepines are called inverse agonists, and the compounds blocking both types of activity
10 have been termed antagonists. This categorization has been developed to emphasize the fact that a wide variety of compounds can produce a spectrum of pharmacological effects, to indicate that compounds can interact at the same receptor to produce opposite effects, and to indicate that beta-carbolines and antagonists with intrinsic anxiogenic effects are not synonymous.


A biochemical test for the pharmacological and behavioral properties of compounds that
15 interact with the benzodiazepine receptor continues to emphasize the interaction with the GABAergic system. In contrast to the benzodiazepines, which show an increase in their affinity due to GABA (Tallman et al., 1978, Nature 274: 383-85, Tallman et al., 1980, Science 207: 274-81), compounds with antagonist properties show little GABA shift (i.e., change in receptor affinity due to GABA) (Mohler & Richards 1981, Nature 294: 763-65), and the inverse agonists
20 actually show a decrease in affinity due to GABA (Braestrup & Nielsen 1981, Nature 294: 472-474). Thus, the GABA shift predicts generally the expected behavioral properties of the compounds.

Various compounds have been prepared as benzodiazepine agonists and antagonists. For Example, U.S. Patents Nos. 3,455,943, 4,435,403, 4,596,808, 4,623,649, and 4,719,210,
25 German Patent No. DE 3,246,932, and Liebigs Ann. Chem. 1986, 1749 teach assorted


204.
192

benzodiazepine agonists and antagonists and related anti-depressant and central nervous system active compounds.



U.S. Patent No. 3,455,943 discloses compounds of the formula:



5 wherein R₁ is a member of the group consisting of hydrogen and lower alkoxy; R₂ is a member of the group consisting of hydrogen and lower alkoxy; R₃ is a member of the group consisting of hydrogen and lower alkyl; and X is a divalent radical selected from the group consisting of




and







10 and the non-toxic acid addition salts thereof.

Other references, such as U.S. Patent No. 4,435,403 and German patent DE 3,246,932 disclose compounds containing the following structural skeleton:




where A is carbon or nitrogen.

15 A variety of indole-3-carboxamides are described in the literature. For example, J. Org. Chem., 42: 1883-1885 (1977) discloses the following compounds.



J. Heterocyclic Chem., 14: 519-520 (1977) discloses a compound of the following

5 formula:



None of these indole-3-carboxamides includes an oxy substituent at the 4-position of the indole ring.

6780-TTE/886


206
194

SUMMARY OF THE INVENTION

This invention provides novel compounds of Formula I which interact with a GABA_A binding site, the benzodiazepine receptor.

The invention provides pharmaceutical compositions comprising compounds of Formula

- 5 I. The invention also provides compounds useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. Accordingly, a broad embodiment of the invention is directed to compounds of general Formula I:



I

or the pharmaceutically acceptable non-toxic salts thereof wherein:

G represents



where

- 15 Q is phenyl, 2- or 3-thienyl, or 2-, 3-, or 4-pyridyl, all of which may be mono or disubstituted with hydroxy or halogen;

T is halogen, hydrogen, hydroxyl, amino or alkoxy having 1-6 carbon atoms;

W is oxygen, nitrogen, sulfur, or CR₇R₈ where R₇ and R₈ are the same or different and represent hydrogen, alkyl, or R₇-R₈ taken together

- 20 represents a cyclic moiety having 3-7 carbon atoms;

X is hydrogen, hydroxyl, or alkyl;

207.1
195

Z is

hydroxy, C₃-C₇ cycloalkylC₁-C₆ alkoxy, or C₁-C₆ alkoxy where the alkyl portion is optionally substituted with amino, mono- or diC₁-C₆alkylamino, or azaC₁-C₆cycloalkyl;

5

amino, or mono- or diC₁-C₆alkylamino, (trifluoromethyl)methyl, C₃-C₇ cycloalkyl optionally substituted with C₁-C₆ alkyl, or C₃-C₇cycloalkylC₁-C₆alkyl;

phenyl or phenyl(C₁-C₆)alkyl where the phenyl portion is optionally substituted with C₁-C₆ alkyl, hydroxy, C₁-C₆ alkoxy, trifluoromethyl, halogen, amino, or mono- or diC₁-C₆ alkylamino;

10

2-, 3-, or 4-pyridyl, 1- or 2-imidazolyl, 1-, 2-, or 3-pyrrolyl, or adamantane-2-yl; each of which may be substituted on a tertiary carbon or a secondary nitrogen with C₁-C₆alkyl;

NR₉COR₁₀ where R₉ and R₁₀ are the same or different and represent


15

hydrogen or C₁-C₆ alkyl or C₃-C₇ cycloalkyl; or

Z is connected, optionally through W, to Q to form a 1-6 membered ring; or

Z represents

a group of the formula:



20

where


p is 1, 2, or 3;

D and D' independently represent oxygen, NR_y or CHR_y, provided that only one of D and D' may be NR_y where each R_y is hydrogen or C₁-C₆ alkyl; or and

09387311.08319

R_1 is hydrogen or C_1-C_6 alkyl;

a group of the formula:




p is 1, 2, or 3;

q is 0, 1, or 2;

R_2 is hydrogen or C_1-C_6 alkyl; or

a group of the formula:




where

s is 0, 1, 2 or 3, and the sum of s and m is not less than 1;

R_3 is hydroxy, C_1-C_6 alkoxy, amino, mono- or di C_1-C_6 alkylamino

where each alkyl is independently optionally substituted with
amino, mono- or di C_1-C_6 alkylamino, or

R_4 is a group of the formula



where p, D, D', and R_4 are as defined above;

, , and  independently represent a carbon chain optionally substituted with hydrogen, halogen, or straight or branched chain lower alkyl having 1-6 carbon atoms;

wherein

k is 0, 1, 2, or 3;

m is 0, 1, 2, or 3; and

n is 0, 1, 2, or 3;

R₃, R₄, R₅, and R₆ are the same or different and are selected from hydrogen, alkyl, -

COR₁₁ or -CO₂R₁₁ where R₁₁ is alkyl or cycloalkyl having 3-7 carbon atoms;

or -CONR₁₂R₁₃ where R₁₂ and R₁₃ are selected independently from hydrogen,

alkyl, cycloalkyl having 3-7 carbon atoms, phenyl, 2-, 3-, or 4-pyridyl, or

NR₁₂R₁₃ forms a heterocyclic group which is morpholinyl, piperidinyl,

pyrrolidinyl, or N-alkyl piperazinyl; or

R₃-R₄ may be taken together to form a cyclic moiety having 3-7 carbon atoms; or

R₅-R₆ may be taken together to form a cyclic moiety having 3-7 carbon atoms; and

where each alkyl group forming an R₃, R₄, R₅, or R₆ substituent or portion

thereof may be substituted independently with hydroxy or mono- or

dialkylamino where each alkyl is independently alkyl or cycloalkyl


having 3-7 carbon atoms.

These compounds are highly selective agonists, antagonists or inverse agonists for GABA_A brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA_A brain receptors. In other words, while the compounds of the invention all interact with GABA_A brain receptors, they do not display identical physiologic activity. Thus, these compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory. For example, these compounds can be used to treat overdoses of benzodiazepine-type drugs as they would competitively bind to the benzodiazepine receptor.

DETAILED DESCRIPTION OF THE INVENTION

The novel compounds encompassed by the instant invention can be described by general formula I set forth above or the pharmaceutically acceptable non-toxic salts thereof.


In addition, the present invention encompasses compounds of Formula II.



II

wherein Y is hydrogen, halogen, or hydroxy; and W, Y, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.

The present invention also encompasses compounds of Formula III




III


wherein Y is hydrogen, halogen, or hydroxy; and W, Y, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.

The present invention also encompasses compounds of Formula IV, IVa, IVb, and IVc.


24 . :
199 .




IV



IVa



IVb




IVc

wherein Y is hydrogen, halogen, or hydroxy; and W, Y, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.

The present invention also encompasses compounds of Formula V.

0 387311-0878E 0

27.6
20




V

wherein Y is hydrogen, halogen, or hydroxy; and W, Y, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.

5

The present invention also encompasses compounds of Formula VI.




VI

wherein Y is hydrogen, halogen, or hydroxy; and W, Y, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.

10

The present invention also encompasses compounds of Formula VII.



VII


wherein W, Z, m, n, R₃, R₄, R₅, and R₆ are defined as above.

15

66780-TFE/BE60

213
201

The present invention also encompasses compounds of Formula VIII.




VIII

wherein W, Z, m, n, R₃, R₄, R₅, and R₆ are defined as above.

5

The present invention also encompasses compounds of Formula IX.




IX

wherein W, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.

10

The present invention also encompasses compounds of Formula X.




X

wherein W, Z, k, m, n, R₃, R₄, R₅, and R₆ are defined as above.


15

0387311-0883199

Preferred compounds of the invention are those where n is 1 or 2. Particularly preferred are those where X and T are both hydrogen. Thus, preferred compounds of the invention have formulas Ia and Ib.




Ia



Ib

Preferred G substituents of the invention include the following:




A

where R_a represents hydrogen or alkyl where the alkyl is optionally halogenated; and e is an integer of 1-3.

More preferred G substituents of formula A include those where e is 1, 2, or 3, and R_a is hydrogen, methyl, ethyl, isopropyl, or cyclopropyl. Particularly preferred G substituents of formula A include those where e is 1, 2, or 3, and R_a is hydrogen or methyl.

Another preferred G substituent is the following formula:




B

where R_a represents hydrogen or alkyl where the alkyl is optionally halogenated; and e is an integer of 1-3.

215
203

More preferred G substituents of formula B include those where e is 1, 2, or 3; and R_a is hydrogen, methyl or ethyl. Particularly preferred G substituents of formula B include those where e is 1 or 2, and R_a is hydrogen or methyl.

5 Another preferred G substituent is the following formula:



where


Hal represents a halogen, preferably fluoro, bromo, or chloro;

R_a and R_b independently represent hydrogen, C₁-C₆ alkyl, C₃-C₇cycloalkyl, C₃-C₇cycloalkylC₁-C₆alkyl where the cycloalkyl group may be substituted with halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or mono- or diC₁-C₆ alkylamino; and e is an integer of 2-3.

Preferred compounds having formula C as the G group include those where Hal is fluoro and e is 2, 3, or 4.

More preferred G substituents of formula C include those where R_a is hydrogen, methyl or ethyl; and R_b is hydrogen. Particularly preferred G substituents of formula C include those where e is 2; R_a is hydrogen or methyl; and R_b is hydrogen.

20 Another preferred G substituent is the following formula:



09387311-08319

where

Hal represents a halogen, preferably fluoro, bromo, or chloro;

R_a and R_b independently represent hydrogen, C₁-C₆ alkyl, C₃-C₇ cycloalkyl, C₇-


C₇ cycloalkyl(C₁-C₆ alkyl where the cycloalkyl group may be substituted with

halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, or mono- or diC₁-C₆ alkylamino; and

e is an integer of 2-3.


Preferred compounds having formula C-1 as the G group include those where Hal is fluoro and e is 2, 3, or 4.

Another preferred G substituent is the following formula:



D

where R_a represents hydrogen, alkyl, or C₃-7 cycloalkyl, or a group of the formula:



where

p is 1, 2, or 3;

D and D' independently represent oxygen, NR_y, or CHR_y, provided that only one of D and D' may be NR_y, where each R_y is hydrogen or C₁-C₆ alkyl; and

R_z is hydrogen or C₁-C₆ alkyl; and

R_b represents hydrogen, alkyl, or acyl;


Y and Y' independently represent hydrogen or halogen; and

27
205

e is an integer of 1-3.

More preferred G substituents of formula D are those where Y is hydrogen or fluorine; and e is 1 or 2. Particularly preferred G substituents of formula D are those where Y is hydrogen or fluorine; e is 1 or 2; R_a is hydrogen, C₁₋₃ alkyl, or cyclopropyl, and R_b is hydrogen, methyl, or acyl. Other particularly preferred G substituents of formula D are those where Y is hydrogen and Y' is fluorine. Still other particularly preferred G groups of Formula D are those where e is 1 or 2; R_a is hydrogen, C₁₋₃ alkyl, cyclopropyl or cyclopropylmethyl, and R_b is hydrogen, methyl, or acyl.

Another preferred G substituent is the following formula:



D-1

where R_a represents hydrogen, alkyl, or C₃₋₇ cycloalkyl; and

R_b represents hydrogen, alkyl, or acyl; or

R_a and R_b independently represent hydrogen, C_{1-C₆} alkyl, C₃₋₇cycloalkylC_{1-C₆}alkyl;

and

Y and Y' independently represent hydrogen or halogen; and


e is an integer of 1-3.

More preferred G substituents of formula D are those where Y is hydrogen or fluorine; and e is 1 or 2. Particularly preferred G substituents of formula D are those where Y is hydrogen or fluorine; e is 1 or 2; R_a is hydrogen, C₁₋₃ alkyl, or cyclopropyl, and R_b is hydrogen, methyl, or acyl. Other particularly preferred G substituents of formula D are those where Y is hydrogen and Y' is fluorine. Still other particularly preferred G groups of Formula

218
206

D are those where e is 1 or 2; R_a is hydrogen, C₁₋₃ alkyl, cyclopropyl or cyclopropylmethyl, and R_b is hydrogen, methyl, or acyl.

5 Another preferred G substituent is the following formula:




E

where Z is oxygen, nitrogen, or methylene; and m is 1 or 2.

10 Particularly preferred G substituents of formula E are those where Z is oxygen, and m is 1 or 2. Other particularly preferred G substituents of formula E are those where Z is nitrogen, and m is 1 or 2.

Another preferred G substituent is the following formula:




F

15 where Z is oxygen or nitrogen; and m is 1 or 2.

Particularly preferred G substituents of formula F are those where Z is nitrogen, and m is 1 or 2.

Another preferred G substituent is the following formula:



H


09387311-08319

219:
207:

where Z is oxygen, nitrogen, or methylene; and m is 1 or 2.

Particularly preferred G substituents of formula H are those where Z is nitrogen, and m is 1 or 2.

5 Another preferred G substituent is the following formula:



J

where R_a represents hydrogen, alkyl, or C₃₋₇ cycloalkyl;


R_b represents hydrogen, alkyl, or acyl;

Y and Y' independently represent hydrogen or halogen; and

e is an integer of 1-3.

More preferred G substituents of formula J are those where Y and Y' are independently hydrogen or fluorine; and e is 1 or 2. Particularly preferred G substituents of formula J are those where Y and Y' are independently hydrogen or fluorine; e is 1 or 2; R_a is hydrogen, C₁₋₃ alkyl, or cyclopropyl, and R_b is hydrogen, methyl, or acyl.

Another preferred G substituent is the following formula:



K

where

0
3
7311-083199

10


15

220
208

R_a and R_b independently represent hydrogen, C_1-C_6 alkyl, C_3-C_6 cycloalkyl, C_3-C_6 cycloalkyl- C_1-C_6 alkyl where the cycloalkyl group may be substituted with halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, or mono- or di- C_1-C_6 alkylamino; and e is an integer of 2-3.

5

Another preferred G substituent is represented by the following formula:




M

where

R_h is hydrogen, halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, or trifluoromethyl; s is 0, 1, 2 or 3, and the sum of s and m is not less than 1;

R_o is hydroxy, C_1-C_6 alkoxy, amino, mono- or di- C_1-C_6 alkylamino where each alkyl is independently optionally substituted with amino, mono- or di- C_1-C_6 alkylamino, or

R_o is a group of the formula



p is 1, 2, or 3;

D and D' independently represent oxygen, NR_z , or CHR_z , provided that only one of D and D' may be NR_z , where each R_z is hydrogen or C_1-C_6 alkyl; or and

R_z is hydrogen or C_1-C_6 alkyl.

0 387311-08319


10

15

20

22T
209

Preferred M groups are those where R_1 is hydrogen or halogen, most preferably fluoro, and R_2 is a group of the formula:



where

5

R_{14} is hydrogen or C_1 - C_6 alkyl;

R_{15} is hydrogen or C_1 - C_6 alkyl;

R_{16} is hydrogen, ethyl, or methyl;


R_{17} is C_1 - C_6 alkyl; and

J is a C_1 - C_4 alkylene group, preferably methylene, ethylene, or propylene.

10

Particularly preferred groups of Formula M include those where s is 1 and R_2 is ethoxy, hydroxy, ethylamino, diethylamino, morpholinyl, piperazinyl, 4-methylpiperazinyl,

66 FEB - TTE/BEE




15


Representative compounds of the invention are shown below in Table 1.

227
210


Table 1




Compound 1




Compound 2




Compound 3




Compound 4




Compound 5




Compound 6




Compound 7



Compound 8




Compound 9




Compound 10


6TES - TFE/BE60




Compound 11




Compound 12




Compound 13




Compound 14




Compound 15




Compound 47




Compound 86



Compound 95




Compound 115




Compound 145

667E80-TTE/8E60


22A :
212




Compound 148




Compound 149




Compound 179




Compound 222




Compound 226



Compound 227





Compound 229



Compound 235

The following numbering conventions are used to identify positions on the ring systems in the compounds of the invention:



Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Table I and their pharmaceutically acceptable salts. Non-toxic pharmaceutically acceptable salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic, alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Table I and their pharmaceutically acceptable salts. The present invention also encompasses the acylated prodrugs of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I.


By "alkyl" or "lower alkyl" in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms, such as, for example, methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl.

By "alkoxy" or "lower alkoxy" in the present invention is meant straight or branched chain alkoxy groups having 1-6 carbon atoms, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, 2-pentyl, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy.

By "diC₁-C₆alkylamino" is meant an amino group carrying two C₁-C₆alkyl groups that are the same or different.

By "benzoxazinyf" as used herein is meant a moiety of the formula:

226
214




A benzoxazin-6-yl group is depicted.

By "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

By "2-hydroxyethoxy" is meant a group of the formula: $-OCH_2CH_2OH$.

5 By a 2-, 3-, or 4-pyridyl, 1- or 2-imidazolyl, 1-, 2-, or 3-pyrrolyl, or adamantane-2-yl group that is substituted on a tertiary carbon or a secondary nitrogen with C_1 - C_6 alkyl is meant any such group in which a hydrogen atom is replaced with an appropriate alkyl group. By way of example, such groups include the following:



10 By "N-alkylpiperazyl" in the invention is meant radicals of the formula:



where R is a straight or branched chain lower alkyl as defined above.

15 The pharmaceutical utility of compounds of this invention are indicated by the following assay for GABA_A receptor binding activity.

Assays are carried out as described in Thomas and Tallman (J. Bio. Chem. 156: 9838-9842, J. Neurosci. 3: 433-440, 1983). Rat cortical tissue is dissected and homogenized in 25 volumes (w/v) of 0.05 M Tris HCl buffer (pH 7.4 at 4°C). The tissue homogenate is centrifuged in the cold (4°C) at 20,000 x g for 20'. The supernatant is decanted and the pellet is rehomogenized in the same volume of buffer and again centrifuged at 20,000 x g. The supernatant is decanted and the pellet is frozen at -20°C overnight. The pellet is then thawed

0 387311-083199

227
215

and rehomogenized in 25 volume (original wt/vol) of buffer and the procedure is carried out twice. The pellet is finally resuspended in 50 volumes (w/vol of 0.05 M Tris HCl buffer (pH 7.4 at 40°C).

Incubations contain 100 ml of tissue homogenate, 100 ml of radioligand 0.5 nM (³H-RO15-1788 [³H-Flumazenil] specific activity 80 Ci/mmol), drug or blocker and buffer to a total volume of 500 ml. Incubations are carried for 30 min at 4°C then are rapidly filtered through GFB filters to separate free and bound ligand. Filters are washed twice with fresh 0.05 M Tris HCl buffer (pH 7.4 at 4°C) and counted in a liquid scintillation counter. 1.0 mM diazepam is added to some tubes to determine nonspecific binding. Data are collected in triplicate determinations, averaged and % inhibition of total specific binding is calculated. Total Specific Binding = Total - Nonspecific. In some cases, the amounts of unlabeled drugs is varied and total displacement curves of binding are carried out. Data are converted to K_i's; results for compounds of this invention are listed in Table 2.

Table 2

Compound Number	K _i (nM)
1	90
2	29
3	49
4	0.24
5	9
6	9
7	30
8	27
9	1.3
10	37
11	7
12	5
13	24

09387311-0831 9

14
15

3
12

The compounds of general formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general formula I and a pharmaceutically acceptable carrier. One or more compounds of general formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For

09387311-083199

229.1
217

example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

230-
216

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

221
219

The compounds of general formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug.

5 Such materials are cocoa butter and polyethylene glycols.

Compounds of general formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can be dissolved in the vehicle.

10 Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from
15 about 1 mg to about 500 mg of an active ingredient.


It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing
20 therapy.

An illustration of the preparation of compounds of the present invention is given in Scheme I.

09387311-08319

232
2201

Scheme I




5 where:

0387311-0831 9

223.
221

Ar is




where Q, W, k, m, n, Z, R₃, R₄, R₅, and R₆ are as defined above.

5 Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention, as demonstrated by the following examples.

In some cases protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general the need for such protecting groups will be apparent to those skilled in the art of organic synthesis as well as the conditions necessary to attach and remove such groups. Representative examples of the preparation of various protected aniline derivatives are shown in Schemes II (1), (2) and (3).

Scheme II


(1)




09387311-083899

23A
23271

(2)



(3)



5

Compounds of Formula I where G is a group of, for example, formulas C, C-1, D, D-1, K or M can be made using the above outlined methods and, e.g., additional ester and amide coupling reactions. In certain situations, protection of the indole ring nitrogen will be necessary.

For example, compounds where R₉ is a dialkylamino group can be prepared from a 2-(4-nitrophenoxy)ethan-1-ol made as described above and oxidation of the hydroxy group, and subsequent formation of an acid chloride or active ester. The active ester or acid chloride may

275
273

then be coupled to an appropriate amine and the resulting nitrophenyl compound used as shown in the above schemes.

The disclosures in this application of all articles and references, including patents, are incorporated herein by reference in their entirety.

5 One skilled in the art will recognize that modifications may be made in the present invention without deviating from the spirit or scope of the invention. The invention is illustrated further by the following examples which are not to be construed as limiting the invention or scope of the specific procedures or compositions described herein.


Example 1

10 Preparation of starting materials and intermediates

The starting materials and various intermediates may be obtained from commercial sources, prepared from commercially available organic compounds, or prepared using well known synthetic methods.

15 Representative examples of methods for preparing intermediates of the invention are set forth below.

1. 4-oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid



4-Oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid is prepared according to the following procedure. Potassium hydroxide (345 g, 6.15 mol) is dissolved in methyl alcohol (1.2 L) then cooled in an ice water bath. A solution of cyclohexanedione (714 g, 6.15 mol) in methyl alcohol (1.2 L), dissolved using gentle heat, is added dropwise to the cold, stirred KOH solution over 2 h. A solution of ethyl bromopyruvate (1200 g, 6.15 mol) in methyl alcohol (1.5 L) is then added dropwise over 3 h. The reaction mixture is allowed to reach ambient temperature

0 387311-08199


226
224

and stirred an additional 14.5 h. While cooling the reaction mixture via a water bath, a solution of sodium hydroxide (492 g, 12.4 mol) in water (984 mL) is added dropwise over 2.5 h. After stirring at ambient temperature for 15.5 h, the reaction mixture is cooled in an ice water bath, 500 g of ice added, and the resulting mixture is then acidified with concentrated hydrochloric acid (ca 1L) to pH 1. The reaction mixture is concentrated *in vacuo*, 1L of ice is added, and the precipitate filtered, washed with ice water (3 X 200 mL), and then dried in a vacuum oven at 75° C to afford 4-oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid (560 g). m.p. 137-138° C.

9387311-08319

237
225


2. 4-oxo-4,5,6,7-tetrahydroindole-3-carboxylate



To a stirred mixture of 4-oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid (640 g, 3.55 mol), potassium carbonate (1.7 kg, 10.65 mol) and cesium carbonate (100 g, 0.32 mol) in N,N-dimethylformamide (9.0 L) is added iodoethane (1250 g, 8.01 mol). The mixture is heated at 60° C for 2 h. After cooling to ambient temperature, the mixture is filtered, the solid is rinsed with ethyl acetate, and the filtrate concentrated *in vacuo*. Water (2 L) is added then extracted with ethyl acetate (2 X 2L); the combined organic extracts are washed with brine, dried over magnesium sulfate, filtered, and concentrated *in vacuo* to give ethyl 4-oxo-4,5,6,7-tetrahydrobenzofuran-3-carboxylic acid (642 g). A mixture of this ester (640 g, 3.07 mol) and ammonium acetate (426 g, 5.53 mol) in N,N-dimethylformamide (320 mL) is heated to 100° C for 2 h. The reaction mixture is concentrated *in vacuo*, ice water (2.5L) is added, and extracted with dichloromethane (2 X 3L); the combined organic extracts are washed with brine, dried over magnesium sulfate, filtered, and concentrated *in vacuo* to give ethyl 4-oxo-4,5,6,7-tetrahydroindole-3-carboxylate (357 g). A mixture of this ester (170 g, 0.82 mol) in ethyl alcohol (250 mL) and a solution of sodium hydroxide (165 g, 4.1 mol) in water (1 L) is heated at reflux for 1 h, then cooled in an ice water bath. Concentrated hydrochloric acid (350 mL) is added dropwise, the precipitate collected by filtration, rinsed with ice water (3 X), and dried in a vacuum oven at 75° C to afford 4-oxo-4,5,6,7-tetrahydroindole-3-carboxylate (125 g). m.p. 269-270° C.

3. 4-[N-trifluoroacetyl-(methylaminomethyl)aniline

228
226



A solution of p-nitrobenzylbromide (5.40 g, 25 mmol) in acetonitrile (60 ml) is added dropwise to a stirred solution of aqueous methylamine (65 mL, 40 wt.%, 0.75 mol) in acetonitrile (50 mL) at 0°. After stirring an additional 15 minutes, the solution is poured into
5 brine and extracted 2X with dichloromethane. The combined organic layers are washed with brine, dried over sodium sulfate, filtered, and concentrated *in vacuo* to give 4-(methylaminomethyl)nitrobenzene (4.04g).

0938731-083199
661E80-TFE/BEE60


10 A solution of trifluoroacetic anhydride (4.46 mL, 31.6 mmol) in dichloromethane (10 mL) is added dropwise to a stirred solution of 4-(methylaminomethyl)nitrobenzene (4.04g, 24.3 mmol) and pyridine (2.16 mL, 26.7 mmol) in dichloromethane (25 mL) at 0°. After stirring an additional 30 minutes, the solution is poured into aqueous 3.6N hydrochloric acid and extracted with dichloromethane. The organic layer is washed with brine, dried over sodium sulfate, filtered, and concentrated *in vacuo* to give 4-[N-trifluoroacetyl-
(methylaminomethyl)]nitrobenzene (6.55 g).

15 Crude 4-[N-trifluoroacetyl-(methylaminomethyl)]nitrobenzene (6.55 g) is dissolved in ethyl alcohol (75 mL), added to 10% Pd/C (655 mg) in a Parr bottle and shaken under Hydrogen (50 PSI) for 4 hours. The mixture is filtered through Celite and concentrated *in vacuo* to give 4-[N-trifluoroacetyl-(methylaminomethyl)aniline (5.75 g).

20 The 3-aminoalkylanilines are prepared in a similar fashion according to the procedure generally set forth in part (1) of Scheme II above.

4. 4-amino-(N-trifluoroacetyl-2-methylaminoethoxy)benzene

229
229.1



A mixture of p-nitrophenol (1.39 g, 10 mmol), 2-chloroethoxytrimethylsilane (3.2 ml, 20 mmol), potassium carbonate (4.15 g, 30 mmol), cesium carbonate (163 mg, 0.5 mmol), and sodium iodide (149 mg, 1 mmol) in N,N-dimethylformamide (10 ml) is heated at 75° for 19.5 hours. After cooling to ambient temperature, the mixture is diluted with ethyl acetate and filtered. The filtrate is washed with saturated aqueous sodium bicarbonate, then washed 2X with water, dried over magnesium sulfate, filtered, concentrated *in vacuo*, and purified on Silica gel (1:1 ethyl acetate / hexanes) to give 4-nitro-(2-Hydroxyethoxy)benzene (1.25 g).

4-Nitro-(2-Hydroxyethoxy)benzene (1.13 g, 6.2 mmol) in thionyl chloride (10 mL) is heated at reflux for 3 hours then concentrated *in vacuo*. After cooling the residue in an ice water bath, saturated aqueous sodium bicarbonate is added and the precipitate collected, rinsed with water, and dried to give 4-nitro-(2-chloroethoxy)benzene (909 mg).


A mixture of 4-nitro-(2-chloroethoxy)benzene (781 mg, 3.9 mmol) and aqueous methylamine (15 mL, 40 wt. %) in isopropyl alcohol (15 mL) is heated in a sealed tube at 100° for 4 hours. After cooling in an ice water bath, the mixture is poured into brine and extracted 2X with dichloromethane, dried over sodium sulfate, filtered, and concentrated *in vacuo* to give 4-nitro-(2-methylaminoethoxy)benzene (697 mg).

To a solution of 4-nitro-(2-methylaminoethoxy)benzene (766 mg, 3.9 mmol) and pyridine (0.35 mL, 4.29 mmol) in dichloromethane (5 mL) at 0° C is added dropwise trifluoroacetic anhydride (0.72 mL, 5.08 mmol). After stirring at 0° C for 3.5 hours, the mixture is poured into aqueous 1.2 N hydrochloric acid and extracted with dichloromethane. The organic layer is washed with saturated aqueous sodium bicarbonate then brine, dried over sodium sulfate, filtered, and concentrated *in vacuo* to give 4-nitro-(N-trifluoroacetyl-2-methylaminoethoxy)benzene (1.06 g). Treatment of this nitro compound with 10% Palladium

240
278

on carbon in ethyl alcohol (18 mL) in a Parr bottle under Hydrogen (55 PSI) for 2.25 hours affords 4-amino-(N-trifluoroacetyl-2-methylaminoethoxy)benzene (709 mg).

Example 2



To a stirred solution of 4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxylic acid (100 mg, 0.6 mmol) and triethylamine (0.15 mL, 1.1 mmol) in N,N-dimethylformamide (5 mL) at 0° C is added ethyl chloroformate (0.1 mL, 1.1 mmol). After stirring an additional 1 hour, 3-(N-trifluoroacetyl-(methylaminomethyl)aniline (0.3 g, 1.3 mmol) is added. The reaction mixture is stirred for 4 hours, then poured into saturated aqueous ammonium chloride and extracted 2X with ethyl acetate. The combined organic layers are washed sequentially with brine, aqueous 2N hydrochloric acid, then brine, dried over sodium sulfate, filtered, and concentrated *in vacuo*. To the residue is added 15% aqueous potassium bicarbonate (5 mL) and methyl alcohol (3 mL), then heated at reflux for 3 hours. After cooling, the reaction mixture is extracted with ethyl acetate, the organic layer dried over sodium sulfate, filtered, and concentrated *in vacuo* to give N-[3-(methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; (Compound 1) m.p. 130-132°C.

Example 3

The following compounds are prepared essentially according to the procedures described in Examples 1-5:

(a) N-[3-(Methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 1); mp 130-132° C.

24
229

(b) N-[4-(Hydroxyethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 16); mp 245-247° C.

(c) N-[4-(Methoxyethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 2).

5 (d) N-[4-(3-Methylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 17); mp 233-236° C.

(e) N-[4-(Methoxymethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 18); mp 164-165° C.

10 (f) N-[4-(Aminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 6); mp >200° C (d).

(g) N-[4-(Methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 19); mp 217-219° C.

15 (h) N-[2-Fluoro-4-(methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 3); mp 186-188° C.

(i) N-[4-[N-acetyl-(methylaminomethyl)phenyl]]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 20); mp 204-206° C.

20 (j) N-[4-(Ethylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 21); mp 194-195° C.

(k) N-[4-(Isopropylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 22); mp 164-166° C.

(l) N-[4-(Cyclopropylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 5); mp 171-173° C.

242
230

(m) N-[4-(Dimethylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 23); mp 216-218°C.

(n) N-[4-(2-Aminoethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 24); mp 85-90° C.

5 (o) N-[4-(2-Methylaminoethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 4); mp 197-200° C.

(p) N-[4-(Methoxymethyl)phenyl]-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 25).

(q) N-[4-(Methylaminomethyl)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 7); mp 173-175° C.

(r) N-{4-[N-acetyl-(methylaminomethyl)phenyl]}-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 26); mp 159-161° C.

(s) N-[4-(Methylaminomethyl)phenyl]-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 27); mp 217-219° C.

15 (t) N-[4-(Hydroxymethyl)phenyl]-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 28); mp 260-262° C.

(u) N-[4-(2-Hydroxyethoxy)phenyl]-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 9); mp 245-247° C.

(v) N-[3-(Methylaminomethyl)phenyl]-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 29); mp 172-174° C.

(w) N-[4-(2-Hydroxyethoxy)phenyl]-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 30); mp 268-270° C.

(x) N-[3-(Hydroxymethyl)phenyl]-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 8); mp 233-235°C.

0 387311-083199
667E80-11E8E

(y) N-[4-(Hydroxymethyl)phenyl]-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 31); mp 245-247°C.

(z) N-[4-(Methylaminomethyl)phenyl]-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 32); mp 230-232°C.

5 (aa) N-(1,3-Benzodioxol-5-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 10); mp 248-249° C.

(bb) N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 11); mp 254-256° C.

10 (cc) N-(3,4-Dihydro-2H-1,4-benzoxazin-6-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 33); mp 216°C.

(dd) N-(2,2-Dimethyl-1,3-benzodioxol-5-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 34).

(ee) N-(2,3-Dihydro-1H-indol-5-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 35); mp 283-286°C.

15 (ff) N-(2,3-Dihydro-1H-indol-6-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 13); mp 322-323°C.

(gg) N-(1,3-Benzodioxol-5-yl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 36).

20 (hh) N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 37); mp 241-243°C.

(ii) N-(4H-1,3-Benzodioxin-7-yl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 38); mp 251-252°C.

(jj) N-(1,3-Benzodioxol-5-yl)-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 39); mp 210-212° C.

(kk) N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 12); mp 222-223° C.

(ll) N-(2,2-Dimethyl-1,3-benzodioxol-5-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 40); mp 155-157°C.

5 (mm) N-(1,3-Benzodioxol-5-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 41); mp 297-299°C.

(nn) N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 42); mp 290-292°C.

10 (oo) N-(1,3-Benzodioxol-5-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 43); mp 245-246° C.

(pp) N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 44).

(qq) N-(4H-1,3-Benzodioxin-7-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 45); mp 234-236° C.

15 (rr) N-[(2-Hydroxyethoxy)pyrid-5-yl]-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 15); mp 221-223° C.

(ss) N-(3,4-Dihydro-2H-1,4-benzoxazin-7-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 46).

20 (tt) N-[3-(2-Pyrrolidinylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; [alternate name: (4-oxo(5,6,7-trihydroindol-3-yl))-N-[4-(2-pyrrolidinylethoxy)phenyl]carboxamide] (Compound 47);

(uu) N-[3-(2-Dimethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide [alternate name: (4-oxo(5,6,7-trihydroindol-3-yl))-N-[4-(2-Dimethylaminoethoxy)phenyl]carboxamide] (Compound 48);

245.
233

(vv) N-[3-(2-n-Propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 49).

(ww) N-[3-(2-n-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 50).

5 (xx) N-[3-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 51) (syrup).

(yy) N-[3-(2-Cyclobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 52).

(zz) N-[3-(2-t-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 53).

(aaa) N-[3-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 54).

(bbb) N-[3-[2-(4-Methylcyclohexyl)aminoethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 55).

15 (ccc) N-[3-[2-(3-Trifluoromethylbenzylamino)ethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 56).

(ddd) N-[3-[3-(3-Trifluoromethylbenzylamino)propoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 57).

20 (eee) N-[4-(2-Dimethylaminoethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 58).

(fff) N-[4-(2-Pyrrolidin-1-ylethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 59); mp 184-186°C.

(ggg) N-[4-(2-Diisopropylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 60).

25 (hhh) N-[4-(2-Methylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 61).

246
234

(iii) N-[4-(2-Ethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 62); mp 140-141°C.

(iii) N-[2-Fluoro-4-(2-ethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 63).

5 (kkk) N-[4-(2-n-Propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 64); mp 130-133°C.

(lll) N-[2-Fluoro-4-(2-n-propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 65).

(mmm) N-[3-Fluoro-4-(2-n-propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 66).

(mmm-a) N-[3-Fluoro-4-(2-n-propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 67); mp 373°C.

(mm) N-[4-(2-Cyclopropylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 68).

15 (ooo) N-[4-(2-Isopropylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 69); mp 284-286°C.

(ppp) N-[4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 70).

20 (ppp-a) N-[4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hemifumarate (Compound 71); mp 234-234°C.

(qqq) N-[2-Fluoro-4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 72); mp 247-250°C.

(rrr) N-[3-Fluoro-4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 73).

25 (rrr-a) N-[3-Fluoro-4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide tosylate (Compound 74); mp 222°C.

242
235

(ss) N-[4-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 75); dust.

(tt) N-[2-Fluoro-4-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 76); mp 152-155°C.

5 (uu) N-[3-Fluoro-4-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 77); mp 147-149°C.

(vv) N-[4-(2-n-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 78).

(vvv-a) N-[4-(2-n-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 79); mp 187-190°C.

(www) N-[3-Fluoro-4-(2-n-butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 80).

(xxx) N-[4-(2-t-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 81); mp 290-292°C.

15 (yyy) N-[3-Fluoro-4-(2-t-butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 82).

(aaa) N-[4-(2-adamant-2-ylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 83); mp 144-149°C.

20 (bbb) N-[4-[(R)-Pyrrolidin-2-ylmethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 84); mp 164-167-170°C.

(ccc) N-[4-[(S)-Pyrrolidin-2-ylmethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 85); mp 165-167°C.

(ddd) N-[4-(Piperidin-3-ylmethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 86).

25 (ddd-a) N-[4-(Piperidin-3-ylmethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 87); mp 196-199°C.

240
236

(eeee) N-[4-(2-Dimethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 88); mp 201°C.

(ffff) N-[3-Fluoro-4-(2-dimethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 89); mp 203°C.

5 (gggg) N-[4-(2-Pyrrolidin-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 90); mp 164-168°C.

(hhhh) N-[4-(2-Imidaz-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 91); mp 226-230°C.

10 (iiii) N-[3-Fluoro-4-(2-morpholin-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 92); mp 200°C.

(jjjj) N-[3-Fluoro-4-(2-pyrrolidin-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 93).

(kkkk) N-[4-(2-Piperidin-2-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 94); mp 281-285°C.

15 (llll) N-{4-[3-(2,2,2-Trifluoroethyl)aminopropoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 95).

(mmmm) N-[4-(3-Isopropylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 96).

20 (nnnn) N-{4-[3-(2-Methylpropyl)aminopropoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 97).

(oooo) N-[4-(3-Isobutylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 98).

(pppp) N-[4-(3-Cyclopropylmethylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 99).

25 (qqqq) N-{4-[3-(3-Ethylpropyl)aminopropoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 100).

09387311
TEE/8560
931
9

249
239

(nnr) N-[4-(3-Cyclopentylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 101).

(sss) N-[4-[3-(N-Cyclopropylmethyl,N-propyl)aminopropoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 102).

5 (ttt) N-[4-(2-Methylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 103).

(uuu) N-[4-(2-Ethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 104).

(uuu-a) N-[4-(2-Ethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 105); mp 178-180°C.

(vvvv) N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 106).

(vvvv-a) N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 107); mp 177-178°C.

15 (www) N-[4-(2-Isopropylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 108).

(www-a) N-[4-(2-Isopropylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 109); mp 167-169°C.

20 (xxx) N-[4-(2-n-Butylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 110).

(xxx-a) N-[4-(2-n-Butylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 111); mp 157-159°C.

(yyy) N-[4-(2-t-Butylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 112); mp 274-278°C.

25 (zzz) N-[4-(2-Benzylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 113)

250
238

(zzzz-a) N-[4-(2-Benzylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 114); mp 143-145°C.

(aaaaa-a) N-[4-(Pyrid-3-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 115).

5 (aaaaa) N-[4-(Pyrid-3-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 116); mp 276-277°C.

(bbbbbb) N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 117).

10 (bbbbbb-a) N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 118); mp 293°C.

(ccccc) N-{4-[(R)-Pyrrolidin-2-ylmethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 119); mp 195-198°C.

(ccccc-a) N-{4-[(R)-Pyrrolidin-2-ylmethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 120); mp 289-291°C.

15 (ddddd) N-{4-[(S)-Pyrrolidin-2-ylmethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 121); mp 138-141°C.

(eeeeee) N-[4-(2-Dimethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 122); mp 163-166°C.

20 (fffff) N-[4-(3-Dimethylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 123); mp 247°C.

(ggggg) N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 124)

(ggggg-a) N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 125); mp 188-245°C (d).

25 (hhhhh) N-[4-(2-Dimethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 126).

09387311.0 319

257
239

(iiii) N-{4-[2-(4-Methyl-piperazin-1-yl)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 127).

(iiii) N-{4-[2-Morpholin-1-ylethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 128).

5 (kkkkk) N-{4-[2-Piperidin-1-ylethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 129).

(kkkkk-a) N-{4-[2-Piperidin-1-ylethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride (Compound 130); mp 208-211°C.

10 (IIII) N-{4-[(1-Methyl-pyrrolidin-3-yl)methoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 131); mp 209-211°C.

(mmmmmm) N-{4-[(1-Ethyl-pyrrolidin-3-yl)methoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 132).

(nnnnnn) N-{4-[2-(1-Methyl-pyrrolidin-2-yl)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 133).

15 (ooooo) N-{4-[2-(1-Methyl-pyrrolidin-2-yl)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrate (Compound 134).

(ppppp) N-[4-(3-n-Propylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 135).

20 (qqqqq) N-[4-(3-Cyclopropylmethylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 136).

(rrrr) N-{4-[3-(2-Ethylbutyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 137).

(sssss) N-[4-(3-Cyclohexylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 138).

25 (ttttt) N-[4-(3-Cyclohexylmethylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 139).

252
240

(uuuuu) N-{4-[3-(Pyrid-4-ylmethyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 140).

(vvvvv) N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 141); mp 148-150°C.

5 (wwwww) N-[4-(3-Di-n-propylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 142).

(xxxxx) N-{4-[3-Di(c-propylmethyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 143).

(yyyyy) N-{4-[3-Di(2-ethylbutyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 144).

(zzzzz) N-{4-[3-Di(pyrid-4-ylmethyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 145).

(aaaaa) N-{4-[2-(2-Pyrrolidin-1-ylethoxy)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 146).

15 (bbbbbb) N-{4-[2-(2,2-Dimethylaminoethylamino)-2-oxoethyl]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 147).

(ccccc) N-{4-[2-(4-Methylaminopiperizin-1-yl)-2-oxoethyl]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 148); oil.

(ddddd) N-{4-[7-azabicyclo(2.2.1)hept-2-yloxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 149).

(eeeee) N-[3-(2-Diethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 150).

(fffff) N-[3-(2-Pyrrolidin-1-ylethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 151).

25 (ggggg) N-[3-(2-Di-Isopropylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 152).

253
241

(hhhhhh) N-[3-(2-n-Propylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 153).

(iiiiii) N-[3-(2-n-Butylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 154).

5 (iiiiii) N-[3-(Methylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 155).

(kkkkkk) N-[3-[3-(N-Ethyl,N-Methyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 156).

(llllll) N-[3-[3-(N-Cyclopropylmethyl,N-n-propyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 157).

(mmmmmm) N-[3-(Azeditinylpropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 158).

(nnnnnn) N-[3-(3-Ethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 159).

15 (oooooo) N-[3-[3-(2,2,2-Trifluoroethyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 160).

(pppppp) N-[3-(3-n-Propylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 161).

20 (rrrrr) N-[3-(3-Isopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 163).

(ssssss) N-[3-(3-Cyclopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 164).

(ttttt) N-[3-(3-Cyclopropylmethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 165).

25 (uuuuuu) N-[3-(3-Cyclobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 166).

25A
242

(vvvvvv) N-[3-(3-Cyclohexylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 167).

(wwwwww) N-[3-[3-(3-Ethylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 168).

5 (xxxxxx) N-[3-[3-(2-Methylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 169).

(yyyyyy) N-[3-(3-Isobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 170).

10 (zzzzzz) N-[3-(3-t-Butylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 171).

(aaaaaa) N-[3-[3-(2-Methylbutyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 172).

(bbbbbbb) N-[3-(3-Isoamylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 173).

15 (ccccc) N-[3-[3-(4-Methylpentyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 174).

(ddddddd) N-[3-[3-(1,1-Dimethylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 175).

(ccccccc) N-[3-[3-(3,3-Dimethylbutyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 176).

(ffffff) N-[3-[3-(2,4-Dimethylpent-3-yl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 177).

(ggggggg) N-[3-[3-(4-Methylcyclohexyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 178).

25 (hhhhhhh) N-[3-[3-(4-t-Butylcyclohexyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 179).

09387311-08319

255-1
243

(iiiiii) N-{3-[3-(2,6-Dimethylcyclohexyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 180).

(iiiiiij) N-{3-[3-(1-Phenylethyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 181).

5 (kkkkkkk) N-[3-(3-Norborn-2-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 182).

(lllllll) N-[3-(3-Adamant-1-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 183); mp 175-176°C.

10 (mmmmmmm) N-[3-(3-Norborn-2-ylmethyaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 184).

(nnnnnn) N-[3-(3-Adamant-2-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 185).

(ooooooo) N-[4-(2-Ethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 186).

15 (ooooooo-a) N-[4-(2-Ethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide hydrochloride (Compound 187); mp 227-228°C.

(ppppppp) N-[2-Fluoro-4-(2-Ethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 188).

20 (qqqqqqq) N-[4-(2-n-Propylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 189).

(rrrrr)N-[4-(2-Cyclopropylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 190).

(ssssss) N-4-(2-n-Butylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 191).

25 (ttttt) N-[4-(3-Ethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 192).

256
244

(uuuuuuu) N-{4-[3-(1-Phenyl-1-methylethyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 193).

(vvvvvvv) N-[4-(Pyrid-3-ylmethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 194); mp 241-243°C.

5 (wwwwwww) N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 195).

(wwwwwww-a) N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide hydrochloride (Compound 196); mp 235-240°C (d).

10 (xxxxxxx) N-[4-(2-Dimethylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 197).

(yyyyyyy) N-[4-(2-Diethylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 198).

15 (zzzzzzz) N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 199).

(zzzzzzz-a) N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide hydrochloride (Compound 200); mp 160-162°C.

(aaaaaaa) N-[4-(2-Piperidin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 201).

20 (bbbbbbbb) N-{4-[2-(1-Methyl-pyrrolidin-2-yl)ethoxy]pyrid-3-yl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 202).

(ccccccc) N-{4-[(1-Ethyl-pyrrolidin-3-yl)methoxy]pyrid-3-yl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 203); oil.

25 (ddddddd) N-[4-(2-Morpholin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 204).

09387311-08319
6TE80-YTE/8860

267
245

(eeeeeee) N-[4-(2-Diethylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 205).

(ffffff) N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 206).

5 (ffffff-a) N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide hydrochloride (Compound 207); mp 210°C.

(ggggggg) N-[4-(2-Isopropylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 208).

10 (hhhhhhh) N-[4-(3-Isopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 209).

(iiiiiii) N-[4-(3-Cyclopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 210).

(iiiiiii) N-[4-(3-Cyclobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 211).

15 (kkkkkkk) N-[4-(3-Cyclopropylmethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 212).

(lllllll) N-[4-(3-Isobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 213).

20 (mmmmmmm) N-[4-[3-(2,2-Dimethylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 214).

(nnnnnnn) N-[4-[3-(3-Ethylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 215).

(oooooooo) N-[4-[3-(2-Methylbutyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 216).

25 (ppppppp) N-[4-[3-(2-Methylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 217).

288
246

(qqqqqqqq) N-[4-(3-i-Pentylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 218).

(mmmm) N-[4-(3-Cyclohexylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 219).

5 (ssssss) N-{4-[3-(N-Cyclopropylmethyl,N-n-propyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 220).

(tttttt) N-[4-(3-Indan-2-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide (Compound 221).

10 (uuuuuuu) N-[3-Fluoro-4-(2-ethoxy-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 222); mp 192-196°C.

(vvvvvvvv) N-[3-Fluoro-4-(2-hydroxy-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 223); mp 246-248°C.

(wwwwwww) N-[3-Fluoro-4-(2-ethylamino-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 224).

15 (xxxxxxx) N-[3-Fluoro-4-(2-diethylamino-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 225); mp 193-196°C.

(yyyyyyyy) N-{3-Fluoro-4-[2-(4-methylpiperizin-1-yl)-2-oxoethoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide (Compound 226).

20 (zzzzzzzz) N-ethyl-N-[2-(ethylamino)ethyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 227).

(aaaaaaaa) N-[2-(dipropylamino)ethyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 228); mp 148-150°C.

(bbbbbbbb) N-[2-(diethylamino)ethyl]-N-methyl-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 229); mp 220-228°C.

09387311-031
FEB 0 1980

258
247

(cccccccc) N-[2-(diethylamino)ethyl]-N-ethyl-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 230); mp 165-167°C.

(dddddddd) N-[4-(2-morpholin-4-yl-2-oxoethoxy)phenyl](4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carboxamide (Compound 231).

5 (cccccccc) N-[3-fluoro-4-(2-morpholin-4-yl-2-oxoethoxy)phenyl](4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carboxamide (Compound 232); mp 110°C.

(ffffff) (4-oxo-(4,5,6,7-tetrahydroindol-3-yl))-N-[4-(2-oxo-2-piperazinyloxy)phenyl]carboxamide (Compound 233)

10 (gggggggg) N-[3-(diethylamino)propyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 234)

(hhhhhhhh) N-[3-(diethylamino)propyl]-2-{2-fluoro-4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 235).


(iiiiiii) N-[4-(diethylamino)-1-methylbutyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 236).

15 (jjjjjjj) N-[4-(diethylamino)-1-methylbutyl]-2-{2-fluoro-4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide (Compound 237).

Example 4

20 Water solubility for various compounds within the invention was determined and compared with that for compounds outside the scope of the invention. The compounds evaluated are encompassed within formula II:

260
248



II	R _x	R _y	n	R
<u>Water Solubility (μg/ml)</u>				
23	H	H	1	
203	H	H	1	
143	H	H	2	
15	H	H	1	
1.0	H	H	1	
0.58	H	H	1	
0.34	H	H	1	
0.26	CH ₃	CH ₃	1	

66780-TTE78E D 387311-083199

261
249


The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred
5 embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

0 387311.08319

262
250

WHAT IS CLAIMED IS:

1. A compound of the formula:



or the pharmaceutically acceptable non-toxic salts thereof wherein:

- 5 G represents



where

Q is phenyl, 2- or 3-thienyl, or 2-, 3-, or 4-pyridyl, all of which may be mono or disubstituted with hydroxy or halogen;

T is halogen, hydrogen, hydroxyl, amino or alkoxy having 1-6 carbon atoms;

W is oxygen, nitrogen, sulfur, or CR7R8 where R7 and R8 are the same or different and represent hydrogen, alkyl, or R7-R8 taken together represents a cyclic moiety having 3-7 carbon atoms;

X is hydrogen, hydroxyl, or alkyl;

15 Z is

hydroxy, C₁-C₆ cycloalkyl, C₁-C₆ alkoxy, or C₁-C₆ alkoxy where the alkyl portion is optionally substituted with amino, mono- or diC₁-C₆alkylamino, or azaC₁-C₆cycloalkyl;

20 amino, or mono- or diC₁-C₆alkylamino, (trifluoromethyl)methyl, C₇-C₇ cycloalkyl optionally substituted with C₁-C₆ alkyl, or C₇-C₇cycloalkylC₁-C₆alkyl;

0 38731-0 FFE/BE 0

263 |
251


phenyl or phenyl(C₁-C₆)alkyl where the phenyl portion is optionally substituted with C₁-C₆ alkyl, hydroxy, C₁-C₆ alkoxy, trifluoromethyl, halogen, amino, or mono- or diC₁-C₆ alkylamino; 2-, 3-, or 4-pyridyl, 1- or 2-imidazolyl, 1-, 2-, or 3-pyrrolyl, or adamantane-2-yl; each of which may be substituted on a tertiary carbon or a secondary nitrogen with C₁-C₆alkyl;

NR₉COR₁₀ where R₉ and R₁₀ are the same or different and represent hydrogen or C₁-C₆ alkyl or C₃-C₇ cycloalkyl; or

Z is connected, optionally through W, to Q to form a 1-6 membered ring; or

Z represents

a group of the formula:




where

p is 1, 2, or 3;

D and D' independently represent oxygen, NR₇ or CHR₇ provided that only one of D and D' may be NR₇ where each R₇ is hydrogen or C₁-C₆ alkyl; or and

R_z is hydrogen or C₁-C₆ alkyl;

a group of the formula:



where

p is 1, 2, or 3;

q is 0, 1, or 2;

0387311-083
6 FEB 80 11 18 AM '80


10

15

20

264-1
252

U. V. D. P.




where

5 **s is 0, 1, 2 or 3, and the sum of s and m is not less than 1;**

R_n is hydroxy, C₁-C₆alkoxy, amino, mono- or diC₁-C₆alkylamino

where each alkyl is independently optionally substituted with amino, mono- or diC₁₋₆alkylamino, or

R_n is a group of the formula



where p , D , D' , and R , are as defined above;

, , and  independently represent a carbon chain optionally substituted with hydrogen, halogen, or straight or branched chain lower alkyl having 1-6 carbon atoms;

15 **wherein**

k is 0, 1, 2, or 3;

m is 0, 1, 2, or 3; and

n is 0, 1, 2, or 3;

R₃, R₄, R₅, and R₆ are the same or different and are selected from hydrogen, alkyl, -

20 COR₁₁ or -CO₂R₁₁ where R₁₁ is alkyl or cycloalkyl having 3-7 carbon atoms;
or -CONR₁₂R₁₃ where R₁₂ and R₁₃ are selected independently from hydrogen,
alkyl, cycloalkyl having 3-7 carbon atoms, phenyl, 2-, 3-, or 4-pyridyl, or

09387311-083199

10

15

20

265
253


NR₁₂R₁₃ forms a heterocyclic group which is morpholinyl, piperidinyl, pyrrolidinyl, or N-alkyl piperazinyl; or

R₃-R₄ may be taken together to form a cyclic moiety having 3-7 carbon atoms; or

R₅-R₆ may be taken together to form a cyclic moiety having 3-7 carbon atoms; and

5 where each alkyl group forming an R₃, R₄, R₅, or R₆ substituent or portion thereof may be substituted independently with hydroxy or mono- or dialkylamino where each alkyl is independently alkyl or cycloalkyl having 3-7 carbon atoms.

10 2. A compound according to claim 1, which is:




wherein Y is hydrogen, halogen, or hydroxy.

0 10 387311-0831 9


266
254

3. A compound according to claim 1, which is:




wherein Y is hydrogen, halogen, or hydroxy.

- 5 4. A compound according to claim 1, which is:




wherein Y is hydrogen, halogen, or hydroxy.

5. A compound according to claim 1, which is:




10 wherein Y is hydrogen, halogen, or hydroxy.


6. A compound according to claim 1, which is:



9. A compound of the formula:




8. A compound of the formula:




7. A compound according to claim 1, which is:

wherein Y is hydrogen, halogen, or hydroxy.



208
256

10. A compound of the formula:



11. A compound according to claim 1, which is

N-[3-(Methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

N-[4-(Hydroxyethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

N-[4-(Methoxyethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

N-[4-(3-Methylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-[4-(Methoxymethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Aminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[2-Fluoro-4-(methylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-[4-[N-acetyl-(methylaminomethyl)phenyl]]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-[4-(Ethylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Isopropylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-[4-(Cyclopropylaminomethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

269
257

N-[4-(Dimethylaminomethyl)phenyl]4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-[4-(2-Aminoethyl)phenyl]4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

5 12. A compound according to claim 1, which is

N-[4-(2-Methylaminoethyl)phenyl]4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Methoxymethyl)phenyl]4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Methylaminomethyl)phenyl]4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-[N-acetyl-(methylaminomethyl)phenyl]]4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Methylaminomethyl)phenyl]4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Hydroxymethyl)phenyl]4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Hydroxyethoxy)phenyl]4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

N-[3-(Methylaminomethyl)phenyl]4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Hydroxyethoxyphenyl)4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[3-(Hydroxymethyl)phenyl]4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

N-[4-(Hydroxymethyl)phenyl]4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

210
258

N-[4-(Methylaminomethyl)phenyl]-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-(1,3-Benzodioxol-5-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

5 carboxamide;

N-(3,4-Dihydro-2H-1,4-benzoxazin-6-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-(2,2-Dimethyl-1,3-benzodioxol-5-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide; or

10 N-(2,3-Dihydro-1H-indol-5-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

13, A compound according to claim 1, which is

N-(2,3-Dihydro-1H-indol-6-yl)-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-(1,3-Benzodioxol-5-yl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-

15 carboxamide;

N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-

indole-3-carboxamide;

N-(4H-1,3-Benzodioxin-7-yl)-4-oxo-5,5-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

20 N-(1,3-Benzodioxol-5-yl)-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-

carboxamide;

N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-1,4,5,6,7,8-hexahydro-

cyclohepta[b]pyrrole-3-carboxamide;

N-(2,2-Dimethyl-1,3-benzodioxol-5-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-

25 3-carboxamide;

271
259

N-(1,3-Benzodioxol-5-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;


5 N-(1,3-Benzodioxol-5-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-(2,3-Dihydro-1,4-benzodioxin-6-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

10 N-(4H-1,3-Benzodioxin-7-yl)-4-oxo-6,6-dimethyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-[(2-Hydroxyethoxy)pyrid-5-yl]-4-oxo-6-methyl-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

14. A compound of the formula:



15 where

R₃, R₅, and R₆ independently represent hydrogen, or alkyl;


R_a represents hydrogen or alkyl where the alkyl is optionally halogenated; and

e is an integer of 1-3.

20

15. A compound of the formula:

272
260




where

R₃, R₅, and R₆ independently represent hydrogen, or alkyl;

R_a and R_b independently represent hydrogen or alkyl; and

e is an integer of 2-3.

16. A compound of the formula:



where

R₃, R₅, and R₆ independently represent hydrogen, or alkyl;

R_a represents hydrogen, alkyl, or C₃₋₇ cycloalkyl;


R_b represents hydrogen, alkyl, or acyl;

Y represents hydrogen or halogen; and

e is an integer of 1-3.

17. A compound of the formula:

233
261-




where

R₃, R₅, and R₆ independently represent hydrogen, or alkyl;


R_a represents hydrogen or alkyl where the alkyl is optionally halogenated; and

e is an integer of 1-3.

18. A compound of the formula:



where G represents



where R_a represents hydrogen, alkyl, or C₃-7 cycloalkyl;

R_b represents hydrogen, alkyl, or acyl;

Y and Y' independently represent hydrogen or halogen; and

e is an integer of 1-3.


19. A compound of the formula:

09387311-083199


10

15

23A
262



where G represents:



where Z is oxygen, nitrogen, or methylene; and m is 1 or 2.

20. A compound according to claim 1, which is
N-[3-(2-Pyrrolidinyloethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-Dimethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-n-Propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-n-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-Cyclobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-t-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;
N-[3-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

09387311.0831
TEBO TTE/8860

275
463

N-{3-[2-(4-Methylcyclohexyl)aminoethoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-{3-[2-(3-Trifluoromethylbenzylamino)ethoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

5

21. A compound according to claim 1, which is

N-{3-[3-(3-Trifluoromethylbenzylamino)propoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Dimethylaminoethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Pyrrolidin-1-ylethyl)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Diisopropylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Methylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Ethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[2-Fluoro-4-(2-ethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-n-Propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[2-Fluoro-4-(2-n-propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-[3-Fluoro-4-(2-n-propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

09387311-083199

276
264

22. A compound according to claim 1 which is

N-[3-Fluoro-4-(2-n-propylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

5 N-[4-(2-Cyclopropylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Isopropylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

10 N-[4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hemifumarate;

N-[2-Fluoro-4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

15 N-[3-Fluoro-4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[3-Fluoro-4-(2-Cyclopropylmethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide tosylate;

N-[4-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

20 N-[2-Fluoro-4-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

23. A compound according to claim 1, which is

25 N-[3-Fluoro-4-(2-Isobutylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

227.
265

N-[4-(2-n-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-n-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

5 N-[3-Fluoro-4-(2-n-butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-t-Butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

10 N-[3-Fluoro-4-(2-t-butylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-adamant-2-ylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-[(R)-Pyrrolidin-2-ylmethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

15 N-[4-[(S)-Pyrrolidin-2-ylmethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-[4-(Piperidin-3-ylmethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

20 24. A compound according to claim 1, which is

N-[4-(Piperidin-3-ylmethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

N-[4-(2-Dimethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

25 N-[3-Fluoro-4-(2-dimethylaminoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

278.
266

N-[4-(2-Pyrrolidin-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Imidaz-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[3-Fluoro-4-(2-morpholin-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

5 carboxamide;

N-[3-Fluoro-4-(2-pyrrolidin-1-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-[4-(2-Piperidin-2-ylethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

10 N-{4-[3-(2,2,2-Trifluoroethyl)aminopropoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-[4-(3-Isopropylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide.

15 25. A compound according to claim 1, which is

N-{4-[3-(2-Methylpropyl)aminopropoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

6 carboxamide;

N-[4-(3-Isobutylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

20 N-[4-(3-Cyclopropylmethylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[3-(3-Ethylpropyl)aminopropoxy]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

carboxamide;

N-[4-(3-Cyclopentylaminopropoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-

25 carboxamide;

0938731-0831
FEB 07 1988

228
267

N-[4-[3-(N-Cyclopropylmethyl,N-propyl)aminopropoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Methylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

5 N-[4-(2-Ethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Ethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride; or

10 N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

26. A compound according to claim 1, which is

N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

15 N-[4-(2-Isopropylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Isopropylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

20 N-[4-(2-n-Butylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-n-Butylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

N-[4-(2-t-Butylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

25 N-[4-(2-Benzylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

280
268

N-[4-(2-Benzylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

N-[4-(Pyrid-3-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

5 N-[4-(Pyrid-3-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride.

27. A compound according to claim 1, which is

10 N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

N-{4-[(R)-Pyrrolidin-2-ylmethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

15 N-{4-[(R)-Pyrrolidin-2-ylmethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

N-{4-[(S)-Pyrrolidin-2-ylmethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

20 N-[4-(2-Dimethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(3-Dimethylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

25 N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride; or

201
269

N-[4-(2-Dimethylaminoethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

28. A compound according to claim 1, which is

N-{4-[2-(4-Methyl-piperazin-1-yl)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[2-Morpholin-1-ylethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[2-Piperidin-1-ylethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[2-Piperidin-1-ylethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrochloride;

N-{4-[(1-Methyl-pyrrolidin-3-yl)methoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[(1-Ethyl-pyrrolidin-3-yl)methoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[2-(1-Methyl-pyrrolidin-2-yl)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[2-(1-Methyl-pyrrolidin-2-yl)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide hydrate;

N-[4-(3-n-Propylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

N-[4-(3-Cyclopropylmethylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

29. A compound according to claim 1, which is

264
270

N-{4-[3-(2-Ethylbutyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(3-Cyclohexylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

5 N-[4-(3-Cyclohexylmethylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[3-(Pyrid-4-ylmethyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

10 N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[4-(3-Di-n-propylaminopropoxy)pyrid-3-yl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[3-Di(cyclopropylmethyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

15 N-{4-[3-Di(2-ethylbutyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-{4-[3-Di(pyrid-4-ylmethyl)aminopropoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

20 N-{4-[2-(2-Pyrrolidin-1-ylethoxy)ethoxy]pyrid-3-yl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

30. A compound according to claim 1, which is

N-{4-[2-(2,2-Dimethylaminoethylamino)-2-oxoethyl]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

25 N-{4-[2-(4-Methylaminopiperizin-1yl)-2-oxoethyl]phenyl}-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

09387311-083188

2831.
271

N-{4-[7-azabicyclo(2.2.1)hept-2-yloxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(2-Diethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

5 N-[3-(2-Pyrrolidin-1-ylethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(2-Di-i-propylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

10 N-[3-(2-n-Propylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(2-n-Butylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(Methylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide; or

15 N-[3-[3-(N-Ethyl,N-Methyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide.

31. A compound according to claim 1, which is

20 N-{3-[3-(N-Cyclopropylmethyl,N-n-propyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(Azeditinylpropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-Ethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

25 N-[3-[3-(2,2,2-Trifluoroethyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

284
272

N-[3-(3-n-Propylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-Isopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

5 N-[3-(3-Cyclopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-Cyclopropylmethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide; or

10 N-[3-(3-Cyclobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide.

32. A compound according to claim 1, which is

N-[3-(3-Cyclohexylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

15 N-[3-(3-(3-Ethylpropyl)aminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

6 N-[3-(3-(2-Methylpropyl)aminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

20 N-[3-(3-Isobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-t-Butylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-(2-Methylbutyl)aminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

25 N-[3-(3-Isoamylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

09387311-0831 6

285
273

N-{3-[3-(4-Methylpentyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-{3-[3-(1,1-Dimethylpropyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide; or

5 N-{3-[3-(3,3-Dimethylbutyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide.

33. A compound according to claim 1, which is

10 N-{3-[3-(2,4-Dimethylpent-3-yl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-{3-[3-(4-Methylcyclohexyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-{3-[3-(4-t-Butylcyclohexyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

15 N-{3-[3-(2,6-Dimethylcyclohexyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-{3-[3-(1-Phenylethyl)aminopropoxy]phenyl}-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

20 N-[3-(3-Norborn-2-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-Adamant-1-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[3-(3-Norborn-2-ylmethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

25 N-[3-(3-Adamant-2-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide; or

286
274

N-[4-(2-Ethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide.

34. A compound according to claim 1, which is

N-[4-(2-Ethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide hydrochloride;

N-[2-Fluoro-4-(2-Ethylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(2-n-Propylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(2-Cyclopropylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(2-n-Butylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(3-Ethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-[3-(1-Phenyl-1-methylethyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(Pyrid-3-ylmethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide; or

N-[4-(Pyrid-4-ylmethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide hydrochloride.

35. A compound according to claim 1, which is

204
275

N-[4-(2-Dimethylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(2-Diethylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

5 N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(2-Pyrrolidin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide hydrochloride;

10 N-[4-(2-Piperidin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-[2-(1-Methyl-pyrrolidin-2-yl)ethoxy]pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-[(1-Ethyl-pyrrolidin-3-yl)methoxy]pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

15 N-[4-(2-Morpholin-1-ylethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(2-Diethylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide; or

20 N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide.

36. A compound according to claim 1, which is

N-[4-(2-n-Propylaminoethoxy)pyrid-3-yl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide hydrochloride;

25 N-[4-(2-Isopropylaminoethoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

286
276

N-[4-(3-Isopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(3-Cyclopropylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

5 N-[4-(3-Cyclobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(3-Cyclopropylmethylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

10 N-[4-(3-Isobutylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-[3-(2,2-Dimethylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-[3-(3-Ethylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide; or

15 N-[4-[3-(2-Methylbutyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide.

37. A compound according to claim 1, which is

20 N-[4-[3-(2-Methylpropyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(3-i-Pentylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

N-[4-(3-Cyclohexylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-
cyclohepta[b]pyrrole-3-carboxamide;

25 N-[4-[3-(N-Cyclopropylmethyl,N-n-propyl)aminopropoxy]phenyl]-4-oxo-1,4,5,6,7,8-
hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

0 387311.0 319
6TE

209
277

N-[4-(3-Indan-2-ylaminopropoxy)phenyl]-4-oxo-1,4,5,6,7,8-hexahydro-cyclohepta[b]pyrrole-3-carboxamide;

N-[3-Fluoro-4-(2-ethoxy-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

5 N-[3-Fluoro-4-(2-hydroxy-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

N-[3-Fluoro-4-(2-ethylamino-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide;

~ N-[3-Fluoro-4-(2-diethylamino-2-oxoethoxy)phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide; or

10 N-[3-Fluoro-4-[2-(4-methylpiperizin-1-yl)-2-oxoethoxy]phenyl]-4-oxo-4,5,6,7-tetrahydro-1H-indole-3-carboxamide.

38. A compound according to claim 1, which is

15 N-ethyl-N-[2-(ethylamino)ethyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide;

N-[2-(dipropylamino)ethyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide;

20 N-[2-(diethylamino)ethyl]-N-methyl-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide;

N-[2-(diethylamino)ethyl]-N-ethyl-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide;

N-[4-(2-morpholin-4-yl-2-oxoethoxy)phenyl](4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carboxamide;

25 N-[3-fluoro-4-(2-morpholin-4-yl-2-oxoethoxy)phenyl](4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carboxamide;

290
279

(4-oxo-(4,5,6,7-trihydroindol-3-yl))-N-[4-(2-oxo-2-piperazinyloxy)phenyl]carboxamide;


N-[3-(diethylamino)propyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide;

5 N-[3-(diethylamino)propyl]-2-{2-fluoro-4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide;

N-[4-(diethylamino)-1-methylbutyl]-2-{4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide; or

10 N-[4-(diethylamino)-1-methylbutyl]-2-{2-fluoro-4-[(4-oxo-(4,5,6,7-tetrahydroindol-3-yl))carbonylamino]phenoxy}acetamide.

39. A compound of the formula:



or the pharmaceutically acceptable non-toxic salts thereof wherein:

15 Q is phenyl or 3-pyridyl, each of which may be mono or disubstituted with hydroxy or halogen;


W is oxygen or nitrogen;

Z is

20 hydroxy, C₁-C₇ cycloalkyl, C₁-C₆ alkoxy, or C₁-C₆ alkoxy where the alkyl portion is optionally substituted with amino, mono- or diC₁-C₆alkylamino, or azaC₁-C₆cycloalkyl;

0387311.0860 FEB 01 11 08 60

292
280




p is 1, 2, or 3;

q is 0, 1, or 2;

R₂ is hydrogen or C₁-C₆ alkyl; or

a group of the formula:




where

s is 0, 1, 2 or 3, and the sum of s and m is not less than 1;

R₆ is hydroxy, C₁-C₆alkoxy, amino, mono- or diC₁-C₆alkylamino where each alkyl is independently optionally substituted with amino, mono- or diC₁-C₆alkylamino, or

R₆ is a group of the formula



where p, D, D', and R₆ are as defined above;

, , and independently represent a carbon chain optionally substituted with hydrogen, halogen, or straight or branched chain lower alkyl having 1-6 carbon atoms;

wherein

k is 0, 1, 2, or 3;

m is 0, 1, 2, or 3; and

n is 1 or 2; and

6 FEB 1986

5

10

15

20

2921
281


R₃, R₄, R₅, and R₆ are the same or different and are selected from hydrogen, alkyl, -
COR₁₁ or -CO₂R₁₁ where R₁₁ is alkyl or cycloalkyl having 3-7 carbon atoms;
or -CONR₁₂R₁₃ where R₁₂ and R₁₃ are selected independently from hydrogen,
alkyl, cycloalkyl having 3-7 carbon atoms, phenyl, 2-, 3-, or 4-pyridyl, or
NR₁₂R₁₃ forms a heterocyclic group which is morpholinyl, piperidinyl,
pyrrolidinyl, or N-alkyl piperazinyl; or

R₃-R₄ may be taken together to form a cyclic moiety having 3-7 carbon atoms; or

R₅-R₆ may be taken together to form a cyclic moiety having 3-7 carbon atoms; and

where each alkyl group forming an R₃, R₄, R₅, or R₆ substituent or portion thereof may be substituted independently with hydroxy or mono- or dialkylamino where each alkyl is independently alkyl or cycloalkyl having 3-7 carbon atoms.

40. A compound of formula A or formula B:




or a pharmaceutically acceptable salt thereof wherein

R₃, R₄, R₅, and R₆ are the same or different and are selected from hydrogen, alkyl, -
COR₁₁ or -CO₂R₁₁ where R₁₁ is alkyl or cycloalkyl having 3-7 carbon atoms;
or -CONR₁₂R₁₃ where R₁₂ and R₁₃ are selected independently from hydrogen,
alkyl, cycloalkyl having 3-7 carbon atoms, phenyl, 2-, 3-, or 4-pyridyl, or
NR₁₂R₁₃ forms a heterocyclic group which is morpholinyl, piperidinyl,
pyrrolidinyl, or N-alkyl piperazinyl; or

281-1
282

R₃-R₄ may be taken together to form a cyclic moiety having 3-7 carbon atoms; or
 R₅-R₆ may be taken together to form a cyclic moiety having 3-7 carbon atoms; and
 where each alkyl group forming an R₃, R₄, R₅, or R₆ substituent or portion thereof may
 be substituted independently with hydroxy or mono- or dialkylamino where each
 alkyl is independently alkyl or cycloalkyl having 3-7 carbon atoms; and

G represents




where

R_h is hydrogen, halogen, C₁-C₆alkyl, C₁-C₆alkoxy, or trifluoromethyl;

s is 0, 1, 2 or 3, and the sum of s and m is not less than 1;

R_g is hydroxy, C₁-C₆alkoxy, amino, mono- or diC₁-C₆alkylamino
 where each alkyl is independently optionally substituted with
 amino, mono- or diC₁-C₆alkylamino, or

R_g is a group of the formula



p is 1, 2, or 3;


D and D' independently represent oxygen, NR_y, or CHR_y, provided
 that only one of D and D' may be NR_y, where each R_y is
 hydrogen or C₁-C₆ alkyl; or and

R_y is hydrogen or C₁-C₆ alkyl.

00367311-08319

215
283


41. A compound according to claim 40, wherein R_2 is hydrogen or halogen, and R_3 is a group of the formula:



where

- 5 R_{14} is hydrogen or C_1-C_6 alkyl;
 R_{15} is hydrogen or C_1-C_6 alkyl;
 R_{16} is hydrogen, ethyl, or methyl;
 R_{17} is C_1-C_6 alkyl; and
 J is a C_1-C_4 alkylene group.


42. A compound according to claim 40, wherein s is 1 and R_4 is ethoxy, hydroxy, ethylamino, diethylamino, morpholinyl, piperazinyl, 4-methylpiperazinyl,



2960
284

Abstract of the Disclosure

Disclosed are compounds of the formula:




or the pharmaceutically acceptable non-toxic salts thereof wherein:

5 G, X, T, n, and R₃-R₆ are as defined herein,

which compounds are highly selective agonists, antagonists or inverse agonists for GABA_A brain receptors or prodrugs of agonists, antagonists or inverse agonists for GABA_A brain receptors. These compounds are useful in the diagnosis and treatment of anxiety, sleep and seizure disorders, overdose with benzodiazepine drugs and for enhancement of memory.

09387311-0831
TEBO-TTE8E60

298
285



LOS ESTADOS UNIDOS DE AMÉRICA

A TODO AQUEL A QUIEN CONCIERNA LO PRESENTE

DEPARTAMENTO DE COMERCIO DE LOS ESTADOS UNIDOS


Oficina de Patentes y Marcas de los Estados Unidos

27 de Septiembre de 2000

LA PRESENTE ES PARA CERTIFICAR QUE ADJUNTA A LA MISMA HAY UNA COPIA FIEL DE LOS REGISTROS DE LA OFICINA DE PATENTES Y MARCAS DE LOS ESTADOS UNIDOS DE LOS DOCUMENTOS DE LA SOLICITUD DE PATENTE IDENTIFICADA EN LO QUE SIGUE, QUE CUMPLEN LOS REQUISITOS PARA OTORGARLE UNA FECHA DE PRESENTACIÓN BAJO 35 USC 111.

NÚMERO DE SOLICITUD: 09/387.311

FECHA DE PRESENTACIÓN: 31 de Agosto de 1999




**Por la Autoridad del
COMISARIO DE PATENTES Y MARCAS**

L. EDELEN

Oficial certificador

REIVINDICACIONES

1. Un compuesto de la fórmula:



o una sal no tóxica farmacéuticamente aceptable del mismo en donde:

G representa



en donde

Q es un fenilo, 2- o 3- tienilo, o 2-, 3-, o 4-piridilo, todos de los cuales pueden ser mono o disustituidos con hidroxilo o halógeno;

T es halógeno, hidrógeno, hidroxilo, amino o alcoxi que tienen 1-6 átomos de carbono;

W es oxígeno, nitrógeno, sulfuro, o CR₂R₂ donde R₁ y R₂ son lo mismo o diferente y representan hidrógeno, alquilo, o R₁-R₂ los cuales juntos representan un mosaico cíclico que tienen 3-7 átomos de carbono;

X es hidrógeno, hidroxilo o alquilo;

Z es

hidroxilo, C₁-C₆ cicloalquiloC₁-C₆(alcoxi), o C₁-C₆ alcoxi donde la porción alquilo es sustituida opcionalmente con amino, mono- o diC₁-C₆alquiloamino, o azaC₁-C₆cicloalquilo;

amino o mono- o di C₁-C₆alquilamino, (trifluorometil)metilo, C₁-C₆ cicloalquilo sustituido opcionalmente con C₁-C₆alquilo o C₁-C₆ cicloalquilC₁-C₆ alquilo;


fenilo o fenil(C₁-C₆)alquilo donde la porción de fenilo es opcionalmente sustituida con C₁-C₆ alquilo, hidroxí, C₁-C₆ alcoxi, trifluorometil, halógeno, amino, o mono- o di C₁-C₆ alquilamino;

2-, 3-, o 4-piridilo, 1- o 2-imidazolilo, 1-, 2-, ó 3-pirrolilo, o adamantan-2-ilo; cada uno de los cuales se puede sustutir en un carbono térciario o un nitrógeno secundario con C₁-C₆alquilo;

NR₉COR₁₀ en donde R₉ y R₁₀ son iguales o diferentes y representan hidrógeno o C₁-C₆ alquilo o C₁-C₆ cicloalquilo, o

Z está conectado, opcionalmente mediante W, a Q para formar un anillo de 1-6 miembros; o

Z representa un grupo de la fórmula:




en donde

p es 1, 2, o 3;

D y D' independientemente representan oxígeno, NR_y o CHR_y siempre que sólo D o D' puede ser NR_y en donde cada R_y es hidrógeno o C₁-C₆ alquilo; o/y R_z is hidrógeno o C₁-C₆ alquilo;

Z representa un grupo de la fórmula:

en donde



201
288


R₁ es hidrógeno o C₁-C₄ alquilo; o

$$\text{---} \left(\text{---} \text{CH}_2 \text{---} \text{C}(=\text{O}) \text{---} \text{R}_0 \right)_n \text{---}$$

en donde

R₆ es hidroxi, C₁-C₆alcoxi, amino, mono- o di-C₁-C₆-alquilamino,

R_0 es un grupo de la fórmula


$$\{ \}_{k}, \{ \}_{m}, \text{ y. } \{ \}_{n}$$

en donde

м е а 0, 1, 2, 6 3; у

n es 0,1,2 o 3;


R₃, R₄, R₅, y R₆ son iguales o diferentes y son independientemente seleccionados en cada caso desde hidrógeno, alquilo, -COR₁₁ o -CO₂R₁₁ en donde R₁₁ es alquilo o cicloalquilo que tiene 3-7 átomos de carbono; o -CONR₁₂R₁₃ en donde R₁₂ y R₁₃ se seleccionan independientemente de hidrógeno, alquilo, cicloalquilo que tiene 3-7 átomos de carbono, fenilo, 2-, 3-, ó 4-piridilo, o NR₁₂R₁₃ forma un grupo heterocíclico de morfolinilo, piperidinilo, pirrolidinilo, o N-alquilo piperazinilo; o

R₃-R₄ se pueden tomar juntos para formar una mitad cíclica que tiene 3-7 átomos de carbono; o

R₅-R₆ se pueden tomar juntos para formar una mitad cíclica que tiene 3-7 átomos de carbono;


en donde cada grupo de alquilo que forma R₃, R₄, R₅, o R₆ sustituyente o mitad del mismo se puede sustituir independientemente con hidroxí o mono- or dialquilamino en donde cada alquilo es independientemente alquilo o cicloalquilo que tiene 3-7 átomos de carbono.

2. Un compuesto con respecto a la reivindicación 1, de fórmula:




donde Y es hidrógeno, halógeno, o hidroxí.

3. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:




en donde Y es hidrógeno, halógeno o hidroxilo.

4. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:




en donde Y es hidrógeno, halógeno o hidroxilo.

5. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:




en donde Y es hidrógeno, halógeno o hidroxilo.


6. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:




7. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:




8. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:



9. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:



10. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:



11. Un compuesto de acuerdo con la Reivindicación 1, el cual es,

N-[3-(Metilaminometil) Fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Hidroxietoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Metoxietoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(3-metilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Metoximetil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Aminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Metilaminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[2-Fluoro-4-(metilaminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[N-acetil-(metilaminometil) fenil]}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(etilaminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(isopropilaminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(ciclopropilaminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(dimetilaminometil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-aminoetil) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

12. Un compuesto de acuerdo con la Reivindicación 1, el cual es,

N-[4-(2-Metilaminoetil) Fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Metoximetil) fenil]-4-oxo-5,5-dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Metilaminometil) fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{4-[N-acetil-(metilaminometil) fenil]}-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Metilaminometil) fenil]-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(hidroximetil) fenil]-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2hidroxietoxi) fenil]-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-(metilaminometil) fenil]-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-(2-hidroxietoxifenil)}-4-oxo-6,6-dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-(hidroximetil) fenil]-4-oxo-6,6 dimetil-4,5,6,7 -tetrahidro-1H-indole-3-carboxamida;

N-[4-(hidroximetil) fenil]-4-oxo-6,6-dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(metilaminometil) fenil]-4-oxo-6,6dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(1,3-benzodioxol-5-yl)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

N-(2,3-dihidro-1,4-benzodioxin-6-yl)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

N-(3,4-dihidro-2H-1,4-benzoxazina-6-yl)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

N-(2,2-dimetil-1,3-benzodioxin-5-yl)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

N-(2,3-dihidro-1H-indol-5-yl)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

13. Un compuesto de acuerdo con la Reivindicación 1, el cual es,

N-(2,3-dihidro-1H-indol-6-yl)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(1,3-benzodioxol-5-yl)-4-oxo-5,5-dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(2,3-dihidro-1,4-benzodioxin-6-yl)-4-oxo-5,5-dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(4H-1,3-benzodioxin-7-yl)-4-oxo-5,5-dimetil-4,5,6,7-tetrahidro-1H-indole-3-carboxamida;

N-(1,3-benzodioxol-5-yl)-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-(2,3,-dihidro-1,4-benzodioxin-6-yl)-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-(2,2-dimetil-1,3-benzodioxol-5-yl)-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(1,3-benzodioxol-5-yl)-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(2,3-dihidro-1,4-benzodioxin-6-yl)-4-oxo-6-metil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-(1,3-benzodioxol-5-yl)-4-oxo-6,6-dimetil-4,5,6,7-tetrahidro-1H-indole-3-carboxamida;

N-(2,3-dihidro-1,4-benzodioxin-6-yl)-4-oxo-6,6 dimetil-4,5,6,7 - tetrahidro-1H-indol-3-carboxamida;

N-(4H-1,3-benzodioxin-7-yl)-4-oxo-6,6-dimetil-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[(2-hidroxi)etoxi]pirido-5-yl]-4-oxo-4,5,6,7-tetrahidro-1H-indole-3-carboxamida;

14. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:

en donde


R₃, R₅, y R₆ representan independientemente hidrógeno, o alquilo;

R_a representa hidrógeno o alquilo en donde el alquilo es halogenado opcionalmente; y

e es un número entero de 1-3.

~~310~~
297


15. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:



donde

R_3 , R_5 , y R_6 representan independientemente hidrógeno o alquilo;
 R_a y R_b representan independientemente hidrógeno o alquilo; y
 e es un número completo entre 2-3.

16. Un compuesto de la fórmula:




donde

R_3 , R_5 , y R_6 representan independientemente hidrógeno o alquilo;
 R_a representa hidrógeno, alquilo, o C_1 -, cicloalquilo;
 R_b representa hidrógeno, alquilo o acilo;
 Y representa hidrógeno o halógeno; y
 e es un número completo de 1-3.

17. Un compuesto de la fórmula:

211
298




donde

R3, R5 y R6 representan independientemente hidrógeno o alquilo;


Ra representa hidrógeno o alquilo donde el alquilo es opcionalmente halogenado; y

e es un número completo de 1-3.

18. Un compuesto de la fórmula:




donde G representa




donde Ra representa hidrógeno, alquilo, o C₃-,cicloalquilo;
Rb representa hidrógeno, alquilo o acilo;
Y y Y' representan independientemente hidrógeno o halógeno; y
e es un número completo de 1-3.

19. Un compuesto de acuerdo con la Reivindicación 1, de la fórmula:



en donde G representa:



en donde Z es oxígeno, nitrógeno, o metileno; y m es 1 ó 2.

20. Un compuesto de acuerdo con la Reivindicación 1, el cual es
N-[3-(2-pirrolidinetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-
carboxamida;
N-[3-(2-Dimetilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-
3-carboxamida;
N-[3-(2-Propilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-
3-carboxamida;
N-[3-(2-n-butilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-
3-carboxamida;

300

N-[3-(2-isobutilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-(2-ciclobutilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-(2-butylaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-(2-n-ciclopropilmetilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-[2-(4-metilciclohexil)aminoetoxi]fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; o

N-[3-[2-(3-trifluorometilbenzilamino)etoxi]fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

21. Un compuesto de acuerdo con la Reivindicación 1, el cual es N-[3-[3-(3-Trifluorometilbencilamino)propoxi]fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Dimetilaminoetil)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Pirrolidin-1-ilet)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Diisopropilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Metilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Etilaminoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

34
307

N-[2-Fluoro-4-(2-etilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol carboxamida;

N-[4-(2-n-Propilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[2-Fluoro-4-(2-n-propilaminoetox) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; o

N-[3-Fluoro-4-(2-n-propilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

22. Un compuesto de acuerdo con la Reivindicación 1, la cual es N-[3-Fluoro-4-(2-n-propilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-[4-(2-Ciclopropilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Isopropilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Ciclopropilmetilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Ciclopropilmetilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida hemifumarato;

N-[2-Fluoro-4-(2-Ciclopropilmetilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-Fluoro-4-(2-Ciclopropilmetilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-Fluoro-4-(2-Ciclopropilmetilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida tosilato;

N-[4-(2-Isobutilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; o

N-[2-Fluoro-4-(2-Isobutilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

23. Un compuesto de acuerdo con la Reivindicación 1, la cual es N-[3-Fluoro-4-(2-Isobutilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-n-Butilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-n-Butilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-[3-Fluoro-4-(2-n-butilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-t-Butilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-Fluoro-4-(2-t-butilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-adamant-2-ilaminoetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[(R)-Pirrolidin-2-ilmetoxi] fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[(S)-Pirrolidin-2-ilmetoxi] fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; or

N-[4-(Piperidin-3-ilmetoxi) fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

24. Un compuesto de acuerdo con la Reivindicación 1, la cual es
N-[4-(Piperidin-3-iletoksi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-
indol-3-clorhidrato de carboxamida;

N-[4-(2-Dimetilaminoetoksi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-
3-carboxamida;

N-[3-Fluoro-4-(2-dimetilaminoetoksi)fenil]-4-oxo-4,5,6,7-tetrahidro-
1H-indol-3-carboxamida;

N-[4-(2-Pirrolidin-1-iletoksi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-
indol-3-carboxamida;

N-[4-(2-Imidaz-1-iletoksi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-
3-carboxamida;

N-[3-Fluoro-4-(2-morofolin-1-iletoksi)fenil]-4-oxo-4,5,6,7-
tetrahidro-1H-indol-3-carboxamida;

N-[3-Fluoro-4-(2-pirrolidin-1-iletoksi)fenil]-4-oxo-,5,6,7-
tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Piperidin-2-iletoksi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-
indol-3-carboxamida;

N-{4-[3-(2,2,2,-Trifluoretil)aminopropoxi]fenil}-4-oxo-4,5,6,7-
tetrahidro-1H-indol-3-carboxamida; o

N-[4-(3-Isopropilaminopropoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-
indol-3-carboxamida.

25. Un compuesto de acuerdo con la Reivindicación 1, la cual es
N-{4-[3-(2-Metilpropil)aminopropoxi]fenil}-4-oxo-4,5,6,7-
tetrahidro-1H-indol-3-carboxamida;

N-[4-(3-Isobutilaminopropoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-
indol-3-carboxamida;

N-[4-(3-Ciclopropilmetilaminopropoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[3-(3-Etilpropil)aminopropoxi]fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(3-Ciclopentilaminopropoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[3-(N-Ciclopropilmetil,N-propil)aminopropoxi]fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Metilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Etilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Etilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida; o

N-[4-(2-n-Propilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

26. Un compuesto de acuerdo con la Reivindicación 1, el cual es N-[4-(2-n-Propilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-[4-(2-Isopropilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Isopropilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-[4-(2-n-Butilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-n-Butilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-[4-(2-t-Butilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Bencilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Bencilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-[4-(Pirid-3-ilmetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; o

N-[4-(Pirid-3-ilmetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida.

27. Un compuesto de acuerdo con la Reivindicación 1, el cual es

N-[4-(Pirid-4-ilmetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(Pirid-4-ilmetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-{4-[(R)-Pirrolidin-2-ilmetoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[(R)-Pirrolidin-2-ilmetoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-{4-[(S)-Pirrolidin-2-ilmetoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Dimetilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(3-Dimetilaminopropoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Pirrolidin-1-iletoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Pirrolidin-1-iletoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida; o

N-[4-(2-Dimetilaminoetoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

28. Un compuesto de acuerdo con la Reivindicación 1, la cual es N-{4-[2-(4-Metil-piperazin-1-ilo)etoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[2-Morfolin-1-iletoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[2-Piperidin-1-iletoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[2-Piperidin-1-iletoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-clorhidrato de carboxamida;

N-{4-[(1-Metil-pirrolidin-3-ilo)metoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[(1-Etil-pirrolidin-3-ilo)metoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[2-(1-Metil-pirrolidin-2-ilo)etoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[2-(1-Metil-pirrolidin-2-ilo)etoxi]pirid-3-ilo}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-hidrato de carboxamida;

270
307

N-[4-(3-n-Propilaminopropoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; o

N-[4-(3-Ciclopropilmetilaminopropoxi)pirid-3-ilo]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

29. Compuesto según la Reivindicación 1, que es

N-{4-[3-(2-etilbutil)aminopropoxi]pirid-3-il}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(3-ciclohexilaminopropoxi)pirid-3-il]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(3-ciclohexilmetilaminopropoxi)pirid-3-il]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[3-(pirid-4-ilmetil)aminopropoxi]pirid-3-yl}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[4-(2-Pirrolidin-1-iletoksi)pirid-3-yl]-4-oxo-4,5,6,7-tetrahidro-1H-indole-3-carboxamida;

N-[4-(3-di-n-propilaminopropoxi)pirid-3-il]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[3-di(ciclopropilmetil)aminopropoxi]pirid-3-il}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[3-di(2-ethylbutil)aminopropoxi]pirid-3-il}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[3-di(pirid-4-ilmetil)aminopropoxi]pirid-3-il}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; or

N-[4-[2-(2-pirrolidin-1-iletoksi)etoxi]pirid-3-il]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

371-
308

30. Compuesto según la Reivindicación, 1 que es

N-{4-[2-(2,2-dimetilaminoetilamino)-2-oxoetil]fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[2-(4-metilaminopiperizin-1il)-2-oxoetil]fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-{4-[7-azabicyclo(2.2.1)hept-2-iloxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(2-dietilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(2-pirrolidin-1-iletoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(2-di-i-propilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(2-n-propilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(2-n-butilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(metilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida; or

N-{3-[3-(N-etil,N-metil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida.

31. Compuesto según la Reivindicación 1, que es

N-{3-[3-(N-ciclopropilmetil,N-n-propil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(azeditinolpropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-etilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(2,2,2-trifluoroetil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-n-propilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-isopropilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-ciclopropilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-ciclopropilmetilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida; o

N-[3-(3-ciclobutilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida.

32. Compuesto según la Reivindicación 1, que es

N-[3-(3-ciclohexilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(3-etilpropil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(2-metilpropil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-isobutilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-t-butilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

323.
310

N-{3-[3-(2-metilbutil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-Isoamilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(4-metilpentil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(1,1-dimetilpropil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida; o

N-{3-[3-(3,3,-dimetilbutil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida.

33. Compuesto según la Reivindicación 1, que es

N-{3-[3-(2,4-dimetilpent-3-il)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(4-metilciclohexil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(4-t-butilciclohexil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(2,6-dimetilciclohexil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{3-[3-(1-feniletil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-norborn-2-ilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-adamant-1-ilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-norborn-2-ilmetilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-(3-adamant-2-ilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida; o

N-[4-(2-etilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida.

34. Compuesto según la Reivindicación 1, que es

N-[4-(2-etilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida clorhidrato;

N-[2-fluoro-4-(2-etilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-n-propilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-ciclopropilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-4-(2-n-butilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-etilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{4-[3-(1-fenil-1-metiletil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(pirid-3-ilmetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(pirid-4-ilmetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida; o

N-[4-(pirid-4-ilmetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida clorhidrato.

35. Compuesto según la Reivindicación 1, que es

N-[4-(2-dimetilaminoetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-dietilaminoetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-pirrolidin-1-ilethoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrole-3-carboxamida;

N-[4-(2-Pirrolidin-1-iletoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrole-3-carboxamida clorhidrato;

N-[4-(2-piperidin-1-iletoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-(1-metil-pirrolidin-2-il)etoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-[(1-etil-pirrolidin-3-il)metoxi]pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-morfolin-1-iletoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(2-dietilaminoetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrole-3-carboxamida; o

N-[4-(2-n-propilaminoetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida.

36. Compuesto según la Reivindicación 1, que es

N-[4-(2-n-propilaminoetoxi)pirid-3-il]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida clorhidrato;

N-[4-(2-isopropilaminoetoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-isopropilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrole-3-carboxamida;

N-[4-(3-ciclopropilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-ciclobutilaminopropox)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-ciclopropilmetilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-isobutilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{4-[3-(2,2-dimetilpropil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{4-[3-(3-etilpropil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida; o

N-{4-[3-(2-metilbutil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrole-3-carboxamida.

37. Compuesto según la Reivindicación 1, que es

N-{4-[3-(2-metilpropil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-i-pentilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

322
314

N-[4-(3-ciclohexilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-{4-[3-(N-ciclopropilmetil, N-n-propil)aminopropoxi]fenil}-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[4-(3-indan-2-ilaminopropoxi)fenil]-4-oxo-1,4,5,6,7,8-hexahidro-ciclohepta[b]pirrol-3-carboxamida;

N-[3-fluoro-4-(2-etoxi-2-oxoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-fluoro-4-(2-hidroxi-2-oxoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-fluoro-4-(2-etilamino-2-oxoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

N-[3-fluoro-4-(2-dietilamino-2-oxoetoxi)fenil]-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida; o

N-{3-fluoro-4-[2-(4-metilpiperizin-1-il)-2-oxoetoxi]fenil}-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida.

38. Compuesto según la Reivindicación 1, que es

N-etil-N-[2-(etilamino)etil]-2-{4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-[2-(dipropilamino)etil]-2-{4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-[2-(dietilamino)etil]-N-metil-2-{4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-[2-(dietilamino)etil]-N-etil-2-{4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

328
315

N-[4-(2-morfolin-4-il-2-oxoetoxi)fenil](4-oxo-(4,5,6,7-tetrahidroindol-3-il))carboxamida;

N-[3-fluoro-4-(2-morfolin-4-il-2-oxoetoxi)fenil](4-oxo-(4,5,6,7-tetrahidroindol-3-il))carboxamida;

(4-oxo-(4,5,6,7-trihidroindol-3-il))-N-[4-(2-oxo-2-piperaziniletoxi)fenil]carboxamida;

N-[3-(dietilamino)propil]-2-{4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-[3-(dietilamino)propil]-2-{2-fluoro-4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-[4-(dietilamino)-1-metilbutil]-2-{4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-[4-(dietilamino)-1-metilbutil]-2-{2-fluoro-4-[(4-oxo-(4,5,6,7-tetrahidroindol-3-il))carbonilamino]fenoxi}acetamida;

N-(2-{[2-(N-metilacetamida)etil]amino}pirid-5-il)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida

N-(2-etoxi-4-metilpirid-5-il)-4-oxo-4,5,6,7-tetrahidro-1H-indol-3-carboxamida;

4-oxo-4,5,6,7-tetrahidro-1H-indol-3-ácido carboxílico {4-[2-(tiofeno-2-sulfonilamino)-etoxi]-fenil}-amida;


4-oxo-4,5,6,7-tetrahidro-1H-indol-3-ácido carboxílico (4-[1,2,4-triazol-1-il-fenil])-amida;

4-oxo-4,5,6,7-tetrahidro-1H-indol-3-ácido carboxílico {4-[3-(1-metil-1H-imidazol-4-sulfonilamino)-propoxi]-fenil}-amida;

4-oxo-4,5,6,7-tetrahidro-1H-indol-3-ácido carboxílico imidazo[1,2-a]piridin-5-ilamida; o

4-oxo-3a,4,5,6,7,7a-hexahidro-1H-indol-3-ácido carboxílico[6-(3-propil-[1,2,4]tidazol-5-ilamino)-piridin-2-il]amida.

39. Compuesto según la Reivindicación 1, de la fórmula:



o las sales no tóxicas farmacéuticamente aceptables de la misma donde:

Q es fenil o 3-piridil, cada uno de los cuales puede ser monosustituido o bisustituido por un hidroxil o un halógeno;

W es oxígeno o nitrógeno;

Z es hidroxil, C₁-C₆, cicloalquil (C₁-C₆ alcoxi), o (C₁-C₆ alcoxi) donde la porción alquilo es sustituida opcionalmente con amino, mono- o di(C₁-C₆ alquil₁)amino, o C₁-C₆ azacicloalquil;

amino o mono- o di(C₁-C₆ alquil₁)amino, o C₃-C₇, cicloalquil opcionalmente sustituido con C₁-C₆ alquil, o C₃-C₇, cicloalquil C₁-C₆ alquil;


Z es fenil o fénil(C₁-C₆)alquil donde la porción fenil es opcionalmente sustituida por C₁-C₆ alquil, hidroxil, C₁-C₆ alcoxi, trifluorometil, trifluorometoxi, halógeno, amino, o mono- o diC₁-C₆ alquilamino, o

Z es 2-, 3-, o 4-piridil, 1- o 2-imidazolil, 1-, 2-, o 3-pirrolil, o adamantano-2-il; cada uno de los cuales puede ser sustituido en un carbono terciario o un nitrógeno secundario por C₁-C₆alquil, o

Z es NR₉COR₁₀ donde R₉ y R₁₀ son iguales o diferentes y representan hidrógeno o C₁-C₆ alquil o C₃-C₇, cicloalquil, o

Z está unido, opcionalmente a través de W a Q desde un anillo de 1 a 6 miembros; o

Z representa un grupo de la fórmula:




donde

p es 1, 2, o 3;

D y D' representan independientemente oxígeno, NR_y o CHR_y siempre y cuando sólo uno de D y D' sea NR_y donde cada R_y es hidrógeno o C₁-C₆ alquil; y

R_z es hidrógeno o C₁-C₆ alquil, o

Z representa un grupo de la fórmula:




donde

p es 1, 2 o 3;

q es 0, 1 o 2;

R_z es hidrógeno o C₁-C₆ alquil; o

un grupo de la fórmula:




donde

s es 0, 1, 2 o 3, y la suma de s y m es no menos de 1;

R₀ es hidroxil, C₁-C₆alcoxi, amino, mono- o diC₁-C₆alquilamino, donde cada alquil es independiente u opcionalmente sustituido por amino, mono o diC₁-C₆alquilamino, o

R₀ es un grupo de la fórmula



donde p, D, D' y R₂ son según se define a continuación;



y



representan independientemente una cadena de carbono opcionalmente sustituida por hidrógeno, halógeno, oxo, ciano, nitro, amino, mono o di(C₁-C₆)alquilamino, una cadena recta o ramificada C₁-C₆ alquil, C₂-C₆ alquenil, C₂-C₆ alquinil, trifluorometil, trifluorometoxi, o cicloC₁-C₆ alquil;

en donde

k es 0, 1, 2 o 3;

m es 0, 1, 2 o 3; y



representa una cadena de carbono opcionalmente sustituida por R₅ y R₆ y n es 0, 1, 2 o 3;

R₃, R₄, R₅, y R₆ son iguales o diferentes e independientemente seleccionados en cada ocurrencia de: hidrógeno, C₁-C₆ alquil, -COR₁₁ o -CO₂R₁₁ donde R₁₁ es C₁-C₆ alquil o C₃-C₇ cycloalquil; o -CONR₁₂R₁₃ donde R₁₂ y R₁₃ son independientemente seleccionados entre hidrógeno, C₁-C₆ alquil, C₃-C₇ cicloalquil, fenil, 2-, 3- o 4-piridil, o NR₁₂R₁₃ forma un grupo heterocíclico que es morfolinil, piperidinil, pirrolidinil o N-alquil piperazinil; o


R₃-R₄ pueden ser tomados juntos para formar una mitad cíclica que contenga 3 a 7 átomos de carbono; o

232
319


R₅-R₆ pueden ser tomados juntos para formar una mitad cíclica que contenga 3 a 7 átomos de carbono; y

Donde cada grupo alquil que forme un sustituyente R₃, R₄, R₅, o R₆ o una porción del mismo pueda ser independientemente sustituido por un hidroxí o mono- o dialquilamino donde cada alquil es independientemente un C₃-C₇ alquil o cicloalquil que contenga 3 a 7 átomos de carbono.

40. Compuesto de la fórmula A o fórmula B:



A



B

o una sal farmacéuticamente aceptable de la misma en donde

R₃, R₄, R₅ y R₆ son iguales o diferentes y son seleccionados de:


hidrógeno, alquil, -COR₁₁ o -CO₂R₁₁ donde R₁₁ es un alquil o cicloalquil que contiene 3 a 7 átomos de carbono; o -CONR₁₂R₁₃ donde R₁₂ y R₁₃ son independientemente seleccionados de hidrógeno, alquil, cicloalquil que contiene 3 a 7 átomos de carbono, fenil, 2-, 3-, o 4-piridil, o NR₁₂R₁₃ forma un grupo heterocíclico que es morfolinil, piperidinil, pirrolidinil, o N-alquil piperazinil; o

R₃-R₄ pueden ser tomados juntos para formar una mitad cíclica que contenga 3 a 7 átomos de carbono; o

R₅-R₆ pueden ser tomados juntos para formar una mitad cíclica que contenga 3 a 7 átomos de carbono; y

Donde cada grupo alquil que forma un sustituyente R₃, R₄, R₅ o R₆ o una porción del mismo puede ser independientemente sustituido por un hidroxil o mono- o dialquilamino donde cada alquil es independiente un alquil o un cicloalquil que contenga 3 a 7 átomos de carbono; y

G representa




donde

R_h es hidrógeno, halógeno, C₁-C₆alquil, C₁-C₆alcoxi o trifluorometil;

s es 0, 1, 2 o 3, y la suma de s y m es no menos de 1;

R_o es hidroxil, C₁-C₆alcoxi, amino, mono- o diC₁-C₆alquilamino donde cada alquil es independiente y opcionalmente sustituido por amino, mono- o diC₁-C₆alquilamino, o

R_o es un grupo de la fórmula




p es 1, 2, or 3;

D y D' representan independientemente oxígeno, NR_y o CHR_y siempre y cuando sólo uno de D y D' sea NR_y donde cada R_y es hidrógeno o C₁-C₆ alquil; y R_z es hidrógeno o C₁-C₆ alquil.

41. Compuesto según la Reivindicación 40, en donde R_h es hidrógeno o halógeno y R_o es un grupo de la fórmula:

325 :
322



donde

R_{14} es hidrógeno o C_1-C_6 alquil;

R_{15} es hidrógeno o C_1-C_6 alquil;

R_{16} es hidrógeno, etil o metil;

R_{17} es C_1-C_6 alquil; y

J es un grupo C_1-C_4 alquilenos.

42. Compuesto según la Reivindicación 40, en donde s es 1 y R_0 es etoxi, hidroxil, etilamino, dietilamino, morfolinil, piperazinil, 4-metilpiperazinil,

