

Señor

SUPERINTENDENTE DE INDUSTRIA Y
DIVISION DE NUEVAS CREACIONES

106
107


SUPERINDUSTRIA Y COMERCIO
Radicación : 00076949 00000001
Fecha (AMB): 2000-12-27 17:13:09 Folios: 70
Trámite : 002 PATENTES D 1 REGISTRO/ 329 COMPLETEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

E. _____ S. _____ D. _____

Re: P1.-F. HOFFMANN-LA ROCHE AG.- Solicitud de Patente de Invención
para "PIRROLES SUSTITUIDOS".-Clase A61.-Expediente número 00-
076.949.-

1. Me refiero a mi solicitud de patente arriba nombrada de fecha 10 de Octubre de 2000.
2. Anexos me permito presentar a usted:
 - (1) Copia autentica, debidamente certificada de la solicitud de patente presentada para el invento arriba nombrado respecto de la cual reclamo prioridad, a saber la solicitud estadinense número 60/158.860 de 12 de Octubre de 1999, junto con su traducción Oficial correspondiente a las legalizaciones y reivindicaciones de dicho documento; y
 - (2) Copia debidamente autenticada por el Notario Sexto del Circulo de Bogotá, de los documentos que acreditan la calidad de Traductor Oficial de la señora María Mercedes Uricoechea quien elaboró la traducción de las legalizaciones y reivindicaciones de la solicitud estadinense número 60/158.860 de 12 de Octubre de 1999.
3. Ruego a usted tomar atenta nota de lo anterior y proceder de conformidad.

Del Señor Superintendente,


JOSE ANTONIO LLOREDA
T.P. 10.784
Código 231

REPUBLICA DE COLOMBIA
MINISTERIO DE RELACIONES EXTERIORES

CERTIFICACION

Se hace constar que el Señor(a) MARIA MERCEDES URICOECHEA TORRES


se encuentra debidamente registrado(a) ante el Ministerio de Relaciones
Exteriores, como traductor e interprete oficial para los idiomas _____

Ingles - Español - Ingles según resolución N° 10607

de fecha 31 de Diciembre de 1981 del Ministerio de Justicia

Se expide a los 6 días del mes de Abril de 1983

Atentamente


RAFAEL HUMBERTO PEÑA FLOREZ
JEFE DE NEGOCIOS GENERALES

COMO NOTARIO S
DE SANTAFÉ DE BOGOTÁ
HAGO CONSTAR
QUE LA FOTOCOPIA COINCIDE CON LA FOTOCOPIA
AUTENTICA QUE HE TENIDO A LA VISTA
SANTAFÉ DE BOGOTÁ
16-11-1998

COMO NOTARIO SEXTO DE
SANTAFÉ DE BOGOTÁ
HAGO CONSTAR
QUE LA FOTOCOPIA COINCIDE CON LA FOTOCOPIA
AUTENTICA QUE HE TENIDO A LA VISTA
SANTAFÉ DE BOGOTÁ
29-III-1983
JUAN MANUEL BOTERO MEDINA
NOTARIO

COMO NOTARIO SEXTO DE
SANTAFÉ DE BOGOTÁ
HAGO CONSTAR
QUE LA FOTOCOPIA COINCIDE CON LA FOTOCOPIA
AUTENTICA QUE HE TENIDO A LA VISTA
SANTAFÉ DE BOGOTÁ
16-IV-1998

COMO NOTARIO SEXTO DE
SANTAFÉ DE BOGOTÁ
HAGO CONSTAR
QUE LA FOTOCOPIA COINCIDE CON LA FOTOCOPIA
AUTENTICA QUE HE TENIDO A LA VISTA
SANTAFÉ DE BOGOTÁ
24-NOV-1998

ESTADOS UNIDOS DE AMÉRICA

A TODOS A QUIENES LLEGUE EL PRESENTE:

DEPARTAMENTO DE COMERCIO DE LOS ESTADOS UNIDOS

Oficina de Marcas y Patentes

26 de junio de 2000.

LA PRESENTE CERTIFICA QUE LA ADJUNTA ES UNA COPIA TOMADA DE LOS REGISTROS DE LA OFICINA DE MARCAS Y PATENTES DE LOS ESTADOS UNIDOS, DE LOS DOCUMENTOS DE LA SOLICITUD DE PATENTE ABAJO IDENTIFICADA Y LA CUAL CUMPLE CON LOS REQUISITOS PARA QUE LE SEA CONFERIDA UNA FECHA DE RADICACIÓN SEGUN 35 USC 111.

NUMERO DE SOLICITUD 60/158.860

FECHA DE RADICACIÓN: 12 de octubre de 1999

Por Autoridad del

COMISIONADO DE MARCAS Y PATENTES

(FIRMA H. L. JACKSON

Oficial de Certificaciones

(SELLO DORADO DE OBLEA) .

Maria Mercen
Res. No. 1
Traducción

Usted Uroschus

109
105

CERTIFICACION BAJO 37 CFR § 1.10

Certifico que esta Remisión de Nueva Solicitud y los documentos que en ella se refieren como adjuntos son depositados en el Servicio Postal de lo EE.UU. el 12 de octubre de 1999 en un sobre como "Oficina Postal al Destinatario" bajo el rótulo de correo No. **EL275412371US** y dirigido al Comisionado Asistente de Patentes, Washington, D.C. 20231.

(FIRMA) PAMELA JOHNSON

CUBIERTA SOLICITUD PROVISIONAL

La presente es una solicitud para la radicación de UNA SOLICITUD PROVISIONAL bajo 37 CFR § 1.53 (b)(2)

INVENTORES SOLICITANTES

Nombre: Fotouhi, Nader
Residencia: Chatham, Morris County, New Jersey
Ciudadano Francés
Dirección Postal 1 Daniel Street, Chatham, New Jersey 07928

Nombre: Kong, Norman
Residencia: West Caldwell, Essex County, New Jersey.
Ciudadano Canadiense
Dirección Postal 12 Klimback Court, West Caldwell, New Jersey 07006


Nombre: Liu, Emily Aijun
Residencia: Nutley, Essex County, New Jersey
Ciudadanía: Estadounidense
Dirección Postal: 191 Alexander Avenue, Nutley, New Jersey 07110

Nombre: Lovey, Allen John
Residencia: North Caldwell, Essex County, New Jersey
Ciudadanía: Estadounidense
Dirección Postal 21 Hickory Drive, North Caldwell, New Jersey 07006

Nombre: Mjullin, John Guilfoyle, Jr
Residencia: Hawthorne, Passaic County, New Jersey
Ciudadanía: Estadounidense
Dirección Postal: 519 Goffle Hill Road, Hawthorne, New Jersey 07506

TITULO DE LA INVENCION

PIRROLOS SUSTITUIDOS ADECUADOS PARA INFUSION CONTNUA



DIRECCION PARA CORRESPONDENCIA

George W. Johnston
Hoffmann-La Roche Inc.
340 Kingsland Street
Nutley, NNJ 07110

SE ADJUNTAN LA SIGUIENTES PARTES DE LA SOLICITUD

- [x] Especificación No. de páginas 44
[x] Reivindicaciones No. de páginas 24
[x] Declaración No. de páginas 4

METODO DE PAGO

- [x] Se autoriza al Comisionado de Patentes Comisión Provisional de radicación
a cargar las comisiones o acreditar
cualquier sobrepago a la Cuenta de \$150.00
Depósito No. 08-2525

La invención fue hecha por una agencia del Gobierno de los Estados Unidos o bajo un contrato con una agencia del Gobierno de los Estados Unidos.

- [x] No.

Respetuosamente,


Apoderado
Patricia S. Rocha-Tramaloni
(Reg. No. 31054)
Tel: (973)(235-2441
Fax: (973) 235-2363
Fecha: 12 de octubre de 1999

Maria Mercedes Urcoeches
Res. No. 10607 Minjusticia
Traductora Juramentada
Urcoc

PIRROLOS SUSTITUIDOS ADECUADOS PARA INFUSIÓN CONTINUA

REIVINDICACIONES:

1. Un compuesto de la fórmula:



y sales farmacéuticamente activas de dichos compuestos, en donde:

R es seleccionado del grupo compuesto por $-\text{CH}_2\text{OPO}_3\text{R}^1\text{R}^2$, $-\text{CH}_2\text{OCOR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{CH}_2\text{OCONHR}^3$, y $-\text{CONHR}^3$;

R^1 y R^2 son seleccionados del grupo compuesto por H, Na y NH_4 y son los mismos salvo que R^1 o R^2 sea H, en cuyo caso el otro puede ser diferente, o alternativamente, R^1 y R^2 juntos representan calcio.

R^3 es seleccionado del grupo compuesto por:

alquil que opcionalmente puede ser sustituido por uno o más sustituyentes seleccionados del grupo compuesto por $-\text{CO}_2\text{R}^4$, $-\text{NR}^5\text{R}^6$, polietilenglicol, $-\text{OPO}_3\text{R}^1\text{R}^2$, hidroxil, alcoxi y aril; alquenil que opcionalmente puede ser sustituido por uno o más sustituyentes seleccionados del grupo compuesto por $-\text{CO}_2\text{R}^4$, $-\text{NR}^5\text{R}^6$, polietilenglicol,

-OPO₃R¹R², hidroxil, alcoxi y aril; cicloalquil que opcionalmente puede ser sustituido por uno o más sustituyentes seleccionados del grupo compuesto por -CO₂R⁴, -NR⁶, polietilenglicol, -OPO₃R¹R³, hidroxil, alcoxi y aril;

heterociclo que cuando incluye N como un heteroátomo, opcionalmente, el N opcionalmente puede ser sustituido con el grupo compuesto por alquil más bajo y -COR⁷,

aril que opcionalmente puede ser sustituido por uno o más sustituyentes seleccionados del grupo compuesto por CO₂R⁴, hidroxil, alcoxi, polietilenglicol, OPO₃R¹R² y alquil que en sí puede ser sustituido por hidroxil alcoxi, carboxil y amino sustituido, siempre y cuando que cuando aril represente piridina, el nitrógeno puede ser sustituido por alquil más bajo;

R⁴ es seleccionado del grupo compuesto por H, Na y alquil más bajo;

R⁵ y R⁶ son cada uno seleccionados independientemente del grupo compuesto por H, alquil más bajo y -COR⁷, o alternativamente, el grupo -NR⁵R⁶ juntos forman un anillo heterocíclico de 5 ó 6 miembros; y

R⁷ es alquil más bajo que opcionalmente puede ser sustituido por carboxil, polietilenglicol polietilen glicol y amino sustituido.

2. El compuesto de la reivindicación 1, en donde R es seleccionado del grupo compuesto por -CH₂OPO₃R¹R² y -CH₂OCOR³.

3. El compuesto de la reivindicación 1, en donde R es -CH₂OCO-piridina, en donde el átomo N en la piridina es sustituido por alquil más bajo.

4. El compuesto de la reivindicación 3, en donde el átomo N en la piridina es sustituido por metil o etil.

5. El compuesto de la reivindicación 1, en donde R¹ y R² son seleccionados independientemente del grupo compuesto por H y Na.

6. El compuesto de la reivindicación 1, en donde R³ es heterociclo que contiene al menos un átomo de nitrógeno que opcionalmente puede ser sustituido por -COR⁷.

7. El compuesto de la reivindicación 1, en donde R³ es aril el cual es sustituido por -OPO₃R¹R² y R¹ y R² son seleccionados independientemente de H y Na.

8. El compuesto de la reivindicación 1, en donde R³ es aril el cual es sustituido por el grupo compuesto por -CO₂Na, polietilenglicol y -CH₂CH₂N(CH₂CH₂)₂.

9. El compuesto de la reivindicación 1, en donde R³ es alquil más bajo el cual es sustituido por -CHO₂Na.

10. El compuesto de la reivindicación 1, en donde el grupo NR⁵R⁶ junto forma un anillo heterocíclico de 5 ó 6 miembros.

11. El compuesto de la reivindicación 10, en donde el grupo NR^5R^6 es seleccionado de piperidina o pirrolidina.

12. El compuesto de la reivindicación 1, en donde R^5 y R^6 son seleccionados cada uno del grupo compuesto por H, metil y etil.

13. El compuesto de la reivindicación 1, en donde R^7 es etil que es sustituido por polietilenglicol.

14. El compuesto de la reivindicación 1, en donde R es $-CH_2OCOR^3$ y R^3 es etil el cual es sustituido por polietilenglicol que tiene un peso molecular aproximado entre 750 a 5000 Daltones.

15. El compuesto de la reivindicación 1, en donde el polietilenglicol tiene un peso molecular aproximado entre 750 y 5000 Daltones.

16. Un compuesto seleccionado del grupo compuesto por

ácido forsfórico mono-[3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-Indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil]sal sódica,

ácido isonicotínico 3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-Indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil ester,

ácido But-2-enedióico mono[3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil] ester,

ácido decanóico 3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil ester,

ácido 3-amino-ciclohexanocarboxílico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil éster trifluoroacetato,

ácido piperidin-4-carboxílico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil sal de acetato estérico,

ácido piperidin-4-carboxílico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil clorhidrato estérico,

ácido 2-amino-2-metil-propiónico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil clorhídrico estérico,

1-Metil-3-[3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetoxycarbonil]-píridinio; trifluoroacetato, y

O-[2-[4-[[[2,5-Dihidro-3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-1H-pirrol-1-ilmetoxi[carbonil]-1-piperidinil]carbonil]etil]-O-metilpolietilenglicol 1000.

17. Un compuesto seleccionado del grupo compuesto por:

ácido 4-fosfonooxi-benzóico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil sal monosódica estérica,

[[3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-il],etoxicarbonil]-4-fenil-O-metilpolietilenglicol 500,

ácido carbónico alil éster 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil éster,

ácido (2-dimetilamino-etil)-carbámico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil éster,

[2-(2-Hidroxi-etoxi)-etil]-ácido carbámico 3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil éster,

ácido (2-Hidroxi-etil)-carbámico 3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil éster,

ácido (2-fosfonooxi-etil)-carbámico 3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil éster,

3-(1-Metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-Indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1- ácido carboxílico (2-dietilamino-etil)-amida,

3-(1-Metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1- ácido carboxílico (2-dimetilamino-hexil)-amida,

3-(1-Metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1- ácido carboxílico (4-hidroxi-butil)-amida, y

ácido fosfórico-(4-{[3-(1-Metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-Indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-carbonil]amino}-butil)sal sódica estérica.

18. Un compuesto seleccionado del grupo compuesto por:

ácido fosfórico mono-[3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetil]sal sódica,

O-[2-[[2,5-dihidro-3-(1-metil-1H-indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-il]metoxycarbonil]etil]-O'-metilpolietilen glicol 2000, y

ácido fosfórico mono-(4-{[3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-carbonil]-amino}-butil)sal sódica estérica, y

1-metil-3-[3-(1-metil-1H-Indol-3-il)-4-(1-metil-6-nitro-1H-indol-3-il)-2,5-dioxo-2,5-dihidro-pirrol-1-ilmetoxycarbonil]-piridinio trifluoroacetato.

19. Una composición farmacéutica que comprende como ingrediente activo una cantidad efectiva de un compuesto de la reivindicación 1, y un vehículo o excipiente farmacéuticamente aceptable.

20. La composición farmacéutica de la reivindicación 19, adecuada para administración parenteral.

21. Un método para el tratamiento de un trastorno proliferativo de célula que consiste en administrar a un paciente que lo requiera, una cantidad terapéuticamente efectiva de un compuesto según la reivindicación 1.

22. El método de la reivindicación 21, en donde el trastorno proliferativo de células es cáncer.

23. El método de la reivindicación 22, en donde el cáncer es un tumor sólido.

24. El método de la reivindicación 22, en donde el cáncer es cáncer de seno, colon o pulmón.

PA 266936

THE UNITED STATES OF AMERICA**TO ALL TO WHOM THESE PRESENTS SHALL COME:****UNITED STATES DEPARTMENT OF COMMERCE****United States Patent and Trademark Office****June 26, 2000**

**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A
FILING DATE UNDER 35 USC 111.**

APPLICATION NUMBER: 60/158,860**FILING DATE: October 12, 1999**

**By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS**

**H. L. JACKSON****Certifying Officer**

116
Approved

CERTIFICATION UNDER 37 CFR 1.18

I hereby certify that this New Application Transmittal and the documents referred to as enclosed therein are being deposited with the United States Postal Service on this date October 12, 1999 in an envelope as "Express Mail Post Office to Addressee" Mailing Label Number EL275412371US addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Pamela Johnston

(Print Name)

Pamela Johnston
(Signature)

JOHN J. U.S. PTO
60/150860

10/12/99

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(b)(2).

Docket No. 1094P

INVENTOR(s)/APPLICANT(s)

Name: Fotouhi, Nader
Residence: Chatham, Morris County, New Jersey
Citizenship: France
Post Office Address: 1 Daniel Street, Chatham, New Jersey 07921

Name: Kong, Norman
Residence: West Caldwell, Essex County, New Jersey
Citizenship: Canada
Post Office Address: 12 Klimback Court, West Caldwell, New Jersey 07006

Name: Liu, Emily Aijun
Residence: Nutley, Essex County, New Jersey
Citizenship: United States of America
Post Office Address: 191 Alexander Avenue, Nutley, New Jersey 07110

Name: Lovey, Allen John
Residence: North Caldwell, Essex County, New Jersey
Citizenship: United States of America
Post Office Address: 21 Hickory Drive, North Caldwell, New Jersey 07006

Name: Mullin, John Guilfoyle, Jr.
Residence: Hawthorne, Passaic County, New Jersey
Citizenship: United States of America
Post Office Address: 519 Goffie Hill Road, Hawthorne, New Jersey 07506

TITLE OF THE INVENTION

SUBSTITUTED PYRROLES SUITABLE FOR CONTINUOUS INFUSION

DEPOSIT ACCOUNT
NO. 08-2525
OUR ORDER NO. 278

10/12/99
JOHN J. U.S. PTO
60/150860

6015 850-101299

120
117

10/12/99
JCS88 U.S. PTO

CORRESPONDENCE ADDRESS

George W. Johnston
Hoffmann-La Roche Inc.
340 Kingsland Street
Nutley, NJ 07110

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification No. of Pages 49 ☒ Declaration No. of Pages 4
☒ Claims No. of Claim 24 ☐ Drawings Sheets of Drawings
☐ Sequence Listing No. of Pages ☐ Other

METHOD OF PAYMENT

☒ The Commissioner is hereby authorized to charge any fees or credit any overpayment to Deposit Account Number: 08-2525 Provisional Filing Fee Amount \$150.00

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No
☐ Yes, the name of the U.S. Government agency and the Government contract number are: _____

Respectfully submitted,



Attorney of Record
Patricia S. Rocha-Tramaleri
(Reg. No. 31054)
Telephone: (973) 235-2441
Telefax: (973) 235-2363
Date: October 12, 1999

60158860-101299

123
118

SUBSTITUTED PYRROLES SUITABLE FOR CONTINUOUS INFUSION

5

Brief Summary of the Invention

10 The present invention is directed to certain substituted pyrroles that are anti-proliferative agents. These compounds and their pharmaceutically acceptable salts are useful in the treatment or control of cell proliferative disorders, in particular cancer. The invention is also directed to pharmaceutical compositions containing such compounds, and to methods for the treatment and/or prevention of cancer, particularly the treatment or control of solid tumors.

15

Background of the Invention

20 Uncontrolled cell proliferation is the hallmark of cancer. Cancerous tumor cells typically have some form of damage to the genes that directly or indirectly regulate the cell-division cycle. Much research has been expended in the study of antiproliferative agents. While many agents having desired antiproliferative activities have been identified, many of these agents have various drawbacks, including poor solubility, molecular complexity, etc., which may render them either unsuitable or inconvenient for therapeutic use in human patients. There continues to be a need
25 for small molecule compounds that may be readily synthesized, are effective as cancer therapeutic agents and are suitable for continuous infusion delivery to patients. It is thus an object of this invention to provide such compounds as well as pharmaceutical compositions containing such compounds.

30

662701-0985109

Definitions

As used herein, the following terms shall have the following definitions.

5 "Alkyl" denotes a straight-chain or branched saturated aliphatic hydrocarbon having 1 to 15, preferably 1 to 10, carbon atoms. Alkyl groups may be substituted as specifically provided *infra*. In addition the alkyl chain may include one or more hetero atoms in lieu of one or more carbon atoms. "Lower alkyl" groups having from 1 to 6, preferably 1 to 4, carbon atoms are preferred. Typical lower alkyl groups include
10 methyl, ethyl, propyl, isopropyl, butyl, t-butyl, 2-butyl, pentyl, hexyl, and the like.

15 "Alkenyl" means a straight-chain or branched aliphatic unsaturated hydrocarbon having 1 to 15 carbon atoms, preferably 1 to 10 carbon atoms, most preferably 1 to 6 carbon atoms.

20 "Alkoxy" means an alkyl group that is attached to the remainder of the molecule by oxygen (e.g. RO-, such as methoxy, ethoxy, etc.).

25 "Aryl" means an aromatic ring having 5 to 10 atoms and consisting of 1 or 2 rings, which optionally may include one or more heteroatoms that are the same or different. For the purposes of this definition, aryl includes heteroaryl. Preferred heteroatoms include nitrogen, sulfur, or oxygen, singly or in any combination, in place of one or more of the carbons. Examples of aryl groups within this definition are phenyl, pyridine, imidazole, pyrrole, triazole, furan, pyrimidine.

30 "Cycloalkyl" means a non-aromatic, partially or completely saturated cyclic aliphatic hydrocarbon group containing 3 to 8 atoms. Examples of cycloalkyl groups include cyclopropyl, cyclopentyl and cyclohexyl.

 "Effective amount" means an amount of at least one compound of Formula I or a pharmaceutically acceptable salt thereof that significantly inhibits proliferation

5015885 - 101299

125
120

and/or prevents differentiation of a human tumor cell, including human tumor cell lines.

"Hetero atom" means an atom selected from nitrogen, sulfur and oxygen.

5 Hetero atoms are independently selected and may replace one or more carbon atoms.

"Heterocycle" means a 3- to 10- membered non-aromatic, partially or completely saturated hydrocarbon group that contains at least one hetero atom.

10 Such ring systems include, morpholine, pyrrolidine, piperidine, piperazine

"IC₅₀" refers to the concentration of a particular compound according to the invention required to inhibit 50% of a specific measured activity. IC₅₀ can be measured, inter alia, as is described in Example 25, *infra*.

15

"Pharmaceutically acceptable salt" refers to conventional acid-addition salts or base-addition salts which retain the biological effectiveness and properties of the compounds of formula I and are formed from suitable non-toxic organic or inorganic acids or organic or inorganic bases. Sample acid-addition salts include those
20 derived from inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid and nitric acid, and those derived from organic acids such as acetic acid, tartaric acid, salicylic acid, methanesulfonic acid, succinic acid, citric acid, malic acid, lactic acid, fumaric acid, and the like. Sample base-addition salts include those derived from ammonium, potassium, sodium and,
25 quaternary ammonium hydroxides, such as for example, tetramethylammonium hydroxide.

"Pharmaceutically acceptable," such as pharmaceutically acceptable carrier, excipient, etc., means pharmacologically acceptable and substantially non-toxic to
30 the subject to which the particular compound is administered.

226

121

"Pharmaceutically active metabolite" means a metabolic product of a compound of formula I which is pharmaceutically acceptable and effective.

5 "Plasma conversion" with respect to compounds of formula I means the degradation (enzymatic and/or non-enzymatic) of such compound in human or rodent plasma at 37° C from 30 minutes to 6 hours to give 3-(1-methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione, a pharmaceutically active metabolite of compounds of formula I, as well as pharmaceutically active metabolites thereof. This conversion is typically given as the percent degradation
10 over a specified time frame.

60158860-101299
15 "Polyethylene glycol" or "PEG" groups represent structures of the general formula $-O(CH_2CH_2)_nOR^8$, where n is on average between 2 and 1500, preferably 15 to 150, with an average molecular weight of 500 to 5000 Daltons, and wherein R^8 is carboxy or lower alkyl, preferably methyl or ethyl.

20 "Prodrug" refers to a compound that may be converted under physiological conditions or by solvolysis to a pharmaceutical active compound. A prodrug may be inactive when administered to a subject but is converted *in vivo* to an active compound.

"Stability" is an overall assessment of the ability of a compound of formula I to withstand degradation in a typical solution used for the administration of drugs intravenously. Specifically, it refers to the ability of any given compound of formula I
25 to release 3-(1-methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione, over a 72 hour period in a mixture of acetonitrile and saline or dextrose water. The "stability" of a compound of formula I is "very good" if the less than 1% 3-(1-Methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione is detected, "good" if less than 2.5% 3-(1-Methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-
30 dione is detected, and "fair" if less than 5% 3-(1-Methyl-3-indolyl)-4-(1-methyl-6-


nitro-3-indolyl)-1H-pyrrole-2,5-dione is detected after 72 hours incubation at room temperature.

"Substituted," as in substituted alkyl, means that the substitution can occur at one or more positions and, unless otherwise indicated, that the substituents at each substitution site are independently selected from the specified options.

"Substituted amino" means an amino group which is mono- or di-substituted with another group, preferably lower alkyl (e.g. methyl, or ethyl).

Detailed Description of the Invention

Specifically, the invention relates to substituted pyrroles having the formula:



and pharmaceutically acceptable salts of the foregoing compounds, wherein :

R is selected from the group consisting of $-CH_2OPO_3R^1R^2$, $-CH_2OCOR^3$, $-CH_2OCO_2R^3$, $-CH_2OCONHR^3$, and $-CONHR^3$,

R^1 and R^2 are selected from the group consisting of H, Na and NH_4 and are the same unless either R^1 or R^2 is H, in which case the other can be different, or alternatively, R^1 and R^2 together represent calcium.

R^3 is selected from the group consisting of

126
123

alkyl which optionally may be substituted by one or more substituents selected from the group consisting of $-\text{CO}_2\text{R}^4$, $-\text{NR}^5\text{R}^6$, polyethylene glycol, $-\text{OPO}_3\text{R}^1\text{R}^2$, hydroxy, alkoxy and aryl; alkenyl which optionally may be substituted by one or more substituents selected from the group consisting of $-\text{CO}_2\text{R}^4$, $-\text{NR}^5\text{R}^6$, polyethylene glycol, $-\text{OPO}_3\text{R}^1\text{R}^2$, hydroxy, alkoxy and aryl; cycloalkyl which optionally may be substituted by one or more substituents selected from the group consisting of $-\text{CO}_2\text{R}^4$, $-\text{NR}^5\text{R}^6$, polyethylene glycol, $-\text{OPO}_3\text{R}^1\text{R}^2$, hydroxy, alkoxy and aryl;

heterocycle, which when including N as a heteroatom, the N optionally may be substituted with lower alkyl and $-\text{COR}^7$,

aryl which optionally may be substituted by one or more substituents selected from the group consisting of CO_2R^4 , hydroxy, alkoxy, polyethylene glycol, $\text{OPO}_3\text{R}^1\text{R}^2$, and alkyl which itself may be substituted with hydroxy alkoxy, carboxy and substituted amino, provided that when aryl represents pyridine, the nitrogen may be substituted with lower alkyl;

R^4 is selected from the group consisting of H, Na and lower alkyl;

R^5 and R^6 are each independently selected from the group consisting of H, lower alkyl, and $-\text{COR}^7$, or alternatively, the group $-\text{NR}^5\text{R}^6$ together form a 5 or 6 membered heterocyclic ring; and

R^7 is lower alkyl which optionally may be substituted with carboxy, polyethylene glycol and substituted amino.

The compounds of formula I have antiproliferative activity, specifically, they inhibit cell division in G2/M phase of the cell cycle and are generally referred to as "G2/M phase cell-cycle" inhibitors. These compounds are stable, soluble prodrugs of

124

an anticancer therapeutic agent within U.S. Patent 5,057,614 and are thus suitable for continuous infusion delivery.

The present invention is further directed to pharmaceutical compositions comprising a pharmaceutically effective amount of any one or more of the above-described compounds and a pharmaceutically acceptable carrier or excipient.

The present invention is also directed to a method for treating solid tumors, in particular breast or colon tumors, by administering to a human patient in need of such therapy an effective amount of a compound of formula I and/or its pharmaceutically acceptable salts.

In a preferred embodiment of the compounds of formula I, R is selected from the group consisting of $-\text{CH}_2\text{OPO}_3\text{R}^1\text{R}^2$, $-\text{CH}_2\text{OCOR}^3$ and $-\text{CONHR}^3$, most preferably $-\text{CH}_2\text{OPO}_3\text{R}^1\text{R}^2$ and $-\text{CH}_2\text{OCOR}^3$.

In another preferred embodiment of the compounds of formula I, R is $-\text{CH}_2\text{OCO}$ -pyridine wherein the N atom on the pyridine is substituted with lower alkyl, most preferably methyl or ethyl, thereby creating a quaternary nitrogen atom.

In another preferred embodiment of the compounds of formula I, R^1 and R^2 are independently selected from the group consisting of H and Na.

In another preferred embodiment of the compounds of formula I, R^3 is heterocycle containing at least one nitrogen atom that optionally may be substituted with $-\text{COR}^7$.

In another preferred embodiment of the compounds of formula I, R^3 is aryl which is substituted with $-\text{OPO}_3\text{R}^1\text{R}^2$, and R^1 and R^2 are independently selected from H or Na.

130
125

In another preferred embodiment of the compounds of formula I, R³ is aryl which is substituted by the group consisting of -CO₂Na, polyethylene glycol and -CH₂CH₂N(CH₂CH₂)₂.

5 In another preferred embodiment of the compounds of formula I, R³ is lower alkyl which is substituted with -CO₂Na.

10 In another preferred embodiment of the compounds of formula I, the group NR⁵R⁶ together forms a 5 or 6 membered heterocyclic ring, preferably piperidine or pyrrolidine.

In another preferred embodiment of the compounds of formula I, R⁵ and R⁶ are each independently selected from the group consisting of H, methyl and ethyl.

15 In another preferred embodiment of the compounds of formula I, R⁷ is ethyl which is substituted with polyethylene glycol.

In another preferred embodiment of the compounds of formula I, the polyethylene glycol has a molecular weight of from about 750 to about 5000 Daltons.

20 In another preferred embodiment of the compounds of formula I, R is -CH₂OCOR³, wherein R³ is ethyl which is substituted with PEG having a molecular weight of from about 750 to about 5000 Daltons.

25 The following are examples of preferred compounds of formula I:

phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl]sodium salt,

30 O-[2-[[2,5-dihydro-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-yl]methoxycarbonyl]ethyl]-O'-methylpolyethylene glycol 2000,

662101-098810S

121
126

phosphoric acid mono-(4-[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino]-butyl) ester sodium salt, and 1-methyl-3-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethoxycarbonyl]-pyridinium trifluoroacetate


The compounds disclosed herein and covered by the above formulae may exhibit tautomerism or structural isomerism. It is intended that the invention encompasses any tautomeric or structural isomeric form of these compounds, or mixtures of such forms, and is not limited to any one tautomeric or structural isomeric form utilized within the formulae drawn above.

Synthesis of Compounds According to the Invention

The compounds of the invention may be prepared by processes known in the art. Suitable processes for synthesizing these compounds are provided in the examples. Generally, these compounds may be prepared according to the following synthesis schemes.

Compounds of formula I, in which R signifies CH_2OCOR^3 , and in which R^3 is as described above, may be prepared as indicated in scheme I below, provided that if R^3 contains a hydroxy, hydroxyalkyl, amino, aminoalkyl, monoalkylamino, or carboxyl, such group is first protected with a conventional protective group known to those skilled in the art.

Scheme I



As indicated in scheme Ia, a chloromethyl ester prepared by reacting a known carboxylic acid or a carboxylic acid prepared by known methods, with ClCH₂OSO₂Cl in methylene chloride and water, in the presence of a base such as sodium carbonate and a phase transfer catalyst such as tetrabutylammoniumhydrogen sulfate, was reacted with 3-(1-methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione [prepared as exemplified in Davis US Patent 5,057,614].

Alternatively, 3-(1-methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione [prepared as exemplified in Davis US Patent 5,057,614] is treated with formaldehyde to yield the hydroxymethyl intermediate 2. This intermediate is then esterified using known procedures. Typically, the hydroxy intermediate 2 is treated

with a known carboxylic acid or a carboxylic acid prepared by known methods, in a solvent such as methylene chloride in the presence of EDC and dimethylaminopyridine for several hours at room temperature. Alternatively, the hydroxy intermediate 2 may be treated with a known acid chloride or an acid chloride prepared from known methods.


To prepare compounds of structure 3 wherein R³ contains a heteroaromatic ring, the heteroatom such as N may be further modified by reaction with an alkyl iodide such as CH₃I in a solvent such as acetonitrile. Alternatively compounds of structure 3 wherein R³ contains a suitably protected hydroxy, hydroxyalkyl, amino, aminoalkyl, monoalkylamino, may be further modified by first removing the protective group by known methods. The amino or hydroxy group can then be modified to the desired amide or ester by methods known in the art.

Compounds of the general formula I in which R signifies -CH₂OPO₃R¹R², and wherein R¹ and R² are as defined above, can be prepared by the following scheme II.

50158860-101299

129

Scheme II




Typically, hydroxymethyl Intermediate 2 is coupled with a suitably protected phosphate by a Mitsunobu reaction using triphenylphosphine and diethyl azodicarboxylate to give compound of structure 4 in which Y represents a suitable protecting group. Removal of the protecting groups may be achieved by any of the standard methods to give the phosphoric acid 5. In particular, when Y represents a benzyl group, the protective groups are removed by using cyclohexadiene and palladium on carbon as a catalyst. Compound 5 can then be converted to its salt, such as a monosodium salt 6, by standard methods.

Compounds of formula I, in which R signifies $-\text{CH}_2\text{OCO}_2\text{R}^3$, and in which R^3 is as described above, were prepared according to scheme III below.

125
130


Scheme III



Typically, hydroxymethyl intermediate 2 is treated with a known chloroformate or a chloroformate prepared using known procedures, in a solvent such as THF at temperatures of 5° to 20° C, in the presence of dimethylaminopyridine and 1,5-diazabicyclo(4.3.0)non-5-ene to afford the desired carbonate,

Compounds of formula I, in which R signifies -CH₂OCONHR³, and in which R³ is as described above, were prepared as described in scheme IV below.

Scheme IV




Typically, hydroxymethyl intermediate 2 is deprotonated using a strong base such as n-butyllithium or lithium bis(trimethylsilyl)amide in a solvent such as THF at 0° C. The anion generated is then treated in the same solvent with bis(p-

nitrophenyl)carbonate, followed by a known amine or an amine prepared using known procedures.

Compounds of formula I, in which R signifies -CONHR³, and in which R³ is as defined above, may be prepared as indicated in scheme V, provided that if R³ contains a hydroxy, hydroxyalkyl, amino, aminoalkyl, monoalkylamino, or carboxyl, such group is protected with a conventional protective group.

Scheme V



Typically, 3-(1-Methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione [prepared as exemplified in Davis U.S. Patent 5,057,614] is deprotonated in an aprotic solvent such as THF at 0° C using a strong base such as n-butyllithium or lithium bis(trimethylsilyl)amide. The resulting anion is then treated with bis(p-nitrophenyl)carbonate, followed by a known amine or an amine prepared by methods known in the art.

The conversion of an acidic compound of formula I into a pharmaceutically acceptable salt can be carried out by treatment with a suitable base in a known manner. Suitable salts are those derived not only from inorganic bases, for example, sodium, potassium or calcium salts, but also from organic bases such as ethylenediamine, monoethanolamine or diethanolamine. The conversion of a basic compound of formula I into a pharmaceutically acceptable salt can be carried out by treatment with a suitable acid in a known manner. Suitable salts are those described on page 3.

137
132

Compositions/Formulations

In an alternative embodiment, the present invention is directed to pharmaceutical compositions comprising at least one compound of formula I or a pharmaceutically acceptable salt thereof.

5

These pharmaceutical compositions can be administered orally, for example, in the form of tablets, coated tablets, dragees, hard or soft gelatin capsules, solutions, emulsions or suspensions. They can also be administered rectally, for example, in the form of suppositories. In particular, however, the compounds of the present invention are suitable for parenteral administration, for example, in the form of injection solutions.

10

The pharmaceutical compositions of the present invention comprising compounds of formula I, prodrugs of such compounds, or the salts thereof, may be manufactured in a manner that is known in the art, e.g. by means of conventional mixing, encapsulating, dissolving, granulating, emulsifying, entrapping, dragee-making, or lyophilizing processes. These pharmaceutical preparations can be formulated with therapeutically inert, inorganic or organic carriers. Lactose, corn starch or derivatives thereof, talc, steric acid or its salts can be used as such carriers for tablets, coated tablets, dragees and hard gelatin capsules. Suitable carriers for soft gelatin capsules include vegetable oils, waxes and fats. Depending on the nature of the active substance, no carriers are generally required in the case of soft gelatin capsules. Suitable carriers for the manufacture of solutions and syrups are water, polyols, saccharose, invert sugar and glucose. Suitable carriers for injection are water, alcohols, polyols, glycerine, vegetable oils, phospholipids and surfactants. Suitable carriers for suppositories are natural or hardened oils, waxes, fats and semi-liquid polyols.

15

20

25

The pharmaceutical preparations can also contain preserving agents, solubilizing agents, stabilizing agents, wetting agents, emulsifying agents, sweetening agents, coloring agents, flavoring agents, salts for varying the osmotic

30

5 15886 98851 101299

138
133

pressure, buffers, coating agents or antioxidants. They can also contain other therapeutically valuable substances, including additional active ingredients other than those of formula I.

5 **Dosages**

As mentioned above, the compounds of the present invention are useful in the treatment or control of cell proliferative disorders, in particular oncological disorders. These compounds and formulations containing said compounds are particularly useful in the treatment or control of solid tumors, such as, for example,
10 breast and colon tumors.

A therapeutically effective amount of a compound in accordance with this invention means an amount of compound that is effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated.
15 Determination of a therapeutically effective amount is within the skill in the art.

The therapeutically effective amount or dosage of a compound according to this invention can vary within wide limits and will be adjusted to the individual requirements in each particular case. In general, in the case of oral or parenteral
20 administration to adult humans weighing approximately 70 Kg, a daily dosage of about 10 mg to about 10,000 mg, preferably from about 200 mg to about 1,000 mg, should be appropriate, although the upper limit may be exceeded when indicated. The daily dosage can be administered as a single dose or in divided doses, or for parenteral administration, it may be given as continuous infusion.

25

Examples

The compounds of the present invention may be synthesized according to known techniques, such as, for example, the general schemes provided above. The following examples illustrate preferred methods for synthesizing the compounds and
30 formulations of the present invention.

0158860-101299

129
134

Example 1: Phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl]sodium salt

- 5 a) A suspension consisting of 0.5 g (1.25 mmole) of 3-(1-methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione (See Davis US Pat. No. 5,057,614), 4.0 ml of formaldehyde solution (37% w/w), and 2.0 ml of water was heated to 125° C with stirring and with a reflux condensed attached for 14 hours. The reaction mixture was cooled and diluted with water. The red solid was filtered, washed with water and dried. The solid was purified by chromatography
- 10 using silica gel and a mixture of EtOAc/Hexane as elutant to afford 1-hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione (mp=209-211° C).
- 15 b) To a cool solution (10° C) of THF containing 0.12 g (0.68 mmole) of diethylazodicarboxylate, 0.20 g (0.46 mmole) of 1-hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione (from Step a above), and 0.40 g (0.74 mmole) of tetrabenzylpyrophosphate was added dropwise 5 ml of a solution containing 0.128 g (0.49 mmole) of
- 20 triphenylphosphine. The resulting mixture was stirred for 14 hours at 20° C. All solvent was evaporated and the residue was purified by chromatography using silica gel and eluting with a mixture of EtOAc/Hexanes to afford phosphoric acid dibenzyl ester 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester as a brick red solid (mp=105-109° C).
- 25 c) A solution of 35 mg (0.051 mmole) of phosphoric acid dibenzyl ester 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (from step b above) in THF (5 ml) with 5 ml ethanol and 0.5 g of 1,4-cyclohexadiene was treated with 35 mg of 10% Palladium/carbon. The mixture was heated to 45° C at which point there was a slight exotherm. After 10
- 30 min. at 50° C, the reaction was filtered hot through celite and the filter cake was washed with THF. All solvent was evaporated and the residue was redissolved in THF, filtered through celite and triturated with hexane to give 14 mg of---

60158860-101299

phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl].

- 5 d) A solution of 14 mg (0.029 mmole) of phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl] (from step c above) in 15 ml of water was treated with dilute sodium hydroxide until the pH of the solution was 7.1. This solution was filtered and lyophilized to afford phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl]sodium salt.

10

Example 2: Isonicotinic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester

- 15 To a solution of dimethylaminopyridine (1.5 equivalents) and 1-ethyl-3-(3-diaminopropyl) carbodiimide hydrochloride (1.5 equivalent) in CH_2Cl_2 was added 4-picolinic acid (1.2 equivalents) and the reaction mixture was stirred a few minutes. 1-Hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione, prepared as in example 1a, was added and the reaction mixture was stirred at room temperature for several hours. The mixture was diluted with
20 dichloromethane, washed with aqueous HCL, saturated NaHCO_3 , water, dried over MgSO_4 and evaporated. The residue was purified by column chromatography to afford isonicotinic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield ~76%).

25 **Example 3**

Using the same procedure as in example 2 the following compounds were prepared:

- 30 a) nicotinic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield ~ 76%);

- b) O-[2-[[2,5-dihydro-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-yl]methoxycarbonyl]ethyl]-O'-methylpolyethylene glycol 2000 (yield - 88%);
- 5 c) pyridin-4-yl acetic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield - 39%);
- d) pyridin-3-yl acetic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield - 84%);
- 10 e) pyridin-2-yl acetic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield - 65%);
- f) succinic acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl] ester (used 3eq EDC and DMAP and 2.4 eq acid, purified by preparative TLC) (yield - 89%);
- 15 g) hexadec-9-enoic 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield - 53%);
- 20 h) [2-[2-(2-carboxymethoxy-ethoxy)-ethoxy]-ethoxy] acetic acid-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester, purified by reverse phase column chromatography (yield -61%).

25 **Example 4: But-2-enedioic acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl] ester**

To a solution of 1-Hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione, prepared as in example 1a, in 1:4 CH₂Cl₂ benzene, were added a large excess of fumaryl chloride and excess diisopropylethylamine. The reaction was stirred at room temperature for 20 min. Acetone and water were added. The reaction mixture was diluted with CH₂Cl₂, washed with water, cold

30

148
137

saturated NaHCO_3 , water and dried over Na_2SO_4 . The resulting but-2-enedioic acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl] ester was purified by flash chromatography (yield 60%).

5 **Example 5: Decanoic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester**

10 a) Decanoic acid (0.2 gm, 1.2 mmol) was treated with 4 eq NaHCO_3 , 0.1 eq Bu_4NHSO_4 and 1.2 eq $\text{ClCH}_2\text{OSO}_2\text{Cl}$ in 1:1 CH_2Cl_2 H_2O . The reaction was stirred vigorously at 0°C for 30 min then room temperature ("RT") for 3 hrs. Aqueous workup yielded chloromethyldecanoate (yield 100%).

15 b) 3-(1-Methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione (20 mg, 0.05 mmol) (See Davis US Pat. No. 5,057,614) was dissolved in DMF and treated with Ca_2CO_3 (49 mg, 0.15 mmol) for a few minutes, the chloromethyl ester (33 mg, 0.15 mmol) prepared in step a) above was added and the reaction was stirred at room temperature for 30 minutes. Aqueous workup, followed by purification by flash chromatography yielded decanoic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield - 58%).

20 **Example 6**

The following compounds were prepared in a similar manner to example 5b above:

25 a) [2-(2-methoxy-ethoxy)-ethoxy]-acetic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (the starting chloromethyl ester was prepared from the corresponding acid as described in example 5a; purified by preparative TLC) (yield - 38%);

30 b) 2,2-dimethyl propionic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (the starting chloromethyl pivalate was purchased from (Aldrich) (yield - 91%).

60158860-101299

143
138

Example 7: 3-amino-cyclohexanecarboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester trifluoroacetate

- 5 a) chloromethyl-N-a-butyloxycarbonyl-3-amino-cyclohexanecarboxylate was prepared in a similar manner (87% yield) as described in example 5a from N-a-butyloxycarbonyl-3-amino-cyclohexanecarboxylic acid which itself was synthesized by the known procedure of BOC protection of 3-amino-
- 10 cyclohexanecarboxylic acid (Aldrich).
- b) 3-(1-Methyl-3-Indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione (68 mg, 0.15 mmol) (see Davis US Pat. No. 5,057,614) was dissolved in DMF and treated with Cs_2CO_3 (0.15 g, 0.45 mmol) for a few minutes. The chloromethyl ester (0.13 gm, 0.45 mmol) prepared in step a) above was added and the reaction mixture was stirred at room temperature for 50 minutes. Aqueous workup, followed by purification by flash chromatography yielded [N-a-butyloxy-3-amino-
- 15 cyclohexanecarboxylic acid]-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (yield - 95%)
- 20 c) The tert-butyloxy protecting group of the compound of step b) above was removed by treatment with TFA in CH_2Cl_2 at room temperature for 50 minutes. The TFA and CH_2Cl_2 were evaporate using a stream of nitrogen. The residue was purified by HPLC to give 3-amino-cyclohexanecarboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-
- 25 ylmethyl ester trifluoroacetate (yield - 57%).

Example 8

- 30 The following compounds were prepared in a similar manner to example 7a,b and c described above:

- a) piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester trifluoroacetate (prepared from N-a-butyloxycarbonyl-piperidine-4-carboxylic acid (Bachem)) (yield - 62%);
- 5 b) 2-amino-2-methyl-propionic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester trifluoroacetate (prepared from N-a-butyloxycarbonyl-aminoisobutyric acid (Bachem)) (yield - 63%).

10 **Example 9: Piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester acetate salt**

Piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester trifluoroacetate, prepared as in
 15 example 8a above, was dissolved in ethyl acetate. The resulting solution was neutralized with cold saturated NaHCO₃. The ethyl acetate layer was concentrated and slurried in H₂O and acidified with excess HOAc and lyophilized to give the desired product (yield - 97%).

20 **Example 10: Piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester hydrochloride**

Piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester acetate salt, prepared as in
 25 example 9 above, was dissolved in ethyl acetate. The ethyl acetate solution was neutralized with cold saturated NaHCO₃, was concentrated, and was taken up in 1:5 CH₃CN: H₂O. The resulting solution was treated with 2 equivalents of 2N aq. HCl and purified by HPLC to give the desired product (yield - 86%).

30

60158860-101299

145
140

Example 11: 2-amino-2-methyl-propionic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester hydrochloride

5 This compound was prepared from 2-amino-2-methyl-propionic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester trifluoroacetate, prepared as in example 8b, using a similar procedure as described in example 10 (yield - 71%).

10 **Example 12:** 1-Methyl-3-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethoxycarbonyl]-pyridinium; trifluoroacetate

15 A solution of nicotinic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester in CH₃CN, prepared as described in example 3a, was treated with 2 eq NaBPh₄ and 6 eq MeI. The reaction was stirred at reflux for 7 hours. The solvent was evaporated and the residue was purified by HPLC to afford 1-methyl-3-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethoxycarbonyl]-pyridinium; trifluoroacetate
20 (yield 52%).

25 **Example 13:** O-[2-[[4-[[[2,6-Dihydro-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-1H-pyrrol-1-yl]methoxy]carbonyl]-1-piperidinyl]carbonyl]ethyl]-O'-methylpolyethylene glycol 1000

3-(1-Methyl-3-Indolyl)-4-(1-methyl-6-nitro-3-Indolyl)-1-(4-piperidinecarboxymethyl)-1H-pyrrole-2,5-dione trifluoroacetate, prepared as in example 8a, was dissolved in ethyl acetate. The resulting solution was neutralized with cold saturated NaHCO₃. The ethyl acetate layer was concentrated and the
30 residue was taken up in CH₃CN. The solution was added dropwise to a solution of O-(2-carboxyethyl)-O'-methylpolyethylene glycol 1000 acid chloride (prepared from O-(2-carboxyethyl)-O'-methylpolyethylene glycol 1000 acid using standard procedure) in CH₃CN. The reaction mixture was stirred for 1 hour. The solvent was evaporated and the residue was purified by HPLC (yield 70%).

35

0158860.101299

146
141

Example 14: 4-Phosphonoxy-benzoic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester monosodium salt

- 5 a) *t*-BuOK (9.6 mL, 9.6 mmol, 1M in THF) was added drop-wise to a solution of 4-hydroxybenzaldehyde (1.10 g, 9 mmol) in THF (40 mL). The mixture was heated at 70° C and tetrabenzyl pyrophosphate (5.05 g, 9.37 mmol) in THF (20 mL) was added. After 1 h, THF (100 mL) and hexanes (200mL) were added. The reaction mixture was filtered and the filtrate was evaporated. Chromatography of the
10 residue over silica using 2%-5%-10% methanol-CH₂Cl₂ gave phosphoric acid dibenzyl ester 4-formyl-phenyl ester (3.38 g, 98%).
- 15 b) Sodium dihydrogenphosphate monohydrate (79 mg, 0.57 mmol) in water (1 mL) was added to a solution of phosphoric acid dibenzyl ester 4-formyl-phenyl ester (0.89 g, 2.32 mmol) (from step a) above) in MeCN (10 mL) at 0° C. Hydrogen peroxide (0.24 mL, 30% in water, 2.4 mmol) was added, followed by sodium chlorite (315 mg, 3.5 mmol) in water (2 mL). The reaction mixture was stirred at 0° C for 1 h, then warmed to room temperature for 2 h. Sodium thiosulfate (100 mg) was added and the reaction mixture was vigorously stirred for 15 min. Water
20 was added and the mixture was extracted with EtOAc. The organic extracts were washed with water, dried (MgSO₄), filtered and concentrated to give phosphoric acid dibenzyl ester 4-carboxyl-phenyl ester (880 mg, 94%) as a pale yellow waxy solid.
- 25 c) A solution of phosphoric acid dibenzyl ester 4-carboxyl-phenyl ester (205 mg, 0.5 mmol) (from step b) above) in CH₂Cl₂ (2 mL) was added to a solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (149 mg, 0.78 mmol) (Aldrich) and Dimethylaminopyridine (139 mg, 1.13 mmol) in CH₂Cl₂ (5 mL). After 5 min., 1-Hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione (215 mg, 0.5 mmol), prepared as in example 1a, in
30 CH₂Cl₂ (2 mL), was added and the mixture was stirred at room temperature for 3 h. The resulting mixture was partitioned between CH₂Cl₂ and water and the

60158860-101299

149
142

organic extracts were washed with water. The aqueous layer was extracted with CH_2Cl_2 . The organic extracts were combined, dried (MgSO_4) and concentrated. Chromatography of the residue over silica gel using 5%-10% EtOAc- CH_2Cl_2 yielded 4-(bis-benzyloxy-phosphoryloxy)-benzoic acid 3-(1,6-dimethyl-1H-indol-3-yl)-4-(1-methyl-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (179 mg, 44%) as an orange foam.

- d) 1,4 Cyclohexadiene (0.12 mL, 1.26 mmol) (Aldrich) was added to a mixture of 4-(bis-benzyloxy-phosphoryloxy)-benzoic acid 3-(1,6-dimethyl-1H-indol-3-yl)-4-(1-methyl-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (100 mg, 0.12 mmol) (from step c) above) and Pd/C (37 mg 10%) in a mixture of THF (8 mL) and ethanol (0.3 mL). The reaction mixture was warmed to 55° C for 20 minutes. The reaction mixture was cooled to room temperature and was filtered over celite, washed with methylene chloride and concentrated. The residue was dissolved in MeCN and water at 0° C. 1N Sodium hydroxide (2.4 mL, 2.4 mmol) was added and the mixture was lyophilized. The resulting mixture was further purified by HPLC with 5 to 50% MeCN - water. Lyophilization yielded 4-phosphonooxy-benzoic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester monosodium salt (80 mg, 74%) as an orange powder.

Example 15: [3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-yl]methoxycarbonyl]-4-phenyl-O-methylpolyethylene glycol 500

- a) Dry PEG-500 monomethyl ether (18.36 g, 33 mmol), ethyl-4-hydroxybenzoate (5.0 g, 30 mmol) and triphenylphosphine (17.34 g, 66 mmol) were dissolved in dry THF (100 mL) and cooled to 0° C under argon. To this was slowly added diethylazodicarboxylate (12.95 mL, 83 mmol) in dry THF (10 mL). The resulting mixture was then warmed to 50° C for 16 h. The mixture was cooled to 0° C under argon and diethylazodicarboxylate (6 mL) in dry THF (5 mL) was added. The mixture was heated at 50° C for 24 h. Evaporation of the solvents and

148
143

chromatography of the residue over silica gel using 1:1 CH₂Cl₂-hexanes, CH₂Cl₂, 50% ether/-CH₂Cl₂, and ether yielded 9 g crude ethyl-4-O-methylpolyethylene glycol 500-benzoate as an oil, 5.4 g orange oil (3/2 mole% triphenylphosphine/product), and 4.9 g (~58 % overall yield) orange oil (5/4 mole% triphenylphosphine/product).

b) To a solution of the crude ethyl-4-O-methylpolyethylene glycol 500-benzoate (4.90g, 5/4 mole% triphenylphosphine/product) (from step a) above) in methanol (10 mL) and water (15 mL) was added 6N NaOH (13.8 mL, 83 mmol), and the resulting solution was heated to 65° C for 2h. The solution was concentrated and partitioned between 30% CH₂Cl₂/ethyl acetate and water. The aqueous layer was cooled to 0° C and acidified to pH 3.0 with 1N HCl. The mixture was extracted with CH₂Cl₂ and the organic extracts were dried (MgSO₄). Evaporation of the organic solvents gave crude 4-O-methylpolyethylene glycol 500-benzoic acid (2.3 g, 49%) as a yellow waxy residue. (sample contained 20-30% molar excess PEG 500).

c) A solution of the crude 4-O-methylpolyethylene glycol 500-benzoic acid (679 mg, 1 mmol) (from step b) above) in dry CH₂Cl₂ (5 mL) was added at room temperature to a solution of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (300 mg, 1.56 mmol) (Aldrich) and dimethylaminopyridine (279 mg, 2.28 mmol) in dry CH₂Cl₂ (5 mL). After 5 min., 1-Hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione (430 mg, 1 mmol), prepared as in example 1a, was added. The mixture was stirred at room temperature for 4 h. Water was added and the mixture was extracted with CH₂Cl₂. The organic extracts were washed with water, dried over (MgSO₄) and concentrated. The residue was purified by HPLC with MeCN / water to give [[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-yl]methoxycarbonyl]-4-phenyl-O-methylpolyethylene glycol 500 (490 mg, 47 %) as an orange oil.

149
144

Example 16: Carbonic acid allyl ester 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester

A solution of 200 mg (0.47 mmole) of 1-hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione, prepared as in the example 1a, was dissolved in 90 ml of THF and treated with 300 mg (2.5 mmol) of 4-dimethylaminopyridine (DMAP). To this solution at 5° C was added 90 mg (0.75 mmole) of allyl chloroformate. While still cool, 300 mg (2.4 mmol) of 1,5-diazabicyclo(4.3.0)non-5-ene (DBN) (Aldrich) was added dropwise. This was stirred for 14 hours at 20° C. The solvent was evaporated and the mixture was purified by chromatography using silica gel and eluting with a mixture of EtOAc/Hexane to afford carbonic acid allyl ester 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester as a red solid (m.p. 191-194° C).

Example 17: (2-Dimethylamino-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester

1-Hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione (200 mg, 0.47 mmol), prepared as in example 1a, was dissolved in THF (10 mL). The resulting red solution was cooled in an ice bath. Lithium bis(trimethylsilyl)amide (0.56 mL, 0.56 mmol, 1 M in THF) was added dropwise to the solution to give a red suspension. After 10 min, bis(p-nitrophenyl)carbonate (200 mg, 0.66 mmol) was added. The resulting solution was stirred at 0° C for 20 min. Diethylaminoethylamine (90 mg, 0.78 mmol) in THF (2 mL) was added. Stirring was continued for 2 hrs. The mixture was evaporated and the residue was purified by flash chromatography to give (2-dimethylamino-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (124 mg 46%).

280
145

Example 18: [2-(2-Hydroxy-ethoxy)-ethyl]-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester

5 This compound was prepared from 1-amino-2-ethoxy-ethanol (Aldrich), using the same procedure as in example 17 (46 mg, 17%).

Example 19: (2-Hydroxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester

10 *n*-BuLi (0.96 mL, 1.54 mmol, 1.6 M in hexanes) was added drop-wise to a solution of 1-hydroxymethyl-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-pyrrole-2,5-dione (600 mg, 1.54 mmol), prepared as in example 1a, in THF (30 mL) at 0°C. After 10 min bis(*p*-nitrophenyl)carbonate (600 mg, 1.96 mmol) was added.
15 The solution was stirred at 0°C for 20 min. A solution of 1-aminoethanol (128 mg, 2.10 mmol) was added. The resulting mixture was stirred at 0°C for 30 min. Aqueous NH₄Cl was added and the mixture was extracted with EtOAc. The organic extract was washed with brine and dried over MgSO₄. Evaporation of the solvents and chromatography of the residue over silica gel using 1:1 -3:2 EtOAc / hexanes
20 gave (2-hydroxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (339 mg, 47%).

Example 20: (2-phosphonoxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester

25 a) Triphenylphosphine (95 mg, 0.36 mmol) and dibenzyl phosphate (101 mg, 0.36 mmol) were added to a solution of (2-Hydroxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester, prepared as in example 19 (150 mg, 0.289 mmol) in THF (6 mL).
30 The mixture was cooled at -78°C. Diethyl azodicarboxylate (0.057 mL, 0.36 mmol) was added dropwise over 5 min. The cooling bath was removed and the mixture was stirred overnight. The mixture was then evaporated. Chromatography of the residue over silica gel using 2:1-4:1 EtOAc / hexanes gave [2-bis-benzyloxy-

60153860-101299

phosphoryloxy)-ethyl]-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (121 mg, 54%).

5 b) Pd / C (33 mg, 10%) and 1,4-cyclohexadiene (0.12 mL, 1.27 mmol) were added to a solution of [2-bis-benzyloxy-phosphoryloxy)-ethyl]-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (115 mg, 0.148 mmol) in a mixture of THF (5 mL) and EtOH (0.25 mL). The mixture was heated to 50-55°C for 30 min. The resulting mixture was filtered through celite, washed with THF and the filtrate was
10 evaporated. The residue was dissolved in a mixture of EtOH (10 mL) and CH₃CN (10 mL). 0.1 N NaOH was added and the pH was adjusted to 8.2. Water (10 mL) was added to dissolve the precipitate. The organic solvents were removed under vacuum and lyophilisation of the red solution gave (2-phosphonooxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester (59 mg, 62%) as
15 an orange powder.

Example 21: 3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (2-diethylamino-ethyl)-amide

20 To a cold solution of 3-(1-Methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione (200 mg, 0.5 mmol) (see Davis U.S. Pat. No. 5,057,614) in THF (7 mL) was added dropwise, lithium bis(trimethylsilyl)amide (0.55 mmol, 0.55 mL, 1 M in THF). The resulting red suspension was stirred for 15 min. Bis(p-nitrophenyl)carbonate (213 mg, 0.7 mmol) was added. The solution was stirred at
25 0°C for 0.5 h. Diethylaminoethylamine (69.7 mg, 0.6 mmol) in THF (2 mL) was added, and stirring was continued for 2 h. The solvent was evaporated and the residue was purified by flash chromatography to give 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (2-diethylamino-ethyl)-amide (28 mg, 10%).
30

182
14.7

Example 22: 3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (6-dimethylamino-hexyl)-amide

This compound was prepared from dimethylaminohexylamine according to the procedure described in example 21 (67 mg, 24%).

Example 23: 3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (4-hydroxybutyl)-amide

To a cold solution of 3-(1-methyl-3-indolyl)-4-(1-methyl-6-nitro-3-indolyl)-1H-pyrrole-2,5-dione (500 mg, 1.25 mmol) (see Davis U.S. Pat. No. 5,057,614) in THF (7 mL) was added dropwise n-butyllithium (0.86 mL, 1.375 mmol, 1.6 M in hexane). The resulting red suspension was stirred for 15 min. Bis(p-nitrophenyl)carbonate (532 mg, 1.75 mmol) was then added. The resulting solution was stirred at 0°C for 1 h. 4-aminobutanol (167 mg, 1.875 mmol) in THF (4 mL) was added, and stirring was continued for 2 h. The solvent was evaporated and the residue was purified by flash chromatography to afford 3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (4-hydroxybutyl)-amide as an orange solid (280 mg, 43%).

Example 24: Phosphoric acid mono-(4-([3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino)-butyl) ester sodium salt

a) Triphenylphosphine (0.86 g, 3.29 mmol) and Dibenzyl phosphate (0.91 g, 3.29 mmol) were added to a solution of 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (4-hydroxybutyl)-amide (prepared as in example 23, 1.41 g, 2.74 mmol) in THF (50 mL). The mixture was cooled to -78°C. Diethyl azodicarboxylate (0.52 mL, 3.29 mmol) was added dropwise over 5 min. The cooling bath was removed and after 2h, the mixture was evaporated. Chromatography of the residue over silica gel using 2:1 EtOAc / hexane then EtOAc gave phosphoric acid dibenzyl ester 4-([3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino)-butyl ester (1.76 g, 83%).

153
148

662TDT-09885TDS
5 b) Pd/C (0.5 g, 10%) and 1,4-cyclohexadiene (1.84 mL, 19 mmol) were added to a solution of phosphoric acid dibenzyl ester 4-[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino)-butyl ester (1.76 g, 2.27 mmol) (from step a) above) in a mixture of THF (75 mL) and EtOH (3.8 mL). The resulting mixture was heated to 50-55°C for 30 min. TLC (EtOAc) showed only a trace of phosphoric acid dibenzyl ester 4-[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino)-butyl ester left. The mixture was filtered through
10 celite and the filtrate was evaporated. The residue was dissolved in a mixture of EtOH (100 mL) and CH₃CN (100 mL) and basified with NaOH (0.2 N) to PH=8.2. Water (50 mL) was added to dissolve the precipitate. The organic solvents were removed under vacuum and lyophilisation of the red solution gave crude product (1.2 g). Purification of the residue using HPLC (10%-90% CH₃CN-H₂O) gave
15 phosphoric acid mono-(4-[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino)-butyl) ester sodium salt (0.49 g, 34%) as a red powder.

20 **Example 25: Antiproliferative Activity**

The antiproliferative activity of the compounds of the invention is demonstrated below. These effects indicate that the compounds of the present invention are useful in treating cancer, in particular solid tumors such as breast and colon tumors.

25 **MDAMB-435 Cell-Based Assay**

The epithelial breast carcinoma cell line (MDAMB-435) was purchased from ATCC (American Type Cell Culture Collection) and was grown in culture in medium
30 as recommended by ATCC. For analysis of the effect of various compounds of formula I on the growth of these cells, the cells were plated at a concentration of

134

149

1500 cells/well in a 96 well tissue culture plate ("test plate"). The day after the cells were plated, the compounds to be analyzed were dissolved in 100% DMSO (dimethyl sulfoxide) to yield at 10mM stock solution. Each compound was diluted in H₂O to 1mM and was added to triplicate wells in the first row of a 96 well master plate containing medium to yield a final concentration of 40μM. The compounds were then serially diluted in medium in the "master plate". The diluted compound(s) were then transferred to test plates containing cells. A row of vehicle "control cells" received DMSO. The final concentration of DMSO in each well was 0.1%. 5 days post drug addition, the plate was analyzed as described below.











MTT (3-(4-5 methyl thiazole-2-yl)-2,5-diphenyl tetrazolium bromide; thiazolyl blue) was added to each well to yield a final concentration of 1mg/ml. The plate was then incubated at 37°C for 2 1/2 - 3 hours. The MTT- containing medium was then removed and 50μl of 100% ethanol was added to each well to dissolve the formazan. The absorbences were then read using an automated plate reader (Bio-tek microplate reader). IC₅₀'s were calculated using the Reed and Munsch equation, see Am. J. Hygiene Vol. 27 pgs. 493-497, 1938.

The results of the foregoing *in vitro* experiments are set forth in Table I below.

Each of the compounds in Table I had an IC₅₀ ≤ 1.00 μM.

185
150












TABLE I

		
Example	R	Scheme
6b		1a
7		1a
8b, 11		1a
8a, 9, 10		1a
13		1b
12		1b
2		1b
3a		1b
3f		1b

60158860.101299












156
151

TABLE I (Continued)

		
Example	R	Scheme
5		1a
3g		1b
4		1b
3h		1b
6a		1a
3b		1a
3c		1b
3d		1b
3e		1b
15		1b
14		1b

152
187












TABLE I (Continued)

		
Example	R	Scheme
16		3
17		4
18		4
19		4
20		4
1	CH2OPO3Na2	2
21		5
22		5
23		5
24		5
		5

662101-09885109

189
154








TABLE II

				
Example	R	Stability	Plasma Conversion	Scheme
8a, 9, 10		Good	30% @30min	1a
13		Good	N/D*	1b
12		Fair	60% @30min	1b
3a		Good	70% @30min	1b
3f		Good	20% @30min	1b
3g		Good	N/D*	1b
3h		Fair	N/D*	1b
3b		Good	80% @30min	1a
3c		Good	60% @30min	1b
3d		Good	70% @30min	1b

60158860-101299

260
155

TABLE II (Continued)

				
Example	R	Stability	Plasma Conversion	Scheme
15		Good	100% @2hrs	1b
14		Good	5% @4hrs	1b
17		Good	15% @6hrs	4
20		Good	N/D*	4
1	CH ₂ OPO ₃ Na ₂	Good	100% @30min	2
21		Fair	100% @1hr	5
22		Good	N/D*	5
24		Good	10% @6hrs	5
		Good	N/D*	5

* N/D = Not Done

662707-09885109

164
156

Example 26: Tablet Formulation

Item	Ingredients	Mg/Tablet					
1	Compound A *	5	25	100	250	500	750
2	Anhydrous Lactose	103	83	35	19	38	57
3	Croscarmellose Sodium	6	6	8	18	32	48
4	Povidone K30	5	5	6	12	24	36
5	Magnesium Stearate	1	1	1	3	6	9
	Total Weight	120	120	150	300	600	900

*Compound A represents a compound of the invention.

5 Manufacturing Procedure:

1. Mix Items 1, 2 and 3 in a suitable mixer for 15 minutes.
2. Granulate the powder mix from step 1 with 20% Povidone K30 Solution (Item 4).
3. Dry the granulation from step 2 at 50°C.
4. Pass the granulation from step 3 through a suitable milling equipment.
5. Add the item 5 to the milled granulation step 4 and mix for 3 minutes.
6. Compress the granulation from step 5 on a suitable press.

Example 27: Capsule Formulation

Item	Ingredients	mg/Capsule				
1	Compound A *	5	25	100	250	500
2	Anhydrous Lactose	159	123	148	—	—
3	Corn Starch	25	35	40	35	70
4	Talc	10	15	10	12	24
5	Magnesium Stearate	1	2	2	3	6
	Total Fill Weight	200	200	300	300	600

* Compound 1 represents a compound of the invention.

162
157

Manufacturing Procedure:

1. Mix items 1, 2 and 3 in a suitable mixer for 15 minutes.
2. Add items 4 & 5 and mix for 3 minutes.
3. Fill into a suitable capsule.

5

Example 28: Injection Solution/Emulsion Preparation

Item	Ingredient	mg/mL
1	Compound A *	1 mg
2	PEG 400	10-50 mg
3	Lecithin	20-50 mg
4	Soy Oil	1-5 mg
5	Glycerol	8-12 mg
6	Water q.s.	1 mL

* Compound A represents a compound of the invention.

10

Manufacturing Procedure:

1. Dissolve item 1 in item 2.
2. Add items 3, 4 and 5 to item 6 and mix until dispersed, then homogenize.
3. Add the solution from step 1 to the mixture from step 2 and homogenize until the dispersion is translucent.
- 15 4. Sterile filter through a 0.2 um filter and fill into vials.

662101.058 ST09

163
158

Example 29: Injection Solution/Emulsion Preparation

Item	Ingredient	mg/mL
1	Compound A *	1 mg
2	Glycofurol	10-50 mg
3	Lecithin	20-50 mg
4	Soy Oil	1-5 mg
5	Glycerol	8-12 mg
6	Water	q.s. 1 mL

* Compound A represents a compound of the invention.

5 Manufacturing Procedure:

1. Dissolve item 1 in item 2.
2. Add items 3, 4 and 5 to item 6 and mix until dispersed, then homogenize.
3. Add the solution from step 1 to the mixture from step 2 and homogenize until the dispersion is translucent.
- 10 4. Sterile filter through a 0.2 um filter and fill into vials.

While the invention has been illustrated by reference to specific and preferred embodiments, those skilled in the art will understand that variations and modifications may be made through routine experimentation and practice of the invention. Thus, the invention is intended not to be limited by the foregoing description, but to be defined by the appended claims and their equivalents.


662101-0985109

164
159

CLAIMS

What is Claimed is:

1. A compound having the formula



and pharmaceutically active salts of said compounds, wherein:

R is selected from the group consisting of $-CH_2OPO_3R^1R^2$, $-CH_2OCOR^3$, $-CH_2OCO_2R^3$, $-CH_2OCONHR^3$, and $-CONHR^3$;

R^1 and R^2 are selected from the group consisting of H, Na and NH_4 and are the same unless either R^1 or R^2 is H, in which case the other can be different, or alternatively, R^1 and R^2 together represent calcium.

R^3 is selected from the group consisting of alkyl which optionally may be substituted by one or more substituents selected from the group consisting of $-CO_2R^4$, $-NR^5R^6$, polyethylene glycol, $-OPO_3R^1R^2$, hydroxy, alkoxy and aryl; alkenyl which optionally may be substituted by one or more substituents selected from the group consisting of $-CO_2R^4$, $-NR^5R^6$, polyethylene glycol, $-OPO_3R^1R^2$, hydroxy, alkoxy and aryl; cycloalkyl which optionally may be substituted by one or more substituents selected from the group consisting of $-CO_2R^4$, $-NR^5R^6$, polyethylene glycol, $-OPO_3R^1R^2$, hydroxy, alkoxy and aryl;

T63
160

heterocycle, which when including N as a heteroatom, the N optionally may be substituted with the group consist of lower alkyl and $-\text{COR}^7$,

aryl which optionally may be substituted by one or more substituents
5 selected from the group consisting of CO_2R^4 , hydroxy, alkoxy, polyethylene glycol, $\text{OPO}_3\text{R}^1\text{R}^2$, and alkyl which itself may be substituted with hydroxy alkoxy, carboxy and substituted amino, provided that when aryl represents pyridine, the nitrogen may be substituted with lower alkyl;

10 R^4 is selected from the group consisting of H, Na and lower alkyl;

R^5 and R^6 are each independently selected from the group consisting of H, lower alkyl, and $-\text{COR}^7$, or alternatively, the group $-\text{NR}^5\text{R}^6$ together form a 5 or 6 membered heterocyclic ring; and

15 R^7 is lower alkyl which optionally may be substituted with carboxy, polyethylene glycol polyethylene glycol, and substituted amino.

20 2. The compound of claim 1 wherein R is selected from the group consisting of $-\text{CH}_2\text{OPO}_3\text{R}^1\text{R}^2$ and $-\text{CH}_2\text{OCOR}^3$.

25 3. The compound of claim 1 wherein R is $-\text{CH}_2\text{OCO}$ -pyridine, wherein the N atom on the pyridine is substituted with lower alkyl.

4. The compound of claim 3 wherein the N atom on the pyridine is substituted with methyl or ethyl.

30

662101-09885109

186
16

5. The compound of claim 1 wherein R^1 and R^2 are independently selected from the group consisting of H and Na.

5 6. The compound of claim 1 wherein R^3 is heterocycle containing at least one nitrogen atom that optionally may be substituted with $-COR^7$.

7. The compound of claim 1 wherein R^3 is aryl which is substituted with
10 $-OPO_3R^1R^2$, and R^1 and R^2 are independently selected from H and Na.

8. The compound of claim 1 wherein R^3 is aryl which is substituted by the
15 group consisting of $-CO_2Na$, polyethylene glycol and $-CH_2CH_2N(CH_2CH_2)_2$.

9. The compound of claim 1 wherein R^3 is lower alkyl which is substituted
20 with $-CO_2Na$.

10. The compound of claim 1 wherein the group NR^5R^6 together forms a 5
or 6 membered heterocyclic ring.

25 11. The compound of claim 10 wherein the group NR^5R^6 is selected from piperidine or pyrrolidine.

12. The compound of claim 1 wherein R^5 and R^6 are each independently
30 selected from the group consisting of H, methyl and ethyl.

662101-09885109

167
162

13. The compound of claim 1 wherein R⁷ is ethyl which is substituted with polyethylene glycol.

5

14. The compound of claim 1 wherein R is -CH₂OCOR³ and R³ is ethyl which is substituted with polyethylene glycol having a molecular weight of from about 750 to about 5000 Daltons.

10

15. The compound of claim 1 wherein the polyethylene glycol has a molecular weight of from about 750 to about 5000 Daltons.

15

16. A compound selected from the group consisting of:

Phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl]sodium salt,

20

Isonicotinic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

But-2-enedioic acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl] ester,

Decanoic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

25

3-amino-cyclohexanecarboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester trifluoroacetate,

60158860.101299

Piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester acetate salt,

Piperidine-4-carboxylic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester hydrochloride,

5 2-amino-2-methyl-propionic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester hydrochloride,

1-Methyl-3-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethoxycarbonyl]-pyridinium; trifluoroacetate, and

10 O-[2-[[4-[[[2,5-Dihydro-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-1H-pyrrol-1-yl]methoxycarbonyl]-1-piperidinyl]carbonyl]ethyl]-O'-methylpolyethylene glycol 1000.

17. A compound selected from the group consisting of:

13 4-Phosphonooxy-benzoic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester monosodium salt,

[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-yl]methoxycarbonyl]-4-phenyl-O-methylpolyethylene glycol 500,

20 Carbonic acid allyl ester 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

(2-Dimethylamino-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

25 [2-(2-Hydroxy-ethoxy)-ethyl]-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

(2-Hydroxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

(2-phosphonooxy-ethyl)-carbamic acid 3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethyl ester,

464
164

3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (2-diethylamino-ethyl)-amide,

3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (6-dimethylamino-hexyl)-amide,

5 3-(1-Methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carboxylic acid (4-hydroxybutyl)-amide, and

Phosphoric acid mono-(4-[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino]-butyl) ester sodium salt.

10

18. A compound selected from the group consisting of:

phosphoric acid mono-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethoxy]sodium salt,

15

O-[2-[[2,5-dihydro-3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-yl]methoxycarbonyl]ethyl]-O'-methylpolyethylene glycol 2000, and

20

phosphoric acid mono-(4-[[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrole-1-carbonyl]-amino]-butyl) ester sodium salt, and 1-methyl-3-[3-(1-methyl-1H-indol-3-yl)-4-(1-methyl-6-nitro-1H-indol-3-yl)-2,5-dioxo-2,5-dihydro-pyrrol-1-ylmethoxycarbonyl]-pyridinium trifluoroacetate.

25

19. A pharmaceutical composition comprising as an active ingredient an effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier or excipient.

30

20. The pharmaceutical composition of claim 19 which is suitable for parenteral administration.

60458860.101299

170
165

21. A method for treating a cell proliferative disorder comprising
administering to a patient in need thereof a therapeutically effective amount of a
5 compound according to claim 1.

22. The method of claim 21 wherein the cell proliferative disorder is
cancer.
10

23. The method of claim 22 wherein the cancer is a solid tumor.


24. The method of claim 22 wherein the cancer is breast, colon, or lung
cancer.
15
20

6015 860-101299

127
166

ABSTRACT

Disclosed are novel substituted pyrroles having the formula



These compounds and their pharmaceutically acceptable salts are suitable for administration to patients as continuous infusion solution and are useful in the treatment and/or control of cell proliferative disorders, in particular cancer. Also disclosed are pharmaceutical compositions containing the foregoing compounds and methods for the treatment and/or control of cancer.

67163

662707-09885709

172
167

Declaration and Power of Attorney for Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

SUBSTITUTED PYRROLES SUITABLE FOR CONTINUOUS INFUSION
the specification of which

(check one)

☒ is attached hereto.

☐ was filed on _____ as

☐ Application Serial No. _____

☐ and was amended on _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Claimed

_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	☐ Yes	☐ No
_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	☐ Yes	☐ No
_____ (Number)	_____ (Country)	_____ (Day/Month/Year Filed)	☐ Yes	☐ No

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

_____ (Application No.)	_____ (Filing Date)	_____ (Application No.)	_____ (Filing Date)
----------------------------	------------------------	----------------------------	------------------------

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.


POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

George W. Johnston	(Reg. No. 28090)	Dennis P. Tramaloni	(Reg. No. 28542)
William H. Epstein	(Reg. No. 20008)	Patricia S. Rocha-Tramaloni	(Reg. No. 31054)
Bileen M. Ebel	(Reg. No. 37316)		

Sole Correspondence to:
George W. Johnston, Esq., Hoffmann-La Roche Inc., 340 Kingsland Street,
Nutley, New Jersey 07110-1199

Direct Telephone Calls to: (name and telephone number)
Patricia S. Rocha-Tramaloni (973) 235-2441

Nader Fotouhi
Full name of sole or first inventor

 10/11/99
Inventor's signature Date

Chatham, Morris County, New Jersey
Residence

France
Citizenship

1 Daniel Street, Chatham, New Jersey 07928
Post Office Address

177
169

Norman Kong

Full name of second joint inventor, if any

Norman Kong
Second Inventor's signature

October 11, 1999
Date

West Caldwell, Essex County, New Jersey
Residence

Canada
Citizenship

12 Klimback Court, West Caldwell, New Jersey 07006
Post Office Address

in
Guangy Aijun Liu

Full name of third joint inventor, if any

Guangy Aijun Liu
Third Inventor's signature

October 11, 1999
Date

in
Nutley, Essex County, New Jersey
Residence

United States of America
Citizenship

193 Alexander Avenue, Nutley, New Jersey 07110
Post Office Address

Allen John Lovey

Full name of fourth joint inventor, if any

Allen John Lovey
Fourth Inventor's signature

October 11, 1999
Date

North Caldwell, Essex County, New Jersey
Residence

United States of America
Citizenship

21 Hickory Drive, North Caldwell, New Jersey 07006
Post Office Address

170

John Guilfoyle Mullin, Jr.

Full name of fifth joint inventor, if any

John Guilfoyle Mullin, Jr.

Fifth Inventor's signature

October 8, 1989

Date

Hawthorne, Passaic County, New Jersey

Residence

United States of America

Citizenship

519 Goffle Hill Road, Hawthorne, New Jersey 07506

Post Office Address

66101-093850-101299

Title 37, Code of Federal Regulations, §1.56, duty to disclose information material to patentability provides, in part, that each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned.

Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim; or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office, or
 - (ii) Asserting an argument of patentability.