

Señores
DIVISION DE NUEVAS CREACIONES
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
Bogotá, D C.

SUPERINDUSTRIA Y COMERCIO
Radicación : 0000427 0000001 Folios: 20
Fecha (R2): 2023-11-30 10:12:03
Trámite : 022 PATENTES 0 1 REGISTRADO 329 CANCELADO
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Re Código 02-01-329
Expediente No 00-83 427
Solicitud de Patente
SMITHKLINE BEECHAM
CORPORATION

En relación con el Expediente de la referencia y con el fin de dar cumplimiento a la preceptuado por el artículo 4o literal d) numeral 3 de la Convención de París adjunto a este escrito sírvanse encontrar la copia certificada de la solicitud de patente de la cual se reivindica prioridad, la cual se encuentra debidamente traducida en sus apartes pertinentes

De Ustedes Atentamente,

GERMAN CASTILLO GRAU
T P No 2978 del C S J



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

October 30, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/163,158

FILING DATE: *November 02, 1999*



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

H. L. Jackson

H. L. JACKSON

Certifying Officer

EXPRESS MAIL CERTIFICATE

Express Mail® Mailing Label Number HL175492139US

Date Of Deposit: 22 November 1999 That This Paper Or Fee Is Being Deposited With The United States Postal Service Express Mail Post Office To Addressee's Service Under 37 CFR 1.10 On The Date Indicated Above And Is Addressed To: Assistant Commissioner for Patents, Washington, D.C. 20231

Name Of Person Mailing Paper Or Fee

(Type Or Print)

Signature

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION for PATENT under 37 CFR 1.53(c).

Docket No.			P51051P
INVENTOR(s) / APPLICANT(s)			
Last Name	First Name	Middle Initial	Residence (City and Either State or Foreign Country)
Goodfellow	Peter	N	Belsize Park, London NW3 2NR
Nieman	Richard		Philadelphia, PA
Torphy	Theodore	J.	Bryn Mawr, PA

TITLE OF THE INVENTION (280 characters max)

METHOD AND COMPOSITION FOR TREATING PULMONARY DISEASES

CORRESPONDENCE ADDRESS

SMITHKLINE BEECHAM CORPORATION

Corporate Intellectual Property - UW2220

709 Swedeland Road

King of Prussia

Telephone No. 610-270-5014

Facsimile No. 610-270-5090

State	PA	Zip Code	19406-2799	Country	United States of America
-------	----	----------	------------	---------	--------------------------

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification	Number of Pages	12	☐ Small Entry Statement
☒ Figure	Number of Sheets	1	☒ Other (specify)
1 page abstract			

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

☒ The Commissioner is hereby authorized to charge filing fees and credit Deposit Account No. 19-2570	PROVISIONAL FILING FEE AMOUNT (\$)	\$150.00
--	------------------------------------	----------

Respectfully submitted,
Signature:

James M. Kanagy

Date:

Registration No.

22 Nov 1999
29,550☐ Additional inventors are being named on separately numbered sheets attached hereto.

PROVISIONAL APPLICATION FILING ONLY

SEND TO: Assistant Commissioner for Patents, Box Provisional Application, Washington, D.C. 20231

NAMEKPATAFP51051provision.doc

80
27

Method and Compositions for Treating Pulmonary Diseases

Area of the Invention

This invention relates compositions and methods for preventing or reducing the onset of symptoms of pulmonary diseases, or treating or reducing the severity of pulmonary diseases. In particular it relates to compositions and methods for treating pulmonary diseases mediated by phosphodiesterase 4 (PDE4) by administering a PDE4 inhibitor with an anti-inflammatory corticosteroid.

Background of the Invention

Identification of novel therapeutic agents for treating pulmonary diseases is made difficult by the fact that multiple mediators are responsible for the development of the disease. Thus, it seems unlikely that eliminating the effects of a single mediator could have a substantial effect on all three components of chronic asthma. An alternative to the "mediator approach" is to regulate the activity of the cells responsible for the pathophysiology of the disease.

One such way is by elevating levels of cAMP (adenosine cyclic 3',5'-monophosphate). Cyclic AMP has been shown to be a second messenger mediating the biologic responses to a wide range of hormones, neurotransmitters and drugs; [Krebs Endocrinology Proceedings of the 4th International Congress Excerpta Medica, 17-29, 1973]. When the appropriate agonist binds to specific cell surface receptors, adenylate cyclase is activated, which converts Mg^{+2} -ATP to cAMP at an accelerated rate. Cyclic AMP modulates the activity of most, if not all, of the cells that contribute to the pathophysiology of extrinsic (allergic) asthma. As such, an elevation of cAMP would produce beneficial effects including: 1) airway smooth muscle relaxation, 2) inhibition of mast cell mediator release, 3) suppression of neutrophil degranulation, 4) inhibition of basophil degranulation, and 5) inhibition of monocyte and macrophage activation. Hence, compounds that activate adenylate cyclase or inhibit phosphodiesterase should be effective in suppressing the inappropriate activation of airway smooth muscle and a wide variety of inflammatory cells. The principal cellular mechanism for the inactivation of cAMP is hydrolysis of the 3'-phosphodiester bond by one or more of a family of isozymes referred to as cyclic nucleotide phosphodiesterases (PDEs).

It has been shown that a distinct cyclic nucleotide phosphodiesterase (PDE) isozyme, PDE IV, is responsible for cAMP breakdown in airway smooth muscle and inflammatory cells. [Torphy, "Phosphodiesterase Isozymes Potential Targets for Novel Anti-asthmatic Agents" in New Drugs for Asthma, Barnes, ed. IBC Technical Services Ltd., 1989]. Research indicates that inhibition of this enzyme not only produces airway smooth muscle relaxation, but also suppresses degranulation of mast cells, basophils and neutrophils along with inhibiting the activation of monocytes and neutrophils. Moreover, the beneficial effects of PDE IV inhibitors are markedly potentiated when adenylate cyclase

2025-05-10

31
28

activity of target cells is elevated by appropriate hormones or autocooids, as would be the case *in vivo*. Thus PDE IV inhibitors would be effective in the lung, where levels of prostaglandin E₂ and prostacyclin (activators of adenylate cyclase) are elevated. Such compounds would offer a unique approach toward the pharmacotherapy of bronchial asthma and possess significant therapeutic advantages over agents currently on the market.

In addition, it could be useful to combine therapies in light of the fact that the etiology of many pulmonary diseases involves multiple mediators. In this invention there is presented the combination of a PDE 4 inhibitor and an anti-inflammatory corticosteroid, particularly one delivered by inhalation, for treating pulmonary diseases. This combination is particularly useful for treating chronic obstructive pulmonary disease (COPD) or asthma.

Summary of the Invention

In a first aspect this invention relates to a method for treating a pulmonary disease in a mammal by administering to a patient in need thereof an effective amount of a PDE 4-specific inhibitor and an effective amount of a steroidal anti-inflammatory agent wherein the drugs are administered concomitantly together or separately and sequentially where the sequential administration is close in time or remote in time.

In a second aspect this invention relates to a composition for treating a pulmonary disease in a mammal comprising an effective amount of a PDE4-specific inhibitor, an effective amount of a steroidal anti-inflammatory agent and a pharmaceutically acceptable excipient.

Detailed Description of the Invention

The combination therapy contemplated by this invention comprises administering a PDE4 inhibitor with a steroidal anti-inflammatory agent to prevent onset of a pulmonary disease event or to treat an existing condition. The compounds may be administered together in a single dosage form. Or they may be administered as two different formulations which may be the same or different. To illustrate, both drugs may be provided separately as oral formulations, or one may be an oral preparation and the other as an inhalant, or both may be provided in an inhaled dose form. They may be administered at the same time. Or they may be administered either close in time or remotely, such as where one drug is administered in the morning and the second drug is administered in the evening.

The combination may be used prophylactically or after the onset of symptoms. In some instances the combination(s) may be used to prevent the progression of a pulmonary disease or to arrest the decline of a function, such as lung function.

The PDE4-specific inhibitor useful in this invention may be any compound that is known to inhibit the PDE4 enzyme or which is discovered to act in as PDE4 inhibitor, and which are only PDE4 inhibitors, not compounds which inhibit other members of the PDE family as well as PDE4. Generally it is preferred to use a PDE4 antagonists which has an IC₅₀ ratio of about 0.1 or greater as regards the IC₅₀ for the PDE IV catalytic form which

0163158.110271

P51051P

binds rolipram with a high affinity divided by the IC_{50} for the form which binds rolipram with a low affinity

PDE inhibitors used in treating inflammation and as bronchodilators, drugs like theophylline and pentoxifyllin, inhibit PDE isozymes indiscriminately in all tissues. These compounds exhibit side effects, apparently because they non-selectively inhibit all 5 PDE isozyme classes in all tissues. The targeted disease state may be effectively treated by such compounds, but unwanted secondary effects may be exhibited which, if they could be avoided or minimized, would increase the overall therapeutic effect of this approach to treating certain disease states. For example, clinical studies with the selective PDE 4 inhibitor rolipram, which was being developed as an antidepressant, indicate it has psychotropic activity and produces gastrointestinal effects, e.g., pyrosis, nausea and emesis.

For purposes of this disclosure, the cAMP catalytic site which binds R and S rolipram with a low affinity is denominated the "low affinity" binding site (LPDE 4) and the other form of this catalytic site which binds rolipram with a high affinity is denominated the "high affinity" binding site (HPDE 4). This term "HPDE4" should not be confused with the term "hPDE4" which is used to denote human PDE4.

Initial experiments were conducted to establish and validate a [^3H]-rolipram binding assay. Details of this work are given in Example 1 below.

To determine whether both the high affinity binding activity and the low affinity binding activity resided in the same gene product, yeast were transformed by known methods and the expression of recombinant PDE 4 was followed over a 6 hour fermentation period. Western blot analysis using an antibody directed against PDE 4 indicated that the amount of PDE 4 expressed increased with time, reaching a maximum after 3 hours of growth. In addition, greater than 90% of the immunoreactive product was in the high speed (100,000 x g) supernatant of yeast lysates. [^3H]R-(-)-Rolipram binding and PDE activity were monitored along with protein expression. PDE 4 activity was co-expressed with rolipram binding activity, indicating that both functions exist on the same gene product. Similar to results with the Western blot analysis, greater than 85% of the rolipram-inhibitable PDE activity and [^3H]-rolipram binding activity was found to be present in the yeast supernatant fraction.

Overall, most of the recombinant PDE 4 expressed in this system exists as LPDE 4 and only a small fraction as HPDE 4. Consequently, inhibition of recombinant PDE 4 catalytic activity primarily reflects the actions of compounds at LPDE 4. Inhibition of PDE 4 catalytic activity can thus be used as an index of the potency of compounds at LPDE 4. The potency of compounds at HPDE 4 can be assessed by examining their ability to compete for [^3H]R-rolipram. To develop SARs for both the low affinity and high affinity rolipram binding sites, the potencies of selected compounds were determined in two assay systems. Results from experiments using standard compounds were tabulated. As

33/9/95
30

expected, certain compounds were clearly more potent in competing with [³H]-rolipram at the site for which rolipram demonstrated high affinity binding as compared with the other site, the one at which rolipram is a low affinity binder. SAR correlation between high affinity binding and low affinity binding was poor and it was concluded that the SAR for inhibition of high affinity [³H]-rolipram binding was distinct from the SAR for binding to the low affinity rolipram binding site. Table I provides results from this SAR work.

It is now known that there are at least two binding forms on human monocyte recombinant PDE 4 (hPDE 4) with which inhibitors interact. One explanation for these observations is that hPDE 4 exists in two distinct forms. One binds the likes of rolipram and denbufylline with a high affinity while the other binds these compounds with a low affinity. The preferred PDE4 inhibitors of use in this invention will be those compounds which have a salutary therapeutic ratio, i.e., compounds which preferentially inhibit cAMP catalytic activity where the enzyme is in the form that binds rolipram with a low affinity, thereby reducing the side effects which apparently are linked to inhibiting the form which binds rolipram with a high affinity. Another way to state this is that the preferred compounds will have an IC₅₀ ratio of about 0.1 or greater as regards the IC₅₀ for the PDE 4 catalytic form which binds rolipram with a high affinity divided by the IC₅₀ for the form which binds rolipram with a low affinity.

A further refinement of this standard is that of one wherein the PDE4 inhibitor has an IC₅₀ ratio of about 0.1 or greater; said ratio is the ratio of the IC₅₀ value for competing with the binding of 1nM of [³H]R-rolipram to a form of PDE 4 which binds rolipram with a high affinity over the IC₅₀ value for inhibiting the PDE IV catalytic activity of a form which binds rolipram with a low affinity using 1 microm[³H]-cAMP as the substrate. A further review explanation with of this test can be found in co-pending U.S. application 08/456274 filed 31 May 1995, the text of which is incorporated herein by reference to the extent that text is necessary to the practice of this invention.

Examples of useful PDE4 inhibitors are.

(R)-(+)-1-(4-bromobenzyl)-4-[(3-cyclopentylloxy)-4-methoxyphenyl]-2-pyrrolidone,

(R)-(+)-1-(4-bromobenzyl)-4-[(3-cyclopentylloxy)-4-methoxyphenyl]-2-pyrrolidone,

3-(cyclopentylloxy-4-methoxyphenyl)-1-(4-N'-[N2-cyano-S-methyl-isothiourea]benzyl)-2-pyrrolidone,

cis-4-cyano-4-(3-cyclopentylloxy-4-methoxyphenyl)cyclohexan-1-carboxylic acid],

cis-[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-

ol],

(R)-(+)-ethyl [4-(3-cyclopentylloxy-4-methoxyphenyl)pyrrolidino-2-ylidene]acetate;

(S)-(-)-ethyl [4-(3-cyclopentylloxy-4-methoxyphenyl)pyrrolidino-2-ylidene]acetate,

31

Most preferred are those PDE4 inhibitors which have an IC₅₀ ratio of greater than 0.5, and particularly those compounds having a ratio of greater than 1.0. Preferred compounds are *is* 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-carboxylic acid, 2-carbomethoxy-4-cyano-4-(3-cyclopropylmethoxy-4-

5 difluoromethoxyphenyl)cyclohexan-1-one, and *is*-[4-cyano-4-(3-cyclopropylmethoxy-4-difluoromethoxyphenyl)cyclohexan-1-ol]; these are examples of compounds which bind preferentially to the low affinity binding site and which have an IC₅₀ ratio of 0.1 or greater.

Compounds set out in U.S. patent 5,552,438 issued 03 September, 1996. This patent and the compounds it discloses are incorporated herein in full by reference. The
10 compound of particular interest, which is disclosed in U.S. patent 5,552,438, is *cis*-4-cyano-4-[3-(cyclopentyloxy)-4-methoxyphenyl]cyclohexane-1-carboxylic acid and its salts, esters, pro-drugs or physical forms.

AWD-12-281 from Astra (Hofgen, N *et al.* 15th EFMC Int Symp Med Chem (Sept 6-10, Edinburgh) 1998, Abst P.98), a 9-benzyladenine derivative nominated NCS-613
15 (INSERM); D-4418 from Chiroscience and Schering-Plough, a benzodiazepine PDE4 inhibitor identified as CI-1018 (PD-168787, Parke-Davis/Warner-Lambert); a benzodioxole derivative Kyowa Hakko disclosed in WO 9916766; V-11294A from Napp (Landella, L.J. *et al.* Eur Resp J [Ann Cong Eur Resp Soc (Sept 19-23, Geneva) 1998] 1998, 12(Suppl 28): Abst P2393); rofinumilast (CAS reference No 162401-32-3) and a phthalazone (WO 9947505) from Byk-Gulden; or a compound identified as T-440 (Tanabe Selyaku; Fuji, K. *et al.* *J Pharmacol Exp Ther*, 1998, 284(1) 162) Any one or all of these compounds may or could benefit from the process described herein.

The several specific compounds set out above which do not have a generic or trade name can be made by the processes described in co-pending U.S. patent applications USSN
25 862,083 filed 30 October 1992, USSN 862, 111 filed 30 October 1992; USSN 862,030 filed 30 October 1992, and USSN 862,114 filed 30 October 1992 or their progeny or U.S. patent(s) claiming priority from one or more of these applications. Each of these applications or related patents is incorporated herein by reference in full as if set out in this document.

30 The steroid agents useful in this invention are oral and inhaled corticosteroids and their pro-drugs which have anti-inflammatory activity. Examples of these steroids are methyl prednisolone, prednisone, dexamethasone, fluticasone, beclomethasone, budesonide, flunisolide, mometasone furoate, and triamcinolone acetonide. Methyl prednisolone and prednisone are oral and
35 injectable forms of anti-inflammatory corticosteroids, they are available from numerous branded and generic pharmaceutical companies. Beclomethasone dipropionate is sold as an aerosol for inhalation under the names Beconase® and Beconase AQ® by Glaxo Wellcome. Fluticasone propionate is sold under the name

35
32

- Flonase® by Glaxo Wellcome as well. Triamcinolone acetonide is sold by Rhone-Poulenc Rorer under the name Nasacort® as a nasal spray and aerosol. Flunisolide is sold as a nasal solution under the name Nasalide® and Nasarel™ by Roche Laboratories. Dexamethasone is sold as the sodium phosphate salt by Medeva Pharmaceuticals, Inc. under the name Dexacort™ Phosphate. Mometasone furoate is sold as the monohydrate as a nasal preparation by Schering Corp under the name Nasonex®. Budesonide is yet another inhaled corticosteroid used in treating pulmonary diseases. It is marketed by Astra Pharmaceuticals, L.P. as a powder in a turbobaler device under the name Pulmicort Turbobaler®. All of these drugs and nasal preparations or oral or injectable formulations can be found in the 1999 edition of the Physicians' Desk Reference® (PDR), published by Medical Economics Corporation, Inc. of New Jersey, USA and is available on the Internet at <http://www.tomescpa.com/frameMain.asp?Menu=0> and linked pages. Additional corticosteroids now under development and which could be used in this invention are set out in Table I.

Table I
Inhaled Corticosteroids

Drug	Company	Indication
mometasone furoate	Schering-Plough	asthma SAR
roflupenide	AstraZeneca	asthma
ciclesonide	Byk-Gulden/Recordati	asthma
butixocort propionate	Warner-Lambert/3M	asthma/rhinitis
RPR-106541	Rhone-Poulenc Rorer	asthma
ST-126	SSP/Torii	asthma

- A preferred combination therapy is that of one or more of dexamethasone, fluticasone, beclomethasone, budesonide, flunisolide, mometasone furoate, and triamcinolone acetonide administered with *cis* 4-cyano-4-(3-cyclopentyloxy-4-methoxyphenyl)cyclohexan-1-carboxylic acid (Aniflo®). A preferred therapy is concomitant administration of the steroid as an inhalant and the acid in an oral dose form, wherein each drug is administered once or twice a day. In regards to the acid, a controlled-release oral tablet is most preferred.

- It is contemplated that both active agents would be administered at the same time, or very close in time. Alternatively, one drug could be taken in the morning and one later in the day. Or in another scenario, one drug could be taken twice daily and the other once daily, either at the same time as one of the twice-a-day dosing occurred, or separately. Preferably both drugs would be taken together at the same time.

50153158-110277

36
33

The present compounds and pharmaceutically acceptable salts which are active when given orally can be formulated as syrups, tablets, capsules, controlled-release preparation or lozenges. A syrup formulation will generally consist of a suspension or solution of the compound or salt in a liquid carrier for example, ethanol, peanut oil, olive oil, glycerin or water with a flavoring or coloring agent.

Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used for preparing solid formulations may be used. Examples of such carriers include magnesium stearate, terra alba, talc, gelatin, acacia, stearic acid, starch, lactose and sucrose. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example using the aforementioned carriers in a hard gelatin capsule shell. Where the composition is in the form of a soft gelatin shell capsule any pharmaceutical carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous gums, celluloses, silicates or oils, and are incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of a compound or salt in a sterile aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil, for example polyethylene glycol, polyvinylpyrrolidone, lecithin, arachis oil or sesame oil.

Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be administered as a dry powder or in the form of an aerosol using a conventional propellant such as fluorinated hydrocarbons such as trichlorofluoromethane.

Preferably the composition for the PDE4 inhibitors is a unit dosage form such as a tablet or capsule. For the steroids a metered aerosol dose, metered dry powder inhaler or nasal spray is preferred.

The active ingredient may be administered from 1 to 6 times a day, sufficient to exhibit the desired activity. Preferably, the active ingredient is administered about once or twice a day, more preferably twice a day.

The present compounds are useful for the treatment of EIA, PIA and CIA, both as chronic conditions as well as intermittently, in anticipation of the stimulus in question. Preferably, the present compounds are used for long-term therapy.

As for the amount of drug administered, it is believed that for the PDE4 inhibitors will be administered in an amount of between 1 and 200 micrograms per day per adult human. Steroids can be administered in conformity with approved labeling.

37
34**Example 1 — Phosphodiesterase and Rolipram Binding Assays****Example 1A**

Isolated human monocyte PDE 4 and hrPDE (human recombinant PDE4) was determined to exist primarily in the low affinity form. Hence, the activity of test compounds against the low affinity form of PDE 4 can be assessed using standard assays for PDE 4 catalytic activity employing 1 microM [^{3} H]cAMP as a substrate (Torphy et al., *J of Biol. Chem.*, Vol. 267, No. 3 pp1798-1804, 1992).

Rat brain high speed supernatants were used as a source of protein and both enantiomers of [^{3} H]-rolipram were prepared to a specific activity of 25.6 Ci/mmol. Standard assay conditions were modified from the published procedure to be identical to the PDE assay conditions, except for the last of the cAMP- 50mM Tris HCl (pH 7.5), 5 mM MgCl₂, 50 microM 5'-AMP and 1 nM of [^{3} H]-rolipram (Torphy et al., *J of Biol. Chem.*, Vol. 267, No. 3 pp1798-1804, 1992). The assay was run for 1 hour at 30° C. The reaction was terminated and bound ligand was separated from free ligand using a Brandel cell harvester. Competition for the high affinity binding site was assessed under conditions that were identical to those used for measuring low affinity PDE activity, except that [^{3} H]-cAMP was not present.

Example 1B**Measurement of Phosphodiesterase Activity**

PDE activity was assayed using a [^{3} H]cAMP SPA or [^{3} H]cGMP SPA enzyme assay as described by the supplier (Amersham Life Sciences). The reactions were conducted in 96-well plates at room temperature, in 0.1 ml of reaction buffer containing (final concentrations) 50 mM Tris-HCl, pH 7.5, 8.3 mM MgCl₂, 1.7 mM EGTA, [^{3} H]cAMP or [^{3} H] cGMP (approximately 2000 dpm/pmol), enzyme and various concentrations of the inhibitors. The assay was allowed to proceed for 1 hr and was terminated by adding 50 ul of SPA yttrium silicate beads in the presence of zinc sulfate. The plates were shaken and allowed to stand at room temperature for 20 min. Radiolabeled product formation was assessed by scintillation spectrometry. Activities of PDE3 and PDE7 were assessed using 0.05 uM [^{3} H]cAMP, whereas PDE4 was assessed using 1 uM [^{3} H]cAMP as a substrate. Activity of PDE1B, PDE1C, PDE2 and PDE5 activities were assessed using 1uM [^{3} H]cGMP as a substrate.

[^{3} H]R-rolipram binding assay

The [^{3} H]R-rolipram binding assay was performed by modification of the method of Schneider and co-workers, see Nicholson, et al., *Trends Pharmacol. Sci.*, Vol. 12, pp.19-27 (1991) and McHale et al., *Mol. Pharmacol.*, Vol. 39, 109-113 (1991). R-rolipram binds to the catalytic site of PDE4 see Torphy et al., *Mol. Pharmacol.*, Vol. 39, pp. 376-384 (1991). Consequently, competition for [^{3} H]R-rolipram binding provides an independent confirmation of the PDE4 inhibitor potencies of unlabeled competitors. The assay was

38 21
35

performed at 30°C for 1 hr in 0.5 ul buffer containing (final concentrations) 50 mM Tris-HCl, pH 7.5, 5 mM MgCl₂, 0.05% bovine serum albumin, 2 nM [³H]-R-rolipram (5.7 x 10⁴ dpm/pmol) and various concentrations of non-radiolabeled inhibitors. The reaction was stopped by the addition of 2.5 ml of ice-cold reaction buffer (without [³H]-R-rolipram) and rapid vacuum filtration (Brandel Cell Harvester) through Whatman GF/B filters that had been soaked in 0.3% polyethylenimine. The filters were washed with an additional 7.5-ml of cold buffer, dried, and counted via liquid scintillation spectrometry

Example 2

Low dose Inhaled Corticosteroid (ICS) Study 005, Dose-ranging Study

Study Design

- R, DB, PC, DR in patients with mild/moderate asthma.
- Ariflo[®] was dosed at 5 mg, 10 mg 15 mg twice daily for 6 weeks.
- Subjects were on a median of 500 mcg of beclomethasone equivalent although the mean dose of ICS was 652 mcg.
- Primary endpoint: change from baseline to endpoint in trough clinic FEV₁, changes in clinic FEV₁ every week and over a 4 hour period following first dose of double-blind medication.
- Secondary endpoints: use of rescue medicines and overnight symptoms.
- Tertiary endpoints: clinic FVC, PEFR, PEF25-75, PEF 75, domiciliary PEFR variability, domiciliary PEFR, summary symptom scores.
- One week run in.

Study Results:

- No statistical improvement in trough FEV₁ at endpoint ITT analysis 15 mg BID vs placebo (0.16 L, p=0.062)
- Response was dose-ordered for trough FEV₁ at endpoint ITT analysis
- Significant improvement in trough FEV₁ (0.21 L; p=0.033) if excluded patients on ICS doses > 500 mcg while on Ariflo 15 mg BID
- Corroborating support for 15 mg BID included 4 hour FEV₁, domiciliary PEFR, PEF25-75 and physician and patient global assessment scores

Example 3

High dose Inhaled Corticosteroid (ICS) Study

Study Design:

- R, DB, PC, DR in patients with mild/moderate asthma on ≥ 800 mcg of beclomethasone
- Ariflo[®] doses were 5, 10 and 15 mg twice daily for 4 weeks.
- 2 week run-in.
- Primary endpoint: trough clinic FEV₁.

39 

36

- Secondary endpoints: morning PEFR, symptoms, PEFR variability, evening PEFR, clinic PEFR, PEF25-75, PEF75, rescue medication used.

Study Results:

- No statistically significant change in clinic trough FEV₁ in any dose group was noted in the ITT analysis
- No dose ordering was observed for the primary endpoint
- Trough FEV₁ statistically improved at 10 mg BID using repeated measures analysis in patients excluded on < 800 ucg of ICS (0.16 L; p=0.009)
- ICS dose ranged from 100 to 4000 ucg and averaged 1136 ucg
- Corroborative support for the 10 mg BID as the effective dose also came from numerically superior results in the 4 hour clinic FEV₁, clinic FVC, PEFR
- No dose-response relationship could be determined, trough concentrations were dose proportional and similar to previous studies.

Table 1

Comparison between compounds in selected studies

Allergen challenge studies

Comparative Data: Allergen Challenge Studies

	1.3	12.3	10.7	22.9	+6.9
Montelukast	NONE	36.4	30.8	32	NO FALL
Budesonide	3.5	63.5	42.4	not done	
Both	NONE	56.9	42.6	not done	

- 20 **Legend:** Data on Ariflo was taken from the protocol analysis from study 005 in asthma patients on < 652 ucg of ICS. Data on montelukast (MT), budesonide (BDP) and both was

P51051P

40-14
37

taken from a single study in the Singular SBA that examined the effect of Singular relative to inhaled corticosteroids.

There are significant differences between the behavior of the Pbo groups in both studies. Conclusion is that Arflo outperformed Singular when added to low dose ICS,
5 despite the fact that the Pbo group also improved in the Arflo study

The foregoing statements and examples are intended to illustrate the invention, not to limit it. Reference is made to the claims for what is reserved to the inventors hereunder.

50153158-110277

AT
38

What is claimed is

1. A method for treating a pulmonary disease by administering to a patient in need thereof an effective amount of a PDE 4 inhibitor and an anti-inflammatory corticosteroid in a combined form, separately, or separately and sequentially where the sequential administration is close in time or remote in time.
2. The method of claim 1 wherein the PDE4 inhibitor is *cis* 4-cyano-4-(3-cyclopentyl-oxy-4-methoxyphenyl)cyclohexan-1-carboxylic acid and the steroid is selected from the group consisting of dexamethasone, fluticasone, beclomethasone, budesonide, flunisolide, mometasone furoate, and triamcinolone acetonide.

SECRET

#2 39 76


ABSTRACT OF THE DISCLOSURE

This invention relates to treating pulmonary diseases such as chronic obstructive pulmonary disease or asthma by administering a phosphodiesterase 4 inhibitor in combination with anti-inflammatory corticosteroid.

50153159.110277

Handwritten marks: a signature and the number 40.

Figure 1
Comparison between Singular® with ICS and Ariflo with ICS
Effect Size Across Studies, Ariflo and Singulair



CC2011-051051P

4/20
41



A TODOS A QUIENES LLEGUE EL PRESENTE:

DEPARTAMENTO DE COMERCIO DE LOS ESTADOS UNIDOS

Oficina de Marcas y Patentes

30 de octubre de 2000

LA PRESENTE CERTIFICA QUE LA ADJUNTA ES UNA COPIA TOMADA DE LOS REGISTROS DE LA OFICINA DE MARCAS Y PATENTES DE LOS ESTADOS UNIDOS DE LOS DOCUMENTOS DE LA SOLICITUD DE PATENTE ABAJO IDENTIFICADA Y LA CUAL CUMPLE CON LOS REQUISITOS PARA QUE LE SEA CONFERIDA UNA FECHA DE RADICACIÓN SEGÚN 35 USC 111

NÚMERO DE SOLICITUD 60/163 158

FECHA DE RADICACIÓN: 2 de noviembre de 1999

Por Autoridad del

COMISIONADO DE MARCAS Y PATENTES

(FIRMA) H L JACKSON

Oficial de Certificaciones

(SELLO DORADO DE OBLEA)

45
92

**MÉTODO Y COMPOSICIONES PARA EL TRATAMIENTO DE
ENFERMEDADES PULMONARES**

REIVINDICACIONES:

1. Un método para el tratamiento de una enfermedad pulmonar mediante la administración a un paciente que lo requiera, de una cantidad efectiva de un inhibidor PDE4 y un corticosteroide antiinflamatorio en forma combinada, separada o separada y secuencial, donde la administración secuencial es cercana en tiempo o remota en tiempo.

2. El método de la reivindicación 1, en el cual el inhibidor PDE4 es *cis* 4-clano-4-(3-ciclopentiloxi-4-metoxifenil)ciclohexan-1-ácido carboxílico y el esteroide es seleccionado del grupo compuesto por dexametasona, fluticasona, beclometasona, budesonida, flunisolida, furoato de mometasona y triamcinilona acetónida