

SEÑOR MINISTRO DE COMERCIO, INDUSTRIA Y TURISMO
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
DIVISION DE NUEVAS CREACIONES
E. S. D.

712
01

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
Rad: 00-083069- -00008-0000
Fec: 2007-04-04 08:52:54 Dep. 2020 NUECREACIONES
Trá. 2 PATENTES
EVE: 1 REGDEPOSITO
Act 444 RESPUESTA REQUE Folios: 3

EXPEDIENTE 00 083069

PATENTE DERIVADOS (TIONA) AMINOQUINAZOLINONA
SUSTITUIDOS O SALES DE LOS MISMOS, Y
CONTROLADORES DE PLAGAS Y UN METODO PARA
UTILIZACIÓN DE LOS MISMOS

CLASE A01N 43/54

SOLICITANTE NIHON NOHYAKU CO., LTD.

TRAMITE 002

EVENTO 001

ACTUACIÓN 444 CONTESTACION A REQUERIMIENTO

En respuesta a su Oficio 0386 de 11 de enero de 2007, dentro del término de ley, completo los antecedentes señalados por su despacho de conformidad con lo pedido en el examen técnico N° 2046 de la siguiente manera:

Presento la siguiente documentación requerida por ese Despacho a saber:

1- UEHARA M ET AL : Synthesis and insecticidal activity of 3-aminoquinazolinone derivatives" Pesticide Science. Vol 55, n° 3, march-1999 pag 359-362

En los términos anteriores dejo contestado el requerimiento.
Cordialmente



GERMAN A. CUBILLOS MENDOZA
c.c. N° 19 ' 397.479 de Bogotá
T.P. N° 38.422 del C.S.J.
Código 3667

S. exigua; (5) chlorfluazuron and tebufenozide and methoxyfenozide in chlorfluazuron-resistant strains of tea tortrix, *H. magnanima*; and (6) flufenoxuron and tebufenozide in flufenoxuron-resistant strains of *S. exigua* and *C. pomonella*. Attempts to select *S. exigua* resistant to tebufenozide in the laboratory have so far been unsuccessful. Adults developing from larvae maintained over 12 generations at LC₂₅ levels of tebufenozide had complete loss of fecundity.¹⁴

Rohm and Haas Company has developed a proactive platform to manage resistance to ecdysone agonist insecticides. Global resistance monitoring programs are currently underway on *C. pomonella*, *S. exigua*, *H. magnanima*, certain Heliothine pests, and certain leafroller species to establish baseline susceptibility data as well to detect any shifts in susceptibility over time. In addition, we pro-actively advocate the use of traditional approaches for managing insecticide resistance, including frequent and repeated rotation with other insecticides with different modes of action to minimize selection pressure, limitations on the number and sequences of applications on certain crops, and incorporation of integrated pest management methods to reduce pesticide use and prolong these products in the marketplace.

REFERENCES

- 1 Steel GB, Insect growth regulators with juvenile hormone activity. *Annu Rev Entomol* 20:417-460 (1975).
- 2 Dhadialla TS, Carlson GR and Le DP, New insecticides with ecdysenoid and juvenile hormone activity. *Annu Rev Entomol* 43:545-569 (1998).
- 3 Yao TP, Forman BM, Jiang Z, Charles L, Chen J-D et al., Functional ecdysone receptor is the product of EcR and ultraspiracle genes. *Nature (London)* 366: 476-479 (1995).
- 4 Song Q, Akmanli ES, Litwack G and Gilbert LI, An immunophilin is a component of the insect ecdysone receptor (EcR) complex. *Insect Biochem Molec Biol* 27: 973-982 (1997).
- 5 Smagghe G, Nonsteroidal ecdysteroid agonists: biological activity and insect specificity PhD chem, University of Ghent, Belgium. 119 pp (1995).
- 6 Remakhan A, Macdonald A, Tomkins WL, Davis CN, Brownright AJ et al, Ultrastructural effects of a non-steroidal ecdysone agonist, RH-5992, on the sixth instar larva of the spruce budworm, *Choristoneura fumiferana*. *J Insect Physiol* 43:55-68 (1997).
- 7 Carlson GR and Yew NK, Tebufenozide, a potent agonist of the insect molting hormone, for the management of important caterpillar pests in Malaysia, in *Crop Prot. Towards 2000: Int Pestic Conf Kuala Lumpur Malaysia*. pp 128-129 (1996).
- 8 Le DP, Thirugnanam M, Lidert Z, Carlson GR and Rypn JB, RH-2485: A new selective insecticide for caterpillar control. *Proc Brighton Crop Prot Conf - Pests & Diseases*. pp 481-486 (1996).
- 9 Oake L, MIMIC® insecticide. *Tech Inf Bull Rohm and Haas Company*. 20 pp (1994).
- 10 Biddinger DJ, Hull LA and McPherson RA, Cross-resistance between azinphosmethyl-resistant and -susceptible strains of tufted apple bud moth (Lepidoptera: Tortricidae) and various insect growth regulators. *J Econ Entomol* 89: 274-287 (1996).
- 11 Wearing CH, Le P and Walker JTS, Concerns about managing organophosphorus-resistant populations of leafrollers in

New Zealand with tebufenozide. *Resistant Pest Management* 9:98-99 (1997).

- 12 Sempacher B and Bouvier JC, Cross-resistance between benzoylureas and benzoylhydrazines in the codling moth, *Cydia pomonella* L. *Pestic Sci* 45:369-379 (1995).
- 13 Regirotti G and Manaresi M, Assenza di resistenza incrociata tra diflubenzuron e tebufenozide in *Cydia pomonella* in due ceppi provenienti dall'Alto Adige. *Inf Fitopatol* 3:60-62 (1997).
- 14 Smagghe G, Dhadialla TS, Deyck S, Terry L and Degheele D, Tebufenozide in susceptible and artificially tolerant beet

XP-002158467

Synthesis and insecticidal activity of 3-aminoquinazolinone derivatives

Masahiro Uehara,* Toshiaki Shimizu, Shinsuke Fujioke, Masayuki Kimura, Kenji Teubata and Akira Seo

Nihon Nohyaku Research Center, 345 Oyamada-cho, Kawachi-nagano, Osaka 586-0084, Japan

Abstract: The discovery, structure-activity relationships and insecticidal properties of R-768 (1-propiony-3-(3-pyridylmethylamino)-1,2,3,4-tetrahydroquinazolin-2-one) and related compounds are described.

© 1999 Society of Chemical Industry

Keywords: 3-aminoquinazolinones; R-768; insecticide; aphicide

In the course of research on methoxyacrylate fungicides, it was found that compounds having a carbamate group instead of a methoxyacrylate group in the relevant compounds also showed fungicidal activity against a broad spectrum of fungi.¹ Structural modification of these compounds was conducted to find a more active compound.² In the synthetic process, 2-halomethylphenyl carbamates (Fig 1; 1) were used as one of the starting materials.³ These proved to be interesting intermediates for synthesizing heterocycles, because they have two different electrophilic functional groups which can react with a primary amine at the 1,1-position of the





Figure 1. Synthesis of 3-substituted quinazolinones.

* Correspondence to: Masahiro Uehara, Nihon Nohyaku Research Center, 345 Oyamada-cho, Kawachi-nagano, Osaka 586-0084, Japan.

Received 29 June 1998; accepted 12 November 1998




R¹: H > CH₃
 R²: alkyl ≠ alkoxy carbonyl ≠ acyl
 R³: H > alkyl, oxo
 X: H ≠ halo ≠ alkyl ≠ alkoxy
 Y: H > halogen, alkoxy etc.
 Z: O ≠ S
 imino form ≠ reduced form

Figure 2. Structure – activity relationship of 3-aminoquinazolinone derivatives.

nucleophilic reactant to give cyclic derivatives. As was expected, they reacted with excess of methylamine on the halomethyl moiety to give first the benzylamino derivative, which was then converted to 3-methylaminoquinazolinone (2d) in excellent yield. The Compound 1 also reacted with hydrazine and hydroxylamine to give 2b and 2c, respectively. Similar reactions had already been reported in the literature, in which the —NCO group of 2-halomethylphenyl isocyanates reacted with a primary amine to give urea derivatives. The ureas were then cyclized by treatment with base to form the *N*-substituted 2-amino-4*H*-3,1-benzoxazines.^{3–5} In our experiments with compound 1, the benzoxazine derivatives were not detected at all as by-products.

Several insecticides having a semicarbazide group have been reported, *N*-phenylcarbamoylpyrazolines being well known to be a highly active class of insecticide.^{6,7} Therefore the 3-amino compound, 2b, having a semicarbazide moiety, and its derivatives were expected to possess insecticidal activity. Of



4 R = H
 R-768, R = propionyl

Figure 3. Structure of the reduced form and R-768.

Table 1. Activity of R-768 against major species of aphid

Chemical	LC ₅₀ (mg litre ⁻¹) ^a		
	<i>Myzus persicae</i>	<i>Aphis gossypii</i>	<i>Brevicoryne brassicae</i>
R-768	0.1–0.3	0.1–0.3	0.3–1
Imidacloprid	0.1–0.3	0.01–0.1	0.1–0.3
Acaphate	10–30	— ^b	— ^b
Lambda-cyhalothrin	0.1–1	0.1–1	0.1–1

^a Assessed six days after treatment.

^b Not Tested.

several derivatives synthesized and tested for insecticidal activity (Fig 1), the imino derivative 3a, synthesized from 2b and pyridine-3-aldehyde, was found to show high insecticidal activity against the Hemipterous insect, *Myzus persicae* Sulz, while the amide 3b and the urea 3c were inactive. From these preliminary results, we selected 3a as a lead compound and synthesized analogues to optimize the insecticidal activity.


Figure 2 summarizes the structure-activity relationships of 3-substituted quinazolinones. Among the compounds having the imino moiety, those of the 3-pyridyl azomethine type showed the highest activity. Replacement of 3-pyridyl by substituted phenyl, heterocycles other than 3-pyridyl or alkyl groups decreased the activity. Although the 2-chloropyridylmethyl group was used successfully to generate nitromethylene insecticides,⁸ halogeno-substituted 3-pyridyl derivatives in the series of compounds with structure 3a were markedly less active. Furthermore, any substitution in the pyridine ring failed to increase the activity.

The activity of quinazolinone derivatives with small alkyl groups at the 1-position in the quinazolinone ring was almost equal to that of 3a, but compounds with long or bulky alkyl substituents showed decreased activity. Derivatives with formyl, lower alkylcarbonyl and lower alkoxy carbonyl groups showed greater activity. The optimum carbon chain length for activity seemed to be two to four. Quinazolinone derivatives with alkyl or oxo at the 4-position were much less active, whereas all quinazolinone derivatives with halogen, alkyl or alkoxy substituents at the 5 to 8 positions showed high activity, differences in activity between them being relatively small. The activity of the 2-thione derivative of 3a was almost equal to that of 3a itself.

The imino moiety of 3a was reduced to give compound 4 (Fig 3). This reduced derivative possessed almost the same activity as 3a in a greenhouse test. However, 3a did not show good activity in field trials, whereas the reduced derivative 4 showed excellent activity. It was found that the poor performance of compound 3a in the field was due to photo-instability, its half-life being <1 h in a sunlight test, while that of 4 was >8 h under the same conditions. Because compound 4 was sufficiently stable to show insecticidal activity in field trials, optimization of the structure and activity was progressed using the reduced form. The effects of substituents R¹, R², R³, X, Y and Z (Fig 2) on the activity of the reduced form were almost the same as in the imino form. Quinazolinone derivatives containing lower alkyl and lower alkoxy carbonyl substituents at the 1-position in the quinazolinone ring were more active than 4, and it was thought that the substituents at the 1-position in the quinazolinone ring play an important role in promoting the penetration of the compound through the cuticular layer of insects. The quinazolinone derivative with a pro-

714

Figure 4. Time-course of mortalities of *Myzus persicae* after spray. Stage: Apterous adult. Diet: Cabbage leaf disc, 80 mm diameter on agar. Treatment: Spraying with 20 ml water solution of compound ($100 \text{ mg litre}^{-1}$) to 20 insects on the leaf disc



pionyl substituent at the 1-position showed the greatest activity. As a result, R-768 (Fig 3) was selected as a candidate for development, taking into account its insecticidal activity and its physico-chemical properties, which would influence the formulation.

Although R-768 had no effect on Lepidopterous and Coleopterous insects and mites, it showed excellent activities against some Hemipterous insect pests.

Its activities against *M. persicae* and *Aphis gossypii* Glover were almost equal to those of imidacloprid, and superior to those of acephate (Table 1).

The lethal action of R-768 against *M. persicae* was slower than that of the conventional insecticides in that aphids sprayed with R-768 needed a few days to die (Fig 4). These affected aphids immediately stopped feeding and abandoned the sprayed leaf. The inhibition of feeding of these affected aphids




Figure 5. Efficacy of R-768 against *Aphis gossypii* on potato in a field trial. Application: 7 August 1998 in Japan. Spray: 800 litre ha^{-1} .




Figure 6. Efficacy of R-768 against *Rhopalosiphum padi* on wheat in a field trial. Application: 17 July 1998 in Japan. Spray: 400 litre ha^{-1} .

Table 2. Activity of R-768 against resistant strains of aphid

Species	Strain	LC_{50} (mg litre^{-1}) ^a
<i>Myzus persicae</i>	US1L (susceptible standard)	0.8-0.9
	784J (resistant to OPs, carbamates and pyrethroids)	0.5
	826B (tolerant to imidacloprid)	0.8
<i>Aphis gossypii</i>	171B (susceptible standard)	0.7
	988E (resistant to OPs, carbamates and pyrethroids)	0.2

^a Assessed six days after treatment.

seemed to be so strong that they never regained the ability to suck even after they were transferred to untreated plants. Although they produced a few offspring before death, the offspring were also unable to feed and died without growing. These actions of R-768 are different from the conventional insecticides, suggesting that R-768 has a novel mode of action. In the laboratory study, R-768 was equally effective against organophosphate-, carbamate- and pyrethroid-resistant strains of aphid as well as against susceptible strains of *M. persicae* and *A. gossypii* (Table 2). This biological performance is also consistent with a new mode of action.

In a field trial on potatoes, the efficacy of R-768 at 50 or 100 g ha⁻¹ against the strain of *A. gossypii* that was resistant to lambda-cyhalothrin and pirimicarb, was comparable to that of imidacloprid at 50 g ha⁻¹ (Fig 5). These results indicated that R-768 possesses no cross-resistance to other insecticides and it will be useful as a candidate in areas where resistance problems become more serious. R-768 also showed excellent control of aphids on wheat (Fig 6), on apple trees (Fig 7) and on various crops, without phytotoxicity, in the field trials.

On the other hand, R-768 was safe to beneficial insects such as *Bombus mori* L., *Osmia cornifrons*, *Bombus terrestris* and *Apis mellifera* L., and non-target arthropods such as *Harmonia axyridis*, *Chrysopa nipponensis*, *Orius* sp., *Aphelinus* sp., *Ephedrus japonicus* and *Minusmenops tricuspidatus* in laboratory studies. This outstanding selectivity was also demonstrated for the pollinators of tomato and strawberry in a greenhouse study, and for some natural enemies in the field trials. Therefore R-768 could be one of a few aphid control agents which can be useful soon after releasing the pollinators and natural enemies in biological control systems.

In conclusion, R-768 is a representative of a new and highly active class of aphid control agent with large margins of safety to beneficial insects and natural enemies. Lack of cross-resistance indicates that R-768 could be useful as a candidate in the

resistant management strategies. Its favorable characters make it especially suitable for IPM programmes.

REFERENCES

- 1 Uehara M, Tajima S, Nishiguchi T, and Tsubata K, N-Substituted phenylcarbamic acid derivatives. *Jpn Kokai Tokkyo Koho JP 8, 188565* (1996).
- 2 Ohnishi O, Tajima S, Nishiguchi T and Tsubata K, N-Substituted phenylcarbamic acid derivatives. *Jpn Kokai Tokkyo Koho JP 8, 225515* (1996).
- 3 Jozef G and Mariani B, Simple and efficient synthesis of 4H-3,1-benzoxazines from 2-bromomethylphenyl isocyanate and amines. *Coll Czech Chem Commun 55:752-760* (1990).
- 4 Brich K and Linthard O, Herstellung von 2-(mono,di,tri-chloromethyl)-phenyl-isocyanaten. *Synthese 376-377* (1978).
- 5 Walter G. and Hans-Joachim K, N-Substituierte 2-amino-4H-3,1-benzoxazine aus 2-chloromethylphenyl-isocyanaten. *Synthese 377-378* (1978).
- 6 Mulder R, Wellings K and van Deelen JJ, A new class of insecticides. *Naturwissenschaften 62:531-532* (1975).
- 7 Wellings K, Grosscurt AC and van Hes R, 1-Phenylcarbamoyl-2-pyrrolines: a new class of insecticides. 1. Synthesis and insecticidal properties of 3-phenyl-1-phenylcarbamoyl-2-pyrrolines. *J Agric Food Chem 25:987-992* (1977).
- 8 Kagabu S, Studies on the synthesis and insecticidal activity of neonicotinoid compound. *Nihon Noyaku Gakkaishi (J Pest Sci) 21:231-239* (1996).

Expression of *Neurospora crassa* β -tubulin, target protein of benzimidazole fungicides, in *Escherichia coli*

Mitsuru Yoshida,^{1*} Yoshihiro Narusaka,¹ Eiichi Minami² and Hideo Ishii¹

¹National Institute of Agro-Environmental Sciences, Tsukuba, Ibaraki 305-8604, Japan

²National Institute of Agro-Biological Resources, Tsukuba, Ibaraki 305-8602, Japan

Abstract: β -Tubulin of a wild-type *Neurospora crassa* strain was expressed using pET-16b, pET-29a(+), and pET-32a(+) expression vectors in *Escherichia coli* BL21 (DE3) strain. Yield of the expressed soluble protein was estimated to be about 0.1 mg ml⁻¹ culture broth. The β -tubulins with S-Tag expressed by pET-29a(+) and pET-32a(+) bound to the S-protein Agarose by affinity binding but the thrombin and enterokinase treatments did not release β -tubulin, suggesting that the protease cleavage sites connecting S-Tag and β -tubulin were not exposed to approach of the proteases. The β -tubulin expressed by pET-16b did not bind to nickel resin, suggesting that its His-Tag was folded into the protein core. The protein expressed by pET-32a(+) was bound to the nickel resin and purified by the column chromatography.

© 1999 Society of Chemical Industry

Keywords: β -tubulin; gene expression; *Neurospora crassa*; *Escherichia coli*; expression vector

* Correspondence to: Mitsuru Yoshida, Food Research Institute, Kannondai 2-1-2, Tsukuba, Ibaraki 305-8642, Japan.

E-mail: mitsuru@nri.affrc.go.jp

(Received 29 June 1998; accepted 12 November 1998)




Figure 7. Efficacy of R-768 against *Aphis citricola* on apple trees in a field trial. Application: 22 May 1998 in Japan. Spray: 1000 litre ha⁻¹.