

Señores
DIVISION DE NUEVAS CREACIONES
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
Bogotá, D.C.

Re.: Código 02-01-329
Expediente No. 00-76.099
Solicitud de Patente
SmithKline Beecham Corporation

En relación con el Expediente de la referencia y con el fin de dar cumplimiento a la preceptuado por el artículo 4o. literal d.) numeral 3 de la Convención de París adjunto a este escrito sírvanse encontrar la copia certificada de la solicitud de patente de la cual se reivindica prioridad, la cual se encuentra debidamente traducida en sus apartes pertinentes.

De Ustedes Atentamente,

GERMAN CASTILLO GRAU
T.P. No. 2978 del C.S.J.

PA 387364



62 48 A

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE


United States Patent and Trademark Office

September 28, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/158,707

FILING DATE: October 08, 1999



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

T. Wallace
T. WALLACE

Certifying Officer

49
A/PROV

EXPRESS MAIL CERTIFICATE

"Express Mail" Mailing Label Number EL085632337US

Date Of Deposit: October 8, 1999

I Herby Certify That This Paper Or Fee Is Being Deposited With The United States Postal Service "Express Mail Post Office To Addressee" Service Under 37 CFR 1.10 On The Date Indicated Above And Is Addressed To: Assistant Commissioner for Patents, Washington, D.C. 20231.

Name Of Person Mailing Paper Or Fee

(Type Or Print)

Signature

Mary E. McCarthy

30/156707

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION for PATENT under 37 CFR 1.53(c).

Docket No.

P51038P

INVENTOR(s) / APPLICANT(s)

Last Name	First Name	Middle Initial	Residence (City and Either State or Foreign Country)
MILLER	William	H.	Collegeville, Pennsylvania
SEEFELD	Mark	A.	Phoenixville, Pennsylvania
NEWLANDER	Kenneth	A.	West Chester, Pennsylvania
UZINSKAS	Irene	N.	Villanova, Pennsylvania

TITLE OF THE INVENTION (280 characters max)

Fab I Inhibitors

CORRESPONDENCE ADDRESS

SMITHKLINE BEECHAM CORPORATION

Corporate Intellectual Property - UW2220

709 Swedeland Road

King of Prussia

Telephone No. 610-270-5022

Facsimile No. 610-270-5090

State

PA

Zip Code

19406-2799

Country

United States of America

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification	Number of Pages	29	☐ Small Entity Statement
☐ Drawings	Number of Sheets		☐ Other (specify)

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

☒ The Commissioner is hereby authorized to charge filing fees and credit Deposit Account No. 19-2570

PROVISIONAL FILING
FEE AMOUNT (\$)

\$150.00

Respectfully submitted,

Signature:

Mary E. McCarthy
Mary E. McCarthy

Date:

October 8, 1999

Registration No.:

32917

☐ Additional inventors are being named on separately numbered sheets attached hereto.

PROVISIONAL APPLICATION FILING ONLY

SEND TO: Assistant Commissioner for Patents, Box Provisional Application, Washington, D.C. 20231.

n:\mms\spec\P5036PProv.doc

5x 50
/

TITLE
Fab I Inhibitors

FIELD OF THE INVENTION

5 This invention relates to pharmaceutically active compounds which inhibit Fab I and are useful for the treatment of bacterial infections.

BACKGROUND OF THE INVENTION

10 While the overall pathway of saturated fatty acid biosynthesis is similar in all organisms, the fatty acid synthase (FAS) systems vary considerably with respect to their structural organization. Vertebrates and yeast possess a FAS in which all the enzymatic activities are encoded on one or two polypeptide chains, respectively, and the acyl carrier protein (ACP) is an integral part of the complex. In contrast, in bacterial FAS, each of the reactions is catalyzed by a distinct, mono-functional enzyme and the ACP is a discrete

15 protein. Therefore, there is considerable potential for the selective inhibition of the bacterial system by antibacterial agents.

20 Fab I (previously designated EnvM) functions as an enoyl-ACP reductase (Bergler, et al, (1994), *J.Biol.Chem.* 269, 5493-5496) in the final step of the four reactions involved in each cycle of bacterial fatty acid biosynthesis. In this pathway, the first step is catalyzed by β -ketoacyl-ACP synthase, which condenses malonyl-ACP with acetyl-CoA (FabH, synthase III). In subsequent rounds, malonyl-ACP is condensed with the growing-chain acyl-ACP (FabB and FabF, synthases I and II, respectively). The second step in the elongation cycle is ketoester reduction by NADPH-dependent β -ketoacyl-ACP reductase (FabG). Subsequent dehydration by β -hydroxyacyl-ACP dehydrase (either FabA or FabZ)

25 leads to trans-2-enoyl-ACP, which in turn is converted to acyl-ACP by NADH-dependent enoyl-ACP reductase (Fab I). Further rounds of this cycle, adding two carbon atoms per cycle, eventually lead to palmitoyl-ACP (16C), where upon the cycle is stopped largely due to feedback inhibition of Fab I by palmitoyl-ACP (Haath, et al, (1996), *J.Biol.Chem.* 271, 1833-1836). Thus, Fab I is a major biosynthetic enzyme and is a key regulatory point in the

30 overall synthetic pathway of bacterial fatty acid biosynthesis. Therefore, Fab I is an ideal target for antibacterial intervention.

35 Studies have shown that diazaborine antibiotics inhibit fatty acid, phospholipid and lipopolysaccharide (LPS) biosynthesis and that the antibacterial target of these compounds is Fab I. For example, derivative 2b18 from Grassberger, et al (1984) *J. Med Chem* 27 947-953 has been reported to be a non-competitive inhibitor of Fab I (Bergler, et al, (1994), *J.Biol.Chem.* 269, 5493-5496). Also, plasmids containing the Fab I gene from diazaborine resistant *S. typhimurium* conferred diazaborine resistance in *E.coli* (Turnowsky, et al,

(1989), *J. Bacteriol.*, 171, 6555-6565). Furthermore, inhibition of Fab I either by diazaborine or by raising the temperature in a Fab I temperature sensitive mutant is lethal. These results demonstrate that Fab I is essential to the survival of the organism (Bergler, et al, (1994), *J. Biol. Chem.* 269, 5493-5496).


- 5 Recent studies have shown that Fab I is also the target for the broad spectrum antibacterial agent triclosan (McMurry, et al, (1998) *Nature* 394, 531-532). A crystal structure of the *E. Coli* Fab I complexed with NAD and triclosan shows that triclosan acts as a site-directed, very potent inhibitor of Fab I by mimicking its natural substrate (Levy, et al, (1999) *Nature* 398, 383-384). Ward, et al ((1999) *Biochem.* 38, 12514-12525) have
- 10 shown that there is no evidence for the formation of a covalent complex between Fab I and triclosan, which would be analogous to the diazaborines; triclosan differs from these compounds in that it is a reversible inhibitor of Fab I. The structural data for the complex of Fab I with NAD and triclosan provides important information about Fab I as a therapeutic target.
- 15 Importantly, it has now been discovered that certain compounds are Fab I inhibitors and have antibacterial activity, and, therefore, may be useful for the treatment of bacterial infections in mammals, particularly in man.

SUMMARY OF THE INVENTION

- 20 This invention comprises compounds of the formula (I), as described hereinafter, which inhibit Fab I and are useful in the treatment of bacterial infections.
- This invention is also a pharmaceutical composition comprising a compound according to formula (I) and a pharmaceutically acceptable carrier.
- This invention is also a method of treating bacterial infections by inhibiting Fab I.
- 25 In a particular aspect, the compounds of this invention are useful as antibacterial agents.

DETAILED DESCRIPTION

This invention comprises compounds of formula (I):




(I)

wherein:

R^1 is C_{1-4} alkyl;

56-5-16

 R^2 is C_{1-4} alkyl; R^3 is Q is a single bond, $-O-$, $-S-$ or $-NH-$, or $-CH_2CH_2-$;

5

 R^4 is C_{1-6} alkyl;

is a five- or six-membered heteroaromatic ring or a six-membered aromatic ring optionally substituted by R^5 ;



is a five- or six-membered heteroaromatic ring or a six-membered aromatic ring optionally substituted by R^6 ;



10

is a five- or six-membered heteroaromatic ring or a six-membered aromatic ring optionally substituted by R^6 ;

 R^5 is OR' or $NR'C(O)R'$; R^6 is OR' or NH_2 ; X is H , C_{1-4} alkyl, OR' , SR' , CN , $N(R')_2$, $CH_2N(R')_2$, NO_2 , CF_3 , CO_2R' ,15 $CON(R')_2$, COR' , $NR'C(O)R'$, F , Cl , Br , I or $-S(O)_rCF_3$; R' is H , C_{1-6} alkyl or $-C_{0-6}$ alkyl-Ar; and r is 0, 1 or 2;

or a pharmaceutically acceptable salt thereof.

20 Also included in this invention are pharmaceutically acceptable addition salts and complexes of the compounds of this invention. In cases wherein the compounds of this invention may have one or more chiral centers, unless specified, this invention includes each unique racemic compound, as well as each unique nonracemic compound.


In cases in which compounds have unsaturated carbon-carbon double bonds, both the *cis* (*Z*) and *trans* (*E*) isomers are within the scope of this invention. In cases wherein

compounds may exist in tautomeric forms, such as keto-enol tautomers, such as  and , each tautomeric form is contemplated as being included within this invention, whether existing in equilibrium or locked in one form by appropriate substitution with R'. The meaning of any substituent at any one occurrence is independent of its meaning, or any other substituent's meaning, at any other occurrence.

Also included in this invention are prodrugs of the compounds of this invention. Prodrugs are considered to be any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo*.

The compounds of formula (I) inhibit Fab I. Inhibition of this enzyme is useful in the treatment of bacterial infections. Also, the compounds of this invention may be useful as antifungal agents. Additionally, the compounds may be useful in combination with known antibiotics.


With respect to formula (I), this invention preferably includes compounds of formula (Ia):



(Ia)

in which R³ is as defined for formula (I) compounds.

Suitably, with respect to formula (I), R³ is:



Preferably, A is 2-, 3- or 4-pyridyl, 2- or 3-furanyl, 2- or 3-thiophenyl, thiazolyl, phenyl or phenyl substituted by OC₁₋₄alkyl or

NHC(O)C₁₋₄alkyl. Most preferably, A is 3- or 4-pyridyl, 3-furanyl, 3-thiophenyl,

phenyl or phenyl substituted by OCH_3 or NHC(O)CH_3 . Also, preferably,



is 2-,


3- or 4-pyridyl, phenyl or phenyl substituted by OH or NH_2 . Most preferably, phenyl or 2-pyridyl.



is

Suitably, with respect to formula (I), R^3 is:


5



in which Q is a single bond, $-\text{O}-$ or $-\text{CH}_2\text{CH}_2-$. Preferably,

Q is a single bond.

Suitably, with respect to formula (I), R^3 is:



10

. Preferably,



is phenyl unsubstituted or

substituted by OH or NH_2 and R^4 is C_{1-4} alkyl. Most preferably,



is phenyl and

R^4 is CH_3 . Alternately,



is 2-pyridyl and R^4 is C_{1-4} alkyl.

Representative of the novel compounds of this invention are the following:

2-[2-(4-Methoxyphenyl)ethyl]-N-methyl-N-[(1-methyl-1H-indol-2-

15 yl)methyl]nicotinamide;

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenylbenzamide;

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-4-yl)benzamide;

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-3-yl)benzamide;

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(thiophen-3-yl)benzamide;

20

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(furan-3-yl)benzamide;

52
SS
9

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenoxybenzamide;
N,2-Dimethyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide; and
2-[3-(Acetylamino)phenyl]-N-methyl-N-[(1-methyl-1H-indol-2-

yl)methyl]benzamide;

5 or a pharmaceutically acceptable salt thereof.

Abbreviations and symbols commonly used in the peptide and chemical arts are used herein to describe the compounds of this invention. In general, the amino acid abbreviations follow the IUPAC-IUB Joint Commission on Biochemical Nomenclature as described in *Eur. J. Biochem.*, 158, 9 (1984).

10 C₁₋₄alkyl as applied herein means an optionally substituted alkyl group of 1 to 4 carbon atoms, and includes methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl. C₁₋₆alkyl additionally includes pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof. C₀₋₄alkyl and C₀₋₆alkyl additionally indicates that no alkyl group need be present (e.g., that a covalent bond is present).

15 Any C₁₋₄alkyl or C₁₋₆alkyl may be optionally substituted with the group R^x, which may be on any carbon atom that results in a stable structure and is available by conventional synthetic techniques. Suitable groups for R^x are C₁₋₄alkyl, OR', SR', CN, N(R')₂, CH₂N(R')₂, -NO₂, -CF₃, -CO₂R', -CON(R')₂, -COR', -NR'C(O)R', F, Cl, Br, I, or -S(O)_xCF₃, wherein R' and r are as defined for formula (I) compounds.

20 Halogen or halo means F, Cl, Br, and I.

Ar, or aryl, as applied herein, means phenyl or naphthyl, or phenyl or naphthyl substituted by one to three substituents, such as those defined above for alkyl, or substituted by methylenedioxy.

25 Het, or heterocycle, indicates an optionally substituted five or six membered monocyclic ring, or a nine or ten-membered bicyclic ring containing one to three heteroatoms chosen from the group of nitrogen, oxygen and sulfur, which are stable and available by conventional chemical synthesis. Illustrative heterocycles are benzofuryl, benzimidazolyl, benzopyranyl, benzothienyl, furyl, imidazolyl, indolyl, morpholinyl, piperidinyl, piperazinyl, pyrrolyl, pyrrolidinyl, tetrahydropyridinyl, pyridinyl, thiazolyl, 30 thienyl, quinolinyl, isoquinolinyl, and tetra- and perhydro- quinolinyl and isoquinolinyl. Any accessible combination of up to three substituents on the Het ring, such as those defined above for alkyl, that are available by chemical synthesis and are stable are within the scope of this invention.


35 Certain radical groups are abbreviated herein. t-Bu refers to the tertiary butyl radical, Boc refers to the t-butyloxycarbonyl radical, Fmoc refers to the fluorenylmethoxycarbonyl radical, Ph refers to the phenyl radical, Cbz refers to the benzyloxycarbonyl radical, Bn refers to the benzyl radical, Me refers to methyl, Et refers to

ethyl, Ac refers to acetyl, Alk refers to C₁₋₄alkyl, Nph refers to 1- or 2-naphthyl and cHex refers to cyclohexyl. Tet refers to 5-tetrazolyl.

Certain reagents are abbreviated herein. DCC refers to dicyclohexylcarbodiimide, DMAP refers to dimethylaminopyridine, EDC refers to 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, hydrochloride, HOBt refers to 1-hydroxybenzotriazole, THF refers to tetrahydrofuran, DIEA refers to diisopropylethylamine, DEAD refers to diethyl azodicarboxylate, PPh₃ refers to triphenylphosphine, DIAD refers to diisopropyl azodicarboxylate, DME refers to dimethoxyethane, DMF refers to dimethylformamide, NBS refers to N-bromosuccinimide, Pd/C refers to a palladium on carbon catalyst, PPA refers to polyphosphoric acid, DPPA refers to diphenylphosphoryl azide, BOP refers to benzotriazol-1-yloxy-tris(dimethyl-amino)phosphonium hexafluorophosphate, HF refers to hydrofluoric acid, TEA refers to triethylamine, TFA refers to trifluoroacetic acid, PCC refers to pyridinium chlorochromate.

Compounds of the formula (I) are prepared by the general methods described in Schemes I and II.

Scheme I




(a) 1-methyl-2-(methylaminomethyl)indole, EDC, HOBt · H₂O, Et₃N, DMF; (b) furan-3-boronic acid, Pd(PPh₃)₄, Cs₂CO₃, DME, H₂O.

A suitable 2-haloarylcarboxylic acid, for instance 2-bromobenzoic acid (I-1), is converted to an activated form using, for example, EDC and HOBt, or SOCl₂, and the activated form is subsequently reacted with an appropriate amine, for instance 1-methyl-2-(methylaminomethyl)indole, in a suitable solvent such as DMF, CH₂Cl₂, or CH₃CN, to afford I-2. Depending on whether acid neutralization is required, an added base, such as triethylamine (Et₃N), diisopropylethylamine ((i-Pr)₂NEt), or pyridine, may be used. Many additional methods for converting a carboxylic acid to an amide are known, and can be found in standard reference books, such as "Compendium of Organic Synthetic Methods", Vol. I - VI (published by Wiley-Interscience), or Bodansky, "The Practice of Peptide Synthesis" (published by Springer-Verlag). The resulting haloarylcarboxamide I-2 is

6A 52

reacted with an appropriate boronic acid derivative, such as furan-3-boronic acid, to afford I-3 in a Suzuki-type reaction (*Synth. Commun.* 1981, 11, 513; for a review, see Martin, A. R.; Yang, Y. *Acta Chem. Scand.* 1993, 47, 221). This reaction is generally mediated by tetrakis(triphenylphosphine)palladium(0), although a wide variety of palladium catalysts can be used, including palladium acetate, palladium acetate in the presence of tri(*ortho*-tolyl)phosphine, bis(triphenylphosphine)palladium (II) chloride, 1,4-bis(diphenylphosphinobutane)palladium(II) chloride, bis(acetonitrile)palladium(II) chloride, bis(dibenzylideneacetone)palladium(0), and palladium on carbon in the presence of triphenylphosphine. The reaction requires a base as an acid scavenger, and generally aqueous sodium carbonate (Na_2CO_3) is used. However, other bases, such as sodium hydrogen carbonate, potassium carbonate, cesium carbonate, cesium fluoride, and triethylamine can be used. Typically, water-miscible neutral solvents, including DME, THF, and DMF, are preferred.

Scheme II



(a) *para*-anisaldehyde, NaH, *tert*-BuOH, DMF; (b) H_2 , Pd/C, NaOH, H_2O ; (c) 1-methyl-2-(methylaminomethyl)indole, EDC, HOBT · H_2O , (*i*-Pr) $_2$ NEt, DMF.

62 58
17

According to the general procedure of Brenner et al. (*J. Het. Chem.* 1982, 19, 897-900), a suitable 2-methylnicotinic ester, for instance ethyl 2-methylnicotinate (II-1), is reacted with an aromatic aldehyde, such as *para*-anisaldehyde, to afford II-2. The reaction is mediated by strong base, typically sodium *tert*-butoxide, and is conducted in a polar solvent, generally DMF. Reduction of the olefin group of I-2 is conveniently accomplished by hydrogenation as described by van der Stelt, et al. (*Arzneim. - Forsch.* 1968, 18, 756 - 758). The reaction is conducted under basic conditions in the presence of a palladium catalyst, generally palladium on activated charcoal. Alkali metal hydroxides, such as sodium hydroxide or potassium hydroxide, are the preferred bases, and the reaction is typically conducted in H₂O, although an aqueous hydroxylic solvent, such as aqueous MeOH, EtOH, or *i*-PrOH, may also be used. The resulting acid, II-3, is converted to II-4 by the methods described in Scheme I.

Amide coupling reagents as used herein denote reagents which may be used to form peptide bonds. Typical coupling methods employ carbodiimides, activated anhydrides and esters and acyl halides. Reagents such as EDC, DCC, DPPA, PPA, BOP reagent, HOBt, N-hydroxysuccinimide and oxalyl chloride are typical.

Acid addition salts of the compounds are prepared in a standard manner in a suitable solvent from the parent compound and an excess of an acid, such as hydrochloric, hydrobromic, hydrofluoric, sulfuric, phosphoric, acetic, trifluoroacetic, maleic, succinic or methanesulfonic. Certain of the compounds form inner salts or zwitterions which may be acceptable. Cationic salts are prepared by treating the parent compound with an excess of an alkaline reagent, such as a hydroxide, carbonate or alkoxide, containing the appropriate cation; or with an appropriate organic amine. Cations such as Li⁺, Na⁺, K⁺, Ca⁺⁺, Mg⁺⁺ and NH₄⁺ are specific examples of cations present in pharmaceutically acceptable salts.

This invention also provides a pharmaceutical composition which comprises a compound according to formula (I) and a pharmaceutically acceptable carrier. Accordingly, the compounds of formula (I) may be used in the manufacture of a medicament. Pharmaceutical compositions of the compounds of formula (I) prepared as hereinbefore described may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially suitable for parenteral administration, but may also be used for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin,

60
59

hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

Alternately, these compounds may be encapsulated, tableted or prepared in a emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. Liquid carriers include syrup, peanut oil, olive oil, saline and water. The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing, when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

For rectal administration, the compounds of this invention may also be combined with excipients, such as cocoa butter, glycerin, gelatin or polyethylene glycols, and molded into a suppository.

For topical administration, the compounds of this invention may be combined with diluents to take the form of ointments, gels, pastes, creams, powders or sprays. The compositions which are ointments, gels, pastes or creams contain diluents, for example, animal and vegetable fats, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures of these substances. The compositions which are powders or sprays contain diluents, for example, lactose, talc, silicic acid, aluminum hydroxide, calcium silicate and polyamide powder, or mixtures of these substances. Additionally, for topical ophthalmologic administration, the typical carriers are water, mixtures of water and water miscible solvents, such as lower alkanols or vegetable oils, and water-soluble non-toxic polymers, for example cellulose derivatives, such as methyl cellulose.

The compounds described herein are inhibitors of Fab I, and are useful for treating bacterial infections. For instance, these compounds are useful for the treatment of bacterial infections, such as, for example, infections of upper respiratory tract (e.g. otitis media, bacterial tracheitis, acute epiglottitis, thyroiditis), lower respiratory (e.g. empyema, lung abscess), cardiac (e.g. infective endocarditis), gastrointestinal (e.g. secretory diarrhoea, splenic abscess, retroperitoneal abscess), CNS (e.g. cerebral abscess), eye (e.g. blepharitis, conjunctivitis, keratitis, endophthalmitis, preseptal and orbital cellulitis, dacryocystitis),

64
60
H

kidney and urinary tract (e.g. epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (e.g. impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis), and bone and joint (e.g. septic arthritis, osteomyelitis). Also, the compounds of this invention may be useful as antifungal agents. Additionally, the compounds may be useful in combination with known antibiotics.

The compounds of this invention are administered to the patient, in a manner such that the concentration of drug is sufficient to treat bacterial infections. The pharmaceutical composition containing the compound is administered at an oral dose of between about 10 mg to about 1000 mg, taken once or several times daily, in a manner consistent with the condition of the patient. Preferably, the oral dose would be about 50 mg to about 500 mg, although the dose may be varied depending upon the age, body weight and symptoms of the patient. For acute therapy, parenteral administration is preferred. An intravenous infusion of the compound of formula (I) in 5% dextrose in water or normal saline, or a similar formulation with suitable excipients, is most effective, although an intramuscular bolus injection is also useful. The precise level and method by which the compounds are administered is readily determined by one skilled in the art.

The compounds may be tested in one of several biological assays to determine the concentration of compound which is required to have a given pharmacological effect.

Cloning of *S. aureus* FabI:

The *fabI* gene was cloned from the chromosomal DNA of *S. aureus* strain WCUH29 using the polymerase chain reaction. Amplification was performed using *Taq* DNA polymerase (BRL) and the following primers: 5'-

CGCCTCGAGATGTTAAATCTTGAAAACAAAACATATGTC-3' and 5'-

CGCGGATCCAATCAAGTCAGGTTGAAATATCCA-3' (*Xho*I and *Bam*HI sites

underlined). The resulting fragment was then digested with *Xho*I and *Bam*HI and ligated into *Xho*I- and *Bam*HI-digested expression vector pET-16b (Novagen), producing pET-His₁₀-*fabI*.

The gene sequence of *fabI* was confirmed by automated cycle sequencing using an Applied Biosystems model 377 machine. The untagged version of pET-*fabI* was constructed by digesting pET-His₁₀-*fabI* with *Nco*I and *Nde*I to remove a 97 bp fragment encoding the His 10 tag, the factor Xa cleavage site and the first 8 amino acids of FabI, and replacing it with a linker encoding the first 8 amino acids of FabI plus a glycine residue between the initiator methionine and the lysine at position 2. This plasmid was called pET-*fabI*. The linker was made by annealing the following two oligonucleotides: 5'-

CATGGGCTTAAATCTTGAAAACAAAACA-3' and 5'-

TATGTTTGTGTTTCAAGATTTAAGCC-3'. The linker sequence in pET-*fabI* was

65
61

confirmed by dideoxy sequencing. Only native FabI was used for compound evaluation. For overproduction of native FabI, plasmid pET-fabI was transformed into BL21(DE3) (Novagen) cells, to form strain BL21(DE3):pET-fabI.

5 Purification of *S. aureus* FabI

S. aureus FabI was expressed as soluble protein to 10% of total cell protein, 400g cells being recovered from 15L fermentation in tryptone phosphate medium. The cells were lysed and the sample centrifuged. The resulting supernatant was filtered and purified using three consecutive chromatography columns: ion-exchange (Source 15Q), dye-affinity (Blue sepharose), and size exclusion chromatography columns (Superose 12). After each column the FabI containing fractions were pooled, concentrated, and checked for purity and biological activity.

15 Cloning of *E. coli* FabI:

A PCR fragment of correct size for *E. coli* FabI was PCR amplified from *E. coli* chromosomal DNA, subcloned into the TOPO TA cloning vector, and verified by colony PCR + restriction endonuclease analysis. The presumptive *E. coli* FabI PCR fragment was subcloned into the expression vector pBluePet. The FabI clone was transformed into *E. coli* strain BL21(DE3). Small Scale expression studies show an over-expressed protein band of correct molecular weight (~28 Kda) for *E. coli* FabI clearly visible following Coomassie staining of SDS PAGE gels. DNA sequencing of the *E. coli* FabI expression constructs illustrated that no errors were apparent. N^o terminal amino acid sequencing has confirmed the over-expressed protein band to be *E. coli* FabI.

Purification of *E. coli* FabI

E. coli FabI was expressed as soluble protein to 15% of total cell protein, 120g cells being recovered from 3L fermentation in shake flasks in modified terrific broth. The cells were lysed and the sample centrifuged. The resulting supernatant was filtered and purified using three consecutive chromatography columns: ion-exchange (Source 15Q), dye-affinity (blue sepharose), and size exclusion (superose 12). After each column the FabI containing fractions were pooled, concentrated and checked for purity and biological activity.

35

668001-2075509

62

S. aureus FabI Enzyme Inhibition Assay (NADH):

Assays were carried out in half-area, 96-well microtitre plates. Compounds were evaluated in 50-uL assay mixtures containing 100 mM NaADA, pH 6.5 (ADA = N-[2-acetamido]-2-iminodiacetic acid), 4 % glycerol, 0.25 mM crotonoyl CoA, 1 mM NADH, and an appropriate dilution of *S. aureus* FabI. Inhibitors were typically varied over the range of 0.01-10 uM. The consumption of NADH was monitored for 20 minutes at 30 °C by following the change in absorbance at 340 nm. Initial velocities were estimated from an exponential fit of the non-linear progress curves represented by the slope of the tangent at t = 0 min. IC₅₀'s were estimated from a fit of the initial velocities to a standard, 4-parameter model and are typically reported as the mean ± S.D. of duplicate determinations. Triclosan, a commercial antibacterial agent and inhibitor of FabI, is currently included in all assays as a positive control. Compounds of this invention have IC₅₀'s from about 1.50 micromolar to about 0.15 micromolar.

S. aureus FabI Enzyme Inhibition Assay (NADPH):

Assays were carried out in half-area, 96-well microtitre plates. Compounds were evaluated in 150-uL assay mixtures containing 100 mM NaADA, pH 6.5 (ADA = N-[2-acetamido]-2-iminodiacetic acid), 4 % glycerol, 0.25 mM crotonoyl CoA, 50 uM NADPH, and an appropriate dilution of *S. aureus* FabI. Inhibitors were typically varied over the range of 0.01-10 uM. The consumption of NADPH was monitored for 20 minutes at 30 °C by following the change in absorbance at 340 nm. Initial velocities were estimated from an exponential fit of the non-linear progress curves represented by the slope of the tangent at t = 0 min. IC₅₀'s were estimated from a fit of the initial velocities to a standard, 4-parameter model and are typically reported as the mean ± S.D. of duplicate determinations. Triclosan, a commercial antibacterial agent and inhibitor of FabI, is currently included in all assays as a positive control.

E. coli FabI Enzyme Inhibition Assay:

Assays were carried out in half-area, 96-well microtitre plates. Compounds were evaluated in 150-uL assay mixtures containing 100 mM NaADA, pH 6.5 (ADA = N-[2-acetamido]-2-iminodiacetic acid), 4 % glycerol, 0.25 mM crotonoyl CoA, 50 uM NADH, and an appropriate dilution of *E. coli* FabI. Inhibitors were typically varied over the range of 0.01-10 uM. The consumption of NADH was monitored for 20 minutes at 30 °C by following the change in absorbance at 340 nm. Initial velocities were estimated from an

60158707.100899

62 63

exponential fit of the non-linear progress curves represented by the slope of the tangent at $t = 0$ min. IC_{50} 's were estimated from a fit of the initial velocities to a standard, 4-parameter model and are typically reported as the mean $\pm$ S.D. of duplicate determinations. Triclosan, a commercial antibacterial agent and inhibitor of FabI, is currently included in all assays as a positive control. Compounds of this invention have IC_{50} 's from about 20.0 micromolar to about 2.0 micromolar.

Antimicrobial Activity Assay:

Whole-cell antimicrobial activity was determined by broth microdilution using the National Committee for Clinical Laboratory Standards (NCCLS) recommended procedure, Document M7-A4, "Methods for Dilution Susceptibility Tests for Bacteria that Grow Aerobically". The compound was tested in serial two-fold dilutions ranging from 0.06 to 64 mcg/mL. Test organisms were selected from the following laboratory strains:

Staphylococcus aureus Oxford, *Staphylococcus aureus* WCUH29, *Streptococcus pneumoniae* ERY2, *Streptococcus pneumoniae* 1629, *Streptococcus pneumoniae* N 1387, *Enterococcus faecalis* 1, *Enterococcus faecalis* 7, *Haemophilus influenzae* Q1, *Haemophilus influenzae* NEMC1, *Moraxella Catarrhalis* 1502, *Escherichia coli* 7623 AcrABEFD+, *Escherichia coli* 120 AcrAB-, *Escherichia coli* MG1655, *Escherichia coli* MG1658. The minimum inhibitory concentration (MIC) was determined as the lowest concentration of compound that inhibited visible growth. A mirror reader was used to assist in determining the MIC endpoint.

One skilled in the art would consider any compound with a MIC of less than 256 μ g/mL to be a potential lead compound. Preferably, the compounds used in the antimicrobial assays of the present invention have a MIC value of less than 128 μ g/mL. Most preferably, said compounds have a MIC value of less than 64 μ g/mL.

The examples which follow are intended in no way to limit the scope of this invention, but are provided to illustrate how to make and use the compounds of this invention. Many other embodiments will be readily apparent to those skilled in the art.

EXAMPLES

General

Proton nuclear magnetic resonance (1H NMR) spectra were recorded at either 250, 300, 360, or 400 MHz, and chemical shifts are reported in parts per million (δ) downfield

60153707-100899

68 64

from the internal standard tetramethylsilane (TMS). Abbreviations for NMR data are as follows: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, dd = doublet of doublets, dt = doublet of triplets, app = apparent, br = broad. J indicates the NMR coupling constant measured in Hertz. CDCl₃ is deuteriochloroform, DMSO-d₆ is

- 5 hexadeteriodimethylsulfoxide, and CD₃OD is tetradeuteriomethanol. Mass spectra were obtained using electrospray (ES) ionization techniques. Elemental analyses were performed by Quantitative Technologies Inc., Whitehouse, NJ. Melting points were obtained on a Thomas-Hoover melting point apparatus and are uncorrected. All temperatures are reported in degrees Celsius. Analtech Silica Gel GF and E. Merck Silica
- 10 Gel 60 F-254 thin layer plates were used for thin layer chromatography. Flash chromatography was carried out on E. Merck Kieselgel 60 (230-400 mesh) silica gel. Analytical HPLC was performed on Beckman chromatography systems. Preparative HPLC was performed using Gilson chromatography systems. ODS refers to an octadecylsilyl derivatized silica gel chromatographic support. YMC ODS-AQ® is an ODS
- 15 chromatographic support and is a registered trademark of YMC Co. Ltd., Kyoto, Japan. PRP-1® is a polymeric (styrene-divinylbenzene) chromatographic support, and is a registered trademark of Hamilton Co., Reno, Nevada. Celite® is a filter aid composed of acid-washed diatomaceous silica, and is a registered trademark of Manville Corp., Denver, Colorado.
- 20

Preparation 1

Preparation of 1-methyl-2-(methylaminomethyl)indole

- 25 a) Ethyl 1-methylindole-2-carboxylate

NaH (60% dispersion in mineral oil, 8.02 g, 200.49 mmole) was washed with hexanes, then was suspended in dry DMF (530 mL). Solid ethyl indole-2-carboxylate (25.29 g, 133.66 mmole) was added portionwise over 5 - 10 min, allowing gas evolution to subside between additions. When the addition was complete, the yellow mixture was

30 stirred for 15 min, then methyl iodide (42 mL, 668.3 mmole) was added all at once. The reaction was exothermic, and the internal temperature rose to 40 - 45°C. After 1 hr, the reaction was quenched with 10% NH₄Cl (100 mL) and concentrated on the rotavap (high vacuum). The residue was partitioned between Et₂O (500 mL) and H₂O (100 mL), and the layers were separated. The Et₂O layer was washed with H₂O (100 mL), dried (MgSO₄),

35 and concentrated to leave the title compound (27.10 g, quantitative) as a light yellow solid. This was used without further purification: TLC (10% EtOAc/hexanes) R_f = 0.39.

60158707.100899

63
62

b) N,1-Dimethylindole-2-carboxamide

A suspension of ethyl 1-methylindole-2-carboxylate (27.10 g, 133.34 mmole) in 40% aqueous CH_3NH_2 (300 mL) and MeOH (30 mL) was stirred at RT. A solid tended to gradually creep up the walls of the flask, and was washed down periodically with MeOH.

- 5 The flask was tightly stoppered to keep the material inside the flask. As the reaction proceeded, the solid dissolved, but eventually the product began to precipitate. The reaction was stirred at RT for 5 days, then was concentrated to remove approximately 200 mL of the solvent. The remaining residue was diluted with H_2O (300 mL), and the solid
- 10 was collected by suction filtration and washed with H_2O . Drying at 50 - 60°C in high vacuum left the title compound (23.45 g, 93%) as a faintly yellow solid: ^1H NMR (300 MHz, CDCl_3) • 7.63 (d, $J = 8.0$ Hz, 1 H), 7.27 - 7.43 (m, 2 H), 7.10 - 7.20 (m, 1 H), 6.80 (s, 1 H), 6.10 - 6.30 (m, 1 H), 4.06 (s, 3 H), 3.01 (d, $J = 4.9$ Hz, 3 H).

c) 1-Methyl-2-(methylaminomethyl)indole

- 15 A 3-liter 3-necked roundbottom flask equipped with overhead stirring was charged with N,1-dimethylindole-2-carboxamide (23.45 g, 124.58 mmole) and anhydrous THF (170 mL). The solution was stirred while a solution of LiAlH_4 in THF (1.0 M, 250 mL, 250 mmole) was added via syringe. Gas was evolved during the addition of the first 50 mL of LiAlH_4 solution. When the addition was complete, the resulting light yellow solution was
- 20 heated at gentle reflux. After 23 hr, the reaction was cooled in ice and quenched by the sequential dropwise addition of H_2O (9.5 mL), 15% NaOH (9.5 mL), and H_2O (28.5 mL). The mixture was stirred for 15 min, then was filtered through celite®, and the filter pad was washed thoroughly with THF. The filtrate was concentrated and the residue was flash chromatographed on silica gel (10% MeOH/ CHCl_3 containing 0.5% conc. NH_4OH). The
- 25 title compound (20.17 g, 93%) was obtained as a light yellow oil: ^1H NMR (300 MHz, CDCl_3) • 7.56 (d, $J = 7.8$ Hz, 1 H), 7.02 - 7.35 (m, 3 H), 6.38 (s, 1 H), 3.88 (s, 2 H), 3.75 (s, 3 H), 2.49 (s, 3 H).

Preparation 2

30

Preparation of 2-[2-(4-methoxyphenyl)ethenyl]nicotinic acid

a) 2-[2-(4-Methoxyphenyl)ethenyl]nicotinic acid

- 35 A mixture of NaH (60% dispersion in mineral oil, 7.4 g, 0.185 mole), *tert*-butanol (11.12 g, 0.15 mole), and anhydrous DMF (100 mL) was carefully warmed at 80 °C until gas evolution ceased, then was cooled to 0 °C under argon. A solution of ethyl 2-methylnicotinate (7.7 mL, 0.050 mole) in anhydrous DMF (17 mL) was added dropwise

70 66

over 3 min, and the reddish-orange mixture was stirred at 0 °C for 1 hr. A solution of *para*-anisaldehyde (7.3 mL, 0.060 mL) in anhydrous DMF (17 mL) was added dropwise over 3 min, and the reaction was allowed to warm to RT. The mixture was stirred overnight, then was poured into ice water (200 mL), and the solution was acidified to pH 6 with glacial acetic acid. The volume was adjusted to 1 L with H₂O, the flask was scratched with a glass rod, and the mixture was allowed to stand at RT for several hr, then in the refrigerator overnight. The solid was collected by suction filtration and washed with H₂O. Recrystallization from boiling absolute EtOH gave the title compound (5.31 g, 42%, two crops) as yellow needles: ¹H NMR (250 MHz, DMSO-d₆) • 8.70 (dd, J = 4.6, 1.7 Hz, 1 H), 8.18 (dd, J = 7.9, 1.7 Hz, 1 H), 7.95 (d, J = 15.8 Hz, 1 H), 7.83 (d, J = 15.8 Hz, 1 H), 7.57 (d, J = 8.7 Hz, 2 H), 7.34 (dd, J = 7.9, 4.6 Hz, 1 H), 6.99 (d, J = 8.7 Hz, 2 H), 3.79 (s, 3 H); MS (ES) *m/e* 256 (M + H)⁺.

b) 2-[2-(4-Methoxyphenyl)ethyl]nicotinic acid
10% Pd/C (2.21 g, 2.08 mmole) was added to a solution of 2-[2-(4-methoxyphenyl)ethenyl]nicotinic acid (5.31 g, 20.80 mmole) and 1.0 N NaOH (23 mL, 23 mmole) in H₂O (81 mL), and the mixture was shaken at RT under H₂ (50 psi) on a Parr apparatus. After 4.5 hr, the mixture was filtered through celite®, and the filtrate was acidified to pH 5 with glacial acetic acid. The solid was collected by suction filtration, washed with H₂O, and dried in high vacuum at 45 °C to afford the title compound (4.50 g, 84%) as an off-white solid: ¹H NMR (400 MHz, DMSO-d₆) • 8.65 (dd, J = 4.7, 1.7 Hz, 1 H), 8.15 (dd, J = 7.8, 1.7 Hz, 1 H), 7.35 (dd, J = 7.8, 4.7 Hz, 1 H), 7.13 (d, J = 8.5 Hz, 2 H), 6.84 (d, J = 8.5 Hz, 2 H), 3.71 (s, 3 H), 3.25 - 3.40 (m, 2 H), 2.82 - 2.93 (m, 2 H).

Preparation 3

Preparation of 2-bromo-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide

a) 2-Bromo-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide - Method I
A solution of 2-bromobenzoyl chloride (1.85 g, 8.45 mmole) in CH₂Cl₂ (10 mL) was added dropwise to a solution of 1-methyl-2-(methylaminomethyl)indole (1.47 g, 8.45 mmole) and Et₃N (3.6 mL, 25.8 mmole) in CH₂Cl₂ (50 mL) at 0 °C. The reaction was stirred at 0 °C for 15 min, then was allowed to warm to RT. After 2 hr, the mixture was concentrated in vacuo, and the residue was diluted with H₂O. The mixture was extracted with Et₂O, and the combined Et₂O layers were dried (MgSO₄) and concentrated in vacuo. The oily residue was triturated with Et₂O/petroleum ether to afford the title compound (2.6 g, 87%) as a white solid: MS (ES) *m/e* 357 (M + H)⁺.

60152707-100899

b) 2-Bromo-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide - Method II

EDC (660 mg, 3.44 mmole) was added to a solution of 2-bromobenzoic acid (693 mg, 3.45 mmole), 1-methyl-2-(methylaminomethyl)indole (600 mg, 3.45 mmole), and HOBt · H₂O (466 mg, 3.45 mmole) in DMF at RT. The reaction was stirred overnight, then was concentrated in vacuo. The residue was diluted with 5% NaHCO₃ and extracted with CH₂Cl₂. The combined organic extracts were washed with brine and dried over MgSO₄. Flash chromatography on silica gel (2% MeOH/CH₂Cl₂) gave the title compound (750 mg, 63%): MS (ES) *m/e* 357 (M + H)⁺.

Example 1

Preparation of 2-[2-(4-methoxyphenyl)ethyl]-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]nicotinamide

a) 2-[2-(4-Methoxyphenyl)ethyl]-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]nicotinamide

EDC (0.60 g, 3.16 mmole) was added to a solution of 2-[2-(4-methoxyphenyl)ethyl]nicotinic acid (0.81 g, 3.16 mmole), 1-methyl-2-(methylaminomethyl)indole (0.50 g, 2.87 mmole), HOBt · H₂O (0.43 g, 3.16 mmole), and diisopropylethylamine (0.55 mL, 3.16 mmole) in DMF (15 mL) at RT. The reaction was stirred overnight, then was diluted with H₂O and extracted with EtOAc. Drying (Na₂SO₄), concentration, and flash chromatography on silica gel (5% MeOH/CHCl₃) gave the title compound (1.07 g, 90%) as a viscous yellow oil: ¹H NMR (300 MHz, DMSO-d₆) δ 8.63 (m, 1 H), 7.22 - 7.61 (m, 8 H), 6.97 (d, J = 8.1 Hz, 1 H), 6.63 (d, J = 8.1 Hz, 1 H), 6.52 (s, 1 H), 3.80 (s, 3 H), 3.78 (m, 2 H), 3.72 (s, 2 H), 3.16 (m, 2 H), 3.03 (s, 3 H), 2.57 (s, 3 H); MS (ES) *m/e* 414 (M + H)⁺. Anal. Calcd for C₂₆H₂₇N₃O₂: C, 75.52; H, 6.58; N, 10.16. Found C, 75.84; H, 6.68; N, 9.76.

Example 2

Preparation of N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenylbenzamide

a) N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenylbenzamide

EDC (0.22 g, 1.14 mmole) was added to a solution of 2-biphenylcarboxylic acid (0.22 g, 1.14 mmole), 1-methyl-2-(methylaminomethyl)indole (0.20 g, 1.15 mmole), and HOBt · H₂O (0.15 g, 1.11 mmole) in DMF (20 mL) at RT. The reaction was stirred

60152707-100899

7F 268

- overnight, then was concentrated in vacuo. The residue was diluted with 5% NaHCO₃ and extracted with CH₂Cl₂. The combined organic extracts were washed with brine and dried over MgSO₄. Flash chromatography on silica gel (3% MeOH/CH₂Cl₂) followed by preparative TLC (3% MeOH/CH₂Cl₂) gave the title compound (0.10 g, 25%) as a light yellow solid: ¹H NMR (360 MHz, CDCl₃) indicated an approximately 5:1 mixture of amide rotamers; for the major rotamer: • 7.03 - 7.58 (m, 13 H), 6.19 (s, 1 H), 4.50 - 5.00 (m, 2 H), 3.57 (s, 3 H), 2.40 (s, 3 H); for the minor rotamer: • 7.03 - 7.58 (m, 13 H), 6.14 (s, 1 H), 4.50 - 5.00 (m, 2 H), 3.38 (s, 3 H), 2.76 (s, 3 H); MS (ES) *m/e* 356 (M + H)⁺. Anal. Calcd for C₂₄H₂₂N₂O · 0.80 H₂O: C, 78.15; H, 6.45; N, 7.59. Found: C, 77.96; H, 6.29; N, 7.35.

Example 3

- Preparation of N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-4-yl)benzamide
- a) N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-4-yl)benzamide
- A solution of 2-bromo-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide (0.30 g, 0.84 mmole), pyridine-4-boronic acid (0.20 g, 1.63 mmole), tetrakis(triphenylphosphine)palladium(0) (29 mg, 0.025 mmole), and Cs₂CO₃ (1.10 g, 3.38 mmole) in 10:1 DME/H₂O (25 mL) was degassed, then was heated under N₂ overnight. The resulting mixture was concentrated in vacuo, and the residue was taken up in 10% NaOH solution. The mixture was extracted with CH₂Cl₂, and the combined organic extracts were washed with brine and dried (MgSO₄). Flash chromatography on silica gel (3% MeOH/CH₂Cl₂) followed by preparative TLC (3% MeOH/CH₂Cl₂) gave the title compound (98 mg, 33%) as a white solid: ¹H NMR (300 MHz, CDCl₃) indicated an approximately 5:1 mixture of amide rotamers; for the major rotamer: • 8.50 - 8.65 (m, 2 H), 7.00 - 7.60 (m, 10 H), 6.27 (s, 1 H), 4.50 - 5.00 (m, 2 H), 3.62 (s, 3 H), 2.49 (s, 3 H); for the minor rotamer: • 8.65 - 8.75 (m, 2 H), 7.00 - 7.60 (m, 10 H), 6.12 (s, 1 H), 4.50 - 5.00 (m, 2 H), 3.41 (s, 3 H), 2.86 (s, 3 H); MS (ES) *m/e* 356 (M + H)⁺. Anal. Calcd for C₂₃H₂₁N₃O · 0.10 H₂O: C, 77.33; H, 5.98; N, 11.76. Found: C, 77.07; H, 5.98; N, 11.56.

60158707.100899

75 69
KJExample 4Preparation of N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-3-yl)benzamide

5 a) N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-3-yl)benzamide

According to the procedure of Example 3 (a), except substituting pyridine-3-boronic acid (173 mg, 1.41 mmole) for the pyridine-4-boronic acid, the title compound (56 mg, 22%) was obtained as a light yellow foam after flash chromatography on silica gel (3% MeOH/CH₂Cl₂) followed by preparative TLC (3% MeOH/CH₂Cl₂): ¹H NMR (360 MHz, CDCl₃) indicated an approximately 5:1 mixture of amide rotamers; for the major rotamer: • 8.68 (br s, 1 H), 8.50 (br s, 1 H), 7.71 - 7.77 (m, 1 H), 7.63 - 7.72 (m, 1 H), 7.31 - 7.57 (m, 4 H), 7.04 - 7.31 (m, 4 H), 6.23 (s, 1 H), 4.60 - 4.85 (m, 2 H), 3.63 (s, 3 H), 2.48 (s, 3 H); for the minor rotamer: • 8.74 (br s, 1 H), 8.68 (br s, 1 H), 7.85 - 7.89 (m, 1 H), 7.31 - 7.57 (m, 5 H), 7.04 - 7.31 (m, 4 H), 6.17 (s, 1 H), 4.30 - 4.45 (m, 2 H), 3.40 (s, 3 H), 2.82 (s, 3 H); MS (ES) m/z 356 (M + H)⁺.

Example 5Preparation of N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(thiophen-3-yl)benzamide

20 a) N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(thiophen-3-yl)benzamide

According to the procedure of Example 3 (a), except substituting thiophene-3-boronic acid (179 mg, 1.40 mmole) for the pyridine-4-boronic acid, the title compound (100 mg, 40%) was obtained following flash chromatography on silica gel (20% EtOAc/hexanes): ¹H NMR (360 MHz, CDCl₃) indicated an approximately 4:1 mixture of amide rotamers; for the major rotamer: • 7.00 - 7.56 (m, 11 H), 6.28 (s, 1 H), 4.95 - 5.15 (m, 1 H), 4.48 - 4.68 (m, 1 H), 3.69 (s, 3 H), 2.41 (s, 3 H); for the minor rotamer: • 7.00 - 7.56 (m, 11 H), 6.15 (s, 1 H), 4.38 (d, J = 15.5 Hz, 1 H), 3.65 (d, J = 15.5 Hz, 1 H), 3.60 (s, 3 H), 2.84 (s, 3 H); MS (ES) m/z 361 (M + H)⁺. Anal. Calcd for C₂₂H₂₀N₂OS • 0.10 H₂O: C, 72.94; H, 5.62; N, 7.73. Found: C, 72.70; H, 5.58; N, 7.46.

66807.100399

Example 6Preparation of N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(furan-3-yl)benzamide

5 a) N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(furan-3-yl)benzamide

According to the procedure of Example 3 (a), except substituting furan-3-boronic acid (156 mg, 1.39 mmole) for the pyridine-4-boronic acid, the title compound (100 mg, 41%) was obtained after preparative TLC (20% EtOAc/hexanes) followed by flash chromatography on silica gel (20% EtOAc/hexanes): ¹H NMR (360 MHz, CDCl₃)

- 10 indicated an approximately 3.5:1 mixture of amide rotamers; for the major rotamer: • 7.61 - 7.65 (m, 1 H), 7.03 - 7.58 (m, 9 H), 6.55 - 6.58 (m, 1 H), 6.37 (s, 1 H), 5.07 - 5.27 (m, 1 H), 4.58 - 4.68 (m, 1 H), 3.74 (s, 3 H), 2.50 (s, 3 H); for the minor rotamer: • 7.68 - 7.71 (m, 1 H), 7.03 - 7.58 (m, 9 H), 6.63 - 6.66 (m, 1 H), 6.24 (s, 1 H), 4.42 (d, J = 15.7 Hz, 1 H), 3.97 (d, J = 15.7 Hz, 1 H), 3.39 (s, 3 H), 2.94 (s, 3 H); MS (ES) *m/e* 345 (M + H)⁺. Anal.
- 15 Calcd for C₂₂H₂₀N₂O₂ · 0.10 H₂O: C, 76.32; H, 5.88; N, 8.09. Found: C, 76.11; H, 5.94; N, 7.79.

Example 720 Preparation of N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenoxybenzamide

a) N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenoxybenzamide

EDC (230 mg, 1.2 mmole) was added to a solution of 2-phenoxybenzoic acid (214.2 mg, 1.0 mmole), 1-methyl-2-(methylaminomethyl)indole (209.1 mg, 1.2 mmole),

25 HOBt · H₂O (162.2 mg, 1.2 mmole), and triethylamine (0.35 mL, 2.5 mmole) in DMF (5 mL) at RT. After 16.5 hr, the reaction was concentrated, and the residue was reconcentrated from xylenes/CHCl₃ to remove residual DMF. Flash chromatography on silica gel (40% EtOAc/hexanes) gave the title compound (393.5 mg, quantitative) as a nearly colorless, very viscous oil: ¹H NMR (360 MHz, CDCl₃) indicated an approximately

- 30 5:1 mixture of amide rotamers; for the major rotamer: • 7.56 (d, J = 7.9 Hz, 1 H), 6.98 - 7.50 (m, 9 H), 6.95 (d, J = 7.6 Hz, 2 H), 6.87 (d, J = 7.5 Hz, 1 H), 6.49 (s, 1 H), 4.40 - 5.40 (m, 2 H), 3.62 (s, 3 H), 2.84 (s, 3 H); for the minor rotamer: 6.80 - 7.50 (m, 13 H), 6.47 (s, 1 H), 4.40 - 5.40 (m, 2 H), 3.56 (s, 3 H), 2.98 (s, 3 H); MS (ES) *m/e* 372 (M + H)⁺.

60158707.100899

71
2

Example 8

Preparation of N,2-dimethyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide

5 a) N,2-Dimethyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide

EDC (0.22 g, 1.15 mmole) was added to a solution of *ortho*-toluic acid (0.15 g, 1.10 mmole), 1-methyl-2-(methylaminomethyl)indole (0.20 g, 1.15 mmole), and HOBT · H₂O (0.15 g, 1.11 mmole) in DMF at RT. The reaction was stirred overnight, then was concentrated in vacuo. The residue was diluted with 5% NaHCO₃ and extracted with
10 CH₂Cl₂. The combined organic extracts were washed with brine and dried over MgSO₄. Flash chromatography on silica gel (20% EtOAc/hexanes) gave a colorless gum, which was triturated with Et₂O to afford the title compound (0.12 g, 37%) as a white solid: ¹H NMR (300 MHz, CDCl₃) indicated an approximately 5:1 mixture of amide rotamers; for the major rotamer: • 7.60 (d, J = 7.7 Hz, 1 H), 7.05 - 7.40 (m, 7 H), 6.52 (s, 1 H), 5.01 (s, 2 H),
15 3.85 (s, 3 H), 2.71 (s, 3 H), 2.30 (s, 3 H); for the minor rotamer: • 7.60 (d, J = 7.7 Hz, 1 H), 7.05 - 7.40 (m, 7 H), 6.44 (s, 1 H), 4.52 (s, 2 H), 3.50 (s, 3 H), 3.14 (s, 3 H), 2.37 (s, 3 H); MS (ES) *m/e* 293 (M + H)⁺. Anal. Calcd for C₁₉H₂₀N₂O · 0.35 H₂O: C, 76.40; H, 6.99; N, 9.38. Found: C, 76.10; H, 6.83; N, 9.33.

20

Example 9

Preparation of 2-[3-(acetylamino)phenyl]-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide

25 a) 2-[3-(Acetylamino)phenyl]-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide

According to the procedure of Example 3 (a), except substituting 3-(acetylamino)phenylboronic acid (250 mg, 1.40 mmole) for the pyridine-4-boronic acid, the title compound (190 mg, 66%) was obtained after flash chromatography on silica gel (3% MeOH/CH₂Cl₂): ¹H NMR (300 MHz, CDCl₃) indicated an approximately 4:1 mixture of
30 amide rotamers; for the major rotamer: • 7.00 - 7.80 (m, 12 H), 6.20 (s, 1 H), 4.65 - 4.90 (m, 2 H), 3.54 (s, 3 H), 2.52 (s, 3 H), 2.04 (s, 3 H); for the minor rotamer: • 7.00 - 7.80 (m, 12 H), 6.12 (s, 1 H), 4.40 (d, J = 15 Hz, 1 H), 3.75 (d, J = 15 Hz, 1 H), 3.38 (s, 3 H), 2.84 (s, 3 H), 2.16 (s, 3 H); MS (ES) *m/e* 412 (M + H)⁺. Anal. Calcd for C₂₆H₂₅N₃O₂ · 0.20 H₂O: C, 74.87; H, 6.42; N, 9.77. Found: C, 74.65; H, 6.36; N, 9.50.

35

668007-100899

72
XO
ADExample 10Parenteral Dosage Unit Composition

- 5 A preparation which contains 20 mg of the compound of Example 1 as a sterile dry powder is prepared as follows: 20 mg of the compound is dissolved in 15 mL of distilled water. The solution is filtered under sterile conditions into a 25 mL multi-dose ampoule and lyophilized. The powder is reconstituted by addition of 20 mL of 5% dextrose in water (D5W) for intravenous or intramuscular injection. The dosage is thereby determined by the
- 10 injection volume. Subsequent dilution may be made by addition of a metered volume of this dosage unit to another volume of D5W for injection, or a metered dose may be added to another mechanism for dispensing the drug, as in a bottle or bag for IV drip infusion or other injection-infusion system.

15

Example 11Oral Dosage Unit Composition

- 20 A capsule for oral administration is prepared by mixing and milling 50 mg of the compound of Example 1 with 75 mg of lactose and 5 mg of magnesium stearate. The resulting powder is screened and filled into a hard gelatin capsule.

Example 12Oral Dosage Unit Composition

- 25 A tablet for oral administration is prepared by mixing and granulating 20 mg of sucrose, 150 mg of calcium sulfate dihydrate and 50 mg of the compound of Example 1 with a 10% gelatin solution. The wet granules are screened, dried, mixed with 10 mg
- 30 starch, 5 mg talc and 3 mg stearic acid; and compressed into a tablet.


- 35 The above description fully discloses how to make and use the present invention. However, the present invention is not limited to the particular embodiments described hereinabove, but includes all modifications thereof within the scope of the following claims. The various references to journals, patents and other publications which are cited herein comprises the state of the art and are incorporated herein by reference as though fully set forth.

60158707.100899

73
AC

What is claimed is:

1. A compound according to formula (I):




wherein:

R¹ is C₁₋₄alkyl;

R² is C₁₋₄alkyl;

R³ is

10



Q is a single bond, -O-, -S- or -NH-, or -CH₂CH₂-;


R⁴ is C₁₋₆alkyl;



is a five- or six-membered heteroaromatic ring or a six-membered

15

aromatic ring optionally substituted by R⁵;



is a five- or six-membered heteroaromatic ring or a six-membered

aromatic ring optionally substituted by R⁶;



is a five- or six-membered heteroaromatic ring or a six-membered

aromatic ring optionally substituted by R⁶;

20

R⁵ is OR' or NR'C(O)R';

R⁶ is OR' or NH₂;

668001-40485F09

70 74


X is H, C₁₋₄alkyl, OR', SR', CN, N(R')₂, CH₂N(R')₂, NO₂, CF₃, CO₂R', CON(R')₂, COR', NR'C(O)R', F, Cl, Br, I or -S(O)_rCF₃;

R' is H, C₁₋₆alkyl or -C₀₋₆alkyl-Ar; and

r is 0, 1 or 2;


5 or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 of formula (Ia):



(Ia).

3. A compound according to claim 1 in which R³ is:




4. A compound according to claim 3 in which  is 2-, 3- or 4-pyridyl, 2- or 3-furanyl, 2- or 3-thiophenyl, thiazolyl, phenyl or phenyl substituted by OC₁₋₄alkyl or NHC(O)C₁₋₄alkyl.

5. A compound according to claim 4 in which  is 3- or 4-pyridyl, 3-furanyl, 3-thiophenyl, phenyl or phenyl substituted by OCH₃ or NHC(O)CH₃.

50158707.100899

75
25


6. A compound according to claim 3 in which  is 2-, 3- or 4-pyridyl, phenyl or phenyl substituted by OH or NH₂.


7. A compound according to claim 6 in which  is phenyl or 2-pyridyl.

8. A compound according to claim 3 in which Q is a single bond.


9. A compound according to claim 3 in which Q is -O- or -CH₂CH₂.

10. A compound according to claim 1 in which R³ is:



11. A compound according to claim 10 in which  is phenyl unsubstituted or substituted by OH or NH₂ and R⁴ is C₁₋₄alkyl.

12. A compound according to claim 11 in which  is phenyl and R⁴ is CH₃.

13. A compound according to claim 10 in which  is 2-pyridyl and R⁴ is C₁₋₄alkyl.

14. A compound according to claim 1 which is:

2-[2-(4-Methoxyphenyl)ethyl]-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]nicotinamide;

60158707.100899

N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenylbenzamide;
N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-4-yl)benzamide;
N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(pyridin-3-yl)benzamide;
N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(thiophen-3-yl)benzamide;
5 N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-(furan-3-yl)benzamide;
N-Methyl-N-[(1-methyl-1H-indol-2-yl)methyl]-2-phenoxybenzamide;
N,2-Dimethyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide; or
2-[3-(Acetylamino)phenyl]-N-methyl-N-[(1-methyl-1H-indol-2-yl)methyl]benzamide;

10 or a pharmaceutically acceptable salt thereof.

15. A pharmaceutical composition which comprises a compound according to claim 1 and a pharmaceutically acceptable carrier.


15 16. A method for inhibiting Fab I which comprises administering to a subject in need thereof a compound according to claim 1.

17. A method of treating bacterial infections which comprises administering to a subject in need thereof a compound according to claim 1.

60153707.100899

7701
MABSTRACT OF THE DISCLOSUREFab I Inhibitors

Compounds of the formula (I) are disclosed which are Fab I inhibitors and are useful in the treatment bacterial infections:




(I)

wherein:

R^1 is C_{1-4} alkyl;

R^2 is C_{1-4} alkyl;

R^3 is



Q is a single bond, -O-, -S- or -NH-, or -CH₂CH₂;

R^4 is C_{1-6} alkyl;



is a five- or six-membered heteroaromatic ring or a six-membered aromatic ring optionally substituted by R^5 ;



is a five- or six-membered heteroaromatic ring or a six-membered aromatic ring optionally substituted by R^6 ;

2278
27

is a five- or six-membered heteroaromatic ring or a six-membered aromatic ring optionally substituted by R^6 ;

R^5 is OR' or $NR'C(O)R'$;

R^6 is OR' or NH_2 ;

X is H , $C_{1-4}alkyl$, OR' , SR' , CN , $N(R')_2$, $CH_2N(R')_2$, NO_2 , CF_3 , CO_2R' , $CON(R')_2$, COR' , $NR'C(O)R'$, F , Cl , Br , I or $-S(O)_2CF_3$;

R' is H , $C_{1-6}alkyl$ or $-C_{0-6}alkyl-Ar$; and

r is 0, 1 or 2;

or a pharmaceutically acceptable salt thereof.

60158707-100899

8579
27

ESTADOS UNIDOS DE AMÉRICA

A TODOS A QUIENES LLEGUE EL PRESENTE:

DEPARTAMENTO DE COMERCIO DE LOS ESTADOS UNIDOS

Oficina de Marcas y Patentes

28 de septiembre de 2000

LA PRESENTE CERTIFICA QUE LA ADJUNTA ES UNA COPIA TOMADA DE LOS REGISTROS DE LA OFICINA DE MARCAS Y PATENTES DE LOS ESTADOS UNIDOS, DE LOS DOCUMENTOS DE LA SOLICITUD DE PATENTE ABAJO IDENTIFICADA Y LA CUAL CUMPLE CON LOS REQUISITOS PARA QUE LE SEA CONFERIDA UNA FECHA DE RADICACIÓN SEGÚN 35 USC 111.

NÚMERO DE SOLICITUD 60/158,707

FECHA DE RADICACIÓN: 08 de octubre de 1999

Por Autoridad del

COMISIONADO DE MARCAS Y PATENTES

(FIRMA) T. WALLACE

Oficial de Certificaciones


(SELLO DORADO DE OBLEA)

80/158
14

INHIBIDORES FAB I

REIVINDICACIONES:

1. Un compuesto según la Fórmula (I):




(I)

en donde:

R¹ es alquil C₁₋₄;

R² es alquil C₁₋₄;

R³ es




Q es un enlace simple, -O-, -S- o -NH-, o -CH₂CH₂-;

R⁴ es alquil C₁₋₆;



2581/17

es un anillo heteroaromático de 5 o 6 miembros o un anillo aromático de seis miembros sustituido opcionalmente por R⁵,



es un anillo heteroaromático de 5 ó 6 miembros o un anillo aromático de 6 miembros sustituido por R⁶,



es un anillo heteroaromático de 5 ó 6 miembros o un anillo aromático de 6 miembros sustituido por R⁶,

R⁵ es OR' o NR'C(O)R,

R⁶ es OR' o NH₂,


X es H, alquil C₁₋₄, OR', SR', CN, N(R')₂, CH₂N(R')₂, NO₂, CF₃, CO₂R', CON(R')₂, COR', NR'C(O)R', F, Cl, Br o -S(O)_rCF₃,

R' es H, alquil C₁₋₆ o alquil C₀₋₆-Ar, y

R es 0, 1 o 2,

o una sal farmacéuticamente aceptable de lo mismo


2 Un compuesto según la reivindicación 1 de la Fórmula (Ia)



(Ia)

82
13


3. Un compuesto según la reivindicación 1, en el cual R³ es:



4. Un compuesto según la reivindicación 3, en el cual
A es 2-, 3- o 4- piridil, 2 o 3-furanil, 2 o 3-tiofenil, tiazolil, fenil o fenil sustituido por alquil OC₁₋₄ o NHC(O)alquilC₁₋₄.
5. Un compuesto según la reivindicación 4 en el cual
A es 3- o 4- piridil, 3-furanil, 3-tiofenil, fenil o fenil sustituido por OCH₃ o NHC(O)CH₃
6. Un compuesto según la reivindicación 3, en el cual
B es 2-, 3- o 4- piridil, fenil o fenil sustituido por OH o NH₂.
7. Un compuesto según la reivindicación 6, en el cual
B es fenil o 2-piridil
8. Un compuesto según la reivindicación 3, en el cual Q es un enlace simple.
9. Un compuesto según la reivindicación 3, en el cual Q es -O- o -CH₂CH₂-.

83
14

10. Un compuesto según la reivindicación 1, en el cual R^3 es:



11. Un compuesto según la reivindicación 10, en el cual  es fenil sustituido o insustituido por OH o NH_2 y R^4 es alquil C_{1-4} .

12. Un compuesto según la reivindicación 11, en el cual  es fenil y R^4 es CH_3 .

13. Un compuesto según la reivindicación 10, en el cual  es 2-piridil y R^4 es alquil C_{1-4} .

14. Un compuesto según la reivindicación 1 que es:

2-[2-(4-Metoxifenil)etil]-N-metil-N-[(1-metil-1H-Indol-2-il)metil]nicotinamida;

N-Metil-N-[(1-metil-1H-Indol-2-il)metil]-2-fenilbenzamida;

N-Metil-N-[(1-metil-1H-Indol-2-il)metil]-2-(piridin-4-il)benzamida;

N-Metil-N-[(1-metil-1H-Indol-2-il)metil]-2-(piridin-3-il)benzamida;

N-Metil-N-[(1-metil-1H-Indol-2-il)metil]-2-(tiofen-3-il)benzamida;

N-Metil-N-[(1-metil-1H-Indol-2-il)metil]-2-(furan-3-il)benzamida;

N-Metil-N-[(1-metil-1H-Indol-2-il)metil]-2-fenoxibenzamida;

N,2-Dimetil-N-[(1-metil-1H-2-il)metil]benzamida, o

2-[3-(Acetilamino)fenil]-N-metil-N-[(1-metil-1H-Indol-2-il)metil]benzamida,

o una sal farmacéuticamente aceptable de lo mismo

15 Una composición farmacéutica que comprende un compuesto según la reivindicación 1, y un portador farmacéuticamente aceptable

16 Un método para Inhibir Fab I que consiste en administrar a un sujeto que lo requiera, un compuesto según la reivindicación 1

17 Un método para tratar infecciones bacterianas que consiste en administrar a un sujeto que lo requiera, un compuesto según la reivindicación 1