

ALVARO CASTELLANOS M. & CIA
ABONDOS
Carrera 9 No. 93 - 09
A.A. 6348
Santa Fe de Bogotá, D.C., COLOMBIA

9/2
56

SUPERINDUSTRIA Y COMERCIO
Indicacion : 01003045 00000002 Folios: 42
Fecha (ADM): 2001-04-10 14:38:16
Tramite : 002 PATENTES D 1 REGISTRAR 329 COMPLETAR
Dependencia: 2028 DIVISION DE NUEVAS CREACIONES

Señora

Jefe de la División de Nuevas Creaciones

Superintendencia de Industria y Comercio

Ministerio de Desarrollo Económico

E.

S.

D.

Ref.: Expediente No. 01.003.045

Solicitud de Patente de Invención titulada : "DROSPIRENONA PARA
TERAPIA DE REEMPLAZO DE HORMONAS".

Solicitante: SCHERING AKTIENGESELLSCHAFT

Yo, MARGARITA CASTELLANOS ABONDANO, mayor de edad,
Identificada y domiciliada como aparece al pie de mi firma, atentamente me
permito presentar a usted la copia certificada del documento de Prioridad
Europeo No. 00200183.2, en donde se certifica así mismo la fecha de
presentación de esa solicitud, junto con su traducción simple de carátula y
reivindicaciones.

En consecuencia atentamente solicito a Ud. se anexe dicho documento al
expediente correspondiente a esta solicitud de Patente de Invención.

Del Señor Jefe,


Margarita Castellanos Abondano
C.C. No. 39.779.654 de Usaquén
T.P.A. No. 70819 del Consejo Sup. de la Judicatura
Carrera 9ª. No. 93 - 09 - de Bogotá
Teléfonos: 610 74 80 y 610 74 60

Santa Fe de Bogotá, D.C.,
xgt

10 ABR. 2001

45
57

LA OFICINA DE PATENTES EUROPEA

Certifica: que los documentos anexos son copias exactas de la solicitud de Patente Europea descrita en la siguiente página, tal como originalmente se presentó.

Número de Solicitud: 00200183.2

Por el presidente de la Oficina de Patentes Europea

I.L.C. HATTEN-HECKMAN

LA HAYA, 18/01/01

Forma/EPO 1014

8
58

LA OFICINA DE PATENTES EUROPEA

Hoja No.2 de la certificación

Número de Solicitud: 00200183.2 Fecha de presentación 18/01/00

Solicitante:
SCHERING AKTIENGESELLSCHAFT
13342 Berlín
ALEMANIA

Título de la invención:
“DROSPIRENONA PARA TERAPIA DE REEMPLAZO DE
HORMONAS”.

Prioridad(es) reivindicadas
País Fecha.....Número de presentación

Clasificación Internacional de Patentes:
A61K31/585, A61P5/30

Contratación de los estados designados a la fecha de presentación:
AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE

Comentarios: VER por título en la página 1 de la descripción

Forma/EPO 1014

6/2
59

REIVINDICACIONES

1. Una composición farmacéutica que comprende como un primer agente activo, un estrógeno (o derivado natural o sintético del mismo) en cantidades suficientes para tratar enfermedades, desórdenes y síntomas asociados con niveles endógenos deficientes de estrógeno en mujeres, y como un segundo agente activo, $6\beta,7\beta;15\beta;16\beta$ -dimetilen-3-oxo-17 α -preg-4-eno-21,17-carbolactona (drospirenona) en cantidades suficientes para proteger el endometrio de los efectos adversos del estrógeno, conjuntamente con un excipiente o portador farmacéuticamente aceptable.
2. Una composición de acuerdo a la reivindicación 1, donde los niveles deficientes de estrógenos son causados por menopausia natural, perimenopausia, post-menopausia, hipogonadismo, castración o falla ovárica primaria.
3. Una composición de acuerdo a cualquiera de las reivindicaciones 1 ó 2, donde las enfermedades, desórdenes y síntomas son seleccionados del grupo que comprende incluyendo acaloramientos, ataques de transpiración, palpitaciones, desórdenes del sueño, cambios de humor, nerviosismo, ansiedad, mala memoria, pérdida de confianza, pérdida de libido, concentración pobre, energía disminuida, empuje disminuido, irritabilidad, atrofia urogenital, atrofia de los senos, enfermedad cardiovascular, cambios en la distribución del vello, espesor del cabello, cambios en el estado de la piel y osteoporosis.
4. Una composición de acuerdo a la reivindicación 3, donde las enfermedades, desórdenes y síntomas son seleccionados del grupo que comprende incluyendo acaloramientos, ataques de transpiración, palpitaciones, desórdenes del sueño, cambios de humor, nerviosismo, ansiedad, atrofia urogenital, atrofia de los senos y osteoporosis.

9/8

60

5. Una composición de acuerdo a la reivindicación 1, donde el estrógeno se selecciona del grupo que consiste de estradiol, sulfamatos de estradiol, valerato de estradiol, benzoato de estradiol, etinil estradiol, estrona, estriol, succinato de estriol y estrógenos conjugados, que incluyen estrógenos equino conjugados tales como sulfato de estrona, sulfato de 17β -estradiol, sulfato de 17α -estradiol, sulfato de equilino, sulfato de 17β -dihidroequilino, sulfato de 17α -dihidroequilino, sulfato de equilenino, sulfato de 17β -dihidroequilenino y sulfato de 17α -dihidroequilenino o mezclas de los mismos.
6. Una composición de acuerdo a la reivindicación 5, donde el estrógeno se selecciona del grupo que consiste de estradiol, sulfamatos de estradiol, valerato de estradiol, benzoato de estradiol, estrona, y sulfato de estrona o mezclas de los mismos.
7. Una composición de acuerdo a la reivindicación 6, donde el estrógeno es estradiol.
8. Una composición de acuerdo a la reivindicación 1, donde la drospirenona (DRSP) está en forma micronizada.
9. Una composición de acuerdo a cualquiera de las reivindicaciones 5 ó 6, donde al menos un estrógeno está en forma micronizada.
10. Una composición de acuerdo a la reivindicación 1, donde la dosis de DRSP corresponde a 15 a 70 mg por ciclo, tal como 20 a 60 mg por ciclo, particularmente 40 a 60 mg por ciclo.
11. Una composición de acuerdo a la reivindicación 1, donde la cantidad de DRSP corresponde a una dosis diaria en el rango de desde 0,25 a 10 mg, tal como aproximadamente 0,25 a 8, 0,25 a 6, 0,25 a 5, 0,5 a 4,5, 1 a 4, y 1,5 a 3,5 mg.

12. Una composición de acuerdo a la reivindicación 7, donde la cantidad de estradiol corresponde a una dosis diaria en el rango de desde 0,1 a 5 mg, tal como aproximadamente 0,2 a 4,5, 0,5 a 4, 1 a 3, en particular 1, 2 ó 3 mg.

13. Una composición farmacéutica que comprende estradiol como un primer agente activo en cantidades que corresponden a una dosis diaria de 1 a 3 mg para tratar enfermedades, desórdenes y síntomas asociados con niveles endógenos deficientes de estrógeno en mujeres, y como un segundo agente activo $6\beta,7\beta;15\beta;16\beta$ -dimetilen-3-oxo-17 α -preg-4-eno-21,17-carbolactona (drospirona) en cantidades suficientes que corresponden a una dosis diaria de 1 a 3,5 mg para proteger el endometrio de los efectos adversos del estrógeno, junto con un excipiente o portador farmacéuticamente aceptable.

14. Uso de una combinación de estrógeno y drospirona para la preparación de un medicamento en donde la cantidad de estrógeno es suficiente para tratar enfermedades, desórdenes y síntomas asociados con niveles endógenos deficientes de estrógeno en mujeres y la cantidad de drospirona es suficiente para proteger el endometrio de los efectos adversos del estrógeno.

15. El uso de acuerdo a la reivindicación 14, donde los niveles deficientes de estrógeno son causados por menopausia natural, peri-menopausia, post-menopausia, hipogonadismo, castración o falla ovárica primaria.

16. El uso de acuerdo a la reivindicación 14, donde la cantidad de estrógeno es suficiente para el tratamiento de acaloramiento, ataque de transpiración, palpitaciones, desórdenes del sueño, cambios de humor, nerviosismo, ansiedad, mala memoria, pérdida de confianza, pérdida de libido, concentración pobre, energía disminuida, empuje disminuido, irritabilidad, atrofia urogenital, atrofia de

los senos, enfermedad cardiovascular, cambios en la distribución del vello, espesor del cabello, cambios en el estado de la piel o para la prevención o control de osteoporosis.

17. El uso de acuerdo a la reivindicación 16, donde la cantidad de estrógeno es suficiente para el tratamiento de acaloramientos, ataques de transpiración, palpitaciones, desórdenes del sueño, cambios de humor, nerviosismo, ansiedad, atrofia urogenital, atrofia de los senos o para la prevención o control de osteoporosis.

18. El uso de acuerdo a la reivindicación 14, donde el estrógeno es seleccionado del grupo que consiste de estradiol, sulfamatos de estradiol, valerato de estradiol, benzoato de estradiol, etinil estradiol, estrona, estriol, succinato de estriol y estrógenos conjugados, que incluyen estrógenos equino conjugados tal como sulfato de estrona, sulfato de 17 β -estradiol, sulfato de 17 α -estradiol, sulfato de equilino, sulfato de 17 β -dihidroequillino, sulfato de 17 α -dihidroequillino, sulfato de equilenino, sulfato de 17 β -dihidroequillenino y sulfato de 17 α -dihidroequillenino o mezclas de los mismos.

19. El uso de acuerdo a la reivindicación 18, donde el estrógeno se selecciona del grupo que consiste de estradiol, sulfamatos de estradiol, valerato de estradiol, benzoato de estradiol, estrona, y sulfato de estrona o mezclas de los mismos.

20. El uso de acuerdo a la reivindicación 19, donde el estrógeno es estradiol.

21. El uso de acuerdo a la reivindicación 14, donde la drospirenona (DRSP) está en forma micronizada.

22. El uso de acuerdo a la reivindicación 14, donde el estrógeno está en forma micronizada.

23. El uso de acuerdo a la reivindicación 20, donde el estradiol está en forma micronizada.

24. El uso de acuerdo a la reivindicación 14, donde la dosis de DRSP corresponde a 15 a 70 mg por ciclo, tal como 20 a 60 mg por ciclo, particularmente 40 a 60 mg por ciclo.

25. El uso de acuerdo a la reivindicación 14, donde la cantidad de DRSP corresponde a una dosis diaria en el rango de desde 0,25 a 10 mg, tal como aproximadamente 0,25 a 8, 0,25 a 6, 0,25 a 5, 0,5 a 4,5, 1 a 4, y 1,5 a 3,5 mg.

26. El uso de acuerdo a la reivindicación 20, donde la cantidad de estradiol corresponde a una dosis diaria en el rango de desde 0,1 a 5 mg, tal como aproximadamente 0,2 a 4,5, 0,5 a 4, 1 a 3, en particular 1, 2 ó 3 mg.

27. El uso de acuerdo a la reivindicación 14, donde el medicamento está en la forma de un número de unidades de dosificación envasadas separadamente e individualmente removibles dispuestas en una unidad de envase y destinadas a administración oral por un período de al menos 28 días;

donde al menos 21 unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y 7 o menos unidades de dosificación diarias comprenden estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg.

28. El uso de acuerdo a la reivindicación 14, donde el medicamento está en la forma de un número de unidades de dosificación envasadas separadamente e individualmente removibles dispuestas en una unidad de envase y destinadas a administración oral por un período de al menos 28 días;

donde al menos 21 unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y 7 o menos unidades de dosificación diarias comprenden un blanco o placebo.

29. El uso de acuerdo a la reivindicación 14, donde el medicamento está en la forma de un número de unidades de dosificación envasadas separadamente e individualmente removibles dispuestas en una unidad de envase y destinadas a administración oral por un período de al menos 28 días,

donde al menos 10 unidades de dosificación diarias comprenden estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg;

y al menos 10 unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg, y 8 o menos unidades de dosificación diarias comprenden un blanco o placebo.

30. El uso de acuerdo a la reivindicación 14, donde el medicamento está en la forma de un número de unidades de dosificación envasadas separadamente e individualmente removibles dispuestas en una unidad de envase y destinadas a administración oral por un período de al menos 28 días;

donde al menos 10 unidades de dosificación diarias comprenden estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg;

y al menos 10 unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg, y 8 o menos

unidades de dosificación diarias comprenden estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg.

31. El uso de acuerdo a la reivindicación 14, donde el medicamento está en la forma de un número de unidades de dosificación envasadas separadamente e individualmente removibles dispuestas en una unidad de envase y destinadas a administración oral por un período de al menos 28 días;

donde 28 unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg.

32. Un método para tratar o prevenir enfermedades, desórdenes y síntomas asociados con niveles endógenos deficientes de estrógeno en mujeres que comprende administrar estrógeno en cantidades suficientes para mitigar dichas enfermedades, desórdenes y síntomas y drospirenona en cantidades suficientes para proteger el endometrio de efectos adversos del estrógeno.

33. Un método de acuerdo a la reivindicación 32, donde los niveles deficientes de estrógeno están causados por menopausia natural, perimenopausia, post-menopausia, hipogonadismo, castración o falla ovárica primaria.

34. Un método de acuerdo a la reivindicación 32, donde las enfermedades, desórdenes y síntomas están seleccionados del grupo que comprende incluyendo acaloramientos, ataques de transpiración, palpitaciones, desórdenes del sueño, cambios de humor, nerviosismo, ansiedad, mala memoria, pérdida de confianza, pérdida de libido, concentración pobre, energía disminuida, empuje disminuido, irritabilidad, atrofia urogenital, atrofia de los senos, enfermedad

cardiovascular, cambios en la distribución del vello, espesor del cabello, cambios en el estado de la piel o para la prevención o control de osteoporosis.

35. Un método de acuerdo a la reivindicación 34, donde las enfermedades y síntomas están seleccionados del grupo que comprende incluyendo acaloramientos, ataques de transpiración, palpitaciones, desórdenes del sueño, cambios de humor, nerviosismo, ansiedad, atrofia urogenital, atrofia de los senos o para la prevención o control de osteoporosis.

36. Un método de acuerdo a la reivindicación 32, donde el estrógeno donde el estrógeno es seleccionado del grupo que consiste de estradiol, sulfamatos de estradiol, valerato de estradiol, benzoato de estradiol, etinil estradiol, estrona, estriol, succinato de estriol y estrógenos conjugados, que incluyen estrógenos equino conjugados tal como sulfato de estrona, sulfato de 17β -estradiol, sulfato de 17α -estradiol, sulfato de equilino, sulfato de 17β -dihidroequilino, sulfato de 17α -dihidroequilino, sulfato de equilenino, sulfato de 17β -dihidroequilenino y sulfato de 17α -dihidroequilenino o mezclas de los mismos.

37. Un método de acuerdo a la reivindicación 36, donde el estrógeno se selecciona del grupo que consiste de estradiol, sulfamatos de estradiol, valerato de estradiol, benzoato de estradiol, estrona, y sulfato de estrona o mezclas de los mismos.

38. Un método de acuerdo a la reivindicación 37, donde el estrógeno es estradiol.

39. Un método de acuerdo a la reivindicación 32, donde la drospirenona (DRSP) está en forma micronizada.

40. Un método de acuerdo a la reivindicación 32, donde el estrógeno está en forma micronizada.

25

69

41. Un método de acuerdo a la reivindicación 38, donde el estradiol está en forma micronizada.
42. Un método de acuerdo a la reivindicación 32, donde la dosis de DRSP corresponde a 15 a 70 mg por ciclo, tal como 20 a 60 mg por ciclo, particularmente 40 a 60 mg por ciclo.
43. Un método de acuerdo a la reivindicación 32, donde la cantidad de DRSP corresponde a una dosis diaria en el rango de desde 0,25 a 10 mg, tal como aproximadamente 0,25 a 8, 0,25 a 6, 0,25 a 5, 0,5 a 4,5, 1 a 4, y 1,5 a 3,5 mg.
44. Un método de acuerdo a la reivindicación 38, donde la cantidad de estradiol corresponde a una dosis diaria en el rango de desde 0,1 a 5 mg, tal como aproximadamente 0,2 a 4,5, 0,5 a 4, 1 a 3, en particular 1, 2 ó 3 mg.
45. Un método para tratar y prevenir enfermedades, desórdenes y síntomas asociados con niveles endógenos deficientes de estrógeno en mujeres que comprende administrar estradiol en cantidades que corresponden a dosis diarias de 1 a 3 mg y drospirenona en cantidades que corresponden a dosis diarias de 1 a 3,5 mg.
46. Un método de acuerdo a la reivindicación 32, que comprende administrar durante 10 a 12 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg; y administrar durante 10 a 12 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg; y

administrar durante 4 a 8 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 5 mg.

47. Un método de acuerdo a la reivindicación 32, que comprende administrar durante 10 a 12 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg; y

administrar durante 10 a 12 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg; y

administrar durante 4 a 8 días una unidad de dosificación diaria que comprende un placebo o blanco.

48. Un método de acuerdo a la reivindicación 32, que comprende administrar durante al menos 21 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg; y

administrar durante no más de 7 días una unidad de dosificación que comprende por placebo o blanco.

49. Un método de acuerdo a la reivindicación 32, que comprende administrar durante al menos 21 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg; y

administrar durante no más de 7 días una unidad de dosificación que comprenda estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg.

50. Un método de acuerdo a la reivindicación 32, que comprende administrar durante 21 a 28 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg.

51. Un método de acuerdo a cualquiera de las reivindicaciones 32 a 45, donde el estrógeno es administrado continuamente.

52. Un método de acuerdo a cualquiera de las reivindicaciones 32 a 45, donde el estrógeno y la drospirenona son administrados continuamente.

53. Un método de acuerdo a cualquiera de las reivindicaciones 32 a 45, donde el estrógeno es administrado continuamente y la drospirenona es administrada secuencialmente.

54. Un método de acuerdo a la reivindicación 53, donde la dosificación de estrógeno es más baja para los 1 a 7 días que siguen inmediatamente a dicha administración secuencial de drospirenona.

55. Un método de acuerdo a la reivindicación 32, donde el estrógeno es administrado continuamente y la drospirenona es administrada en una forma interrumpida.

56. Un método de acuerdo a la reivindicación 55, donde el estrógeno es administrado continuamente durante 21 a 30 días y la drospirenona es administrada en un ciclo de 3 días sí y 3 días no.

78
70

57. Un método de acuerdo a la reivindicación 56, donde la drospirenona es administrada en días 4 a 6, 10 a 12, 16 a 18, 22 a 24, y 28 a 30.

58. Un método de acuerdo a cualquiera de las reivindicaciones 32 a 45, donde el estrógeno y la drospirenona son cada una administrada secuencialmente con un intervalo libre de tratamiento.

59. Un método de acuerdo a la reivindicación 32, que comprende administrar durante 20 a 24 días una unidad de dosis diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg,

y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg durante los últimos 10 a 12 días de dichos 20 a 24 días, y administrar durante 4 a 8 días una unidad de dosificación diaria que no comprende un ingrediente activo.

60. Un método de acuerdo a la reivindicación 32, que comprende administrar durante 20 a 24 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg, y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg durante los últimos 10 a 12 días de dichos 20 a 24 días, y

administrar durante 4 a 8 días una unidad de dosificación diaria que comprende estradiol en cantidades menores que la unidad de dosificación diaria tomada durante dichos 20 a 24 días de administración de estradiol.

61. Un método de acuerdo a la reivindicación 32, que comprende

79
71

administrar durante 20 a 40 días una unidad de dosificación diaria que comprende estradiol en cantidades que corresponden a dosis diarias en el rango de desde 0,1 a 5 mg,

y drospirenona en cantidades que corresponden a dosis diarias en el rango de desde 0,25 a 6 mg durante los últimos 10 a 12 días de dichos 20 a 24 días, y no administrar ninguna unidad de dosificación durante 4 a 8 días.

62. Un método de acuerdo a cualquiera de las reivindicaciones 32 a 61, donde el estrógeno y/o drospirenona son administrados en formulación oral, desde un parche, desde un implante o combinaciones de los mismos.

63. Un método de acuerdo a la reivindicación 62, donde el estrógeno y/o drospirenona son administrados en formulación oral.

64. Un método de acuerdo a cualquiera de las reivindicaciones 46 a 61, donde las unidades de dosificación diarias son administradas durante 1 a 12, preferiblemente 2 a 8, tal como 2, 3, 4, 5, 6, 7, y 8 múltiplos de 28 días.

65. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente removibles dispuestas dentro de una unidad de envase destinadas a administración oral durante un período de al menos 21 días donde dichas unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg.

66. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente removibles dispuestas dentro de una unidad de envase destinadas a administración oral durante un período de 28 días donde dichas unidades de

dosificación diarias comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg.

67. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente removibles dispuestas dentro de una unidad de envase y destinadas a administración oral durante un período de 28 días,

donde al menos 21 de dichas unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y

no más de 7 de dichas unidades de dosificación comprenden un placebo o blanco.

68. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente removibles dispuestas dentro de una unidad de envase y destinadas a administración oral durante un período de 28 días

donde al menos 21 de dichas unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y

no más de 7 de dichas unidades de dosificación comprenden estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg.

69. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente

removibles dispuestas dentro de una unidad de envase y destinadas a administración oral durante un período de al menos 28 días consecutivos, donde al menos 10 de dichas unidades de dosificación diarias comprenden estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg; y al menos 10 de dichas unidades de dosificación diaria comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y no más de 8 de dichas unidades de dosificación comprenden un placebo o blanco.

70. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente removibles dispuestas dentro de una unidad de envase destinadas a administración oral durante un período de al menos 28 días consecutivos, donde al menos 10 de dichas unidades de dosificación diarias comprenden estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg; y al menos 10 de dichas unidades de dosificación diaria comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y no más de 8 de dichas unidades de dosificación comprenden estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg.

71. Una preparación farmacéutica multi-fásica que consiste en un número de unidades de dosificación diarias separadamente envasadas e individualmente

removibles dispuestas dentro de una unidad de envase destinadas a administración oral durante un período de 21 a 30 días consecutivos,

donde 10 a 15 de dichas unidades de dosificación diarias comprenden una combinación de estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg y drospirenona en una cantidad en el rango de desde aproximadamente 0,25 a 6 mg; y

10 a 15 de dichas unidades de dosificación diarias comprenden estradiol en una cantidad en el rango de desde aproximadamente 0,1 a 5 mg.

72. Una preparación de acuerdo con cualquiera de las reivindicaciones 65 a 71, en donde el número de unidades de dosificación diarias es al menos 21 o un múltiplo de 21 tales como 2 a 12, particularmente 2 a 8, tales como 2 a 6.

73. Una preparación de acuerdo con cualquiera de las reivindicaciones 65 a 71, en donde el número de unidades de dosificación diarias es 28 o un múltiplo de 28 tales como 2 a 12, particularmente 2 a 8 tales como 12 a 6.

74. Una preparación de acuerdo con cualquiera de las reivindicaciones 65 a 71, en donde dichas unidades de dosificación diarias comprenden estradiol y/o drospirenona en forma micronizada o rociadas a partir de una solución sobre partículas de un portador inerte.

75. Un uso de un estrógeno (y/o derivado natural o sintético del mismo) en cantidades suficientes para el tratamiento de síntomas menopáusicos y post-menopáusicos y la prevención de osteoporosis en mujeres menopáusicas y post-menopáusicas

y $6\beta,7\beta;15\beta;16\beta$ -dimetilen-3-oxo-17 α -preg-4-eno-21,17-carbo-lactona (drospirenona) en cantidades suficientes para proteger el endometrio de los efectos adversos del estrógeno para la fabricación de un medicamento.

73
75

76. Un uso de un estrógeno (y/o derivado natural o sintético del mismo) en cantidades suficientes para el tratamiento de síntomas menopáusicos y postmenopáusicos y la prevención de osteoporosis en mujeres menopáusicas y postmenopáusicas y 6 β ,7 β ;15 β ;16 β -dimetilen-3-oxo-17 α -preg-4-eno-21,17-carbolactona (drospirenona) en cantidades suficientes para proteger el endometrio de los efectos adversos del estrógeno para la fabricación de un medicamento para terapia de reemplazo de hormonas.



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

4802 8/x
76

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

00200183.2

**Der Präsident des Europäischen Patentamts:
Im Auftrag**

For the President of the European Patent Office

**Le Président de l'Office européen des brevets
p.o.**

I.L.C. HATTEN-HECKMAN

**DEN HAAG, DEN
THE HAGUE, 18/01/01
LA HAYE, LE**



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

85
77

**Blatt 2 der Bescheinigung
Sh et 2 of the certificate
Page 2 de l'attestation**

Anmeldung Nr.:
Application no.:
Demande n°: **00200183.2**

Anmeldetag:
Date of filing:
Date de dépôt: **18/01/00**

Anmelder:
Applicant(s):
Demandeur(s):
**SCHERING AKTIENGESELLSCHAFT
13342 Berlin
GERMANY**

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:
Drospirenone for hormone replacement therapy

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:
A61K31/585, A61P5/30

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

78

1

Drospirenone for Hormone Replacement Therapy**FIELD OF INVENTION**

- 5 The present invention relates to a pharmaceutical composition comprising drospirenone and estrogen, and to methods of hormone replacement therapy by administration of drospirenone and estrogen for estrogen-deficient women.

GENERAL BACKGROUND OF THE INVENTION

10

- The rise in life expectancy and consequent rise in the number of peri- and post-menopausal women has led to an increase in public and medical awareness of the climacteric period of transition in the reproduction phase of women. Menopause, the last menstruation, occurs between the ages of 45 and 55 in most women. Many factors, including race, genetics, nutrition, altitude, smoking, number of live births, the use of hormonal contraception, length of menstrual cycle and the age of onset of puberty have all been attributed, rightly or wrongly, to affect the age of the last menstrual period.

15

- During these phases of life, female endocrine activity undergoes a series of changes, with the result that the physical and psychological well being of many women is adversely affected. Hormone replacement therapy has aimed to improve the quality of life of women during this natural ageing process to alleviate symptoms associated with this time of transition and to reduce the likelihood or slow the progression of disorders and diseases associated with reduced hormonal activity.

25

Drospirenone is known from DE 26 52 761 in which its use as a diuretic is disclosed.

The gestagen-like activity of drospirenone and its consequent utility as a contraceptive agent at dosage levels of 0.5-50 mg is disclosed in DE 30 22 337.

30

The use and role of progestogens in opposed forms of hormone replacement therapy has been studied by the scientific community (Lobo R.A., 1992; Sobel N.B., 1994) as have been regimens comprising estrogens and progestogens (Corson S.L., 1993; Jones K.P., 1992).

35

The use of a preparation for substitution therapy and for oral contraception comprising at least one progestagen and at least one estrogen wherein the estrogen dose varies with a periodicity such that blood loss is substantially avoided is disclosed in PCT/EP94/02897.

5 SUMMARY OF THE INVENTION

The invention relates in a first aspect to a pharmaceutical composition comprising as a first active agent, an estrogen (or naturally or synthetic derivative thereof) in sufficient amounts to treat diseases, disorders and symptoms associated with deficient endogenous
10 levels of estrogen in women, and as a second active agent, $6\beta,7\beta;15\beta;16\beta$ -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone) in sufficient amounts to protect the endometrium from the adverse effects of estrogen, together with pharmaceutically acceptable excipients or carriers.

15 In a second aspect, the invention relates to a pharmaceutical composition comprising as a first active agent estradiol in amounts corresponding to a daily dose of 1 to 3 mg to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women, and as a second active agent $6\beta,7\beta;15\beta;16\beta$ -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone) in amounts corresponding to a daily dose
20 of 1 to 3.5 mg to protect the endometrium from the adverse effects of estrogen, together with a pharmaceutically acceptable excipient or carrier.

Another aspect of the invention relates to the use of a combination of estrogen and drospirenone for the preparation of a medicament wherein the amount of estrogen is
25 sufficient to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in and the amount of drospirenone is sufficient to protect the endometrium from the adverse effects of estrogen.

In a further aspect, the invention relates to a method of treating diseases, disorders and
30 symptoms associated with deficient endogenous levels of estrogen in women comprising administering estrogen in sufficient amounts to alleviate said symptoms and drospirenone in sufficient amounts to protect the endometrium from adverse effects of estrogen.

Furthermore, the invention relates to a method of treating and preventing diseases,
35 disorders and symptoms associated with deficient endogenous levels of estrogen in

79

3

women comprising administering estradiol in amounts corresponding to daily doses of 1 to 3 mg, such as 1, 2 or 3 mg, and drospirenone in amounts corresponding to daily doses of 1 to 3.5 mg, such as 1, 1.5, 2, 2.5, 3, or 3.5 mg.

5 DETAILED DISCLOSURE OF THE INVENTION

In the present context, the term cycle itself or when associated with the term menstrual is intended to mean the number of days between menses in a woman. It can range from 21-31 days, typically 28 days.

10

In the present context, the term menopause is understood as the last natural (ovary - induced) menstruation. It is a single event and a result of an age-dependent dysfunction of the ovarian follicles. Menopause results from the ovaries decreasing their production of the sex hormones estrogen and progesterone. When the number of follicles falls below a certain threshold (a bleeding threshold), the ovaries can no longer produce mature follicles and sex hormones. The ability to reproduce capability ends with menopause.

15

The peri-menopausal phase begins with the onset of climacteric symptoms when the cycle becomes irregular and ends one year after menopause. The end of peri-menopausal phase can be identified after a protracted period of time without bleeding. Post-menopause is the phase that begins at menopause and continues until death.

20

One principal aim of hormone replacement therapy is to restore levels of the sex steroid hormones in naturally or prematurely pre-menopausal, menopausal and post-menopausal women or to establish these levels in hypogonadal females.

25

Monotherapy, also referred to as unopposed therapy, is the treatment with estrogens alone. Exogenous estrogens stimulate the proliferation of the endometrium. In estrogen monotherapy, the opposing effect of progesterone, which terminates proliferation, is absent. The desquamation phase, during which the top layers of the endometrium are shed, does not occur and proliferation of the endometrium occurs to a greater extent than in the phases up to and including the pre-menopausal phase. The result is hyperplasia, a risk factor for endometrial cancer.

30

Combination therapy, also referred to as opposed therapy, is a treatment where a progestagen is added to protect the endometrium from hyperplasia.

The use of natural progesterone in combination therapy is limited by the low bioavailability of natural progesterone, even in micronized form. Significantly, it has been found that combination therapy comprising the use of drospirenone as a progestagen, is remarkably effective. Drospirenone (DRSP), a 17- α -spiro lactone derivative, is a synthetic progestagen that has a surprisingly similar physiological profile to progesterone yet notably better bioavailability. It is the first synthetic progestagen to have a progesterone-like pharmacological profile in that it is antiestrogenic, antiandrogenic, and has antimineralocorticoid activity.

The invention embodies a pharmaceutical composition comprising an estrogen, or naturally or synthetic derivative thereof, in sufficient amounts to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women, and as a second active agent, 6 β ,7 β ;15 β ;16 β -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone) in sufficient amounts to protect the endometrium from the adverse effects of estrogen, together with pharmaceutically acceptable excipients or carriers.

20

Apart from the active substances themselves, it is envisaged that an ester or prodrug of drospirenone may be employed in the present composition, e.g. an oxylminopregnane carbolactone as disclosed in WO 96/24801.

Deficient levels of estrogen can occur for a variety of reasons. The composition can be such that it is adequate for deficient levels of estrogen, regardless of the cause. Causes anticipated by the therapy are, but not limited to, natural menopause, peri-menopause, post-menopause, hypogonadism, castration or primary ovarian failure.

Low levels of estrogen, irrespective of the cause, lead to an overall decreased quality of life for women. Symptoms, diseases and disorders range from merely being inconvenient to life threatening. The composition of this therapy anticipates the effective alleviation of all physiological and psychological signs of estrogen deficiency.

80

5

Transient symptoms, such as vasomotor signs and psychological symptoms are certainly embodied with the realm of therapy. Vasomotor signs comprise but are not limited to hot flushes, sweating attacks such as night sweats, and palpitations. Psychological symptoms of estrogen deficiency comprise, but are not limited to, insomnia and other sleep disorders, poor memory, loss of confidence, mood changes, anxiety, loss of libido, difficulties in concentration, difficulty in making decisions, diminished energy and drive, irritability, and crying spells.

The treatment of the aforementioned symptoms can be associated with the perimenopausal phase of a woman's life or after, sometimes long after menopause. It is anticipated that the invention is applicable to these and other transient symptoms during the perimenopausal phase, menopause, or postmenopausal phase. Moreover, the aforementioned symptoms can be alleviated if the cause of the estrogen deficiency is hypogonadism, castration or primary ovarian failure.

In another embodiment of the invention, the therapy is used for the treatment of permanent effects of estrogen deficiency. Permanent effects comprise physical changes such as urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition and osteoporosis.

Urogenital atrophy, conditions associated with it such as vaginal dryness, increase in vaginal pH and subsequent changes in flora, or events which lead to such atrophy, such as decreases in vascularity, fragmentation of elastic fibres, fusion of collagen fibres, or decreases in cell volume are symptoms thought to be particularly relevant to this therapy.

Furthermore, the invention is thought to be relevant to other urogenital changes associated estrogen deficiency such as decreases in the length and/or diameter of the vagina, decreases mucus production, changes in cell population, decreases in glycogen production, decreases in growth of lactobacilli or increases in growth of streptococci, staphylococci, or coliform bacilli. Other associated changes that are thought to be preventable by the invention are those that may render the vagina susceptible to injury or infection, such as exudative discharges, vaginitis, and dyspareunia. Furthermore, infections of the urinary tract and incontinence are other common symptoms associated with lowered estrogen levels.

Other embodiments of the invention include the prevention or alleviation of physical changes associated with estrogen deficiency, such as changes in the skin, changes in hair distribution, thickness of hair, atrophy of the breasts, or osteoporosis.

- 5 The prevention and management of osteoporosis, most notably post-menopausal osteoporosis, is a particularly interesting embodiment of the invention. Furthermore, bone demineralisation, reduction of bone mass and density, thinning and interruption of trabeculae, and/or consequent increase in bone fractures or bone deformations are thought to be particularly relevant. The prophylactic treatment of osteoporosis is an
10 interesting therapeutic application of the invention.

- A particularly interesting embodiment of the invention comprises the use of the composition for lessening the frequency, persistence, duration and/or severity of hot
flushes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness,
15 anxiety, poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive, irritability, urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition and osteoporosis, most notably hot flushes, sweating attacks, palpitations, sleep
disorders, mood changes, nervousness, anxiety, urogenital atrophy, atrophy of the
20 breasts or for the prevention or management of osteoporosis.

- The pharmaceutical composition for HRT disclosed herein comprises estrogen. In preferred embodiments, the estrogen is selected from the group consisting of estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate, ethinyl estradiol, estrone,
25 estriol, estriol succinate and conjugated estrogens, including conjugated equine estrogens such as estrone sulfate, 17 β -estradiol sulfate, 17 α -estradiol sulfate, equilin sulfate, 17 β -dihydroequilin sulfate, 17 α -dihydroequilin sulfate, equilenin sulfate, 17 β -dihydroequilenin sulfate and 17 α -dihydroequilenin sulfate or mixtures thereof. Particularly interesting estrogens are selected from the group consisting of estradiol, estradiol sulfamates,
30 estradiol valerate, estradiol benzoate, estrone, and estrone sulfate or mixtures thereof, notably estradiol, estradiol valerate, estradiol benzoate and estradiol sulfamates. Most preferred are estradiol or estradiol sulfamates, particularly estradiol.

- Thought to be particularly relevant are micronized forms of estrogens, such as micronized
35 estradiol, micronized estradiol sulfamates, micronized estradiol valerate, micronized

estradiol benzoate, micronized estrone, or micronized estrone sulfate or mixtures thereof, notably micronized estradiol, micronized estradiol valerate, micronized estradiol benzoate or micronized estradiol sulfamates. Most preferred is micronized estradiol or micronized estradiol sulfamates, particularly micronized estradiol.

5

In certain embodiments of the invention, wherein the composition comprises more than one estrogen, one or more estrogen may be in micronized form, such as 2 or all of the estrogen.

- 10 Moreover, an interesting embodiment of the invention comprises a composition wherein the drospirenone (DRSP) is in micronized form, such that one or both estrogen and DRSP are in micronized form, preferably both estrogen and DRSP are in micronized form.

- Drospirenone, which may be prepared substantially as described in, e.g., US 4,129,584 or
15 WO 98/06738, is a sparingly soluble substance in water and aqueous buffers at various pH values. Furthermore, drospirenone is rearranged to an inactive isomer under acid conditions and hydrolysed under alkaline conditions. To ensure good bioavailability of the compound, it is therefore advantageously provided in a form that promotes rapid dissolution thereof.

20

- It has been found that when drospirenone is provided in micronized form in a pharmaceutical composition, rapid dissolution of the active compound from the composition occurs *in vitro*. A micronized substance is such that a test batch (ca. 200 mg) of the particles, herein drospirenone particles, has a surface area of more than 10,000
25 cm²/g, and has the following particle size distribution for drospirenone as determined under the microscope: not more than 2 particles in a given batch (ca. 200 mg) with a diameter of more than 30 µm, and preferably ≤ 20 particles with a diameter of ≥ 10 µm and ≤ 30 µm. The term "rapid dissolution" is defined as the dissolution of at least 70% over about 30 minutes, in particular at least 80% over about 20 minutes, of drospirenone
30 from a tablet preparation containing 3 mg of drospirenone in 900 ml of water at 37°C determined by the USP XXIII Paddle Method using a USP dissolution test apparatus 2 at 50 rpm.

- As an alternative to providing the drospirenone in micronized form, it is possible to
35 dissolve it in a suitable solvent, e.g. methanol or ethyl acetate, and spray it onto the

Tuesday 18 of Jan 2000, PVLP 33838800

->00 31 70 3403018

Page 1

8

surface of inert carrier particles followed by incorporation of the particles containing drospirenone on their surface in the composition.

Without wishing to be limited to any particular theory, it appears that the *in vitro* dissolution rate of drospirenone is connected to the dissolution rate *in vivo* resulting in rapid absorption of drospirenone *in vivo* on oral administration of the compound. This is an advantage because isomerization of the compound in the gastric environment and/or hydrolysis in the intestine is substantially reduced, leading to a high bioavailability of the compound.

10

With respect to the estrogen which may also be a sparingly soluble substance, though usually less sensitive to degradation than drospirenone under conditions prevailing in the gastrointestinal tract, it is also an advantage to provide it in micronized form or sprayed from a solution, e.g. in ethanol, onto the surface of inert carrier particles. This has the added advantage of facilitating a more homogenous distribution of the estrogen throughout the composition. When the estrogen is provided in micronized form, it preferably has the following particle size distribution as determined under the microscope: 100% of the particles have a diameter of $\leq 15.0 \mu\text{m}$, 99% of the particles have a diameter of $\leq 12.5 \mu\text{m}$, 95% of the particles have a diameter of $\leq 10.0 \mu\text{m}$, and 50% of the particles have a diameter of $\leq 3.0 \mu\text{m}$. Furthermore, no particle is larger than $20 \mu\text{m}$, and ≤ 10 particles have a diameter of $\geq 15 \mu\text{m}$ and $\leq 20 \mu\text{m}$.

The particle size distribution for estradiol or estradiol hemihydrate is preferably such that 100% of the particles in a given batch have a diameter less than $15.0 \mu\text{m}$, 99% have a diameter less than $12.5 \mu\text{m}$, 95% have a diameter less than $10.0 \mu\text{m}$, 50% have a diameter less than $3.0 \mu\text{m}$, or 40% have a diameter less than $1.1 \mu\text{m}$.

To obtain a more rapid rate of dissolution, it is preferred to include carriers or excipients, which act to promote dissolution of both active substances. Examples of such carriers and excipients include substances that are readily soluble in water such as cellulose derivatives, carboxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, gelled starch, gelatin or polyvinylpyrrolidone. In particular, it is anticipated that polyvinylpyrrolidone might be particularly helpful to promote dissolution.

82

9

The dose of the drospirenone in each composition is preferably such as to protect the endometrium from the adverse effects of estrogen. DRSP, in sufficient doses, may be used as an opponent to estrogen to protect the endometrium from hyperplasia or cancer.

5 In some cases, however, the dose of DRSP may be sufficient so as to stabilise the menstrual cycle and bleeding pattern. In such cases, doses of estrogen may be low or nil. Adipose tissue production of estrogen may be such that no or very low doses of estrogen are required to stabilise the menstrual cycle and bleeding pattern. Moreover, in certain
10 of an exogenous source of estrogen (such as for the absolute contraindications of severe liver diseases and pregnancy or for the relative contraindications endometrial carcinoma and endometriosis, mammary carcinoma, venous thromboembolism, hypertension, diabetes mellitus, osteoclerosis and melanoma) no or very low amounts of estrogen may be in the composition. In such embodiments, the dose of DRSP may be such that the
15 disorder, disease, or symptom is alleviated.

In preferred embodiments, the dose of DRSP corresponds to 15 to 70 mg per cycle, such as 20 to 60 mg per cycle, particularly 40 to 60 mg per cycle. The length of the cycle, as stated *supra* may vary from 21 to 31 days. Viewed otherwise, a composition may comprise
20 of an amount of DRSP corresponding to a daily dose ranging from 0.25 to 10, such as about 0.25 to 8, 0.25 to 6, 0.25 to 5, 0.5 to 4.5, 1 to 4, and 1.5 to 3.5 mg.

The dose of estrogen may vary from woman to woman, depending on the phase of her life (peri-menopausal or post-menopausal), endogenous levels of estrogen, the severity of the
25 symptom(s), disorder or disease, the disorder, disease or symptom targeted, the use by the woman of other medicaments for other purposes, and other pharmacokinetic variables.

In other embodiments, the dose of estrogen and/or DRSP is sufficient to treat hot flashes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety,
30 poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive, irritability, urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition or for the prevention or management of osteoporosis.

35

Tuesday 18 of Jan 2000, PVLP 33838800

->00 31 70 3403018

Page 1

10

The dose of estrogen and/or DRSP may depend on the treatment, be it for endometrial protection such as bleeding pattern, for osteoporosis prevention or management, for the treatment of menopausal symptoms, such as the for the reduction in the number, frequency, and severity of hot flashes, night sweats, mood swings palpitations, insomnia
5 and other sleep disorders, mood changes, nervousness, anxiety, poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive and irritability.

In embodiments wherein the estrogen is estradiol, the amount of estradiol corresponds to
10 a daily dose ranging from 0.1 to 5 mg, such as about 0.2 to 4.5, 0.5 to 4, 1 to 3, in particular 1, 2, or 3 mg.

With regards to the doses of estradiol derivatives with greater potency, e.g. estradiol valerate, the comparable dosage may be calculated by adjusting the above-mentioned
15 dosages in accordance to the relative potency.

In embodiments where the woman is peri-menopausal the dose of estrogen and/or DRSP may depend on the day of the cycle, that is, whether she is in the preovulatory or postovulatory phase of the cycle, and how advanced within each phase. In embodiments
20 where the woman is post-menopausal or even pre-menopausal, the dose of estrogen and/or DRSP may depend on the time since the last menstruation.

In certain embodiments of the invention, the medicament is administered in the form of a number of separately packaged and individually removable dosage units placed in a
25 packaging unit and intended for oral administration for a period of at least 21, such as at least 28 days, such as at least 30 or 31 days. In such embodiments, the dose of DRSP and/or estrogen may be the same within each dosage unit or may vary. In embodiments where the amount of DRSP and/or estrogen in the dosage unit varies according to the phase or day within the period of at least 21 days, such as at least 28 days that said
30 dosage unit is to be administered, such compositions, methods of treatments and preparations are termed multi-phased.

The dose proportionality may vary according to its use. In preferred embodiments the dose proportionality of estrogen and drospirenone for the preparation of a medicament is
35 such that estrogen doses are to treat diseases, disorders and symptoms associated with

deficient endogenous levels of estrogen and the amount of drospirenone is sufficient to protect the endometrium from the adverse effects of estrogen.

In preferred embodiments, the dose proportionality is sufficient to treat hot flashes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive, irritability, urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition or for the prevention or management of osteoporosis.

10

The invention relates to a method of treating diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women comprising administering estrogen in sufficient amounts to alleviate said symptoms and drospirenone in sufficient amounts to protect the endometrium from adverse effects of estrogen.

15 Preferably, the deficient levels of estrogen to which the method applies are caused by natural menopause, peri-menopause, post-menopause, hypogonadism, castration or primary ovarian failure.

The diseases, disorders and symptoms to which the method applies comprise hot flashes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive, irritability, urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition or for the prevention or management of osteoporosis. Specifically, the method is anticipated to apply to hot flashes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, urogenital atrophy, atrophy of the breasts or for the prevention or management of osteoporosis.

The method applies to the administration of an estrogen, preferably estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate, ethinyl estradiol, estrone, estriol, estriol succinate and conjugated estrogens, including conjugated equine estrogens such as estrone sulfate, 17 β -estradiol sulfate, 17 α -estradiol sulfate, equilin sulfate, 17 β -dihydroequilin sulfate, 17 α -dihydroequilin sulfate, equilenin sulfate, 17 β -dihydroequilenin sulfate and 17 α -dihydroequilenin sulfate or mixtures thereof, most estradiol, estradiol

sulfamates, estradiol valerate, estradiol benzoate, estrone, and estrone sulfate or mixtures thereof, particularly estradiol.

A particularly attractive embodiment of the invention comprises the administration of drospirenone (DRSP) and/or the estrogen in micronized form. Furthermore, of particular interest is where the estrogen is estradiol in micronized form. In such embodiments, one or both active ingredients are administered in micronized form.

In certain embodiments, the method comprises the administration of a dose of DRSP corresponding to 15 to 70 mg per cycle, such as 20 to 60 mg per cycle, particularly 40 to 60 mg per cycle. The method preferably comprises the administration a dose of DRSP the amount of DRSP corresponding to a daily dose ranging from 0.25 to 10 mg, such as about 0.25 to 8, 0.25 to 6, 0.25 to 5, 0.5 to 4.5, 1 to 4, or 1.5 to 3.5 mg.

The amount of estradiol administered may correspond to a daily dose ranging from 0.1 to 5 mg, such as about 0.2 to 4.5, 0.5 to 4, 1 to 3, in particular 1, 2 or 3 mg.

A particularly pertinent embodiment relates to a pharmaceutical composition comprising, as a first active agent, estradiol in amounts corresponding to a daily dose of 1 to 3 mg to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women and as a second active agent $6\beta,7\beta;15\beta;16\beta$ -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone) in amounts corresponding to a daily dose of 1 to 3.5 mg sufficient amounts to protect the endometrium from the adverse effects of estrogen together with a pharmaceutically acceptable excipient or carrier.

25

Certain preferred combinations with regards to the active ingredients in the composition, wherein the estrogen is estradiol comprise 1 mg of estradiol with 0.5 mg of DRSP, 1 mg of estradiol with 1 mg of DRSP, 1 mg of estradiol with 1.5 mg of DRSP, 1 mg of estradiol with 2 mg of DRSP, 1 mg of estradiol with 2.5 mg of DRSP, 1 mg of estradiol with 3 mg of DRSP, 2 mg of estradiol with 1 mg of DRSP and 2 mg of estradiol with 4 mg of DRSP.

In a preferred embodiment, a preferred embodiment of the invention relates to a pharmaceutical composition comprising as a first active agent estradiol in amounts corresponding to a daily dose of 1 to 3 mg, such as 1, 1.5, 2, 2.5, or 3 mg of estradiol, and as a second active agent $6\beta,7\beta;15\beta;16\beta$ -dimethylene-3-oxo-17 α -preg-4-ene-21,17-

84

13

carbolactone (drospirenone) in amounts corresponding to a daily dose of 1 to 3.5 mg, such as 1, 1.5, 2, 2.5, 3, or 3.5 mg of DRSP, together with a pharmaceutically acceptable excipient or carrier.

- 5 Accordingly, a preferred embodiment of the invention relates to a method of treating and preventing diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women comprising administering estradiol in amounts corresponding to daily doses of 1 to 3 mg, such as 1, 2 or 3 mg, and drospirenone in amounts corresponding to daily doses of 1 to 3.5 mg, such as 1, 1.5, 2, 2.5, 3, or 3.5 mg.

10

Given that deficient endogenous levels of estrogens may be associated with, amongst other conditions, natural menopause, peri-menopause, post-menopause, hypogonadism, castration or primary ovarian failure, the method of treating and preventing associated diseases, disorders and symptoms may persist until the death of the individual. That is to say that the composition may be administered from the diagnosis of the disease, disorder or symptoms for the entirety of the individual's life. In certain embodiments, the method and the composition may be based on rhythm of the menstrual cycle. In other embodiments, the method may totally disregard the natural cycle.

15

- 20 The method is preferably multi-phased. The method may comprise administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg; and further administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and further administering for 4 to 8 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.25 to 5 mg.

25

Similarly, the multi-phased method may comprise administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg; and further administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and further administering for 4 to 8 days a daily dosage unit comprising of a placebo or blank.

30

18-01-2000

EP00200183.2

SPEC

Tuesday 18 of Jan 2000, PV&P 33639600

->00 31 70 9403016

Page

14

A regimen of the method may comprise administering for at least 21 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and further administering for no more than 7 days a daily dosage unit comprising of a placebo or blank. A similar regimen of method may comprise administering for at least 21 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and further administering for no more than 7 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg.

Alternatively, a regimen of the method may comprise administering for at least 21 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and not administering a dosage unit for no more than 7 days.

An alternative embodiment of the method comprises administering for 21 to 28 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg.

The regimen may comprise of a continuous administration; that is to say that throughout the 21 to 28 days, a daily dose of estrogen is administered. Likewise, drospirenone may administered continuously, such that either one or both of estrogen and DRSP is/are administered continuously.

In another embodiment, estrogen is administered continuously and drospirenone is administered sequentially. In such an embodiment, throughout the continuous administration of estrogen, DRSP is administered at regular intervals, for 1 to 20 days, such as for 3 to 15 days, 5 to 14 days, particularly for 6 to 14 days. In another interesting embodiment of the regimen of the method, the estrogen dosage is lower for the 1 to 7 days immediately following said sequential administration of drospirenone.

Furthermore, in one embodiment of the invention, the method of treating and preventing diseases, disorders and symptoms associated with deficient endogenous levels of

85

15

estrogen in women comprises continuous administration of estrogen and interrupted administration of progestin. Specifically the method may estrogen being administered continuously for 21 to 30 days and drospirenone being administered in a 3-day-on-3-day-off cycle. A particularly attractive embodiment within this alternative drospirenone being administered on days 4 through 6, 10 through 12, 16 through 18, 22 through 24, and 28 through 30, whilst estrogen, such as estradiol is administered continuously.

The composition of the dosage unit may be formulated in any way conventional in the pharmaceutical art. In particular, as indicated above, the composition may be formulated by a method comprising providing drospirenone and, if desired, an estrogen such as estradiol in micronized form in said unit dosage form, or sprayed from a solution onto particles of an inert carrier in admixture with one or more pharmaceutically acceptable excipients that promote dissolution of the drospirenone and an estrogen so as to promote rapid dissolution of drospirenone and preferably estradiol on oral administration.

Examples of suitable excipients include fillers, e.g. sugars such as lactose, glucose or sucrose, sugar alcohols such as mannitol, starch such as corn or potato starch or modified starch, lubricants such as talc or magnesium stearate and binders such as polyvinylpyrrolidone, cellulose derivatives, carboxymethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, methyl cellulose, or gelatin for making oral dosage forms such as tablets, pills or capsules. Tablets may conveniently be coated with a suitable film-forming agent, e.g. hydroxypropylmethylcellulose. The present composition may also be formulated in liquid form, e.g. as solutions, suspensions or emulsions, together with conventional diluents or excipients in a manner known *per se* in the pharmaceutical art.

25

A particularly interesting regimen of a multi-phased pharmaceutical preparation comprises the sequential administration of both estrogen and drospirenone. An attractive embodiment of such a regimen will comprise a treatment free interval wherein no dosage unit is administered such as comprising administering for 20 to 24 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg, and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg for the last 10 to 12 days of said 20 to 24 day, and not administering any dosage units for 4 to 8 days.

16

Alternatively, any interval wherein neither DRSP nor estrogen is administered, a placebo or blank is administered. Such a regimen of hormone replacement therapy can effectively comprise a method comprising administering for 20 to 24 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg, and further administering drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg for the last 10 to 12 days of said 20 to 24 day, and further administering for 4 to 8 days a daily dosage unit comprising no active ingredient.

An alternative regimen comprises administering for 20 to 24 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg, and further administering drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg for the last 10 to 12 days of said 20 to 24 day, followed by administering for 4 to 8 days a daily dosage of unit comprising estradiol in amounts less than daily dosage unit taken for said 20 to 24 day administration of estradiol.

15

In embodiments where the medicament is in the form of a number of separately packaged and individually removable dosage units placed in a packaging unit and intended for oral administration for a period of at least 21, such as at least 28 days and wherein the estrogen is estradiol, preferably at least 21 daily dosage units comprise a combination of estradiol in an amount from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and 7 or less daily dosage units comprise estradiol in an amount from about 0.1 to 5 mg.

Alternatively, the medicament may be in the form of a number of separately packaged and individually removable dosage units placed in a packaging unit and intended for oral administration for a period of at least 28 days and at least 21 daily dosage units comprise a combination of estradiol in an amount from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg and 7 or less daily dosage units comprise a blank or placebo.

30

It easily follows that the medicament may be in the form of a number of separately packaged and individually removable dosage units placed in a packaging unit and intended for oral administration for a period of at least 28 days and at least 21 daily dosage units comprise a combination of estradiol in an amount from about 0.1 to 5 mg

86

17

and drospirenone in an amount ranging from about 0.25 to 6 mg. There may not be any dosage units for the last 7 or less days of the regimen.

Patient compliance of the many embodiments of the regimen may be aided by means of a pharmaceutical preparation tailored to the patients needs or habits. One such preparation may consist of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of at least 21 days, such as at least 28 days, wherein said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg.

Furthermore, patient compliance may be aided by a regimen comprising a multi-phase pharmaceutical preparation.

Every regimen may be easily be adhered to by a multi-phased pharmaceutical preparation such as one consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of 28 days wherein said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg. In such a preparation, the amount of active ingredient varies through the 28-day period.

An attractive embodiment relates to multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of 21 to 30 consecutive days wherein 10 to 15 said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and 10 to 15 said daily dosage units comprise estradiol in an amount ranging from about 0.1 to 5 mg. This embodiment is particularly suited for preparations wherein estrogen is administered continuously for 21 to 30 days and drospirenone is administered in a 3-day-on-3-day-off cycle. Preferably, within this embodiment, the preparation is designed so that drospirenone is administered on days 4 through 6, 10 through 12, 16 through 18, 22 through 24, and days 28 through 30.

18

Moreover, a multi-phased pharmaceutical preparation wherein the number of daily dosage units is 21 or 28, or a multiple of 21 or 28, such as 2 to 24, such as 2 to 12, particularly 2 to 8, such as multiples of 2 to 6.

- 5 Likewise, the invention relates to a method of treating diseases, disorders, and symptoms associated with estrogen deficiency comprising a multi-phased pharmaceutical preparation comprising daily dosage units which are administered for 1 to 12, preferably 2 to 8, such as 2, 3, 4 5, 6, 7, and 8 multiples of 28 days.
- 10 A packaging unit comprising the daily dosage units described above may be prepared in a manner analogous to that of making oral contraceptives or hormone replacement regimens. This may for instance be a conventional blister pack or any other form known for this purpose, for instance a pack comprising the appropriate number of dosage units (In this case at least 28, or for particular applications, a multiple of 28) in a sealed blister
- 15 pack with a cardboard, paperboard, foil or plastic backing and enclosed in a suitable cover. Each blister container may conveniently be numbered or otherwise marked.

It is also envisaged that the present composition may be in the form of a parenteral formulation such as a subcutaneous implant or transdermal formulation. For making

20 implants, the active agents may suitably be formulated together with one or more polymers that are gradually eroded or degraded when in use, e.g. silicone polymers, ethylene vinylacetate, polyethylene or polypropylene.

Where transdermal formulations are concerned, they may be prepared in the form of

25 matrices or membranes or as fluid or viscous formulations in oil or hydrogels. For transdermal patches, an adhesive which is compatible with the skin should be included, such as polyacrylate, a silicone adhesive or polyisobutylene, as well as a foil made of, e.g. polyethylene, polypropylene, ethylene vinylacetate, polyvinylchloride, polyvinylidene chloride or polyester, and a removable protective foil made from, e.g., polyester or paper

30 coated with silicone or a fluoropolymer. For the preparation of transdermal solutions or gels, water or organic solvents or mixtures thereof may be used. Transdermal gels may furthermore contain one or more suitable gelling agents or thickeners such as silicone, tragacanth, starch or starch derivatives, cellulose or cellulose derivatives or polyacrylic acids or derivatives thereof. Transdermal formulations may also suitably contain one or

35 more substances that enhance absorption through the skin, such as bile salts or

19

derivatives thereof and/or phospholipids. Suitable transdermal formulations may, for instance, be made in a manner analogous to that described in WO 84/04157 for 3-ketodesogestrel. Alternatively, transdermal formulations may be prepared according to a method disclosed in, e.g., BW Barry, "Dermatological Formulations, Percutaneous Absorption", Marcel Dekker Inc., New York - Basel, 1983, or YW Chien, "Transdermal Controlled Systemic Medications", Marcel Dekker Inc., New York - Basel, 1987.

The present invention is further described in the following examples which are not in any way intended to limit the scope of the invention as claimed.

10

18-01-2000

EP00200183.2

SPEC

Tuesday 18 f Jan 2000, PV&P 33839800

->00 31 70 3403018

Page :

20

EXPERIMENTAL

Example 1

5 *Preparation of tablets containing drospirenone and estradiol may be performed in the following manner*

Tablet cores of the following composition

10	micronized drospirenone	3.00 mg
	micronized estradiol	1.00, 2.00, 3.00 mg
	lactose monohydrate	45.2, 46.2, 47.2 mg
	corn starch	14.40 mg
	modified starch	9.60 mg
15	polyvinylpyrrolidone 25,000	4.00 mg
	magnesium stearate	0.80 mg

are prepared by charging a fluidised bed granulator with 31.68 kg corn starch, 21.12 kg modified starch, 6.60 kg micronized drospirenone, 2.20, 4.40 or 6.6 kg of micronized
 20 estradiol (for 1 mg, 2 mg, and 3 mg dose, respectively) and 99.44, 101.64, or 103.84 kg of lactose monohydrate (for 3 mg, 2 mg, and 1 mg dose, respectively) and activating the fluidised bed. An aqueous solution of 8.80 kg polyvinylpyrrolidone 25,000 in 46.20 kg purified water is sprayed continuously onto the fluidised bed while drying by heating the air stream of the fluidised bed. At the end of the process 1.76 kg magnesium stearate is
 25 sucked into the granulator and mixed with the granules by maintaining the fluidised bed. The resulting granulate is pressed into tablet cores by compression using a rotary tablet press.

For tablets comprising 1 mg of estradiol, 2.22464 kg of hydroxypropylmethylcellulose and
 30 0.44528 macrogol 6000 are dissolved in 14.67 kg purified water. 0.44528 kg talc, 1.25906 kg titanium dioxide and 0.02575 kg ferric oxide pigment are suspended in 10.26 kg purified water with stirring and homogenised. The solution and suspension are combined and used to coat the tablet cores by continuous application of the coating suspension in a coater. For tablets comprising 2 or 3 mg of estradiol, the specific weights of the reagents
 35 for the preparation of the coating can easily be calculated.

An alternative formulation for the coatings of 1 mg tablets comprises 2.22484 kg of hydroxypropylmethylcellulose, 0.44528 kg of macrogol 600, 0.44528 kg of talc, 1.17326 kg of titanium dioxide, 0.07834 kg of ferric oxide pigment, yellow, and 0.03520 kg of ferric oxide pigment, red. A possible formulation for the coatings of 2 mg tablets comprises of 2.22484 kg of hydroxypropylmethylcellulose, 0.44528 kg of macrogol 600, 0.44528 kg of talc, 1.19636 kg of titanium dioxide and 0.08844 kg of ferric oxide pigment, red. A possible formulation for the coatings of 3 mg tablets comprises of 2.22484 kg of hydroxypropylmethylcellulose, 0.44528 kg of macrogol 600, 0.44528 kg of talc, 1.25906 kg of titanium dioxide and 0.02574 kg of ferric oxide pigment, red.

Example 2: Relative bioavailability

The evaluation of relative bioavailability of estradiol (E2) and drospirenone (DRSP), in an open randomised 2-way cross over study with volunteers. E2/DRSP after treatment with either 2 mg of E2 + 2 mg of DRSP, 2 mg of E2 + 6 mg of DRSP coated tablets p.o versus 2 mg of E2 + 2 mg of DRSP oral solution.

Example 3: Repeated Dose

The evaluation of repeated dose pharmacokinetics (accumulation) and potential interaction between estradiol and drospirenone was performed. This open-label, randomised, intra-individual cross-over study of two dose level combinations with an intervening wash-out phase (4 weeks), and multiple application over 28 days was done with 4 dose combinations. A 4-week observation period was performed after the last dose.

Treatment A: 1 mg E2 + 1 mg DRSP, daily, p.o.

Treatment B: 1 mg E2 + 4 mg DRSP, daily, p.o.

Treatment C: 2 mg E2 + 1 mg DRSP, daily, p.o.

30 Treatment D: 2 mg E2 + 4 mg DRSP, daily, p.o.

Assessment: No statistically significant interaction was observed between the two active ingredients.

18-01-2000

EP00200183.2

SPEC

Tuesday 18 of Jan 2000, PV&P 33839800

->00 31 70 3403018

Page 2

22

Example 4: Endometrial Protection

Primary Objective: To evaluate the efficacy of thirteen, 28 day cycles of continuous E2-DRSP combinations compared to continuous E2 by analysis of protection against
5 hyperplasia in post-menopausal women.

Secondary Objectives: To evaluate the effect of endometrial morphology, bleeding patterns, metabolic and hemostatic laboratory parameters. Well-being of post-menopausal women as assessed by the Women's Health Questionnaire (WHQ) and The
10 Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36). Effect on the frequency and severity of hot flashes, and on the relief of urogenital symptoms. Drug levels of DRSP and E2. Detailed evaluation of the metabolic parameters in subgroup.

Synopsis:

15 **Dose Groups:** E2 1 mg; E2 1 mg + DRSP 0.5 mg; E2 1 mg + DRSP 1 mg; E2 1 mg + DRSP 2 mg; E2 1 mg + DRSP 3 mg.

Post-menopausal women with or without menopausal symptoms will be assigned 1 of 5 regimens. Endometrial effects will be evaluated by biopsy to determine the incidence of
20 endometrial hyperplasia. Symptoms and bleeding patterns will be evaluated from subject diaries. General safety, and effects on specific biochemical and hematological parameters will be evaluated. A post-menopausal quality of life assessment and evaluation of patient satisfaction will also be made.

25 Two-hour glucose tolerance tests, and 15-minute insulin tolerance tests were performed at certain sites.

Data Analysis

Endometrial biopsy at Visit 1 and Final Visit. Bleeding pattern information is recorded in a
30 daily diary throughout the study.

Two analyses will be performed for the primary efficacy variable: i) 2-sided comparison of treatments and ii) 1-sided, within group interval estimation of dose effect.

Frequency tables will be generated for visits at which endometrial biopsies are performed.
35 These tables will display the number and percent of subjects in each endometrial

89

23

response category for each treatment group (and by centre in case of a treatment-by-centre interaction). Overall and between group comparison of the incidence of hyperplasia among the treatment groups will be performed. This analysis will be intent-to-treat and will test the null hypothesis of no difference in response to treatments

5 (unopposed vs. opposed estradiol) when adjusted for centre.

Interval estimation of dose response

The probability π_i will be estimated for each dose of DRSP (t 0.0, 0.5, 1.0, 2.0, 3.0). Additionally, the upper limit of the one-sided 95% confidence interval will be calculated for

10 each π_i separately. This means that the confidence intervals will be non-simultaneous.

Example 5: Osteoporosis Prevention

15 Primary Objective: Bone mineral density of the hip after 104 weeks of treatment.

Secondary Objectives: Bone mineral density (BMD) of the hip after 12, 28, 52, and 80 weeks of treatment. BMD of the lumbar spine, of the midshaft of the radius, of the total body. Effects on parameters for bone metabolism. Bleeding pattern. General safety.

20 The study will be performed as a double blind, placebo-controlled trial with 240 healthy, post-menopausal women randomly assigned to one of 4 groups of 60 after giving their informed consent.

The groups will be as follows:

25 1 mg E2 + 1 mg DRSP

 1 mg E2 + 2 mg DRSP

 1 mg E2 + 3 mg DRSP

Synopsis

Forty osteopenic patients (BMD hip T-score between -1 and -2.5) and 20 non-osteopenic patients should be enrolled in each group. All treatments will be applied daily *per os*

30 during the whole treatment of 2 years without a treatment-free interval. In addition, the patients will be supplied with calcium tablets (500 mg of calcium daily). Measurements of bone mineral density of the hip will be measured on the left side utilizing dual-energy x-ray absorptiometry at screening, baseline, and after 12, 28, 52, 80, and 104 weeks of treatment. Biochemical markers of bone remodelling will be measured at intervals.

35

18-01-2000

EP00200183.2

SPEC

Tuesday 18 of Jan 2000, PV&P 33639800

->00 31 70 3403016

Page 3

24

Serum bone specific alkaline phosphatase, serum N-mid osteocalcin, urinary calcium/creatinine (second morning void), and urinary CrossLaps® /creatinine (second morning void) will additionally be evaluated.

5

Example 8: Menopausal Symptoms

Objective: To demonstrate that the therapeutic efficacy of E2-DRSP regarding menopausal symptoms is superior to placebo

10

Primary Objective: Hot flushes

Secondary Objectives: Sweating episodes, sleep problems, depressed moods, nervousness, urogenital symptoms (vaginal dryness, pollakiuria, nocturia), bleeding patterns. General safety.

15

Synopsis

Double-blind, placebo controlled trial with healthy, post-menopausal women randomly assigned to one of 4 groups

20

1 mg E2 + 1 mg DRSP

1 mg E2 + 2 mg DRSP

1 mg E2 + 3 mg DRSP

placebo

25 Synopsis:

Main inclusion criterion: A minimum of 5 moderate to severe hot flushes per day on at least 7 days of the 2-week pre-treatment period.

The duration of the treatment will be 16 weeks (4 28-day cycles).

30

Hot flushes will be assessed by means of recording the frequency and severity (mild, moderate and severe) and comparing these between the verum groups and placebo. The patient will record the incidence and severity of the hot flushes and diary cards. In addition, the investigator will ask the patient at each visit about other typical menopausal symptoms (sweating episodes, sleep problems, depressed moods, nervousness,

35

Printed:22-01-2001

RVELSE/24300EP1ALI/PV&P/18-01-00

Last printed 18/01/00 11:23

90

25

urogenital symptoms. The intensity of the symptoms will be assessed as mild, moderate or severe.

If the patient has an intact uterus, the occurrence of bleeding will be recorded every day during the study on diary cards. The patient will make daily entries in the diary card according to the bleeding intensity definitions given below

Code	Category	Definition
0	no	no vaginal bleeding
1	spotting	less than associated with normal menstruation relative to the subject's experience with no need for sanitary protection (except for panty liners)
2	light	less than associated with normal menstruation relative to the subject's experience with no need sanitary protection
3	normal	like normal menstruation relative to the subject's experience.
4	heavy	more than normal menstruation relative to the subject's experience

For urogenital symptoms, the following parameter and categories will be used:

10

Parameter	Categories
vaginal dryness	yes/no/ not applicable
increased frequency of urination	no /yes
nocturia	no/yes ("yes" means at least 2 urinations during 2 or more nights within the preceding week)

Data Analysis

For the efficacy parameters, both a valid-case analysis (VCA) and an intention to treat (ITT) analysis will be performed. The primary target variable is the individual relative change (C) in the number of hot flushes. C is defined as $(T-B) / B$, where T and B are individual means. T is the mean number of hot flushes per week, calculated by the observations during the weeks 3 through 16 of the treatment phase. B is the mean number of hot flushes per week, calculated from the observations of the 2-week pre-treatment phase.

20

35 Tuesday 18 of Jan 2000, PV&P 33838800

->00 31 70 3403018

Page 3

26

Example 7: Lipid Profile

Objective: The primary goal of this study is to compare two E2/DRSP treatments and Premique®/Premelle® regarding lipid profile. Lipid profile is generally accepted as the
5 surrogate endpoint to estimate cardiovascular risk.

Primary Objectives: Lipid pattern including HDL cholesterol, LDL cholesterol

Secondary Objectives: Lipid pattern including total cholesterol, triglycerides, HDL2
10 cholesterol, HDL3 cholesterol, VLDL cholesterol, apolipoproteins (A-1, A-2, B, E), Lp(a);
Menopausal symptoms; Bleeding pattern; Endometrial safety; General safety

Design: The study is performed as an open-label, multicentre, comparative study in 300
post-menopausal women randomly assigned to one of 3 groups of 100 after giving their
15 informed consent.

Groups:

1 mg E2 + 2 mg DRSP

1 mg E2 + 3 mg DRSP

20 Premique®/Premelle®: 0.625 mg conjugated estrogens + 5 mg MPA

All treatments are applied daily orally during the whole treatment period of 2 years without a treatment free interval. Measurements of lipid profile are performed at screening and after 12, 28, 52, and 104 weeks of treatment as well as six weeks after treatment.

25

91

27

CLAIMS

1: A pharmaceutical composition comprising

5 as a first active agent, an estrogen (or naturally or synthetic derivative thereof) in sufficient amounts to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women, and

as a second active agent, 6 β ,7 β ;15 β ;16 β -dimethylene-3-oxo-17 α -preg-4-ene-21,17-
10 carbolactone (drospirenone) in sufficient amounts to protect the endometrium from the adverse effects of estrogen,

together with a pharmaceutically acceptable excipient or carrier.

15 2. A composition according to claim 1, wherein the deficient levels of estrogen are caused by natural menopause, peri-menopause, post-menopause, hypogonadism, castration or primary ovarian failure.

3. A composition according to any of claims 1 or 2, wherein the diseases, disorders and
20 symptoms are selected from the group comprising including hot flushes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive, irritability, urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition and osteoporosis.

25 4. A composition according to claim 3, wherein the diseases, disorders and symptoms are selected from the group comprising including hot flushes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, urogenital atrophy, atrophy of the breasts and osteoporosis.

30 5. A composition according to claim 1, wherein the estrogen is selected from the group consisting of estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate, ethinyl estradiol, estrone, estriol, estriol succinate and conjugated estrogens, including conjugated equine estrogens such as estrone sulfate, 17 β -estradiol sulfate, 17 α -estradiol

18-01-2000

EP00200183.2

SPEC

Tuesday 18 of Jan 2000, PV&P 33639800

->00 31 70 3403018

Page 3

28

sulfate, equilin sulfate, 17 β -dihydroequilin sulfate, 17 α -dihydroequilin sulfate, equilenin sulfate, 17 β -dihydroequilenin sulfate and 17 α -dihydroequilenin sulfate or mixtures thereof.

6. A composition according to claim 5, wherein the estrogen is selected from the group consisting of estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate, estrone, and estrone sulfate or mixtures thereof.

7. A composition according to claim 6, wherein the estrogen is estradiol.

8. A composition according to claim 1, wherein drospirenone (DRSP) is in micronized form.

9. A composition according to any of claims 5 or 6, wherein at least one estrogen is in micronized form.

15

10. A composition according to claim 1, wherein the dose of DRSP corresponds to 15 to 70 mg per cycle, such as 20 to 60 mg per cycle, particularly 40 to 60 mg per cycle.

11. A composition according to claim 1, wherein the amount of DRSP corresponds to a daily dose ranging from 0.25 to 10 mg, such as about 0.25 to 8, 0.25 to 6, 0.25 to 5, 0.5 to 4.5, 1 to 4, and 1.5 to 3.5 mg.

12. A composition according to claim 7, wherein the amount of estradiol corresponds to a daily dose ranging from 0.1 to 5 mg, such as about 0.2 to 4.5, 0.5 to 4, 1 to 3, in particular 1, 2, or 3 mg.

13. A pharmaceutical composition comprising as a first active agent estradiol in amounts corresponding to a daily dose of 1 to 3 mg to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women, and as a second active agent 6 β ,7 β ;15 β ;16 β -dimethylans-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone) in amounts corresponding to a daily dose of 1 to 3.5 mg to protect the endometrium from the adverse effects of estrogen, together with a pharmaceutically acceptable excipient or carrier.

35

92

29

14. Use of a combination of estrogen and drospirenone for the preparation of a medicament wherein the amount of estrogen is sufficient to treat diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women and the amount of drospirenone is sufficient to protect the endometrium from the adverse effects
5 of estrogen.

15. The use according to claim 14, wherein the deficient levels of estrogen are caused by natural menopause, peri-menopause, post-menopause, hypogonadism, castration or primary ovarian failure.

10

16. The use according to claim 14, wherein the amount of estrogen is sufficient for the treatment of hot flashes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, poor memory, loss of confidence, loss of libido, poor concentration, diminished energy, diminished drive, irritability, urogenital atrophy, atrophy of the breasts,
15 cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition or for the prevention or management of osteoporosis.

17. The use according to claims 16, wherein the amount of estrogen is sufficient for the treatment of hot flashes, sweating attacks, palpitations, sleep disorders, mood changes,
20 nervousness, anxiety, urogenital atrophy, atrophy of the breasts or for the prevention or management of osteoporosis.

18. The use according to claim 14, wherein the estrogen is selected from the group consisting of estrogen is selected from the group consisting of estradiol, estradiol
25 sulfamates, estradiol valerate, estradiol benzoate, ethinyl estradiol, estrone, estriol, estriol succinate and conjugated estrogens, including conjugated equine estrogens such as estrone sulfate, 17 β -estradiol sulfate, 17 α -estradiol sulfate, equilin sulfate, 17 β -dihydroequilin sulfate, 17 α -dihydroequilin sulfate, equilenin sulfate, 17 β -dihydroequilenin sulfate and 17 α -dihydroequilenin sulfate or mixtures thereof.

30

19. The use according to claim 18, wherein the estrogen is selected from the group consisting of estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate, estrone, and estrone sulfate or mixtures thereof.

35 20. The use according to claim 19, wherein the estrogen is estradiol.

37 Tuesday 18 of Jan 2000, PV&P 33839800

->00 31 70 3403018

Page 3

30

21. The use according to claim 14, wherein drospirenone (DRSP) is in micronized form.

22. The use according to claim 14, wherein the estrogen is in micronized form.

5

23. The use according to claim 20, wherein the estradiol is in micronized form.

24. The use according to claim 14, wherein the dose of DRSP corresponds to 15 to 70 mg per cycle, such as 20 to 60 mg per cycle, particularly 40 to 60 mg per cycle.

10

25. The use according to claim 14, wherein the amount of DRSP corresponds to a daily dose ranging from 0.25 to 10 mg, such as about 0.25 to 8, 0.25 to 6, 0.25 to 5, 0.5 to 4.5, 1 to 4, and 1.5 to 3.5 mg.

15 26. The use according to claim 20, wherein the amount of estradiol corresponds to a daily dose ranging from 0.1 to 5 mg, such as about 0.2 to 4.5, 0.5 to 4, 1 to 3, in particular 1, 2, or 3 mg.

27. The use according to claim 14, wherein the medicament is in the form of a number
20 of separately packaged and individually removable dosage units placed in a packaging unit and intended for oral administration for a period of at least 28 days;
wherein at least 21 daily dosage units comprise a combination of estradiol in an amount from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg;
and 7 or less daily dosage units comprise estradiol in an amount from about 0.1 to 5 mg.

25

28. The use according to claim 14, wherein the medicament is in the form of a number of
separately packaged and individually removable dosage units placed in a packaging unit
and intended for oral administration for a period of at least 28 days;
wherein at least 21 daily dosage units comprise a combination of estradiol in an amount
30 from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg;
and 7 or less daily dosage units comprise a blank or placebo.

29. The use according to 14, wherein the medicament is in the form of a number of
separately packaged and individually removable dosage units placed in a packaging unit
35 and intended for oral administration for a period of at least 28 days,

1-2000

EP00200183.2

SPEC

Monday 18 of Jan 2000, PV&P 33838800

->00 31 70 3403018

Page 38 of

93

31

wherein at least 10 daily dosage units comprise estradiol in an amount from about 0.1 to 5 mg;

and at least 10 daily dosage unit comprise a combination of estradiol in an amount from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg,

5 and 8 or less daily dosage units comprise a blank or placebo.

30. The use according to 14, wherein the medicament is in the form of a number of separately packaged and individually removable dosage units placed in a packaging unit and intended for oral administration for a period of at least 28 days;

10 wherein at least 10 daily dosage units comprise estradiol in an amount from about 0.1 to 5 mg,

and at least 10 daily dosage units comprise a combination of estradiol in an amount from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg;

and 8 or less daily dosage units comprise estradiol in an amount from about 0.1 to 5 mg.

15

31. The use according to claim 14, wherein the medicament is in the form of a number of separately packaged and individually removable dosage units placed in a packaging unit and intended for oral administration for a period of at least 28 days;

wherein 28 daily dosage units comprising a combination of estradiol in an amount from

20 about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg.

32. A method of treating and preventing diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women comprising administering estrogen in sufficient amounts to alleviate said diseases, disorders and symptoms and

25 drospirenone in sufficient amounts to protect the endometrium from adverse effects of estrogen.

33. A method according to claim 32, wherein the deficient levels of estrogen are caused by natural menopause, peri-menopause, post-menopause, hypogonadism, castration or

30 primary ovarian failure.

34. A method according to claim 32, wherein the diseases, disorders and symptoms are selected from the group comprising including hot flashes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, poor memory, loss of confidence,

35 loss of libido, poor concentration, diminished energy, diminished drive, irritability.

32

urogenital atrophy, atrophy of the breasts, cardiovascular disease, changes in hair distribution, thickness of hair, changes in skin condition or for the prevention or management of osteoporosis.

5 35. A method according to claim 34, wherein the diseases, disorders and symptoms are selected from the group comprising including hot flushes, sweating attacks, palpitations, sleep disorders, mood changes, nervousness, anxiety, urogenital atrophy, atrophy of the breasts or for the prevention or management of osteoporosis.

10 36. A method according to claim 32, wherein the estrogen is selected from the group consisting of estrogen is selected from the group consisting of estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate, ethinyl estradiol, estrone, estriol, estriol succinate and conjugated estrogens, including conjugated equine estrogens such as estrone sulfate, 17 β -estradiol sulfate, 17 α -estradiol sulfate, equilin sulfate, 17 β -
15 dihydroequilin sulfate, 17 α -dihydroequilin sulfate, equilenin sulfate, 17 β -dihydroequilenin sulfate and 17 α -dihydroequilenin sulfate or mixtures thereof.

37. A method according to claim 36, wherein the estrogen is selected from the group consisting of estradiol, estradiol sulfamates, estradiol valerate, estradiol benzoate,
20 estrone, and estrone sulfate or mixtures thereof.

38. A method according to claim 37, wherein the estrogen is estradiol.

39. A method according to claim 32, wherein drospirenone (DRSP) is in micronized form.
25

40. A method according to claim 32, wherein the estrogen is in micronized form.

41. A method according to claim 38, wherein the estradiol is in micronized form.

30 42. A method according to claim 32, wherein the dose of DRSP corresponds to 15 to 70 mg per cycle, such as 20 to 60 mg per cycle, particularly 40 to 60 mg per cycle.

43. A method according to claim 32, wherein the amount of DRSP corresponds to a daily dose ranging from 0.25 to 10 mg, such as about 0.25 to 8, 0.25 to 6, 0.25 to 5, 0.5 to 4.5,
35 1 to 4, and 1.5 to 3.5 mg.

94

33

44. A method according to claim 38, wherein the amount of estradiol corresponds to a daily dose ranging from 0.1 to 5 mg, such as about 0.2 to 4.5, 0.5 to 4, 1 to 3, in particular 1, 2 or 3 mg.

5

45. A method of treating and preventing diseases, disorders and symptoms associated with deficient endogenous levels of estrogen in women comprising administering estradiol in amounts corresponding to daily doses of 1 to 3 mg and drospirenone in amounts corresponding to daily doses of 1 to 3.5 mg.

10

46. A method according claim 32, comprising administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg; and administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and administering for 4 to 8 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.25 to 5 mg.

15

47. A method according to claim 32, comprising administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg; and administering for 10 to 12 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and administering for 4 to 8 days a daily dosage unit comprising of a placebo or blank.

20

48. A method according to claim 32, comprising administering for at least 21 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and administering for no more than 7 days a daily dosage unit comprising of a placebo or blank.

30

49. A method according to claim 32, comprising

35

34

administering for at least 21 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg; and administering for no more than 7 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg.

50. A method according to claim 32, comprising administering for 21 to 28 days a daily dosage unit comprising estradiol in amounts corresponding to daily doses ranging from 0.1 to 5 mg and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg.

51. A method according to any of claims 32 to 45, wherein the estrogen is administered continuously.

52. A method according to any of claims 32 to 45, wherein the estrogen and drospirenone are administered continuously.

53. A method according to any of claims 32 to 45, wherein the estrogen is administered continuously and drospirenone is administered sequentially.

54. A method according to claim 53, wherein the estrogen dosage is lower for the 1 to 7 days immediately following said sequential administration of drospirenone.

55. A method according to claim 32, wherein estrogen is administered continuously and drospirenone is administered in an interrupted manner.

56. A method according to claim 55, wherein estrogen is administered continuously for 21 to 30 days and drospirenone is administered in a 3-day-on-3-day-off cycle.

57. A method according to claim 56, wherein drospirenone is administered on days 4 through 6, 10 through 12, 16 through 18, 22 through 24, and 28 through 30.

58. A method according to any of claims 32 to 45, wherein the estrogen and the drospirenone are each administered sequentially with a treatment-free interval.

35

95

35

59. A method according to claim 32, comprising
administering for 20 to 24 days a daily dosage unit comprising estradiol in amounts
corresponding to daily doses ranging from 0.1 to 5 mg,
and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg for
5 the last 10 to 12 days of said 20 to 24 days, and
administering for 4 to 8 days a daily dosage unit comprising no active ingredient.

60. A method according to claim 32, comprising
administering for 20 to 24 days a daily dosage unit comprising estradiol in amounts
10 corresponding to daily doses ranging from 0.1 to 5 mg, and drospirenone in amounts
corresponding to daily doses ranging from 0.25 to 6 mg for the last 10 to 12 days of said
20 to 24 day, and
administering for 4 to 8 days a daily dosage of unit comprising estradiol in amounts less
than daily dosage unit taken for said 20 to 24 day administration of estradiol.

15

61. A method according to claim 32, comprising
administering for 20 to 24 days a daily dosage unit comprising estradiol in amounts
corresponding to daily doses ranging from 0.1 to 5 mg,
and drospirenone in amounts corresponding to daily doses ranging from 0.25 to 6 mg for
20 the last 10 to 12 days of said 20 to 24 day, and
not administering any dosage units for 4 to 8 days.

62. A method according to any of claims 32 to 61, wherein the estrogen and/or
drospirenone are administered in oral formulation, from a patch, from an implant or
25 combinations thereof.

63. A method according to claim 62, wherein the estrogen and/or drospirenone are
administered in oral formulation.

30 64. A method according to any of claims 46 to 61, wherein the daily dosage units are
administered for 1 to 12, preferably 2 to 8, such as 2, 3, 4 5, 6, 7, and 8 multiples of 28
days.

65. A multi-phased pharmaceutical preparation consisting of a number of separately
35 packaged and individually removable daily dosage units placed into a packaging unit and

36

intended for oral administration for a period of at least 21 days wherein said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg.

- 5 66. A multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of 28 days wherein said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg.

10

67. A multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of 28 days, wherein at least 21 said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and
15 no more than 7 said dosage units comprise a placebo or a blank.

68. A multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of 28 days, wherein at least 21 said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and
20 no more than 7 said dosage units comprise estradiol in an amount ranging from about 0.1 to 5 mg.

69. A multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of at least 28 consecutive days, wherein at least 10 said daily dosage units comprise estradiol in an amount ranging from about 0.1 to 5 mg; and
30 at least 10 said daily dosage units comprises a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and
35 6 mg; and

37

no more than 8 of said daily dosage units comprise a placebo or blank.

70. A multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of at least 28 consecutive days, wherein at least 10 said daily dosage units comprise estradiol in an amount ranging from about 0.1 to 5 mg; and at least 10 said daily dosage units comprises a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and no more than 8 of said daily dosage units comprise estradiol in an amount ranging from about 0.1 to 5 mg.

71. A multi-phased pharmaceutical preparation consisting of a number of separately packaged and individually removable daily dosage units placed into a packaging unit and intended for oral administration for a period of 21 to 30 consecutive days, wherein 10 to 15 said daily dosage units comprise a combination of estradiol in an amount ranging from about 0.1 to 5 mg and drospirenone in an amount ranging from about 0.25 to 6 mg; and 10 to 15 said daily dosage units comprise estradiol in an amount ranging from about 0.1 to 5 mg.

72. A preparation according to any of claims 65 to 71, wherein the number of daily dosage units is at least 21 or a multiple of 21 such as 2 to 12, particularly 2 to 8, such as 2 to 6.

73. A preparation according to any of claims 65 to 71, wherein the number of daily dosage units is 28 or a multiple of 28 such as 2 to 12, particularly 2 to 8 such as 2 to 6.

74. A preparation according to any of claims 65 to 71, wherein said daily dosage units comprise estradiol and/or drospirenone in micronized form or sprayed from a solution onto particles of inert carrier

75. A use of an estrogen (and/or naturally or synthetic derivative thereof) in sufficient amounts for the treatment of menopausal and post-menopausal symptoms and the prevention of osteoporosis in menopausal and post-menopausal women

18-01-2000

gen: 18/ 1/00 15:38;

EP00200183.2

JetFax M830; Pagina 48

SPEC

41

Tuesday 18 of Jan 2000, PV&P 33838600

->00 31 70 3403018

Page 4

38

and 6 β ,7 β ;15 β ;16 β -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone)
in sufficient amounts to protect the endometrium from the adverse effects of estrogen
for the manufacture of a medicament.

- 5 76. A use of an estrogen (and/or naturally or synthetic derivative thereof) in sufficient
amounts for the treatment of menopausal and post-menopausal symptoms and the
prevention of osteoporosis in menopausal and post-menopausal women
and 6 β ,7 β ;15 β ;16 β -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone)
in sufficient amounts to protect the endometrium from the adverse effects of estrogen
10 for the manufacture of a medicament for hormone replacement therapy.

ABSTRACT

Drospirenone for Hormone Replacement Therapy

A pharmaceutical composition comprising as a first active ingredient an estrogen, such as estradiol or estradiol valerate, in sufficient amounts to treat disorders and symptoms associated with deficient endogenous levels of estrogen in women, and as a second active ingredient $6\beta,7\beta;15\beta;16\beta$ -dimethylene-3-oxo-17 α -preg-4-ene-21,17-carbolactone (drospirenone, DRSP) in sufficient amounts to protect the endometrium from the adverse effects of estrogen is useful for, amongst others, treating peri-menopausal, menopausal and post-menopausal women. This composition may be used for hormone replacement therapy and may be administered as a multi-phased pharmaceutical preparation. This combination therapy may comprise continuous, sequential or interrupted administration, or combinations thereof, of DRSP and estrogen, each optionally in micronized form.