

Señores

DIVISION DE NUEVAS CREACIONES

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO

Bogotá, D.C.

SUPERINDUSTRIA Y COMERCIO
Radicación: 01000200 00000001 Folios:
Fecha (RAD): 2001-03-30 12:58:15
Trámite: 002 PATENTES D 1 REGISTRO 329 COMPLEMEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Re.: Expediente No. 01 - 20.200

Solicitud de Patente

Debidamente traducida, adjunto acompaño la copia oficial de la solicitud de patente básica estadounidense No. 60/189.175 la cual le solicito agregar a este expediente para los efectos legales respectivos.

De Ustedes atentamente,

GERMAN CASTILLO GRAU

T.P. No. 2978 del C.S.J.

64
60



A TODOS A QUIENES LLEGUE EL PRESENTE:

DEPARTAMENTO DE COMERCIO DE LOS ESTADOS UNIDOS

Oficina de Marcas y Patentes

26 de enero de 2001

LA PRESENTE CERTIFICA QUE LA ADJUNTA ES UNA COPIA TOMADA DE LOS REGISTROS DE LA OFICINA DE MARCAS Y PATENTES DE LOS ESTADOS UNIDOS, DE LOS DOCUMENTOS DE LA SOLICITUD DE PATENTE ABAJO IDENTIFICADA Y LA CUAL CUMPLE CON LOS REQUISITOS PARA QUE LE SEA CONFERIDA UNA FECHA DE RADICACIÓN SEGÚN 35 USC 111.

NÚMERO DE SOLICITUD 60/189.175

FECHA DE RADICACIÓN: 14 de marzo de 2000

Por Autoridad del

COMISIONADO DE MARCAS Y PATENTES

(FIRMA) N. WILLIAMS


Oficial de Certificaciones

(SELLO DORADO DE OBLEA)

ANTAGONISTAS DEL RECEPTOR IL-8

REIVINDICACIONES:

1. Un compuesto de la fórmula (I):



en donde

R_b es independientemente hidrógeno, NR_6R_7 , OH, OR_a , alquilo C_{1-5} , arilo, arilo-alquilo C_{1-4} , arilo, alqueno C_{2-4} ; cicloalquilo, cicloalquilo alquilo C_{1-5} , heteroarilo, heteroarilo-alquilo C_{1-4} , heteroarilo-alqueno C_{2-4} , heterocíclico, heterocíclico-alquilo C_{1-4} o un medio heterocíclico-alqueno C_{2-4} , todos cuyos medios pueden ser sustituido opcionalmente una o tres veces independientemente por halógeno; alquilo C_{1-4} sustituido por halo; alquilo C_{1-4} ; amino, amina sustituida por mono o di-alquilo C_{1-4} ; OR_a ; $C(O)R_a$; $NR_aC(O)OR_a$; $OC(O)NR_6R_7$; hidroxilo; $NR_6C(O)R_a$; $S(O)_mR_a$; $C(O)NR_6R_7$; $C(O)OH$; $C(O)OR_a$; $S(O)NR_6R_7$; $NHS(O)_2R_a$. Alternativamente, los dos sustituyentes R_b pueden unirse para formar un anillo de 3 a 10 miembros, sustituido opcionalmente y que contengan además de carbono, independientemente, de 1 a 3 NR_a , O, S, SO, o SO_2 medios que opcionalmente pueden ser insaturados;

R_a es un alquilo, arilo arilo-alquilo C_{1-4} , heteroarilo, heteroarilo-alquilo C_{1-4} , heterocíclico, $COOR_{13}$, o un medio heterocíclico-alquilo C_{1-4} , todos los cuales pueden ser sustituido opcionalmente;

m es un entero con un valor de 1 a 3;

66
62

m' es 0 o un entero con un valor de 1 o 2;

n es un entero con valor de 1 a 3;

q es 0, o un entero con un valor de 1 a 10;

t es 0 o un entero con un valor de 1 o 2;

s es un entero con un valor de 1 a 3;

R_1 es seleccionado independientemente de hidrógeno, halógeno, nitro, ciano, alquilo C_{1-10} , alquilo C_{1-10} sustituido por halo, alquenilo C_{2-10} , alcoxi C_{1-10} , alcoxi C_{1-10} sustituido por halo, azida, $S(O)_tR_4$, $(CR_8R_9)_qS(O)_tR_4$, hidroxí, alquilo C_{1-4} sustituido por hidroxí, arilo, arilo-alquilo C_{1-4} , arilo-alquenilo C_{2-10} , ariloxi, arilo-alquiloxi C_{1-4} , heteroarilo, heteroarilalquilo, heteroarilo-alquenilo C_{2-10} , heteroarilo-alquiloxi C_{1-4} , heterocíclico, heterociclo-alquilo C_{1-4} , heterocíclico-alquiloxi C_{1-4} , heterocíclico-alquenilo C_{2-10} , $(CR_8R_9)_qNR_4R_5$, $(CR_8R_9)_qC(O)NR_4R_5$, alquenilo C_{2-10} $C(O)NR_4R_5$, $(CR_8R_9)_qC(O)NR_4R_{10}$, $S(O)_3R_8$, $(CR_8R_9)_qC(O)R_{11}$, alquenilo C_{2-10} $C(O)R_{11}$, alquenilo C_{2-10} $C(O)OR_{11}$, $(CR_8R_9)_qC(O)OR_{11}$, $(CR_8R_9)_qOC(O)R_{11}$, $(CR_8R_9)_qNR_4C(O)R_{11}$, $(CR_8R_9)_qC(NR_4)NR_4R_5$, $(CR_8R_9)_qNR_4C(NR_5)R_{11}$, $(CR_8R_9)_qNHS(O)_2R_{13}$, $(CR_8R_9)_qS(O)SNR_4R_5$, o dos medios R_1 juntos pueden formar $O-(CH_2)_sO$ o un anillo saturado o insaturado de 5 a 6 miembros, y en donde los medios alquilo, arilo, arilalquilo, heteroarilo, heterocíclico pueden ser sustituidos opcionalmente;

R_4 y R_5 son independientemente hidrógeno, alquilo C_{1-4} sustituido opcionalmente, arilo sustituido opcionalmente, arilo-alquilo C_{1-4} sustituido opcionalmente, heteroarilo sustituido opcionalmente, heteroarilo-alquilo C_{1-4} sustituido opcionalmente, heterocíclico, heterocíclico-alquilo C_{1-4} , o R_4 y R_5 junto con el nitrógeno al que están fijos forman un anillo de 5 a 7 miembros que opcionalmente pueden comprender un heteroátomo adicional seleccionado de O, N y S;

62
63

R_6 y R_7 son independientemente hidrógeno, o un alquilo C_{1-4} , heteroarilo, arilo, alquilarilo, alquilo-heteroalquilo C_{1-4} o R_6 y R_7 junto con el nitrógeno al que están fijos forman un anillo de 5 a 7 miembros que opcionalmente puede contener un heteroátomo adicional seleccionado de oxígeno, nitrógeno o azufre, y cuyo anillo puede ser sustituido opcionalmente;

Y es $CR_{14}C_{15}$, NR_{14} , O, CO, $S(O)_i$,

R_8 es hidrógeno o alquilo C_{1-4} ;

R_9 es alquilo C_{1-4} ;

R_{10} es alquilo C_{1-10} $C(O)_2R_6$;

R_{11} es hidrógeno, alquilo C_{1-4} sustituido opcionalmente, arilo sustituido opcionalmente, ariloalquilo C_{1-4} sustituido opcionalmente, heteroarilo sustituido opcionalmente, heteroarilo-alquilo C_{1-4} sustituido opcionalmente, heterocíclico sustituido opcionalmente, o heterocíclico-alquilo C_{1-4} sustituido opcionalmente; y

R_{13} y R_{14} son independientemente hidrógeno, alquilo C_{1-4} sustituido opcionalmente, alquínilo C_{1-4} sustituido opcionalmente, alquénilo C_{1-4} sustituido opcionalmente, heteroarilo sustituido opcionalmente, heteroarilo-alquilo C_{1-4} sustituido opcionalmente, heterocíclico sustituido opcionalmente, heterocíclico-alquilo C_{1-4} sustituido opcionalmente, nitro OR_a , NR_6R_7 , $C(O)R_a$, $NR_{11}C(O)OR_a$, $OC(O)NR_6R_7$, $NR_{11}C(O)R_a$, $S(O)_mR_a$, NR_6R_7 , $C(O)R_a$, NR_{11} $C(O)OR_a$, $OC(O)NR_6R_7$, $NR_{11}C(O)R_a$, $S(O)_mR_a$, $C(O)NR_6R_7$, $C(O)OH$, $C(O)OR_a$, $S(O)_2NR_6R_7$, o $NR_{11}S(O)_2R_1$,

R_{20} es hidrógeno o alquilo, alquínilo, alquénilo, heterocíclico, heteroaromático todo lo cual puede ser sustituido opcionalmente,

o una sal farmacéuticamente aceptable de lo mismo.

2. El compuesto según la Reivindicación 1, en donde R_1 es sustituido en la posición 4 por un electrón que retira el medio.

5

68
64

3. El compuesto según la Reivindicación 3, en donde R_1 es halógeno, ciano o nitro.

4. El compuesto según la Reivindicación 3, en donde R_1 es halógeno.

5. El compuesto según la Reivindicación 4, en donde R_1 es independientemente, flúor, cloro o bromo.

6. El compuesto según la Reivindicación 1, en donde R_b es hidrógeno, alquilo C_{1-4} o alquilo C_{1-4} sustituido por $C(O)OH$, o $C(O)OR_a$.

7. El compuesto según la Reivindicación 1, seleccionado del grupo compuesto por:

N-alilo-N'-(3-aminosulfonilo-4-cloro-2-hidroxifenilo)úrea;

N-(3-aminosulfonilo-4-cloro-3-hidroxifenilo)-N'-isopropilurea;

N-(3-aminosulfonilo-4-cloro-3-hidroxifenilo)-N'-etilurea;

N-(3-aminosulfonilo-4-cloro-3-hidroxifenilo)-N'-propilurea; y

N-(3-aminosulfonilo-4-cloro-3-hidroxifenilo)-N'-(etoxicarbonil)metilurea.

8. Un compuesto según la Reivindicación 7 que es:

N-alilo-N'-(3-aminosulfonilo-4-cloro-2-hidroxifenilo)úrea;

N-(3-aminosulfonilo-4-cloro-3-hidroxifenilo)-N'-isopropilurea.

9. Un compuesto según la Reivindicación 7, en donde el compuesto es en su forma de sal sódica.

10. Un compuesto según la Reivindicación 7, en donde el compuesto es en su forma de sal potásica.

6

65

11. Una composición farmacéutica que comprende un compuesto según cualquiera de las Reivindicaciones 1 a 10, y un vehículo o diluyente farmacéuticamente aceptable.

12. Un método para tratar una enfermedad mediada por quimiocina, en donde la quimiocina se liga a un receptor IL-8 a o b, cuyo método consiste en administrar a dicho mamífero una cantidad efectiva de un compuesto de la fórmula según cualquiera de las Reivindicaciones 1 a 14.

13. El método según la Reivindicación 12, en donde el mamífero sufre de una enfermedad mediada por quimiocina seleccionada del grupo compuesto por; psoriasis, dermatitis atópica, osteoartritis, artritis reumatoide, asma, enfermedad pulmonar obstructiva crónica, síndrome de agotamiento respiratorio en adultos, enfermedad de intestinos inflamados, enfermedad de Crohn, colitis ulcerativa, ataque, shock séptico, esclerosis múltiple, shock entodóxico, sepsis gram negativa, síndrome de shock tóxico, lesión de reperfusión renal y cardíaca, glomerulonefritis, trombosis, reacción injerto-anfitrión, enfermedad de Alzheimer, rechazos a injertos, malaria, restenosis, angiogenesis, aterosclerosis, osteoporosis, gingivitis y liberación no deseada de células hematopoiéticas y enfermedades causadas por virus respiratorios, virus de herpes y virus de hepatitis, meningitis, fibrosis cística, trabajo de parto antes de tiempo, tos, prurito, disfunción de órganos múltiples, trauma, tensiones, desgarres, contusiones, artritis psoriática, herpes, encefalitis, vasculitis SNC, lesión cerebral traumática, tumores del SNC, hemorragia subaracnoide, trauma posquirúrgico, neumonitis intersticial, hipersensibilidad, artritis inducida por cristales, pancreatitis crónica, hepatitis alcohólica, enterocolitis necrotizante, sinusitis crónica, uveitis, polimiositis, vasculitis, acné, úlceras gástricas y duodenales, enfermedad celiaca, esofagitis, glositis, obstrucción de las vías respiratorias, hiperrespuesta de las vías respiratorias, bronquiolitis, obliterantes de bronquiolitis, neumonía, bronquiectasis, bronquiolitis, obliterantes de bronquilitis, bronquitis crónica, cor pulmonae, disnea, enfisema, hipercapnea, hiperinflación, hipoxemia, inflamaciones inducidas por hiperoxia, hipoxia, reducción en el volumen pulmonar por cirugía, fibrosis

70
66

pulmonar, hipertensión pulmonar, hipertropía ventricular derecha, sarcoidosis, enfermedad de las vías respiratorias pequeñas, no concordancia ventilación-perfusión, resuello, gripes y lupus.

8

27
67

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

January 26, 2001

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/189,175

FILING DATE: *March 14, 2000*



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

A handwritten signature in cursive script, appearing to read "N. Williams", is written over the printed name.

N. WILLIAMS
Certifying Officer

03/14/00
1c759 U.S. PTO

EXPRESS MAIL CERTIFICATE

"Express Mail" Mailing Label Number EL421195569US

Date Of Deposit: March 14, 2000

I Herby Certify That This Paper Or Fee Is Being Deposited With The United States Postal Service "Express Mail Post Office To Addressee" Service Under 37 CFR 1.18 On The Date Indicated Above And Is Addressed To: Assistant Commissioner for Patents, Washington, D.C. 20231.

Name Of Person Mailing Paper Or Fee

(Type Or Print) USAN M. Galen

Signature [Signature]

53-15-n

A/PNO

72
68

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION for PATENT under 37 CFR 1.53(c).

Docket No.

P51111P

INVENTOR(s) / APPLICANT(s)

Last Name	First Name	Middle Initial	Residence (City and Either State or Foreign Country)
WIDDOWSON QI	Katherine Jin	L.	King of Prussia, Pennsylvania King of Prussia, Pennsylvania

TITLE OF THE INVENTION (280 characters max)

IL-8 RECEPTOR ANTAGONISTS

CORRESPONDENCE ADDRESS

SMITHKLINE BEECHAM CORPORATION

Corporate Intellectual Property - UW2220

709 Swedeland Road

King of Prussia

Telephone No. 610-270-5019

Facsimile No. 610-270-5090

State

PA

Zip Code

19406-2799

Country

United States of America

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification	Number of Pages	36	☐ Small Entity Statement
☐ Drawings	Number of Sheets		☐ Other (specify)

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

☒ The Commissioner is hereby authorized to charge filing fees and credit Deposit Account No. 19-2570

PROVISIONAL FILING
FEE AMOUNT (\$)

\$150.00

Respectfully submitted,
Signature:

[Signature]
Soma G. Simoni

Date:

3/14/00

Registration No.:

37,444

☐ Additional inventors are being named on separately numbered sheets attached hereto.

PROVISIONAL APPLICATION FILING ONLY

SEND TO: Assistant Commissioner for Patents, Box Provisional Application, Washington, D.C. 20231.

N:\SOMA\CASES\VP51111PATProv.doc

60189175-031400

U.S. PTO
60/189175
03/14/00

IL-8 RECEPTOR ANTAGONISTS**FIELD OF THE INVENTION**

This invention relates to novel sulfonamide substituted diphenyl urea compounds, pharmaceutical compositions, processes for their preparation, and use thereof in treating IL-8, GRO α , GRO β , GRO γ , NAP-2, and ENA-78 mediated diseases.

BACKGROUND OF THE INVENTION

Many different names have been applied to Interleukin-8 (IL-8), such as neutrophil attractant/activation protein-1 (NAP-1), monocyte derived neutrophil chemotactic factor (MDNCF), neutrophil activating factor (NAF), and T-cell lymphocyte chemotactic factor. Interleukin-8 is a chemoattractant for neutrophils, basophils, and a subset of T-cells. It is produced by a majority of nucleated cells including macrophages, fibroblasts, endothelial and epithelial cells exposed to TNF, IL-1 α , IL-1 β or LPS, and by neutrophils themselves when exposed to LPS or chemotactic factors such as FMLP. M. Baggiolini et al., *J. Clin. Invest.* 84, 1045 (1989); J. Schroder et al., *J. Immunol.* 139, 3474 (1987) and *J. Immunol.* 144, 2223 (1990); Strieter, et al., *Science* 243, 1467 (1989) and *J. Biol. Chem.* 264, 10621 (1989); Cassatella et al., *J. Immunol.* 148, 3216 (1992).

GRO α , GRO β , GRO γ and NAP-2 also belong to the chemokine family. Like IL-8 these chemokines have also been referred to by different names. For instance GRO α , β , γ have been referred to as MGSA α , β and γ respectively (Melanoma Growth Stimulating Activity), see Richmond et al., *J. Cell Physiology* 129, 375 (1986) and Chang et al., *J. Immunol.* 148, 451 (1992). All of the chemokines of the α -family which possess the ELR motif directly preceding the CXC motif bind to the IL-8 B receptor (CXCR2).

IL-8, GRO α , GRO β , GRO γ , NAP-2, and ENA-78 stimulate a number of functions in vitro. They have all been shown to have chemoattractant properties for neutrophils, while IL-8 and GRO α have demonstrated T-lymphocytes, and basophilic chemotactic activity. In addition IL-8 can induce histamine release from basophils from both normal and atopic individuals. GRO- α and IL-8 can in addition, induce lysosomal enzyme release and respiratory burst from neutrophils. IL-8 has

00189175-031400

70

also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils without de novo protein synthesis. This may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many known diseases are characterized by massive neutrophil infiltration. As IL-8, GRO α , GRO β , GRO γ and NAP-2 promotes the accumulation and activation of neutrophils, these chemokines have been implicated in a wide range of acute and chronic inflammatory disorders including psoriasis and rheumatoid arthritis, Baggiolini et al., FEBS Lett. 307, 97 (1992); Miller et al., Crit. Rev. Immunol. 12, 17 (1992); Oppenheim et al., Annu. Rev. Immunol. 9, 617 (1991); Seitz et al., J. Clin. Invest. 87, 463 (1991); Miller et al., Am. Rev. Respir. Dis. 146, 427 (1992); Donnelly et al., Lancet 341, 643 (1993). In addition the ELR chemokines (those containing the amino acids ELR motif just prior to the CXC motif) have also been implicated in angiostasis, Strieter et al., Science 258, 1798 (1992).

In vitro, IL-8, GRO α , GRO β , GRO γ and NAP-2 induce neutrophil shape change, chemotaxis, granule release, and respiratory burst, by binding to and activating receptors of the seven-transmembrane, G-protein-linked family, in particular by binding to IL-8 receptors, most notably the IL-8 β receptor (CXCR2). Thomas et al., J. Biol. Chem. 266, 14839 (1991); and Holmes et al., Science 253, 1278 (1991). The development of non-peptide small molecule antagonists for members of this receptor family has precedent. For a review see R. Freidinger in: Progress in Drug Research, Vol. 40, pp. 33-98, Birkhauser Verlag, Basel 1993. Hence, the IL-8 receptor represents a promising target for the development of novel anti-inflammatory agents.

Two high affinity human IL-8 receptors (77% homology) have been characterized: IL-8R α , which binds only IL-8 with high affinity, and IL-8R β , which has high affinity for IL-8 as well as for GRO α , GRO β , GRO γ and NAP-2. See Holmes et al., *supra*; Murphy et al., Science 253, 1280 (1991); Lee et al., J. Biol. Chem. 267, 16283 (1992); LaRosa et al., J. Biol. Chem. 267, 25402 (1992); and Gayle et al., J. Biol. Chem. 268, 7283 (1993).

There remains a need for treatment, in this field, for compounds, which are capable of binding to the IL-8 α or β receptor. Therefore, conditions associated with an increase in IL-8 production (which is responsible for chemotaxis of neutrophil

12

60139175-031400

25
H

and T-cells subsets into the inflammatory site) would benefit by compounds, which are inhibitors of IL-8 receptor binding.


SUMMARY OF THE INVENTION

5 This invention provides for a method of treating a chemokine mediated disease, wherein the chemokine is one which binds to an IL-8 a or b receptor and which method comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In particular the chemokine is IL-8.

10 This invention also relates to a method of inhibiting the binding of IL-8 to its receptors in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I).

The present invention also provides for the novel compounds of Formula (I), and pharmaceutical compositions comprising a compound of Formula (I), and a
15 pharmaceutical carrier or diluent.

Compounds of Formula (I) useful in the present invention are represented by the structure:



20

wherein

R_b is independently hydrogen, NR₆R₇, OH, OR_a, C₁₋₅alkyl, aryl, arylC₁₋₄alkyl, aryl

C₂₋₄alkenyl; cycloalkyl, cycloalkyl C₁₋₅ alkyl, heteroaryl, heteroarylC₁₋₄alkyl,

heteroarylC₂₋₄ alkenyl, heterocyclic, heterocyclic C₁₋₄alkyl, or a heterocyclic

25 C₂₋₄alkenyl moiety, all of which moieties may be optionally substituted one to three

times independently by halogen; nitro; halosubstituted C₁₋₄ alkyl; C₁₋₄ alkyl;

amino, mono or di-C₁₋₄ alkyl substituted amine; OR_a; C(O)R_a; NR_aC(O)OR_a;

OC(O)NR₆R₇; hydroxy; NR₉C(O)R_a; S(O)_mR_a; C(O)NR₆R₇; C(O)OH;

96
72

$C(O)OR_a$; $S(O)_2NR_6R_7$; $NHS(O)_2R_a$. Alternatively, the two R_b substituents can join to form a 3-10 membered ring, optionally substituted and containing, in addition to carbon, independently, 1 to 3 NR_a , O, S, SO, or SO_2 moieties which can be optionally unsaturated;

- 5 R_a is an alkyl, aryl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclic, $COOR_g$, or a heterocyclic C_{1-4} alkyl moiety, all of which moieties may be optionally substituted;

m is an integer having a value of 1 to 3;

m' is 0, or an integer having a value of 1 or 2;

- 10 n is an integer having a value of 1 to 5;

q is 0, or an integer having a value of 1 to 10;

t is 0, or an integer having a value of 1 or 2;

s is an integer having a value of 1 to 3;

R_1 is independently selected from hydrogen, halogen, nitro, cyano, C_{1-10} alkyl,

- 15 halosubstituted C_{1-10} alkyl, C_{2-10} alkenyl, C_{1-10} alkoxy, halosubstituted C_{1-10} alkoxy, azide, $S(O)_tR_4$, $(CR_gR_g)_q S(O)_tR_4$, hydroxy, hydroxy substituted C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-10} alkenyl, aryloxy, aryl C_{1-4} alkyloxy, heteroaryl, heteroarylalkyl, heteroaryl C_{2-10} alkenyl, heteroaryl C_{1-4} alkyloxy, heterocyclic, heterocyclic C_{1-4} alkyl, heterocyclic C_{1-4} alkyloxy, heterocyclic C_{2-10}
- 20 alkenyl, $(CR_gR_g)_q NR_4R_5$, $(CR_gR_g)_q C(O)NR_4R_5$, C_{2-10} alkenyl $C(O)NR_4R_5$, $(CR_gR_g)_q C(O)NR_4R_{10}$, $S(O)_3R_8$, $(CR_gR_g)_q C(O)R_{11}$, C_{2-10} alkenyl $C(O)R_{11}$, C_{2-10} alkenyl $C(O)OR_{11}$, $(CR_gR_g)_q C(O)OR_{11}$, $(CR_gR_g)_q OC(O)R_{11}$, $(CR_gR_g)_q NR_4C(O)R_{11}$, $(CR_gR_g)_q C(NR_4)NR_4R_5$, $(CR_gR_g)_q NR_4C(NR_5)R_{11}$, $(CR_gR_g)_q NHS(O)_2R_a$, $(CR_gR_g)_q S(O)_2NR_4R_5$, or two R_1 moieties together may

- 25 form $O-(CH_2)_sO$ or a 5 to 6 membered saturated or unsaturated ring, and wherein the alkyl, aryl, arylalkyl, heteroaryl, heterocyclic moieties may be optionally substituted;

R_4 and R_5 are independently hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-4} alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl C_{1-4} alkyl, heterocyclic, heterocyclic C_{1-4} alkyl, or R_4 and R_5 together with the nitrogen to which they are attached form a 5

30

14

00189175-031400

to 7 member ring which may optionally comprise an additional heteroatom selected from O, N and S;

R₆ and R₇ are independently hydrogen, or a C₁₋₄ alkyl, heteroaryl, aryl, alkyl aryl, alkyl C₁₋₄ heteroalkyl or R₆ and R₇ together with the nitrogen to which they are

5 attached form a 5 to 7 member ring which ring may optionally contain an additional heteroatom is selected from oxygen, nitrogen or sulfur, and which ring may be optionally substituted;

R₈ is hydrogen or C₁₋₄ alkyl;

R₉ is C₁₋₄ alkyl;

10 R₁₀ is C₁₋₁₀ alkyl C(O)₂R₈;

R₁₁ is hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl;

15 R₁₃ and R₁₄ are independently hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted C₁₋₄ alkynyl, optionally substituted C₁₋₄ alkenyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄ alkyl, optionally substituted heterocyclic, optionally substituted heterocyclicC₁₋₄alkyl, halogen, nitro, OR_a, NR₆R₇, C(O)R_a, NR₁₁C(O)OR_a, OC(O)NR₆R₇, NR₁₁C(O)R_a, S(O)_mR_a,

20 C(O)NR₆R₇, C(O)OH, C(O)OR_a, S(O)₂NR₆R₇, or NR₁₁S(O)₂R_a.

R₂₀ is hydrogen or alkyl, alkynyl, alkenyl, heterocyclic, heteroaromatic all of which can be optional substituted

or a pharmaceutically acceptable salt thereof.

60189175-031400

DETAILED DESCRIPTION OF THE INVENTION

The compounds of Formula (I), may also be used in association with the veterinary treatment of mammals, other than humans, in need of inhibition of IL-8 or other chemokines which bind to the IL-8 α and β receptors. Chemokine mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted herein in the Methods of Treatment section.

Suitably, R_b is independently hydrogen, NR_6R_7 , OH, OR_a , C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{2-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, or a heterocyclic C_{2-4} alkenyl moiety, all of which moieties may be optionally substituted one to three times independently by halogen, nitro, halosubstituted C_{1-4} alkyl, C_{1-4} alkyl, amino, mono or di- C_{1-4} alkyl substituted amine, cycloalkyl, cycloalkyl C_{1-5} alkyl, OR_a , $C(O)R_a$, $NR_aC(O)OR_a$, $OC(O)NR_6R_7$, aryloxy, aryl C_{1-4} oxy, hydroxy, C_{1-4} alkoxy, $NR_9C(O)R_a$, $S(O)_mR_a$, $C(O)NR_6R_7$, $C(O)OH$, $C(O)OR_a$, $S(O)_2NR_6R_7$, $NHS(O)_2R_a$. Alternatively, the two R_b substituents can join to form a 3-10 membered ring, optionally substituted and containing, in addition to carbon, independently, 1 to 3 NR_9 , O, S, SO, or SO_2 moities which can be optionally substituted.

Suitably, R_a is an alkyl, aryl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclic, or a heterocyclic C_{1-4} alkyl moiety, all of which moieties may be optionally substituted.

Suitably, R_1 is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C_{1-10} alkyl, such as CF_3 , C_{1-10} alkyl, such as methyl, ethyl, isopropyl, or n-propyl, C_{2-10} alkenyl, C_{1-10} alkoxy, such as methoxy, or ethoxy; halosubstituted C_{1-10} alkoxy, such as trifluoromethoxy, azide, $(CR_gR_g)_q S(O)_tR_4$, wherein t is 0, 1 or 2, hydroxy, hydroxy C_{1-4} alkyl, such as methanol or ethanol, aryl, such as phenyl or naphthyl, aryl C_{1-4} alkyl, such as benzyl, aryloxy, such as phenoxy, aryl C_{1-4} alkyloxy, such as benzyloxy; heteroaryl, heteroarylalkyl, heteroaryl C_{1-4} alkyloxy; aryl C_{2-10} alkenyl, heteroaryl C_{2-10} alkenyl, heterocyclic C_{2-10} alkenyl,

$(CR_8R_9)_qNR_4R_5$, C₂₋₁₀ alkenyl C(O)NR₄R₅, $(CR_8R_9)_qC(O)NR_4R_5$,
 $(CR_8R_9)_qC(O)NR_4R_{10}$, S(O)₃H, S(O)₃R₈, $(CR_8R_9)_qC(O)R_{11}$, C₂₋₁₀ alkenyl C(O)
R₁₁, C₂₋₁₀ alkenyl C(O)OR₁₁, $(CR_8R_9)_qC(O)R_{11}$, $(CR_8R_9)_qC(O)OR_{11}$,
 $(CR_8R_9)_qOC(O)R_{11}$, $(CR_8R_9)_qNR_4C(O)R_{11}$, $(CR_8R_9)_qC(NR_4)NR_4R_5$,
5 $(CR_8R_9)_qNR_4C(NR_5)R_{11}$, $(CR_8R_9)_qNHS(O)_2R_8$, $(CR_8R_9)_qS(O)_2NR_4R_5$. All of
the aryl, heteroaryl, and heterocyclic-containing moieties may be optionally
substituted as defined herein below.

For use herein the term "the aryl, heteroaryl, and heterocyclic containing
moieties" refers to both the ring and the alkyl, or if included, the alkenyl rings, such
10 as aryl, arylalkyl, and aryl alkenyl rings. The term "moieties" and "rings" may be
interchangeably used throughout.

Suitably, R₄ and R₅ are independently hydrogen, optionally substituted C₁₋₄
alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally
substituted heteroaryl, optionally substituted heteroaryl C₁₋₄alkyl, heterocyclic,
15 heterocyclicC₁₋₄ alkyl, or R₄ and R₅ together with the nitrogen to which they are
attached form a 5 to 7 member ring which may optionally comprise an additional
heteroatom selected from O, N and S.

Suitably, R₈ is independently hydrogen or C₁₋₄ alkyl.

Suitably, R₉ is hydrogen or a C₁₋₄ alkyl;

20 Suitably, q is 0 or an integer having a value of 1 to 10.

Suitably, R₁₀ is C₁₋₁₀ alkyl C(O)₂R₈, such as CH₂C(O)₂H or
CH₂C(O)₂CH₃.

Suitably, R₁₁ is hydrogen, C₁₋₄ alkyl, aryl, aryl C₁₋₄ alkyl, heteroaryl,
heteroaryl C₁₋₄alkyl, heterocyclic, or heterocyclic C₁₋₄alkyl.

25 Suitably, R₁₂ is hydrogen, C₁₋₁₀ alkyl, optionally substituted aryl or
optionally substituted arylalkyl.

Suitably, R₈ is an alkyl, aryl C₁₋₄ alkyl, heteroaryl, heteroaryl-C₁₋₄alkyl,
heterocyclic, or a heterocyclicC₁₋₄ alkyl, wherein all of these moieties may all be
30 optionally substituted.

80
76

R₁₃ and R₁₄ are suitably hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄ alkyl, optionally substituted heterocyclic, optionally substituted heterocyclicC₁₋₄alkyl, halogen, nitro, OR_a, NR₆R₇,
 5 NR₁₁C(O)OR_a, OC(O)NR₆R₇, NR₁₁C(O)R_a, C(O)NR₆R₇, S(O)₂NR₆R₇, or NR₁₁S(O)₂R_a.

R₂₀ is suitably C₁₋₄ alkyl, alkenyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclic, or heterocyclicC₁₋₄alkyl all of which may be optionally substituted.

As used herein, "optionally substituted" unless specifically defined shall
 10 mean such groups as halogen, such as fluorine, chlorine, bromine or iodine, hydroxy; hydroxy substituted C₁₋₁₀alkyl, C₁₋₁₀ alkoxy, such as methoxy or ethoxy, S(O)_{m'}C₁₋₁₀ alkyl, wherein m' is 0, 1 or 2, such as methyl thio, methyl sulfinyl or methyl sulfonyl; amino, mono & di-substituted amino, such as in the NR₄R₅ group, NHC(O)R₁₁, C(O)NR₈R₁₁, C(O)OH, S(O)₂NR₈R₁₁, NHS(O)₂R₉, C₁₋₁₀ alkyl,
 15 such as methyl, ethyl, propyl, isopropyl, or t-butyl, halosubstituted C₁₋₁₀ alkyl, such as CF₃, an optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl, optionally substituted heterocyclic, optionally substituted heterocyclicalkyl, optionally substituted heteroaryl, optionally substituted heteroaryl alkyl, wherein these aryl, heteroaryl, or heterocyclic moieties
 20 may be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl, C₁₋₁₀ alkoxy; S(O)_mC₁₋₁₀ alkyl; amino, mono & di-substituted alkyl amino, such as in the NR₄R₅ group; C₁₋₁₀ alkyl, or halosubstituted C₁₋₁₀ alkyl, such as CF₃.

Suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of inorganic and organic acids, such as hydrochloric
 25 acid, hydrobromic acid, sulphuric acid, phosphoric acid, methane sulphonic acid, ethane sulphonic acid, acetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid and mandelic acid. In addition, pharmaceutically acceptable salts of compounds of Formula (I) may also be formed with a pharmaceutically
 30 acceptable cation. Suitable pharmaceutically acceptable cations are well known to

18

60189175-031400

21
77

those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations.

The following terms, as used herein, refer to:

- "halo" - all halogens, that is chloro, fluoro, bromo and iodo.
- 5 • "C₁₋₁₀alkyl" or "alkyl" - both straight and branched chain moieties of 1 to 10 carbon atoms, unless the chain length is otherwise limited, including, but not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl and the like.
- 10 • "cycloalkyl" is used herein to mean cyclic moiety, preferably of 3 to 8 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, and the like.
- 15 • "alkenyl" is used herein at all occurrences to mean straight or branched chain moiety of 2-10 carbon atoms, unless the chain length is limited thereto, including, but not limited to ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like.
- 20 • "aryl" - phenyl and naphthyl;
- "heteroaryl" (on its own or in any combination, such as "heteroaryloxy", or "heteroaryl alkyl") - a 5-10 membered aromatic ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O or S, such as, but not limited, to pyrrole, pyrazole, furan, thiophene, quinoline, isoquinoline, quinazolinyl, pyridine, pyrimidine, oxazole, tetrazole, thiazole, thiadiazole, triazole, imidazole, or benzimidazole.
- 25 • "heterocyclic" (on its own or in any combination, such as "heterocyclicalkyl") - a saturated or partially unsaturated 4-10 membered ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O, or S; such as, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, tetrahydropyran, thiomorpholine, or imidazolidine. Furthermore, sulfur may be optionally oxidized to the sulfone or the sulfoxide.
- 30 • "arylalkyl" or "heteroarylalkyl" or "heterocyclicalkyl" is used herein to mean C₁₋₁₀ alkyl, as defined above, attached to an aryl, heteroaryl or heterocyclic moiety, as also defined herein, unless otherwise indicated.

82
78

• "sulfinyl" - the oxide S (O) of the corresponding sulfide, the term "thio" refers to the sulfide, and the term "sulfonyl" refers to the fully oxidized S(O)₂ moiety.

• "wherein two R₁ moieties may together form a 5 or 6 membered saturated or unsaturated ring" is used herein to mean the formation of an aromatic ring system, such as naphthalene, or is a phenyl moiety having attached a 6 membered partially saturated or unsaturated ring such as a C₆ cycloalkenyl, i.e. hexene, or a C₅ cycloalkenyl moiety, such as cyclopentene.

Illustrative compounds of Formula (I) include:


- 10 N-allyl-N'-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)urea;
N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-isopropylurea;
N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-ethylurea;
N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-propylurea; and
N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-(ethoxycarbonyl)methylurea.

15

METHODS OF PREPARATION

The compounds of Formulas (I) may be obtained by applying synthetic procedures, some of which are illustrated in the Schemes below. The synthesis provided for in these Schemes is applicable for the producing compounds of Formulas (I), having a
20 variety of different R, R_b, and R₂₀ groups which are reacted, employing optional substituents which are suitably protected, to achieve compatibility with the reactions outlined herein. Subsequent deprotection, in those cases, then affords compounds of the nature generally disclosed. Once the urea nucleus has been established, further compounds of these formulas may be prepared by applying standard techniques for
25 functional group interconversion, well known in the art.

Scheme 1



a) i) NCS, AcOH, H₂O, ii NRR'H, pyr b) H₂SO₄, HNO₃ c) NaOAc, 18-crown-6


5 d) H₂SO₄, MeOH e) Pd/C, H₂ f) RCNO, DMF

The desired 4-chloro N-(3-sulfonamido-2-hydroxy phenyl)-N'-alkyl urea can synthesized from the commercially available 2, 6-dichloro thiophenol using the procedure elaborated above in Scheme 1. The thiol can be oxidized to the
10 corresponding sulfonyl halide using a halogenating agent, such as NCS, NBS, Cl₂ or Br₂, in the presence of a protic solvent, such as water, acetic acid, or an alcohol or combination. The yield may be increased if a buffering agent, such as sodium or potassium acetate is included in the reaction mixture, and the reaction is conducted at or below room temperature. The corresponding sulfonyl halide can then be
15 condensed with an amine in presence of a base such as pyridine, triethyl amine, potassium carbonate or sodium hydride to form the analogous sulfonamide 2-scheme 1. The dichlorosulfonamide 2-scheme 1 can be nitrated using strong nitrating conditions such as nitric acid in sulfuric acid to form the aromatic nitro compound 3-scheme 1. The chlorine ortho to the nitro group can be selectively hydrolyzed using

84
30

- acetate salt such as sodium acetate in the presence of a crown ether, such as 18-crown-6, to form the acetate 4-scheme 1. The acetate group can be hydrolyzed under acidic conditions in an alcohol solvent such as methanol or ethanol with a catalytic amount of acid to form the phenol 5-scheme 1. The nitro can be reduced by
- 5 conditions well known in the art such as hydrogen and palladium on carbon, tin chloride in methanol, zinc in acetic acid or thiol to form the corresponding aniline 5-scheme 1. The aniline can then be coupled with a commercially available isocyanate or thioisocyanate to form the desired urea or thio urea. Alternatively the desired isocyanates can be made by condensing the amine with triphosgene in the presence
- 10 of base (such as potassium carbonate) or by reacting the carboxylic acid with diphenyl phosphoryl azide in the presence of a base (such as triethyl amine).

Scheme 2




- 15 a) NaH , $\text{R}'\text{X}$ b) NaH , $\text{R}''\text{X}$

- If the sulfonamide 1-scheme 2 (3-scheme 1) is unfunctionalized $\text{R}'=\text{R}''=\text{H}$ then it can be functionalized as required herein, by alkylation. The sulfonamide is deprotonated using a base such as sodium hydride and then alkylated using an alkyl halide such as benzyl bromide or methyl iodide form 2-scheme 2. The sulfonamide can then be
- 20 alkylated a second time using sodium hydride and another alkyl halide to form 3-scheme 2. This compound can then be converted to the desired urea using the process elaborated in scheme 1.

85
81

Scheme 3



a) i) NCS, AcOH, H₂O ii) NaOH MeOH b) H₂SO₄, HNO₃ c) NaOH MeOH d) POCl₅,

5 POCl₃ e) HNR'R'', Et₃N

An alternative route to 5-scheme 3 (3-scheme 1) is outlined above, in scheme 3 wherein the commercially available 2,6-dichloro thiol can be oxidized to the sulfonyl halide using a halogenating agent such as NCS, NBS, chlorine or bromine in the presence of a protic solvent such as alcohol, acetic acid or water. The sulfonyl halide can be hydrolyzed by using a metal hydroxide such as sodium or potassium hydroxide to form the corresponding sulfonic acid salt. The sulfonic acid salt can then be nitrated under nitration conditions such as nitric acid in a solvent of strong acid such as sulfuric acid to form the nitro phenyl sulfonic acid 3-scheme 3. The sulfonic acid 3-scheme 3 can be converted to the sulfonamide 5-scheme 3 using a three step procedure involving the formation of the metal salt using a base such as sodium hydroxide, sodium hydride or sodium carbonate to form 4-scheme 3. The sulfonic acid salt is then converted to the sulfonyl chloride using POCl₅ with POCl₃ as a solvent. The sulfonyl chloride can then be converted to the corresponding sulfonamide using the desired amine HNR'R'' in triethyl amine at temperatures ranging from -78 °C to 60 °C to form the corresponding sulfonamide 5-scheme 3 (3-scheme 1). The sulfonamide 5-scheme 3 can be further elaborated by the methods contained in scheme 1. This method is not limited to the 2,6-dichloro thiol it can also be applied to the 2,6-difluoro thiol, 2,6-dibromo thiol and the 2,6-diiodo thiol. The halogens in these compounds can be converted to the corresponding cyano,

23

60189175.031400

B6
82

amino, thiol, or alkoxy compounds by nucleophilic displacement reactions using nucleophiles such as alkyl thiolates, alkoxides, amine and cyanides. The halogens can also be further functionalized by palladium coupling and carbonylation reactions, well known in the art, to form the corresponding amido, carbonyl, alkenyl, alkyl, phenyl and heterocyclic substituted products as required by Formula (I).

SYNTHETIC EXAMPLES

The invention will now be described by reference to the following examples, which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. All temperatures are given in degrees centigrade, all solvents are highest available purity and all reactions run under anhydrous conditions in an argon atmosphere unless otherwise indicated.

In the Examples, all temperatures are in degrees Centigrade (°C). Mass spectra were performed upon a VG Zab mass spectrometer using fast atom bombardment, unless otherwise indicated. ¹H-NMR (hereinafter "NMR") spectra were recorded at 250 MHz using a Bruker AM 250 or Am 400 spectrometer. Multiplicities indicated are: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet and br indicates a broad signal. Sat. indicates a saturated solution, eq indicates the proportion of a molar equivalent of reagent relative to the principal reactant.

General synthesis of 2-hydroxy-3-amino-6-chlorobenzenesulfonamide

a) 2,6-dichlorobenzenesulfonyl chloride

Into a mixture of 200 milliliters (hereinafter "mL") of acetic acid, water and dichloromethane (3/1/4, v/v/v), 2,6 dichlorobenzenethiol (10.0 grams (hereinafter "g"), 55.8 millimoles (hereinafter "mmol"), N-chlorosuccinimide (37.28 g, 279 mmol) and potassium acetate (2.29 g, 27.9 mmol) were added. The resulting mixture was stirred at 0°C, then warmed to room temperature overnight. The mixture was then diluted with 200 mL of dichloromethane, and washed with water (100 mL x 3). The organic layer was dried (Na₂SO₄) and concentrated to give the desired product (11 g, 80%). ¹H NMR (CDCl₃): δ 7.57 (d, 2H), 7.47 (t, 1H).

24

00400-031400-52168109

b) 2,6-dichlorobenzenesulfonamide

A solution of 2,6-dichlorobenzenesulfonyl chloride (10.50 g, 42.77 mmol) in 100 mL of pyridine was added into 100 mL of pyridine dropwise while anhydrous ammonia gas was passing through the solution simultaneously for 4 hours at 0°C.

- 5 The mixture was acidified to pH >1 with 6N aq. HCl, then extracted with ethyl acetate. The combined organic layer was then dried (Na_2SO_4) and concentrated to give the desired product (8.69 g, 90%). EI-MS (m/z) 225.0, 227.1 (M).

c) 2,6-dichloro-3-nitrobenzenesulfonamide

- 10 Into a solution of 2,6-dichlorobenzenesulfonamide (7.8 g, 34.5 mmol) in 30 mL of concentrated sulfuric acid at 0°, nitric acid (1.74 mL, 41.4 mmol) was added dropwise. The mixture was stirred at 0°C for 2 hours, then 200 mL of water was added to produce a precipitate. The resulting mixture was filtered. The white solid was collected, washed with water and dried *in vacuo* to give the desired product
- 15 (7.17 g, 76%). ^1H NMR ($\text{DMSO}-d_6$): δ 8.25 (s, 2H), 8.20 (d, 1H), 7.92 (d, 1H).

d) 2-acetoxy-3-nitro-6-chlorobenzenesulfonamide

- A solution of 2,6-dichloro-3-nitrobenzenesulfonamide (2.04 g, 7.5 mmol), potassium acetate (2.21 g, 22.5 mmol) and 18-crown-6 (5.95 g, 22.5 mmol) in 50 mL of
- 20 dimethyl sulfoxide was heated to 45°C for 7 days. The mixture was acidified with 1N aq. HCl, and extracted with ethyl acetate. The organic layer was concentrated to give the crude. Column chromatography on silica gel, eluting with ethyl acetate/hexane/acetic acid (50/49/1, v/v/v), gave the desired product (1.67 g, 76%).
- EI-MS (m/z) 293.1, 295.1 (M).

25

e) 2-hydroxy-3-nitro-6-chlorobenzenesulfonamide

- A solution of 2-acetoxy-3-nitro-6-chlorobenzenesulfonamide (1.72 g, 5.83 mmol), chlorotrimethylsilane (2 mL) and fuming sulfuric acid (0.5 mL) in methanol was heated to reflux for 20 hours. The solvent was evaporated. The residue was diluted
- 30 with ethyl acetate and washed with water. The organic layer was then dried (Na_2SO_4) and concentrated to give the desired product (1.0 g, 68%). EI-MS (m/z) 251.1, 253.2 (M).

88
84

f) 2-hydroxy-3-amino-6-chlorobenzene sulfonamide

To a solution of 2-hydroxy-3-nitro-6-chlorobenzenesulfonamide (1.1 g, 4.36 mmol) in ethyl acetate, was added 10 % Pd/C (500 mg). The mixture was flushed with hydrogen, and then stirred under a hydrogen atmosphere at balloon pressure for 4 hours at room temperature. The mixture was filtered through celite and the celite was washed with methanol. The solvent was evaporated to give the desired product (91%). EI-MS (m/z) 221.1, 223.1 (M⁺).

Example 1Preparation of N-allyl-N'-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)urea

- 10 A solution of 3-amino-6-chloro-2-hydroxybenzenesulfonamide (40 mg, 0.18 mmol) and allyl isocyanate (19 μ L, 0.22 mmol) in 1 mL of N,N-dimethyl-formamide was stirred at room temperature for 20 hours. The mixture was diluted with ethyl acetate and washed with water to give the crude. Purification upon column chromatograph on silica gel, eluting with ethyl acetate/hexane (70/30, v/v), gave the desired product
- 15 (15 mg, 27%). EI-MS (m/z) 304.1, 306.4 (M⁺).

Example 2Preparation of N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-isopropylurea

- A solution of 3-amino-6-chloro-2-hydroxybenzenesulfonamide (150 mg, 0.67 mmol) and isopropyl isocyanate (80 μ L, 0.81 mmol) in 1 mL of N,N-dimethyl-formamide was stirred at room temperature for 20 hours. The mixture was diluted with ethyl acetate and washed with water to give the crude. Purification upon column chromatograph on silica gel, eluting with ethyl acetate/hexane (50/50, v/v), followed by recrystallization from acetone and hexane, gave the desired product
- 25 (111 mg, 54%). LC-MS (m/z) 308.0 (M⁺).

Example 3Preparation of N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-ethylurea

- A solution of 3-amino-6-chloro-2-hydroxybenzenesulfonamide (150 mg, 0.67 mmol) and ethyl isocyanate (64 μ L, 0.81 mmol) in 1 mL of N,N-dimethyl-formamide was stirred at room temperature for 20 hours. The mixture was diluted with ethyl acetate and washed with water to give the crude. Purification upon

column chromatograph on silica gel, eluting with ethyl acetate/hexane (50/50, v/v), followed by recrystallization from acetone and hexane, gave the desired product (88 mg, 44%). LC-MS (m/z) 294.0 (M⁺).

5

Example 4

Preparation of N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-propylurea

A solution of 3-amino-6-chloro-2-hydroxybenzenesulfonamide (150 mg, 0.67 mmol) and propyl isocyanate (76 μ L, 0.81 mmol) in 1 mL of N,N-dimethylformamide was stirred at room temperature for 20 hours. The mixture was diluted with ethyl acetate and washed with water to give the crude. Purification upon column chromatograph on silica gel, eluting with ethyl acetate/hexane (50/50, v/v), followed by recrystallization from acetone and hexane, gave the desired product (92 mg, 44%). LC-MS (m/z) 308.2 (M⁺).

10

Example 5

Preparation of N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-(ethoxycarbonyl)methylurea

A solution of 3-amino-6-chloro-2-hydroxybenzenesulfonamide (170 mg, 0.76 mmol) and ethyl isocyanatoacetate (103 μ L, 0.92 mmol) in 1.5 mL of N,N-dimethylformamide was stirred at room temperature for 20 hours. The mixture was diluted with ethyl acetate and washed with water to give the crude. Purification upon column chromatograph on silica gel, eluting with ethyl acetate/hexane (60/40, v/v), followed by Gilson HPLC separation, gave the desired product (20 mg, 7%). LC-MS (m/z) 352.0 (M⁺).

15

METHOD OF TREATMENT

The compounds of Formula (I), or a pharmaceutically acceptable salt thereof can be used in the manufacture of a medicine for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by excessive or unregulated IL-8 cytokine production by such mammal's cell, such as but not limited to monocytes and/or macrophages, or other chemokines

20

90
86

which bind to the IL-8 α or β receptor, also referred to as the type I or type II receptor.

Accordingly, the present invention provides a method of treating a chemokine mediated disease, wherein the chemokine is one which binds to an IL-8 α or β receptor and which method comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In particular, the chemokines are IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78.

The compounds of Formula (I) are administered in an amount sufficient to inhibit cytokine function, in particular IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78, such that they are biologically regulated down to normal levels of physiological function, or in some case to subnormal levels, so as to ameliorate the disease state. Abnormal levels of IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 for instance in the context of the present invention, constitute: (i) levels of free IL-8 greater than or equal to 1 picogram per mL; (ii) any cell associated IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 above normal physiological levels; or (iii) the presence of IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 above basal levels in cells or tissues in which IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 respectively, is produced.

The compounds of Formula (I), in generally have been shown to have a longer $t_{1/2}$ and improved oral bioavailability over the compounds disclosed in WO 96/25157 and WO 97/29743 whose disclosures are incorporated herein by reference.

There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. Chemokine mediated diseases include psoriasis, atopic dermatitis, osteo arthritis, rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, adult respiratory distress syndrome, inflammatory bowel disease, Crohn's disease, ulcerative colitis, stroke, septic shock, multiple sclerosis, endotoxic shock, gram negative sepsis, toxic shock syndrome, cardiac and renal reperfusion injury, glomerulonephritis, thrombosis, graft vs. host reaction, Alzheimer's disease, allograft rejections, malaria, restenosis, angiogenesis, atherosclerosis, osteoporosis, gingivitis and undesired hematopoietic stem cells release and diseases caused by respiratory viruses, herpes viruses, and hepatitis viruses, meningitis, cystic fibrosis, pre-term labor, cough, pruritus, multi-

28

E0189175.031400

gl
87

organ dysfunction, trauma, strains, sprains, contusions, psoriatic arthritis, herpes, encephalitis, CNS vasculitis, traumatic brain injury, CNS tumors, subarachnoid hemorrhage, post surgical trauma, interstitial pneumonitis, hypersensitivity, crystal induced arthritis, acute and chronic pancreatitis, acute alcoholic hepatitis,

5 necrotizing enterocolitis, chronic sinusitis, uveitis, polymyositis, vasculitis, acne, gastric and duodenal ulcers, celiac disease, esophagitis, glossitis, airflow obstruction, airway hyperresponsiveness, bronchiolitis obliterans organizing pneumonia, bronchiectasis, bronchiolitis, bronchiolitis obliterans, chronic bronchitis, cor pulmonae, dyspnea, emphysema, hypercapnea, hyperinflation, hypoxemia,

10 hyperoxia-induced inflammations, hypoxia, surgical lung volume reduction, pulmonary fibrosis, pulmonary hypertension, right ventricular hypertrophy, sarcoidosis, small airway disease, ventilation-perfusion mismatching, wheeze, colds and lupus.

These diseases are primarily characterized by massive neutrophil infiltration,

15 T-cell infiltration, or neovascular growth, and are associated with increased IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 production which is responsible for the chemotaxis of neutrophils into the inflammatory site or the directional growth of endothelial cells. In contrast to other inflammatory cytokines (IL-1, TNF, and IL-6), IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 have the unique property of

20 promoting neutrophil chemotaxis, enzyme release including but not limited to elastase release as well as superoxide production and activation. The α -chemokines but particularly, GRO α , GRO β , GRO γ , NAP-2 or ENA-78, working through the IL-8 type I or II receptor can promote the neovascularization of tumors by promoting the directional growth of endothelial cells. Therefore, the inhibition of IL-8 induced

25 chemotaxis or activation would lead to a direct reduction in the neutrophil infiltration.

Recent evidence also implicates the role of chemokines in the treatment of HIV infections, Littleman *et al.*, Nature 381, pp. 661 (1996) and Koup *et al.*, Nature 381, pp. 667 (1996).

30 Present evidence also indicates the use of IL-8 inhibitors in the treatment of atherosclerosis. The first reference, Boisvert *et al.*, J. Clin. Invest. 1998, 101:353-363 shows, through bone marrow transplantation, that the absence of IL-8 receptors

29

60189175-031400

JPC
88

on stem cells (and, therefore, on monocytes/macrophages) leads to a reduction in the development of atherosclerotic plaques in LDL receptor deficient mice. Additional supporting references are: Apostolopoulos, et al., Arterioscler. Thromb. Vasc. Biol. 1996, 16:1007-1012; Liu, et al., Arterioscler. Thromb. Vasc. Biol. 1997, 17:317-323; Rus, et al., Atherosclerosis 1996, 127:263-271.; Wang et al., J. Biol. Chem. 1996, 271:8837-8842; Yue, et al., Eur. J. Pharmacol. 1993, 240:81-84; Koch, et al., Am. J. Pathol. 1993, 142:1423-1431.; Lee, et al., Immunol. Lett. 1996, 53, 109-113.; and Terkeltaub et al., Arterioscler. Thromb. 1994, 14:47-53.

The present invention also provides for a means of treating, in an acute setting, as well as preventing, in those individuals deemed susceptible to, CNS injuries by the chemokine receptor antagonist compounds of Formula (I).

CNS injuries as defined herein include both open or penetrating head trauma, such as by surgery, or a closed head trauma injury, such as by an injury to the head region. Also included within this definition is ischemic stroke, particularly to the brain area.

Ischemic stroke may be defined as a focal neurologic disorder that results from insufficient blood supply to a particular brain area, usually as a consequence of an embolus, thrombi, or local atheromatous closure of the blood vessel. The role of inflammatory cytokines in this area has been emerging and the present invention provides a mean for the potential treatment of these injuries. Relatively little treatment, for an acute injury such as these has been available.

TNF- α is a cytokine with proinflammatory actions, including endothelial leukocyte adhesion molecule expression. Leukocytes infiltrate into ischemic brain lesions and hence compounds which inhibit or decrease levels of TNF would be useful for treatment of ischemic brain injury. See Liu *et al.*, Stroke, Vol. 25., No. 7, pp. 1481-88 (1994) whose disclosure is incorporated herein by reference.

Models of closed head injuries and treatment with mixed 5-LO/CO agents is discussed in Shohami *et al.*, J. of Vasc. & Clinical Physiology and Pharmacology, Vol. 3, No. 2, pp. 99-107 (1992) whose disclosure is incorporated herein by reference. Treatment, which reduced edema formation, was found to improve functional outcome in those animals treated.

The compounds of Formula (I) are administered in an amount sufficient to inhibit IL-8, binding to the IL-8 alpha or beta receptors, from binding to these receptors, such as evidenced by a reduction in neutrophil chemotaxis and activation. The discovery that the compounds of Formula (I) are inhibitors of IL-8 binding is based upon the effects of the compounds of Formulas (I) in the *in vitro* receptor binding assays which are described herein. The compounds of Formula (I) have been shown to be inhibitors of type II IL-8 receptors.

As used herein, the term "IL-8 mediated disease or disease state" refers to any and all disease states in which IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 plays a role, either by production of IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 themselves, or by IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 causing another monokine to be released, such as but not limited to IL-1, IL-6 or TNF. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to IL-8, would therefore be considered a disease state mediated by IL-8.

As used herein, the term "chemokine mediated disease or disease state" refers to any and all disease states in which a chemokine which binds to an IL-8 α or β receptor plays a role, such as but not limited to IL-8, GRO- α , GRO- β , GRO γ , NAP-2 or ENA-78. This would include a disease state in which, IL-8 plays a role, either by production of IL-8 itself, or by IL-8 causing another monokine to be released, such as but not limited to IL-1, IL-6 or TNF. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to IL-8, would therefore be considered a disease stated mediated by IL-8.

As used herein, the term "cytokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule, which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines, regardless of which cells produce them. For instance, a monokine is generally referred to as being produced and secreted by a mononuclear cell, such as a macrophage and/or monocyte. Many other cells however also produce monokines, such as natural killer cells, fibroblasts, basophils, neutrophils, endothelial cells, brain astrocytes, bone

Gd
90

marrow stromal cells, epidermal keratinocytes and B-lymphocytes. Lymphokines are generally referred to as being produced by lymphocyte cells. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF- α) and Tumor Necrosis Factor beta (TNF- β).

As used herein, the term "chemokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response, similar to the term "cytokine" above. A chemokine is primarily secreted through cell transmembranes and causes chemotaxis and activation of specific white blood cells and leukocytes, neutrophils, monocytes, macrophages, T-cells, B-cells, endothelial cells and smooth muscle cells. Examples of chemokines include, but are not limited to IL-8, GRO- α , GRO- β , GRO- γ , NAP-2, ENA-78, IP-10, MIP-1 α , MIP- β , PF4, and MCP 1, 2, and 3.

In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof in therapy, it will normally be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice. This invention, therefore, also relates to a pharmaceutical composition comprising an effective, non-toxic amount of a compound of Formula (I) and a pharmaceutically acceptable carrier or diluent.

Compounds of Formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions incorporating such may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally, topically, parenterally or by inhalation. The compounds of Formula (I) may be administered in conventional dosage forms prepared by combining a compound of Formula (I) with standard pharmaceutical carriers according to conventional procedures. The compounds of Formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active compound. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable character or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The

carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25mg to about 1g. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

Compounds of Formula (I) may be administered topically, that is by non-systemic administration. This includes the application of a compound of Formula (I) externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the Formulation. It may however comprise as much as 10% w/w but preferably will comprise less than 5% w/w, more preferably from 0.1% to 1% w/w of the Formulation.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those

for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

5 Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage;
10 an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or a macrogel. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surfactant such as a sorbitan ester or a polyoxyethylene derivative thereof. Suspending agents such
15 as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent
20 and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-100°C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of
25 bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted alcohol and propylene glycol.

Compounds of formula (I) may be administered parenterally, that is by
30 intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are generally preferred. Appropriate dosage forms for such

60189175-031400

9A
93

administration may be prepared by conventional techniques. Compounds of Formula (I) may also be administered by inhalation that is by intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques.

For all methods of use disclosed herein for the compounds of Formula (I) the daily oral dosage regimen will preferably be from about 0.01 to about 80 mg/kg of total body weight. The daily parenteral dosage regimen about 0.001 to about 80 mg/kg of total body weight. The daily topical dosage regimen will preferably be from 0.1 mg to 150 mg, administered one to four, preferably two or three times daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1 mg/kg per day. It will also be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a compound of Formula (I) or a pharmaceutically acceptable salt thereof will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of a compound of Formula (I) or a pharmaceutically acceptable salt thereof given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

The invention will now be described by reference to the following biological examples, which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

BIOLOGICAL EXAMPLES

The IL-8, and GRO- α chemokine inhibitory effects of compounds of the present invention are determined by the following *in vitro* assay:

Receptor Binding Assays:

[¹²⁵I] IL-8 (human recombinant) is obtained from Amersham Corp., Arlington Heights, IL, with specific activity 2000 Ci/mmol. GRO- α is obtained from

35

50189175-031400

NEN- New England Nuclear. All other chemicals are of analytical grade. High levels of recombinant human IL-8 type α and β receptors were individually expressed in Chinese hamster ovary cells as described previously (Holmes, *et al.*, Science, 1991, 253, 1278). The Chinese hamster ovary membranes were

5 homogenized according to a previously described protocol (Haour, *et al.*, J. Biol. Chem., 249 pp 2195-2205 (1974)). Except that the homogenization buffer is changed to 10mM Tris-HCL, 1mM MgSO₄, 0.5mM EDTA (ethylene-diaminetetra-acetic acid), 1mM PMSF (α -toluenesulphonyl fluoride), 0.5 mg/L Leupeptin, pH 7.5. Membrane protein concentration is determined using Pierce Co. micro-assay kit

10 using bovine serum albumin as a standard. All assays are performed in a 96-well micro plate format. Each reaction mixture contains ¹²⁵I IL-8 (0.25 nM) or ¹²⁵I GRO- α and 0.5 μ g/mL of IL-8R α or 1.0 μ g/mL of IL-8R β membranes in 20 mM Bis-Trispropane and 0.4 mM Tris HCl buffers, pH 8.0, containing 1.2 mM MgSO₄, 0.1 mM EDTA, 25 mM Na and 0.03% CHAPS. In addition, drug or compound of

15 interest is added which has been pre-dissolved in DMSO so as to reach a final concentration of between 0.01nM and 100 μ M. The assay is initiated by addition of ¹²⁵I-IL-8. After 1 hour at room temperature the plate is harvested using a Tomtec 96-well harvester onto a glass fiber filtermat blocked with 1% polyethylenimine/ 0.5% BSA and washed 3 times with 25 mM NaCl, 10 mM TrisHCl, 1 mM MgSO₄,

20 0.5 mM EDTA, 0.03 % CHAPS, pH 7.4. The filter is then dried and counted on the Betaplate liquid scintillation counter. The recombinant IL-8 R α , or Type I, receptor is also referred to herein as the non-permissive receptor and the recombinant IL-8 R β , or Type II, receptor is referred to as the permissive receptor.

Representative compounds of Formula (I), Examples 1 to 106 have

25 exhibited positive inhibitory activity in this assay at IC₅₀ levels < 30 μ M.

Chemotaxis Assay :

The *in vitro* inhibitory properties of these compounds are determined in the neutrophil chemotaxis assay as described in Current Protocols in Immunology, vol.

30 I, Suppl 1, Unit 6.12.3., whose disclosure is incorporated herein by reference in its entirety. Neutrophils were isolated from human blood as described in Current

99
95

Protocols in Immunology Vol. I, Suppl 1 Unit 7.23.1, whose disclosure is incorporated herein by reference in its entirety. The chemoattractants IL-8, GRO- α , GRO- β , GRO- γ and NAP-2 are placed in the bottom chamber of a 48 multiwell chamber (Neuro Probe, Cabin John, MD) at a concentration between 0.1 and 100 nM. The two chambers are separated by a 5 μ M polycarbonate filter. When compounds of this invention are tested, they are mixed with the cells (0.001 - 1000 nM) just prior to the addition of the cells to the upper chamber. Incubation is allowed to proceed for between about 45 and 90 min at about 37°C in a humidified incubator with 5% CO₂. At the end of the incubation period, the polycarbonate

membrane is removed and the top side washed, the membrane then stained using the Diff Quick staining protocol (Baxter Products, McGaw Park, IL, USA). Cells which have chemotaxed to the chemokine are visually counted using a microscope. Generally, four fields are counted for each sample, these numbers are averaged to give the average number of cells which had migrated. Each sample is tested in triplicate and each compound repeated at least four times. To certain cells (positive control cells) no compound is added, these cells represent the maximum chemotactic response of the cells. In the case where a negative control (unstimulated) is desired, no chemokine is added to the bottom chamber. The difference between the positive control and the negative control represents the chemotactic activity of the cells.

Elastase Release Assay:

The compounds of this invention are tested for their ability to prevent Elastase release from human neutrophils. Neutrophils are isolated from human blood as described in Current Protocols in Immunology Vol. I, Suppl 1 Unit 7.23.1. PMNs 0.88 x 10⁶ cells suspended in Ringer's Solution (NaCl 118, KCl 4.56, NaHCO₃ 25, KH₂PO₄ 1.03, Glucose 11.1, HEPES 5 mM, pH 7.4) are placed in each well of a 96 well plate in a volume of 50 μ L. To this plate is added the test compound (0.001 - 1000 nM) in a volume of 50 μ L, Cytochalasin B in a volume of 50 μ L (20 μ g/ml) and Ringers buffer in a volume of 50 μ L. These cells are allowed to warm (37 °C, 5% CO₂, 95% RH) for 5 min before IL-8, GRO α , GRO β , GRO γ or NAP-2 at a final concentration of 0.01 - 1000 nM was added. The reaction is allowed to proceed for

3x

60189175-031400

45 min before the 96 well plate is centrifuged (800 xg 5 min.) and 100 ul of the supernatant removed. This supernatant is added to a second 96 well plate followed by an artificial elastase substrate (McOSuc-Ala-Ala-Pro-Val-AMC, Nova Biochem, La Jolla, CA) to a final concentration of 6 ug/ml dissolved in phosphate buffered saline. Immediately, the plate is placed in a fluorescent 96 well plate reader (Cytofluor 2350, Millipore, Bedford, MA) and data collected at 3 min intervals according to the method of Nakajima et al J. Biol. Chem. 254 4027 (1979). The amount of Elastase released from the PMNs is calculated by measuring the rate of McOSuc-Ala-Ala-Pro-Val-AMC degradation.

10

TNF- α in Traumatic Brain Injury Assay

The present assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions, which follow experimentally, induced lateral fluid-percussion traumatic brain injury (TBI) in rats. Adult Sprague-Dawley rats (n=42) were anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and subjected to lateral fluid-percussion brain injury of moderate severity (2.4 atm.) centered over the left temporoparietal cortex (n=18), or "sham" treatment (anesthesia and surgery without injury, n=18). Animals are sacrificed by decapitation at 1, 6 and 24 hr. post injury, brains removed, and tissue samples of left (injured) parietal cortex (LC), corresponding area in the contralateral right cortex (RC), cortex adjacent to injured parietal cortex (LA), corresponding adjacent area in the right cortex (RA), left hippocampus (LH) and right hippocampus (RH) are prepared. Total RNA are isolated and Northern blot hybridization is performed and quantitated relative to an TNF- α positive control RNA (macrophage = 100%). A marked increase of TNF- α mRNA expression is observed in LH (104 $\pm$ 17% of positive control, p < 0.05 compared with sham), LC (105 $\pm$ 21%, p < 0.05) and LA (69 $\pm$ 8%, p < 0.01) in the traumatized hemisphere 1 hr. following injury. An increased TNF- α mRNA expression is also observed in LH (46 $\pm$ 8%, p < 0.05), LC (30 $\pm$ 3%, p < 0.01) and LA (32 $\pm$ 3%, p < 0.01) at 6 hr which resolves by 24 hr following injury. In the contralateral hemisphere, expression of TNF- α mRNA is increased in RH (46 $\pm$ 2%, p < 0.01), RC (4 $\pm$ 3%) and RA (22 $\pm$ 8%) at 1 hr and in RH (28 $\pm$ 11%), RC (7 $\pm$ 5%) and RA (26 $\pm$ 6%, p < 0.05) at 6 hr but not at 24 hr following injury. In sham (surgery without injury) or naive animals, no consistent changes in expression of TNF- α mRNA are observed in any

101
97

of the 6 brain areas in either hemisphere at any times. These results indicate that following parasagittal fluid-percussion brain injury, the temporal expression of TNF- α mRNA is altered in specific brain regions, including those of the non-traumatized hemisphere. Since TNF- α is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated astrocytes, this post-traumatic alteration in gene expression of TNF- α plays an important role in both the acute and regenerative response to CNS trauma.

CNS Injury model for IL-1 β mRNA

This assay characterizes the regional expression of interleukin-1 β (IL-1 β) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Adult Sprague-Dawley rats (n=42) are anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and subjected to lateral fluid-percussion brain injury of moderate severity (2.4 atm.) centered over the left temporoparietal cortex (n=18), or "sham" treatment (anesthesia and surgery without injury). Animals are sacrificed at 1, 6 and 24 hr. post injury, brains removed, and tissue samples of left (injured) parietal cortex (LC), corresponding area in the contralateral right cortex (RC), cortex adjacent to injured parietal cortex (LA), corresponding adjacent area in the right cortex (RA), left hippocampus (LH) and right hippocampus (RH) are prepared. Total RNA is isolated and Northern blot hybridization was performed and the quantity of brain tissue IL-1 β mRNA is presented as percent relative radioactivity of IL-1 β positive macrophage RNA which was loaded on the same gel. At 1 hr following brain injury, a marked and significant increase in expression of IL-1 β mRNA is observed in LC (20.0 $\pm$ 0.7% of positive control, n=6, p < 0.05 compared with sham animal), LH (24.5 $\pm$ 0.9%, p < 0.05) and LA (21.5 $\pm$ 3.1%, p < 0.05) in the injured hemisphere, which remained elevated up to 6 hr. post injury in the LC (4.0 $\pm$ 0.4%, n=6, p < 0.05) and LH (5.0 $\pm$ 1.3%, p < 0.05). In sham or naive animals, no expression of IL-1 β mRNA is observed in any of the respective brain areas. These results indicate that following TBI, the temporal expression of IL-1 β mRNA is regionally stimulated in specific brain regions. These regional changes in cytokines, such as IL-1 β play a role in the post-traumatic.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual

39

60189175-031400

P51111P

publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore the Examples herein are to be construed as merely illustrative and not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.


60189125.031400

103
99

What is Claimed Is:

1. A compound of the formula (I):

5



wherein

R_b is independently hydrogen, NR_6R_7 , OH, OR_a , C_{1-5} alkyl, aryl, aryl C_{1-4} alkyl, aryl

- 10 C_{2-4} alkenyl; cycloalkyl, cycloalkyl C_{1-5} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{2-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, or a heterocyclic C_{2-4} alkenyl moiety, all of which moieties may be optionally substituted one to three times independently by halogen; nitro; halosubstituted C_{1-4} alkyl; C_{1-4} alkyl; amino, mono or di- C_{1-4} alkyl substituted amine; OR_a ; $C(O)R_a$; $NR_aC(O)OR_a$;
 15 $OC(O)NR_6R_7$; hydroxy; $NR_9C(O)R_a$; $S(O)_mR_a$; $C(O)NR_6R_7$; $C(O)OH$; $C(O)OR_a$; $S(O)_2NR_6R_7$; $NHS(O)_2R_a$. Alternatively, the two R_b substituents can join to form a 3-10 membered ring, optionally substituted and containing, in addition to carbon, independently, 1 to 3 NR_a , O, S, SO, or SO_2 moieties which can be optionally unsaturated;

- 20 R_a is an alkyl, aryl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclic, $COOR_a$, or a heterocyclic C_{1-4} alkyl moiety, all of which moieties may be optionally substituted;

m is an integer having a value of 1 to 3;

m' is 0, or an integer having a value of 1 or 2;

- 25 n is an integer having a value of 1 to 5;

q is 0, or an integer having a value of 1 to 10;

t is 0, or an integer having a value of 1 or 2;

41

104
100

s is an integer having a value of 1 to 3;

R₁ is independently selected from hydrogen, halogen, nitro, cyano, C₁₋₁₀ alkyl, halosubstituted C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxy, halosubstituted C₁₋₁₀ alkoxy, azide, S(O)₁R₄, (CR₈R₉)_q S(O)₁R₄, hydroxy, hydroxy substituted C₁₋₄alkyl, aryl, aryl C₁₋₄ alkyl, aryl C₂₋₁₀ alkenyl, aryloxy, aryl C₁₋₄ alkyloxy, heteroaryl, heteroarylalkyl, heteroaryl C₂₋₁₀ alkenyl, heteroaryl C₁₋₄ alkyloxy, heterocyclic, heterocyclic C₁₋₄alkyl, heterocyclic C₁₋₄alkyloxy, heterocyclic C₂₋₁₀ alkenyl, (CR₈R₉)_q NR₄R₅, (CR₈R₉)_q C(O)NR₄R₅, C₂₋₁₀ alkenyl C(O)NR₄R₅, (CR₈R₉)_q C(O)NR₄R₁₀, S(O)₂R₈, (CR₈R₉)_q C(O)R₁₁, C₂₋₁₀ alkenyl C(O)R₁₁, C₂₋₁₀ alkenyl C(O)OR₁₁, (CR₈R₉)_q C(O)OR₁₁, (CR₈R₉)_q OC(O)R₁₁, (CR₈R₉)_q NR₄C(O)R₁₁, (CR₈R₉)_q C(NR₄)NR₄R₅, (CR₈R₉)_q NR₄C(NR₅)R₁₁, (CR₈R₉)_q NHS(O)₂R₈, (CR₈R₉)_q S(O)₂NR₄R₅, or two R₁ moieties together may form O-(CH₂)₈O or a 5 to 6 membered saturated or unsaturated ring, and wherein the alkyl, aryl, arylalkyl, heteroaryl, heterocyclic moieties may be optionally substituted;

R₄ and R₅ are independently hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl C₁₋₄alkyl, heterocyclic, heterocyclic C₁₋₄ alkyl, or R₄ and R₅ together with the nitrogen to which they are attached form a 5 to 7 member ring which may optionally comprise an additional heteroatom selected from O, N and S;

R₆ and R₇ are independently hydrogen, or a C₁₋₄ alkyl, heteroaryl, aryl, alkyl aryl, alkyl C₁₋₄ heteroalkyl or R₆ and R₇ together with the nitrogen to which they are attached form a 5 to 7 member ring which ring may optionally contain an additional heteroatom is selected from oxygen, nitrogen or sulfur, and which ring may be optionally substituted;

R₈ is hydrogen or C₁₋₄ alkyl;

R₉ is a C₁₋₄ alkyl;

R₁₀ is C₁₋₁₀ alkyl C(O)₂R₈;

R₁₁ is hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl,

optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally

42

60189175-031400

105
101

substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl; and

R₁₃ and R₁₄ are independently hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted C₁₋₄ alkynyl, optionally substituted C₁₋₄ alkenyl optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄ alkyl, optionally substituted heterocyclic, optionally substituted heterocyclicC₁₋₄alkyl, halogen, nitro, OR_a, NR₆R₇, C(O)R_a, NR₁₁C(O)OR_a, OC(O)NR₆R₇, NR₁₁C(O)R_a, S(O)_mR_a, C(O)NR₆R₇, C(O)OH, C(O)OR_a, S(O)₂NR₆R₇, or NR₁₁S(O)₂R_a.

R₂₀ is hydrogen or alkyl, alkynyl, alkenyl, heterocyclic, heteroaromatic all of which maybe optionally substituted or a pharmaceutically acceptable salt thereof.

2. The compound according to Claim 1 wherein R₁ is substituted in the 4- position by an electron withdrawing moiety.

3. The compound according to Claim 2 wherein R₁ is halogen, cyano or nitro.

4. The compound according to Claim 3 wherein R₁ is halogen.

5. The compound according to Claim 4 wherein R₁ is independently, fluorine, chlorine, or bromine.

6. The compound according to Claim 1 wherein R_b is hydrogen, C₁₋₄ alkyl, or C₁₋₄ alkyl substituted with C(O)OH, or C(O)OR_a.

7. The compound according to Claim 1 which is selected from the group consisting of:

N-allyl-N'-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)urea;

60189175-031400

106
162

N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-isopropylurea;
 N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-ethylurea;
 N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-propylurea; and
 N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-(ethoxycarbonyl)methylurea.

5

8. A compounds according to claim 7 which are

N-allyl-N'-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)urea;
 N-(3-aminosulfonyl-4-chloro-2-hydroxyphenyl)-N'-isopropylurea;

10 9. A compound according to claim 7 wherein the compound is in its sodium salt form.

10. A compound according to claim 7 wherein the compound is in its potassium salt form.

15 11. A pharmaceutical composition comprising a compound according to any of Claims 1 to 10 and a pharmaceutically acceptable carrier or diluent.

12. A method of treating a chemokine mediated disease, wherein the chemokine binds to an IL-8 a or b receptor in a mammal, which method comprises administering to
 20 said mammal an effective amount of a compound of the formula according to any one of Claims 1 to 14.

13. The method according to Claim 12 wherein the mammal is afflicted with a chemokine mediated disease selected from the group consisting of:
 25 psoriasis, atopic dermatitis, osteo arthritis, rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, adult respiratory distress syndrome, inflammatory bowel disease, Crohn's disease, ulcerative colitis, stroke, septic shock, multiple sclerosis, endotoxic shock, gram negative sepsis, toxic shock syndrome, cardiac and renal reperfusion injury, glomerulonephritis, thrombosis, graft vs. host reaction,
 30 Alzheimer's disease, allograft rejections, malaria, restenosis, angiogenesis, atherosclerosis, osteoporosis, gingivitis and undesired hematopoietic stem cells release and diseases caused by respiratory viruses, herpes viruses, and hepatitis

448

50189175-031400

107
163

viruses, meningitis, cystic fibrosis, pre-term labor, cough, pruritus, multi-organ
dysfunction, trauma, strains, sprains, contusions, psoriatic arthritis, herpes,
encephalitis, CNS vasculitis, traumatic brain injury, CNS tumors, subarachnoid
hemorrhage, post surgical trauma, interstitial pneumonitis, hypersensitivity, crystal
5 induced arthritis, acute and chronic pancreatitis, acute alcoholic hepatitis,
necrotizing enterocolitis, chronic sinusitis, uveitis, polymyositis, vasculitis, acne,
gastric and duodenal ulcers, celiac disease, esophagitis, glossitis, airflow obstruction,
airway hyperresponsiveness, bronchiolitis obliterans organizing pneumonia,
bronchiectasis, bronchiolitis, bronchiolitis obliterans, chronic bronchitis, cor
10 pulmonae, dyspnea, emphysema, hypercapnea, hyperinflation, hypoxemia,
hyperoxia-induced inflammations, hypoxia, surgical lung volume reduction,
pulmonary fibrosis, pulmonary hypertension, right ventricular hypertrophy,
sarcoidosis, small airway disease, ventilation-perfusion mismatching, wheeze, colds
and lupus.

60189175-031400

+08
KX

ABSTRACT

This invention relates to novel compounds of Formula (I) to (VII), and compositions thereof, useful in the treatment of disease states mediated by the chemokine, Interleukin-8 (IL-8).

60189175.031400

46

Señores

DIVISION DE NUEVAS CREACIONES

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO

Bogotá, D.C.

SUPERINDUSTRIA Y COMERCIO
Radicación : 01022200 00000001 Folios:
Fecha (Año): 2001-03-30 12:50:15
Trámite : 002 PATENTES D 1 REGISTRO/ 329 COMPLEMEN
Dependencia: 2020 DIVISION DE NUEVAS CREACIONES

Re.: Expediente No. 01 - 20.200

Solicitud de Patente

Debidamente traducida, adjunto acompaño la copia oficial de la solicitud de patente básica estadounidense No. 60/189.175 la cual le solicito agregar a este expediente para los efectos legales respectivos.

De Ustedes atentamente,

GERMAN CASTILLO GRAU

T.P. No. 2978 del C.S.J.