

**SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
DIRECCIÓN DE NUEVAS CREACIONES**

2020
Bogotá, D.C.,

Abogado
JUAN PABLO CADENA SARMIENTO
jcadena@bc.com.co

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO



No. 09-149835- -00005-0000

Fecha: 2012-08-23 16:39:26 Dep. 2020 DIR.NUEVASCR
Tra. 11 PATENTEIYII Eve: 378 FASENACIONALI
Act. 519 TRASLADOINFORME Folios: 1

ASUNTO	Radicación	09 149 835
	Trámite	011
	Evento	378
	Actuación	519
	Oficio	
	Folios	006

15167

Patente de Invención



Patente de Modelo de Utilidad



Vía Nacional

Vía PCT (Fase Nacional)



Capítulo I



Capítulo II



No. Solicitud Internacional

PCT/GB08/02194

Fecha de Presentación Internacional

24/Junio/2008

Como resultado del Examen de Fondo realizado, con fundamento en el Artículo 45 de la Decisión 486 de la Comisión de la Comunidad Andina, se comunica:

1. Documentos estudiados

Descripción:	Folios 7 a 51	31/Diciembre/2009
Reivindicaciones:	Folios 52 a 54	31/Diciembre/2009
Prioridad:	PCT/GB2007/002390	26/Junio/2007

2. Análisis de la Prioridad

Se acepta la reivindicación de Prioridad porque esta solicitud fue presentada dentro del plazo establecido (Art. 9 D 486), y su contenido técnico corresponde con el de la prioridad reivindicada.

3. Objeto de la invención

Título de la solicitud: "DERIVADOS DE TIAZOL FUSIONADOS COMO INHIBIDORES DE QUINASA".

El problema planteado en esta solicitud consiste en la necesidad de suministrar compuestos útiles como inhibidores selectivos de PI3 quinasa.

Para resolver este problema técnico la solicitud en estudio proporciona un compuesto de la fórmula (I) sus sales y composiciones que lo contienen.

El objeto de la presente Solicitud se relaciona con la Clasificación Internacional de Patentes (CIP): C07D 513/04, A61K 31/437, C07D 519/00.

Expediente 09 149 835 1er Art 45

4. Excepción a la Patentabilidad (Art 20 D 486 literal (d))

El objeto de la reivindicación 15 no es patentable de acuerdo con el Art citado.

5. Reivindicaciones relativas a un uso o a un segundo uso (Art 14 y 21 D 486)

El uso de las reivindicaciones 12 – 14 no es patentable, de acuerdo con el Art 14.

6. De la Solicitud de Patente

Reivindicaciones (Art 30 D 486, Art 22 PCT)

La reivindicación 1 y sus dependientes 2 a 9 y 11, reclaman también los solvatos de los cuales no existe soporte en la descripción.

En las Reivindicaciones 1 y sus dependientes, se sugiere una razonable generalización del tipo de compuestos que ha sido sintetizado ó probado, con la actividad biológica que se les atribuye, teniendo en cuenta que sólo están soportadas la sustituciones de R11 y R12 por un grupo metilo, R24 por H, V y W son siempre un átomo de C (según los ejemplos de la descripción).

La reivindicación 10 no esta caracterizada por esta correctamente caracterizada.

7. Determinación del Estado de la Técnica

El Estado de la Técnica se determinó con base en la fecha de presentación de la Solicitud de Prioridad: Junio 26 de 2007 y se encontraron los siguientes documentos que anticipan el objeto de esta solicitud:

Nº	Documento	Título	Fecha de Publicación*
D1	WO2006114606	Fused Thiazole derivatives as kinase inhibitors	Noviembre 2 del 2006
D2	Libro	Foye's principles of Medicinal Chemistry	2002

D1¹, enseña el compuesto derivado de tiazol fusionado con actividad como inhibidores de quinasa de fórmula (I), sus sale y composiciones que lo contienen.

D2² libro se refiere al bioisosterismo como herramienta para la modificación de compuestos.

8. Examen de Patentabilidad (Art 14 D486)






Nivel Inventivo (Art18 D486)

La reivindicación 1 consiste en un compuesto de fórmula (I), o una sal o un solvato farmacéuticamente aceptable del mismo.

1 http://worldwide.espacenet.com/publicationDetails/biblio?DB=EPODOC&II=0&ND=3&adjacent=true&locale=en_EP&FT=D&date=20061102&CC=WO&NR=2006114606A1&KC=A1

2 Foye's principles of Medicinal Chemistry sixth Edition Thomas L. LEMKE, M David A. WILLIAMS, Victoria F. ROCHE, S. William Zito; Classical Bioisosteres, Pág. 50, 2002, (págs. Portada y 49, Fls. 111 y 112).

Comparación de las características técnicas **esenciales**, con las del Estado de la Técnica:

Características esenciales	D1
 Fórmula I	 Fórmula IIE
 C(O) N-H R11 y R12 son: -CH3	 X1= C(O), C-H R21 y R22 son: alquil C₁₋₆
 T=N-alquilC1-6 W=CH V=CH R23=hidrógeno, halógeno, ciano, alquilo C1-6, hidroxialquilo C1-6, trifluorometilo, arilalquiloC1-6, hidroxil, alcoxiC1-6, trifluorometoxi, ariloxi, arilalcoxiC1-6, alquilC1-6tio, alquilC1-6sulfoniloxi, amino, alquilC2-6carbonilamino.	 N-H W=CH CH R43= hidrógeno, halógeno, ciano, alquilo C1-6, hidroxialquilo C1-6, trifluorometilo, arilalquiloC1-6, hidroxil, alcoxiC1-6, trifluorometoxi, ariloxi, arilalcoxiC1-6, alquilC1-6tio, alquilC1-6sulfoniloxi, amino, alquilC2-6carbonilamino.

Los documentos D1 y D2 constituyen el estado de la técnica cercano a la invención porque D1, da a conocer el compuesto de fórmula IIE (pág.31) sus sales, solvatos y enantiómeros (págs.2, 5 y 6), además de sus composiciones farmacéuticas (reivindicación 16, pág.140).

El compuesto, objeto de esta solicitud se diferencia del conocido en D1, en que este incluye un átomo de N en el anillo de ciclohexanona para formar la piridinona.

El hecho de incluir un átomo de N en el anillo de ciclohexanona para formar la piridinona, en el compuesto que se daba a conocer en D1, no causa un efecto sorprendente.

Por tanto, el Problema Técnico Objetivo consiste en ¿cómo modificar el compuesto según D1, para lograr un compuesto alternativo con la misma actividad que el conocido?.

Es importante mencionar que en la descripción de la presente solicitud se observan valores de IC50 para la inhibición de la actividad de PI3K α y/o PI3K β , y/o PI3K γ y/o PI3K δ humanas de 50 μ M o mejores (pág.18, Fl.24), que al compararlos con los resultados de actividad del compuesto de D1, que también son inhibidores de la actividad de PI3K α y/o PI3K β y/o PI3K γ y/o PI3K δ humanas, se comprueba que son exactamente iguales por tener un IC50 de 50 μ M o mejores (enzyme inhibition assays pág.40).

No obstante D2 muestra que es posible cambiar un grupo CH2 por un grupo NH (tabla 2.8, pág.49) con la expectativa de mantener la misma actividad en el Expediente 09 149 835 1er Art 45

compuesto reivindicado.

En consecuencia, la reivindicación es obvia, porque la persona del oficio normalmente versada en la materia enfrentada al problema de "¿cómo modificar el compuesto según D1, para lograr un compuesto alternativo con la misma actividad que el conocido?", incluiría el grupo NH que se revela en D2 en el compuesto que se enseña en D1, logrando de esta forma el compuesto reivindicado.

Por lo cual el compuesto de la Reivindicación 1 carece de Nivel Inventivo.

Puesto que las reivindicaciones dependientes 2 a 9 y 11 no mencionan características que las hagan inventivas, tampoco tienen nivel inventivo.


9. Conclusión

Los requisitos de Patentabilidad de la presente Solicitud están afectados así:

Nº	Documento	Reivindicaciones afectadas	Requisito que afecta
D1	WO2006114606	1-9 y 11	Nivel Inventivo
D2	Libro	1-9 y 11	Nivel Inventivo

En cumplimiento del artículo 45 de la Decisión 486 de la Comisión de la Comunidad Andina, se le indica que debe dar respuesta dentro del término de sesenta (60) días contados a partir de la fecha de notificación de la presente comunicación. En caso de no dar respuesta al presente requerimiento, o si a pesar de la respuesta subsistieran los impedimentos para la concesión, se denegará la patente.

Notifíquese el presente oficio conforme a lo dispuesto en el numeral 5.2, literal e), del capítulo quinto del título primero de la Circular Única No. 10 de 2001.


JOSÉ LUIS SALAZAR LÓPEZ
Director de Nuevas Creaciones (C)

El anterior requerimiento fue notificado por fijación en lista, de fecha


_____ 20 Abr. 2012

Elaborado por: Lyna del Carmen Mordecay Dunoyer
Revisado Por: Gloria Jacqueline Alfonso Rozo
Aprobado por: José Luis Salazar López. 

FOYE'S PRINCIPLES OF

MEDICINAL CHEMISTRY

SIXTH EDITION



Thomas L. LEMKE
David A. WILLIAMS
Victoria F. ROCHE
S. William ZITO

Walters Kluwer | Lippincott
Williams & Wilkins



FOYE'S Principles of Medicinal Chemistry

SIXTH EDITION

Edited By

THOMAS L. LEMKE, PhD

*Professor Emeritus
College of Pharmacy
University of Houston
Houston, Texas*

DAVID A. WILLIAMS, PhD

*Professor Emeritus of Chemistry
Massachusetts College of Pharmacy and
Health Sciences
Boston, Massachusetts*

Assistant Editors

VICTORIA F. ROCHE, PhD

*Senior Associate Dean
School of Pharmacy and Health Professions
Creighton University
Omaha, Nebraska*

S. WILLIAM ZITO, PhD

*Professor of Pharmaceutical Sciences
College of Pharmacy and
Allied Health Professions
St. John's University
Jamaica, New York*



Walters Kluwer | Lippincott Williams & Wilkins

Health
Philadelphia • Baltimore • New York • London
Buenos Aires • Hong Kong • Sydney • Tokyo

Copyrighted material

Apurva Editor: David Troy
Managing Editor: Meredith Britton
Marketing Manager: Christian Murphy
Production Editor: Hearstside Publishing Services
Designer: Sunflower Publishing Services
Graphic Artist: Nicole Williams
Compositor: Aptara

Cover image courtesy of Wovofunction, Inc. The ribbon display (green and red) is tyrosine kinase. The "active site" for tyrosine kinase is shown as tube display with a mesh-style electrostatic potential map (a charge distribution map) with the drug (tamoxifen) shown in the "active site" with a solid-color style electrostatic potential map (where the colors (red and blue) represent negative and positive charges, respectively). Tamoxifen (a 4-phenyl amino pyrimidine derivative, a.k.a. CGP71488, ST1571 or Glivec®) is a selective inhibitor of several tyrosine kinases that binds in the ATP-binding pocket of tyrosine kinase and blocks the activity of Abl, c-kit, and PDGFR. It is used for treating chronic myelogenous leukemia (CML), gastrointestinal stromal tumors (GIST), and a number of other malignancies.

South Edition

Copyright © 2008, 2002, 1995, 1989, 1981, 1974
Lippincott Williams & Wilkins, a Wolters Kluwer business.

351 West Camden Street 530 Walnut Street
Baltimore, MD 21201 Philadelphia, PA 19106

Printed in the USA

All rights reserved. This book is protected by copyright. No part of this book may be reproduced or transmitted in any form or by any means, including as photocopies or scanned in or other electronic copies, or utilized by any information storage and retrieval system without written permission from the copyright owner, except for brief quotations embodied in critical articles and reviews. Materials appearing in this book prepared by individuals as part of their official duties as U.S. government employees are not covered by the above-mentioned copyright. To request permission, please contact Lippincott Williams & Wilkins at 530 Walnut Street, Philadelphia, PA 19106, via email at permissions@lww.com, or via website at lww.com (products and services).

9 8 7 6 5 4 3 2 1

Library of Congress Cataloging-in-Publication Data

Foye's principles of medicinal chemistry / edited by Thomas L. Lenzke. . . [et al]. — 6th ed.

p. : cm.
Includes bibliographical references and index.

ISBN 07817-6879-5

I. Pharmaceutical chemistry. I. Foye, William O. II. Lenzke, Thomas L.

III. Title: Principles of medicinal chemistry.

[DNLM: 1. Chemistry, Pharmaceutical. QV 744 F7962 2008]

RS403 P73 2008

616.0756—dc22

2007036236

DISCLAIMER

Great care has been taken to confirm the accuracy of the information presented and to describe generally accepted practices. However, the authors, editors, and publisher are not responsible for errors or for any consequences arising from the application of the information in this book and make no warranty, expressed or implied, with respect to the accuracy, completeness, or accuracy of the contents of the publication. Application of the information in a particular situation remains the professional responsibility of the practitioner. The clinical decisions described and recommended may not be considered suitable and universal recommendations.

The authors, editors, and publisher have exerted every effort to ensure that drug selection and dosage in this text are in accordance with the current recommendations and practice at the time of publication. However, in view of rapid drug research, changes in government regulations, and the constant flow of information relating to drug efficacy and drug reactions, the reader is urged to check the package insert for each drug for any change in indications and dosage and for added cautions and precautions. This is particularly important when the recommended agent is a new or infrequently employed drug.

Some drugs and medical devices presented in this publication have Food and Drug Administration (FDA) clearance for limited use in controlled research settings. It is the responsibility of the health care provider to ascertain the FDA status of each drug or device proposed for use in their clinical practice.

To purchase additional copies of this book, call our customer service department at (800) 638-3030 or fax orders to (301) 223-2330. International customers should call (301) 223-2300.

Visit Lippincott Williams & Wilkins on the Internet: <http://www.lww.com>. Lippincott Williams & Wilkins customer service representatives are available from 8:30 am to 6:00 pm, EST.

Table 2.8. Grimm's Hydride Displacement "Law"

C	N	O	F	Na
	CH	NH	OH	PH
		CH	NH	OH
			CH	NH

valence electrons for each group of atoms within a column. Within each column, the groups are considered to be "pseudoatoms" with respect to one another. Thus, NH_2 is considered to be isosteric to CH_3 and so on. This early view of isosterism did not consider the actual location, motion, and resonance of electrons within the orbitals of these functional group replacements. Careful observation of this table reveals that some groups do share similar physicochemical properties, but others have very different properties, despite having the same number of valence electrons. For example, OH and NH_2 share similar hydrogen-bonding properties and, therefore, should be interchangeable if that is the only criterion necessary. The NH_2 group is basic, however, whereas the OH is neutral. Hence, at physiological pH, the NH_2 group would be in its protonated, or conjugate acid, form and give the molecule a positive charge. If OH is being substituted by NH_2 , the additional positive charge could have a significant effect on the overall physicochemical properties of the molecule in which it is being introduced. The difference in physicochemical properties of the CH_3 group relative to the OH and NH_2 groups is even greater. In addition to basicity and acidity, this "law" fails to take into account other important physicochemical parameters, such as electronegativity, polarizability, bond angles, size, shape of molecular orbitals, electron density, and partition coefficients, all of which contribute significantly to the overall physicochemical properties of a molecule.

Instead of considering only partial structures, Hinsberg (54) applied the concept of isosterism to entire molecules. He developed the concept of "ring equivalents"—that is, groups that can be exchanged for one another in aromatic ring systems without drastic changes in physicochemical properties relative to the parent structure. Benzene, thiophene, and pyridine illustrate this concept (Fig. 2.33). A $-\text{CH}=\text{CH}-$ group in benzene is replaced by the divalent sulfur, S , in thiophene, and a $-\text{CH}=\text{CH}-$ is replaced by the trivalent $-\text{N}-$ to give pyridine. The physical properties of benzene and thiophene

are very similar. For example, the boiling point of benzene is 81.1°C , and that of thiophene is 84.4°C (at 760 mm Hg). Pyridine, however, deviates, with a boiling point of 115 to 116°C . Hinsberg therefore concluded that divalent sulfur (S - or thioether) must resemble $-\text{C}=\text{C}-$ in shape, and these groups were considered to be isosteric. Note that hydrogen atoms are ignored in this comparison. Today, this isosteric relationship is seen in many drugs (e.g., H_2 -receptor antagonists) (Fig. 2.33).

Bioisosterism

It is difficult to relate biological properties to physicochemical properties of individual atoms, functional groups, or entire molecules, because many physicochemical parameters are involved simultaneously and, therefore, are difficult to quantitate. Simple relationships as described above often do not hold up across the many types of biological systems seen with medicinal agents. That is, what may work as an isosteric replacement in one biological system (for a given drug receptor, enzyme, etc.) may not work in another system. Because of this, it was necessary to introduce the term "bioisosterism" to describe functional groups related in structure and having similar biological effects. Friedman (55) introduced the term bioisosterism and defined it as follows: "Bioisosteres are (functional) groups or molecules that have chemical and physical similarities producing broadly similar biological properties." Recently, Burger (56) expanded this definition to take into account biochemical views of biological activity: "Bioisosteres are compounds or groups that possess near equal molecular shapes and volumes, approximately the same distribution of electrons, and which exhibit similar physical properties such as hydrophobicity. Bioisosteric compounds affect the same biochemical associated systems as agonists or antagonists and thereby produce biological properties that are related to each other." The key point is that the same pharmacologic target is influenced by bioisosteres as agonists or antagonists. What may work as a bioisosteric group in one biological system may not have similar effects on another.

Classical and Nonclassical Bioisosteres

Bioisosteric groups can be subdivided into two categories: classical and nonclassical bioisosteres. Functional groups that satisfy the original conditions of Langmuir and Grimm are referred to as classical bioisosteres. Nonclassical bioisosteres do not obey steric and electronic definitions of classical bioisosteres and do not necessarily have the same number of atoms as the substituent they replace. A wider set of compounds and functional groups are encompassed by nonclassical bioisosteres that produce, at the molecular level, qualitatively similar agonistic or antagonistic responses. In animals, many hormones, neurotransmitters, and so on with very similar structures and biological actions can be classified as bioisosteres. An example would be the insulins isolated from various mammalian species. Even though these insulins may differ by several amino acid residues, they




Fig. 2.33. Isosteric substitution of thiophene for benzene and benzene for pyridine.

122