

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO			
RAD:	12-41581- -8	FECHA:	2014-08-13 11:13:27
DEP:	2020 DIRECCION DE NUEVAS CREACIONES	EVE:	379 FASE NACIONAL INCLUIDO CAPITULO II
TRA:	11 PCT-PATENTE DE INVENCION CAPITULOS I Y II	FOLIOS:	13
ACT:	519 PRESENTACION		

DIRECCIÓN DE NUEVAS CREACIONES

2020
Bogotá, D.C.,

Abogado(a)/Señor(a)
LUIS FELIPE CASTILLO GIBSON

Asunto: Radicación: 12-41581- -8
 Trámite: 11
 Evento: 379
 Actuación: 519
 Oficio: 9982
 Folios: 13

Patente de Invención	X			Patente de Modelo de Utilidad	
Vía Nacional					
Vía PCT (Fase Nacional)	X	Capítulo I		Capítulo II	X
No. Solicitud Internacional	PCT/KR2010/005169				
Fecha de Presentación Internacional	06/08/2010				

Como resultado del examen de fondo realizado, con fundamento en el Artículo 45 de la Decisión 486 de la Comisión de la Comunidad Andina, se comunica:

1. Documentos estudiados

Descripción:	No. 12-041581-00000-0000	09/Marzo/2012
Reivindicaciones:	No. 12-041581-00000-0000	09/Marzo/2012
	No. 12-041581-00007-0000	17/Enero/2014
Dibujos:	No. 12-041581-00000-0000	09/Marzo/2012
Secuencias:	No. 12-041581-00000-0000	09/Marzo/2012
Prioridad:	10-2009-0075404	14/Agosto/2009
	10-2010-0075514	05/Agosto/2010

2. Análisis de la Prioridad

Se aceptan las reivindicaciones de prioridad porque ésta solicitud fue presentada dentro del plazo establecido (Art. 9 D 486) y su contenido técnico corresponde con el de las prioridades reivindicadas.

3. Objeto de la invención

De acuerdo con lo establecido en el numeral 1.2.2.2 del capítulo primero del Título 10 de la Circular Única para efectos del examen de patentabilidad, se tendrá en cuenta la

tasa pagada por el solicitante al momento de comenzar el examen para las reivindicaciones adicionales; esto es que cuando la solicitud contenga más de diez (10) reivindicaciones y no se haya pagado la tasa complementaria, sólo se estudiarán las cubiertas por la tasa pagada. En el presente caso se estudiará la totalidad del capítulo No. 12-041581-00007-0000 (reivindicaciones 1-4).

Título de la solicitud: “Método para producir ostras tetraploides viables”.

El problema planteado en esta solicitud consiste en la necesidad de proporcionar un método de producción de ostras tetraploides viables con alta eficiencia.

Para resolver este problema técnico la solicitud en estudio proporciona un método para producir ostras tetraploides viables usando una combinación de citocalasina B y 6-dimetilaminopurina.

El objeto de la presente solicitud se relaciona con la Clasificación Internacional de Patentes (CIP): A01K 61/00.

4. De la solicitud de patente

Reivindicaciones (Art 30 D 486, Art 22 PCT)

- 1– En la reivindicación 1 no es clara debido a que no se mencionan todas las características esenciales del método. Se sugiere incluir los parámetros de temperatura, salinidad y tiempo de las reivindicaciones 2-4 en la reivindicación independiente 1.
- 2– Por otra parte, la reivindicación 1 no se ajusta a la estructura: PREAMBULO-ENLAZANTE-PARTE CARACTERÍSTICA por cuanto incluye dos tipos de enlazantes, el primero: “*que comprende las etapas de*” y el segundo “*que se caracteriza porque*”. Al presentar la reivindicación 1 de esta manera no es posible establecer con total certeza cuál es el aporte al estado de la técnica, es decir si corresponde a la etapa de: “*fertilizar los huevos de ostras tetraploides hembras con espermatozoides de ostras diploides machos*”, lo cual es un proceso esencialmente biológico ya que de acuerdo con la divulgación esta etapa consiste en poner en agua de mar espermatozoides y oocitos a fecundar, en donde el reconocimiento oocito-espermatozoide ocurre naturalmente a nivel intra-específico, o bien a la etapa de: “*impedir que los primeros cuerpos polares sean eyectados de los huevos feertilizados que incluye poner en contacto los huevos fertilizados con citocalasina B y luego poner en contacto los huevos fertilizados con 6-dimetilmercaptapurina (DMAP)*” que es materia elegible para estudio.

Se requiere al solicitante hacer las aclaraciones necesarias.

5. Determinación del Estado de la Técnica

El estado de la técnica se determinó con base en la fecha de presentación de la solicitud de prioridad: Agosto 14 de 2009, y se encontraron los siguientes documentos que anticipan el objeto de esta solicitud:

Ítem	Documento	Título	Fecha de Publicación
D1	WO9519703	“Tetraploid Shellfish”	27/Junio/1995
D2	Yi and Park, 2004. Animal Reproduction Science 86: 297–304	“Parthenogenetic development of porcine oocytes treated by ethanol, cycloheximide, cytochalasin B and 6-dimethylaminopurine”	20/Julio/2004

El documento D1¹ divulga moluscos tetraploides, y un método novedoso para su producción (Resumen).

El documento D2 divulga un método para la producción de oocitos porcinos que incluye poner en contacto citocalasina B y 6-dimetilaminopurina (Pág. 299).

Nivel inventivo (Art18 D486)

La reivindicación 1 consiste en *“un método para producir ostras tetraploides que comprende (...)”*

Comparación de las características técnicas esenciales, con las del estado de la técnica:

Características esenciales	D1	D2
Método para producir ostras	Método para producir ostras (Pág. 3, Párr. 1)	Método para producir oocitos porcinos (Resument)
Impedir que los primeros cuerpos polares sean expulsados de los huevos fertilizados (incluye poner en contacto citocalasina B con 6-dimetilaminopurina)	Impedir que los primeros cuerpos polares sean expulsados de los huevos fertilizados (incluye poner en contacto citocalasina B ó 6-dimetilaminopurina); (Pág. 6, Párr. 2)	Activar los oocitos poniendo en contacto citocalasina B y 6-dimetilaminopurina (Pág. 299)
Producir ostras tetraploides viables	Producir ostras tetraploides viables (Resumen)	No menciona

El documento D1 se considera el estado de la técnica cercano a la invención definida en la reivindicación 1 porque divulga moluscos tetraploides, y un método novedoso para su producción (Resumen).

La diferencia entre la invención, definida en la reivindicación 1 y el procedimiento divulgado en el documento D1 consiste en la utilización de citocalasina B y 6-dimetilaminopurina al mismo tiempo para impedir que los cuerpos polares sean expulsados.

El efecto que logra el incluir el poner en contacto citocalasina B y 6-dimetilaminopurina al mismo tiempo es que impide la expulsión de los primeros cuerpos polares, aumentando la viabilidad de las ostras tetraploides.

Por lo tanto, el problema técnico objetivo que pretende resolver esta invención se puede formular así: “cómo modificar la manera de impedir que los primeros cuerpos polares sean expulsados para aumentar la viabilidad de las ostras tetraploides y hacer el proceso más eficiente”.

Sin embargo, la utilización de citocalasina B ó 6-dimetilaminopurina para inhibir la expulsión de los cuerpos polares, ya está divulgada en el documento D1 que se refiere a un procedimiento para la producción de ostras tetraploides viables usando citocalasina B ó 6-dimetilaminopurina para impedir que los primeros cuerpos polares sean expulsados (Pág. 6, Párr. 2) y en el documento D2 que se refiere a un procedimiento para producir oocitos viables porcinos (Pág. 299) en donde se pone en contacto al mismo tiempo citocalasina B y 6-dimetilaminopurina (6-DMAP).

En consecuencia, la persona del oficio normalmente versada en la materia, incluiría el poner en contacto la citocalasina B y 6-dimetilaminopurina de acuerdo al procedimiento del documento D2 en el procedimiento divulgado por el documento D1, al conocer el efecto que se produce al poner en contacto los dos compuestos en el medio de cultivo, específicamente lograr la inhibición de la expulsión de los cuerpos polares para llegar

¹.[http://worldwide.espacenet.com/publicationDetails/biblio?](http://worldwide.espacenet.com/publicationDetails/biblio?CC=WO&NR=9519703A1&KC=A1&FT=D&ND=3&date=19950727&DB=EPODOC&locale=en_EP)
CC=WO&NR=9519703A1&KC=A1&FT=D&ND=3&date=19950727&DB=EPODOC&locale=en_EP

así al objeto de la reivindicación 1 de la solicitud. Por lo cual se considera obvia.

Las reivindicaciones dependientes 2-4 no cumplen con el requisito de nivel inventivo (Art. 18).

6. Conclusión

Los requisitos de Patentabilidad de la presente solicitud están afectados así:

Ítem	Documento	Reivindicaciones afectadas	Requisito que afecta
D1	WO9519703	1-4	Nivel Inventivo
D2	Yi and Park, 2004. Animal Reproduction Science 86: 297–304	1-4	Nivel Inventivo

7. Respuesta a los argumentos del señor solicitante

Argumentos con respecto al estudio de nivel inventivo

El señor solicitante argumenta que cuando se ponen en contacto los huevos fertilizados con citocalasina B y luego con 6-dimetilaminopurina (DMAP), se obtiene un efecto sinérgico en comparación con las enseñanzas del documento D1. Al respecto es pertinente señalar que en la tabla 1 existen diferencias significativas entre usar los compuestos solos o en combinación, sin embargo, la combinación y su aplicación fue anticipada por el documento D1, y por otra parte a partir de los hallazgos divulgados en la memoria descriptiva frente a usar un compuesto después del otro, es evidente que no hay diferencias significativas. Adicionalmente, en la tabla 1 el número de oocitos usados en los grupos experimentales 1 y 2 comparados con el grupo experimental 3 y 4 es diferente por lo cual no se puede establecer una diferencia significativa.

En cumplimiento del artículo 45 de la Decisión 486 de la Comisión de la Comunidad Andina, se le indica que debe dar respuesta dentro del término de sesenta (60) días contados a partir del día siguiente de la fecha de notificación de la presente comunicación. En caso de no dar respuesta al presente requerimiento, o si a pesar de la respuesta subsistieran los impedimentos para la concesión, se denegará la patente.

Se invita al solicitante se sirva indicar, si así lo considera, su renuncia al término faltante de sesenta (60) días para dar respuesta a este requerimiento y presentarla por escrito.

Finalmente, se le recuerda al solicitante que si como respuesta a este requerimiento llegara a modificar el capítulo reivindicatorio y éste contenga más de diez (10) reivindicaciones, no pagará la tasa de modificación voluntaria pero sí la correspondiente al pago por reivindicación adicional.

Notifíquese el presente oficio conforme a lo dispuesto en el numeral 5.2, literal e), del capítulo quinto del título primero de la Circular Única No. 10 de 2001.



JOSÉ LUIS SALAZAR LÓPEZ
Director de Nuevas Creaciones (c)

El anterior requerimiento fue notificado por fijación en lista, de fecha 2014-08-19

Elaboró: Lina Maria Lizaraso Trespalacios
Revisó: Consuelo Leguizamon Leguizamon



ELSEVIER

Animal Reproduction Science 86 (2005) 297–304

ANIMAL
REPRODUCTION
SCIENCE

www.elsevier.com/locate/anireprosci

Parthenogenetic development of porcine oocytes treated by ethanol, cycloheximide, cytochalasin B and 6-dimethylaminopurine

Y.J. Yi, C.S. Park*

*Division of Animal Science and Resources, Research Center for Transgenic Cloned Pigs,
Chungnam National University, 305-764 Daejeon, Korea*

Received 2 February 2004; received in revised form 6 July 2004; accepted 20 July 2004

Abstract

This study was carried out to investigate the various concentrations and exposure times of ethanol, one of many intracellular calcium elevating agents, and a sequential combination of ethanol (8%), cycloheximide (CHX, 10 μ g/ml), cytochalasin B (CCB, 7.5 μ g/ml) and 6-dimethylaminopurine (6-DMAP, 2 mM) to improve parthenogenetic activation and development of in vitro matured porcine oocytes. Cumulus-oocyte complexes (COCs) were matured in tissue culture medium (TCM) 199 for 44 h at 38.5 °C, 5% CO₂ in air. Cumulus-free oocytes showing first polar body were activated by concentrations of 0, 5, 6, 7, 8, 9 and 10% ethanol for 10 min and exposure times of 0, 5, 8, 10, 12 and 15 min with 8% ethanol in HEPES buffered (25 mM) NCSU-23 medium. Also, oocytes were activated with the NCSU-23 medium containing 8% ethanol for 10 min. After that, oocytes were incubated in the NCSU-23 medium supplemented with CHX, CCB, 6-DMAP, CHX + CCB, CHX + 6-DMAP, CCB + 6-DMAP and CHX + CCB + 6-DMAP for 3 h, respectively. Following activation, oocytes were transferred into the NCSU-23 medium containing 0.4% BSA for further culture of 20 and 144 h at 38.5 °C, 5% CO₂ in air. The activation rates of oocytes were higher in 6, 7 and 8% ethanol concentrations compared with 0, 5, 9 and 10% ethanol concentrations. Significantly, more oocytes (29.3–33.7%) were activated in the exposure for 8, 10, 12 and 15 min than those in the exposure for 0 and 5 min, but there was no difference due to exposure to 8% ethanol for 8–15 min. Oocytes treated by chemical agents (40.5–70.5%) after exposure to ethanol significantly improved the rate of oocyte activation compared with ethanol alone (31.2%). The percentage of cleaved oocytes was higher in the ethanol + CHX + CCB + 6-DMAP treatment (66.4%) than in other treatments (24.9–57.6%). Also,

* Corresponding author. Tel.: +82 42 821 5777; fax: +82 42 822 6712.

E-mail address: parkcs@cnu.ac.kr (C.S. Park).

the rate of blastocyst formation was higher in the ethanol + CHX + CCB + 6-DMAP treatment (25.0%) than in other treatments (0.0–19.3%). In conclusion, the optimal activation treatment of ethanol exposure alone for the in vitro matured porcine oocytes was 8% ethanol for 8–15 min. Oocytes activated by 8% ethanol for 10 min and incubated in the NCSU-23 medium supplemented with CHX, CCB and 6-DMAP for 3 h were more efficient for parthenogenetic development of in vitro matured porcine oocytes.

© 2004 Elsevier B.V. All rights reserved.

Keywords: Porcine oocyte; Parthenogenesis; Ethanol; Activation

1. Introduction

Activation of oocytes matured in vitro is essential for the success of animal cloning by nuclear transfer. Artificial stimuli elevating the cytoplasmic levels of calcium ions can induce activation of the oocytes even without penetration of the sperm into the oocyte. Chemical activating stimuli such as ethanol (Nagai, 1987), calcium ionophore and ionomycin (Jones et al., 1995; Tesarik and Sousa, 1995) are used to induce the artificial activation of mammalian oocytes. Ethanol was shown to induce oocyte activation in the mouse (Cuthberston et al., 1981; Cuthberston, 1983) and cattle (Nagai, 1987, 1992; Yang et al., 1994). Yang et al. (1994) reported that the combined treatment with 7% ethanol for 10 min and cycloheximide (CHX), the protein synthesis inhibitor, improved the rate of activation of the young oocytes of cattle. The histone kinase inhibitor, 6-dimethylaminopurine (6-DMAP), has been reported to accelerate and enhance the formation of pronuclei in non-age metaphase II oocytes from mice and cattle (Leal and Liu, 1998). Also, cytochalasin B (CCB) prevents the release of the second polar body after activation of the oocytes, which would result in diploid development (Fukui et al., 1992), and may also help prevent fragmentation of embryos (Collas and Robl, 1990; Yang et al., 1992). Beux et al. (2003) reported that 6-DMAP, cycloheximide, roscovitine and butyrolactone I could be used to block meiotic resumption in porcine oocytes in NCSU-23 culture medium. Although, many reports on porcine oocyte activation have been described, limited information is available on directly comparing the activation efficiency of various chemical agents mentioned above.

The purpose of this study was to investigate the effects of the different concentrations and exposure times of ethanol, and a sequential combination of ethanol, cycloheximide, cytochalasin B and 6-DMAP to improve parthenogenetic activation and development of in vitro matured porcine oocytes.

2. Materials and methods

2.1. Oocyte collection and in vitro maturation

Ovaries were collected from prepubertal gilts at a local slaughter house, and transported to the laboratory in 0.9% NaCl solution containing 75 µg/ml potassium penicillin G, 50 µg/ml streptomycin sulfate at 30–35 °C. Cumulus-oocyte complexes (COCs) were aspirated from

antral follicles (3–6 mm in diameter) using a 18-gauge needle fixed to a 10 ml disposable syringe. COCs were washed three times in Tyrode-lactate-pyruvate-polyvinylalcohol (mTLP-PVA) as described by Bavister (1989), and were washed twice with a maturation medium. Thirty-fourty COCs were transferred to 500 μ l of the same medium that had been covered with mineral oil in a 4-well multidish (Nunc, Roskilde, Denmark) and equilibrated at 38.5 °C, 5% CO₂ in air. The medium used for oocyte maturation was tissue culture medium (TCM) 199 supplemented with 26.19 mM sodium bicarbonate, 0.9 mM sodium pyruvate, 10 μ g/ml insulin, 2 μ g/ml vitamin B₁₂, 25 mM HEPES, 10 μ g/ml bovine apotransferrin, 500 μ M cysteamine, 10 IU/ml PMSG (Folligon, Intervet International B.V. Boxmeer, Holland), 10 IU/ml hCG (Chorulon, Intervet International B.V. Boxmeer, Holland), 10 ng/ml EGF (Sigma E-4127, USA), 0.4% BSA (Fraction V, Sigma A-8022, USA), 75 μ g/ml sodium penicillin G, 50 μ g/ml streptomycin sulfate and 10% pFF. After about 22 h of culture, oocytes were cultured without cysteamine and hormones for 22 h at 38.5 °C, 5% CO₂ in air.

2.2. Activation and culture of oocytes

After the completion of culture of oocytes for IVM, cumulus cells were removed with 0.1% hyaluronidase in mTLP-PVA, and washed three times with HEPES buffered (25 mM) NCSU-23 medium. Cumulus-free oocytes showing first polar body were selected and activated by various concentrations and exposure times of ethanol in the NCSU-23 medium.

Also, oocytes were activated with the NCSU-23 medium containing 8% ethanol for 10 min. After that, oocytes were incubated in the NCSU-23 medium supplemented with 10 μ g/ml cycloheximide, 7.5 μ g/ml cytochalasin B and 2 mM 6-dimethylaminopurine, with a single treatment or combination treatments for 3 h, respectively. Following activation, oocytes were washed three times with the NCSU-23 culture medium containing 0.4% BSA, and were transferred into 500 μ l of the same medium for further culture of 20 and 144 h at 38.5 °C, 5% CO₂ in air.

2.3. Examination of oocytes

At 20 h after activation, oocytes were fixed for 48 h in 25% acetic acid (v:v) in ethanol at room temperature, and stained with 1% (w:v) orcein in 45% (v:v) acetic acid to examine activated oocytes under a phase-contrast microscope at $\times 400$ magnification. Blastocysts at 144 h were stained with Hoechst 33342, and nuclei were counted under fluorescent microscope (Olympus, Japan).

Unless otherwise mentioned, all chemicals used in this study were purchased from Sigma Chemical Co. (St. Louis, MO, USA).

2.4. Experimental design

Experiment 1 was conducted to investigate activation of in vitro matured oocytes after exposure to ethanol for 10 min on the concentrations of 0, 5, 6, 7, 8, 9 and 10% ethanol, respectively.

Experiment 2 investigated activation of *in vitro* matured oocytes after 8% ethanol treatment for varying of 0, 5, 8, 10, 12 and 15 min exposure time, respectively.

Experiment 3 was carried out to investigate the effects of chemical activation agents. All oocytes were activated with the NCSU-23 medium containing 8% ethanol for 10 min. After that, oocytes were incubated in the NCSU-23 medium supplemented with CHX, CCB, 6-DMAP, CHX + CCB, CHX + 6-DMAP, CCB + 6-DMAP and CHX + CCB + 6-DMAP for 3 h, respectively.

2.5. Statistical analysis

Analyses of variance (ANOVA) were carried out using the [SAS software package \(1996\)](#). Duncan's multiple range test was used to compare mean value of individual treatments, when the *F*-value was significant ($P < 0.05$).

3. Results

3.1. Effects of ethanol concentrations on activation of porcine oocytes

As shown in [Table 1](#), seven different concentrations of ethanol were used to compare activation of porcine oocytes matured *in vitro*. The activation rates of oocytes were higher in 6, 7 and 8% ethanol concentrations compared with 0, 5, 9 and 10% ethanol concentrations. However, more oocytes were activated in the 8% ethanol treatment than in the other treatments. This concentration was therefore chosen for the successive experiments.

3.2. Effects of 8% ethanol exposure times on activation of porcine oocytes

As shown in [Table 2](#), significantly more oocytes (29.3–33.7%) were activated in the exposure for 8, 10, 12 and 15 min than those in the exposure for 0 and 5 min, but there was no difference due to exposure to 8% ethanol for 8–15 min.

Table 1
Effects of ethanol concentrations on activation of porcine oocytes matured *in vitro*^a

Concentration of ethanol (%)	No. of oocytes examined ^b	No. and percentage activated
0	119	7 (5.6 ± 3.6 ^c)
5	111	29 (25.8 ± 2.8 ^b)
6	116	34 (29.7 ± 1.8 ^{ab})
7	121	37 (30.7 ± 0.9 ^{ab})
8	118	41 (34.9 ± 1.6 ^a)
9	120	32 (26.4 ± 1.6 ^b)
10	121	14 (11.3 ± 3.1 ^c)

^a Oocytes were exposed to ethanol for 10 min after 44 h maturation in tissue culture medium (TCM) 199.

^b Experiments were repeated four times.

^{abc} Means ± S.E. in the same column with different superscripts differ significantly ($P < 0.05$).

Table 2

Effect of activation of porcine oocytes matured in vitro according to length of exposure by 8% ethanol

Length of exposure (min)	No. of oocytes examined ^a	No. and percentage activated
0	106	9 (8.7 ± 3.3^c)
5	100	24 (24.0 ± 1.7^b)
8	102	30 (29.3 ± 2.0^{ab})
10	103	35 (33.7 ± 2.1^a)
12	102	32 (31.5 ± 1.9^{ab})
15	107	33 (30.1 ± 2.0^{ab})

^a Experiments were repeated four times.^{abc} Means $\pm$ S.E. in the same column with different superscripts differ significantly ($P < 0.05$).

Table 3

Effects on activation of porcine oocytes matured in vitro treated by ethanol or ethanol plus various chemical agents

Treatment	No. of oocytes examined ^a	No. and percentage activated
Ethanol	103	36 (31.2 ± 3.3^e)
	Ethanol plus	
CHX	109	52 (46.6 ± 2.7^{cd})
CCB	107	55 (50.5 ± 2.4^c)
6-DMAP	108	46 (40.5 ± 3.8^d)
CHX + CCB	108	69 (62.7 ± 3.0^{ab})
CHX + 6-DMAP	108	52 (47.8 ± 3.1^{cd})
CCB + 6-DMAP	107	59 (54.3 ± 3.3^{bc})
CHX + CCB + 6-DMAP	105	75 (70.5 ± 3.3^a)

CHX: 10 μ g/ml Cycloheximide, CCB: 7.5 μ g/ml Cytochalasin B, 6-DMAP: 2 mM 6-dimethylaminopurine.^a Experiments were repeated five times.^{abcde} Means $\pm$ S.E. in the same column with different superscripts differ significantly ($P < 0.05$).

3.3. Effect of chemical agents for parthenogenetic activation and development

As shown in Table 3, the chemical agent treatments followed by exposure to ethanol significantly improved ($P < 0.05$) the rate of oocyte activation compared with that of the ethanol alone. The ethanol + CHX + CCB + 6-DMAP treatment significantly improved the rate of oocyte activation compared with that of the other seven treatments.

As shown in Table 4, the percentages of cleaved oocytes and blastocyst formation were higher with the ethanol + CHX + CCB + 6-DMAP treatment than in the other seven treatments. The chemical agent treatments improved ($P < 0.05$) the cell numbers per blastocyst compared with those of ethanol alone, but there were no differences among the chemical agent treatments.

4. Discussion

Oocyte activation was induced by ethanol in cattle (Nagai, 1987, 1992; Yang et al., 1994). Liu et al. (1998) reported that both sequential combination activation procedures of

Table 4

Effect on development of porcine oocytes matured in vitro treated by ethanol or ethanol plus various chemical agents

Treatment	No. of oocytes examined ^a	% of cleaved oocytes	% of blastocyst from cleaved oocytes	Cell no. per blastocyst
Ethanol	100	24.9 ± 4.3 ^f	0.0 ± 0.0 ^f	0.0 ± 0.0 ^b
Ethanol plus				
CHX	98	43.8 ± 3.4 ^{de}	8.5 ± 2.3 ^{de}	21.4 ± 0.8 ^a
CCB	98	48.0 ± 2.6 ^{cd}	12.2 ± 1.1 ^{cd}	23.3 ± 0.9 ^a
6-DMAP	97	39.3 ± 1.0 ^e	4.4 ± 1.9 ^{ef}	21.3 ± 1.3 ^a
CHX + CCB	107	57.6 ± 1.5 ^b	19.3 ± 1.5 ^b	24.6 ± 1.2 ^a
CHX + 6-DMAP	106	45.7 ± 2.5 ^{cde}	10.0 ± 2.6 ^{cd}	21.5 ± 0.8 ^a
CCB + 6-DMAP	104	52.6 ± 1.4 ^{bc}	14.5 ± 0.7 ^{bc}	24.5 ± 1.2 ^a
CHX + CCB + 6-DMAP	108	66.4 ± 2.0 ^a	25.0 ± 2.3 ^a	25.6 ± 1.3 ^a

CHX: 10 µg/ml Cycloheximide, CCB: 7.5 µg/ml Cytochalasin B, 6-DMAP: 2 mM 6-dimethylaminopurine.

^a Experiments were repeated five times.

^{abcdef} Means ± S.E. in the same column with different superscripts differ significantly ($P < 0.05$).

calcium ionophore A23187 (CaA) + 6-dimethylaminopurine, and CaA + cycloheximide + cytochalasin D were effective for the activation and development of young bovine oocytes. In this study, we were interested in testing the effects of the optimal concentration and length of exposure to ethanol for activating porcine oocytes matured in vitro. The synergistic effect of ethanol with CHX, CCB and 6-DMAP treatments was also investigated. Our results indicated that exposure to 8% ethanol for 10 min activated 33.7–34.9% of the porcine oocytes matured in vitro. This rate of activation was similar to other reports on activation of bovine oocytes with ethanol (Nagai, 1987; Yang et al., 1994). It is known that ethanol can induce a single intracellular free-calcium increase in oocytes which can cause activation of some oocytes (Cuthberston et al., 1981). Liu et al. (1998) reported that optimal parthenogenetic development of newly matured bovine oocytes can be obtained by calcium ionophore treatment by incubation with either 6-DMAP or cycloheximide + cytochalasin D. This study confirmed that treatment with ethanol alone resulted in low activation, cleavage and development rates compared with the combined treatments of ethanol plus various chemical agents.

A protein synthesis inhibitor, cycloheximide has been used to induce pronuclear development in matured mouse oocytes (Siracusa et al., 1978; Clarke and Masui, 1983) and in cattle (Sirard et al., 1989). Presicce and Yang (1994a,b) reported that activation rates achieved with the combined ethanol and cycloheximide treatment were high for both young and aging oocytes. Ethanol-induced Ca^{2+} elevation would inactivate the existing cytosstatic factor (CSF), and the subsequent cycloheximide exposure would prevent renewal of CSF synthesis in the oocyte. A protein serine/threonine kinase (or phosphorylation) inhibitor, 6-dimethylaminopurine has been shown to enhance the activation stimulus and to accelerate pronuclear formation and parthenogenetic development in the young mouse and bovine oocytes (Moses and Masui, 1994; Moses et al., 1995; Takahashi et al., 1996). After the combined treatments with ethanol plus various chemical agents, we also observed an increased rate of parthenogenetic development in activated oocytes compared with the treatment with

ethanol alone. However, the activation, cleavage and blastocyst formation rates of oocytes with the treatments with ethanol + CHX + CCB, ethanol + CCB + 6-DMAP and ethanol + CHX + CCB + 6-DMAP were significantly increased compared with the treatments with ethanol + CHX, ethanol + CCB, ethanol + 6-DMAP and ethanol + CHX + 6-DMAP. Cytochalasin was reported to inhibit extrusion of the second polar body; in the presence of cytochalasin, segregation of the chromosomes occurred, but cytokinesis did not take place. This resulted in diploid zygotes with two pronuclei (Presicce and Yang, 1994a,b; Liu et al., 1998). The protein kinase inhibitor 6-DMAP, on the other hand, is known to cause the second meiotic spindle to disintegrate and the oocytes to pass directly into interphase (Navara et al., 1994). In our experiment, the activation, cleavage and blastocyst formation rates were more effective following incubation in the treatments with cytochalasin B. In conclusion, the optimal activation treatment for ethanol exposure alone for the porcine oocytes matured in vitro was 8% ethanol for 8–15 min. Also, we can conclude that the combined treatment of in vitro matured porcine oocytes with ethanol + CHX + CCB + 6-DMAP significantly increased the activation, cleavage and blastocyst formation rates. Further comparative investigations will be needed to show what effects each inhibitor has on oocyte quality and further developmental competence.

Acknowledgements

This work was supported by grant No. R11-2002-100-00000-0 from ERC program of the Korea Science & Engineering Foundation.

References

- Bavister, B.D., 1989. A consistently successful procedure for in vitro fertilization of golden hamster eggs. *Gamete Res.* 23, 139–158.
- Beux, G.L., Richard, F.J., Sirard, M.A., 2003. Effect of cycloheximide, 6-DMAP, roscovitine and butylolactone I on resumption of meiosis in porcine oocytes. *Theriogenology* 60 (6), 1049–1058.
- Clarke, H.J., Masui, Y., 1983. The induction of reversible and irreversible chromosome decondensation by protein synthesis inhibition during meiotic maturation of mouse oocytes. *Dev. Biol.* 97, 291–301.
- Collas, P., Robl, J.M., 1990. Factors affecting the efficiency of nuclear transplantation in the rabbit embryo. *Biol. Reprod.* 43, 877–884.
- Cuthberston, K.S.R., Whittingham, D.G., Cobbold, P.H., 1981. Free calcium increase in exponential phases during mouse oocyte activation. *Nature* 294, 754–757.
- Cuthberston, K.S.R., 1983. Parthenogenetic activation of mouse oocytes in vitro with ethanol and benzyl alcohol. *J. Exp. Zool.* 226, 311–314.
- Fukui, Y., Sawai, K., Furudate, M., Sato, N., Iwazumi, Y., Ohsaki, K., 1992. Parthenogenetic development of bovine oocytes treated with ethanol and cytochalasin B after in vitro maturation. *Mol. Reprod. Dev.* 33, 357–362.
- Jones, K.T., Carroll, J., Whittingham, D.G., 1995. Ionomycin, thapsigargin, ryanodine and sperm induced Ca^{2+} release increase during meiotic maturation of mouse oocytes. *J. Biol. Chem.* 270, 6671–6677.
- Leal, C.L., Liu, L., 1998. Differential effects of kinase inhibitor and electrical stimulus on activation and histone H kinase activity in pig oocytes. *Anim. Reprod. Sci.* 52 (1), 51–61.
- Liu, L., Ju, J.C., Yang, X.Z., 1998. Differential inactivation of maturation-promoting factor and mitogen-activated protein kinase following parthenogenetic activation of bovine oocytes. *Biol. Reprod.* 59, 537–545.

- Moses, R.M., Masui, Y., 1994. Enhancement of mouse egg activation by the kinase inhibitor, 6-dimethylaminopurine (6-DMAP). *J. Exp. Zool.* 270, 211–218.
- Moses, R.M., Kline, D., Masui, Y., 1995. Maintenance of metaphase in colcemid-treated mouse eggs by distinct calcium- and 6-dimethylaminopurine (6-DMAP)-sensitive mechanisms. *Dev. Biol.* 167, 829–837.
- Nagai, T., 1987. Parthenogenetic activation of cattle follicular oocytes in vitro with ethanol. *Gam. Res.* 16, 243–249.
- Nagai, T., 1992. Development of bovine in vitro-matured follicular oocytes activated with ethanol. *Theriogenology* 37, 869–875.
- Navara, C.S., First, N.L., Schatten, G., 1994. Microtubule organization in the cow during fertilization, polyspermy, parthenogenesis and nuclear transfer: the role of the sperm aster. *Dev. Biol.* 162, 29–40.
- Presicce, G.A., Yang, X., 1994a. Nuclear dynamics of parthenogenesis of bovine oocytes matured in vitro for 20 and 40 h and activated with combined ethanol and cycloheximide treatment. *Mol. Reprod. Dev.* 37, 61–68.
- Presicce, G.A., Yang, X., 1994b. Parthenogenetic development of bovine oocytes matured in vitro for 24 h and activated by ethanol and cycloheximide. *Mol. Reprod. Dev.* 38 (4), 380–385.
- SAS Institute Inc., 1996. The SAS system for Windows. Release 6.12, Cary, NC.
- Siracusa, G., Whittingham, D.G., Molinaro, M., Vivarelli, E., 1978. Parthenogenetic activation of mouse oocytes induced by inhibitors of protein synthesis. *J. Embryol. Exp. Morphol.* 43, 157–166.
- Sirard, M.A., Florman, H.M., Leibfried-Rutledge, M.L., Barnes, F.L., First, N.L., 1989. Timing of nuclear progression and protein synthesis necessary for meiotic maturation of bovine oocytes. *Biol. Reprod.* 40, 1257–1263.
- Takahashi, S., Kubota, C., Ogata, Y., Tokunaga, T., Imai, H., 1996. Parthenogenetic activation and development of bovine oocytes treated with protein synthesis or protein phosphorylation inhibitors. *Theriogenology* (Suppl. 1) 45 (1), 156.
- Tesarik, J., Sousa, M., 1995. More than 90% fertilization rates after intracytoplasmic sperm injection and artificial induction of oocyte activation with calcium ionophore. *Fertil. Steril.* 63, 343–349.
- Yang, X., Jiang, S., Shi, Z., 1992. Improved activation by combined cycloheximide and electrical pulse treatment of bovine follicular oocytes matured in vitro for 23–24 h. *Biol. Reprod. (Suppl.)* 46, 117 (abstract).
- Yang, X., Presicce, G.A., Moraghan, L., Jiang, S.E., Foote, R.H., 1994. Synergistic effect of ethanol and cycloheximide on activation of freshly matured bovine oocytes. *Theriogenology* 41, 395–403.