

FORMULARIO ÚNICO DE OPOSICIÓN

①

OPOSICIÓN A:

- | | |
|--|--|
| ☒ Patente de Invención. | ☐ Marca Colectiva. |
| ☐ Patente de Modelo de Utilidad. | ☐ Marca de Certificación. |
| ☐ Diseño Industrial. | ☐ Lema Comercial. |
| ☐ Esquema de Trazado para Circuitos Integrados. | |
| ☐ Marcas de Productos o Servicios. | |

②

Número del expediente: 06-038472

Solicitante: BRISTOL-MYERS SQUIBB COMPANY

**SOLICITUD ·
PUBLICADA**

Apoderado: JUAN PABLO CADENA SARMIENTO

Título de la nueva creación o denominación del signo: PROCESO PARA LA ELABORACIÓN DE ARIPIPAZOL ESTERIL DE TAMAÑO DE PARTÍCULA PROMEDIO DESEADO.

Gaceta: **567**

③ OPOSICIÓN PRESENTADA POR:

SOLICITANTE	Nombre: PROCAPS S.A Dirección: CALLE 80 No. 78B -201 Nacionalidad o Domicilio: COLOMBIA Lugar de Constitución: BARRANQUILLA	IDENTIFICACIÓN C.C. ☐ NT ☒ C.E. ☐ Otro ☐ Cual _____ Número <u>890.106.527-5</u>
	Teléfono: (571)3719900 Fax: (571)3730818 E-mail:	
REPRESENTANTE O APODERADO	Nombre: REINALDO PINILLA MORENO Dirección: TRANSVERSAL 13ª No. 119-95 OF 104 Teléfono: (571)2131213 Fax: (571)6121979 E-mail:	IDENTIFICACIÓN C.C. ☐ NT ☒ C.E. ☐ Otro ☐ Cual _____ Número <u>80.425.609</u> TP <u>91563</u>

④ FUNDAMENTOS DE LA OPOSICIÓN:

Art.14, 16, 18, 20 y 30 de la Decisión 486 de la CAN

Causal: No cumple con los requisitos de patentabilidad.

Signo opozitor:

Justificación:

La presente solicitud no cumple con los requisitos de patentabilidad dado que lo revelado, ya se encontraba en el estado de la técnica o es posible derivarlo de la misma tal y como se demostrará en el capítulo "Fundamento de Hechos".

⑤ Comprobante de pago No. _____ Fecha _____

⑥ ANEXOS

- ☒ Comprobante de pago de la tasa.
- ☒ Poderes, si fuere el caso.
- ☒ Pruebas de los fundamentos de la oposición.
- ☐ Documentos que acrediten el legítimo interés, si fuere el caso.
- ☒ Documento que acredita la existencia y representación legal de las partes, cuando sean personas jurídicas.

⑦

NOMBRE: REINALDO PINILLA MORENO

FIRMA: _____

C.C. 80425609

TP. 91563 CSJ

PRESENTACION PERSONAL

En Bogotá, D.C. a 28 NOV 2006 se presentó

en la Notaría Decana, cuyo titular es SERGIO FRANCO LEON

REINALDO PINILLA MORENO


Con cédula No. 80425609

Expedida en WUQUEN y dijo que

el anterior documento es cierto y que la firma es de su puño y

letra. En constancia se firma _____





ptimum

Derecho de la Competencia, Propiedad Industrial y Comercio Exterior

Señora Jefe,

DIVISIÓN DE NUEVAS CREACIONES

Superintendencia de Industria y Comercio

E. S. D.

Referencia : Expediente N° 06 038.472
Asunto : Oposición a la solicitud de patente de Invención titulada "PROCESO PARA LA ELABORACIÓN DE ARIPIPRAZOL ESTÉRIL DE TAMAÑO DE PARTÍCULA PROMEDIO DESEADO".
Solicitante : BRISTOL-MYERS SQUIBB COMPANY.
Opositora : PROCAPS S.A.

Publicada en la Gaceta de la Propiedad Industrial


N° 567 del 31 de Agosto de 2006

Yo, **REINALDO PINILLA MORENO**, abogado con tarjeta profesional N° 91563 del consejo superior de la judicatura, en mi condición de apoderado de la sociedad **PROCAPS S.A.** conformada bajo las leyes de la República de Colombia y domiciliada en la ciudad de Barranquilla, Colombia, atentamente manifiesto que estando dentro del término de ley y por orden expresa de mi representada, presento oposición a la concesión del registro de la patente de Invención titulada "**PROCESO PARA LA ELABORACIÓN DE ARIPIPRAZOL ESTÉRIL DE TAMAÑO DE PARTÍCULA PROMEDIO DESEADO**", solicitada por **BRISTOL-MYERS SQUIBB COMPANY.**, y pido a Usted declarar fundada esta oposición y en consecuencia negar el registro de la patente solicitada.

Transversal 13° N° 119-95, Of. 104 Tel. (571) 2 13 12 13 - 6 20 50 21 - Fax. (571) 6 12 19 79
E-mail: negocios@optimum-co.com - www.optimum-co.com
Bogotá Colombia

2.2. La solicitud colombiana 06 038.472 objeto de la presente oposición, se refiere a un proceso para preparar aripiprazol estéril cristalino de tamaño de partícula pequeño

Transversal 13° N° 119-95, Of. 104 Tel. (571) 2 13 12 13 - 6 20 50 21 - Fax. (571) 6 12 19 79
E-mail: negocios@optimum-co.com - www.optimum-co.com
Bogotá Colombia



Optimum

137 4
Derecho de la Competencia, Propiedad Industrial y Comercio Exterior

1. INTERÉS PARA ACTUAR


Conforme a lo dispuesto en el artículo 42 de la Decisión 486 de 2000 de la CAN, estando dentro del término estipulado por la ley, manifiesto el legítimo interés de nuestra representada para oponerse al registro de la patente de la referencia, atendiendo a que la solicitud pretendida no cumple con los requisitos dispuestos en la ley como se analizará más adelante y de ser concedida afectaría directamente la comercialización de algunos productos de mi representada, que contienen, en particular Aripiprazol. Por lo tanto, solicito respetuosamente a ese Despacho se sirva tener en cuenta la presente oposición dentro del trámite administrativo que se surte en esta Superintendencia.

2. FUNDAMENTOS DE HECHOS

2.1. La solicitud objeto de la presente fue presentada el 24 de abril de 2006, reivindicando prioridad bajo las solicitudes US60/513,886 del 23 de octubre de 2003 y fue publicada en Colombia, en la Gaceta de la Propiedad Industrial N° 567 del 31 de agosto de 2006, página 376, N° publicación 354, cuya publicación internacional corresponde a la WO2005041970 del 12 de mayo de 2005.

2.2. La solicitud colombiana 06 038.472 objeto de la presente oposición, se refiere a un proceso para preparar aripiprazol estéril cristalino de tamaño de partícula pequeño

Transversal 13ª N° 119-95, Of. 104 Tel. (571) 2 13 12 13 - 6 20 50 21 - Fax. (571) 6 12 19 79
E-mail: negocios@optimum-co.com - www.optimum-co.com
Bogotá Colombia



deseado y distribución de tamaño de partícula estrecho, caracterizado porque comprende: (a) proporcionar una corriente de chorro de una solución de aripiprazol en un solvente orgánico; (b) proporcionar una corriente de chorro de anti-solvente la cual es capaz de iniciar la precipitación de aripiprazol de la solución; (c) ocasionar que la corriente de chorro de la solución de aripiprazol en solvente y la corriente de chorro de anti-solvente choquen entre sí y colisionen entre sí para crear turbulencia elevada en su punto de impacto, cada corriente de chorro que tiene suficiente velocidad lineal logra micromezclado de intensidad elevada de cada corriente previo a la nucleación para producir una suspensión de cristales de monohidrato de aripiprazol; y (d) recuperar cristales de monohidrato de aripiprazol de tamaño de partícula pequeño deseado y distribución de tamaño de partícula estrecho.


2.3. Ahora bien, el artículo 14 de la Decisión 486 de la CAN consagra en materia de nuevas creaciones que: *"Los países miembros otorgarán patentes para las invenciones, sean de producto o procedimiento, en todos los campos de la tecnología, siempre que sean nuevas, tengan nivel inventivo y sean susceptibles de aplicación industrial"*. Así las cosas, las invenciones objeto de patente deben ser nuevas y tener nivel inventivo. Bien, pues antes de la fecha de prioridad de esta solicitud se encuentran el siguiente documento que afecta la altura inventiva de la presente solicitud:

- D1: WO03026659, 2003-04-03.

2.4. Por su parte, el documento D1 en la descripción con sus ejemplos y reivindicaciones, particularmente a partir de la página 4 línea 6, se refiere a un proceso para preparar una forma de aripiprazol, que tiene una reducida higroscopicidad, de manera que no se conviertan en un hidrato, ni perdiera su

solubilidad original aún cuando se prepare un compuesto medicinal que contenga los cristales de anhídrido de aripiprazol y permanezca durante un periodo prolongado. Según dicha anterioridad, se revela que la reducida higroscopicidad de aripiprazol es una sustancia cristalina definida con un anhídrido B. Según dicha anterioridad, se define que dentro del proceso de preparación de dicho anhídrido B se forma un anhídrido A intermedio. Allí se define también, que los cristales anhídridos de aripiprazol son obtenidos por procesos de calentamiento a una temperatura específica, con el uso de solventes específicos. Según la descripción dichos solventes pueden ser solventes orgánicos que contiene agua. Según ello, el solvente orgánico debe ser miscible con el agua, tal como un alcohol como el metanol, etanol, propilalcohol o isopropanol, una quetona como la acetona, un éter como el tetrahidrofurano, dimetilformamida, o una mezcla de los mismos. Según la descripción, el proceso de manufactura del hidrato A, define medios de mezclado, calentamiento y medios de hidratación convencional tal como atomizado. Para la manufactura del hidrato B, se definen medios de calentamiento entre 90 y 125°C. Así las cosas, a partir del presente documento, cualquier persona versada en la materia, puede llegar a definir la conformación de un proceso sobre el cual se combine una solución de arpiprazole y un solvente orgánico con una solución antisolvente y esta a su vez produzca cristales de monohidrato de aripiprazol. Son tan cercanos los procesos, que mediante el proceso definido en la anterioridad D1 de manufactura de aripiprazole cristalino, se pueden establecer tamaños de partícula entre 14 y 50um. Por lo tanto, a los ojos de este documento D1, la presente solicitud no comprende altura o nivel inventivo.

2.5. En consecuencia y de acuerdo a lo descrito anteriormente, se deriva que la solicitud colombiana 06 038.472 no puede pretender un monopolio por patente toda vez



140 7

que lo que pretende no cumple con uno de los tres requisitos necesarios para su protección.

2.6. Que por consiguiente, respecto a la solicitud pretendida se tiene que **CARECE DE ALTURA INVENTIVA**, dado que según la Decisión 486 de la Comunidad Andina de Naciones, Artículo 18.- *"Se considerará que una invención tiene nivel inventivo, si para una persona del oficio normalmente versada en la materia técnica correspondiente, esa invención no hubiese resultado obvia ni se hubiese derivado de manera evidente del estado de la técnica."* A la fecha de prioridad, 23 de octubre de 2003, a partir del documento D1, es posible deducir el proceso de preparación de aripiprazol de tamaño de partícula promedio deseado según lo definido en las reivindicaciones de la presente solicitud.

2.7. Dado lo anterior, es posible deducir que la solicitud y lo reivindicado no es objeto de patentabilidad pues es contrario a lo que dicta la Decisión 486 de la Comunidad Andina de Naciones, Artículo 14.- *"Los Países Miembros otorgarán patentes para las invenciones, sean de producto o de procedimiento, en todos los campos de la tecnología, siempre que sean nuevas, tengan nivel inventivo y sean susceptibles de aplicación industrial."* La solicitud pretendida no cumple con uno de los tres requisitos de patentabilidad.

2.8. Por todo lo anterior, atentamente solicito a usted que se sirva declarar fundada la presente oposición y negar el registro Patente de invención denominada **"PROCESO PARA LA ELABORACIÓN DE ARIPIPAZOL ESTÉRIL DE TAMAÑO DE PARTÍCULA PROMEDIO DESEADO"**, solicitada por **BRISTOL-MYERS SQUIBB COMPANY**, solicitud Colombiana que apareció publicada en la Gaceta de la Propiedad Industrial N°-567.

3. PRUEBAS

De conformidad con lo dispuesto por las normas vigentes, solicito a usted tener como prueba en este asunto los siguientes:

- D1: WO03026659, 2003-04-03.

4. FUNDAMENTO DE DERECHO

Fundamento esta observación en las normas legales siguientes:


- 4.1. Numeral 1 del artículo 586 del Código de Comercio.
- 4.2. Artículo 42 de la Decisión 486 de la Comisión del Acuerdo de Cartagena.
- 4.3. En las demás normas concordantes.

5. ANEXOS

Para los efectos correspondientes anexo al presente escrito los siguientes documentos:

- D1: WO03026659, 2003-04-03.

6. NOTIFICACIONES



ptimum

142-9
Derecho de la Competencia, Propiedad Industrial y Comercio Exterior

Para efectos de notificaciones que por disposición del artículo 50 del Decreto 01 de 1984 debe hacer su Despacho, informo que mi representada y la suacrita recibiremos notificaciones en la Secretaría de su Despacho o en mis oficinas situadas en la Transversal 13ª N° 119-95, Of. 104 de esta ciudad de Bogotá.

Señor Superintendente,

REINALDO PINILLA MORENO

C.C. 80'425.609 de Bogotá

T.P.A. 91563 del C.S. de la J.

PRESENTACION PERSONAL

En Bogotá, D.C. a **28 NOV 2006** se presento


en la Notaría Décima, cuyo titular es **SERGIO FRANCO LEON**

REINALDO PINILLA MORENO

Con cédula No. **80425609 - TPA 91563**

Expedida en **USAKEN** y dijo que el anterior documento es cierto y que la firma es de su puño y

letra. En constancia se firma



143

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO
Rad: 06-038472-00003-0000
Tm: 11 PATENTENYII
Lvs: 3/8 PASSEMAJUALI
M: 400 OPOSICIONES
Folios: 187

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO

MIT : 800176089-2

RECIBO OFICIAL DE CAJA : 06 - 91,397
FECHA : NOVIEMBRE 28 DE 2006


***** CONSIGNACION *****

POSITANTE	TIPO PAGO	BANCO	CUENTA	No. PAGO	FECHA PGO	Vr. PAGO
GYNOFARM S.A.	CONSIGNACION	BANCO POPULAR	050-00110-6	44837800	27/09/2006	260,500.00
LAB. PROCAPS	CONSIGNACION	BANCO POPULAR	050-00110-6	60060415	31/10/2006	246,000.00

***** CONCEPTO *****

CANT. REMITISTICO	CONCEPTO	TOTAL CONCEPTO
1 50005-01-04 PRESENTACION DE OBSERVACIONES	12 MARCAS Y LENAS COMERCIALES	246,000.00
	TOTAL :	\$ 246,000.

SON: DOSCIENTOS CUARENTA Y SEIS MIL PESOS

RESPONSABLE : 



PODER ESPECIAL

Yo, **WALTER TANAKA TATEKAWA**, mayor de edad y vecino de la ciudad de Barranquilla (Colombia), identificado como aparece al pie de mi firma, quien en el presente documento actúa en calidad de Representante Legal de la sociedad **PROCAPS S.A.** sociedad legalmente constituida y domiciliada en la ciudad de Barranquilla (Colombia), por el presente documento otorgo a **REINALDO PINILLA MORENO**, abogado titulado, identificado como aparece al pie de su firma, con tarjeta profesional No. 91563 expedida por el Consejo Superior de la Judicatura, poder especial, amplio y suficiente, para formular oposiciones u observaciones a la solicitud de patente No. 06 038.472, titulada **PROCESO PARA LA ELABORACIÓN DE ARIPIPAZOL ESTERIL DE TAMAÑO DE PARTÍCULA PROMEDIO DESEADO** de **BRISTOL-MYERS SQUIBB COMPANY**, publicada en la gaceta No. 567 del 31 de Agosto de 2006.

Nuestro apoderado queda facultado, de manera expresa, para realizar todo tipo de actos necesarios para llevar a cabo el trámite de la oposición, incluyendo, sin limitación, preparar y presentar solicitudes, hacer declaraciones, pagar tasas, contribuciones e impuestos, recabar los títulos o certificados. Asimismo quedan autorizados para impugnar administrativa y judicialmente todo lo que fuere resuelto en el expediente a que se hace expresa referencia en el presente poder, con todas las facultades generales y específicas que fuera requerido para ella.

De la misma manera, además de las facultades que le confiere la naturaleza del presente mandato, se encuentran facultados para recibir, desistir, transigir, sustituir, y reasumir el presente poder, incoar acciones, interponer recursos y demás facultades conferidas por la ley.

Dado y Firmado en Barranquilla (Colombia) a los 20 días del mes de Noviembre de 2006.


PROCAPS S.A.

WALTER TANAKA TATEKAWA
C.C. N° 16.251.923 de Palmira
Representante Legal - atrás

Acepto,


REINALDO PINILLA MORENO

C.C. 80'425.609 de Bogotá
T.P.A. 91563 del C.S. de la J.

ADY IN-888. NÚMERO SECURIA

NOTARIA SEGUNDA DE PRIMERA DE BARRANQUILLA

CERTIFICA

Que la(s) firma(s) que aparece(n) en el documento que antecede se corresponde(n) a la(s) de WALTER
TORRES TATEKOW

Que aparece(n) registrada(s) en esta Notaría previa confrontación de la(s) manual(es).

(Decreto 98871)

Barranquilla 22 de NOV de 2006

[Handwritten signature]

[Circular notary seal: REPUBLICA DE COLOMBIA, LEI NOTARIA SEGUNDA DE PRIMERA DE BARRANQUILLA, CIRCULO DE BARRANQUILLA]

[Handwritten signature]



***** Pag. 1
C A M A R A D E C O M E R C I O
D E B A R R A N Q U I L L A
05*****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

EL SUSCRITO SECRETARIO DE LA CAMARA DE COMERCIO DE BARRANQUILLA,

C E R T I F I C A

Que por Escritura Pública No. 1.190 del 22 de Julio de 1976, otorgada en la Notaria Tercera de Barranquilla, inscrito(as) en esta Cámara de Comercio, el 28 de Julio de 1976 bajo el No. 6.070 del libro respectivo, fue constituida la sociedad----- limitada denominada "PRODUCTORA DE CAPSULAS DE GELATINAS LIMITADA "PROCAPS LIMITADA".-----

C E R T I F I C A

Que por Escritura Pública No. 1.307 del 28 de Junio de 1979, otorgada en la Notaria Tercera de Barranquilla, inscrito(as) en esta Cámara de Comercio, el 23 de Julio de 1979 bajo el No. 10.039 del libro respectivo, la sociedad antes mencionada----- se transformo en sociedad anonima bajo la denominacion social de "PRODUCTORA DE CAPSULAS DE GELATINA S.A." PROCAPS S.A.".-----

C E R T I F I C A

Que por Escritura Pública No. 3.810 del 31 de Dic/bre de 1980, otorgada en la Notaria Primera de Barranquilla, inscrito(as) en esta Cámara de Comercio, el 30 de Enero de 1981 bajo el No. 12.705 del libro respectivo, la sociedad antes mencionada----- se transformo en sociedad de responsabilidad limitada bajo al razon social de "PRODUCTORA DE CAPSULAS DE GELATINA LIMITADA "PROCAPS LI - MITADA".-----

C E R T I F I C A

Que por Escritura Pública No. 3.755 del 10 de Nov/bre de 1983, otorgada en la Notaria Unica de Soledad, inscrito(as) en esta Cámara de Comercio, el 22 de Nov/bre de 1983 bajo el No. 17.922 del libro respectivo, la sociedad antes mencionada----- se transformo en sociedad anonima bajo la denominacion social de "PRODUCTORA DE CAPSULAS DE GELATINA S.A." "PROCAPS S.A.".-----

C E R T I F I C A

Que por Escritura Pública No. 3.615 del 06 de Dic/bre de 1996, otorgada en la Notaria 2a. de Barranquilla, inscrito(as) en esta Cámara de Comercio, el 20 de Dic/bre de 1996 bajo el N. 67.218 del libro respectivo, la sociedad antes menci nada-----

***** C O N T I N U A *****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

cambio de razon social, por la denominacion LABORATORIOS PROCAPS S.A

C E R T I F I C A

Que por Escritura Pública No. 3.775 del 18 de Dic/bre de 1996, otorgada en la Notaria 2a. de Barranquilla, inscrito(as) en esta Cámara de Comercio, el 23 de Enero de 1997 bajo el No. 67.704 del libro respectivo, la sociedad antes mencionada-----
cambio de razon social, por la denominacion PROCAPS S.A. -----

C E R T I F I C A

Que dicha sociedad ha sido reformada por las siguientes escrituras y/o documentos privados:

Numero	aaaa/mm/dd	Notaria	No. Insc o Reg	aaaa/mm/dd
39	1979/01/18	Notaria 3a. de Barranquilla.	9.452	1979/01/24
992	1981/04/30	Notaria 1a. de Barranquilla.	13.112	1981/05/06
1.464	1982/08/27	Notaria Unica de Soledad	15.533	1982/09/08
2.744	1983/08/18	Notaria Unica de Soledad	17.416	1983/08/26
3.826	1984/07/16	Notaria Unica de Soledad	19.554	1984/07/27
3.835	1985/08/26	Notaria Unica de Soledad	22.230	1985/09/04
2.260	1986/09/10	Notaria Unica de Soledad.	24.995	1986/09/15
1.849	1989/07/11	Notaria 2a. de Barranquilla.	33.909	1989/07/13
444	1990/02/21	Notaria 2a. de Barranquilla.	36.101	1990/03/16
3.090	1991/12/17	Notaria 2a. de Barranquilla	43.993	1992/01/28
269	1993/02/01	Notaria 2a. de Barranquilla	48.428	1993/02/16
2.342	1993/07/28	Notaria 2a. de Barranquilla	50.582	1993/08/10
3.652	1993/11/09	Notaria 2a. de Barranquilla	51.798	1993/11/22
3.615	1996/12/06	Notaria 2a. de Barranquilla	67.218	1996/12/20
89	1997/01/16	Notaria 2a. de Barranquilla	67.704	1997/01/23
2.070	1997/07/17	Notaria 2a. de Barranquilla	70.627	1997/07/31
2.281	1997/08/04	Notaria 2a. de Barranquilla	70.813	1997/08/14
765	1999/04/22	Notaria 3a. de Barranquilla	80.628	1999/04/23
459	2001/02/21	Notaria 2a. de Barranquilla	91.502	2001/02/23
2.608	2001/10/05	Notaria 2a. de Barranquilla	95.771	2001/11/08
1.730	2004/07/29	Notaria 2. de Barranquilla	112.999	2004/08/26
416	2005/03/03	Notaria 2. de Barranquilla	116.364	2005/03/10
1.613	2005/07/28	Notaria 2. de Barranquilla	119.322	2005/08/18

C E R T I F I C A

Que de acuerdo con la(s) escritura(s) arriba citada(s), la sociedad se rige por las siguientes disposiciones:

DENOMINACION O RAZON SOCIAL:

PROCAPS S.A.-----

SIGLA: PROCAPS.

DOMICILIO PRINCIPAL: Barranquilla.

NIT No: 890.106.527-5.

MATRICULA MERCANTIL: 24.802.

C E R T I F I C A

***** C O N T I N U A *****



***** Pag. 2
CAMARA DE COMERCIO
DE BARRANQUILLA
05*****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

Direccion para notificaciones judiciales:

CL 80 No 78 B - 201.

De la ciudad de Barranquilla.

C E R T I F I C A

DURACION: El término de duración de la sociedad se fijó hasta el 18 de Agosto de 2073.

C E R T I F I C A

OBJETO SOCIAL: El objeto principal de la sociedad será: a)-La fabricacion de capsulas y otros productos similares a partir de materiales tales como gelatinas u otros de distinta naturaleza, que puedan ser utilizados como envases o recipientes de productos y sustancias quimicas, farmaceuticas y cosmeticas, para uso y consumo humano y veterinario; b)-La investigacion, desarrollo y fabricacion de toda clase de productos y sustancias quimicas, farmaceuticas y cosmeticas para uso y consumo humano, animal y vegetal; c)-La compra y venta de materias primas, insumos y productos intermedios necesarios para los procesos de fabricacion de sustancias quimicas, farmaceuticas, alimenticias y cosmeticas, asi como la comercializacion interna y externa de los productos terminados, d) La compra, venta, importacion, exportacion y distribucion de toda clase de elementos instrumentales y equipos medicos, quirurgicos, odontologicos, y hospitalarios para la salud humana y animal. e) La compra, venta, exportacion y distribucion de dulces, confites, galletas y todo tipo de golosinas o alimentos para el consumo humano. f) Actuar como representante o agente de personas naturales y juridicas, nacionales y extranjeras en la realizacion de operaciones o transacciones relacionadas con la industria farmaceutica y en especial con cualquiera de las actividades descritas en los literales a), b), c). d) y e) antes descritos; g) Prestar los servicios de asesoria y consultoria en las areas administrativas, de mercadeo, analisis clinico, asistencia tecnica, investigacion, costos y produccion, exportaciones, importaciones, ventas, relacionadas con iguales o similares actividades a las que constituyen el objeto de PROCAPS S.A., asi como los de publicidad y promocion de productos quimicos, cosmeticos y farmaceuticos para uso humano, animal o industrial; h)-Adquirir, operar, administrar y explotar, a cualquier titulo, empresa o establecimientos dedicados a la investigacion, desarrollo, fabricacion o comercializacion de productos quimicos, farmaceuticos, hospitalarios o cosmeticos. En desarrollo de su objeto principal, la sociedad podra tomar intereses cial como accionista, soci fundador de empresas que tengan igual similar objeto, com-

***** C O N T I N U A *****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----
NIT: 890.106.527-5.

prar, vender, importar y exportar maquinarias, insumos y repuestos requeridos para el desarrollo de sus actividades; tomar dinero en prestamo, financiar y garantizar las operaciones comerciales que celebre con terceros relativas a su objeto; celebrar y ejecutar todo tipo de actos y contratos necesarios para desarrollar sus operaciones; girar, aceptar, otorgar y protestar todo tipo de titulos valores e instrumentos negociables; comprar, vender y constituir gravámenes sobre bienes muebles e inmuebles; designar representantes o apoderados en el pais y en el exterior y, en general, realizar cualquier acto o negocio que sea necesario o conveniente para el desarrollo de su objeto social y para la mejor proteccion de sus intereses corporativos o la de sus socios. La sociedad podra garantizar el cumplimiento de obligaciones adquiridas por terceros o por sus socios, para lo cual podra inclusive constituir gravámenes reales sobre sus bienes de cualquier naturaleza.-----

C E R T I F I C A

CAPITAL	Nro Acciones	Valor Acción
Autorizado		
\$*****2.000.000.000	*****2.000.000	*****1.000
Suscrito		
\$*****1.220.000.000	*****1.220.000	*****1.000
Pagado		
\$*****1.220.000.000	*****1.220.000	*****1.000

C E R T I F I C A

ADMINISTRACION: La direccion y administracion de la sociedad correspondera a la Asamblea General de Accionistas, a la Junta Directiva, al Presidente, al Vicepresidente y a un Gerente General. Corresponde a la Junta Directiva las siguientes funciones y atribuciones entre otras: Autorizar cualquier acto o contrato cuya cuantia sea superior al equivalente en pesos Colombianos a doscientos cincuenta mil dolares de los Estados Unidos de America (US\$250.000), liquidados a la tasa de cambio representativa del mercado vigente el dia en que se surta la transacion. Cuando se trate de operaciones en virtud de las cuales la sociedad garantice obligaciones de terceros o de los socios, se requerira que la operacion sea aprobada con el voto de por lo menos 2/3 partes de los integrantes de la Junta Directiva. La sociedad tendra un Presidente , quien ejercera la representacion legal de la sociedad y tendra las siguientes atribuciones entre otras: Ejecutar las decisiones de la Asamblea General de Accionistas y de la Junta Directiva. Ejercer la Representacion Legal de la sociedad en todos los actos y contratos que se requieran para el desarrollo del objeto social, de conformidad con lo previsto en las leyes y en los presentes estatutos. Constituir apoderados judiciales y extrajudiciales, recibir notificaciones, absolver interrogatorios de parte, confesar obligaciones de la sociedad, transigir, disistir y allanar a la sociedad en cualquier asunto judicial. Aprobar y celebrar en nombre de la sociedad todo acto o contrato cuya cuantia sea igual o inferior al equivalente en pesos colombianos a doscientos cincuenta mil dolares de los

***** C O N T I N U A *****



***** Pag. 3
CAMARA DE COMERCIO
DE BARRANQUILLA
05*****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

Estados Unidos de America (US\$250.000), liquidados a la tasa de cambio representativa del mercado vigente el dia en que se surta la transacion o celebracion del acto. Las demas funciones que le correspondan como Representante Legal por disposicion de la ley, de estos estatutos o de la Junta Directiva. El Presidente tendra un suplente, designado por la Junta Directiva, quien reemplazara al Presidente en sus faltas temporales, absolutas o meramente accidentales, con entidades facultades y sin que sea necesario para ello acreditar la ausencia del principal. La sociedad tendra un Vicepresidente designado por la Junta Directiva, quien tambien ejercera la representacion legal de la sociedad, pudiendo ejercerla conjunta o separadamente con el Presidente. El vicepresidente tendra las siguientes facultades entre otras. Ejecutar dentro de su competencia las decisiones de la Asamblea General de Accionistas, de la Junta Directiva y del Presidente. Ejercer la representacion legal de la sociedad en todos los actos y contratos que se requieran para el desarrollo del objeto social, de conformidad con lo previsto en las leyes y en los presentes estatutos. Constituir apoderados judiciales y extrajudiciales, recibir notificaciones, absolver interrogatorios de parte, confesar obligaciones de la sociedad, transigir, desistir y allanar a la sociedad en cualquier asunto judicial. Aprobar y celebrar en nombre de la sociedad todo acto o contrato cuya cuantia sea igual o inferior al equivalente en pesos colombianos a ciento cincuenta mil dolares de los Estados Unidos de America (US\$150.000), liquidados a la tasa de cambio representativa del mercado vigente el dia en que se surta la aprobacion o celebracion del acto, siempre que no se trate de enajenar bienes inmuebles o cualquier otro activo fijo de la sociedad, inclusive de naturaleza muebles, acciones, cuotas o partes de interes inversiones y todo tipo de derechos de la sociedad, o de constituir gravámenes sobre cualquiera de las anteriores, facultades que radican exclusiva y privativamente en cabeza del Presidente. Las demas funciones que le corresponda como Representante legal por disposicion de la ley, de estos estatutos de la Junta Directiva. El Vicepresidente tendra (1) suplente, designado por la Junta Directiva, quien lo reemplazara conjunta o separadamente en sus faltas temporales, absolutas o meramente accidentales, con identicas facultades y sin que sea necesario para ello acreditar la ausencia del principal. La sociedad tendra un (1) Gerente General, con su respectivo Suplente, designados por la Junta Directiva. El Suplente reemplazara al Gerente General, en sus faltas temporales, absolutas o meramente accidentales, quien junto con el gerente general, llevaran la representacion legal de la sociedad, con identicas facultades, pudiendolas ejercer conjunta separadamente con el Presidente y Vicepresidente, con identicas facultades que el Gerente General y sin que sea necesario para ello acreditar

***** C O N T I N U A *****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

la ausencia del titular. El Gerente General tendra las siguientes facultades entre otras: Ejercer la representacion legal de la sociedad, en todos los actos y contratos que se requieran para el desarrollo del objeto social, de conformidad con lo previsto en las leyes y en los presentes estatutos. Aprobar y celebrar en nombre de la sociedad todo acto o contrato cuya cuantia sea igual o inferior al equivalente en pesos Colombianos a Cincuenta Mil dolares de los Estados Unidos de America (US\$50.000), liquidados a la tasa de cambio representativa del mercado vigente el dia en que se surta la aprobacion o celebracion del acto, siempre que no se trate de enajenar a cualquier titulo bienes inmuebles o cualquier otro activo fijo de la sociedad, inclusive de naturaleza mueble, acciones, cuotas o partes de interes, inversiones y todo tipo de derechos de la sociedad o de constituir gravámenes sobre cualquiera de las anteriores facultades que radican exclusiva y privativamente en cabeza del Presidente.

C E R T I F I C A

Que según Acta No. 116 del 27 de Junio de 1997 correspondiente a la Asamblea de Accionistas en Barranquilla, de la sociedad:

PROCAPS S.A.-----

cuya parte pertinente se inscribió en esta Cámara de Comercio, el 15 de Agosto de 1997 bajo el No. 70.838 del libro respectivo, y según Acta No. 127 del 22 de Marzo de 2000 correspondiente a la Asamblea de Accionistas en Barranquilla, de la sociedad:

PROCAPS S.A.-----

cuya parte pertinente se inscribió en esta Cámara de Comercio, el 29 de Mayo de 2000 bajo el No. 87.250 del libro respectivo, fueron hechos los siguientes nombramientos:

CLASE:

JUNTA DIRECTIVA

Principales

1. Minski Gontovnik Ruben	CC.*****7.463.982
2. Minski Gontovnik Meyer	CC.*****7.461.324
3. Minski Gontovnik Jose	CC.*****8.671.834

Suplentes

1. Minski Zilberblum Elliot	CC.*****72.219.541
2. Minski Deitel Daniel	.*****
3. Minski Sharon	PA.*990.212.082.805

C E R T I F I C A

Que según Acta No. 116 del 27 de Junio de 1997 correspondiente a la Asamblea de Accionistas en Barranquilla, de la sociedad: PROCAPS S.A. cuya parte pertinente se inscribió en esta Cámara de Comercio, el 15 de Agosto de 1997 bajo el No. 70.838 del libro respectivo, fueron hechos los siguientes nombramientos:

Cargo/Nombre	Identificación
Presidente.	
Minski Gontovnik Ruben	CC.*****7.463.982

******* C O N T I N U A *******



***** Pag. 4
CAMARA DE COMERCIO
DE BARRANQUILLA
05*****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

Suplente del Presidente

Minski Minski Abraham

CC.*****802.575

Vicepresidente

Minski Gontovnik Meyer

CC.*****7.461.324

Suplente del vicepresidente

Minski Gontovnik Jose

CC.*****8.671.834

C E R T I F I C A

Que según Acta No. 11 del 01 de Febrero de 2001 correspondiente a la Junta Directiva en Barranquilla, de la sociedad: PROCAPS S.A. cuya parte pertinente se inscribió en esta Cámara de Comercio, el 12 de Febrero de 2001 bajo el No. 91.273 del libro respectivo, fueron hechos los siguientes nombramientos:

Cargo/Nombre

Identificación

Gerente General.

Uribe Ochoa Guillermo Luis

CC.*****70.073.754

C E R T I F I C A

Que según Acta No. 406 del 02 de Sep/bre de 2004 correspondiente a la Junta Directiva en Barranquilla, de la sociedad: PROCAPS S.A. cuya parte pertinente se inscribió en esta Cámara de Comercio, el 09 de Sep/bre de 2004 bajo el No. 113.263 del libro respectivo, fueron hechos los siguientes nombramientos:

Cargo/Nombre

Identificación

Suplente del Gerente

Tanaka Walter

CC.*****16.251.923

C E R T I F I C A

Que según Acta No. 124 del 29 de Marzo de 1999 correspondiente a la Asamblea de Accionistas en Barranquilla, de la sociedad: PROCAPS S.A. cuya parte pertinente se inscribió en esta Cámara de Comercio, el 04 de Agosto de 1999 bajo el No. 82.226 del libro respectivo, fueron hechos los siguientes nombramientos:

Cargo/Nombre

Identificación

Revisor Fiscal Ppal.

Vargas Pertuz Ana Isabel

CC.*****32.696.645

C E R T I F I C A

Que según Acta No. 127 del 22 de Marzo de 2000 correspondiente a la Asamblea de Accionistas en Barranquilla, de la sociedad: PROCAPS S.A. cuya parte pertinente se inscribió en esta Cámara de Comercio,

***** C O N T I N U A *****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----
NIT: 890.106.527-5.

el 29 de Mayo de 2000 bajo el No. 87.250 del libro respectivo, fueron hechos los siguientes nombramientos:

Cargo/Nombre	Identificación
Suplente del Revisor Fiscal.	
Lozada Villareal Milena	CC.*****32.748.111

C E R T I F I C A

Que por escritura publica No. 2.608 del 5 de Octubre del 2.001, torpada en la Notaria 2a. de Barranquilla, inscrita en esta Camara de Comercio el 8 de Noviembre del 2.001, bajo los Nos. 95.771, 2.056, 2.057, 2.058, 2.059, 2.060, 2.061, 2.062 y 2.064 del libro respectivo, se reforma totalmente el articulo 65, contenido en la escritura publica No. 3.705 de Noviembre 6 de 1.998, de la Notaria Segunda del Circuito de Barranquilla, y se constituyen los siguientes Apoderados Generales, quienes tendran las funciones que a continuacion se indican en razon de su cargo en la sociedad: 1.-Se confiere Poder Especial, Amplio y Suficiente a los senores: WALTER TANAKA TATEKAWA, C.C. No. 16.251.923; WILLIAM ALBERTO BARROS CERRA, C.C. No. 8.715.941, para que cada uno en forma independiente en nombre y representacion de PROCAPS S.A., suscriba todas las declaraciones tributarias, aduaneras y cambiarias a que este obligada la sociedad, asi como para que responda todo tipo de requerimiento o solicitudes de informacion de las autoridades tributarias, aduaneras o cambiarias, en los ordenes Distrital, Departamental y Nacional. 2.-Se confiere Poder Especial, amplio y suficiente, a los siguientes senores que conforman el Grupo A y el Grupo B para que conjuntamente suscriban contratos de suministros de mercancías y servicios con entidades oficiales, hasta por la suma de US\$800.000.00 (OCHOCIENTOS MIL DOLARES DE LOS ESTADOS UNIDOS DE AMERICA), liquidados a la tasa representativa del mercado vigente en el dia que se realice la respectiva transaccion. GUPO A: GUILLERMO LUIS URIBE OCHO C.C. No. 70.073.754, LINO LAZALA SILVA VARGAS C.C. No. 19.260.410, JAIRO MOLINA MEJIA C.C. No. 8.706.380, CLAUDIA TANGARIFE CASTILLO C.C. No. 41.660.293. BRUPO B: WALTER TANAKA TATEKAWA C.C. No. 16.251.923, HENRY CARO DIAZ C.C. No. 19.395.298, CARMEN ALICIA FLOREZ C.C. No. 22.434.044, CARMEN ALICIA HERRERA DIAZGRANADOS C.C. No. 22.377.540. Para el ejercicio de estas facultades se requiera por lo menos la firma de uno (1) de los apoderados especiales del Grupo A y uno de los apoderados especiales del Grupo B; 3.-Se otorga Poder General, Amplio y Suficiente a la Doctora JOSEFINA MIELES BUELVAS C.C. No. 33.138.497, a la Doctora MARCELA CARVAJALINO PAGANO C.C. No. 22.579.748, y al Doctor DANIEL MORENO VILLALBA, C.C. No. 19.113.210; para que cada uno representen a PROCAPS S.A., ante todo tipo de autoridades administrativas y judiciales, en asuntos de naturaleza cambiaria y fiscal, tales como: Contestar e interponer demandas y requerimientos, comparecer en juicios, pedir la practica de pruebas, recibir notificaciones, interponer recursos y ejercer acciones contencioso administrativas y de tutela, transigir, conciliar, recibir en nombre de la sociedad desistir, sustituir y reasumir este mandato. Para la celebracion de cualquier acuerdo conciliatorio o transaccional, judicial o extrajudicial, cualquiera que sea su cuantía, los Apoderados deberan aportar la aut rizacion escrita del Representante Legal de la socie-

***** C O N T I N U A *****



***** Pag. 5
CAMARA DE COMERCIO
DE BARRANQUILLA
05*****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

dad. Los Apoderados podran presentar solicitudes de devoluci n y/o compensacion de impuestos y recibir los pagos por devoluciones de impuestos, firmar, notificarse y realizar todo acto necesari para cumplir con el mandato encomendado.-----

C E R T I F I C A

Que segun documento privado de fecha 7 de junio de 2000, inscrito en esta Camara de Comercio el 8 de junio de 2000, bajo el No.1.786 del libro respectivo, consta que RUBEN MINSKI GONTOVNIK, C.C. No. 7.463.982 de Barranquilla, actuando como Presidente y Representante Legal de la sociedad PROCAPS S.A, confiere poder especial, amplio y suficiente a KAREN FIGUEROA GARCIA, C.C.No.32.732.070 de Barranquilla, para que en su nombre y representacion de la sociedad suscriba todas declaraciones cambiarias a que este obligada PROCAPS S.A, en el giro normal de sus negocios y operaciones sociales.--

C E R T I F I C A

Que segun Documento Privado del 13 de diciembre del 2.002, inscrito en esta Camara de Comercio el 15 de enero del 2.003 bajo el No.2.290 del libro respectivo, el senor RUBEN MINSKI GONTOVNIK, C.C.No.7.463.982, en calidad de REPRESENTANTE LEGAL, y en condicion de PRESIDENTE de PROCAPS S.A., mediante el presente documento otorgo PODER ESPECIAL al Sr. MARLON TATIS PAREDES, C.C.No.72.176.874, para que en representacion de PROCAPS S.A., suscriba declaraciones, rep rtes y cualquier tipo de documentos relacionados con tramites o registros de Comercio Exterior' ante el Ministerio de Comercio Exteri r y/ cualquier entidad del Estado que haga sus veces. En consecuencia, el Sr. MARLON TATIS PAREDES, podra suscribir en representaci n de PROCAPS S.A. declaraciones, reportes y cualquier tipo de document s relacionados con tramites o registros de Comercio Exterior ante el Ministerio de Comercio Exterior y/o cualquier entidad del Estado que haga sus veces.-----

C E R T I F I C A

Que segun documento privado del 10 de noviembre del 2004, inscrito en esta Camara de Comercio el 1 de diciembre del 2.004, bajo el No. 2.753 del libro respectivo consta que RUBEN MINSNSKI GONTOVNIK, C.C. 7.463.982 de Barranquilla, actuando como Presidente y Representante Legal de la sociedad PROCAPS S.A, confiere poder especial, amplio y suficiente a ANTONIO VARELA DIAZ, C.C.No.72.004.669 de Barranquilla, para que en nombre y representacion de la s ciedad firme las declaraciones cambiarias a que este obligada PROCAPS S.A., a

***** C O N T I N U A *****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

suscribir en el giro normal de sus negocios y operaciones sociales. Por lo anterior, el señor ANTONIO VARELA DIAZ, queda facultado para firmar las declaraciones comabiarías y demás documentos relacionados con las declaraciones, notificaciones, aportar y solicitar las pruebas que sean necesarias, desistir en general a realizar todo acto necesario de tal forma que pueda cumplir con el mandato dado.-

C E R T I F I C A

Que según documento privado del 22 de Febrero de 2005, inscrito en esta Cámara de Comercio el 28 de Marzo de 2.005, bajo el No. 2.820 del libro respectivo, consta que WALTER TANAKA TATEKAWA C.C. 16.251-923 de Palmira, quien actúa como representante legal de la sociedad PROCAPS S.A, otorga Poder Especial, Amplio y Suficiente al señor MARLON TATIS PAREDES C.C. No. 72.176.874, para que en representación de PROCAPS S.A., suscriba declaraciones aduaneras, reportes y cualquier tipo de documentos relacionados con los trámites o registros de comercio exterior ante la dirección de impuestos y aduanas nacionales y/o cualquier entidad que haga sus veces, y para que responda todo tipo de requerimientos y/o solicitudes de información de las autoridades aduaneras en los órdenes distrital, departamental y nacional.-----

C E R T I F I C A

Que por Escritura Pública No.1.613 del 28 de Junio del 2.005, otorgada en la Notaría Segunda de Barranquilla, inscrita en esta Cámara de Comercio el 18 de agosto del 2.005 bajo el No.2.913 del libro respectivo, consta que RUBEN MINSKI GONTOVNIK, C.C.No.7.463.928, en su condición de Presidente y Representante Legal de la sociedad PROCAPS S.A., confiere poder especial amplio y suficiente a los siguientes señores que conforman el Grupo A y el Grupo B, para que conjuntamente suscriban contratos de suministro de mercancías y servicios con entidades oficiales, hasta por la suma de OCHOCIENTOS MIL DOLARES DE LOS ESTADOS DE AMERICA (US\$800.000), liquidados a la tasa representativa del mercado vigente en el día que se realice la respectiva transacción. GRUPO A: MIRIAM RODRIGUEZ DIAZ, C.C.No. 41.473.569. GRUPO B: NELSON SUAREZ CHAVEZ, C.C.No.79.417.971.-----

C E R T I F I C A

Que en esta Cámara de Comercio no aparecen inscripciones posteriores de documentos referentes a reforma, disolución, liquidación o nombramientos de representantes legales de la expresada sociedad.

C E R T I F I C A

Que su última Renovación fue el: 29 de Marzo de 2006.

La información sobre embargos de establecimiento se suministra en Certificados de Matrícula, la de contratos sujetos a registro, en Certificados Especiales.



***** Pag. 6
C A M A R A D E C O M E R C I O
D E B A R R A N Q U I L L A
05*****

CERTIFICADO DE EXISTENCIA Y REPRESENTACION LEGAL O DE INSCRIPCION DE DOCUMENTOS.

PROCAPS S.A.-----

NIT: 890.106.527-5.

m. fajon

ANEXO 1

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 April 2003 (03.04.2003)

PCT

(10) International Publication Number
WO 03/026659 A1

(51) International Patent Classification⁷: A61K 31/496,
C07D 215/22, A61P 25/18

(21) International Application Number: PCT/JP02/09858

(22) International Filing Date:
25 September 2002 (25.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2001-290645 25 September 2001 (25.09.2001) JP
2001-348276 14 November 2001 (14.11.2001) JP
2,379,005 27 March 2002 (27.03.2002) CA

(71) Applicant (for all designated States except US): OT-
SUKA PHARMACEUTICAL CO., LTD. [JP/JP];
9, Kanda-Tsukasa-cho 2-chome, Chiyoda-ku, Tokyo
101-8535 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BANDO, Takuji
[JP/JP]; 31-3, Minamishomachi 2-chome, Tokushima-shi,
Tokushima 770-0045 (JP). AOKI, Satochi [JP/JP];
132-6, Muiyachokurotsuki Miyazu, Naruto-shi, Tokushima
772-0001 (JP). KAWASAKI, Junichi [JP/JP]; 112-2,
Tainohama Aza Kaya, Kitajimacho, Itano-gun, Tokushima
771-0204 (JP). ISHIGAMI, Makoto [JP/JP]; 61-3, Shozui
Higashikatsuchi, Aizumicho, Itano-gun, Tokushima
771-1273 (JP). TANIGUCHI, Youichi [JP/JP]; 5-70-701,
Sundiyoshi 4-chome, Tokushima-shi, Tokushima 770-0861
(JP). YABUUCHI, Tsuyoshi [JP/JP]; 900-25, Kawachi-
cho Omatsu, Tokushima-shi, Tokushima 771-0131 (JP).
FUJIMOTO, Kiyoshi [JP/JP]; 43-74, Otsuchoyagura
Aza Rokunokoshi, Naruto-shi, Tokushima 772-0035 (JP).
NISHIOKA, Yoshihiro [JP/JP]; 5-3, Nakagiri Aza
Ushikadnotigashinokoshi, Matsushigecho, Itano-gun,
Tokushima 771-0212 (JP). KOBAYASHI, Noriyuki
[JP/JP]; 21-5, Nakashowacho 3-chome, Tokushima-shi,
Tokushima 770-0944 (JP). FUJIMURA, Tadamu
[JP/JP]; 71, Otsuchoyagura Aza Kita, Naruto-shi,
Tokushima 772-0035 (JP). TAKAHASHI, Masanori

[JP/JP]; 7-1, Sangenyachosoto, Tokushima-shi, Tokushima
770-8025 (JP). ABE, Kaoru [JP/JP]; 2-23-808, Sue-
hiro 1-chome, Tokushima-shi, Tokushima 770-0866
(JP). NAKAGAWA, Tomonori [JP/JP]; Grace78-205,
78, Sasagino Aza Hachiyamakaitaku, Matsushigecho,
Itano-gun, Tokushima 771-0219 (JP). SHINAHAMA,
Kolehi [JP/JP]; 9-15, Kitajosanjimacho 3-chome,
Tokushima-shi, Tokushima 770-0812 (JP). UTSUMI,
Naoto [JP/JP]; B202, Muiyacho Benzaiten Aza Mitsudo,
Naruto-shi, Tokushima 772-0014 (JP). TOMINAGA,
Michiaki [JP/JP]; 310-6, Takaiso, Kamitacho, Itano-gun,
Tokushima 771-1346 (JP). OI, Yoshihiro [JP/JP]; 8-33,
Kitatamiya 1-chome, Tokushima-shi, Tokushima 770-0003
(JP). YAMADA, Shobei [JP/JP]; 5-11, Tomiyoshi Aza
Toyoyoshi, Aizumicho, Itano-gun, Tokushima 771-1231
(JP). TOMIKAWA, Kenji [JP/JP]; 31-11, Kokufuchoido
Aza Maeno, Tokushima-shi, Tokushima 779-3118 (JP).

(74) Agents: ASAMURA, Kiyoshi et al.; Room 331, New
Ohtemachi Bldg., 2-1, Ohtemachi 2-chome, Chiyoda-ku,
Tokyo 100-0004 (JP).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GR, GH,
GM, HR, HU, ID, IL, IN, IS, KE, KG, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
European patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CL, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: LOW HYGROSCOPIC ARIPIPRAZOLE DRUG SUBSTANCE AND PROCESSES FOR THE PREPARATION THEREOF

(57) Abstract: The present invention provides low hygroscopic forms of aripiprazole and processes for the preparation thereof which will not convert to a hydrate or lose their original solubility even when a medicinal preparation containing the aripiprazole anhydride crystals is stored for an extended period.



WO 03/026659 A1

- 1 -

DESCRIPTION

LOW HYGROSCOPIC ARIPIPRAZOLE DRUG SUBSTANCE AND
PROCESSES FOR THE PREPARATION THEREOF

DETAILED DESCRIPTION OF THE INVENTION

Field of the Invention

The present invention relates to an improved form of aripiprazole having reduced hygroscopicity and processes for the preparation of this improved form.

Background of the Invention

Aripiprazole, 7-(4-[4-(2,3-dichlorophenyl)-1-piperazinyl]-butoxy)-3,4-dihydro carbostyryl or 7-(4-[4-(2,3-dichlorophenyl)-1-piperazinyl]-butoxy)-3,4-dihydro-2(1H)-quinolinone, is an atypical antipsychotic agent useful for the treatment of schizophrenia (U.S. 4,734,416 and U.S. 5,006,528). Schizophrenia is a common type of psychosis characterized by delusions, hallucinations and extensive withdrawal from others. Onset of schizophrenia typically occurs between the age of 16 and 25 and affects 1 in 100 individuals worldwide. It is more prevalent than Alzheimer's disease, multiple sclerosis, insulin-dependent diabetes and muscular dystrophy. Early diagnosis and treatment can lead to significantly improved recovery and outcome. Moreover, early therapeutic intervention can avert costly hospitalization.

20
154

- 2 -

According to Example 1 of Japanese Unexamined Patent Publication No. 191256/1990, aripiprazole anhydride crystals are manufactured for example by reacting 7-(4-bromobutoxy)-3,4-dihydrocarbostyryl with 1-(2,3-dichlorophenylpiperadine and recrystallizing the resulting raw aripiprazole anhydride with ethanol. Also, according to the Proceedings of the 4th Japanese-Korean Symposium on Separation Technology (October 6-8, 1996), aripiprazole anhydride crystals are manufactured by heating aripiprazole hydrate at 80°C. However, the aripiprazole anhydride crystals obtained by the aforementioned methods have the disadvantage of being significantly hygroscopic.

The hygroscopicity of these crystals makes them difficult to handle since costly and burdensome measures must be taken in order ensure they are not exposed to moisture during process and formulation. Exposed to moisture, the anhydrous form can take on water and convert to a hydrous form. This presents several disadvantages. First, the hydrous forms of aripiprazole have the disadvantage of being less bioavailable and less dissoluble than the anhydrous forms of aripiprazole. Second, the variation in the amount of hydrous versus anhydrous aripiprazole drug substance from batch to batch could fail to meet specifications set by drug regulatory agencies. Third, the milling may cause the drug substance, Conventional Anhydride, to adhere to manufacturing equipment which

31
BS

- 3 -

may further result in processing delay, increased operator involvement, increased cost, increased maintenance and lower production yield. Fourth, in addition to problems caused by introduction of moisture during the processing of these hygroscopic anhydrides, the potential for absorbance of moisture during storage and handling would adversely affect the dissolubility of aripiprazole drug substance. Thus shelf-life of the product could be significantly decreased and/or packaging costs could be significantly increased. It would be highly desirable to discover a form of aripiprazole that possessed low hygroscopicity thereby facilitating pharmaceutical processing and formulation operations required for producing dosage units of an aripiprazole medicinal product having improved shelf-life, suitable dissolubility and suitable bioavailability.

Also, Proceedings of the 4th Japanese-Korean Symposium on Separation Technology (October 6-8, 1996) state that, aripiprazole anhydride crystals exist as type-I crystals and type-II crystals; the type-I crystals of aripiprazole anhydride can be prepared by recrystallizing from an ethanol solution of aripiprazole, or by heating aripiprazole hydrate at 80°C; and the type-II crystals of aripiprazole anhydride can be prepared by heating the type-I crystals of aripiprazole anhydride at 130 to 140°C for 15 hours.

- 4 -

By the aforementioned methods, aripiprazole anhydride type-II crystals having high purity can not be easily prepared in an industrial scale with good repeatability.

5

SUMMARY OF THE INVENTION

Thus according to the present invention is provided a form of aripiprazole having reduced hygroscopicity and which is more amenable to pharmaceutical processing and formulation. The inventors of the present invention have discovered that this reduced-hygroscopic form of Aripiprazole is a crystalline substance defined herein as Anhydride B. A particular process for the preparation of this anhydrous crystalline substance has also been discovered and comprises yet another aspect of the present invention. Particularly, it was discovered as part of the present invention that in order to produce Anhydride B having the desired pharmaceutical properties and utilizing the most efficient process, Hydrate A, as defined herein, would have to serve as the intermediate. It was also discovered that a particular sequence of processing had to be implemented in order to form Hydrate A. It was discovered that the preparation of Hydrate A required milling what is defined herein as Conventional Hydrate. Then, Hydrate A can be transformed into Anhydride B through suitable heating as defined herein. Surprisingly, if the

10

15

20

25

72
150

23
157

- 5 -

Conventional Hydrate is first heated and then milled, serious agglomeration sets in rendering the processing commercially unsuitable.

An object of the present invention is to
5 provide novel aripiprazole anhydride crystals.

Moreover, another object of the present invention is to provide aripiprazole anhydride crystals which neither easily convert into hydrates nor substantially decrease the original solubility, even
10 when a pharmaceutical composition comprising aripiprazole anhydride is stored for a long period of time.

Further object of the present invention is to provide preparation methods, in order to obtain
15 aripiprazole anhydride crystals having high purity in an industrial scale with good repeatability.

The present inventors have conducted research works aimed to attain the aforementioned objects. In the course of the research, they have found that the
20 desired aripiprazole anhydride crystals can be obtained when a well-known aripiprazole anhydride is heated at the specific temperature. Further, the present inventors have found that the desired aripiprazole anhydride crystals can be obtained from
25 recrystallization of a well-known aripiprazole anhydride by using the specific solvents. Moreover, the present inventors found that the desired aripiprazole anhydride crystals can be obtained by

150 24

- 6 -

suspending a well-known aripiprazole anhydride in the specific solvent, and heating thus obtained suspension.

The present invention thus completed on the basis of these findings and knowledge.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a thermogravimetric/differential thermogram of the Aripiprazole Hydrate A obtained in Example 1.

10

Figure 2 shows the ^1H -NMR spectrum ($\text{DMSO}-d_6$, TMS) of the Aripiprazole Hydrate A obtained in Example 1.

Figure 3 is a powder x-ray diffraction diagram of the Aripiprazole Hydrate A obtained in

15

Example 1.

Figure 4 shows the ^1H -NMR spectrum ($\text{DMSO}-d_6$, TMS) of the Aripiprazole Anhydride Crystals B obtained in Example 2.

Figure 5 is a powder x-ray diffraction diagram of the Aripiprazole Anhydride Crystals B obtained in Example 2.

20

Figure 6 is a thermogravimetric/differential thermogram of the aripiprazole hydrate obtained in Reference Example 3.

25

Figure 7 is a powder x-ray diffraction diagram of the aripiprazole hydrate obtained in Reference Example 3.

Figure 8 shows thermogravimetric/differential

- 7 -

thermal analysis endothermic curve of the type C crystals of aripiprazole anhydride obtained in Example 11.

Figure 9 shows an ^1H -NMR spectrum (DMSO- d_6 , TMS) of the type C crystals of aripiprazole anhydride obtained in Example 11.

Figure 10 shows a powder X-ray diffraction spectrum of the type C crystals of aripiprazole anhydride obtained in Example 11.

Figure 11 shows an IR spectrum of the type C crystals of aripiprazole anhydride obtained in Example 11.

Figure 12 shows a solid ^{13}C -NMR spectrum of the type C crystals of aripiprazole anhydride obtained in Example 11.

Figure 13 shows a thermogravimetric/differential thermal analysis endothermic curve of the type D crystals of aripiprazole anhydride obtained in Example 12 or Example 13.

Figure 14 shows an ^1H -NMR spectrum (DMSO- d_6 , TMS) of the type D crystals of aripiprazole anhydride obtained in Example 12 or Example 13.

Figure 15 shows a powder X-ray diffraction spectrum of the type D crystals of aripiprazole anhydride obtained in Example 12 or Example 13.

Figure 16 shows an IR spectrum of the type D crystals of aripiprazole anhydride obtained in Example 12 or Example 13.

26
160

- 8 -

Figure 17 shows a solid ^{13}C -NMR spectrum of the type D crystals of aripiprazole anhydride obtained in Example 12 or Example 13.

Figure 18 shows a thermogravimetric/
5 differential thermal analysis endothermic curve of the type E crystals of aripiprazole anhydride obtained in Example 14.

Figure 19 shows an ^1H -NMR spectrum (DMSO-d_6 , TMS) of the type E crystals of aripiprazole anhydride
10 obtained in Example 14.

Figure 20 shows a powder X-ray diffraction spectrum of the type E crystals of aripiprazole anhydride obtained in Example 14.

Figure 21 shows an IR spectrum of the type E
15 crystals of aripiprazole anhydride obtained in Example 14.

Figure 22 shows a thermogravimetric/
differential thermal analysis endothermic curve of the type F crystals of aripiprazole anhydride obtained in
20 Example 15.

Figure 23 shows an ^1H -NMR spectrum (DMSO-d_6 , TMS) of the type F crystals of aripiprazole anhydride obtained in Example 15.

Figure 24 shows a powder X-ray diffraction
25 spectrum of the type F crystals of aripiprazole anhydride obtained in Example 15.

Figure 25 shows an IR spectrum of the type F crystals of aripiprazole anhydride obtained in Example

77
61

- 9 -

15.

Figure 26 shows thermogravimetric/
differential thermal analysis endothermic curve of the
type G crystals of aripiprazole anhydride obtained in
5 Example 16-b).

Figure 27 shows an ¹H-NMR spectrum (DMSO-d₆,
TMS) of the type G crystals of aripiprazole anhydride
obtained in Example 16-b).

Figure 28 shows a powder X-ray diffraction
10 spectrum of the type G crystals of aripiprazole
anhydride obtained in Example 16-b).

Figure 29 shows an IR spectrum of the type G
crystals of aripiprazole anhydride obtained in Example
16-b).

15 Figure 30 shows a thermogravimetric/
differential thermal analysis endothermic curve of the
glass form of aripiprazole anhydride obtained in
Example 16-a).

Figure 31 shows a powder X-ray diffraction
20 spectrum of the glassy state of aripiprazole anhydride
obtained in Example 16-a).

DETAILED DESCRIPTION OF THE INVENTION

According to first embodiment of the first
25 aspect of the present invention is provided Hydrate A
of aripiprazole wherein said Hydrate has a powder x-ray
diffraction spectrum which is substantially the same as
the powder x-ray diffraction spectrum shown in Figure

- 10 -

3.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has powder x-ray diffraction characteristic peaks at $2\theta = 12.6^\circ$, 15.4° , 17.3° , 18.0° , 18.6° , 22.5° and 24.8° .

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has particular infrared absorption bands at 2951, 2822, 1692, 1577, 1447, 1378, 1187, 963 and 784 cm^{-1} on the IR (KBr) spectrum.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has an ^1H -NMR spectrum which is substantially the same as the ^1H -NMR spectrum (DMSO-d_6 , TMS) shown in Figure 2.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has an ^1H -NMR spectrum (DMSO-d_6 , TMS) having characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, $J = 7.4\text{ Hz}$, 2H), 2.97 ppm (brt, $J = 4.6\text{ Hz}$, 4H), 3.92 ppm (t, $J = 6.3\text{ Hz}$, 2H), 6.43 ppm (d, $J = 2.4\text{ Hz}$, 1H), 6.49 ppm (dd, $J = 8.4\text{ Hz}$, $J = 2.4\text{ Hz}$, 1H), 7.04 ppm (d, $J = 8.1\text{ Hz}$, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

- 11 -

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has an endothermic curve which is substantially the same as the thermogravimetric/differential thermal analysis (heating rate 5°C/min) endothermic curve shown in Figure 1.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size of 50 µm or less.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size of 40 µm or less.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size of 35 µm or less.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size of 30 µm or less.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size of 25 µm or less.

According to another embodiment of the first

76
164

- 12 -

aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size of 20 μm or less.

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size range of 40 to 10 μm .

According to another embodiment of the first aspect of the present invention is provided Hydrate A of aripiprazole wherein said Hydrate has a mean particle size range of 36 to 14 μm .

According to a second aspect of the present invention is provided a process for the preparation of Hydrate A wherein said process comprises the steps of milling Conventional Hydrate.

According to a first embodiment of the second aspect of the present invention is provided a process for the preparation of Hydrate A comprising the steps of milling Conventional Hydrate wherein said milling is performed by a milling machine.

According to another embodiment of the second aspect of the present invention is provided a process for the preparation of Hydrate A comprising the steps of milling Conventional Hydrate wherein said milling machine is an atomizer, pin mill, jet mill or ball mill.

According to another embodiment of the second aspect of the present invention is provided a process

165 31

- 13 -

for the preparation of Hydrate A comprising the steps of milling Conventional Hydrate wherein said milling machine is an atomizer.

According to another embodiment of the second aspect of the present invention is provided a process for the preparation of Hydrate A comprising the steps of milling Conventional Hydrate wherein said milling machine is an atomizer using a rotational speed of 5000-15000 rpm for the main axis, a feed rotation of 10-30 rpm and a screen hole size of 1-5 mm.

According to various embodiments of a third aspect of the present invention is provided Hydrate A defined according to one or more of the embodiments described herein wherein said Hydrate is made by a process as described herein.

According to a fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity.

According to a first embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.5% or less after placing said drug substance for 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

According to a first embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity

32
166

- 14 -

wherein said low hygroscopicity is a moisture content of 0.4% or less after placing said drug substance for 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

5 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.25% or less after placing said drug substance for
10 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity
15 wherein said low hygroscopicity is a moisture content of 0.15% or less after placing said drug substance for 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

 According to another embodiment of the fourth
20 aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.10% or less after placing said drug substance for 24 hours in a dessicator maintained at a temperature of
25 60°C and a humidity level of 100%.

 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity

33
67

- 15 -

wherein said low hygroscopicity is a moisture content of 0.05% or less after placing said drug substance for 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

5 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.04% or less after placing said drug substance for
10 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity
15 wherein said drug substance is Aripiprazole Anhydride Crystals B as defined herein.

 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity
20 wherein said drug substance has a powder x-ray diffraction spectrum which is substantially the same as the powder x-ray diffraction spectrum shown in Figure 5.

 According to another embodiment of the fourth
25 aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said drug substance has a powder x-ray diffraction spectrum having characteristic peaks at $2\theta =$

34
168

- 16 -

11.0°, 16.6°, 19.3°, 20.3° and 22.1°.

According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity

5 wherein said drug substance has particular infrared absorption bands at 2945, 2812, 1678, 1627, 1448, 1377, 1173, 960 and 779 cm^{-1} on the IR (KBr) spectrum.

According to another embodiment of the fourth aspect of the present invention is provided

10 aripiprazole drug substance of low hygroscopicity wherein said drug substance has an ^1H -NMR spectrum which is substantially the same as the ^1H -NMR spectrum (DMSO-d_6 , TMS) shown in Figure 4.

According to another embodiment of the fourth aspect of the present invention is provided

15 aripiprazole drug substance of low hygroscopicity wherein said drug substance has an ^1H -NMR spectrum (DMSO-d_6 , TMS) having characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, $J = 7.4$ Hz, 2H), 2.97 ppm (brt, $J = 4.6$ Hz, 4H), 3.92 ppm (t, $J = 6.3$ Hz, 2H), 6.43 ppm (d, $J = 2.4$ Hz, 1H), 6.49 ppm (dd, $J = 8.4$ Hz, $J = 2.4$ Hz, 1H), 7.04 ppm (d, $J = 8.1$ Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H)
20 and 10.00 ppm (s, 1H).
25

According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity

83
169

- 17 -

wherein said drug substance exhibits an endothermic peak near about 141.5°C in thermogravimetric/differential thermal analysis (heating rate 5°C/min).

According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said drug substance exhibits an endothermic peak near about 140.7°C in differential scanning calorimetry (heating rate 5°C/min).

10 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said drug substance is Aripiprazole Anhydride Crystals B and will not substantially convert to a
15 hydrous form of aripiprazole when properly stored even for an extended period. For instance, said Aripiprazole Anhydride Crystals B can be stored under a relative humidity (RH) of 60 % and at a temperature of 25°C, even for a period not less than 1 year.

20 According to another embodiment of the fourth aspect of the present invention is provided aripiprazole drug substance of low hygroscopicity wherein said drug substance is Aripiprazole Anhydride Crystals B and will not substantially convert to a
25 hydrous form of aripiprazole when properly stored even for an extended period. For instance, said Aripiprazole Anhydride Crystals B can be stored under a relative humidity (RH) of 60 % and at a temperature of

36
170

- 18 -

25°C, even for a period not less than 4 years.

According to another embodiment of the fourth aspect of the present invention is provided

aripiprazole drug substance of low hygroscopicity

- 5 wherein said drug substance is Aripiprazole Anhydride Crystals B and will not substantially convert to a hydrous form of aripiprazole when properly stored even for a period not less than 0.5 year under a relative humidity (RH) of 75 % and at a temperature of 40°C.

- 10 According to another embodiment of the fourth aspect of the present invention is provided

aripiprazole drug substance of low hygroscopicity

wherein said drug substance has a mean size of 50µm or less when small particle size is required for the

- 15 formulation such as Tablet and other solid dose formulations including for example flashmelt formulations.

According to another embodiment of the fourth aspect of the present invention is provided

- 20 aripiprazole drug substance of low hygroscopicity

wherein said drug substance has a mean size of 40µm or less if small particle size is required for the

formulation such as Tablet and other solid dose formulations including for example flashmelt

- 25 formulations.

According to another embodiment of the fourth aspect of the present invention is provided

aripiprazole drug substance of low hygroscopicity

37
171

- 19 -

wherein said drug substance has a mean size of 30µm or less if small particle size is required for formulation such as Tablet and other solid dose formulations including for example flashmelt formulations.

5 According to a fifth aspect of the present invention is provided a process for the preparation of Aripiprazole Anhydride Crystals B.

 According to a first embodiment of the fifth aspect of the present invention is provided a process
10 for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A.

 According to a first embodiment of the fifth aspect of the present invention is provided a process
15 for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 90-125°C for about 3-50 hours.

 According to another embodiment of the fifth aspect of the present invention is provided a process
20 for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 100°C for about 18 hours.

 According to another embodiment of the fifth aspect of the present invention is provided a process
25 for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 100°C for about 24 hours.

 According to another embodiment of the fifth

~~28~~
172

- 20 -

aspect of the present invention is provided a process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 120°C for about 3 hours.

5 According to another embodiment of the fifth aspect of the present invention is provided a process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A for about 18 hours at 100°C followed by
10 additional heating for about 3 hours at 120°C.

 According to a sixth aspect of the present invention is provided Aripiprazole Anhydride Crystals B defined according to one or more of the embodiments described herein and made by a process as provided
15 herein.

 According to a seventh aspect of the present invention is provided Aripiprazole Anhydride Crystals B formulated with one or more pharmaceutically acceptable carriers.

20 Other embodiments of the present invention may comprise suitable combinations of two or more of the embodiments and/or aspects disclosed herein.

 Yet other embodiments and aspects of the invention will be apparent according to the description
25 provided below.

 Yet another aspect of the present invention comprised discovering that when aripiprazole hydrate (Conventional Hydrate as defined herein) is milled, it

39
123

- 21 -

converts to an aripiprazole hydrate (Hydrate A as defined herein) with a different powder x-ray diffraction spectrum by different peak intensities. Moreover, it was found that Hydrate A loses the sharp
5 dehydration endothermic peak of 123.5°C which characterizes unmilled Conventional Hydrate in thermogravimetric/differential thermal analysis. Thus, the Conventional Hydrate is transformed into Hydrate A after milling Conventional Hydrate and exhibits a
10 gradual dehydration endothermic peak between about 60°C and 120°C with a weak peak at about 71°C.

Yet another aspect of the invention comprised discovering that when heated to a specific temperature of 90-125°C for 3-50hr, this novel aripiprazole hydrate
15 dehydrates gradually avoiding the aggregation phenomenon thought to be caused in conventional aripiprazole hydrate by rapid dehydration, and that aripiprazole anhydride crystals obtained by heating of the novel aripiprazole hydrate to a specific
20 temperature are aripiprazole anhydride crystals with the desired properties.

Characterization of Hydrate A

Particles of "Hydrate A" as used herein have
25 the physicochemical properties given in (1)-(5) below:

(1) It has an endothermic curve which is substantially the same as the thermogravimetric/differential thermal analysis (heating rate 5°C/min)

46
174

- 22 -

endothermic curve shown in Figure 1. Specifically, it is characterized by the appearance of a small peak at about 71°C and a gradual endothermic peak around 60°C to 120°C.

5 (2) It has an ¹H-NMR spectrum which is substantially the same as the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 2. Specifically, it has characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm
10 (m, 4H + DMSO), 2.78 ppm (t, J = 7.4 Hz, 2H), 2.97 ppm (brt, J = 4.6 Hz, 4H), 3.92 ppm (t, J = 6.3 Hz, 2H), 6.43 ppm (d, J = 2.4 Hz, 1H), 6.49 ppm (dd, J = 8.4 Hz, J = 2.4 Hz, 1H), 7.04 ppm (d, J = 8.1 Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm
15 (s, 1H).

(3) It has a powder x-ray diffraction spectrum which is substantially the same as the powder x-ray diffraction spectrum shown in Figure 3. Specifically, it has characteristic peaks at 2θ = 12.6°,
20 15.4°, 17.3°, 18.0°, 18.6°, 22.5° and 24.8°.

(4) It has clear infrared absorption bands at 2951, 2822, 1692, 1577, 1447, 1378, 1187, 963 and 784 cm⁻¹ on the IR (KBr) spectrum.

(5) It has a mean particle size of 50 μm or
25 less.

Process for Manufacturing Hydrate A

Hydrate A is manufactured by milling

41
125

- 23 -

Conventional Hydrate. Conventional milling methods can be used to mill Conventional Hydrate. For example, Conventional Hydrate can be milled in a milling machine. A widely used milling machine can be used, such as an atomizer, pin mill, jet mill or ball mill. Of these, the atomizer is preferred.

Regarding the specific milling conditions when using an atomizer, a rotational speed of 5000-15000 rpm could be used for the main axis, for example, with a feed rotation of 10-30 rpm and a screen hole size of 1-5 mm.

The mean particle size of the Aripiprazole Hydrate A obtained by milling should normally be 50 μm or less, preferably 30 μm or less. Mean particle size can be ascertained by the particle size measurement method described hereinafter.

Characterization of Aripiprazole Anhydride Crystals B

"Aripiprazole Anhydride Crystals B" of the present invention as used herein have the physicochemical properties given in (6)-(12) below.

(6) They have an ^1H -NMR spectrum which is substantially the same as the ^1H -NMR spectrum (DMSO- d_6 , TMS) shown in Figure 4. Specifically, they have characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, $J = 7.4$ Hz, 2H), 2.97 ppm (brt, $J = 4.6$ Hz, 4H), 3.92 ppm (t, $J = 6.3$ Hz, 2H),

42
176

- 24 -

6.43 ppm (d, $J = 2.4$ Hz, 1H), 6.49 ppm (dd, $J = 8.4$ Hz, $J = 2.4$ Hz, 1H), 7.04 ppm (d, $J = 8.1$ Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

5 (7) They have a powder x-ray diffraction spectrum which is substantially the same as the powder x-ray diffraction spectrum shown in Figure 5. Specifically, they have characteristic peaks at $2\theta = 11.0^\circ$, 16.6° , 19.3° , 20.3° and 22.1° .

10 (8) They have clear infrared absorption bands at 2945, 2812, 1678, 1627, 1448, 1377, 1173, 960 and 779 cm^{-1} on the IR (KBr) spectrum.

 (9) They exhibit an endothermic peak near about 141.5°C in thermogravimetric/differential thermal
15 analysis (heating rate $5^\circ\text{C}/\text{min}$).

 (10) They exhibit an endothermic peak near about 140.7°C in differential scanning calorimetry (heating rate $5^\circ\text{C}/\text{min}$).

 (11) Aripiprazole Anhydride Crystals B of the
20 present invention have low hygroscopicity. For example, Aripiprazole Anhydride Crystals B of the present invention maintain a water content of 0.4% or less after 24 hours inside a dessicator set at a temperature of 60°C and a humidity of 100%. Well-known
25 methods of measuring water content can be used as long as they are methods commonly used for measuring the water content of crystals. For example, a method such as the Karl Fischer method can be used.

- 25 -

(12) When the small particle size is required for the formulation such as tablet and other solid dose formulations including for example flashmelt formulations, the mean particle size is preferably 50
5 μm or less.

Process for Manufacturing Anhydride B

In case of the formulation for which small particle size (less than 50 μm) is required, the
10 milling is necessary for the preparation. However, when a large amount of Conventional Aripiprazole Anhydride or Anhydride Crystals B having large particle size is milled, the milled substances adhere with each other in the milling machine. Accordingly, there is a
15 disadvantage that it is difficult to industrially prepare Aripiprazole Anhydride Crystals B having small particle size.

Under the circumstances, the inventors of the present invention have found that Conventional hydrate
20 can be easily milled, and Aripiprazole Anhydride B having small particle size can be obtained in high yield with good-operability by heating the milled hydrate A thus obtained.

The Aripiprazole Anhydride Crystals B of the
25 present invention are prepared for example by heating the aforementioned Aripiprazole Hydrate A at 90-125°C. The heating time is generally about 3-50 hours, but cannot be stated unconditionally since it differs

44
178

- 26 -

depending on heating temperature. The heating time and heating temperature are inversely related, so that for example the heating time will be longer the lower the heating temperature, and shorter the higher the heating temperature. Specifically, if the heating temperature of Aripiprazole Hydrate A is 100°C, the heating time should normally be 18 hours or more or preferably about 24 hours. If the heating temperature of Aripiprazole Hydrate A is 120°C, on the other hand, the heating time can be about 3 hours. The Aripiprazole Anhydride Crystals B of the present invention can be prepared with certainty by heating Aripiprazole Hydrate A for about 18 hours at 100°C, and then heating it for about 3 hours at 120°C. The Aripiprazole Anhydride Crystals B of the present invention can also be obtained if the heating time is extended still further, but this may not be economical.

When small particle size is not required for the formulation, e.g., when drug substance is being manufactured for injectable or oral solution formulations, Aripiprazole Anhydride Crystal B can be also obtained the following process.

The inventors also discovered that it is possible to obtain aripiprazole anhydride crystals by heating conventional aripiprazole hydrate or conventional aripiprazole anhydride crystals to a specific temperature but this process does not yield Anhydride B crystalline substance suitable for

95
129

- 27 -

commercial use in the formulation of solid oral dose formulations.

Furthermore, the Aripiprazole Anhydride Crystals B of the present invention are prepared for example by heating conventional aripiprazole anhydride crystals at 90-125°C. The heating time is generally about 3-50 hours, but cannot be stated unconditionally since it differs depending on heating temperature. The heating time and heating temperature are inversely related, so that for example the heating time will be longer the lower the heating temperature, and shorter the higher the heating temperature.

Specifically, if the heating temperature of the aripiprazole anhydride crystals is 100°C, the heating time can be about 4 hours, and if the heating temperature is 120°C the heating time can be about 3 hours.

In addition to Aripiprazole Hydrate A and Aripiprazole Anhydride Crystals B mentioned above, the present invention provides Aripiprazole Anhydride Crystals C to G as follows.

1. The present invention relates to aripiprazole anhydride crystals (hereinafter referred to as "type C crystals of aripiprazole anhydride") having the following physicochemical properties (1) to (5):

(1) an endothermic curve which is substantially identical to the thermogravimetric/

A6
100

- 28 -

differential thermal analysis (heating rate: 5°C/min.)
endothermic curve shown in Figure 8;

(2) an ¹H-NMR spectrum which is substantially
identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in
5 Figure 9;

(3) a powder X-ray diffraction spectrum which
is substantially identical to the powder X-ray
diffraction spectrum shown in Figure 10;

(4) an IR spectrum which is substantially
10 identical to the IR (KBr) shown in Figure 11; and

(5) a solid ¹³C-NMR spectrum which is
substantially identical to the solid ¹³C-NMR spectrum
shown in Figure 12.

2. The present invention relates to
15 aripiprazole anhydride crystals (hereinafter referred
to as "type D crystals of aripiprazole anhydride")
having the following physicochemical properties (6) to
(10):

(6) an endothermic curve which is
20 substantially identical to the thermogravimetric/
differential thermal analysis (heating rate: 5°C/min.)
endothermic curve shown in Figure 13;

(7) an ¹H-NMR spectrum which is substantially
identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in
25 Figure 14;

(8) a powder X-ray diffraction spectrum which
is substantially identical to the powder X-ray
diffraction spectrum shown in Figure 15;

47
B1

- 29 -

(9) an IR spectrum which is substantially identical to the IR (KBr) shown in Figure 16; and

(10) a solid ^{13}C -NMR spectrum which is substantially identical to the ^{13}C -NMR spectrum shown in
5 Figure 17.

3. The present invention relates to aripiprazole anhydride crystals (hereinafter referred to as "type E crystals of aripiprazole anhydride") having the following physicochemical properties (11) to
10 (14):

(11) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate: $5^{\circ}\text{C}/\text{min.}$) endothermic curve shown in Figure 18;

15 (12) an ^1H -NMR spectrum which is substantially identical to the ^1H -NMR spectrum ($\text{DMSO}-d_6$, TMS) shown in Figure 19;

(13) a powder X-ray diffraction spectrum which is substantially identical to the powder X-ray
20 diffraction spectrum shown in Figure 20; and

(14) an IR spectrum which is substantially identical to the IR (KBr) shown in Figure 21.

4. The present invention relates to aripiprazole anhydride crystals (hereinafter referred to as "type F crystals of aripiprazole anhydride")
25 having the following physicochemical properties (15) to (18):

(15) an endothermic curve which is

48
182

- 30 -

substantially identical to the thermogravimetric/
differential thermal analysis (heating rate: 5°C/min.)
endothermic curve shown in Figure 22;

(16) an ¹H-NMR spectrum which is substantially
5 identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in
Figure 23;

(17) a powder X-ray diffraction spectrum
which is substantially identical to the powder X-ray
diffraction spectrum shown in Figure 24; and

10 (18) an IR spectrum which is substantially
identical to the IR (KBr) shown in Figure 25.

5. The present invention relates a process
for preparing aripiprazole anhydride crystals stated in
the aforementioned item 1, characterized by heating
15 aripiprazole anhydride crystals at a temperature being
higher than 140°C and lower than 150°C.

6. The present invention relates a process
for preparing aripiprazole anhydride crystals stated in
the aforementioned item 2, characterized by
20 recrystallizing from toluene.

7. The present invention relates to a process
for preparing aripiprazole anhydride crystals stated in
the aforementioned item 3, characterized by heating and
dissolving aripiprazole anhydride crystals in
25 acetonitrile, and cooling it.

8. The present invention relates to a process
for preparing aripiprazole anhydride crystals stated in
the aforementioned item 4, characterized by heating a

49
183

- 31 -

suspension of aripiprazole anhydride crystals in acetone.

9. The present invention relates to a pharmaceutical composition containing at least one
5 aripiprazole anhydride crystals selected from the group consisting of the aripiprazole anhydride crystals stated in the aforementioned item 1, the aripiprazole anhydride crystals stated in the aforementioned item 2, the aripiprazole anhydride crystals stated in the
10 aforementioned item 3, the aripiprazole anhydride crystals stated in the aforementioned item 4, and the aripiprazole anhydride crystals stated in the after-mentioned item 10, together with pharmaceutically acceptable carriers.

15 10. The present invention relates to aripiprazole anhydride crystals (hereinafter referred to as "type G crystals of aripiprazole anhydride") having the following physicochemical properties (19) to (22):

20 (19) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate; 5°C/min.) endothermic curve shown Figure 26.

25 (20) an ¹H-NMR spectrum which is substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 27.

(21) a power X-ray diffraction spectrum which

56
184

- 32 -

is substantially identical to the power X-ray diffraction spectrum shown in Figure 28; and

(22) an IR spectrum which is substantially identical to the IR (Kbr) shown in Figure 29.

5 11. The present invention relates to a process for preparing aripiprazole anhydride crystals stated in the aforementioned item 10, characterized by putting glassy state of Aripiprazole Anhydride in a sealed vessel and keeping it at room temperature for at
10 least 2 weeks.

 12. The present invention relates to a process for the preparation of granules, characterized by wet granulating conventional Aripiprazole Anhydride Crystals or Aripiprazole Anhydride Crystals B, C, D, E,
15 F or G, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

 13. The present invention relates to a process for the pharmaceutical solid oral preparation,
20 characterized by drying a pharmaceutical solid oral preparation comprising conventional Aripiprazole Anhydride Crystals or Aripiprazole Anhydride Crystals B, C, D, E, F or G, and one or more pharmaceutically acceptable carriers at 70 to 100°C.

25 14. The present invention relates to a pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B, C, D, E, F or G and one or more pharmaceutically acceptable carriers,

51
185

- 33 -

wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at
5 pH 5.0 after 60 minutes.

15. The present invention relates to a pharmaceutical solid oral preparation having at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at
10 pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

16. The present invention relates to a pharmaceutical solid oral preparation obtained by wet granulating conventional Aripiprazole Anhydride
15 Crystals, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again, and the pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after
20 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

17. The present invention relates to a pharmaceutical solid oral preparation obtained by drying a pharmaceutical solid oral preparation
25 comprising conventional Aripiprazole Anhydride Crystals and one or more pharmaceutically acceptable carriers at 70 to 100°C, and the pharmaceutical solid oral preparation has at least one dissolution rate selected

- 34 -

from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

18. The present invention relates to a
5 process for the preparation of granules, characterized by wet granulating conventional Aripiprazole Hydrate Crystals, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

10 19. The present invention relates to a process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising conventional Aripiprazole Hydrate Crystals and one or more pharmaceutically
15 acceptable carriers at 70 to 100°C.

20 20. The present invention relates to a pharmaceutical solid oral preparation obtained by wet granulating conventional Aripiprazole Hydrate Crystals, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again, and the pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and
25 55% or more at pH 5.0 after 60 minutes.

21. The present invention relates to a pharmaceutical solid oral preparation obtained by drying a pharmaceutical solid oral preparation

53
187

- 35 -

comprising conventional Aripiprazole Hydrate Crystals and one or more pharmaceutically acceptable carriers at 70 to 100°C, and the pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

The Type C to F crystals of aripiprazole anhydride of the present invention correspond to the Type-III to VI crystals of aripiprazole anhydride disclosed in JP-2001-348276.

Type C crystals of aripiprazole anhydride

Type C crystals of aripiprazole anhydride of the present invention have the following physicochemical properties (1) to (5):

(1) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate: 5°C/min.) endothermic curve shown in Figure 8, more particularly, it has an endothermic peak around 150.2°C;

(2) an ¹H-NMR spectrum which is substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 9. Specifically, it has characteristic peaks at 1.55 - 1.63 ppm (m, 2H), 1.68 - 1.78 ppm (m, 2H), 2.35 - 2.46 ppm (m, 4H), 2.48 - 2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, J=7, 4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4Hz,

54
188

- 36 -

1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11 - 7.17 ppm (m, 1H), 7.28 - 7.32 ppm (m, 2H) and 10.00 ppm (s, 1H);

(3) a powder X-ray diffraction spectrum which is substantially identical to the powder X-ray diffraction spectrum shown in Figure 10. Specifically, it has characteristic peaks at $2\theta = 12.6^\circ, 13.7^\circ, 15.4^\circ, 18.1^\circ, 19.0^\circ, 20.6^\circ, 23.5^\circ$ and 26.4° ;

(4) an IR spectrum which is substantially identical to the IR (KBr) spectrum shown in Figure 11. Specifically, it has clear infrared absorption bands at 2939, 2804, 1680, 1375 and 780 cm^{-1} ; and

(5) a solid ^{13}C -NMR spectrum which is substantially identical to the solid ^{13}C -NMR spectrum shown in Figure 12, specifically, it has characteristic peaks at 32.8 ppm, 60.8 ppm, 74.9 ppm, 104.9 ppm, 152.2 ppm, 159.9 ppm and 175.2 ppm.

Preparation method of type C crystals of aripiprazole anhydride

Type C crystals of aripiprazole anhydride of the present invention is prepared, for example by heating an aripiprazole anhydride at a temperature of higher than 140°C and lower than 150°C .

Aripiprazole anhydride used as the raw material may be conventional aripiprazole anhydride crystals, for example, type-I crystals of aripiprazole anhydride, type-II crystals of aripiprazole anhydride

- 37 -

crystals and the like, and these anhydrides may be either purified products or crude materials. Alternatively, type B crystals of aripiprazole anhydride, type D crystals of aripiprazole anhydride, type E crystals of aripiprazole anhydride, type F crystals of aripiprazole anhydride, or type G crystals of aripiprazole anhydride being prepared in the present invention can be used as the raw material of aripiprazole anhydrides. These aripiprazole anhydrides can be used singly or in combination of at least 2 kinds thereof.

Heating temperature is generally higher than 140°C and lower than 150°C, preferably at 142 - 148°C, and heating time is generally for 15 minutes to 3 hours, preferably for 30 minutes to 1 hour.

When, an aripiprazole anhydride is heated at the above-mentioned temperature, then type C crystals of aripiprazole anhydride are formed.

Thus obtained type C crystals of aripiprazole anhydride can be isolated and purified by well-known methods. For example, after heating the aripiprazole anhydride at the above-mentioned temperature, and by cooling to a room temperature, then type C crystals of aripiprazole anhydride, having 100 % of purity can be obtained.

25

Type D crystals of aripiprazole anhydride

Type D crystals of aripiprazole anhydride of the present invention have the following

58
190

- 38 -

physicochemical properties (6) to (10):

(6) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate: 5°C/min.)

5 endothermic curve shown in Figure 13; more particularly, it has an endothermic peak around 136.8 °C and 141.6 °C;

(7) an ¹H-NMR spectrum which is substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in
10 Figure 14. Specifically, it has characteristic peaks at 1.55 - 1.63 ppm (m, 2H), 1.68 - 1.78 ppm (m, 2H), 2.35 - 2.46 ppm (m, 4H), 2.48 - 2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, J=7, 4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d,
15 J=2.4Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11 - 7.17 ppm (m, 1H), 7.28 - 7.32 ppm (m, 2H) and 10.00 ppm (s, 1H);

(8) a powder X-ray diffraction spectrum which is substantially identical to the powder X-ray
20 diffraction spectrum shown in Figure 15. Specifically, it has characteristic peaks at 2θ = 8.7°, 11.6°, 16.3°, 17.7°, 18.6°, 20.3°, 23.4° and 25.0°;

(9) an IR spectrum which is substantially identical to the IR (KBr) spectrum shown in
25 Figure 16. Specifically, it has clear infrared absorption bands at 2946, 1681, 1375, 1273, 1175 and 862 cm⁻¹; and

(10) a solid ¹³C-NMR spectrum which is

57
191

- 39 -

substantially identical to the solid ^{13}C -NMR spectrum shown in Figure 17, specifically, it has characteristic peaks at 32.1 ppm, 62.2 ppm, 66.6 ppm, 104.1 ppm, 152.4 ppm, 158.4 ppm, and 174.1 ppm.

5

Preparation method of type D crystals of aripiprazole anhydride

Type D crystals of aripiprazole anhydride of the present invention is prepared, for example, by
10 recrystallization of aripiprazole anhydride from toluene. Specifically, an aripiprazole anhydride is added to toluene, further heated and dissolved, then thus obtained solution is cooled. By such procedures, type D crystals of aripiprazole anhydride of the
15 present invention is separated out as crystals in toluene.

Aripiprazole anhydride to be used as the raw materials may be conventional aripiprazole anhydride, for example type-I crystals of aripiprazole anhydride,
20 type-II crystals of aripiprazole anhydride and the like, and these anhydrides may be either purified products or crude materials. Alternatively, type B crystals of aripiprazole anhydride, type C crystals of aripiprazole anhydride, type E crystals of aripiprazole
25 anhydride, type F crystals of aripiprazole anhydride, or type G crystals of aripiprazole anhydride being prepared in the present invention can be used as the raw material for aripiprazole anhydrides. These

58
192

- 40 -

aripiprazole anhydrides can be used singly or in combination of at least 2 kinds thereof.

When the solution obtained by heating and dissolving is cooled, type D crystals of aripiprazole
5 may be added as a seed crystal to said solution.
Further, the seed crystal may be formed by cooling gradually said solution being obtained by heating and dissolving. In the presence of the seed crystal, type D crystals of aripiprazole anhydride may be separated
10 out.

Thus separated out type D crystals of aripiprazole anhydride can be isolated and purified in accordance with well-known methods. By such procedures, type D crystals of aripiprazole anhydride,
15 having the purity of 100 % can be obtained.

Type E crystals of aripiprazole anhydride

Type E crystals of aripiprazole anhydride of the present invention have the following
20 physicochemical properties (11) to (14):

(11) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate: 5°C/min.) endothermic curve shown in Figure 18, specifically, it
25 has an endothermic peak around 146.5°C;

(12) an ¹H-NMR spectrum which is substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 19. Specifically, it has characteristic peaks

59
193

- 41 -

at 1.55 - 1.63 ppm (m, 2H), 1.68 - 1.78 ppm (m, 2H),
2.35 - 2.46 ppm (m, 4H), 2.48 - 2.56 ppm (m, 4H +
DMSO), 2.78 ppm (t, J=7, 4 Hz, 2H), 2.97 ppm (brt,
J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d,
5 J=2.4Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H),
7.04 ppm (d, J=8.1 Hz, 1H), 7.11 - 7.17 ppm (m, 1H),
7.28 - 7.32 ppm (m, 2H) and 10.00 ppm (s, 1H);

(13) a powder X-ray diffraction spectrum
which is substantially identical to the powder X-ray
10 diffraction spectrum shown in Figure 20. Specifically,
it has characteristic peaks at $2\theta = 8.0^\circ$, 13.7° , 14.6° ,
 17.6° , 22.5° and 24.0° ; and

(14) an IR spectrum which is substantially
identical to the IR (KBr) spectrum shown in Figure 21.
15 Specifically, it has clear infrared absorption bands at
2943, 2817, 1686, 1377, 1202, 969 and 774 cm^{-1} .

Preparation method of type E crystals of aripiprazole
anhydride

20 Type E crystals of aripiprazole anhydride of
the present invention is prepared, for example by
recrystallization of the aripiprazole anhydride from
acetonitrile. Specifically, by adding a well-known
aripiprazole anhydride to acetonitrile, heating and
25 dissolving, then the solution thus obtained may be
cooled. In accordance with such procedures, type E
crystals of aripiprazole anhydride of the present
invention are separated out in the acetonitrile.

- 42 -

When a conventional aripiprazole anhydride is added to acetonitrile, type-I crystals of aripiprazole anhydride, type-II crystals of aripiprazole anhydride and type D crystals of aripiprazole anhydride are separated out, other than type E crystals of aripiprazole anhydride. Plate crystals being separated out from the acetonitrile solution at 70°C are type-I crystals, type-II crystals and type D crystals, while type E crystals are precipitated out as needle crystals. When the acetonitrile solution after separated out of these crystals is heated again (for example, heated at over 75°C), the plate crystals (type-I crystals, type-II crystals and type D crystals) are quickly dissolved, on the contrary, the needle form crystals (type E crystals) do not dissolved. Additionally, when the acetonitrile solution is cooled again, then needle form crystals (type E crystals) are further separated out around the needle form crystals (type E crystals) previously precipitated as the seed crystals. Thus, type E crystals of aripiprazole anhydride can be precipitated in the acetonitrile solution.

Aripiprazole anhydrides used as the raw materials may be conventional aripiprazole anhydrides, for example any one of type-I crystals of aripiprazole anhydride and type-II crystals of aripiprazole anhydride and the like, and these anhydrides may be either purified products or crude materials.

65
195

- 43 -

Alternatively, type B crystals of aripiprazole anhydride, type C crystals of aripiprazole anhydride, type D crystals of aripiprazole anhydride, type F crystals of aripiprazole anhydride, or type G crystals of aripiprazole anhydride can be used as the raw materials for aripiprazole anhydrides. These aripiprazole anhydrides can be used singly or in combination of at least 2 kinds thereof.

When the acetonitrile solution obtained by heating (heating and dissolving) is cooled, the type E crystals of aripiprazole may be added as a seed crystal to said solution. Further, the seed crystal may be formed by cooling gradually said acetonitrile solution which was obtained by heating.

Thus separated out type E crystals of aripiprazole anhydride can be isolated and purified in accordance with well-known methods. By such procedures, type E crystals of aripiprazole anhydride, having the purity of 100 % can be obtained.

20

Type F crystals of aripiprazole anhydride

Type F crystals of aripiprazole anhydride of the present invention have the following physicochemical properties (15) to (18):

(15) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate: 5°C/min.) endothermic curve shown in Figure 22, specifically,

62
196

- 44 -

it has an endothermic peaks around 137.5°C and 149.8°C;

(16) an ¹H-NMR spectrum which is substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 23. Specifically, it has characteristic peaks at 1.55 - 1.63 ppm (m, 2H), 1.68 - 1.78 ppm (m, 2H), 2.35 - 2.46 ppm (m, 4H), 2.48 - 2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, J=7, 4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11 - 7.17 ppm (m, 1H), 7.28 - 7.32 ppm (m, 2H) and 10.00 ppm (s, 1H);

(17) a powder X-ray diffraction spectrum which is substantially identical to the powder X-ray diffraction spectrum shown in Figure 24. Specifically, it has characteristic peaks at 2θ = 11.3°, 13.3°, 15.4°, 22.8°, 25.2° and 26.9°, and

(18) Having an IR spectrum which is substantially identical to the IR (KBr) spectrum shown in Figure 25. Specifically, it has clear infrared absorption bands at 2940, 2815, 1679, 1383, 1273, 1177, 1035, 963 and 790 cm⁻¹.

Preparation method of type F crystals of aripiprazole anhydride

Type F crystals of aripiprazole anhydride of the present invention is prepared, for example by suspending an aripiprazole anhydride in acetone, and

68
197

- 45 -

thus obtained acetone suspension is heated.

Aripiprazole anhydrides used as the raw materials may be conventional aripiprazole anhydride, for example any one of type-I crystals of aripiprazole anhydride and type-II crystals of aripiprazole anhydride and the like, and these anhydrides may be either purified products or crude materials. Alternatively, type B crystals of aripiprazole anhydride, type C crystals of aripiprazole anhydride, type D crystals of aripiprazole anhydride, type E crystals of aripiprazole anhydride, or type G crystals of aripiprazole anhydride prepared in the present invention can be used as the raw materials for aripiprazole anhydrides. These aripiprazole anhydrides can be used singly or in combination of at least 2 kinds thereof.

Heating temperature of the acetone suspension may be generally about the boiling point of acetone, and heating time is generally 5 to 10 hours. When the acetone suspension is heated about the boiling point of acetone, then type F crystals of aripiprazole anhydride is formed, the crystals are isolated by filtration with heating. Isolation of the crystals may be carried out in accordance with well-known methods. By such procedures, type F crystals of aripiprazole anhydride, having the purity of 100 % can be obtained.

64
198

- 46 -

Type G crystals of aripiprazole anhydride

Type G crystals of aripiprazole anhydride of the present invention have the following physicochemical properties (19) to (22):

- 5 (19) an endothermic curve which is substantially identical to the thermogravimetric/differential thermal analysis (heating rate: 5°C/min.) endothermic curve shown in Figure 26, more particularly, it has an endothermic peak around 141.0°C
- 10 and an exothermic peak around 122.7°C;
- (20) an ¹H-NMR spectrum which is substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 27. Specifically, it has characteristic peaks at 1.55 - 1.63 ppm (m, 2H), 1.68 - 1.78 ppm (m, 2H),
- 15 2.35 - 2.46 ppm (m, 4H), 2.48 - 2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, J=7.4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1
- 20 Hz, 1H), 7.11 - 7.17 ppm (m, 1H), 7.28 - 7.32 ppm (m, 2H) and 10.00 ppm (s, 1H);
- (21) a powder X-ray diffraction spectrum which is substantially identical to the powder X-ray diffraction spectrum shown in Figure 28. Specifically,
- 25 it has characteristic peaks at 2θ = 10.1°, 12.8°, 15.2°, 17.0°, 17.5°, 19.1°, 20.1°, 21.2°, 22.4°, 23.3°, 24.5° and 25.8°; and
- (22) an IR spectrum which is substantially

85
199

- 47 -

identical to the IR (KBr) spectrum shown in Figure 29. Specifically, it has clear infrared absorption bands at 2942, 2813, 1670, 1625, 1377, 1195, 962 and 787 cm^{-1} .

5 Preparation method of type G crystals of aripiprazole anhydride

Type G crystals of aripiprazole anhydride of the present invention can be prepared, for example by putting glassy state of aripiprazole anhydride in a
10 sealed vessel and leaving to stand it at room temperature for at least two weeks, preferably two weeks to six months. Further, glassy state of aripiprazole anhydride as starting material can be obtained by heating and melting aripiprazole anhydride
15 at around 170°C, then cooling it to room temperature.

Aripiprazole anhydride used as the raw material may be well-known aripiprazole anhydride crystals, for example, any one of type-I crystals of aripiprazole anhydride and type-II crystals of
20 aripiprazole anhydride and the like, and these anhydrides may be either purified products or crude materials. Alternatively, type B crystals of aripiprazole anhydride, type C crystals of aripiprazole anhydride, type D crystals of aripiprazole anhydride,
25 type E crystals of aripiprazole anhydride, or type F crystals of aripiprazole anhydride being prepared in the present invention can be used as the raw material of aripiprazole anhydrides. These aripiprazole

86
290

- 48 -

anhydrides can be used singly or in combination of at least 2 kinds thereof.

Thus obtained type G crystals of aripiprazole anhydride can be isolated and purified by well-known
5 methods. For example, glassy state of aripiprazole anhydride leave to stand according to the above-mentioned method, then type G crystals of aripiprazole anhydride, having 100% of purity can be obtained.

10 Type C crystals of aripiprazole anhydride, type D crystals of aripiprazole anhydride, type E crystals of aripiprazole anhydride, type F crystals of aripiprazole anhydride and type G crystals of aripiprazole anhydride of the present invention neither
15 easily convert into hydrates thereof, nor substantially decrease the original solubility, even when they are stored for a long period of time.

In accordance with the present invention, methods for preparing aripiprazole anhydride crystals
20 having high purity, which can apply in an industrial scale with a good repeatability is provided.

In accordance with the present invention, pharmaceutical compositions comprising aripiprazole anhydride crystals are provided, of which the
25 solubility does not decrease, and of which the stability can keep excellent, even if they are stored for long time.

- 49 -

The aripiprazole anhydride crystals which are the raw material for preparing the Aripiprazole Anhydride Crystals B to G of the present invention are prepared for example by Method a or b below.

5

"Method a": Process for Preparing Crude Aripiprazole Crystals

Conventional Aripiprazole Anhydride crystals are prepared by well-known methods, as described in Example 1 of Japanese Unexamined Patent Publication No. 191256/1990.

A suspension of 47 g of 7-(4-bromobutoxy)-3,4-dihydrocarbostyryl, 35 g of sodium iodide with 600 ml of acetonitrile was refluxed for 30 minutes. To this suspension was added 40 g of 1-(2,3-dichlorophenyl)piperazine and 33 ml of triethylamine and the whole mixture was further refluxed for 3 hours. After the solvent was removed by evaporation, the residue thus obtained was dissolved in chloroform, washed with water then dried with anhydrous magnesium sulfate. The solvent was removed by evaporation, and the residue thus obtained was recrystallized from ethanol twice, to yield 57.1 g of 7-{4-[4-(2,3-dichlorophenyl)-1-piperazinyl]butoxy}-3,4-dihydrocarbostyryl.

Colorless flake crystals

Melting point: 139.0-139.5°C

68
202

- 50 -

"Method b": Process for Preparing Conventional Anhydride

The Method b is described in the Proceedings of the 4th Japanese-Korean Symposium on Separation
5 Technology (October 6-8, 1996).

Furthermore, the Aripiprazole Anhydride Crystals B of the present invention are prepared for example by heating conventional aripiprazole hydrate at 90-125°C. The heating time is generally about 3-50
10 hours, but cannot be stated unconditionally since it differs depending on heating temperature. The heating time and heating temperature are inversely related, so that for example the heating time will be longer the lower the heating temperature, and shorter the higher
15 the heating temperature. Specifically, if the heating temperature of the aripiprazole hydrate is 100°C, the heating time can be about 24 hours, while if the heating temperature is 120°C, the heating time can be about 3 hours.

20 The aripiprazole hydrate which is the raw material for preparing the Aripiprazole Anhydride Crystals B of the present invention is prepared for example by Method c below.

25 "Method c": Process for Preparing Conventional Hydrate

Aripiprazole hydrate is easily obtained by dissolving the aripiprazole anhydride crystals obtained by Method a above in a hydrous solvent, and heating and

89
107

- 51 -

then cooling the resulting solution. Using this method, aripiprazole hydrate is precipitated as crystals in the hydrous solvent.

An organic solvent containing water is usually used as the hydrous solvent. The organic solvent should be one which is miscible with water, such as for example an alcohol such as methanol, ethanol, propanol or isopropanol, a ketone such as acetone, an ether such as tetrahydrofuran, dimethylformamide, or a mixture thereof, with ethanol being particularly desirable. The amount of water in the hydrous solvent can be 10-25% by volume of the solvent, or preferably close to 20% by volume.

15 Medicinal Composition

A medicinal composition of the present invention will contain Aripiprazole Anhydride Crystals B, C, D, E, F and G in a pharmaceutically acceptable carrier or combination of carriers.

20 Carriers which are pharmaceutically acceptable include diluents and excipients generally used in pharmaceuticals, such as fillers, extenders, binders, moisturizers, disintegrators, surfactants, and lubricants.

25 The medicinal composition of the present invention may be formulated as an ordinary medicinal preparation, for example in the form of tablets, flashmelt tablets, pills, powder, liquid, suspension,

204/20

- 52 -

emulsion, granules, capsules, suppositories or as an injection (liquid, suspension, etc.).

When a tablet formulation is used, a wide variety of carriers that are known in the field can be used. Examples include lactose, saccharose, sodium chloride, glucose, xylitol, mannitol, erythritol, sorbitol, urea, starch, calcium carbonate, kaolin, crystal cellulose, silic acid and other excipients; water, ethanol, propanol, simple syrup, glucose liquid, starch liquid, gelatin solution, carboxymethyl cellulose, shellac, methyl cellulose, potassium phosphate, polyvinyl pyrrolidone and other binders; dried starch, sodium alginate, agar powder, laminaran powder, sodium bicarbonate, calcium carbonate, polyoxyethylene sorbitan fatty acid esters, sodium lauryl sulfate, monoglyceride stearate, starch, lactose and other disintegrators; saccharose, stearin, cacao butter, hydrogenated oil and other disintegration inhibitors; quaternary ammonium salt, sodium lauryl sulfate and other absorption promoters; glycerine, starch and other moisture retainers; starch, lactose, kaolin, bentonite, colloidal silic acid and other adsorbents; and refined talc, stearate, boric acid powder, polyethylene glycol and other lubricants and the like. Tablets can also be formulated if necessary as tablets with ordinary coatings, such as sugar-coated tablets, gelatin-coated tablets, enteric coated tablets and film coated tablets, as well as double tablets and

205 71

- 53 -

multilayered tablets.

When a pill formulation is used, a wide variety of carriers that are known in the field can be used. Examples include glucose, lactose, starch, cacao
5 butter, hardened vegetable oil, kaolin, talc and other excipients; gum arabic powder, traganth powder, gelatin, ethanol and other binders; and laminaran, agar and other disintegrators and the like.

When a suppository formulation is used, a
10 wide variety of carriers that are known in the field can be used. Examples include polyethylene glycol, cacao butter, higher alcohol, esters of higher alcohol, gelatin semi-synthetic glyceride and the like.

Capsules are prepared according to ordinary
15 methods by mixing aripiprazole anhydride crystals with the various carriers described above and packing them in hard gelatin capsules, soft capsules, hydroxypropylmethyl cellulose capsules (HPMC capsules) and the like.

20 In addition, colorants, preservatives, perfumes, flavorings, sweeteners and the like as well as other drugs may be included in the medicinal composition.

In case of forming the pharmaceutical solid
25 oral preparation in the form of granules, it can be prepared by wet granulating a mixed powder of granulating ingredients comprising, aripiprazole anhydride crystals (conventional aripiprazole anhydride

72
106

- 54 -

crystals or aripiprazole anhydride crystals selected from the group consisting of aripiprazole anhydride type B, C, D, E, F and G crystals) and various carriers which are heretofore well-known in this field, such as

5 excipients, disintegrators, disintegration inhibitors, humectants, absorption accelerators, adsorbents, lubricants, colorants and the like (for the examples of these agents, those of previously mentioned can be referred to) by adding a liquid (generally, water or an

10 aqueous solution containing binding agents). As for the wet granulation, there are various methods are included, for example, fluidized bed granulation, kneading granulation, extruding granulation, rotating granulation and the like can be mentioned. Among these

15 methods, in case of conducting the fluidized bed granulation, the granulating ingredients containing various carriers are mixed with inlet air, then upon continued fluidizing the granulating ingredients and the liquid is sprayed to conduct granulation. In case

20 of conducting the kneading granulation, the granulating ingredients containing various carriers are mixed by agitation, then upon continued agitating the granulating ingredients, granulation is conducted by adding the liquid. After the granulation, if

25 necessary, the obtained granules are sized to make them to the desired size by use of a suitable sieve or a mill having suitable screen size. The granules thus obtained by such a method are dried again in addition

73
202

- 55 -

to usual drying being conducted when preparing the granules. As for the drying methods, various methods can be applied, for example, methods by use of a fluidized bed dryer, a fan dryer, a vacuum dryer and the like can be mentioned. Generally, drying methods can be conducted under conventional conditions, for example, in case of using the fluidized bed dryer, drying procedure is conducted with an air flow of 0.5 m³/min to 50 m³/min, an inlet air temperature at 70 to 100°C for 10 min to 1 hour. After dried, the granules are subjected to size, then further dried. In case of using the fluidized bed dryer or fan dryer or the like, the drying procedure is conducted under the conditions with an air flow of 0.5 m³/min to 50 m³/min, an inlet air temperature at 70 to 100°C for 1 to 6 hours. In case of using the vacuum dryer, the drying procedure is conducted under the conditions of reduced pressure of about at 0-10 torr of degree of vacuum at 70 to 100°C of jacket temperature for 1 to 6 hour.

The thus prepared granules may be used as they are for the pharmaceutical solid oral preparations, or if necessary, they may be shaped in the form of tablets. Further, the dried granules dried by usual manner are shaped in the form of tablets, then they may be dried again.

The thus prepared pharmaceutical solid oral preparation comprising aripiprazole anhydride crystals hardly changes to hydrates even if they are stored for

34
88

- 56 -

a long period of time, therefore the pharmaceutical solid oral preparation, of which dissolution rate does not hardly lowered (dissolution rate to maintain maximum drug concentration (C_{max}): 60% or higher
5 dissolution rate obtained after 30 minutes at pH 4.5, 70% or higher dissolution rate obtained after 60 minutes at pH 4.5, or 55% or higher dissolution rate obtained after 60 minutes at pH 5.0) can be provided.

Another pharmaceutical solid oral preparation
10 can be provided by granulating a conventional aripiprazole hydrate crystals by a method similar to that of mentioned above, and dried by usual manner under similar conditions, then dried again.

Alternatively, the dried granules dried by usual manner
15 are shaped to tablets form, then they are dried again, then pharmaceutical solid oral preparations of which dissolution rate does not lowered (dissolution rate to maintain maximum drug concentration (C_{max}): 60% or higher dissolution rate obtained after 30 minutes at pH
20 4.5, 70% or higher dissolution rate obtained after 60 minutes at pH 4.5 or 55% or higher dissolution rate obtained after 60 minutes at pH 5.0) can be provided.

These facts can be understood that, the conventional aripiprazole anhydride crystals or the aripiprazole
25 hydrate crystals contained in the pharmaceutical solid oral preparation are changed to "B type crystals" of aripiprazole anhydride by drying twice.

The amount of Aripiprazole Anhydride Crystals

28
29

- 57 -

B, C, D, E, F and G that should be included in the medicinal composition of the present invention can be selected from a wide range suitable for the indication sought to be treated. Generally, the Aripiprazole
5 Anhydride Crystals B should be present in about 1-70% by weight or particularly about 1-30% by weight based on the medicinal composition.

The method of administration of the medicinal composition of the present invention may be adjusted to
10 suit, for example, the formulation of the drug product, the age, gender and other conditions (including the severity thereof) of the patient. In the case of tablets, pills, liquids, suspensions, emulsions, granules and capsules, for example, administration is
15 oral. In the case of an injection, it is administered intravenously either by itself or mixed with an ordinary replenisher such as glucose or amino acids, or may also be administered by itself intramuscularly, intracutaneously, subcutaneously or intraperitoneally,
20 as necessary. In the case of a suppository, administration is intrarectal.

The dosage of the medicinal composition of the present invention is selected depending on the usage, the age, gender and other conditions of the
25 patient, the severity of the condition and so forth, but ordinarily the amount of aripiprazole anhydride crystals can be about 0.1-10 mg per 1 kg of body weight per day. The preparation which is the unit of

28
210

- 58 -

administration should contain in the range of about 1-100 mg of Aripiprazole Anhydride Crystals B, more particularly 1-30 mg per unit dose.

The medicinal composition of the present invention is extremely stable, with substantially no decrease in solubility even when stored for long periods of time.

The medicinal composition of the present invention is effective in the prevention and treatment of central nervous system disorders such as schizophrenia and may also be effective in the treatment of intractable (drug-resistant, chronic) schizophrenia with cognitive impairment and intractable (drug-resistant, chronic) schizophrenia without cognitive impairment, anxiety including mild anxiety, mania including bipolar disorder acute mania and acute mania, bipolar disorder, depression including bipolar disorder depression, autism, Down's syndrome, attention deficit hyperactivity disorder (ADHD), Alzheimer's disease, Parkinson's disease and other neurodegenerative diseases, panic, obsessive compulsive disorder (OCD), sleep disorders, sexual dysfunction, alcohol and drug dependency, vomiting, motion sickness, obesity, multiparticulate headache and cognitive impairment.

Analytical Methods

(1) The ¹H-NMR spectrum was measured in DMSO-d₆ using TMS as the standard.

77
211

- 59 -

(2) Powder X-ray Diffraction

Using a Rigaku Denki RAD-2B diffraction meter, the powder x-ray diffraction pattern was measured at room temperature using a Cu Ka filled tube
5 (35 kV 20mA) as the x-ray source with a wide-angle goniometer, a 1° scattering slit, an 0.15 mm light-intercepting slit, a graphite secondary monochromator and a scintillation counter. Data collection was done in 2θ continuous scan mode at a scan speed of 5°/minute
10 in scan steps of 0.02° in the range of 3° to 40°.

(3) The IR spectrum was measured by the KBr method.

(4) Thermogravimetric/Differential Thermal Analysis

15 Thermogravimetric/differential thermal analysis was performed using a Seiko SSC 5200 control unit and a TG/DTA 220 simultaneous differential thermal/thermogravimetric measurement unit. 5-10 mg samples were placed in open aluminum pans and heated
20 from 20°C to 200°C in a dry nitrogen atmosphere at a heating rate of 5°C/minute. α-alumina was used as the standard substance.

(5) Differential Scanning Calorimetry

Thermogravimetric/differential thermal
25 analysis was performed using a Seiko SSC 5200 control unit and a DSC 220C differential scanning calorimeter. 5-10 mg samples were placed in crimped aluminum pans and heated from 20°C to 200°C in a dry nitrogen

- 60 -

atmosphere at a heating rate of 5°C/minute. α -alumina was used as the standard substance.

(6) Particle Size Measurement

0.1 g of the particles to be measured were
5 suspended in a 20 ml n-hexane solution of 0.5 g soy lecithin, and particle size was measured using a size distribution meter (Microtrack HRA, Microtrack Co.).

(7) Hygroscopicity Test Method

One g of the sample was accurately weighed in
10 a weighing bottle (diameter 5 cm), covered with kimwipes and left to rest in a 60°C/100% RH environment (water/dessicator). 24 hours later, the weighing bottle was removed, transferred to an environment of a room temperature and about 30% RH (magnesium chloride
15 hexahydrate saturated water solution/dessicator) and left to rest for 24 hours and the water content of the sample was measured by the Karl Fischer method.

(8) Solid ^{13}C -NMR Spectrometry

Solid ^{13}C -NMR spectrum was measured under the
20 conditions as follows.

Measuring apparatus: CMX-360 Solid State NMR Spectrometer (manufactured by Chemagnetic Inc.)

Computer: SPARC Station 2 (manufactured by SUN Microsystem, Inc.)

25 OS, Software: Solalis 1.1.1 Rev. B
(Registered trademark: UNIX), Spinsight Ver. 2.5

Name of measured pulse: TOSS method (TOSS is a program name of the apparatus) among CP/MAS method.

213 79

- 61 -

Width of measured puls: 90° puls was used
under the condition of CP.

Measuring sample tube: Test tube made of
zirconia, having the outside diameter of 7.5 mm, and
5 inside capacity of 0.8 ml

Revolution: 4250 Hz (Revolution per second
Contact time: 1 msec.

Waiting time: 20 sec.

Integrated times: 512 times

10 Measuring temperature: About 25°C temperature
of outside of test tube)

External standard: Methyl group (δ 17.3) of
hexamethylbenzene was used as the external standard.

15 The present invention is explained in more
detail below using reference examples, examples, sample
preparations and formulation examples.

Reference Example 1

20 19.4 g of 7-(4-chlorobutoxy)-3,4-
dihydrocarbostyryl and 16.2 g of 1-(2,3-dichlorophenyl)
piperadine 1 hydrochloride were added to 8.39 g of
potassium carbonate dissolved in 140 ml of water, and
circulated for 3 hours under agitation. After reaction
25 the mixture was cooled and the precipitated crystals
filtered out. These crystals were dissolved in 350 ml
of ethyl acetate, and about 210 ml of water/ethyl
acetate azeotrope removed under reflux. The remaining

86
214

- 62 -

solution was cooled, and the precipitated crystals filtered out. The resulting crystals were dried for 14 hours at 60°C to produce 20.4 g (74.2%) of raw aripiprazole.

5 30 g of the raw aripiprazole obtained above was recrystallized from 450 ml of ethanol according to the methods described in Japanese Unexamined Patent Publication No. 191256/1990, and the resulting crystals dried for 40 hours at 80°C to obtain aripiprazole
10 anhydride crystals. The yield was 29.4 g (98.0%).

The melting point (mp) of these aripiprazole anhydride crystals was 140°C, matching the melting point of the aripiprazole anhydride crystals described in Japanese Unexamined Patent Publication No.

15 191256/1990.

When these crystals were left for 24 hours in a dessicator set at humidity 100%, temperature 60°C, they exhibited hygroscopicity of 3.28% (see Table 1 below).

20

Reference Example 2

6930 g of the intermediate raw aripiprazole obtained in Reference Example 1 was heat dissolved in 138 liters of hydrous ethanol (water content 20%)

25 according to the method presented at the 4th Japanese-Korean Symposium on Separation Technology, gradually (2-3 hours) cooled to room temperature, and then chilled to near 0°C. The precipitated crystals were

25

- 63 -

filtered out, producing about 7200 g of aripiprazole hydrate (wet state).

The wet-state aripiprazole hydrate crystals obtained above were dried for 30 hours at 80°C to obtain 6480 g (93.5%) of conventional aripiprazole anhydride crystals. The melting point (mp) of these crystals was 139.5°C. These crystals were confirmed by the Karl Fischer method to be anhydrous, with a moisture value of 0.03%.

When left for 24 hours in a dessicator set at humidity 100%, temperature 60°C, these crystals exhibited hygroscopicity of 1.78% (see Table 1 below).

Reference Example 3

820 g of the intermediate wet-state aripiprazole hydrate obtained in Reference Example 2 was dried for 2 hours at 50°C to obtain 780 g of aripiprazole hydrate crystals. These crystals had a moisture value of 3.82% according to the Karl Fischer method. As shown in Figure 6, thermogravimetric/differential thermal analysis revealed endothermic peaks at 75.0, 123.5 and 140.5°C. Because dehydration began near 70°C, there was no clear melting point (mp).

As shown in Figure 7, the powder x-ray diffraction spectrum of aripiprazole hydrate obtained by this method exhibited characteristic peaks at $2\theta = 12.6^\circ, 15.1^\circ, 17.4^\circ, 18.2^\circ, 18.7^\circ, 24.8^\circ$ and 27.5° .

The powder x-ray diffraction spectrum of this

216/82

- 64 -

aripiprazole hydrate was identical to the powder x-ray diffraction spectrum of aripiprazole hydrate presented at the 4th Joint Japanese-Korean Symposium on Isolation Technology.

5

Reference Example 4

Preparation of 15 mg tablets containing type I crystals of aripiprazole anhydride obtained in Reference Example 2.

10 Type-I crystals of aripiprazole anhydride (525 g), lactose (1,995 g), corn starch (350 g) and crystalline cellulose (350 g) were charged in a fluidized bed granulating dryer (Flow coater FLO-5, manufactured by FREUND INDUSTRIAL CO., LTD.), and these
15 granulating ingredients were mixed by fluidizing for about 3 minutes with an inlet air temperature at 70°C and air flow rate of 3 m³/min. Further, the granulating ingredients were upon continued fluidizing under the same condition and sprayed about 1,400 g of the aqueous
20 solution to obtained wet granules. The wet granules were dried under inlet air at temperature at 80°C, for about 15 minutes. The obtained dried granules contained 4.3% of water. (Yield: 99%). The dried granules were subjected to sizing by passing to a sieve
25 of 710 μm.

About 1% by weight of magnesium stearate was added to the sized granules and mixed, then the granules were supplied to a tablet machine (Rotary

- 65 -

single tablet press 12HUK: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), there were obtained tablets, each having 95 mg of weight.

Water content of the tablets was measured according to volumetric titration method (Karl-Fischer method) described in water content measuring method in Japanese Pharmacopoea or the electrical quantity titration method.

10 Water content measuring method:

Sample (0.1 to 0.5 g) (in case of a tablet, 1 tablet was used) was weighed precisely, and the water content was measured by use of a water content measuring equipment.

15 Volumetric titration:

Automated water content measuring equipment

Model: KF-06 (manufacture by MITSUBISHI CHEMICAL CORP.)

Electrical quantity titration method:

20 Automated micro-water content measuring equipment

Model: AQ-7F (manufactured by HIRANUMA SANGYO CO., LTD.)

Automated water vaporization equipment Model:

LE-20S (manufactured by HIRANUMA SANGYO CO., LTD.)

Heating temperature: $165 \pm 10^{\circ}\text{C}$

Nitrogen gas flow rate: about 150 ml/min.

84
2/3

- 66 -

Reference Example 5

Preparation of 15 mg tablets containing type B crystals of aripiprazole anhydride

Type B crystals of aripiprazole anhydride

5 (4,500 g), lactose (17,100 g), corn starch (3,000 g) and crystalline cellulose (3,000 g) were charged in a fluidized bed granulating dryer (NEW-MARUMERIZER Model: NQ-500, manufactured by FUJI PAUDAL CO., LTD.), and these granulating ingredients were mixed by fluidizing
10 for about 3 minutes with an inlet air temperature at 70°C, air flow rate of 10 to 15 m³/min. Further, the granulating ingredients were upon continued fluidizing under the same condition, and sprayed about 12,000 g of 5% aqueous solution of hydroxypropyl cellulose to
15 obtained wet granules. The wet granules were dried under inlet air at temperature at 85°C, for about 28 minutes. The thus obtained dried granules contained 3.8% of water (measured by the method according to Reference Example 4). (Yield: 96%). The dried
20 granules were subjected to sizing by passing to a sieve of 850 μm.

About 1% by weight of magnesium stearate was added to the sized granules and mixed, then the granules were supplied to a tablet machine (Rotary
25 single tablet press 12HUK: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), there were obtained tablets, each having 95 mg of weight.

85
219

- 67 -

Example 1

500.3 g of the aripiprazole hydrate crystals obtained in Reference Example 3 were milled using a sample mill (small atomizer). The main axis rotation rate was set to 12,000 rpm and the feed rotation rate to 17 rpm, and a 1.0 mm herringbone screen was used. Milling was completed in 3 minutes, resulting in 474.6 g (94.9%) of Aripiprazole Hydrate A powder.

The Aripiprazole Hydrate A (powder) obtained in this way had a mean particle size of 20-25 μm . The melting point (mp) was undetermined because dehydration was observed beginning near 70°C.

The Aripiprazole Hydrate A (powder) obtained above exhibited an ^1H -NMR (DMSO- d_6 , TMS) spectrum which was substantially the same as the ^1H -NMR spectrum shown in Figure 2. Specifically, it had characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, $J = 7.4$ Hz, 2H), 2.97 ppm (brt, $J = 4.6$ Hz, 4H), 3.92 ppm (t, $J = 6.3$ Hz, 2H), 6.43 ppm (d, $J = 2.4$ Hz, 1H), 6.49 ppm (dd, $J = 8.4$ Hz, $J = 2.4$ Hz, 1H), 7.04 ppm (d, $J = 8.1$ Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

The Aripiprazole Hydrate A (powder) obtained above had a powder x-ray diffraction spectrum which was substantially the same as the powder x-ray diffraction spectrum shown in Figure 3. Specifically, it had characteristic peaks at $2\theta = 12.6^\circ$, 15.4° , 17.3° , 18.0° ,

86
20

- 68 -

18.6°, 22.5° and 24.8°. This pattern is different from the powder x-ray spectrum of unmilled aripiprazole hydrate shown in Figure 7.

The Aripiprazole Hydrate A (powder) obtained above had infrared absorption bands at 2951, 2822, 1692, 1577, 1447, 1378, 1187, 963 and 784 cm⁻¹ on the IR (KBr) spectrum.

As shown in Figure 1, the Aripiprazole Hydrate A (powder) obtained above had a weak peak at 71.3°C in thermogravimetric/differential thermal analysis and a broad endothermic peak (weight loss observed corresponding to one water molecule) between 60-120°C--clearly different from the endothermic curve of unmilled aripiprazole hydrate (see Figure 6).

15

Example 2

450 g of the Aripiprazole Hydrate A (powder) obtained in Example 1 was dried for 24 hours at 100°C using a hot air dryer to produce 427 g (yield 98.7%) of Aripiprazole Anhydride Crystals B.

These Aripiprazole Anhydride Crystals B had a melting point (mp) of 139.7°C.

The Aripiprazole Anhydride Crystals B obtained above had an ¹H-NMR spectrum (DMSO-d₆, TMS) which was substantially the same as the ¹H-NMR spectrum shown in Figure 4. Specifically, they had characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm

25

87
221

- 69 -

(m, 4H + DMSO), 2.78 ppm (t, J = 7.4 Hz, 2H), 2.97 ppm (brt, J = 4.6 Hz, 4H), 3.92 ppm (t, J = 6.3 Hz, 2H), 6.43 ppm (d, J = 2.4 Hz, 1H), 6.49 ppm (dd, J = 8.4 Hz, J = 2.4 Hz, 1H), 7.04 ppm (d, J = 8.1 Hz, 1H), 7.11-
5 7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

The Aripiprazole Anhydride Crystals B obtained above had a powder x-ray diffraction spectrum which was substantially the same as the powder x-ray
10 diffraction spectrum shown in Figure 5. Specifically, they had characteristic peaks at $2\theta = 11.0^\circ$, 16.6° , 19.3° , 20.3° and 22.1° .

The Aripiprazole Anhydride Crystals B obtained above had remarkable infrared absorption bands
15 at 2945, 2812, 1678, 1627, 1448, 1377, 1173, 960 and 779 cm^{-1} on the IR (KBr) spectrum.

The Aripiprazole Anhydride Crystals B obtained above exhibited an endothermic peak near about 141.5°C in thermogravimetric/differential thermal
20 analysis.

The Aripiprazole Anhydride Crystals B obtained above exhibited an endothermic peak near about 140.7°C in differential scanning calorimetry.

Even when the Aripiprazole Anhydride Crystals
25 B obtained above were left for 24 hours in a dessicator set at humidity 100%, temperature 60°C , they did not exhibit hygroscopicity exceeding 0.4% (See Table 1 below).

88
22

- 70 -

Example 3

44.29 kg of the Aripiprazole Hydrate A (powder) obtained in Example 1 was dry heated for 18 hours in a 100°C hot air dryer and then heated for 3 hours at 120°C to produce 42.46 kg (yield 99.3%) of Aripiprazole Anhydride Crystals B.

The physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 2.

The Aripiprazole Anhydride Crystals B obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see Table 1 below).

Example 4

40.67 kg of the Aripiprazole Hydrate A (powder) obtained in Example 1 was dry heated for 18 hours in a 100°C hot air dryer and then heated for 3 hours at 120°C to produce 38.95 kg (yield 99.6%) of Aripiprazole Anhydride Crystals B.

The physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 2.

89
20

- 71 -

The Aripiprazole Anhydride Crystals B obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see 5 Table 1 below).

Examples 5-10 are useful for injectable or oral solution formulations but not solid dose formulations since they were made by heating Conventional Anhydride or Conventional Hydrate instead 10 of Hydrate A.

Example 5

The hygroscopic aripiprazole anhydride crystals obtained in Reference Example 1 were heated 15 for 50 hours at 100°C using the same methods as in Example 2. The physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 20 2.

The Aripiprazole Anhydride Crystals B obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see 25 Table 1 below).

Example 6

The hygroscopic aripiprazole anhydride

90
224

- 72 -

crystals obtained in Reference Example 1 were heated for 3 hours at 120°C using the same methods as in Example 2. The physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 2.

The Aripiprazole Anhydride Crystals B obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see Table 1 below).

Example 7

The hygroscopic aripiprazole anhydride crystals obtained in Reference Example 2 were heated for 50 hours at 100°C using the same methods as in Example 2. The physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 2.

The Aripiprazole Anhydride Crystals B obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see Table 1 below).

9/25

- 73 -

Example 8

The hygroscopic aripiprazole anhydride crystals obtained in Reference Example 2 were heated for 3 hours at 120°C using the same methods as in

5 Example 2. The physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example

2.

10 The Aripiprazole Anhydride Crystals B obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see Table 1 below).

15

Example 9

The aripiprazole hydrate crystals obtained in Reference Example 3 were heated for 50 hours at 100°C using the same methods as in Example 2. The

20 physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 2.

The Aripiprazole Anhydride Crystals B

25 obtained in this way did not exhibit hygroscopicity of more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see Table 1 below).

92
206

- 74 -

Example 10

The aripiprazole hydrate crystals obtained in Reference Example 3 were heated for 3 hours at 120°C using the same methods as in Example 2. The
5 physicochemical properties of the resulting Aripiprazole Anhydride Crystals B were the same as the physicochemical properties of the Aripiprazole Anhydride Crystals B obtained in Example 2.

The Aripiprazole Anhydride Crystals B
10 obtained in this way exhibited hygroscopicity of no more than 0.4% even when left for 24 hours in a dessicator set at humidity 100%, temperature 60°C (see Table 1 below).

15 Example 11 (Preparation of type C crystals of aripiprazole anhydride)

100 Miligrams of type-I crystals of aripiprazole anhydride obtained in Reference Example 2 were heated about 145°C ($\pm 3^\circ\text{C}$). In this occasion,
20 there was observed the phenomena that the crystals were once melted, then again crystallized. After that, 100 mg (yield: 100%) of Type C crystals of aripiprazole anhydride were obtained. The melting point of the crystals was 150°C. The crystals were colorless prism
25 form.

The type C crystals of aripiprazole anhydride obtained above had an endothermic curve which was substantially identical to the endothermic curve of

93
207

- 75 -

thermogravimetric/differential thermal analysis

(heating rate: 5°C/minute) shown in Figure 8.

Specifically, it showed the endothermic curve around 150.2°C.

5 The type C crystals of aripiprazole anhydride thus obtained exhibited an ¹H-NMR spectrum (DMSO-d₆, TMS) which was substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 9.

Specifically, it had the characteristic peaks at 1.55-

10 1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H+DMSO), 2.78 ppm (t, J=7.4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 15 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H), and 10.00 ppm (s, 1H).

 The type C crystals of aripiprazole anhydride obtained above had a powder X-ray diffraction spectrum which was substantially identical to the powder X-ray
20 diffraction spectrum shown in Figure 10. Specifically, it had the characteristic peaks at 2θ = 12.6°, 13.7°, 15.4°, 18.1°, 19.0°, 20.6°, 23.5° and 26.4°.

 The type C crystals of aripiprazole anhydride obtained above had an IR spectrum which was
25 substantially identical to the IR (KBr) spectrum shown in Figure 11. Specifically, it had the characteristic infrared absorption bands at 2939, 2804, 1680, 1375 and 780 cm⁻¹.

228

- 76 -

The type C crystals of aripiprazole anhydride obtained above exhibited a solid ^{13}C -NMR spectrum, which was substantially identical to the solid ^{13}C -NMR spectrum shown in Figure 12. Specifically, it had the 5 characteristic peaks at 32.8 ppm, 60.8 ppm, 74.9 ppm, 104.9 ppm, 152.2 ppm, 159.9 ppm and 175.2 ppm.

According to the above-mentioned data on endothermic curve of thermogravimetric/differential thermal analysis (heating rate: 5°C/minute) and powder 10 X-ray diffraction spectrum, the formation of the type C crystals of aripiprazole anhydride was confirmed.

When the type C crystals of aripiprazole anhydride crystals obtained above were left for 24 hours in a dessicator where the conditions were set at 15 humidity 100%, and temperature 60°C, then the crystals did not exhibit hygroscopicity higher than 0.4% (see, Table 1 below).

Example 12 (Preparation of type D crystals of 20 aripiprazole anhydride)

The type-I crystals of aripiprazole anhydride obtained in Reference Example 2 were added in 200 ml of toluene, and dissolved by heating at 74°C. After confirmed that it was dissolved completely, the toluene 25 solution was cooled to 7°C, and the precipitated crystals were collected by filtration. The crystals were subjected to air-drying as they were so as to obtain 17.9 g (yield: 89.5%) of type D crystals of

229

- 77 -

aripiprazole anhydride.

The type D crystals of aripiprazole anhydride obtained above had an endothermic curve substantially identical to the endothermic curve of

5 thermogravimetric/differential thermal analysis (heating rate: 5°C/minute) shown in Figure 13.

Specifically, it had the endothermic peaks at about 136.8°C and about 141.6°.

The type D crystals of aripiprazole anhydride
10 obtained above exhibited ¹H-NMR spectrum (DMSO-d₆, TMS) which was substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 14. Specifically, they had the characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56
15 ppm (m, 4H+DMSO), 2.78 ppm (t, J=7.4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H), and 10.00 ppm (s, 1H).

20 The type D crystals of aripiprazole anhydride obtained above had a powder X-ray diffraction spectrum which was substantially identical to the powder X-ray diffraction spectrum shown in Figure 15. Specifically, it had the characteristic peaks at 2θ = 8.7°, 11.6°,
25 16.3°, 17.7°, 18.6°, 20.3°, 23.4° and 25.0°.

The type D crystals of aripiprazole anhydride obtained above had an IR spectrum which was substantially identical to the IR (KBr) spectrum shown

96
230

- 78 -

in Figure 16. Specifically, it had the characteristic infrared absorption bands at 2946, 1681, 1375, 1273, 1175 and 862 cm^{-1} .

The type D crystals of aripiprazole anhydride
5 obtained above exhibited a solid ^{13}C -NMR spectrum which was substantially identical to the solid ^{13}C -NMR spectrum shown in Figure 17. Specifically, it had the characteristic peaks at 32.1 ppm, 62.2 ppm, 66.6 ppm, 104.1 ppm, 152.4 ppm, 158.5 ppm and 174.1 ppm.

10 According to the above-mentioned data on the endothermic curve of thermogravimetric/differential thermal analysis (heating rate: 5°C/minute) and powder X-ray diffraction spectrum, the formation of type D crystals of aripiprazole anhydride was confirmed.

15 When the type D crystals of aripiprazole anhydride crystals obtained above were left for 24 hours in a dessicator where the conditions were set at humidity 100%, and temperature 60°C, the crystals did not have hygroscopicity higher than 0.4% (see, Table 1
20 below).

Example 13 (Preparation of type D crystals of
aripiprazole anhydride)

1,200 Grams of the type-I crystals of
25 aripiprazole anhydride obtained in Reference Example 2 were dissolved in 18 liters of toluene, with heating. This toluene solution was cooled to 40°C, and 36 g of type-D crystals of aripiprazole anhydride obtained in

231 97

- 79 -

Example 12 were added as seed crystals, then the solution was cooled to 10°C and allowed to stand as it is. The precipitated crystals were collected by filtration, dried at 60°C for 18 hours to obtain 1,073 g (yield: 86.8%) of type D crystals of aripiprazole anhydride (purity: 100%). The crystals were colorless plate form.

The type D crystals of aripiprazole anhydride had an endothermic curve substantially identical to the endothermic curve of thermogravimetric/differential thermal analysis (heating rate: 5°C/minute) shown in Figure 13. Specifically, it had the endothermic peaks around about 136.8°C and about 141.6°.

The type D crystals of aripiprazole anhydride obtained above exhibited an ¹H-NMR spectrum (DMSO-d₆, TMS) which was substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 14.

Specifically, it had the characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H+DMSO), 2.78 ppm (t, J=7.4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H), and 10.00 ppm (s, 1H).

The type D crystals of aripiprazole anhydride obtained above had a powder X-ray diffraction spectrum which was substantially identical to the powder X-ray

232-98

- 80 -

diffraction spectrum shown in Figure 15. Specifically, it had the characteristic peaks at $2\theta = 8.7^\circ$, 11.6° , 16.3° , 17.7° , 18.6° , 20.3° , 23.4° and 25.0° .

The type D crystals of aripiprazole anhydride
5 obtained above had an IR spectrum which was substantially identical to the IR (KBr) spectrum shown in Figure 16. Specifically, it had characteristic infrared absorption bands at 2946, 1681, 1375, 1273, 1175 and 862 cm^{-1} .

10 The type D crystals of aripiprazole anhydride obtained above had a solid ^{13}C -NMR spectrum which was substantially identical to the solid ^{13}C -NMR spectrum shown in Figure 17. Specifically, it had the characteristic peaks at 32.1 ppm, 62.2 ppm, 66.6 ppm,
15 104.1 ppm, 152.4 ppm, 158.5 ppm and 174.1 ppm.

According to the above-mentioned data on the endothermic curve of thermogravimetric/differential thermal analysis (heating rate: $5^\circ\text{C}/\text{minute}$) and powder X-ray diffraction spectrum, the formation of type D
20 crystals of aripiprazole anhydride was confirmed.

When the type D crystals of aripiprazole anhydride crystals obtained above were left for 24 hours in a dessicator where the conditions were set at humidity 100%, and temperature 60°C , the crystals did
25 not exhibit hygroscopicity higher than 0.4% (see, Table 1 below).

99
23

- 81 -

Example 14 (Preparation of type E crystals of
aripiprazole anhydride)

40 Grams of type-I crystals of aripiprazole
anhydride obtained in Reference Example 2 was dissolved
5 in 1000 ml of acetonitrile with heating at 80°C. This
acetonitrile solution was cooled to about 70°C by
taking for about 10 minutes, and was kept at this
temperature for about 30 minutes to precipitate the
seed crystals. Next, the temperature of said solution
10 was slowly risen to 75°C, and the crystals were grown
up by keeping this temperature for 1 hour. Then, the
solution was cooled to 10°C by taking about 4 hours,
and the precipitated crystals were collected by
filtration. Thus obtained crystals were subjected to
15 air-drying overnight, there were obtained 37.28 g
(yield: 93.2%) of type E crystals of aripiprazole
anhydride (purity: 100%). The melting point of these
crystals was 145°C, and the crystals were colorless
needle form.

20 The type E crystals of aripiprazole
anhydride had an endothermic curve substantially
identical to the endothermic curve of
thermogravimetric/differential thermal analysis
(heating rate: 5°C/minute) shown in Figure 18.
25 Specifically, it had endothermic peak at about 146.5°.

The type E crystals of aripiprazole anhydride
obtained above exhibited an ¹H-NMR spectrum (DMSO-d₆,
TMS) which was substantially identical to the ¹H-NMR

- 82 -

spectrum (DMSO- d_6 , TMS) shown in Figure 19.

Specifically, it had the characteristic peaks at 1.55-1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H+DMSO), 2.78 ppm (t, $J=7.4$ Hz, 2H), 2.97 ppm (brt, $J=4.6$ Hz, 4H), 3.92 ppm (t, $J=6.3$ Hz, 2H), 6.43 ppm (d, $J=2.4$ Hz, 1H), 6.49 ppm (dd, $J=8.4$ Hz, $J=2.4$ Hz, 1H), 7.04 ppm (d, $J=8.1$ Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H), and 10.00 ppm (s, 1H).

10 The type E crystals of aripiprazole anhydride obtained above had a powder X-ray diffraction spectrum which was substantially identical to the powder X-ray diffraction spectrum shown in Figure 20. Specifically, it had the characteristic peaks at $2\theta = 8.0^\circ$, 13.7° ,
15 14.6° , 17.6° , 22.5° and 24.0° .

 The type E crystals of aripiprazole anhydride obtained above had an IR spectrum which was substantially identical to the IR (KBr) spectrum shown in Figure 21. Specifically, it had the characteristic
20 infrared absorption bands at 2943, 2817, 1686, 1377, 1202, 969 and 774 cm^{-1} .

 According to the data on the endothermic curve of thermogravimetric/differential thermal analysis (heating rate: $5^\circ\text{C}/\text{minute}$) and powder X-ray
25 diffraction spectrum, the formation of type E crystals of aripiprazole anhydride was confirmed.

 When the type E crystals of aripiprazole anhydride obtained above were left for 24 hours in a

101
035

- 83 -

dessicator where the conditions were set at humidity 100%, and temperature 60°C, the crystals did not exhibit hygroscopicity higher than 0.4% (see, Table 1 below).

5

Example 15 (Preparation of type F crystals of aripiprazole anhydride)

140 Grams of type-I crystals of aripiprazole anhydride obtained in Reference Example 2 were
10 suspended in 980 ml of acetone and continued to reflux for 7.5 hours with stirring. Next, the suspension was filtered in hot condition, and crystals separated out were subjected to air-drying for 16 hours at room temperature, there was obtained 86.19 g (yield: 61.6%)
15 of type F crystals of aripiprazole anhydride (purity: 100%). The crystals were colorless prism form.

The type F crystals of aripiprazole anhydride had an endothermic curve substantially identical to the endothermic curve of
20 thermogravimetric/differential thermal analysis (heating rate: 5°C/minute) shown in Figure 22. Specifically, it had the exothermic peaks at about 137.5°C and about 149.8°C.

The type F crystals of aripiprazole anhydride
25 obtained above exhibited an ¹H-NMR spectrum (DMSO-d₆, TMS) which was substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 23. Specifically, it had the characteristic peaks at 1.55-

102
236

- 84 -

1.63 ppm (m, 2H), 1.68-1.78 ppm (m, 2H), 2.35-2.46 ppm (m, 4H), 2.48-2.56 ppm (m, 4H+DMSO), 2.78 ppm (t, J=7.4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11-7.17 ppm (m, 1H), 7.28-7.32 ppm (m, 2H), and 10.00 ppm (s, 1H).

The type F crystals of aripiprazole anhydride obtained above had a powder X-ray diffraction spectrum which was substantially identical to the powder X-ray diffraction spectrum shown in Figure 24. Specifically, it had the characteristic peaks at $2\theta = 11.3^\circ$, 13.3° , 15.4° , 22.8° , 25.2° and 26.9° .

The type F crystals of aripiprazole anhydride obtained above had an IR spectrum which was substantially identical to the IR (KBr) spectrum shown in Figure 25. Specifically, it had the characteristic infrared absorption bands at 2940, 2815, 1679, 1383, 1273, 1177, 1035, 963 and 790 cm^{-1} .

According to the data on endothermic curve of thermogravimetric/differential thermal analysis (heating rate: $5^\circ\text{C}/\text{minute}$) and powder X-ray diffraction spectrum, the formation of type F crystals of aripiprazole anhydride was confirmed.

When the type F crystals of aripiprazole anhydride crystals obtained above were left for 24 hours in a dessicator where the conditions were set at humidity 100%, and temperature 60°C , the crystals did

193
237

- 85 -

not exhibit hygroscopicity higher than 0.4% (see, Table 1 below).

Table 1

Sample	Initial Moisture Content (%)	Moisture Content After 24 hrs (%)
Reference Example 1	0.04	3.28
Reference Example 2	0.04	1.78
Example 2	0.04	0.04
Example 3	0.02	0.02
Example 4	0.02	0.02
Example 5	0.04	0.04
Example 6	0.04	0.04
Example 7	0.04	0.03
Example 8	0.04	0.03
Example 9	0.03	0.01
Example 10	0.05	0.05
Example 11	0.03	0.03
Example 12	0.04	0.03
Example 13	0.04	0.03
Example 14	0.06	0.09
Example 15	0.04	0.04

5 Example 16

- a) Type I crystals of aripiprazole anhydride (10 g) obtained in Reference Example 2 was charged in a stainless steel round tray (diameter: 80 mm), and heated to about 170°C so as to melted completely. When this melted liquid was cooled, then it solidified

104
238

- 86 -

clarity with pale brown in color, the solid was peeled off from the stainless steel round tray, there was obtained 9.8 g (yield: 98%) of glassy state of aripiprazole anhydride. The obtained glassy state product was characterized by having no significant peak observed in a powder X-ray determination. (cf. Figure 31).

According to the thermogravimetric/differential thermal analysis (heating rate: 5°C/minute), as shown in Figure 30, an exothermic peak of type B crystals of aripiprazole anhydride was observed at around 86.5°C. While, an endothermic peak of type B crystals of aripiprazole anhydride owing to melting was observed at around 140.1°C.

b) When the glassy state of aripiprazole anhydride obtained in Example 16-a) were charged in a sealed vessel and left to stand at room temperature for about 6 months, then type G crystals of aripiprazole anhydride having white in color was obtained by changing the color from pale brown (25 g, yield: 100%). Melting point: 138 to 139°C.

The type G crystals of aripiprazole anhydride had an endothermic curve which was substantially identical to the thermogravimetric/differential thermal analysis (heating rate: 5°C/min.) endothermic curve shown in Figure 26, more particularly, it has an endothermic peak around 141.0°C and an exothermic peak around 122.7°C.

108
39

- 87 -

The type G crystals of aripiprazole anhydride obtained as above exhibited an ¹H-NMR spectrum which was substantially identical to the ¹H-NMR spectrum (DMSO-d₆, TMS) shown in Figure 27. Specifically, it has

5 characteristic peaks at 1.55 - 1.63 ppm (m, 2H), 1.68 - 1.78 ppm (m, 2H), 2.35 - 2.46 ppm (m, 4H), 2.48 - 2.56 ppm (m, 4H + DMSO), 2.78 ppm (t, J=7.4 Hz, 2H), 2.97 ppm (brt, J=4.6 Hz, 4H), 3.92 ppm (t, J=6.3 Hz, 2H), 6.43 ppm (d, J=2.4 Hz, 1H), 6.49 ppm (dd, J=8.4 Hz, J=2.4 Hz, 1H), 7.04 ppm (d, J=8.1 Hz, 1H), 7.11 - 7.17 ppm (m, 1H), 7.28 - 7.32 ppm (m, 2H) and 10.00 ppm (s, 1H).

The type G crystals of aripiprazole anhydride obtained as above had a powder X-ray diffraction

15 spectrum which was substantially identical to the powder X-ray diffraction spectrum shown in Figure 28. Specifically, it has characteristic peak at 2θ = 10.1°, 12.8°, 15.2°, 17.0°, 17.5°, 19.1°, 20.1°, 21.2°, 22.4°, 23.3°, 24.5° and 25.8°.

20 The type G crystals of aripiprazole anhydride obtained above had an IR spectrum which was substantially identical to the IR (KBr) spectrum shown in Figure 29. Specifically, it has clear infrared absorption bands at 2942, 2813, 1670, 1625, 1377, 1195, 25 962 and 787 cm⁻¹.

Example 17

a) Preparation of granules of 30 mg tablets containing

100
240

- 88 -

type B crystals of aripiprazole anhydride for additional drying

Type B crystals of aripiprazole anhydride (1,500 g), lactose (5,700 g), corn starch (1,000 g) and 5 crystalline cellulose (1,000 g) were charged in a fluidized bed granulating dryer (Flow Coater Model FLO-5M; manufactured by FROINT SANGYO KABUSHIKI KAISHA), and these granulating ingredients were mixed by fluidizing for about 3 minutes with an inlet air 10 temperature at 60°C, air flow rate of 3 to 4 m³/min. Further, the granulating ingredients were continued fluidizing under the same condition, and sprayed with about 4,000 g of 5% aqueous solution of hydroxypropyl cellulose to obtain wet granules. The wet granules were 15 dried under an inlet air temperature at 85°C, for about 20 minutes. The obtained dried granules contained 3.8% of water (measured by the method according to Reference Example 4).

b) The dried granules (4 kg) obtained in Example 20 17-a) were sized by use of a mill (FIORE F-0: manufactured by TOKUJU CORPORATION).

The sized granules (3 kg) were charged in a fluidized bed granulating dryer (Flow Coater Model FLO-5M; manufactured by FREUND INDUSTRIAL CO., LTD.), and 25 these granulating ingredients were dried under an inlet air temperature at 85°C, and air flow rate of 2 m³/min for 2 hours. The obtained dried granules contained 3.6% of water (measured by the method according to

107
241

- 89 -

Reference Example 4).

About 1% by weight of magnesium stearate was added to the sized granules and mixed, then the granules were supplied to a tabletting machine (a

- 5 Rotary single tablet press, Model VIRGO: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and there were obtained tablets, each having 190 mg of weight.

- c) The dried granules (3 kg) obtained in Example 17-a) were charged in a vacuum dryer (vacuum
10 granulating dryer model; VG-50: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and dried at 70°C of a jacket temperature, under a reduced pressure at 5 torr of degree of vacuum for 1 hour. The thus obtained dried
15 granules contained 3.1% of water (measured by the method according to Reference Example 4). The dried granules were subjected to sizing by passing to a sieve of 850 μ m.

- About 1% by weight of magnesium stearate was added to the sized granules and mixed, then the
20 granules were supplied to a tablet machine (Rotary single tablet press, Model VIRGO: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and there were obtained tablets, each having 190 mg of weight.

25 Example 18

- a) Preparation of 30 mg tablets containing type B crystals of aripiprazole anhydride

Aripiprazole anhydride (type B crystals)

108
242

- 90 -

(4,500 g), lactose (17,100 g), corn starch (3,000 g) and crystalline cellulose (3,000 g) were charged in a fluidized bed granulating dryer (NEW-MARUMERIZER Model: NQ-500, manufactured by FUJI PAUDAL CO., LTD.), and
5 these granulating ingredients were mixed by fluidizing for about 3 minutes with an inlet air temperature at 70°C, air flow rate of 10 - 15 m³/min. Further, the granulating ingredients were continued fluidizing under the same condition, and were sprayed with about 12,000
10 g of 5% aqueous solution of hydroxypropyl cellulose to obtain wet granules. The wet granules were dried under inlet air at temperature of 85°C, for about 30 minutes. The obtained dried granules contained 3.6% of water (measured by the method according to Reference Example
15 4). (Yield: 96%). The dried granules were sized by passing to a mill (FIOLE F-0: manufactured by TOKUJU CORPORATION).

About 1% by weight of magnesium stearate was added to the sized granules and mixed, then the
20 granules were supplied to a tablet machine (a Rotary single tablet press, VIRGO: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and there were obtained tablets, each having 190 mg of weight.

b) The tablets (5 kg) obtained in Example 18-a)
25 were charged in a fan dryer (AQUA COATER AQC-48T, manufactured by FREUND INDUSTRIAL CO., LTD.), and dried under inlet air at temperature of 90°C, air flow rate of 2 m³/min for 6 hours. The obtained dried granules

b9
243

- 91 -

contained 3.3% of water (measured by the method according to Reference Example 4).

- c) The dried tablets (3 kg) obtained in Example 18-a) were charged in a vacuum dryer (vacuum granulating dryer, VG-50: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and dried at 80°C of a jacket temperature, under reduced pressure of 5 torr of degree of vacuum for 4 hours. The obtained dried tablets contained 2.7% of water (measured by the method according to Reference Example 4).

Example 19

- a) By the procedures similar to those of Example 18-a), there were obtained tablets (containing type I crystals of aripiprazole anhydride obtained in Reference Example 2), each having 190 mg of weight,
- b) The tablets were dried by the procedures similar to those of Example 18-b), except that air inlet temperature was 100°C and dried for 1 hour.
- c) The tablets were dried by the procedures similar to those of Example 18-b), except that inlet air temperature was 100°C and dried for 3 hours.

Example 20

- By the procedures similar to those of Example 18-a), there were obtained tablets, each having 190 mg of weight, containing type C crystals of aripiprazole anhydride.

- 92 -

Example 21

By the procedures similar to those of Example 18-a), there were obtained tablets, each having 190 mg of weight, containing type D crystals of aripiprazole anhydride.

Example 22

a) Aripiprazole hydrate crystals (156 g) obtained in Reference Example 3, lactose (570 g), corn starch (100 g) and crystalline cellulose (100 g) were charged in a fluidized bed granulating dryer (NEW-MARUMERIZER, NQ-160: manufactured by FUJI POWDAL CO., LTD.), and these granulating ingredients were mixed under fluidizing for about 3 minutes with an inlet air temperature at 60°C, air flow rate of 1.0 to 1.5 m³/min, and rotating disc with rotary speed of 400 rpm.

Further, the granulating ingredients were continued fluidizing under the same condition, and sprayed about 500 g of 4% aqueous solution of hydroxypropyl cellulose to obtain wet granules. The inlet air temperature was elevated up to 85°C, and dried until the temperature of the product was reached to 46°C. The obtained dried granules were sized by passing to a sieve of 850 μm. The dried granules contained 4.37% of water (measured by the method according to Reference Example 4).

b) The dried granules (200 g) obtained in Example 22-a) were charged in a fluidized bed dryer (multiplex, MP-01: manufactured by POWREX CORPORATION),

14
245

- 93 -

and dried at 85°C of inlet air temperature, air flow rate of 0.5 m³/min for 2 hours. The dried granules contained 3.50% of water (measured by the method according to Reference Example 4).

5 c) The dried granules (100 g) obtained in Example 22-a) were charged in a vacuum dryer (vacuum granulating dryer LCV-232: manufactured by TABAI CO., LTD.), and dried 80°C of tray temperature, about 760 mmHg of degree of vacuum for 2 hours. The dried
10 granules were further dried similarly for 6 hours. The dried granules contained 3.17% of water (the product being dried for 2 hours: measured by the method according to Reference Example 4). The further dried granules contained 2.88% of water (the product being
15 dried for 6 hours: measured by the method according to Reference Example 4).

d) About 1% by weight of magnesium stearate was added to the sized granules being obtained in Example 22-b) and mixed, then the mixed granules were supplied
20 to a tablet machine (Single type Tablet machine No. 2B: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and tableted with punch, there were obtained tablets, each having 191 mg of weight.

e) About 1% by weight of magnesium stearate was
25 added to the sized granules being obtained in Example 22-c) and mixed, then the mixed granules were supplied to a tablet machine (Single type Tablet machine No. 2B: manufactured by KIKUSUI SEISAKUSHO CO., LTD.), and

112
246

- 94 -

tabletted with punch, there were obtained tablets, each having 191 mg of weight.

Dissolution Test

- 5 Each tablets of the pharmaceutical solid oral preparations obtained previously was kept, repectively under the open at 25°C/60% RH for 6 months, and at 40°C/75% RH for 1 week, then their dissolution rates were measured by the following methods. The
- 10 dissolution rates obtained from 60 minutes after the exposure are shown in Tables 2 and 3. The dissolution rates after 60 minutes, using the tablets kept under the open at 40°C/75% RH for 2 weeks, are shown in Tables 4 and 5. The dissolution rates after 60
- 15 minutes, using the tablets kept under the open condition at 40°C/75% RH for 1 week, are shown in Table 6.
- Dissolution test equipment: USP
- Model: NTR-6100 (manufactured by TOYAMA SANGYO CO., LTD.)
- 20 Model: DT-610 (manufactured by JASCO CORPORATION)

a) Method of dissolution test of the 15 mg tablet

- One tablet (containing 15 mg each of aripiprazole anhydride or hydrate) was tested by using
- 25 900 ml of acetic acid buffer solution (pH 5.0) (Note: 1) as the test solution, and by rotating a paddle at 100 rpm according to the method of USP (United States Pharmacopoea) (Note: 2).

112
249

- 95 -

The test solutions obtained respectively from 10 minutes, 20 minutes, 30 minutes, 45 minutes and 60 minutes after the start of test are named as T10, T20, T30, T45 and T60.

5 On the other hand, about 0.05 g of standard sample of aripiprazole was weighed accurately, dissolved in ethanol (95%) so as to make exactly 50 ml of ethanol solution. Twenty (20) ml of this ethanol solution was taken accurately, and to prepared exactly
10 1000 ml of the standard solution by adding 0.01 mol/liter of hydrochloric acid reagent solution (Note: 3).

The test solutions and the standard solution were subjected to filtration, respectively by using a filter having micropores of 10 to 20 μ m in diameters,
15 then each of the filtrates were introduced to a spectrophotometer installed with flow cell (cell length: 10 mm), and to measure the absorbance of wave length at 249 nm and absorbance of wave length at 325 nm and determined the differences between absorbances
20 to named as At10, At20, At30, At45, At60 and As, respectively.

After the measurements, the test solutions of T10, T20, T30 and T45 were put back to the test vessels respectively. Further, similar procedures were
25 conducted to other 5 samples of the test solutions.

Dissolution rate (%) relating to the indicated amount of aripiprazole =

114
248

- 96 -

Amount of the standard sample of aripiprazole (mg)

$\times A_t \times A_s \times 9/5 \times 20/C$

wherein, A_t : A_{t10} , A_{t20} , A_{t30} , A_{t45} or A_{t60}

A_s : standard solution

5 C : Indicated amount of aripiprazole (mg)

(Note:1) Water was added to 1.97 g of acetic acid (100)
and 9.15 g of sodium acetate trihydrate to
make 1000 ml of solution (0.1 mol/l).

(Note:2) Paddle method

10 (Note:3) Water was added to 100 ml of 0.1 mol/l
hydrochloric acid (Note:4) to make 1000 ml of
solution.

(Note:4) Water was added to 0.9 ml of hydrochloric acid
to make 1000 ml of solution.

15

b) Method of dissolution test of the 30 mg tablet

One tablet each of the pharmaceutical solid
oral preparations (containing 30 mg each of
aripiprazole anhydride or hydrate) was tested by using
20 900 ml of acetic acid buffer solution (pH 4.5) (Note:
5) as the test solution, and to conduct the test by
rotating a paddle at 75 rpm in accordance with the
method of USP (United States Pharmacopoeia) (Note: 6).

The test solutions obtained respectively from
25 10 minutes, 20 minutes, 30 minutes 45 minutes and 60
minutes after the start of test, were named as T_{10} ,
 T_{20} , T_{30} , T_{45} and T_{60} .

On the other hand, about 0.05 g of the

- 97 -

standard sample of aripiprazole was weighed accurately,
and dissolved in ethanol (95%) so as to made exactly 50
ml of the ethanol solution. Twenty (20) ml of the
ethanol solution was taken accurately, and prepared
5 exactly 1000 ml of the standard solution by adding 0.01
mol/liter of hydrochloric acid reagent solution (Note:
7).

The test solutions and standard solution were
subjected to filtration, respectively by using a filter
10 having micropores of 10 to 20 μ m in diameters, then
each of the filtrates were introduced to a
spectrophotometer in which a flow cell (cell length: 10
mm) was installed, and measured the absorbance of wave
length at 249 nm and absorbance of wave length at 325
15 nm, and the difference between these absorbances were
named as At10, At20, At30, At45, At60 and As,
respectively.

After the measurements, the test solutions of
T10, T20, T30 and T45 were put back respectively to the
20 test vessels. Further, similar procedures were
conducted to other 5 samples of the test solutions.

Dissolution rate (%) relating to the indicated
amount of aripiprazole =
25 Amount of the standard sample of aripiprazole (mg)
 $\times At \times As \times 9/5 \times 20/C$
wherein, At: At10, At20, At30, At45 or At60
As: standard solution

- 98 -

C: Indicated amount of aripiprazole (mg)

(Note:5) Water was added to 1.91 g of acetic acid (100) and 2.99 g of sodium acetate trihydrate to made 1000 ml of solution (0.05 mol/l).

5 (Note:6) Paddle method

(Note:7) Water is added to 100 ml of 0.1 mol/l hydrochloric acid (Note:8) to made 1000 ml of solution.

(Note:8) Water was added to 0.9 ml of hydrochloric acid
10 to make 1000 ml of solution.

Table 2

Samples used	Open at 25°C/60% RH		Open at 40°C/75% RH	
	Initial	After 6 months	Initial	After 1 week
Tablet (15 mg) of Reference Example 4	83.4%	44.3%	83.4%	44.1%
Tablet (15 mg) of Reference Example 5	90.1%	61.9%	90.1%	65.2%

15

Table 3

Samples used	Open at 25°C/60% RH		Open at 40°C/75% RH	
	Initial	After 6 months	Initial	After 1 week
Tablet (30 mg) of Example 18-a)	96.7%	77.1%	96.7%	75.9%
Tablet (30 mg) of Example 17-b)	96.5%	93.6%	95.0%	92.2%
Tablet (30 mg) of Example 17-c)	97.0%	96.3%	94.7%	94.8%
Tablet (30 mg) of Reference Example 18-b)	97.2%	95.3%	97.2%	97.8%
Tablet (30 mg) of Reference Example 18-c)	97.8%	96.3%	97.8%	96.9%

117
251

- 99 -

Table 4

Samples used	Initial	After 2 weeks
Samples used Tablet (30 mg) of Example 19-a)	89.8%	66.9%
Tablet (30 mg) of Example 19-b)	-	79.8%
Tablet (30 mg) of Example 19-c)	-	85.9%

5

Table 5

Samples used	Initial	After 2 weeks
Tablet (30 of Example 18-a)	94.8%	94.7%
Tablet (30 mg) of Example 20	93.7%	93.1%
Tablet (30 mg) of Example 21	94.8%	90.9%

10

Table 6

Samples used	Initial	After 1 weeks
Tablet (30 mg) of Example 22-d)	96.5%	84.5%
Tablet (30 mg) of Example 22-e) (dried for 2 hours)	92.5%	74.4%
Tablet (30 mg) of Example 22-e) (dried for 6 hours)	96.2%	83.4%

(Note: Dissolution tests in Table 5 were conducted

15

similarly to the procedures in the above-mentioned "b) Method of dissolution test of the 30 mg tablet" except that by using 900 ml of acetic acid buffer solution (pH 4.0) as the test solution, and by rotating a paddle at 50 rpm.

48
52

- 100 -

As can be seen clearly from the data shown in Table 2, in comparison with the 15 mg tablet containing conventional aripiprazole anhydride crystals (Reference Example 4), the 15 mg tablet containing type B crystals of aripiprazole anhydride (Reference Example 5) had the dissolution rate to maintain maximum drug concentration (C_{max}), at pH 5.0 after 60 minutes, even though such tablet was kept under the open at 25°C/60%RH for 6 months and under the open at 40°C/75%RH for 1 week.

As can be seen clearly from the data shown in Table 3, even though 30 mg tablets (Examples 17-b) and 17-c)) prepared from twice dried granules of type B crystals of aripiprazole anhydride, and 30 mg tablets (Examples 18-b) and 18-c)) prepared from further dried pharmaceutical solid oral preparation containing type B crystals of aripiprazole anhydride were subjected to keep under the open at 25°C/60%RH for 6 months or 40°C/75%RH for 1 week, the dissolution rates of these tablets obtained 60 minutes after the test at pH 4.5 were not substantially lowered.

As can be seen clearly from the data shown in Table 4, when 30 mg tablets (Examples 19-a), 19-b) and 19-c)) containing conventional aripiprazole anhydride crystals were further dried and subjected to keep under open at 40°C/75%RH for 2 weeks, then the dissolution rates of the tablets obtained 60 minutes after the test at pH 4.5 were the dissolution rates to maintain

119
253

- 101 -

maximum drug concentration (C_{max}).

As can be seen clearly from the data shown in Table 5, when 30 mg tablet (Example 18-a)) containing type B crystals of aripiprazole anhydride, 30 mg tablet (Example 20) containing type C crystals of aripiprazole anhydride and 30 mg tablet (Example 21) containing type D crystals of aripiprazole anhydride were subjected to keep under open at 40°C/75%RH for 2 weeks, then the dissolution rates of the tablets obtained 60 minutes after the test at pH 4.0 were not substantially lowered.

As can be seen clearly from the data shown in Table 6, when 30 mg tablets (Examples 22-d) and 22-e)) prepared from granules of conventional aripiprazole hydrate being twice dried, and subjected to keep under open at 40°C/75%RH for 1 week, then the dissolution rates of the tablets obtained 60 minutes after the test at pH 4.5 were the dissolution rates to maintain maximum drug concentration (C_{max}).

20

Sample Preparation 1

	Aripiprazole anhydride crystals B	5 mg
	Starch	131 mg
	Magnesium stearate	4 mg
25	<u>Lactose</u>	<u>60 mg</u>
	Total	200 mg

120
64

- 102 -

Tablets containing the above ingredients in each tablet were prepared by formulation methods known to one skilled in the art of pharmaceutical formulation.

5

Sample Preparation 2

	Type C crystals of aripiprazole anhydride	5 mg
	Starch	131 mg
	Magnesium stearate	4 mg
10	<u>Lactose</u>	<u>60 mg</u>
	Total	200 mg

In accordance with an ordinary method, tablet preparation, containing the above-mentioned ingredients per 1 tablet was prepared.

15

Sample Preparation 3

	Type D crystals of aripiprazole anhydride	5 mg
	Starch	131 mg
20	Magnesium stearate	4 mg
	<u>Lactose</u>	<u>60 mg</u>
	Total	200 mg

In accordance with an ordinary method, tablet preparation, containing the above-mentioned ingredients per 1 tablet was prepared.

25

121
255

- 103 -

Sample Preparation 4

	Type E crystals of aripiprazole anhydride	5 mg
	Starch	131 mg
	Magnesium stearate	4 mg
5	<u>Lactose</u>	<u>60 mg</u>
	Total	200 mg

In accordance with an ordinary method, tablet
preparation, containing the above-mentioned ingredients
10 per 1 tablet was prepared.

Sample Preparation 5

	Type F crystals of aripiprazole anhydride	5 mg
	Starch	131 mg
15	Magnesium stearate	4 mg
	<u>Lactose</u>	<u>60 mg</u>
	Total	200 mg

In accordance with an ordinary method, tablet
20 preparation, containing the above-mentioned ingredients
per 1 tablet was prepared.

Sample Preparation 6

	Type G crystals of aripiprazole anhydride	5 mg
25	Starch	131 mg
	Magnesium stearate	4 mg
	<u>Lactose</u>	<u>60 mg</u>
	Total	200 mg

122
56

- 104 -

In accordance with an ordinary method, tablet preparation, containing the above-mentioned ingredients per 1 tablet was prepared.

5 Formulation Example

The following examples used aripiprazole drug substance made by first milling or pulverizing the conventional hydrate of aripiprazole and then heating it to form the anhydrous form (anhydride B).

10

Formulation Example 1

Flash-melt tablets were prepared as follows:

Intragranulation:

15

Ingredient	Percent w/w	Mg. per tablet
Xylitol (300) Xylisorb	26	52
Avicel® PH 102	12	24
Calcium Silicate	43.35	86.7
Crospovidone	3	6
Amorphous silica	2	4
Aspartame	2	4
Wild cherry flavor	0.15	0.3
Tartaric acid	2	4
Acesulfame K	2	4
Magnesium stearate	0.25	0.5
Total weight	92.75	185.5

123
25X

- 105 -

The ingredients except for the magnesium stearate were blended in a commercial V-blender in geometric proportions for 5 minutes each until all were added. The magnesium stearate was then added and the mixture blended for an additional three minutes. The blended formulation was compacted at a pressure of 30-35 kgF/cm² in a commercial compactor equipped with an orifice such that the compacts therefrom are in the form of ribbons. The ribbons were passed through a 30 mesh (600 microns) screen to form stable granules of about 150 to 400 microns.

Extragranulation Ingredients:

Ingredient	Percent w/w	Mg. per tablet
Intragranulation	92.75	185.5
Avicel® PH 200	3	6
Crospovidone	4	8
Magnesium stearate	0.25	0.5
Total weight	100	200

15

The intragranulation was placed in the blender and the Avicel® PH 200 and crospovidone added thereto and blended for five minutes. The magnesium stearate was then added and the mixture blended for an additional three minutes to form the final blend. Tablets compressed therefrom had a breaking force of 2.3 kP (3.5 SCU) and disintegrated in 10 seconds in 5

124
258

- 106 -

ml of water. The final blend formulation demonstrated excellent flow and was free of other problems such as chipping, capping and sticking. It has been found that utilizing Avicel® PH 102 for the intragranulation and 5 Avicel® PH 200 for the extragranulation ingredient enhanced the quality of the resultant tablets.

Formulation Example 2

Flash-melt tablets containing a combination 10 of two grades of calcium silicate were prepared as follows:

Intragranulation:

Ingredient	Percent w/w	Mg. per tablet
Xylitol (300) Xylisorb	26	52
Avicel® PH 102	12	24
Calcium Silicate (crystalline, alpha triclinic)	33.35	66.7
Hubersorb 600 NF (amorphous calcium silicate)	10	20
Crospovidone	3	6
Amorphous silica	2	4
Aspartame	2	4
Wild cherry flavor	0.15	0.3
Tartaric acid	2	4
Acesulfame K	2	4
Magnesium stearate	0.25	0.5
Total weight	92.75	185.5

125
259

- 107 -

The ingredients except for the magnesium stearate were blended in a commercial V-blender in geometric proportions for 5 minutes each until all were added. The magnesium stearate was added and the mixture blended for an additional three minutes. The blended formulation was compacted, and screened to form stable granules in accordance with the procedure of Formulation Example 1.

10 Extragranulation Ingredients:

Ingredient	Percent w/w	Mg. per tablet
Intragranulation	92.75	185.5
Avicel® PH 200	3	6
Crospovidone	4	8
Magnesium stearate	0.25	0.5
Total weight	100	200

The intragranulation was placed in the blender and the Avicel® PH 200 and crospovidone added thereto and blended for five minutes. The magnesium stearate was then added and the mixture blended for an additional three minutes to form the final blend. Tablets compressed therefrom had a breaking force of 2.0 kP (3.1 SCU) and disintegrated in 10 seconds in 5 ml of water.

- 108 -

Formulation Example 3

Flash-melt tablets containing aripiprazole,
an antischizophrenic drug, were prepared as follows:

5 Intragranulation

Ingredient	Percent w/w	Mg. per tablet
Aripiprazole	15	30
Xylitol (300) Xylisorb	25	50
Avicel® PH 102	6	12
Calcium Silicate	37	74
Crospovidone	3	6
Amorphous silica	2	4
Aspartame	2	4
Wild cherry flavor	0.15	0.3
Tartaric acid	2	4
Acesulfame K	2	4
Magnesium stearate	0.25	0.5
Total weight	94.4	188.8

The ingredients except for the magnesium
stearate were blended in a commercial V-blender in
10 geometric proportions for 5 minutes each until all were
added. The magnesium stearate was added and the
mixture blended for an additional three minutes. The
blended formulation was compacted, and screened to form
stable granules in accordance with the procedure of
15 Formulation Example 1.

127
261

- 109 -

Extragranulation Ingredients:

Ingredient	Percent w/w	Mg. per tablet
Intragranulation	94.4	188.8
Avicel® PH 200	1.1	2.2
Crospovidone	4	8
Magnesium stearate	0.5	1
Total weight	100	200

The intragranulation was placed in the
5 blender and the Avicel® PH 200 and crospovidone added
thereto and blended for five minutes. The magnesium
stearate was then added and the mixture blended for an
additional three minutes to form the final blend.
Tablets compressed therefrom had a breaking force of
10 2.0 kP (3.1 SCU) and disintegrated in 10 seconds in 5
ml of water.

Formulation Example 4

Flash-melt tablets containing aripiprazole
15 were prepared as follows:

- 110 -

Intragranulation:

Ingredient	Percent w/w	Mg. per tablet
Aripiprazole	0.5	1
Xylitol (300) Xylisorb	27	54
Avicel® PH 102	12	24
Calcium Silicate	42	84
Crospovidone	3	6
Amorphous silica	2	4
Aspartame	2	4
Wild cherry flavor	0.15	0.3
Tartaric acid	2	4
Acesulfame K	2	4
Magnesium stearate	0.25	0.5
Total weight	92.9	185.8

The ingredients except for the magnesium
5 stearate were blended in a commercial V-blender in
geometric proportions for 5 minutes each until all were
added. The magnesium stearate was added and the
mixture blended for an additional three minutes. The
blended formulation was compacted, and screened to form
10 stable granules in accordance with the procedure of
Formulation Example 1.

129
263

- 111 -

Extragranulation Ingredients:

Ingredient	Percent w/w	Mg. per tablet
Intragranulation	92.9	185.8
Avicel® PH 200	2.6	5.2
Crospovidone	4	8
Magnesium stearate	0.5	1
Total weight	100	200

The intragranulation was placed in the
5 blender and the Avicel® PH 200 and crospovidone added
thereto and blended for five minutes. The magnesium
stearate was then added and the mixture blended for an
additional three minutes to form the final blend.
Tablets compressed therefrom had a breaking force of
10 2.3 kP (3.5 SCU) and disintegrated in 10 seconds in 5
ml of water.

130
64

- 112 -

CLAIMS

1. Hydrate A of aripiprazole wherein said Hydrate has a powder x-ray diffraction spectrum which is substantially the same as the following powder x-ray
5 diffraction spectrum shown in Figure 3.
2. Hydrate A of aripiprazole wherein said Hydrate has powder x-ray diffraction characteristic peaks at $2\theta = 12.6^\circ, 15.4^\circ, 17.3^\circ, 18.0^\circ, 18.6^\circ, 22.5^\circ$ and 24.8° .
- 10 3. Hydrate A of aripiprazole wherein said Hydrate has particular infrared absorption bands at 2951, 2822, 1692, 1577, 1447, 1378, 1187, 963 and 784 cm^{-1} on the IR (KBr) spectrum.
4. Hydrate A of aripiprazole wherein said
15 Hydrate has an endothermic curve which is substantially the same as the thermogravimetric/differential thermal analysis (heating rate $5^\circ\text{C}/\text{min}$) endothermic curve shown below shown in Figure 1.
5. Hydrate A of aripiprazole wherein said
20 Hydrate has a mean particle size of 50 μm or less.
6. Hydrate A of aripiprazole wherein said Hydrate has a mean particle size range of 36 to 14 μm .
7. Hydrate A of aripiprazole wherein said Hydrate has
25 a powder x-ray diffraction spectrum which is substantially the same as the following powder x-ray diffraction spectrum shown in Figure 3;
particular infrared absorption bands at 2951,

187
265

- 113 -

2822, 1692, 1577, 1447, 1378, 1187, 963 and 784 cm^{-1} on the IR (KBr) spectrum;

an endothermic curve which is substantially the same as the thermogravimetric/differential thermal analysis (heating rate 5°C/min) endothermic curve shown below shown in Figure 1; and

a mean particle size of 50 μm or less.

8. A process for the preparation of Hydrate A wherein said process comprises milling Conventional Hydrate to a mean particle size of 50 μm or less.
9. A process according to claim 8, wherein said milling is performed by an atomizer using a rotational speed of 5000-15000 rpm for the main axis, a feed rotation of 10-30 rpm and a screen hole size of 1-5 mm.
10. The Hydrate A according to claim 8 made by a process comprising milling Conventional Hydrate to a mean particle size of 50 μm or less.
11. The Hydrate A according to claim 8 made by a process comprising milling Conventional Hydrate to a mean particle size of 50 μm or less wherein said milling is performed by an atomizer using a rotational speed of 5000-15000 rpm for the main axis, a feed rotation of 10-30 rpm and a screen hole size of 1-5 mm.
12. Aripiprazole drug substance of low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.40% or less after placing said drug substance for 24 hours in a dessicator maintained at a temperature of 60°C and a humidity level of 100%.

132
266

- 114 -

13. Aripiprazole Anhydride Crystals B having low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.40% or less after placing said drug substance for 24 hours in a dessicator maintained
5 at a temperature of 60°C and a humidity level of 100%.
14. Aripiprazole drug substance of low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.10% or less after placing said drug substance for 24 hours in a dessicator maintained
10 at a temperature of 60°C and a humidity level of 100%.
15. Aripiprazole Anhydride Crystals B having low hygroscopicity wherein said low hygroscopicity is a moisture content of 0.10% or less after placing said drug substance for 24 hours in a dessicator maintained
15 at a temperature of 60°C and a humidity level of 100%.
16. Aripiprazole Anhydride Crystals B having a powder x-ray diffraction spectrum which is substantially the same as the following powder x-ray diffraction spectrum shown in Figure 5.
- 20 17. Aripiprazole Anhydride Crystals B having a powder x-ray diffraction spectrum having characteristic peaks at $2\theta = 11.0^\circ$, 16.6° , 19.3° , 20.3° and 22.1° .
18. Aripiprazole Anhydride Crystals B having a particular infrared absorption bands at 2945, 2812,
25 1678, 1627, 1448, 1377, 1173, 960 and 779 cm^{-1} on the IR (KBr) spectrum.
19. Aripiprazole Anhydride Crystals B exhibiting an endothermic peak near about 141.5°C in

123
267

- 115 -

thermogravimetric/differential thermal analysis
(heating rate 5°C/min).

20. Aripiprazole Anhydride Crystals B exhibiting
an endothermic peak near about 140.7°C in differential
5 scanning calorimetry (heating rate 5°C/min).

21. Aripiprazole Anhydride Crystals B wherein
said Crystals will not substantially convert to a
hydrous form of aripiprazole when properly stored under
a relative humidity (RH) of 60 % and at a temperature
10 of 25°C, even for an extended period being not less
than 4 years.

22. Aripiprazole Anhydride Crystal B wherein said
crystals has a mean particle size of 50 µm or less.

23. Aripiprazole Anhydride Crystal B wherein said
15 crystals has a mean particle size of 30 µm or less.

24. Aripiprazole Anhydride Crystals B having all
physicochemical properties defined in claims 16 and 18
to 22.

25. Aripiprazole Anhydride Crystals B having all
20 physicochemical properties defined in claims 17 to 22.

26. Aripiprazole Anhydride Crystals B having all
physicochemical properties defined in claims 13, 16, 18
to 20 and 22.

27. Aripiprazole Anhydride Crystals B having all
25 physicochemical properties defined in claims 15, 16, 18
to 20 and 22.

28. Aripiprazole Anhydride Crystals B having all
physicochemical properties defined in claims 13, 17 to

134
268

- 116 -

20 and 22.

29. Aripiprazole Anhydride Crystals B having all physicochemical properties defined in claims 15, 17 to 20 and 22.

5 30. A process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A.

31. A process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises
10 heating Aripiprazole Hydrate A at 90-125°C for about 3-50 hours.

32. A process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 100°C for about 18
15 hours.

33. A process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 100°C for about 24 hours.

20 34. A process for the preparation of Aripiprazole Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A at 120°C for about 3 hours.

35. A process for the preparation of Aripiprazole
25 Anhydride Crystals B wherein said process comprises heating Aripiprazole Hydrate A for about 18 hours at 100°C followed by additional heating for about 3 hours at 120°C.

125
289

- 117 -

36. The Aripiprazole Anhydride Crystals B according to any one of claims 24-29 made by a process comprising heating Aripiprazole Hydrate A for about 18 hours at 100°C followed by additional heating for about 5 3 hours at 120°C.

37. The Aripiprazole Anhydride Crystals B according to any one of claims 24-29 formulated with one or more pharmaceutically acceptable carriers.

38. The Aripiprazole Anhydride Crystals B 10 according to any one of claims 24-29 formulated with one or more pharmaceutically acceptable carriers to form a solid oral tablet.

39. The Aripiprazole Anhydride Crystals B according to any one of claims 24-29 formulated with 15 one or more pharmaceutically acceptable carriers to form an oral flashmelt tablet.

40. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B defined in claim 26 and one or more pharmaceutically 20 acceptable carriers, wherein said process comprises heating Aripiprazole Hydrate A defined in claim 7.

41. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B defined in claim 26 and one or more pharmaceutically 25 acceptable carriers, wherein said process comprises heating Aripiprazole Hydrate A defined in claim 7 at 90-125°C for about 3-50 hours.

42. A process for the pharmaceutical solid oral

136
27

- 118 -

preparation comprising Aripiprazole Anhydride Crystals B defined in claim 27 and one or more pharmaceutically acceptable carriers, wherein said process comprises heating Aripiprazole Hydrate A defined in claim 7.

5 43. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B defined in claim 27 and one or more pharmaceutically acceptable carriers, wherein said process comprises heating Aripiprazole Hydrate A defined in claim 7 at
10 90-125°C for about 3-50 hours.

44. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B defined in claim 28 and one or more pharmaceutically acceptable carriers, wherein said process comprises
15 heating Aripiprazole Hydrate A defined in claim 7.

45. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B defined in claim 28 and one or more pharmaceutically acceptable carriers, wherein said process comprises
20 heating Aripiprazole Hydrate A defined in claim 7 at 90-125°C for about 3-50 hours.

46. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals B defined in claim 29 and one or more pharmaceutically
25 acceptable carriers, wherein said process comprises heating Aripiprazole Hydrate A defined in claim 7.

47. A process for the pharmaceutical solid oral preparation comprising Aripiprazole Anhydride Crystals

187
271

- 119 -

B defined in claim 29 and one or more pharmaceutically acceptable carriers, wherein said process comprises heating Aripiprazole Hydrate A defined in claim 7 at 90-125°C for about 3-50 hours.

- 5 48. Aripiprazole Anhydride Crystals B wherein said Crystals will not substantially convert to a hydrous form of aripiprazole when properly stored under a relative humidity (RH) of 60 % and at a temperature of 25°C, even for an extended period being not less
10 than 1 year.
49. Aripiprazole Anhydride Crystals B wherein said Crystals will not substantially convert to a hydrous form of aripiprazole when properly stored under a relative humidity (RH) of 75 % and at a temperature
15 of 40°C, even for an extended period being not less than 0.5 year.
50. Aripiprazole Anhydride Crystals B having all physicochemical properties defined in claims 16, 18 to 20, 22 and 48.
- 20 51. Aripiprazole Anhydride Crystals B having all physicochemical properties defined in claims 17 to 20, 22 and 48.
52. Aripiprazole Anhydride Crystals B having all physicochemical properties defined in claims 16, 18 to
25 20, 22 and 49.
53. Aripiprazole Anhydride Crystals B having all physicochemical properties defined in claims 17 to 20, 22 and 49.

138
272

- 120 -

54. The Aripiprazole Anhydride Crystals B according to any one of claims 50-53 formulated with one or more pharmaceutically acceptable carriers.

55. The Aripiprazole Anhydride Crystals B
5 according to any one of claims 50-53 formulated with one or more pharmaceutically acceptable carriers to form a solid oral tablet.

56. The Aripiprazole Anhydride Crystals B according to any one of claims 50-53 formulated with
10 one or more pharmaceutically acceptable carriers to form an oral flashmelt tablet.

57. The use of aripiprazole anhydride crystals B for the treatment of central system disorder.

58. The use of aripiprazole anhydride crystals B
15 for the treatment of schizophrenia.

59. The use of aripiprazole anhydride crystals B for the treatment of bipolar disorder.

60. The use of aripiprazole anhydride crystals B for the treatment of intractable (drug-resistant,
20 chronic) schizophrenia with cognitive impairment or intractable (drug-resistant, chronic) schizophrenia without cognitive impairment.

61. The use of aripiprazole anhydride crystals B for the treatment of autism, Down's syndrome or
25 attention deficit hyperactivity disorder (ADHD).

62. The use of aripiprazole anhydride crystals B for the treatment of Alzheimer's disease, Parkinson's disease or other neurodegenerative diseases.

139
273

- 121 -

63. The use of aripiprazole anhydride crystals B for the treatment of panic, obsessive compulsive disorder (OCD), sleep disorders, sexual dysfunction, alcohol and drug dependency, vomiting, motion sickness, obesity, miparticlee headache or cognitive impairment.

64. The use of aripiprazole anhydride crystals B for the treatment of anxiety, depression or mania.

65. The use of aripiprazole anhydride crystals B to prepare a medicament to treat or prevent schizophrenia and the symptoms associated with schizophrenia.

66. A drug for treating schizophrenia or symptoms associated with schizophrenia, which comprises aripiprazole anhydride crystals B in an amount effective to treat schizophrenia or the symptoms thereof, in admixture with a pharmaceutically acceptable diluent.

67. The drug as claimed in claim 66, which is contained in a commercial package carrying instructions that the drug should be used for treating schizophrenia, or symptoms thereof.

68. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals B defined in claim 26, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

69. A process for the preparation of granules, characterized by wet granulating the Aripiprazole

140
274

- 122 -

Anhydride Crystals B defined in claim 27, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

70. A process for the preparation of granules,
5 characterized by wet granulating the Aripiprazole Anhydride Crystals B defined in claim 28, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

71. A process for the preparation of granules,
10 characterized by wet granulating the Aripiprazole Anhydride Crystals B defined in claim 29, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

72. A process for the pharmaceutical solid oral
15 preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 26 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

20 73. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 27 and one or more pharmaceutically acceptable carriers at 70 to
25 100°C.

74. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole

141
275

- 123 -

Anhydride Crystals B defined in claim 28 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

75. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 29 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

76. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 26 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

77. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 27 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

78. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 28 and one or more pharmaceutically

142
276

- 124 -

acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

79. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals B defined in claim 29 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

80. Aripiprazole Anhydride Crystals C having a powder X-ray diffraction spectrum shown in Figure 10.

81. Aripiprazole Anhydride Crystals C having a powder X-ray diffraction spectrum having characteristics peaks at $2\theta = 12.6^\circ, 13.7^\circ, 15.4^\circ, 18.1^\circ, 19.0^\circ, 20.6^\circ, 23.5^\circ$ and 26.4° .

82. Aripiprazole Anhydride crystals C having a particular infrared absorption bands at 2939, 2804, 1680, 1375 and 780 cm^{-1} on the IR (KBr) spectrum.

83. Aripiprazole Anhydride crystals C exhibiting an endothermic peak near about 150.2°C in thermogravimetric/differential thermal analysis (heating rate 5°C/min).

84. Aripiprazole Anhydride crystals C having a solid ^{13}C -NMR spectrum having characteristic peaks at

143
277

- 125 -

32.8 ppm, 60.8 ppm, 74.9 ppm, 104.9 ppm, 152.2 ppm and 175.2 ppm.

85. Aripiprazole Anhydride crystals C having all physicochemical properties defined in claims 80, 82 to 84.

86. Aripiprazole Anhydride crystals C having all physicochemical properties defined in claims 81 to 84.

87. Aripiprazole Anhydride crystals D having a powder x-ray diffraction spectrum shown in Figure 15.

88. Aripiprazole Anhydride crystals D having a powder x-ray diffraction spectrum having characteristic peaks at $2\theta = 8.7^\circ$, 11.6° , 16.3° , 17.7° , 18.6° , 20.3° , 23.4° and 25.0° .

89. Aripiprazole Anhydride crystals D having a particular infrared absorption bands at 2946, 1681, 1375, 1273, 1175 and 862 cm^{-1} on the IR (KBr) spectrum.

90. Aripiprazole Anhydride crystals D exhibiting an endothermic peak near about 136.8°C and 141.6°C in thermogravimetric/differential thermal analysis (heating rate 5°C/min).

91. Aripiprazole Anhydride crystals D having a solid ^{13}C -NMR spectrum having characteristic peaks at 32.1 ppm, 62.2 ppm, 66.6 ppm, 104.1 ppm, 152.4 ppm, 158.4 ppm, and 174.1 ppm.

92. Aripiprazole Anhydride crystals D having all physicochemical properties defined in claims 87, 89 to 91.

93. Aripiprazole Anhydride crystals D having all

144
278

- 126 -

physicochemical properties defined in claims 88 to 91.

94. Aripiprazole Anhydride crystals E having a powder x-ray diffraction spectrum shown in Figure 20.

95. Aripiprazole Anhydride crystals E having a
5 powder x-ray diffraction spectrum having characteristic peaks at $2\theta = 8.0^\circ$, 13.7° , 14.6° , 17.6° , 22.5° and 24.0° .

96. Aripiprazole Anhydride crystals E having a particular infrared absorption bands at 2943, 2817,
10 1686, 1377, 1202, 969 and 774 cm^{-1} on the IR (KBr) spectrum.

97. Aripiprazole Anhydride crystals E exhibiting an endothermic peak near about 146.5°C in thermogravimetric/differential thermal analysis
15 (heating rate $5^\circ\text{C}/\text{min}$).

98. Aripiprazole Anhydride crystals E having all physicochemical properties defined in claims 94, 96 to 97.

99. Aripiprazole Anhydride crystals E having all
20 physicochemical properties defined in claims 95 to 97.

100. Aripiprazole Anhydride crystals F having a powder x-ray diffraction spectrum shown in Figure 24.

101. Aripiprazole Anhydride crystals F having a powder x-ray diffraction spectrum having characteristic
25 peaks at $2\theta = 11.3^\circ$, 13.3° , 15.4° , 22.8° , 25.2° and 26.9° .

102. Aripiprazole Anhydride crystals F having a particular infrared absorption bands at 2940, 2815,

1545
279

- 127 -

1679, 1383, 1273, 1177, 1035, 963 and 790 cm^{-1} on the IR (KBr) spectrum.

103. Aripiprazole Anhydride crystals F exhibiting an endothermic peak near about 137.5°C and 149.8°C in thermogravimetric/differential thermal analysis (heating rate 5°C/min).

104. Aripiprazole Anhydride crystals F having all physicochemical properties defined in claims 100, 102 to 103.

105. Aripiprazole Anhydride crystals F having all physicochemical properties defined in claims 101 to 103.

106. Aripiprazole Anhydride crystals G having a powder x-ray diffraction spectrum shown in Figure 28.

107. Aripiprazole Anhydride crystals G having a powder x-ray diffraction spectrum having characteristic peaks at $2\theta = 10.1^\circ, 12.8^\circ, 15.2^\circ, 17.0^\circ, 17.5^\circ, 19.1^\circ, 20.1^\circ, 21.2^\circ, 22.4^\circ, 23.3^\circ, 24.5^\circ$ and 25.8° .

108. Aripiprazole Anhydride crystals G having a particular infrared absorption bands at 2942, 2813, 1670, 1625, 1377, 1195, 962 and 787 cm^{-1} on the IR (KBr) spectrum.

109. Aripiprazole Anhydride crystals G exhibiting an endothermic peak near about 141.0°C and an exothermic peak around 122.7°C in thermogravimetric/differential thermal analysis (heating rate 5°C/min).

110. Aripiprazole Anhydride crystals G having all physicochemical properties defined in claims 106, 108

146
780

- 128 -

to 109.

111. Aripiprazole Anhydride crystals G having all physicochemical properties defined in claims 107 to 109.

5 112. A process for preparing aripiprazole anhydride crystals C according to claim 85 or 86, characterized by heating aripiprazole anhydride crystals at a temperature higher than 140°C and lower than 150°C.

10 113. A process for preparing aripiprazole anhydride crystals D according to claim 92 or 93, characterized by recrystallizing it from toluene.

114. A process for preparing aripiprazole anhydride crystals E according to claim 98 or 99,
15 characterized by heating and dissolving aripiprazole anhydride crystals in acetonitrile, then cooling it.

115. A process for preparing aripiprazole anhydride crystals F according to claim 104 or 105, characterized by heating a suspension of aripiprazole
20 anhydride in acetone.

116. A process for preparing aripiprazole anhydride crystals G according to claim 110 or 111, characterized by leaving to stand aripiprazole anhydride glassy state in a sealed vessel at room
25 temperature for at least 2 weeks.

117. A pharmaceutical composition comprising at least one aripiprazole anhydride crystals selected from the group consisting of the aripiprazole anhydride

147
281

- 129 -

crystals C according to claim 85, the aripiprazole anhydride crystals D according to Claim 92, the aripiprazole anhydride crystals E according to Claim 98, the aripiprazole anhydride crystals F according to
5 Claim 104 and the aripiprazole anhydride crystals G according to Claim 110, together with pharmaceutically acceptable carriers.

118. A pharmaceutical composition comprising at least one aripiprazole anhydride crystals selected from
10 the aripiprazole anhydride crystals C according to claim 86, the aripiprazole anhydride crystals D according to claim 93, the aripiprazole anhydride crystals E according to claim 99, the aripiprazole anhydride crystals F according to Claim 105, and the
15 aripiprazole anhydride crystals G according to Claim 111, together with pharmaceutically acceptable carriers.

119. A process for the preparation of granules, characterized by wet granulating the Aripiprazole
20 Anhydride Crystals C defined in claim 85, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

120. A process for the preparation of granules, characterized by wet granulating the Aripiprazole
25 Anhydride Crystals C defined in claim 86, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

121. A process for the preparation of granules,

148
282

- 130 -

characterized by wet granulating the Aripiprazole Anhydride Crystals D defined in claim 92, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

- 5 122. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals D defined in claim 93, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.
- 10 123. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals E defined in claim 98, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.
- 15 124. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals E defined in claim 99, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.
- 20 125. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals F defined in claim 104, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.
- 25 126. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals F defined in claim 105, drying the obtained granules at 70 to 100°C and sizing it, then

149
283

- 131 -

drying the sized granules at 70 to 100°C again.

127. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals G defined in claim 110, drying the
5 obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

128. A process for the preparation of granules, characterized by wet granulating the Aripiprazole Anhydride Crystals G defined in claim 111, drying the
10 obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

129. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole
15 Anhydride Crystals C defined in claim 85 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

130. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical
20 solid oral preparation comprising the Aripiprazole Anhydride Crystals C defined in claim 86 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

131. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical
25 solid oral preparation comprising the Aripiprazole Anhydride Crystals D defined in claim 92 and one or more pharmaceutically acceptable carriers at 70 to

156
284

- 132 -

100°C.

132. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole

5 Anhydride Crystals D defined in claim 93 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

133. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical

10 solid oral preparation comprising the Aripiprazole Anhydride Crystals E defined in claim 98 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

134. A process for the pharmaceutical solid oral

15 preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals E defined in claim 99 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

20 135. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals F defined in claim 104 and one or more pharmaceutically acceptable carriers at 70 to

25 100°C.

136. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole

101
235

- 133 -

Anhydride Crystals F defined in claim 105 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

137. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals G defined in claim 110 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

138. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals G defined in claim 111 and one or more pharmaceutically acceptable carriers at 70 to 100°C.

139. A process for the preparation of granules, characterized by wet granulating conventional Aripiprazole Anhydride Crystals, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

140. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical solid oral preparation comprising conventional Aripiprazole Anhydride Crystals and one or more pharmaceutically acceptable carriers at 70 to 100°C.

141. A pharmaceutical solid oral preparation having at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30

152
285

- 134 -

minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

142. The pharmaceutical solid oral preparation prepared by the process of claim 139, wherein said
5 pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

10 143. The pharmaceutical solid oral preparation prepared by the process of claim 140, wherein said pharmaceutical solid oral preparation having at least one dissolution rate selected from the group consisting
60% or more at pH 4.5 after 30 minutes, 70% or more at
15 pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

144. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals C defined in claim 85 and one or more pharmaceutically
20 acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

25 145. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals C defined in claim 86 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid

153
287

- 135 -

oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

5 146. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals D defined in claim 92 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate
10 selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

147. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals D
15 defined in claim 93 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60
20 minutes, and 55% or more at pH 5.0 after 60 minutes.

148. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals E defined in claim 98 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid
25 oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

184
283

- 136 -

149. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals E defined in claim 99 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid
5 oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

150. The pharmaceutical solid oral preparation
10 comprising the Aripiprazole Anhydride Crystals F defined in claim 104 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH
15 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

151. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals F defined in claim 105 and one or more pharmaceutically
20 acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

25 152. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals G defined in claim 110 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid

165
289

- 137 -

oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

5 153. The pharmaceutical solid oral preparation comprising the Aripiprazole Anhydride Crystals G defined in claim 111 and one or more pharmaceutically acceptable carriers, wherein said pharmaceutical solid oral preparation has at least one dissolution rate
10 selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

154. A process for the preparation of granules, characterized by wet granulating conventional

15 Aripiprazole Hydrate Crystals, drying the obtained granules at 70 to 100°C and sizing it, then drying the sized granules at 70 to 100°C again.

155. A process for the pharmaceutical solid oral preparation, characterized by drying a pharmaceutical
20 solid oral preparation comprising conventional Aripiprazole Hydrate Crystals and one or more pharmaceutically acceptable carriers at 70 to 100°C.

156. The pharmaceutical solid oral preparation prepared by the process of claim 154, wherein said
25 pharmaceutical solid oral preparation has at least one dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after

156
290

- 138 -

60 minutes.

157. The pharmaceutical solid oral preparation prepared by the process of claim 155, wherein said pharmaceutical solid oral preparation has at least one
5 dissolution rate selected from the group consisting 60% or more at pH 4.5 after 30 minutes, 70% or more at pH 4.5 after 60 minutes, and 55% or more at pH 5.0 after 60 minutes.

157

FIG.1





FIG.2



159

FIG.3




FIG.4





FIG.5



6/31

182

FIG.6






FIG.7



8/31

169

FIG.8




9/31

PCT/JP02/09658

158


FIG.9



10/31


156

FIG.10



11/31

FIG.11




167

168

12/31

FIG.12



32

W/O 13/026639

PCT/JP02/09658

13/31

169

FIG.13





FIG.14



58


WO 03/026659

PCT/JP02/09651

15/31

131

FIG 15



16/31

172

FIG.16






FIG.17

17/31



18/31


184



19/31


125

FIG.19



126


FIG.20



21/31


177

FIG.21




126

FIG.22



23/31


FIG.23



24/31

/8


FIG.24



25/31

18x

FIG.25



26/31

182

FIG.26






FIG.27



28/31


184

FIG 28



29/31


FIG.29



30/31

186

FIG.30



187

FIG.31

31/31

