

As. 123.233

Señor

**DIRECTOR DIVISIÓN DE NUEVAS CREACIONES
SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO**

E.

S.

D.

REF: Expediente No. **13 015.329-** solicitud de patente de invención a favor de **ADETHERAPEUTICS INC.** Título: **COMPOSICIÓN EN FORMA DE ORGANOGEL QUE COMPRENDE UNA FUENTE DE GLUTAMINA.**

RESPUESTA A EXAMEN DE FONDO

JUAN PABLO CADENA SARMIENTO, portador de la tarjeta profesional No. 57.492 del C.S.J., obrando como apoderado de la sociedad solicitante me permito dar respuesta al Oficio No. 10357 relacionado con la patentabilidad de la presente invención y notificado por fijación en lista el 8 de Septiembre de 2015.

En primera instancia, nos permitimos manifestar nuestro total desacuerdo frente a todas y cada una de las objeciones y comentarios manifestados por el señor Examinador en contra de la novedad y nivel inventivo de la presente invención, requisitos que hemos defendido de manera pertinente a través de argumentos y evidencia documental a lo largo del presente trámite.

Pese a esto, ponemos a su consideración una nueva versión del pliego reivindicatorio que consta de tres (3) reivindicaciones, las cuales han sido modificadas pertinentemente con la intención de acentuar aún más las diferencias existentes entre el objeto y alcance de la presente invención y las divulgaciones del arte previo.

Por esta razón, y para contextualizar el análisis que prosigue, presentamos a continuación la nueva reivindicación 1:

“Una composición farmacéutica para administración tópica que comprende una fuente de glutamina y uno o más vehículos farmacéuticamente aceptables, en donde el vehículo farmacéuticamente aceptable comprende un vehículo tópico que es un organogel capaz de proporcionar administración sitio-específica de compuestos bioactivos a la piel.”

Adaptado del nuevo pliego reivindicatorio, reivindicación 1

Así las cosas, notará el señor Examinador que el vehículo farmacéuticamente aceptable que hace parte de la composición reivindicada se define ahora como un ***“un vehículo tópico que es un organogel capaz de proporcionar administración sitio-específica de compuestos bioactivos a la piel”***, expresión que será analizada en detalle más adelante.

Por lo pronto, concluimos que las modificaciones llevadas a cabo no amplían el alcance de la invención previamente reclamada y se mantienen enteramente sustentadas por las enseñanzas del capítulo descriptivo, razón por la que solicitamos sea reconocida la total validez de este nuevo pliego reivindicatorio.

Antes de abordar el análisis de dicha nueva reivindicación 1 y la defensa de su patentabilidad, es preciso aclarar que en lugar de referirse al posible uso de la presente invención, la expresión *“para administración tópica”* representa una **característica técnica esencial** de la misma pues permite (i) definir su objeto y alcance; (ii) conocer la solución al problema técnico que esta proporciona, y (iii) diferenciarla del arte previo.

Más específicamente, afirmamos que dicha expresión permite comprender el **tipo de composición farmacéutica** al cual hace referencia la presente solicitud sin aludir en ningún momento a los posibles usos o aplicaciones de la misma, resultando entonces en una caracterización completamente adecuada y pertinente.

Inclusive, destacamos que gracias a la presencia de dicha expresión es posible comprender de manera precisa que la materia reivindicada corresponde a una **composición farmacéutica tópica** y no a cualquier otro tipo de composición farmacéutica, hecho que además restringir el campo técnico al cual pertenece la invención, permite diferenciarla del arte previo.

Con el fin de demostrar la pertinencia de la materia objetada, incluimos a continuación un fragmento del Instructivo de Examen de Solicitudes de Patentes de Invención y Modelo de Utilidad, en donde se define qué tipo de información debe contener una reivindicación:

“Las reivindicaciones deben contener todas las características técnicas esenciales de la invención las cuales definen la invención y la hacen (o deberían hacerla) diferente del estado de la técnica y, por tanto, constituyen la solución al problema técnico que intenta resolver la invención.”

Adaptado del Instructivo de Patentabilidad
Numeral 2.6.1 – Contenido de las reivindicaciones

Debido a que la materia incluida en la reivindicación 1, o más específicamente la expresión “*para administración tópica*” cumple satisfactoriamente con los lineamientos del Instructivo de Patentabilidad citado, solicitamos sea reconocido que la misma no sugiere ningún uso o posible uso de la invención sino que caracteriza de manera apropiada el objeto y alcance de la presente invención.

De hecho, es de notar que dicha expresión adquiere un rol muy importante al momento de comparar el objeto de la presente solicitud frente al estado de la técnica ya que, como bien reconoce el señor Examinador, los documentos D3 y D4 dan a conocer composiciones farmacéuticas de administración intraperitoneal provistas de componentes similares a los aquí reivindicados.

En este sentido, resaltamos que a pesar de que exista similitud a nivel de sus componentes técnicos esenciales, la expresión “*para administración tópica*” permite establecer una clara diferencia entre el campo técnico relativo a la presente solicitud y su alcance, el cual resulta en todo caso distinto al concerniente a los citados documentos D3 y D4.

Teniendo en cuenta entonces que el **tipo de composición farmacéutica** representa un atributo distintivo de la presente invención con respecto al arte previo, solicitamos sea reconocido que la expresión “*para administración tópica*” no hace referencia a posibles usos o aplicaciones de la composición aquí descrita, sino representa una característica técnica esencial de la misma.

En conexión con lo anterior, nos permitimos señalar que la caracterización expuesta en la nueva reivindicación 1 permite acentuar aún más el hecho de que la presente invención corresponde a una **composición farmacéutica tópica** dada la inclusión de la expresión “*en donde el vehículo farmacéuticamente aceptable comprende un vehículo tópico que es un organogel capaz de proporcionar administración sitio-específica de compuestos bioactivos a la piel*” en la parte caracterizante de la misma.

De esta manera, solicitamos al señor Examinador reconocer que la presente invención representa una **composición farmacéutica tópica** conformada por los siguientes elementos:

- Una fuente de glutamina;
- Uno o más vehículos farmacéuticamente aceptables; el cual comprende un **vehículo tópico de organogel** que ejerce de funcionalidades específicas,

los cuales definen una composición completamente novedosa, inventiva y sin precedentes en el estado de la técnica.

Novedad y Nivel inventivo

Acudiendo a las enseñanzas de los documentos D3 (*Reducing post-operative adhesion formation with intraperitoneal glutamine*, WO2008/098364) y D4 (*Reducing post-operative adhesion formation with intraperitoneal glutamine*, WO2007/016791), afirma el señor Examinador que la invención descrita en la reivindicación 1 carece de novedad, mientras que la divulgada en la reivindicación 3 representa una derivación obvia del estado de la técnica.

Partiendo de nuestro total desacuerdo frente a tales objeciones, procedemos a demostrar que tanto la nueva reivindicación 1, como sus respectivas dependientes, cumplen a cabalidad con los citados requisitos de novedad y nivel inventivo.

Como primera medida, nos permitimos mencionar que el examen conducido por el señor Examinador evidencia una interpretación errónea de la materia divulgada por dichos documentos D3 y D4, referencias que ignoran completamente la utilización e inclusión de organogeles como parte de las composiciones allí descritas.

Así pues, tras llevar a cabo una revisión exhaustiva de las citadas anterioridades, encontramos que ninguna de estas anticipa de manera exacta la presencia de un **organogel** como parte del vehículo de las composiciones farmacéuticas allí descritas.

Muy por el contrario, dichos documentos D3 y D4 se relacionan con el desarrollo de composiciones que incorporan **hidrogeles** como parte de sus respectivos vehículos, tal y como revelan los siguientes fragmentos:

"Un agente mejorador de viscosidad se usa en esta invención tal que pueda formarse un gel al hidratar. Tales composiciones de gel pueden comprender una fuente de glutamina soluble en agua y un agente formador de gel adecuado en forma seca."

Adaptado del documento D3

"Una formulación particularmente preferida para uso en esta invención es un gel o una malla que contiene una fase acuosa en la que se disuelve una fuente de glutamina."

Adaptado del documento D4

"Agentes espesantes farmacéuticamente aceptables adecuados son conocidos y pueden ser empleados. Preferiblemente, tal agente formará un hidrogel cuando se hidrate o formará un hidrogel cuando se someta a un agente de reticulación adecuado y se hidrate. Tales componentes que forman gel son seleccionados por su biocompatibilidad y pueden ser reabsorbibles. Ejemplos de espesantes adecuados y agentes formadores de gel que han sido empleados en formulaciones farmacéuticas incluyen polímeros que tienen un componente hidrófilo, tal como colágeno; polímeros de polioxialquileno tales como óxidos de polietileno, alcoholes de polivinilo, polivinilpirrolidonas, y metacrilato de hidroxietilo;

hialuronato; y varias proteínas como la albúmina, etc. geles hemostáticos, incluyendo aquellos que contienen fibrinógeno o fibrina también se pueden usar.”

Adaptado de los documentos D3 y D4

Con base en estos fragmentos, concluimos que las divulgaciones del arte previo se limitan únicamente a describir la utilización e incorporación de **hidrogeles**; materia que no resulta de ninguna manera comparable con el objeto y alcance de la presente invención, la cual se caracteriza por comprender un **vehículo tópico de organogel**.

Dicho esto, concluimos que ninguno de los documentos D3 y D4 anticipa la caracterización precisa que da origen a la presente invención pues los mismos desconocen por completo la posibilidad de incluir un **vehículo tópico de organogel** como componente esencial de la misma.

Por esta razón, solicitamos de manera respetuosa sea reconocido que no hay lugar a atribuirle falta de novedad a la materia descrita en el presente pliego reivindicatorio y en consecuencia, sea retirada la respectiva objeción.

Adicional a lo anterior, es pertinente mencionar que en nuestro concepto el señor Examinador recae en un claro error conceptual al asumir de manera deliberada que los hidrogeles descritos en los documentos D3 y D4 pueden ser semejantes o equivalentes al **vehículo tópico de organogel** que caracteriza las composiciones de la presente invención.

Como muestra de ello, nos permitimos adjuntar al presente memorial el artículo de revisión elaborado por Tarun *et al.* titulado “*Organogels: Advanced and novel drug delivery system*” en donde se exponen abiertamente las diferencias existentes entre un “*organogel*” y un “*hidrogel*”.

Concretamente, Tarun *et al.* establecen la siguiente caracterización con respecto a los sistemas de geles:

“Un gel puede definirse como una formulación semi-sólida provista de una fase externa de solvente apolar (organogeles) o polar (hidrogeles), inmovilizada al interior de los espacios disponibles de una estructura en red tridimensional”

Adaptado de Anexo 1, Introducción

Asimismo, y de conformidad con tal clasificación inicial, Tarun *et al.* definen así el concepto de **organogel**:

“Un organogel es un material sólido termorreversible (termoplástico) no-cristalino, no-vidrioso y un sistema visco-elástico que puede ser reconocido como una preparación semi-sólida provista de una fase apolar externa.”

Adaptado de Anexo 1, Resumen

“En contraste con los hidrogeles, en donde el agente gelificante normalmente es un polímero, la mayoría de los organo-gelificantes son moléculas relativamente pequeñas que son reconocidas como organo-gelificantes de bajo peso molecular.”

Adaptado de Anexo 1, Introducción

De esta manera, es evidente que existe una clara diferencia conceptual entre un **“organogel”** y un **“hidrogel”**, razón por la que insistimos en que las objeciones del señor Examinador carecen de total fundamento en la medida que los documentos D3 y D4 abordan únicamente el desarrollo de **composiciones de hidrogel**, las cuales difieren considerablemente del objeto de la presente invención.

Dentro de las diferencias existentes entre dichos dos tipos de geles, vale la pena destacar que los **hidrogeles** representan sistemas polares que forman semisólidos al ser dejados en reposo, mientras que los **organogeles** representan sistemas apolares que contrario a los hidrogeles, no son capaces de formar tales semisólidos al ser dejados en reposo.

Teniendo en cuenta que a pesar de que ambos son reconocidos como geles, los hidrogeles y organogeles gozan de **características químicas y físicas propias y diferenciales** como bien expone el Anexo 1 y como resultará evidente para cualquier persona normalmente versada en la materia, concluimos que no hay ninguna razón

por la cual la novedad de la presente invención pueda ser perjudicada por las enseñanzas D3 y D4.

Consecuentemente, solicitamos respetuosamente sea retirada la objeción que cuestiona la novedad de la presente invención en vista de los documentos D3 y D4, los cuales insistimos no representan soporte documental suficiente que anticipe de manera exacta la caracterización expuesta a lo largo del actual pliego reivindicatorio.

Abordando ahora la defensa del nivel inventivo de la presente invención, nos permitimos señalar que los documentos D3 y D4 carecen de instrucciones que motiven al versado en la materia a alcanzar de manera obvia la caracterización actualmente reivindicada, razón por la que consideramos que la objeción del señor Examinador no tiene ningún sustento.

Como punto de partida, es de notar que el **campo técnico** al cual pertenecen las referencias citadas es sustancialmente distinto al que concierne a la presente invención por cuanto se refiere al desarrollo de composiciones de administración intraperitoneal.

En contraste con ello, el objeto de la presente invención corresponde a una **composición farmacéutica de administración tópica**, la cual constituye ciertamente una forma farmacéutica diferente a la divulgada por los documentos D3 y D4, la cual cumple con requerimientos específicos y se enfrenta a problemáticas distintas a las abordadas por dicho arte previo.

En este sentido, llamamos la atención sobre el hecho de que las consideraciones biofarmacéuticas, farmacotécnicas y farmacocinéticas relativas al desarrollo de composiciones farmacéuticas varían de acuerdo al blanco terapéutico o, en este caso, al sitio de aplicación o utilización de la misma, siendo evidente la diferencia que existe entre una **composición intraperitoneal** y una **composición tópica** como la presente invención.

Debido a esto, afirmamos que el versado en la materia interesado en el desarrollo de una **composición tópica** como la aquí reivindicada no acudiría a las enseñanzas de los documentos D3 y D4 con la expectativa de hallar en ellos información de utilidad puesto que dichas anterioridades se relacionan con un objeto totalmente diferente y alejado del campo técnico de su interés.

Ahora bien, considerando que el versado en la materia consultara por alguna razón dichas referencias, es evidente que este no encontraría en ellas información relevante que lo incite a seleccionar cada una de las características técnicas esenciales de la presente invención y desarrollar una **composición farmacéutica de administración tópica** como la aquí descrita.

Más concretamente, insistimos en que dichos documentos D3 y D4 ignoran completamente el diseño e inclusión de un **vehículo tópico de organogel**, razón por la que solicitamos sea reconocido que dicha característica técnica representa un aspecto distintivo e inventivo de la presente invención que de ninguna manera habría podido ser previsto a partir de las referencias en mención.

Inclusive, encontramos que las enseñanzas de los documentos D3 y D4 sugieren fuertemente la utilización de **hidrogeles** como parte del vehículo de tales composiciones, razón por la que consideramos que el versado en la materia optaría por conservar dicho vehículo de hidrogel en lugar de pensar en emplear un organogel, alejándose en definitiva del alcance de la presente invención.

Dicho esto, concluimos que dichas referencias no hacen otra cosa que **alejar** al versado en la materia del objeto de la presente invención al divulgar composiciones que comprenden un **vehículo de hidrogel**, ignorando por completo la caracterización y concepto técnico que da lugar a la presente invención, la cual reiteramos se encuentra definida de manera clara como una composición que comprende un **vehículo tópico de organogel**.

Teniendo en cuenta que los documentos D3 y D4 citados por el señor Examinador no contienen ninguna información relacionada con (i) el desarrollo de una **composición farmacéutica de administración tópica**, y/o (ii) la inclusión de un **vehículo tópico de organogel** como parte de la misma, concluimos que los mismos **ignoran** aspectos distintivos de la presente invención y por tanto no podrían anticipar su desarrollo.

Partiendo de dichas carencias, concluimos también que dichos documentos D3 y D4 no representan el debido soporte documental a la objeción manifestada por el señor Examinador en contra del nivel inventivo de la presente solicitud, razón por la que solicitamos respetuosamente sea retirada.

Con base en lo anterior, concluimos que el desarrollo de la invención expuesta a lo largo de este nuevo pliego reivindicatorio no se ve anticipado y mucho menos sugerido o guiado por las enseñanzas de los citados documentos D3 y D4, los cuales resultan **alejados** de la misma y por tanto, fallan en desvirtuar su patentabilidad.

En complemento de esta argumentación, consideramos oportuno manifestar nuestro desacuerdo frente al examen de patentabilidad conducido por el señor Examinador, el cual acude a dos referencias del arte previo que guardan muy poca relación con el objeto de la presente solicitud y que por tanto, resultan completamente irrelevantes a ojos del versado en la materia.

En nuestro concepto, este hecho demuestra que dicho examen de patentabilidad fue llevado a cabo desde una visión altamente subjetiva del arte previo en donde se ignoran las evidentes deficiencias de las anterioridades citadas y se presume sin ningún fundamento una supuesta relación entre las mismas y el objeto de la presente invención.

Sumado a lo anterior, es preciso señalar que las dos referencias citadas por el señor Examinador como arte previo relevante para el objeto de la presente solicitud corresponden a **solicitudes análogas** pertenecientes al mismo titular y enfocadas en divulgar la misma materia, tal y como indica el siguiente fragmento:

“La solicitud equivalente del solicitante, publicada ahora como WO 2007016791 describe formulaciones que actúan como fuente de glutamina y se espesan con el fin de tener mayor viscosidad que formulaciones que son adecuadas para inyección intravenosa.”

Adaptado del documento D3

De este modo, es evidente que dichos documentos D3 y D4 no son de ninguna manera complementarios ni constituyen en conjunto una aproximación al objeto de la presente solicitud, sino que estos revelan prácticamente la misma materia y por ende resultan redundantes.


Más aún, consideramos que el señor Examinador se limitó a identificar de manera aislada e inconexa los diferentes elementos que definen la composición reivindicada en dichos dos documentos del estado de la técnica, sin que estos brindaran instrucciones concretas respecto a **cómo** conseguiría el versado en la materia desarrollar una invención como la reivindicada sin la necesidad de aplicar su capacidad inventiva.

En este orden de ideas, afirmamos que la presente invención fue juzgada bajo un análisis en retrospectiva que estudia de manera superficial las divulgaciones del estado de la técnica y presume sin ningún fundamento concreto que la misma podría haber sido derivada de manera obvia, cuando en realidad ninguno de los documentos D3 o D4 anticipa o sugiere cada uno de los elementos característicos de la composición reclamada en este nuevo pliego reivindicatorio.

Considerando entonces el conjunto de modificaciones y argumentos presentados a lo largo de este memorial, concluimos que la novedad y nivel inventivo de la presente invención se mantienen vigentes a pesar de las objeciones y comentarios expresados en el Oficio técnico, las cuales hemos solucionado y aclarado de manera pertinente.

En vista de lo anteriormente expuesto, respetuosamente solicitamos el retiro de las objeciones de la presente solicitud ya que en opinión del solicitante cumple con todos los requisitos establecidos en la Decisión 486.

Atentamente,



JUAN PABLO CADENA SARMIENTO

T.P. No. 57.492 del C.S.J.

C.C. 80.410.337 de Usaquén

Calle 70 A No. 4 - 41

REIVINDICACIONES

1. Una composición farmacéutica para administración tópica que comprende una fuente de glutamina y uno o más vehículos farmacéuticamente aceptables, en donde el vehículo farmacéuticamente aceptable comprende un vehículo tópico que es un organogel capaz de proporcionar administración sitio-específica de compuestos bioactivos a la piel.

2. La composición de acuerdo con la reivindicación 1, en donde la fuente de glutamina es un dipéptido que comprende L-alanil-L-glutamina.

3. La composición de acuerdo con la reivindicación 2, en donde el dipéptido está desde 5% hasta 25% en peso de la composición.

ORGANOGELES: ADVANCED AND NOVEL DRUG DELIVERY SYSTEM

Garg Tarun*, Bilandi Ajay, Kapoor Bhawana, Kumar Sunil, Joshi Ravi
Seth G. L. Bihani S. D. College of Technical Education, Sriganaganagar, India

Article Received on: 12/10/11 Revised on: 20/11/11 Approved for publication: 14/12/11

*Email: tarun_garg9@gmail.com

ABSTRACT

Organogel, is a non crystalline, non-glassy thermoreversible (thermoplastic) solid material and viscoelastic system, can be regarded as a semi-solid preparation which has an immobilized external apolar phase. The apolar phase gets immobilized within spaces of the three-dimensional networked structure formed due to the physical interactions amongst the self assembled structures of compounds regarded as gelators. Often, these systems are based on self-assembly of the structural molecules. In general, organogels are thermodynamically stable in nature and have been explored as matrices for the delivery of bioactive agents. Organogels have potential for use in a number of applications, such as in pharmaceuticals, cosmetics, art conservation, and food. An example of formation of an undesired thermoreversible network is the occurrence of wax crystallization in petroleum. In the current manuscript, attempts have been made to understand the properties of organogels, various types of organogelators and some applications of the organogels in controlled delivery.

KEYWORDS: Organogels, gelators, cosmetics, controlled delivery

INTRODUCTION

A gel may be defined as a semi-solid formulation having an external solvent phase, apolar (organogels) or polar (hydrogel), immobilized within the spaces available of a three dimensional networked structure¹. The organogel do not form semisolids on standing. Because an organogel may consists of macromolecules existing as twisted matted strands. The units are of bound together by strong types of vanderwaal forces so as to form crystalline amorphous regions throughout the entire system². Gels can also be classified according to the bonds present in the gelator network: physical gels are held by weaker physical forces of attraction such as vanderwaals interactions and hydrogen bonds, whereas chemical gels are held by covalent bonds. Hydrogels (composed of water held by a three-dimensional polymeric network) have been extensively studied as vehicles for a wide range of drugs. The organogels may be regarded as bi-continuous systems consisting of gelators and apolar solvent, which may or may not contain water-molecules entrapped within the self-assembled structures of the gelator. The organogels have lower hydrations, the drug dissolving polymer and is transported between the chains. Cross linking increases hydrophobicity of gels & diminishes the diffusion rate of drug. The gelators, when used in concentration < 15 % (approx.), may undergo physical or chemical interactions so as to form self-assembled fibrous structures which get entangled with each other resulting in the formation of a three-dimensional networked structure. The three-dimensional networked structure, hence formed, prevents the flow of external apolar phase. Some common examples of gelators include sterol, sorbitan monostearate, lecithin and cholesteryl anthraquinone derivatives. They have been fabricated in a variety of different shapes (e.g., rods, disks, films and microparticles) depending on the intended applications and sites of administration. In addition, some thermoresponsive gels can be administered parenterally as a liquid, which forms a gel in situ at body temperature. The thermo-reversible property of the organogels has generated much interest for the potential use of the organogels as drug delivery system. The thermodynamic stable nature of the organogels has been attributed to the spontaneous formation of fibrous structure by virtue of which the organogels reside in a low energy state. The occurrence of the gel-to-sol transition above room-temperature indicates that external energy has to be supplied to the organogels so as to disrupt the three-dimensional structure and subsequent transformation of the gelled state to the sol state. Apart from the temperature sensitivity, organogels are also sensitive to the presence of moisture which has also been explored to develop controlled delivery systems. Examples

of gellable organic solvents include aliphatic and aromatic hydrocarbons, alcohols, silicone oil, dimethyl sulfoxide and vegetable oils. In contrast to hydrogels, in which the gelator is normally a polymer, most of the organogelators are relatively small molecules and they have been called low molecular weight organogelators. Various organogel-based formulations have been designed to administer of the bioactive agents by different routes administration¹.

Advantages Of Organogels

- Organogel can't form semisolids preparation on standing.
- Organogels are moisture insensitive.
- They enhance the skin penetration and transport of the molecules.
- They are organic in character also resist microbial contamination.
- Organogel can diminish the diffusion rate of drug because the drug is dissolved in polymer & transported between chains.
- Structural integrity of organogels is maintained for longer time periods.
- Process very simple and easy to handle.
- Use of biocompatible, biodegradable and non-immunogenic materials makes them safe for long term applications.
- Organogels provide opportunities for incorporation of wide range of substances with different physicochemical characters³.

Limitations Of Organogels

- Less stable to temperature.
- When a gel stands for some time, it often shrinks naturally, & some of its liquid is pressed out, known as syneresis.
- If impurity present then no gelling will occur.
- Expensive in production.
- Raw materials like lecithin are not available on large scale.
- It should be stored in a proper condition.
- When the gel is taken up of liquid with an increasing volume known as swelling⁴.

Structure Of Organogels & Mechanism Of Organogelling

The incorporation of a polar solvent, the organogelling or the gelation of the lecithin solutions in organic solvents is induced. The aggregate transformation (i.e. sphere-to-cylinder transformations) are determined by a change of a curvature for the amphiphile monolayer and this approaches developed by Israelachvili et al.⁵ In particular, the effects of polar solvent introduced into spherical lecithin micelles may be associated in which the solvent with an increase in the cross-sectional area of the lecithin polar region is

arranged. The shape of the hydrated molecules is close to a cylinder. This shape leads to packing constraints in the spherical micelles that are diminished through the transition into the cylindrical ones with a smaller curvature.

Gelator Self-Assembly

A variety of gelator aggregates such as platelets, tubules, fibres, rods, worm-like chains, ribbons and fan-like structures have been reported. Aggregate thickness ranges from a fraction of a nanometre to microns. The manner in which gelator molecules self-assemble and the nature of the gelator aggregate depends to a large extent on the gelator, whose component groups dictate the forces of interactions involved in gelator self-assembly. For example, molecules of the non-ionic surfactant sorbitan monostearate are thought to assemble into bilayers, which are then organised into tubules. 12-Hydroxyoctadecanoic acid is thought to form fibrillar aggregates via extensive axial hydrogen bonding and dipolar interactions⁶. Aggregation forces between gelator molecules thus include hydrogen bonding, dipole-dipole interactions, π -stacking, electron transfer, London dispersion forces, solvophobic effects, ionic interactions and so on, depending on the chemical structures of the gelators. The manner in which the gelator molecules are packed in aggregates must not be assumed to be the same as in the neat gelator solid, and the different packing and forces of interactions in gelator aggregates and in the neat solid are often reflected in the different melting points of the two. The liquid phase of the gel plays a fundamental role in gelation and affects both macroscopic (e.g., opacity) and microscopic (e.g., aggregate size, shape, cross-sectional nature, helicity and gel network) properties of the gels. The fluid phase must provide the correct solubility/insolubility balance so that the gelator dissolves or is dispersed at high temperatures but comes out of solution (as aggregates) following cooling. Molecular shape of the solvent molecule can also have a profound effect on gelation; for example, steroidal nitroxide forms stable gels in trans-decalin and in cyclohexane but not in cis-decalin, methyl-cyclohexane and n-alkane. The higher melting point was thought to be due to the additional attractive forces provided by hydrogen bonding between 1-octanol and CAB in octanol gels. The dimensions of the container in which gelation proceeds can also have an impact on the gel network. Furman and Weiss reported that if the size of the container

was smaller than the mesh size of a particular gel network, gelation was either inhibited or a different type of gel network (mesh) was formed⁷.

Types Of Organogelators

The organogelators may be categorized into two groups based on their capability to form hydrogen bonding. The examples of organogelators the hydrogen bond forming organogelators include amino acids, amide, urea moieties and carbohydrates whereas which do not form hydrogen hydrogen bonding include anthracene, anthraquinone and steroid based molecules. The simplest organogelators are n-alkanes (C = 24, 28, 32, 36), which gel other relatively short chain n-alkanes such as hexadecane and other organic liquids. Other examples of organogelators include substituted fatty acids (e.g., 12-hydroxyoctadecanoic acid); 1,3:2,4-di-O-benzylidene-Dsorbitol (D-DBS), sorbitan monostearate, a non-ionic surfactant, steroids and their derivatives, anthryl derivatives (e.g., 2,3-bis-n-decyloxyanthracene, macrocyclic gelators (e.g., calixarenes, A1S compounds (an aromatic moiety attached to a steroidal group by a linker segment), cyclo(dipeptide)s, bisurea compounds, bisamides, bolaform amides derived from amino acids, n-alkyl perfluoroalkanamides, carbohydrate derivatives, perfluoroalkanes, which gel liquid carbon dioxide, a mixture of highly reactive methyl 2,6-diisocyanatohexanoate and alkylamines in an organic solvent, which react when mixed to form a product that gels the organic solvent, primary alkyl amines, which gel organic solvents following the uptake of CO₂, NO₂, SO₂ or CS₂, light-responsive gelators, which produce gels whose sol-to-gel transition may be switched by irradiation with UV and visible light, oxadiazole-based benzene 1,3,5- tricarboxamide, a nonfluorescent gelator, which produces highly fluorescent organogel, cobalt (II) triazole complexes, which, unlike most organogels, form gels at high temperatures and solutions at low temperatures, and fatty acid derivative of L-alanine, which selectively gels the organic solvent but not the aqueous phase when added to an oil/water mixture. Certain compounds can only gel organic solvents in the presence of other compounds, for example, aminopyrimidine and dialkylbarbituric acid gel cyclohexane when present at 1:1 molar mixtures⁸. Different types of organogelators with their property and uses are described in Table 1

Table 1. Types of organogelators with their advantages and applications^{2,10}

S.No.	Types and property of organogelators	Advantages and applications of organogelators
1	4-tertbutyl-1-aryl cyclohexanols derivative organogelators These gelators are solid at room temperature having low solubility in apolar solvents viz cyclohexane, benzene and carbon tetra chloride	4-tertbutyl-1-aryl cyclohexanols, categorized under arylcyclohexanol derivatives, helps in designing thermo-reversible organogels.
2	Polymer organogelators - Polymer organogelators have been found to induce organogelation even at very low concentrations and their gelling capability of these gelators may be tailored by modifying the chemical structure of the polymer backbone - Some common examples of polymeric organogelators include L-lysine derivatives apart from the conventional polymers like poly(ethylene glycol), polycarbonate, polyesters, and poly(alkaline)	The gels developed by polymeric organogelators generally have lower gel-sol transition temperature and comparatively higher gel strength when compared with organogels developed with low-molecular weight organogelators.
3	Gemini organogelators The Gemini organogelators which had two L-lysine derivatives connected with alkaline spacer chains, of varying chain lengths, by amide bonds.	- bis(Nε-lauroyl-L-lysine ethyl ester) oxalyl amide organogelator was able to immobilize a variety of apolar solvents. - Various other oxalyl amide derivatives containing various alkyl ester groups (e.g. hexyl, decyl, dodecyl, 2-ethyl-1-hexyl and 3,5,5-trimethylhexyl) have also showed relatively good organogelation property.
4	Boc-Ala(1)-Aib(2)-β-Ala(3)-OMe organo gelator Boc-Ala(1)-Aib(2)-β-Ala(3)-OMe is a synthetic tripeptide which has the capability to undergo self-association so as to form thermoreversible transparent gels in the presence of various apolar solvents viz 1, 2-dichlorobenzene (DCB), monochloro benzene and benzene	Formation of thermoreversible transparent gels
5	Low Molecular Weight (LMW) organo gelators These gelators may produce either solid-fiber matrix or fluidfiber	

- matrix depending upon the physical intermolecular interactions. - Solid-fiber matrix may be formed when a heated mixture of the organogelators in apolar solvent is cooled down below the solubility limit of the organogelators. This results in the precipitation of the organogelators as fiber-like structures which undergoes physical interaction so as to form a gelled structure. These solid fiber-like structures align themselves into bundles. - fluid-fiber matrix is formed by the addition of polar solvent into a solution of amphiphiles in apolar solvents. Amphiphiles in apolar solvents are present as reverse micelles, which on addition of minute quantity of water forms tubular reverse micellar structures.	- Solid-fiber matrix organogels have improved mechanical properties as compared with the fluid-fiber matrix organogels. This can be attributed to the highly ordered structures present in the solid-fiber matrix organogels as compared to the simple chain entanglements in the fluid-fiber matrix organogels. - Apart from the above-mentioned organogels, various amphiphiles having the ability to form self-assembled structures in the presence of apolar solvents have also been tried.
--	--

Organogel Preparation¹¹

Most organogels are prepared by heating a mixture of the gelator and the liquid component to form an organic solution/ dispersion, followed by cooling of the latter, which sets into a gel. Heating allows dissolution of the gelator in the liquid. Following cooling, the solubility of the gelator in the liquid phase decreases, and gelator-solvent interactions are reduced, which results in the gelator molecules 'coming out' of solution. Gelator-gelator interactions lead to gelator self-assembly into well-defined aggregates such as tubules, rods and fibres. The physical organogels, held together by noncovalent forces, are thermoreversible; that is, following heating, the gel melts to the sol phase as the gelator aggregates dissolve in the organic liquid, whereas cooling the hot sol phase results in gelation. Mainly 3 methods are used for preparation of organogels like:

(1) Fluid-filled fiber mechanism

Firstly surfactants and co-surfactants mixtures were dissolved in apolar solvent and formation of reverse micelles was formed. After the addition of water, tubular reverse micelles were formed. Elongated tubular reverse micelle gets entangled to form a 3-

dimensional network, which immobilizes apolar solvent after the addition of water into tubular reverse micelles.

(2) Solid fiber mechanism

Apolar solvent and solid organogelator were heated and formation of apolar solution of organogelator. After cooling to room temperature, organogelator precipitates out as fibers which undergo physical interactions amongst each other thereby forming a 3-dimensional network structure, which immobilizes apolar solvent. To achieve a fine degree of subdivision of the particle & gelatinous character of those particles.

(3) Hydration method

Gel may be prepared by directly hydrating the inorganic chemical, which produces dispersed phase of the dispersion. In addition of water vehicle, other agents as propylene glycol, propyl gallate and hydroxyl propyl cellulose may be used to enhance gel formation.

Types Of Organogels

Different types of organogels are used for delivery of drugs and other pharmaceutical ingredients. Table 2 describe types, property, advantages and applications of different types of organogels.

Table 2 Types and property of organogels¹²⁻¹⁵.

S.No.	Types and property of organogels	Advantages and applications of organogels
1	Lecithin organogels - Extracted from various plants and animal tissues apart from the egg yolk - Experimental results indicate that the lecithin fails to initiate the process of gelification of the apolar solvent if the lecithin contains < 95% phosphatidyl content. - The organogels prepared using lecithin has been found to have an isotropic structure	- The lecithin-based organogels have been found to be thermodynamically stable, thermo reversible (sol-to-gel transition temperature at 40°C), transparent, viscoelastic, biocompatible and non-irritant - The lecithin organogels help either in the solubilization or accommodation of various guest molecules within its structure. These properties of the lecithin organogels have generated great potential for the use of the same as a controlled delivery vehicle.
2	Pluronic Lecithin Organogel (PLO) - PLO is a soy lecithin-based organogels which consists of isopropyl palmitate or isopropyl myristate, water and Pluronic F127 (also known as Poloxamer 407). - The apolar phase in the PLO constitutes 22 % (v/v) and hence is often regarded as micro-emulsion-based gel	- PLO is thermostable, viscoelastic and biocompatible in nature - PLO has also been found to produce minimal skin irritation. It has been used as a delivery vehicle for both hydrophobic and hydrophilic molecules for topical and transdermal applications
3	Premium lecithin organogels (PrLO) - The PrLO is a second generation lecithin organogel and has got higher thermostability apart from its non-greasy and non-tacky nature, which provides a cosmetically pleasing acceptability. - This gel do not have pluronic derivative, which results in the avoidance of the skin-irritation and thereby local skin-intolerance reactions.	- The use of PrLO as a carrier for drug delivery has indicated that the gel help in achieving improved bioavailability in the tissues by improving the penetration of the bioactive agents - This gel has been successfully used to accommodate various bioactive agents, viz. diclofenac, ibuprofen, ketoprofen and progesterone, and has been regarded as vehicle of choice for intradermal drug delivery.
4	Limonene GP1/PG organogel - The GP1 (dibutyl lauryl glutamide) / PG (propylene glycol) organogels can be prepared by mixing the appropriate amounts of GP1, limonene and PG with the subsequent incubation of the same at 120°C. When the mixture is cooled down, it forms a white gel. - It was found that the presence of limonene within the GP1/PG organogels resulted in the alteration of the rheological properties of the organogels though there was no significant change in the chemical stability of the organogels.	- Limonene, a terpene, has been found to be an excellent penetration enhancer and hence has been incorporated within various transdermal formulations for the improving the penetration of the bioactive agent across the transdermal layer, thereby improving the bioavailability of the bioactive agent within the dermal tissue - Apart from limonene, various other terpene-based penetration enhancers (e.g. linalool, farnesol and cineole) have also been incorporated successfully in GP1/PG organogels. The presence of penetration enhancers within the organogels results in the improvement of the rate permeation of the bioactive agents
5	Gelatin stabilized microemulsion based organogel (MBG) Gelatin is a protein which has been used as a structuring agent in various food preparations having excess of aqueous phase. It forms a gelled structure when a concentrated heated solution of gelatin having temperature in excess of 40°C is cooled down to a	Microemulsions are preferred for the development of gelatin stabilized organogels because of the thermostable nature and the ease of preparation of the same.

	- temperature below 35°C. - The addition of gelatin to the water-in-oil microemulsion results in the gelation of the whole micellar solution and the gel formed is transparent in nature.	The MBGs have been used to device topical and/or transdermal controlled delivery vehicle for hydrophobic bioactive agents.
6	Fatty acid derived sorbitan Organogels - These gelators are hydrophobic non-ionic molecules having surface active properties and have the ability to immobilize various solvents viz. isopropyl myristate, and vegetable oils. - These gelators form solid-fiber matrix when the heated solution of gelator in apolar solvent is cooled down. - The formation of the gel has been attributed to the formation of toroidal reverse micelles as the temperature is lowered. - The toroidal reverse micelles reorganize themselves to form rod-shaped tubules which subsequently undergo physical interaction amongst each other thereby forming a three-dimensional networked structure.	- The gels developed by using these gelators are opaque, thermoreversible and thermostable at room-temperature for weeks. - Organogels using fatty acid gelators may also be prepared by dissolving the gelators in a water-in-oil emulsion at a higher temperature followed by the decrease of the emulsion temperature. The decrease in the temperature results in the decrease in the solubility of the gelator with the subsequent precipitation and self-assembly of the gelators into network of tubules, which gets entangled so as to form a gelled structure.
7	Poly (ethylene) organogels The polyethylene organogels are colourless in nature, which are formed when the low molecular weight polyethylene is dissolved in mineral oil at a temperature >130°C and subsequently shocked cooled.	These organogels have been extensively used as ointment bases. The formation of gelled structure may be attributed to the physical interactions of the solid-fibers formed due to the precipitation of the polyethylene molecules.
8	Eudragit organogels Eudragit organogels are different from the organogels they are the mixtures of Eudragit (L or S) and polyhydric alcohols such as glycerol, propylene glycol and liquid polyethylene glycol containing high concentrations of Eudragit.	They show high gel rigidity and stability when drug concentration was low.
9	Sorbitan Monostearate Organogels - Sorbitan monostearate (span 60) and sorbitan monopalmitate (span 40) have been found to gel a number of organic solvents at low concentrations. - Prepared by heating the gelator/ liquid mixer in a water bath at 60°C and cooling of the resulting suspension.	They are used for delivery of hydrophilic vaccines and sorbitan monostearate.
10	L-alanine derivative organogels - Prepared from N-lauroyl-L-alanine methyl ester which gels in the organic solvents as soybean oil and triglycerides. - The system exists in the gel state at room temperature.	- It could act as a sustained release implant. - Used for delivery of rivastigmine and leuprolide drug.

Mechanism Of Drug Release

A drug may be regarded as a random network permeated by pores that are filled with a liquid component; substances that are soluble in the liquid component will tend to permeate through the gel by diffusion in solution through the space in the network. The rate of diffusion (the spontaneous transfer of solute from concentration regions in the solution where the concentration is lower, until there is a uniform distribution throughout) of substances through gels by this means will therefore be affected by those factors that normally affect simple diffusion in solution and by additional factors that are

associated with the presence of the gel network¹⁶. Fick's first law expresses the rate of diffusion of a solute, which is given by the equation.

$dm/dt = -DA(dc/dx)$, Where dm = amount of solute diffusing, dt = time, A = Area, dc/dx = concentration gradient and D = diffusion coefficient.

Physicochemical Property Of Organogels

In the present section, attempts will be made to discuss about the various physicochemical properties of the organogels. Table 3 describe physicochemical property of organogels

Table 3. Physicochemical property of organogels^{17,18,19}

S.No	Physicochemical property	Details about Physicochemical property
1	Viscoelasticity	The organogels behaves like a solid at lower shear rates and hence shows an elastic property. As the shear stress is increased, the physical interacting points amongst the fiber structures start getting weakened until the shear stress is high enough to disrupt the interactions amongst the fiber structures, when the organogels starts flowing. This behaviour may be best explained with the plastic flow behaviour.
2	Non-birefringence	The organogels when viewed under polarized light appears as a dark matrix. This can be accounted to the isotropic nature of the organogels which does not allow the polarized light to pass through the matrix. This property of the organogels of not allowing the polarized light to pass through it's matrix is regarded as non-birefringent.
3	Thermoreversibility	As the organogels are heated up above a critical temperature, the organogels loses its solid matrix-like structure and starts flowing. This has been attributed to the disruption in the physical interactions amongst the gelator molecules due to the increase in the thermal energy within the organogels. But as the heated organogels systems are subsequently cooled down, the physical interaction amongst the organogelators prevail and the organogels revert back to the more stable configuration.
4	Thermostability	The organogels are inherently thermostable in nature. Then stability of the organogels may be attributed to the ability of the gelators to undergo self-assembly, under suitable conditions, so as to form organogels. As the gelators undergo self-assembly, it results in the decrease in the total free energy of the system and renders the organogels as low-energy thermostable system.
5	Optical clarity	Depending on the composition of the organogels, the organogels may be transparent or opaque in nature. The lecithin organogels are transparent in nature while the sorbitan monostearate organogels are opaque in nature.
6	Chirality effects	- The presence of chirality in the LMW gelators has been found to affect the growth and the stability of the solid-fiber networks. Thermo reversibility of the gels formed due to the formation of the self assembled solid-fiber network has also been associated with the chirality. - The presence of chiral centers within the gelators helps in the formation of a compact molecular packing, which provides a thermodynamic and kinetic stability to the organogels system. Crown ether

7	Biocompatibility	phthalocyanine organogels are the excellent example of chiral organogels Initially, organogels were developed using various nonbiocompatible organogels which rendered the organogels nonbiocompatible. Of late, research on organogels using various biocompatible constituents has opened up new dimensions for the use of the same in various biomedical applications
---	------------------	---

Factor Affecting Organogels

Various factors related to nature of chemicals, solvents, environmental conditions and other parameters affect the preparation and stability of organogels. Table 4 describe factor affecting parameters which affect the preparation of organogel.

Table 4. Various parameters affect the property of organogels^{20,21}

S.No	Factors affecting parameters	Details about Factors affecting parameters
1	Organic solvent - Polar solvent - Non-aqueous solvent	- The effect of polar solvent introduces into spherical lecithin micelles may be associated with an increase in cross-sectional area of the lecithin polar region, in which the solvent is arranged - A non-aqueous solvent is not particularly limited as long as it replaces water of the bacterial cellulose hydrogel completely without destroying its shape for example- Polyethylene glycol, dimethyl ether
2	Phase Transition Temperature	It gives an insight into nature of microstructures that form the gelling cross linked network. For ex- a narrow PTI range is indicative of homogenous microstructures within the gel.
3	Salt addition	Salt may attract part of water of hydration of the polymer allowing more formation inter molecular secondary bond, this is known as salting out
4	Temperature	The effect of temperature depends on the chemistry of the polymer and its mechanism of interaction with the medium. If the temperature is reduced once the gel is in the solution, degree of hydration is reduced and gelation occurs. Gel resulting from the chemical cross linking often cannot be liquefied by dilution or temperature changes.
5	Molecular weight	Low molecular weight polymers require a high concentration to build up viscosity and to set to gel possibly
6	Surfactants	Gel characteristics can be varied by adjusting the proportion and concentration of the ingredients. Poloxamer 407 is a polyoxyethylene that function as a surfactant.
7	Physicochemical properties - Charge - Solubility - Molecular weight/ spatial configuration	- The presence of charged groups on a polymer favours mucoadhesion. Polyanions particularly, polycarboxy lates, are preferred to polycations - Mucoadhesives swell on contact with moisture, increasing the mobility of polymer molecules at the interface and exposing more sites for bond formation. - It favours change in entanglement and interaction after the polymer and mucins have interpenetrated.

Characterization Of Organogels

Evaluation and characterization is very important steps after the formation of organogels. Various characterization parameters are used for confirm the purity and stability of prepared organogels. Table 5 describe various characterization parameters and different techniques which are used for evaluation of these parameters.

Table 5. Characterization parameters and their evaluation techniques²²

S.No.	Characterization parameters	Techniques for Characterization parameters
1	Physicochemical properties - The isotopic nature and optical clarity organ gel study. - Establishing the hydrogen bonding as one of the major driving force for the self assembly of organogelator molecules in organic solvent - The knowledge of molecular packing within the organogel network	- Spectroscopic techniques, namely NMR and FTIR spectroscopy. - FTIR spectroscopy - Scanning and transmission electron microscopies, dynamic and static light scattering (elastic or quasielastic light scattering technique) small angle neutron scattering (SANS)
2	Rheological behaviour - Viscoelasticity - Swelling - Water content	- Scartazzini and Luisi performed the dynamic shear viscosity prepared using different types of organ gel solvent (eg. linear and cyclic alkenes, amines). The higher values obtained using linear alkenes were related to the higher state of structural organization organogels. - Gels can swell by absorbing liquid with an increase in volumes. Solvent penetrates the gel matrix, so that gel-gel interactions are replaced by gel solvent interaction. - Near infra red spectroscopy studies on lecithin/IPP/water organogel system by measuring the water absorption in the NIR region (1800-2200nm). In this region, water shows a strong absorption peaks at 918nm due to H-O-H stretching overtones, which are easily detectable and quantifiable
3	Phase transition temperature Phase transition temperature (PTT) (i.e. sol to gel or gel to sol) gives an insight into the nature of micro structure that form gelling cross linked network	Hot stage microscopy and high sensitivity differential scanning calorimetry have been reported to be useful as accurate and sensitive techniques.
4	Gelation Kinetics - Gel-sol and sol-Gel transitions - Gelation kinetics	- Inverse method - Turbidimetry method

5	In vitro drug release	• Franz diffusion cell
6	Safety and skin compatibility studies	• The irritation potential of loss has been assessed by carrying out human skin irritation study
7	Structural features • Molecular architecture of organogels • Hydrogen bonding	• NMR spectroscopy • FTIR spectroscopy

Applications And Uses Of Organogels

Organogels are used for delivery of drugs and many pharmaceutical ingredients through various routes like oral, parenteral, topical, transdermal etc. Table 6 describe various applications and uses of organogels.

Table 6. Applications and uses of organogels^{23,27}

S.No.	Application and uses	Detail study of Applications and uses
1	Topical drug delivery • Cosmetic • Ophthalmic • Ointments	• Therapeutic compounds of different chemical and physicochemical background such as muscle relaxants, steroids hormones, analgesics, antiemetic, and cardio vascular agents have been incorporated in the organogel with some encouraging results. • Used in the cosmetic and personal care markets. • Drug product like normal lachrymal turnover causes rapid clearance of solution and suspension dosage forms. • It is of various advantages like good tolerability, formation of a protective film over the cornea, protection from conjunctival adhesion. Methazolamide ineffective as an ophthalmic solution has been incorporated into carbomer and poloxamer gels for treatment of glaucoma.
2	Dermal drug delivery	• The muscle relaxants administered in lecithin –Isopropyl myristate organogel is shown to provide immediate relief of pain resulting from bruxism (tooth grinding) and tooth clenching. • Effective delivery of antipsoriatic agents and for drugs used in eczema. • Phospholipids organogel containing anti inflammatory macromolecule bromelain (15%) along with capsaicin (0.025%) has been found to be effective anti inflammatory composition.
3	Transdermal drug delivery	• Organogel systems have also been used as a matrix for transdermal transport of different therapeutic compounds. • The solubility of various drugs such as nifedipine, clonidine, scopolamine. And broxaterol was noted to be increase in lecithin –IPP solution compared with drug solubility in IPP alone, suggesting the solubility enhancing properties of the organogels. • Nicardipine, a calcium channel blocker, because of its low dose, short half live and extensive first pass metabolism, has been incorporated in the system in order to achieve systemic absorption through topical route.
4	Parenteral delivery	• L-alanine based injectable in situ forming organogels may be used for the delivery of labile macromolecular bioactive agents. • Experimental results indicates that the organogels system, when injected subcutaneously in rats releases the bioactive agents (e.g leuprolide) for a period of 14-25 days with subsequent degradation of the gelled structure.
5	Oral delivery	• Ibuprofen, a NSAID (non-steroidal anti-inflammatroly drug), was incorporated within the gelled structure. The release studies indicated that with the increase in the organogelator concentration within the organogel, there was a subsequent decrease in the release rate of the organogels. • In vivo studies in rats showed that the organogels may be used a controlled delivery vehicle for oral delivery of lipophilic compounds.
6	Organogels in drug delivery	• Organogels that have been studied for drug delivery include in situ forming organogels from L-alanine derivatives, Eudragit gels, lecithin gels, microemulsion-based gels (MBGs) and sorbitan monostearate gels.
7	Bioadhesives	• Bioadhesives of pharmaceutical interest are mucoadhesives this implies that the substrate for adhesion is the mucus itself. Many of the alternate routes of administration (buccal, ophthalmic, nasal, vaginal, etc) lend themselves bioadhesives because of the presence of mucosal tissue.
8	Cosmetic	• Gels have been employed in a variety of products including shampoo, fragrance products, dentifrices and skin and hair care preparation.
9	Dosage forms	• Glycogelatin gels are frequently used as a basis for medicated pastilles. They are used in the formulation of some suppositories e.g. Glycerin suppositories BP 1968.
10	Gelatin gels	• They are employed in the preparation of hard and soft capsules that may be used to mask the unpleasant tastes of solids and liquids.
11	Microbiological media	• Agar and gelatin gels are used as a solid media for the culture of microorganisms. • The diffusion of antibiotics, antiseptics, vitamins and enzymes through the culture media is used in the microbiological assays of these materials. Such diffusion produces zones of either retarded or enhanced growth on seeded agar plates depending on the activity of the diffusing substance.

CONCLUSION

There has been an exponential rise in exploring the possibility of the use of organogels as a drug delivery vehicle. This has been greatly motivated due to the longer shelf life, ease of preparation and thermo-reversible nature of the organogels-based formulations. Moreover, the topical delivery of new biotech- generated proteinaceous molecules in the protective non polar micro environment of these systems may help to protect these sensitive micromolecules from degradation during transport to the desired site. The few organogels that have been investigated for drug delivery have yielded interesting results, and it is hoped that some of

these will make it to the market and improve drug therapy for the benefit of patients.

REFERENCES

1. Vintilou A, Leroux JC. Organogels and their use in drug delivery -- A review. Journal of Controlled Release 2008; 125(3):179-192.
2. Aulton ME. The science of dosage form design, 2nd Edition, Churchill Livingstone New York 2002;89.
3. Tamilvanan S. Oil in water lipid emulsion: implications for parenteral and ocular delivering systems. Progress in lipid research 2004;489-533.
4. Suzuki M. L-Lysine based Gemini organogelators: their organogelation property. Org Biomol chem. 2003;22:4124-4134.
5. Lin Y, Kachar B, Weiss RG. Novel family of gelators of organic fluids and the structure of their gels. J Am Chem Soc. 1989; 111:5542-5551.

6. TERECH P. 12-D-Hydroxyoctadecanoic acid organogels: a small angle neutron scattering study. *J. Phys II France* 1992; 2:2181-2195.
7. FURMAN I, WEISS RG. Factors influencing the formation of thermally reversible gels comprised of cholesteryl 4-(2 anthryloxy)butanoate in hexadecane, 1-octanol, or their mixtures. *Langmuir* 1993; 9:2084-2088.
8. Terech P, Weiss RG. Low Molecular Mass Gelators of Organic Liquids and the Properties of Their Gels. *Chemical Reviews* 1997;97(8):3133-3160.
9. Malik S. A synthetic tripeptide as organogelator: elucidation of gelation mechanism. *J. Chem. Soc., Perkin Trans. 2* 2002; 2:1177 - 1186.
10. Maji SK. A synthetic tripeptide as a novel organo-gelator: a structural investigation. *Tetrahedron Letters* 2003;44(21):4103-4107.
11. Sahoo S. Organogels: Properties and Applications in drug delivery. *Designed Monomers and Polymers* 2011;14:95-108.
12. Liu XY, Sawant PD. Determination of the Fractal Characteristic of Nanofiber-Network Formation in Supramolecular Materials. *ChemPhysChem* 2002; 3(4): 374-377.
13. Sawant PD, Liu XY. Formation and Novel Thermomechanical Processing of Biocompatible Soft Materials. *Chemistry of Materials* 2002; 14(9): 3793-3798.
14. Kang L. SMGA gels for the skin permeation of haloperidol. *Journal of Controlled Release* 2005; 106(1-2):88-98.
15. Haering G, Luisi PL. Hydrocarbon gels from water-in-oil microemulsions. *The Journal of Physical Chemistry* 1986; 90(22): 5892-5895.
16. Rawling EA. Text book of pharmaceutics, 8th edition; baillier Tindal london, 120.
17. Abdallah DJ, Sirchio SA, Weiss RG. Hexatriacontane Organogels. The First Determination of the Conformation and Molecular Packing of a Low-Molecular-Mass Organogelator in Its Gelled State. *Langmuir* 2000;16(20):7558-7561.
18. Esch JH, Feringa BL. New Functional Materials Based on Self-Assembling Organogels: From Serendipity towards Design. *Angewandte Chemie International Edition* 2000; 39(13): 2263-2266.
19. Shechipunov YA. Lecithin organogels: rheological properties of polymer-like micelles formed in the presence of water. *Colloid J.* 1995;57:556-560.
20. Pal K, Banthia AK, Majumdar DK. Effect of heat treatment of starch on the properties of the starch hydrogels. *Materials Letters* 2008; 62(2):215-218.
21. Pal K, Banthia A, Majumdar D. Biomedical evaluation of polyvinyl alcohol-gelatin esterified hydrogel for wound dressing. *Journal of Materials Science: Materials in Medicine* 2007; 18(9): 1889-1894.
22. Wright A, Marangoni A. Formation, structure, and rheological properties of ricinoleic acid-vegetable oil organogels. *Journal of the American Oil Chemists' Society* 2006; 83(6):497-503.
23. Remington, *The Science & practice of pharmacy*, 21st edition, Williams & Wilkins, New York, 884.
24. Martin Alfred, *physical pharmacy*, 4th edition, Williams & Wilkins, New York, 496.
25. Pisal SS, Reddy P, Ketkar AR, Patil VB, Paradkar AR. Transnasal absorption of melatonin from thermo-reversible gels of pluronic PF 127 gels. Paper presented at 62nd World Pharmaceutical Congress, September 3, 2002, Nice, France, DDT-P-039.
26. Pisal SS, Paradkar AR, Mahadik KR, Kadam SS. Pluronic gels for nasal delivery of vitamin B12. Part I. Preformulation study. *Int. J. Pharm.* 2004;270:37-45.