

AS: 104.013

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO



No. 07-068704- -00001-0000

Fecha: 2008-06-06 16:44:02 Dep. 2020 NUECREACIONE
Tra. 11 PATENTE/YII Evs: 378 FASENACIONAL
Art. 538 PETICION Folios: 2

Señores

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO

DIVISION DE NUEVAS CREACIONES

E. S. D.

REF.: Expediente No. 07 068704 - Solicitud de Patente FASE NACIONAL

Corresp. a la Solicitud Internacional PCT/GB2005/004726


a nombre de **ASTRAZENECA AB**

Titulada: DERIVADOS DE OXADIAZOL COMO INHIBIDORES DE DGAT

JUAN PABLO CADENA SARMIENTO, vecino de Bogotá, portador de la tarjeta profesional No. 57492 del Consejo Superior de la Judicatura, obrando como apoderado de la sociedad **ASTRAZENECA AB**, según lo acredita el documento de poder presentado en esa División, de acuerdo con el Artículo 44 de la Decisión 486 de la Comisión de la Comunidad Andina, me permito solicitar que se examine si la invención contenida en la solicitud en referencia es patentable.

De acuerdo con lo anterior, me permito anexar el recibo oficial de caja No. 08-39.197 por valor de \$420.000.

De la Señora Jefe atentamente,


JUAN PABLO CADENA SARMIENTO

C.C. No. 80.410.337 de Usaquén

T.P. No. 57492 del C.S.J.

Cra 7 No. 71-21 Torre B, Piso 4

Bogotá, 06 de junio de 2008

MEV/CAG/aac

45/104/19

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO

NIT : 900176089-2

RECIBO OFICIAL DE CAJA : 08 - 39,197
FECHA : MAYO 16 DE 2008

***** CONSIGNACION *****

DEPOSITANTE	TIPO PAGO	BANCO	CUENTA	No. PAGO	FECHA PGO	Vr. PAGO
BRIGARD & CASTRO LINA	CONSIGNACION	BANCO DE BOGOTA	062754387	760622	16/05/2008	31,380,000.00

***** CONCEPTO *****


CANT.	RENTISTAS	CONCEPTO	TOTAL CONCEPTO
50005-01-01	SOLICITUDES	85 EXAMEN DE PATENTABILIDAD DE INVEN	420,000.00
		TOTAL :	\$ 420,000.00

SON: CUATROCIENTOS VEINTE MIL PESOS

RESPONSABLE :




RECIBO DE CAJA APLICADO AL EXPEDIENTE No. _____

 INDUSTRIA Y COMERCIO DIVISION DE NUEVAS CREACIONES TARJETA ARCHIVO TEMÁTICO		PATENTE DE INVENCION ☒ PCT ☒ MODELO DE UTILIDAD ☐ DISEÑO INDUSTRIAL ☐	
(21) N° de solicitud:		(22) Fecha de solicitud:	
(51) Clasificación Internacional:			
(71) Solicitante(s) ASTRAZENECA AB			
(72) Inventor(es): Alan Martin BIRCH, Suzanne Saxon BOWKER, Roger John BUTLIN, Craig Samuel DONALD, William MCCOULL, Thorsten NOWAK, Allyn PLOWRIGHT.			
(74) Agente: JUAN PABLO CADENA SARMIENTO			
(30) Prioridad	(31) No. Prioridad	(32) Fecha	(33) País
	0427328.0	14-12-2004	GB
	0507403.4	13-04-2005	GB
	0520383.1	07-10-2005	GB
(85) Fecha inicio fase nacional: 14-07-2007		(87) Publicación internacional:	
(86) Datos relativos a la presentación de la sol. PCT		Fecha: 22-06-2006	
Fecha de presentación de la solicitud: 09-12-2005		No. Publicación: WO 2006/064189 A1	
No. de solicitud PCT/GB2005/004726			
(54) Título: DERIVADOS DE OXADIAZOL COMO INHIBIDORES DE DGAT			
(55) Reivindicación(es): 1. Un compuesto de fórmula (I)			
 (I)			
CONTINUA AL RESPALDO ...			

(1) N° de solicitud:					(51) Int Cl:	
(22) Fecha de solicitud:					(71) Solicitante(s): ASTRAZENECA AB	
30	Prioridad	(31)	No. Prioridad	(32)	Fecha	(33) Pais
			0427328.0		14-12-2004	GB
			0507403.4		13-04-2005	GB
			0520383.1		07-10-2005	GB
(85) Fecha inicio fase nacional: 14-07-2007					(87) Publicación internacional	
(86) Datos relativos a la presentación de la solicitud PCT					Fecha: 22-06-2006	
Fecha de presentación de la solicitud: 09-12-2005					N° publicación: WO 2006/064189 A1	
No. de solicitud PCT/GB2005/004726						
(54) Título: DERIVADOS DE OXADIAZOL COMO INHIBIDORES DE DGAT						

Reivindicación(es):

1. Un compuesto de fórmula (I):



o su sal o profármaco farmacéuticamente aceptable, en donde:

R¹ es un grupo arilo opcionalmente sustituido o heteroarilo opcionalmente sustituido, en donde los sustituyentes opcionales son uno o más grupos seleccionados de un grupo -Z^a, un grupo -X²-(CR⁵²R⁵³)_w-Z^a, un grupo -X²-(CR⁵²R⁵³)_a-X³-Z^a, un grupo -(CR⁵²R⁵³)_a-X³-Z^a o un grupo funcional (el cual es distinto de un grupo -X²-(CR⁵²R⁵³)_w-Z^a o un grupo -X²-(CR⁵²R⁵³)_a-X³-Z^a);

Y es un enlace directo, o un grupo (CR⁴⁰R⁴¹)_s o -X⁶(CR⁴⁰R⁴¹)_t - donde cada R⁴⁰ y R⁴¹ se seleccionan independientemente de hidrógeno, (1-4C)alquilo, hidroxilo, halo, halo(1-4C)alquilo, amino, ciano, (1-4C)alcoxi, (1-4C)haloalcoxi o ((1-3C)alquil)CONH-, s es un entero de 1 a 6 y t es un entero de 1 a 6, siempre que el átomo X⁶ del grupo -X⁶(CR⁴⁰R⁴¹)_t se una al grupo R² y que un solo átomo de carbono hibridado sp³ no lleve dos o más enlaces a un heteroátomo, a menos que el heteroátomo es un halo;

R² es un arilo opcionalmente sustituido, un cicloalquilo opcionalmente sustituido o un grupo heterocíclico opcionalmente sustituido, en donde sustituyentes opcionales son uno o más grupos seleccionados de un grupo -Z, un grupo -X-(CR⁴²R⁴³)_u-Z, un grupo -X-(CR⁴²R⁴³)_v-X¹-Z o un grupo -(CR⁴²R⁴³)_v-X¹-Z o un grupo funcional (el cual es distinto de un grupo -X-(CR⁴²R⁴³)_u-Z o un grupo -X-(CR⁴²R⁴³)_v-X¹-Z);

en donde Z y Z^a se seleccionan independientemente de un grupo hidrocarbilo o un grupo heterocíclico o una combinación de los mismos, en donde el grupo Z y Z^a se sustituyen opcionalmente en cualquier átomo disponible por uno o más grupos funcionales, o por un grupo -X⁷-(CR⁶²R⁶³)_b-R⁶⁴;

X, X¹, X², X³, X⁶ y X⁷ son grupos enlazantes seleccionados independientemente de -C(O)x-, -O-, -S(O)y-, -NR⁴⁴-, -C(O)NR⁴⁴-, -OC(O)NR⁴⁴-, -CH=NO-, -NR⁴⁴C(O)x-, -NR⁴⁴CONR⁴⁵-, -S(O)2NR⁴⁴- o -NR⁴⁴S(O)2- donde x es un entero de 1 ó 2, y es 0, 1 ó 2, y R⁴⁴ y R⁴⁵ se seleccionan independientemente de hidrógeno o (1-6C)alquilo,

u y w se seleccionan independientemente de 0 o un entero de desde 1 a 6;

v, a y b se seleccionan independientemente de un entero de desde 1 a 6;

cada R⁴², R⁴³, R⁵², R⁵³, R⁶² y R⁶³ se selecciona independientemente de hidrógeno, (1-4C)alquilo, hidroxilo, halo, halo(1-4C)alquilo, amino, ciano, (1-4C)alcoxi, (1-4C)haloalcoxi, ((1-3C)-alquil)CONH-, carboxi o una réplica de ácido carboxílico o bioisómero del mismo, y R⁶⁴ es un grupo funcional.

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
22 June 2006 (22.06.2006)

PCT

(10) International Publication Number
WO 2006/064189 A1

(51) International Patent Classification:

C07D 271/10 (2006.01) A61P 3/04 (2006.01)
C07D 413/12 (2006.01) A61P 3/10 (2006.01)
A61K 31/4245 (2006.01)

(21) International Application Number:

PCT/GH2005/004726

(22) International Filing Date:

9 December 2005 (09.12.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0427328.0 14 December 2004 (14.12.2004) GB
0507403.4 13 April 2005 (13.04.2005) GB
0520383.1 7 October 2005 (07.10.2005) GB

(71) Applicant (for all designated States except MG, US): AS-
TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).(71) Applicant (for MG only): ASTRAZENECA UK LIM-
ITED [GB/GB]; 15 Stanhope Gate, London, Greater, Lon-
don W1K 1LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BIRCH, Alan,
Martin [GB/GB]; AstraZeneca R & D Alderley, Alder-
ley Park, Macclesfield, Cheshire SK10 4TG (GB).
BOWKER, Suzanne, Saxon [GB/GB]; AstraZeneca
R & D Alderley, Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB). BUTLIN, Roger, John [GB/GB]; As-
traZeneca R & D Alderley, Alderley Park, Macclesfield,
Cheshire SK10 4TG (GB). DONALD, Craig, Samuel
[GB/GB]; AstraZeneca R & D Alderley, Alderley Park,

Macclesfield, Cheshire SK10 4TG (GB). MCCOULL,
William [GB/GB]; AstraZeneca R & D Alderley, Alderley
Park, Macclesfield, Cheshire SK10 4TG (GB). NOWAK,
Thorsten [DE/GB]; AstraZeneca R & D Alderley,
Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
PLOWRIGHT, Allyn [GB/GB]; AstraZeneca R & D
Alderley, Alderley Park, Macclesfield, Cheshire SK10
4TG (GB).

(74) Agent: GLOBAL INTELLECTUAL PROPERTY; As-
traZeneca AB, S-151 85 Södertälje (SE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AB, AG, AI, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, GU, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MY, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: OXADIAZOLI DERIVATIVES AS DGAT INHIBITORS



(I)

(57) Abstract: Compounds of formula (I), and salts and pro-drugs thereof: Formula (I) wherein for example R^1 is optionally sub-
stituted aryl or heteroaryl; Y is a linking group selected from, for example, a direct bond, and a (substituted) alkyl chain; R^2 is an
optionally substituted aryl, an optionally substituted cycloalkyl or an optionally substituted heterocyclic group; are described. Pro-
cesses to make such compounds and their use as DGAT inhibitors, for example in the treatment of obesity, are also described.

OXADIAZOLE DERIVATIVE AS DGAT INHIBITORS

The present invention relates to compounds which inhibit acetyl CoA(acetyl coenzyme A):diacylglycerol acyltransferase (DGAT1) activity, processes for their preparation, pharmaceutical compositions containing them as the active ingredient, methods for the treatment of disease states associated with DGAT1 activity, to their use as medicaments and to their use in the manufacture of medicaments for use in the inhibition of DGAT1 in warm-blooded animals such as humans. In particular this invention relates to compounds useful for the treatment of type II diabetes, insulin resistance, impaired glucose tolerance and obesity in warm-blooded animals such as humans, more particularly to the use of these compounds in the manufacture of medicaments for use in the treatment of type II diabetes, insulin resistance, impaired glucose tolerance and obesity in warm-blooded animals such as humans.

Acyl CoA:diacylglycerol acyltransferase (DGAT) is found in the microsomal fraction of cells. It catalyzes the final reaction in the glycerol phosphate pathway, considered to be the main pathway of triglyceride synthesis in cells by facilitating the joining of a diacylglycerol with a fatty acyl CoA, resulting in the formation of triglyceride. Although it is unclear whether DGAT is rate-limiting for triglyceride synthesis, it catalyzes the only step in the pathway that is committed to producing this type of molecule [Lehner & Kuksis (1996) Biosynthesis of triacylglycerols. Prog. Lipid Res. 35: 169-201].

Two DGAT genes have been cloned and characterised. Both of the encoded proteins catalyse the same reaction although they share no sequence homology. The DGAT1 gene was identified from sequence database searches because of its similarity to acyl CoA:cholesterol acyltransferase (ACAT) genes. [Cases *et al* (1998) Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. Proc. Natl. Acad. Sci. USA 95: 13018-13023]. DGAT1 activity has been found in many mammalian tissues, including adipocytes.

Because of the previous lack of molecular probes, little is known about the regulation of DGAT1. DGAT1 is known to be significantly up-regulated during adipocyte differentiation.

Studies in gene knockout mice has indicated that modulators of the activity of DGAT1 would be of value in the treatment of type II diabetes and obesity. DGAT1

- 2 -

knockout (*Dgat1*^{-/-}) mice, are viable and capable of synthesizing triglycerides, as evidenced by normal fasting serum triglyceride levels and normal adipose tissue composition.

Dgat1^{-/-} mice have less adipose tissue than wild-type mice at baseline and are resistant to diet-induced obesity. Metabolic rate is ~20% higher in *Dgat1*^{-/-} mice than in wild-type mice on both regular and high-fat diets [Smith *et al* (2000) Obesity resistance and multiple mechanisms of triglyceride synthesis in mice lacking DGAT. *Nature Genetics* 25: 87-90]. Increased physical activity in *Dgat1*^{-/-} mice partially accounts for their increased energy expenditure. The *Dgat1*^{-/-} mice also exhibit increased insulin sensitivity and a 20% increase in glucose disposal rate. Leptin levels are 50% decreased in the *Dgat1*^{-/-} mice in line with the 50% decrease in fat mass.


When *Dgat1*^{-/-} mice are crossed with *ob/ob* mice, these mice exhibit the *ob/ob* phenotype [Chen *et al* (2002) Increased insulin and leptin sensitivity in mice lacking acyl CoA:diacylglycerol acyltransferase *J. Clin. Invest.* 109:1049-1055] indicating that the *Dgat1*^{-/-} phenotype requires an intact leptin pathway. When *Dgat1*^{-/-} mice are crossed with *Agouti* mice a decrease in body weight is seen with normal glucose levels and 70% reduced insulin levels compared to wild type, *agouti* or *ob/ob* *Dgat1*^{-/-} mice.

Transplantation of adipose tissue from *Dgat1*^{-/-} mice to wild type mice confers resistance to diet-induced obesity and improved glucose metabolism in these mice [Chen *et al* (2003) Obesity resistance and enhanced glucose metabolism in mice transplanted with white adipose tissue lacking acyl CoA:diacylglycerol acyltransferase *J. Clin. Invest.* 111: 1715-1722].

International Patent Applications WO2004/047755 (Tularik and Japan Tobacco) and WO2005/013907 (Japan Tobacco and Amgen) describe fused bicyclic nitrogen-containing heterocycles which are inhibitors of DGAT-1. JP2004-67635 (Otsuka Pharmaceuticals) describes thiazoleamido substituted phenyl compounds which are further substituted with alkylphosphonates and which inhibit DGAT-1. WO2004/100881 (Bayer) describes biphenylamino compounds substituted with imidazole, oxazole or thiazole which inhibit DGAT-1.

Accordingly, the present invention provides a compound of formula (I)

- 3 -



(I)

or a pharmaceutically-acceptable salt or prodrug thereof, wherein:

R^1 is an optionally substituted aryl or optionally substituted heteroaryl group,

5 wherein the optional substituents are one or more groups selected from a group $-Z^a$, a group $-X^2-(CR^{52}R^{53})_w-Z^a$, a group $-X^2-(CR^{52}R^{53})_a-X^3-Z^a$, a group $-(CR^{52}R^{53})_aX^3-Z^a$ or a functional group (which is other than a group $-X^2-(CR^{52}R^{53})_w-Z^a$ or a group $-X^2-(CR^{52}R^{53})_a-X^3-Z^a$);

10 Y is a direct bond, or a group $(CR^{40}R^{41})_s$ or $-X^6(CR^{40}R^{41})_t-$ where each R^{40} and R^{41} is independently selected from hydrogen, (1-4C)alkyl, hydroxy, halo, halo(1-4C)alkyl, amino, cyano, (1-4C)alkoxy, (1-4C)haloalkoxy or ((1-3C)alkyl)CONH-, s is an integer of from 1 to 6 and t is an integer of from 1 to 6, provided that the X^6 atom of the group $-X^6(CR^{40}R^{41})_t-$ is attached to the R^2 group and that a single sp^3 hybridised carbon atom does not carry two or more bonds to a heteroatom unless the heteroatom is a halo;

15 R^2 is an optionally substituted aryl, an optionally substituted cycloalkyl or an optionally substituted heterocyclic group, wherein optional substituents are one or more groups selected from a group $-Z$, a group $-X-(CR^{42}R^{43})_u-Z$, a group $-X-(CR^{42}R^{43})_v-X^1-Z$ or a group $-(CR^{42}R^{43})_vX^1-Z$ or a functional group (which is other than a group $-X-(CR^{42}R^{43})_u-Z$ or a group $-X-(CR^{42}R^{43})_v-X^1-Z$);

20 wherein Z and Z^a are independently selected from a hydrocarbyl group or a heterocyclic group or a combination thereof, wherein the group Z and Z^a is optionally substituted on any available atom by one or more functional groups, or by a group $-X^7-(CR^{62}R^{63})_bR^{64}$;

25 X , X^1 , X^2 , X^3 , X^6 and X^7 are linking groups independently selected from $-C(O)_x-O-$, $-S(O)_y-$, $-NR^{44}-$, $-C(O)NR^{44}-$, $-OC(O)NR^{44}-$, $-CH=NO-$, $-NR^{44}C(O)-$, $-NR^{44}CONR^{45}-$, $-S(O)_2NR^{44}-$ or $-NR^{44}S(O)_2-$ where x is an integer of 1 or 2, y is 0, 1 or 2, and R^{44} and R^{45} are independently selected from hydrogen or (1-6C)alkyl,

u and w are independently selected from 0 or an integer of from 1 to 6;

v , a and b are independently selected from an integer of from 1 to 6;

- 4 -

each R^{42} , R^{43} , R^{52} , R^{53} , R^{62} and R^{63} is independently selected from hydrogen, (1-4C)alkyl, hydroxy, halo, halo(1-4C)alkyl, amino, cyano, (1-4C)alkoxy, (1-4C)haloalkoxy, ((1-3C)alkyl)CONH-, carboxy or a carboxylic acid mimic or bioisostere thereof, and R^{64} is a functional group.

5 As used herein, the term "functional group" includes halo, halo(1-6C)alkyl, cyano, nitro, $-C(O)_nR^{20}$, a carboxylic acid mimic or bioisostere thereof, $-OR^{20}$, $-S(O)_mR^{20}$, $-OS(O)_2R^{20}$, $-NR^{21}R^{22}$, $-C(O)NR^{21}R^{22}$, $-OC(O)NR^{21}R^{22}$, $-CH=NOR^{20}$, $-NR^{21}C(O)_nR^{20}$, $-NR^{20}CONR^{21}R^{22}$, $-N=CR^{21}R^{22}$, $-S(O)_2NR^{21}R^{22}$ or $-NR^{21}S(O)_2R^{22}$ where R^{20} , R^{21} and R^{22} are independently selected from hydrogen or optionally substituted hydrocarbonyl or
10 optionally substituted heterocyclyl, or R^{21} and R^{22} together with the nitrogen atom to which they are attached form an optionally substituted ring having from 3 to 10 atoms, which optionally contains further heteroatoms such as $S(O)_m$, oxygen and nitrogen, n is an integer of 1 or 2, m is 0 or an integer of 1-2.

Suitable optional substituents for hydrocarbonyl groups or heterocyclic groups R^{20} ,
15 R^{21} and R^{22} (including for rings formed by $NR^{21}R^{22}$) include halo, halo(1-4C)alkyl (such as trifluoromethyl, difluoromethyl or fluoromethyl), mercapto, hydroxy, alkoxy, oxo, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyalkoxy, aryloxy (where the aryl group may be substituted by halo, cyano, nitro, hydroxy(1-4C)alkyl, halo(1-4C)alkyl, amino, (1-4C)alkoxy, (1-4C)haloalkoxy, ((1-3C)alkyl)CONH-, carboxy or a carboxylic acid mimic or
20 bioisostere thereof), cyano, nitro, amino, mono- or di-alkyl amino, alkylamido, oximino (for example hydroxyimino or alkyloxyimino), carbamoyl, carboxy or a carboxylic acid mimic or bioisostere thereof, or $-S(O)_mR^{23}$ where m is as defined above and R^{23} is alkyl (optionally substituted by one or more groups selected from hydroxy, halo, amino, cyano, ((1-3C)alkyl)CONH-, carboxy or a carboxylic acid mimic or bioisostere thereof), (1-
25 6C)alkoxy, (1-6C)alkoxycarbonyl, carbamoyl, N-((1-6C)alkyl)carbamoyl, halo(1-6C)alkyl (such as trifluoromethyl), (1-6C)alkylsulphonyl, (1-6C)alkylsulphinyl. Heterocyclic groups R^{20} , R^{21} and R^{22} may also be optionally substituted by one or more hydrocarbonyl groups such as (1-4C)alkyl.

In this specification the term "alkyl" includes both straight and branched chain
30 alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms. Unless otherwise stated the term "alkyl" advantageously refers to chains with 1-10 carbon

atoms, suitably from 1-6 carbon atoms, preferably 1-4 carbon atoms.

In this specification the term "alkoxy" means an alkyl group as defined hereinbefore linked to an oxygen atom.

It is to be understood that optional substituents on any group may be attached to
5 any available atom as appropriate unless otherwise specified, including heteroatoms provided that they are not thereby quaternised.

In this specification the term "heteroatom" refers to non-carbon atoms such as oxygen, nitrogen or sulphur atoms. In addition, where the heteroatom may have a single valency, it may comprise a halo. The terms "alkenyl" and "alkynyl" refer to unsaturated
10 straight or branched structures, which unless specified otherwise, contain for example from 2 to 10, preferably from 2 to 6 carbon atoms. Cyclic moieties such as cycloalkyl and cycloalkenyl are similar in nature but have at least 3 carbon atoms. Examples of alkyl, alkenyl and cycloalkyl groups are given hereinafter, such as examples of (1-6C)alkyl, (3-8C)cycloalkyl etc.

15 References to aryl groups include aromatic carbocyclic groups such as phenyl and naphthyl, as well as partially aromatic groups such as indenyl and indanyl. The term "aralkyl" refers to aryl substituted alkyl groups such as benzyl.

The term "heterocyclyl" or "heterocyclic" includes saturated or unsaturated rings, which may be aromatic, non-aromatic rings or partially aromatic, for example containing
20 from 3 to 20, suitably from 4 to 10 ring atoms, at least one of which is a heteroatom such as oxygen, sulphur or nitrogen. They may be mono- or bicyclic ring systems, wherein one or both rings may be saturated or unsaturated, for example they may be aromatic. In particular, bicyclic ring systems will comprise fused 5,6-membered or 6,6-membered rings.

"Heteroaryl" refers to those heterocyclic groups described above which have an
25 aromatic character. Where "heteroaryl" is a bi-cyclic ring system, then at least one ring is aromatic and one or both rings contain ring heteroatoms.

In general, examples of heterocyclyl rings include furyl, thienyl, pyrrolyl, pyrrolidinyl, imidazolyl, triazolyl, thiazolyl, tetrazolyl, oxazolyl, isoxazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, indolyl, quinolinyl, isoquinolinyl,
30 quinoxalinyl, benzthiazolyl, benzoxazolyl, benzothienyl and benzofuryl. Further examples include thiadiazolyl, oxadiazolyl and isothiazolyl, morpholino, piperazinyl and piperidinyl. Further examples of monocyclic heteroaryl rings are those listed hereinafter as suitable

examples of HET-1^{a-f}. Further examples of bicyclic heterocyclic rings are those listed hereinafter as suitable examples of HET-2^{a-b}. Further examples of monocyclic heterocyclic (non heteroaryl) rings are those listed hereinafter as suitable examples of HET-3^{b,c,e}. Suitable examples of bicyclic heteroaryl rings include indolyl, quinoliny, isoquinoliny, quinoxaliny, benzthiazolyl, benzoxazolyl, benzothienyl, benzofuryl, benzimidazolyl, benzodioxolanyl, pyrrolopyridyl, quinazoliny and naphthyridiny. It will be understood that structures such as 2-oxo-2,3-dihydro-1H-benzimidazolyl and oxothiadiazolyl which fall within the definition of the term heteroaryl, retain their aromatic characteristics in both rings by virtue of tautomerism.

Other expressions used in the specification include "hydrocarbyl" which refers to any structure comprising carbon and hydrogen atoms. These may be arranged in rings or chains or combinations in which rings are joined to chains or to further rings, or a fused to further rings. Generally, hydrocarbyl groups will contain from 1 to 20, for instance from 1-12 carbon atoms. These may be alkyl, alkenyl, alkynyl, aryl, aralkyl, aralkenyl, aralkynyl, cycloalkyl or cycloalkenyl, wherein any cyclic moiety such as aryl, aralkyl, cycloalkyl or cycloalkenyl are optionally substituted with alkyl, alkenyl, alkynyl and/or with further cyclic moieties, and where any alkyl, alkenyl or alkynyl groups are optionally substituted with cycloalkyl, or cycloalkenyl. The term cycloalkyl also includes bi- and tri-cycloalkyl rings, such as adamantyl and bicyclo[2.2.2]octanyl

Suitable combinations of rings and chains which are comprised by the term hydrocarbyl include

a) cycloalkyl linked to a (1-6C)alkyl group (in particular cyclohexylmethyl, cyclopentylmethyl, cyclobutylmethyl, cyclohexylethyl), or to two (1-6C)alkyl groups (for example methylcyclobutylmethyl);

b) cyclohexyl linked to a second cyclohexyl or a cyclopentyl group by a direct bond, or with a (1-6C)alkyl group linker;

c) a phenyl group linked to a second phenyl group by a direct bond, or with a (1-6C)alkyl group linker;

d) a (3-8C)cycloalkyl group (such as cyclohexyl or cyclopentyl) linked to a phenyl group by a direct bond or with a (1-6C)alkyl linker;

e) a benzyl or methylphenyl (such as tolyl) group.

References to a "combination" of hydrocarbyl and heterocyclic groups refer to moieties which contain one or more heterocyclic groups joined together, or one or more heterocyclic groups joined to one or more hydrocarbyl groups.

Suitable combinations of hydrocarbyl and heterocyclic groups include a
5 heterocyclyl group (such as pyridyl, morpholino, thiomorpholino, piperazinyl or piperidinyl) linked to (or substituted by) a hydrocarbyl group (such as a (1-6C)alkyl group and/or a (3-8C)cycloalkyl group; in particular a (1-6C)alkyl group). For example methylpyridyl (wherein the methyl may be further substituted by a functional group such as carboxy), benzylpiperazine, (methyl)oxopyridazine, (methyl)oxothiadiazole, (optionally
10 carboxy substituted)methylpiperidyl, (optionally carboxy substituted)methylpiperidylmethyl, (optionally carboxy substituted)dimethylpiperidyl, (optionally carboxy substituted)ethylpiperidyl and (cyclopropylmethyl)piperazinyl.

Unless specified otherwise, the expression "haloalkyl" refers to alkyl groups which carry at least one halo substituent. This includes perhalo groups where all hydrogen atoms
15 are replaced by halo such as fluoro.

It is to be understood that optional substituents on any group may be attached to any available atom as appropriate unless otherwise specified, including heteroatoms provided that they are not thereby quaternised.

Within this specification composite terms are used to describe groups comprising
20 more than one functionality such as -(1-6C)alkylNHSO₂(1-6C)alkyl. Such terms are to be interpreted in accordance with the meaning which is understood by a person skilled in the art for each component part. For example -(1-6)alkylNHSO₂(1-6C)alkyl includes -methylaminosulfonylmethyl, -methylaminosulfonylethyl, -ethylaminosulfonylmethyl, and -propylaminosulfonylbutyl.

25 Where optional substituents are chosen from "0, 1, 2 or 3" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups. An analogous convention applies to substituents chosen from "0, 1 or 2" groups and "1 or 2" and any other analogous groups.

30 Substituents may be present at any suitable position on, for example, an alkyl group. Therefore, hydroxy substituted (1-6C)alkyl includes hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl and 3-hydroxypropyl.

Examples of (1-4C)alkyl include methyl, ethyl, propyl and isopropyl; examples of (1-6C)alkyl include methyl, ethyl, propyl, isopropyl, t-butyl, pentyl, iso-pentyl, 1-2-dimethylpropyl and hexyl; examples of (2-6C)alkenyl include ethenyl, propenyl, isopropenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-methylpropenyl and hexenyl; examples of (2-6C)alkynyl include ethynyl, propynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl and hexynyl; examples of (1-4C)alkoxy include methoxy, ethoxy, propoxy, isopropoxy and tert-butoxy; examples of (1-6C)alkoxy include methoxy, ethoxy, propoxy, isopropoxy, tert-butoxy and pentoxy; examples of (1-6C)alkoxy(1-6C)alkyl include methoxymethyl, ethoxymethyl, methoxyethyl, propoxymethyl, isopropoxymethyl, pentoxyethyl, methoxyhexyl and tert-butoxybutyl; examples of (3-8C)cycloalkyl include (3-6C)cycloalkyl (cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl), cycloheptyl and cyclooctyl; examples of (3-8C)cycloalkoxy include cyclopropoxy, cyclobutoxy, cyclopentoxo, cyclohexyloxy, cyclopentyloxy and cyclooctyloxy; examples of (3-8C)cycloalkyl(1-6C)alkyl include cyclopropylmethyl, cyclopropylethyl, cyclopropylbutyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopentylethyl and cyclooctylpropyl; examples of (3-8C)cycloalkoxy(1-6C)alkyl include cyclopropoxymethyl, cyclopropoxyethyl, cyclopropoxybutyl, cyclobutoxymethyl, cyclopentoxymethyl, cyclohexyloxymethyl, cyclopentoxyethyl and cyclooctyloxypropyl; examples of (3-8C)cycloalkoxy(1-6C)alkoxy include cyclopropoxymethoxy, cyclopropoxyethoxy, cyclopropoxybutoxy, cyclobutoxymethoxy, cyclopentoxymethoxy, cyclohexyloxymethoxy, cyclopentoxyethoxy and cyclooctyloxypropoxy; examples of (3-8C)cycloalkoxy(1-6C)alkoxy(1-6C)alkyl include cyclopropoxymethoxymethyl, cyclopropoxyethoxymethyl, cyclopropoxybutoxymethyl, cyclobutoxymethoxyethyl, cyclopentoxymethoxypropyl, cyclohexyloxymethoxymethyl, cyclopentoxyethoxymethyl and cyclooctyloxypropoxymethyl; examples of halo are chloro, bromo, iodo and fluoro; examples of halo(1-6C)alkyl include halo(1-4C)alkyl such as chloromethyl, fluoroethyl, fluoromethyl, fluoropropyl, fluorobutyl, dichloromethyl, difluoromethyl, 1,2-difluoroethyl and 1,1-difluoroethyl as well as perhalo(1-6C)alkyl (including perhalo(1-4C)alkyl) such as trifluoromethyl, pentafluoroethyl, and heptafluoropropyl; examples of halo(1-6C)alkoxy include halo(1-4C)alkoxy such as chloromethoxy, fluoroethoxy and fluoromethoxy, difluoromethoxy, as well as perhaloalkoxy such as pentafluoroethoxy, trifluoromethoxy and heptafluoropropoxy; examples of hydroxy(1-6C)alkyl include hydroxy(1-4C)alkyl

such as hydroxy methyl, 1-hydroxyethyl, 2-hydroxyethyl and 3-hydroxybutyl; example of **carboxy(1-6C)alkyl** include carboxy(1-4C)alkyl, such as carboxymethyl, carboxyethyl, carboxypropyl and carboxybutyl; examples of **amino(1-6C)alkyl** include aminomethyl, aminoethyl, 2-aminopropyl, 3-aminopropyl, 2-aminoiso-propyl, aminobutyl and

5 2-aminotert-butyl; examples of **(1-6C)alkylamino** include (1-4C)alkylamino such as methylamino, ethylamino and propylamino; examples of **di-((1-6C)alkyl)amino** include di-(1-4C)alkylamino such as dimethylamino, *N*-ethyl-*N*-methylamino, diethylamino, *N*-methyl-*N*-propylamino and di-isopropylamino; examples of **(1-6C)alkylcarbonyl** include (1-4C)alkylcarbonyl such as methylcarbonyl, ethylcarbonyl, propylcarbonyl,

10 iso-propylcarbonyl and tert-butylcarbonyl; examples of **(1-6C)alkylcarbonyloxy** include (1-4C)alkylcarbonyloxy such as methylcarbonyloxy, ethylcarbonyloxy, propylcarbonyloxy, iso-propylcarbonyloxy and tert-butylcarbonyloxy; examples of **(1-6C)alkoxycarbonyl** (*N*-(1-6C)alkylcarbonyl) include (1-4C)alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, iso-propoxycarbonyl and

15 tert-butoxycarbonyl; examples of **(1-6C)alkoxycarbonylamino** include (1-4C)alkoxycarbonylamino such as methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, iso-propoxycarbonylamino and tert-butoxycarbonylamino; examples of **(1-6C)alkoxycarbonyl(N-methyl)amino** include (1-4C)alkoxycarbonyl(N-methyl)amino such as methoxycarbonyl(N-methyl)amino,

20 ethoxycarbonyl(N-methyl)amino, propoxycarbonyl(N-methyl)amino, iso-propoxycarbonyl(N-methyl)amino and tert-butoxycarbonyl(N-methyl)amino; examples of **(1-6C)alkylthio** include methylthio, ethylthio, propylthio, isopropylthio and butylthio; examples of **(1-6C)alkylsulfinyl** include methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl and butylsulfinyl; examples of **(1-6C)alkylsulfonyl** include

25 methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl and butylsulfonyl; examples of **(1-6C)alkoxysulfonyl** include methoxysulfonyl, ethoxysulfonyl, propoxysulfonyl, isopropoxysulfonyl and butoxysulfonyl; examples of **(1-6C)alkylcarbonylamino** include (1-4C)alkylcarbonylamino such as (1-3C)alkylCONH) (methylcarbonylamino, ethylcarbonylamino, propylcarbonylamino, iso-propylcarbonylamino) and tert-butylcarbonylamino; examples of

30 **(1-6C)alkylaminocarbonyl** include (1-4C)alkylaminocarbonyl such as methylaminocarbonyl, ethylaminocarbonyl, propylaminocarbonyl,

- iso-propylaminocarbonyl and tert-butylaminocarbonyl; examples of di(1-6C)alkylaminocarbonyl include di(1-4C)alkylaminocarbonyl such as dimethylaminocarbonyl, *N*-methyl-*N*-ethylaminocarbonyl, diethylaminocarbonyl, *N*-methyl-*N*-propylaminocarbonyl and di-isopropylaminocarbonyl; examples of
- 5 (1-6C)alkylaminocarbonyloxy include (1-4C)alkylaminocarbonyloxy such as methylaminocarbonyloxy, ethylaminocarbonyloxy, propylaminocarbonyloxy, iso-propylaminocarbonyloxy and tert-butylaminocarbonyloxy; examples of -S(O)_p(1-4C)alkyl (wherein p is 0, 1 or 2) include (1-6C)alkylthio, (1-6C)alkylsulfinyl and (1-6C)alkylsulfonyl; examples of (1-6C)alkylaminosulfonyl include
- 10 -SO₂NH(1-4C)alkyl such as methylaminosulfonyl, ethylaminosulfonyl, propylaminosulfonyl, *iso*-propylaminosulfonyl and *tert*-butylaminosulfonyl; examples of di(1-6C)alkylaminosulfonyl include di(1-4C)alkylaminosulfonyl such as dimethylaminosulfonyl, *N*-methyl-*N*-ethylaminosulfonyl, diethylaminosulfonyl, *N*-methyl-*N*-propylaminosulfonyl and di-isopropylaminosulfonyl; examples of
- 15 (1-6C)alkylsulfonylamino include (1-4C)alkylsulfonylamino such as methylsulfonylamino, ethylsulfonylamino, propylsulfonylamino, *iso*-propylsulfonylamino and *tert*-butylsulfonylamino.

Particular values of variable groups in compounds of formula (I) are as follows. Such values may be used where appropriate with any of the other values, definitions,

20 claims or embodiments defined hereinbefore or hereinafter.

In a particular embodiment, R¹ is an optionally substituted aryl group such as optionally substituted phenyl or naphthyl. R¹ as an optionally substituted aryl group may also be indanyl. It will be understood that when R¹ is a partially saturated aryl group, such as indanyl, it is the aromatic ring portion of R¹ which is directly attached to the linking nitrogen

25 atom.

Alternatively, R¹ is an optionally substituted heteroaryl group, and in particular is an optionally substituted monocyclic heteroaryl group such as pyridyl. Suitable values for R¹ as a heteroaryl ring include pyrimidinyl, pyridyl, pyrazolyl, pyrazinyl, thiazolyl, oxadiazolyl, isoxazolyl and thiadiazolyl.

30 It will be understood that when R¹ is a partially saturated bicyclic heteroaryl group, such as benzodioxolanyl, it is the aromatic ring portion of R¹ which is directly attached to the linking nitrogen atom.

Suitable values for R^1 as a bicyclic heteroaryl ring include pyrrolopyridyl, benzodioxolanyl, benzthiazolyl, benzimidazolyl and quinolyl.

More suitable values for R^1 include phenyl, naphthyl, indanyl, pyrimidinyl, pyridyl, pyrazolyl, pyrazinyl, thiazolyl, oxadiazolyl, isoxazolyl, thiadiazolyl, pyrrolopyridyl, 1,3-benzodioxan-5-yl, benzthiazolyl, benzimidazolyl and quinolyl.

In one embodiment, R^1 may not be pyrrolo(1,2-b)pyridazine.

Preferred optional substituents for R^1 include functional groups or (1-6C)alkyl groups such as methyl. Particular functional groups for substituents on R^1 include halo, $-C(O)_nR^{20}$ or $-OR^{20}$, where R^{20} is as defined above, and in particular is an aryl or aralkyl group.

Suitable functional groups as substituents on R^1 include halo, $-OR^{20}$ (wherein R^{20} is hydrogen, phenyl or (1-4C)alkyl, optionally substituted by one or more halo, such that for example R^{20} is difluoromethyl or trifluoromethyl, or optionally substituted by (1-4C)alkoxy), cyano, halo(1-4C)alkyl, $-S(O)_mR^{20}$ (wherein R^{20} is phenyl or (1-4C)alkyl, particularly methyl or ethyl, m is 0, 1 or 2, particularly 0 or 2), trifluoromethylthio, $-NR^{20}CONR^{21}R^{22}$ (wherein R^{20} , R^{21} and R^{22} are suitably all hydrogen), $-C(O)_nR^{20}$ (wherein n is 1 or 2, particularly 2 and R^{20} is (1-4C)alkyl or phenyl), $-OSO_2R^{20}$ (wherein R^{20} is suitably (1-4C)alkyl), $-SO_2NR^{21}R^{22}$ (wherein R^{21} and R^{22} are suitably both hydrogen), $-NR^{21}C(O)_nR^{20}$ (wherein n is 1 or 2, particularly 1, R^{21} is suitably hydrogen and R^{20} is suitably phenyl or (1-4C)alkyl), and $-CONR^{21}R^{22}$ (wherein R^{21} and R^{22} are suitably hydrogen).

Suitable values for Z^a include phenyl (optionally substituted by a functional group as hereinbefore defined, for example by $-CO_2Me$, or carboxy), benzyl, cyclohexyl, pyridyl, pyrimidinyl (optionally substituted by (1-4C)alkyl), triazolyl, morpholino, (2-4C)alkynyl (for example ethynyl) and (1-4C)alkyl (optionally substituted by a substituent selected from $-CO_2Me$, carboxy, methoxy, hydroxy and cyano).

Where R^1 is substituted by a group $-X^2-(CR^{52}R^{53})_w-Z^a$, suitably w is 0 or 1; Z^a is selected from the suitable values given above, particularly hydrocarbyl (such as optionally substituted alkyl, phenyl or benzyl) or pyridyl, and is more suitably optionally substituted phenyl; X^2 is suitably $-SO_2-$, $-CO-$, $NHCO-$, $-NH-$, $-O-$, and R^{52} and R^{53} are suitably both hydrogen.

In another aspect, preferred optional substituents on R^1 are 1, 2 or 3 substituents independently selected from alkyl (for example (1-6C)alkyl such as methyl or ethyl), halo, haloalkyl (such as halo(1-6C)alkyl, such as halomethyl, for example trifluoromethyl), haloalkoxy (such as halo(1-6C)alkoxy, such as halomethoxy, for example trifluoromethoxy) and cyano.

In another aspect, preferred optional substituents on R^1 are 1, 2 or 3 substituents independently selected from halo, halo(1-4C)alkyl, (1-4C)alkoxy, halo(1-4C)alkoxy, cyano, (1-4C)alkyl, (2-4C)alkynyl, $\text{Ph}(\text{CH}_2)_{0-1}\text{O}-$ (wherein the phenyl group is optionally substituted by halo), phenyl, benzoyl and anilino.

In another aspect, preferred optional substituents on R^1 are 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino.

In one aspect, R^1 is mono-substituted in the 3-position relative to the bond to NH; in another aspect R^1 is mono-substituted in the 4-position. In a further aspect R^1 is 2,4-disubstituted, 2,6-disubstituted, 3,4-disubstituted, 2,4-disubstituted, or 2,5-disubstituted by any of the possible substituents hereinbefore or hereinafter, but particularly those preferred optional substituents above, and more particularly di-halo, for example di-fluoro. In a further aspect R^1 is trisubstituted, for example 2,4,5-trisubstituted, such as 2,4,5-trihalo (for example 2,4,5-trifluoro).

When R^1 is di- or tri-substituted, the substituents are suitably independently selected from a functional group, Z^a and $-\text{X}^2-(\text{CR}^{52}\text{R}^{53})_w-\text{Z}^a$, for example any of those values given herein for these groups. For example, R^1 may be substituted by di-halo (such as difluoro, dichloro, mono-fluoro mono-chloro and mono-chloro mono-bromo), tri-halo (such as trifluoro), mono-halo mono-alkyl (such as mono-methyl, mono-chloro), mono-halo (such as fluoro or chloro) mono-trifluoromethyl, mono-alkyl (such as methyl) mono-cyano, di-methoxy, mono-chloro mono-methoxy, di-halo mono-hydroxy (such as 2-F, 4-Cl, 5-OH), or may be for example di-halo mono $-\text{O}-\text{Z}^a$ (such as $-\text{OCH}_2\text{CO}_2\text{Me}$). When R^1 is di-substituted, in one aspect at least one of the substituents is selected from halo, (1-4C)alkyl, (1-4C)alkoxy, trifluoromethyl and cyano. When R^1 is tri-substituted, in one aspect at least one, for example at least two, of the substituents are selected from halo, (1-4C)alkyl, (1-4C)alkoxy, trifluoromethyl and cyano.

- 13 -

Where R^1 is substituted by a group $-X^2-(CR^{52}R^{53})_w-Z^a$, a group $-X^2-(CR^{52}R^{53})_a-X^3-Z^a$ or a group $-(CR^{52}R^{53})_vX^3-Z^a$, R^{52} and R^{53} are suitably hydrogen.

When R^1 is substituted by Z^a , wherein Z^a is a heterocyclyl ring, such as a morpholino ring, preferably Z^a is not attached to the carbon atom of R^1 which is in an ortho position to the bond to the linking nitrogen atom.

In one embodiment, R^1 is selected from phenyl, naphthyl, indanyl, pyrimidinyl, pyridyl, pyrazolyl, pyrazinyl, thiazolyl, oxadiazolyl, isoxazolyl, thiadiazolyl, pyrrolopyridyl, 1,3-benzodioxan-5-yl, benzthiazolyl, benzimidazolyl and quinolyl; optionally substituted with 1, 2 or 3 substituents independently selected from halo, (1-4C)alkyl, ethynyl, (1-4C)alkoxy, hydroxy, (1-4C)alkoxy(1-4C)alkoxy, methoxymethyl, cyanomethyl, hydroxy(1-4C)alkyl, trifluoromethyl, difluoromethoxy, trifluoromethoxy, trifluoromethylthio, cyano, methylthio, methylsulfonyloxy, methylsulfonyl, ethylsulfonyl, aminocarbonylamino, methoxycarbonylamino, methylcarbonylamino, (1-4C)alkoxycarbonyl, methoxycarbonylmethoxy, benzyloxy, pyridylmethoxy, phenoxy (optionally substituted by methoxy or halo), phenyl (optionally substituted by methoxycarbonyl or carboxy), benzyl, anilino, anilincarbonyl, aminocarbonyl, benzoyl, benzoylamino, phenylsulfonyl, aminosulfonyl, cyclohexyl, methylpyrimidinyl, triazolyl and morpholino.

In a preferred embodiment, Y is a direct bond.

Where Y is a group $-X^6(CR^{40}R^{41})_t$, X^6 is suitably oxygen and t is preferably an integer of from 2 to 6.

Alternatively, Y is a group $(CH_2)_s$ or more preferably $-O(CH_2)_t-$ where s is an integer of from 1 to 6 and t is an integer of from 2 to 6, and in particular s or t are 3.

When R^2 is unsubstituted aryl or unsubstituted cycloalkyl, Y is preferably other than a direct bond.


R^2 is a suitably a substituted phenyl or a substituted heteroaryl group (for example any of those heteroaryl groups listed hereinbefore). Suitable examples of R^2 include phenyl, pyridyl, pyrimidinyl, indanyl, cyclohexyl, piperidinyl and benzthiazolyl.

When R^2 is an optionally substituted cycloalkyl group, it is preferably a monocyclic group such as (3-8C)cycloalkyl or (3-6C)cycloalkyl.

When R^2 is a substituted group, it is suitably substituted by at least one and optionally more than one substituent group -Z, a group $-X-(CR^{42}R^{43})_u-Z$, a group -

- $X-(CR^{42}R^{43})_n-X^1-Z$ or a group $-(CR^{42}R^{43})_n-X^1-Z$, where one or more further substituents may be selected from halo, cyano, nitro, amino, hydroxy or halo(1-6C)alkyl. Preferably R^2 is substituted by 1 or 2 groups independently selected from those defined hereinbefore or hereinafter, more preferably by 1 group. When R^2 is substituted by 2 groups, preferably one is a functional group as hereinbefore defined, such as halo, $-CO_2R^{20}$ (wherein R^{20} is hydrogen, (1-4C)alkyl or allyl) or cyano, or one substituent is (1-4C)alkyl.


Particular examples of groups Z or Z^a include groups of sub formula (x), (y) or (z).



- wherein each ring A or A' is independently selected from an optionally substituted heterocyclic ring, an optionally substituted cycloalkyl ring or an optionally substituted aryl ring, each R^{60} is an optionally substituted (1-6C)alkyl, an optionally substituted (2-6C)alkenyl or an optionally substituted (2-6C)alkynyl, and R^{61} is an optionally substituted (1-6C)alkylene, an optionally substituted (2-6C)alkenylene or an optionally substituted (2-6C)alkynylene.


- Suitably optional substituents for groups A , A' , R^{60} and R^{61} are functional groups.

A further particular example of groups Z or Z^a includes groups of sub formula (zz), wherein A , R^{60} and R^{61} , and suitable optional substituents therein are as defined above for sub formulae (x), (y) and (z).



- In a particularly preferred embodiment, Z is a group of sub-formula (x) above. In one embodiment, R^2 is a 5- or 6-membered aromatic ring of sub-structure (a):

- 15 -



(a)

Z^1 , Z^2 , Z^3 and Z^4 are independently selected from $-\text{CH}-$, $-\text{CR}^6-$ or a heteroatom selected from $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^{50})_r-$, where r is 0 or 1 depending upon the requirements of the aromatic ring, and R^{50} is hydrogen or (1-6C)alkyl, and Z^4 may additionally be a direct bond;

R^4 is a group $-\text{Z}$, a group $-\text{X}-(\text{CR}^{42}\text{R}^{43})_u-\text{Z}$, a group $-\text{X}-(\text{CR}^{42}\text{R}^{43})_v-\text{X}^1-\text{Z}$ or a group $-(\text{CR}^{42}\text{R}^{43})_v\text{X}^1-\text{Z}$, wherein Z , X , X^1 , R^{42} , R^{43} , u and v are as defined above;

each R^6 is independently selected from halo, cyano, nitro, amino, hydroxy, halo C_{1-6} alkyl, a group $-\text{Z}$, a group $-\text{X}-(\text{CR}^{42}\text{R}^{43})_u-\text{Z}$, a group $-\text{X}-(\text{CR}^{42}\text{R}^{43})_v-\text{X}^1-\text{Z}$ or a group $-(\text{CR}^{42}\text{R}^{43})_v\text{X}^1-\text{Z}$, wherein Z , X , X^1 , R^{42} , R^{43} , u and v are as defined above.

Suitably, when Z^4 is a direct bond, one of Z^1 or Z^2 is a heteroatom, in particular oxygen or sulphur.

Preferably Z^4 is other than a direct bond.

Suitably in this case, Z^2 and Z^3 are independently selected from $-\text{CH}-$, $-\text{CR}^{24}-$ or a nitrogen atom.

Suitably Z^1 is a $-\text{CH}-$ group.


Preferred groups R^6 are listed below as R^{6a} .

Suitably, Z^1 , Z^2 , Z^3 and Z^4 are $-\text{CH}-$.

Suitably R^{42} and R^{43} are hydrogen.

Where one of Z^1 to Z^4 is $\text{N}(\text{R}^{50})_r$, preferably it is Z^2 or Z^3 . Where one of Z^1 to Z^4 is $-\text{CR}^6$, preferably it is Z^2 or Z^3 .

In an alternative embodiment, R^2 is a cycloalkyl group such as cyclohexyl of sub-formula (b)



(b)

where R^4 is as defined above, and R^a , R^b , R^c and R^d are independently selected from hydrogen or a group R^6 as defined above.

In yet a further embodiment, R^2 is a bicyclic ring, which may be a bicyclic aryl ring or a bicyclic heterocyclic ring. For instance, R^2 comprises fused 6,6-membered rings, or fused 5,6-membered rings, one or both of said rings may be unsaturated. Examples of such rings include benzimidazole (preferably linked to the group-Y-NH- by way of the benzene ring), indanyl, indenyl. Particularly suitable bicyclic rings are partially unsaturated, such that the ring linked to the group-Y-NH- is saturated and this is fused to an aromatic ring. Particular examples of such rings are indanyl rings, such as 2-indanyl. In one embodiment, R^2 may not be pyrrolo(1,2-b)pyridazine.

In particular, R^4 is a group Z.

Suitably Z is an aryl, heterocyclyl or cycloalkyl group, any of which are optionally substituted by a functional group or an (1-6C)alkyl, (2-6C)alkenyl or (2-6C)alkynyl group.

Preferably Z is substituted by a functional group or by a (1-6C)alkyl group which is substituted by a functional group. Particular examples of such functional groups include $-C(O)_2R^{20}$ or a carboxylic acid mimic or bioisostere thereof, $-C(O)NR^{21}R^{22}$ and $-NR^{21}C(O)_nR^{20}$, where R^{20} , R^{21} and R^{22} are as defined above.

In another embodiment, R^2 is substituted by Z and Z is an optionally substituted heterocyclyl group. Suitable examples of Z as an optionally substituted heterocyclyl group include any of the suitable values for heterocyclyl given hereinbefore and in particular include pyrrolidinyl, piperazinyl, piperidinyl, pyridyl, morpholino, thiomorpholino, homopiperazinyl, thiadiazolyl, (oxo)pyridazinyl and (oxo)thiadiazolyl.

In another embodiment, R^2 is substituted by Z and Z is an optionally substituted hydrocarbyl group. Suitable examples of Z as an optionally substituted hydrocarbyl group include (all optionally substituted) (1-6C)alkyl (such as (1-4C)alkyl), phenyl, cycloalkyl (such as adamantyl, cyclobutyl, cyclopentyl and cyclohexyl), cycloalkyl combined with (1-4C)alkyl (such as methylcyclohexyl, ethylcyclohexyl, isopropylcyclohexyl, cyclohexylmethyl, ethylcyclobutyl, cyclobutylmethyl and methylcyclopentyl) and phenyl combined with (1-4C)alkyl (such as benzyl and methylphenyl (such as tolyl)).

In another embodiment, R^2 is substituted by Z and Z is an optionally substituted combination of hydrocarbyl and heterocyclyl groups. Suitable examples of Z as an optionally substituted combination of hydrocarbyl and heterocyclyl groups include non

aromatic heterocycles such as piperazinyl or piperidyl substituted by (1-4C)alkyl (for example methyl, ethyl and isopropyl), benzyl or cycloalkyl(1-4C)alkyl (for example cyclopropylmethyl); oxidised heterocycles such as oxopyridazine or oxothiadiazine substituted by one or two (1-4C)alkyl (such as methyl); aromatic heterocycles (such as
5 pyridyl) substituted by one or two (1-4C)alkyl (such as methyl). For example pyridylmethyl (wherein the methyl may be further substituted by a functional group such as carboxy), benzylpiperazinyl, (methyl)oxopyridazinyl, (methyl)oxothiadiazolyl, (optionally carboxy substituted)methylpiperidyl, (optionally carboxy substituted)methylpiperidylmethyl, (optionally carboxy substituted)dimethylpiperidyl,
10 (optionally carboxy substituted)ethylpiperidyl and (cyclopropylmethyl)piperazinyl.

In another embodiment R^2 is substituted by Z and Z is an optionally substituted combination of two heterocyclyl groups, for example pyridyl in combination with piperazinyl.

Suitable substituents on a group Z include halo, hydroxy, carboxy, $-CO_nR^{20}$ [wherein
15 R^{20} is hydrogen, optionally substituted hydrocarbyl (such as (1-4C)alkyl, benzyl, phenyl, methylphenyl, phenethyl) or optionally substituted heterocyclyl (such as pyridyl) and wherein n is 1 or 2], $-CONH_2$, $-CONHR^{21}$ (wherein R^{21} is selected from hydrogen, alkyl and benzyl), cyano, amino, $-NHCO_2(1-4C)alkyl$, and $-CONR^{21}R^{22}$ (wherein $NR^{21}R^{22}$ forms an optionally substituted heterocyclyl ring).

20 Suitably a ring formed by $NR^{21}R^{22}$ contains 0 or 1 further heteroatom selected from O, N and S and may be for example piperidiny, piperazinyl, pyrrolidiny or morpholino. A ring formed by $NR^{21}R^{22}$ may also be fused to another ring, for example thereby comprise a pyrrolidiny ring fused with dioxolan.

In general, suitably R^{20} is hydrogen or is selected from (all optionally substituted) (1-
25 4C)alkyl, phenyl, pyridyl, benzyl, phenethyl, methylphenyl and allyl.

In general, R^{21} and R^{22} are suitably are each independently hydrogen or are selected from (optionally substituted) phenyl, (1-4C)alkyl, and benzyl.

Suitably R^{20} , R^{21} and R^{22} (and rings formed by $NR^{21}R^{22}$) are unsubstituted or are substituted by 1 or 2 substituents. Suitable optional substituents for R^{20} , R^{21} and R^{22} include
30 halo, cyano, hydroxy, (1-4C)alkoxy, carboxy and $-CO_2(1-4C)alkyl$. A particular substituent for R^{21} and R^{22} is hydroxy. Particular substituents for rings formed by $NR^{21}R^{22}$ are hydroxy, carboxy and $-CO_2(1-4C)alkyl$.

In another embodiment R^2 is substituted by $-X-(CR^{42}R^{43})_uZ$, wherein X is preferably O, -NH-, -NMe-, or -SO₂NH-, u is 0, 1 or 2, R^{42} and R^{43} are each hydrogen and Z is selected from any of the values mentioned hereinbefore, particularly morpholino or optionally substituted phenyl (such as methoxyphenyl) or methylphenyl.

- 5 In another embodiment, R^2 is substituted only by a functional group as hereinbefore defined. In particular, the functional group may be selected from (1-4C)alkoxy, (1-4C)alkylthio and (1-4C)alkylsulfonyl, wherein the aforementioned (1-4C)alkyl groups may optionally be substituted by carboxy or (1-4C)alkoxycarbonyl.


In one aspect, R^2-Y is selected from:

- 10 4-(piperazino)phenyl (substituted on the available piperazine nitrogen by a substituent selected from (1-4C)alkyl, acyl, benzyl, ethoxycarbonyl, tert-butoxycarbonyl, pyridyl, cyclopropylmethyl and methoxyethyl),
4-(piperazino)-3-methyl-phenyl (substituted on the available piperazine nitrogen by a substituent selected from acyl and tert-butoxycarbonyl),
15 4-(piperazino)pyridyl (substituted on the available piperazine nitrogen by a substituent selected from (1-4C)alkyl, acyl, ethoxycarbonyl, tert-butoxycarbonyl, benzoyl, fluorobenzoyl, cyanobenzoyl, methylbenzoyl, pyridylcarbonyl, methylsulfonyl, methoxymethylcarbonyl, phenethylcarbonyl, anilino carbonyl, benzylaminocarbonyl, phenoxycarbonyl, benzyloxycarbonyl and aminocarbonyl),
20 4-piperidinophenyl, benzyloxyphenyl, (1-4C)alkylphenyl, pyridylmethoxyphenyl, N-methylanilinophenyl, biphenyl, (chloro)biphenyl, (methoxycarbonyl)biphenyl, (ethoxycarbonylmethyl)biphenyl, (carboxy)biphenyl, morpholinophenyl, dimethylmorpholinophenyl, (morpholino)(chloro)phenyl, (morpholino)(dichloro)phenyl, (morpholino)(bromo)phenyl, (morpholino)(methyl)phenyl, (morpholino)(fluoro)phenyl,
25 (morpholino)(cyano)phenyl, (morpholino)(allyloxycarbonyl)phenyl, (morpholinoethoxy)phenyl, morpholinopyridyl, (morpholino)(chloro)pyridyl, (morpholino)(fluoro)pyridyl, phenoxypyridyl, anilinopyridyl, methoxypyridyl, butoxypyridyl, pyridyloxypyridyl, hydroxybutylpyridyl, thiomorpholinopyridyl, phenoxypropyl, (fluoro)(trifluoromethyl)phenyl, (N-acyl)homopiperazinophenyl, (N-acyl)piperazinopyrimidinyl, methylbenzthiazolyl, butylthiophenyl,
30 (methoxythiadiazolyl)aminosulfonylphenyl, (difluorophenoxy)phenyl, (dimethyl-oxo-pyridazinyl)phenyl, (methyl-oxo-thiadiazinyl)phenyl, methoxycarbonylpropylphenyl,


- (methoxycarbonyl)methylthiophenyl, (methoxycarbonyl)adamantyl, carboxyadamantyl,
 (methoxycarbonylmethyl)bi-cyclohexyl, (carboxy)bi-cyclohexyl,
 (ethoxycarbonyl)ethylindanyl, (carboxyethyl)indanyl, (methoxycarbonyl)phenoxypropyl,
 (carboxy)phenoxypropyl, (carboxypropyl)phenyl, dicarboxybutylphenyl,
 5 carboxybutylphenyl, propylsulfonylphenyl, (carboxy)isopropylsulfonylphenyl,
 phenoxyphenyl, (methoxyphenoxy)phenyl, methylphenoxyphenyl, hydroxypropoxyphenyl,
 ethoxyethoxyphenyl, (dimethylaminocarbonyl)isopropoxyphenyl,
 (carboxymethyl)piperidinophenyl, (methoxycarbonylmethyl)piperidinopyridyl,
 (carboxymethyl)piperidinopyridyl, (carboxy)piperidinopyridyl,
 10 (methoxycarbonyl)piperidinopyridyl, (carboxy)(methyl)piperidinopyridyl,
 (methoxycarbonyl)(methyl)piperidinopyridyl, (carboxymethyl)(methyl)piperidinopyridyl,
 (methoxycarbonyl)piperidyl[(methyl)pyridyl], (carboxy)piperidyl[(methyl)pyridyl],
 (methoxycarbonylmethyl)piperidyl[(methyl)pyridyl],
 (carboxymethyl)piperidyl[(methyl)pyridyl], (carboxyethyl)piperidyl[(methyl)pyridyl],
 15 (ethoxycarbonyl)ethylpiperidyl[(methyl)pyridyl],
 [(methoxycarbonyl)piperidinocarbonylmethyl]cyclohexylphenyl,
 [(carboxy)piperidinocarbonylmethyl]cyclohexylphenyl,
 [(methoxycarbonyl)isopropylaminocarbonylmethyl]cyclohexylphenyl,
 [(carboxy)isopropylaminocarbonylmethyl]cyclohexylphenyl, ~
 20 [(methoxycarbonyl)pyrrolidinylcarbonylmethyl]cyclohexylphenyl,
 [(carboxy)pyrrolidinylcarbonylmethyl]cyclohexylphenyl,
 [(methoxycarbonyl)(hydroxy)pyrrolidinylcarbonylmethyl]cyclohexylphenyl,
 [(carboxy)(hydroxy)pyrrolidinylcarbonylmethyl]cyclohexylphenyl,
 (hydroxyisobutylaminocarbonylmethyl)cyclohexylphenyl,
 25 [(tetrahydrodioxolopyrrolyl)carbonyl]methylcyclohexylphenyl,
 (methoxycarbonylpyridyl)piperidino, (carboxypyridyl)piperidino,
 (aminocarbonylpyridyl)piperidino, (carboxymethylpyridyl)piperidino,
 (ethoxycarbonylmethyl)cyclohexylphenyl, (methoxycarbonylmethyl)cyclohexylphenyl,
 cyclohexylphenyl, (carboxymethyl)cyclohexylphenyl, (hydroxyethyl)cyclohexylphenyl,
 30 (aminocarbonylmethyl)cyclohexylphenyl, (cyanomethyl)cyclohexylphenyl,
 carboxycyclohexylphenyl, (dimethylaminocarbonylmethyl)cyclohexylphenyl, [(N-
 hydroxyethyl-N-methylaminocarbonylmethyl]cyclohexylphenyl, [N-

- (dihydroxypropyl)aminocarbonylmethylcyclohexylphenyl,
 (aminomethyl)cyclohexylphenyl, (tert-butoxycarbonylaminomethyl)cyclohexylphenyl,
 (methoxycarbonyl)cyclohexyloxyphenyl, (ethoxycarbonyl)cyclohexyloxyphenyl,
 carboxycyclohexyloxyphenyl, (methoxycarbonyl)cyclopentylphenyl,
 5 (ethoxycarbonylmethyl)cyclopentylphenyl, (carboxy)cyclopentylphenyl
 (carboxymethyl)cyclopentylphenyl, (methoxycarbonyl)cyclobutylphenyl,
 (ethoxycarbonyl)cyclobutylmethylphenyl, (carboxy)cyclobutylphenyl,
 (carboxy)cyclobutylmethylphenyl, (carboxyethyl)cyclobutylphenyl,
 (ethoxycarbonyl)ethylcyclobutylphenyl, (carboxymethyl)phenylcyclohexyl, 1-
 10 (ethoxycarbonyl)ethylcyclohexylphenyl, 1-carboxyethylcyclohexylphenyl, (2-
 methoxycarbonyl)isopropylcyclohexyl, (2-carboxy)isopropylcyclohexyl,
 pyrrolidinylphenyl, (methoxycarbonylmethyl)cyclohexylpyridyl and
 (carboxymethyl)cyclohexylpyridyl.


- 15 As used herein, the reference to carboxylic acid mimic or bioisostere includes groups as defined in The Practice of Medicinal Chemistry, Wermuth C.G. Ed.: Academic Press: New York, 1996, p203. Particular examples of such groups include $-\text{SO}_3\text{H}$, $\text{S}(\text{O})_2\text{NHR}^{13}$, $\text{S}(\text{O})_2\text{NHC}(\text{O})\text{R}^{13}$, $-\text{CH}_2\text{S}(\text{O})_2\text{R}^{13}$, $-\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}^{13}$, $-\text{C}(\text{O})\text{NHOH}$, $-\text{C}(\text{O})\text{NHCN}$, $-\text{CH}(\text{CF}_3)\text{OH}$, $\text{C}(\text{CF}_3)_2\text{OH}$, $-\text{P}(\text{O})(\text{OH})_2$ and groups of sub-formula (a)-(i') below




(a)




(b)




(c)




(d)




(e)




(f)




(g)




(h)




(i)




(j)




(k)




(l)




(m)




(n)




(o)




(p)




(q)




(r)




(s)




(t)




(u)




(v)




(w)




(x)




(y)




(z)




(a')




(b')




(c')



(d')



(e')




where p is 1 or 2, R^{27} and R^{28} are independently selected from hydrogen, hydroxy, (1-6C)alkoxy, thiol, (1-6C)alkylthio, $-C(O)R^{29}$, $-S(O)R^{30}$, $-SO_2R^{31}$, $-NR^{32}R^{33}$, $-NHCN$, halogen and trihalomethyl, where R^{29} , R^{30} and R^{31} are $-OR^{34}$, (1-6C)alkyl, $-NR^{32}R^{33}$ or

5 trihalomethyl,

R^{32} and R^{33} are independently selected from hydrogen, (1-6C)alkyl, $-SO_2R^{34}$ and $-COR^{35}$, where R^{35} is (1-6C)alkyl or trihalomethyl, and R^{34} is hydrogen, (1-6C)alkyl or trihalomethyl.

Particular examples of R^{27} or R^{28} are hydroxy.

10 In a further aspect of the invention, there is provided a compound of formula (IZA), or a pharmaceutically-acceptable salt or pro-drug thereof,



(IZA)

wherein R^1 is selected from phenyl (optionally substituted with 1, 2 or 3 substituents
15 independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino), 2-pyridyl (optionally substituted by chlorophenoxy, chlorobenzyloxy or methoxyphenoxy, and/or substituted with a substituent selected from halo, trifluoromethyl, (1-4C)alkyl, (1-4C)alkoxy
20 and cyano), 3-pyridyl (optionally substituted as for 2-pyridyl), halopyrimidinyl and trifluoromethylthiazolyl;

Z^2 is N or CH;

R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen or methyl;

R^{6ZA} is hydrogen, fluoro, chloro or methyl;

A is N or CH;

X^{ZA} is a direct bond, $-CH_2-$ or $-O-$ (except when A is N);

5 m is 0, 1 or 2;

n is 0 or 1;

provided that $m + n = 0, 1$ or 2;

p is 0 or 1.

10 In another aspect of this embodiment, in the compound of formula (IZA), R^1 is phenyl, optionally substituted as defined above.

In a further aspect of this embodiment, in the compound of formula (IZA), R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl.

15 In a further aspect of this embodiment, in the compound of formula (IZA), R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

Z^2 is CH;

R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen;

20 R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

A is CH;

X^{ZA} is a direct bond or $-CH_2-$;

m is 0, 1 or 2;

n is 0 or 1;

25 provided that $m + n = 0, 1$ or 2;

p is 0 or 1.

In a further aspect of this embodiment, in the compound of formula (IZA), R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

30 Z^2 is N;

R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen;

R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

A is CH;

X^{ZA} is a direct bond or $-CH_2-$;

m is 0, 1 or 2;

5 n is 0 or 1;

provided that $m + n = 0, 1$ or 2;

p is 0 or 1.

In a further aspect of this embodiment, in the compound of formula (IZA),

10 R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

Z^2 is CH;

R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen;

R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

15 A is N;

X^{ZA} is a direct bond or $-CH_2-$;

m is 0, 1 or 2;

n is 0 or 1;

provided that $m + n = 0, 1$ or 2;

20 p is 0 or 1.

In a further aspect of this embodiment, in the compound of formula (IZA),

R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

Z^2 is N;

25 R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen;

R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

A is N;

X^{ZA} is a direct bond or $-CH_2-$;

30 m is 0, 1 or 2;

n is 0 or 1;

provided that $m + n = 0, 1$ or 2;



p is 0 or 1.

In a further aspect of this embodiment, in the compound of formula (IZA),

R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

5 Z^2 is N;

R^{ZA3} is methyl;

R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

A is N;

X^{ZA} is a direct bond or $-CH_2-$;

10 m is 1;

n is 1;

p is 0.

In a further aspect of this embodiment, in the compound of formula (IZA),

R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

15 Z^2 is CH;

Z^2 is CH;

R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen;

R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

20 A is CH;

X^{ZA} is $-O-$;

m is 0, 1 or 2;

n is 0 or 1;

provided that $m + n = 0, 1$ or 2;

25 p is 0 or 1.

In a further aspect of this embodiment, in the compound of formula (IZA),

R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl;

Z^2 is N;

30 R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;

R^{ZA3} is hydrogen;

R^{6ZA} is hydrogen, chloro, fluoro or methyl, particularly hydrogen;

A is CH;

X^{ZA} is -O-;

m is 0, 1 or 2;

n is 0 or 1;


5 provided that $m + n = 0, 1$ or 2;

p is 0 or 1.

In one aspect, for a compound of formula (IZA), wherein A is -CH-, the substituents on the ring containing A (ie the X^{ZA} -pyridyl/phenyl group and the carboxy(alkyl) group) are trans relative to each other.

10 In one aspect, for a compound of formula (IZA), wherein A is -CH-, the substituents on the ring containing A (ie the X^{ZA} -pyridyl/phenyl group and the carboxy(alkyl) group) are cis relative to each other.

In a further aspect of the invention, there is provided a compound of formula (IZB), or a pharmaceutically-acceptable salt or pro-drug thereof,



(IZB)

15 wherein R^1 is selected from phenyl (optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino), 2-pyridyl (optionally substituted by chlorophenoxy, chlorobenzyloxy or methoxyphenoxy, and/or substituted with a substituent selected from halo, trifluoromethyl, (1-4C)alkyl, (1-4C)alkoxy and cyano), 3-pyridyl (optionally substituted as for 2-pyridyl), halopyrimidinyl and trifluoromethylthiazolyl;

25 Z^2 is N or CH;

R^{6ZB} is hydrogen, fluoro, chloro or methyl;


X^{ZB} is O or S, particularly O.

34

In another aspect of this embodiment, in the compound of formula (IZB), R^1 is phenyl, optionally substituted as defined above.

In a further aspect of this embodiment, in the compound of formula (IZB), R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from
 5 fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl.

In a further aspect of the invention, there is provided a compound of formula (IZC), or a pharmaceutically-acceptable salt or pro-drug thereof,



(IZC)

10 wherein R^1 is selected from phenyl (optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino), 2-pyridyl (optionally substituted by chlorophenoxy, chlorobenzyloxy or methoxyphenoxy, and/or
 15 substituted with a substituent selected from halo, trifluoromethyl, (1-4C)alkyl, (1-4C)alkoxy and cyano), 3-pyridyl (optionally substituted as for 2-pyridyl), halopyrimidinyl and trifluoromethylthiazolyl;

Z^2 is N or CH;

R^{6ZC} is hydrogen, fluoro, chloro or methyl;


20 R^{ZC} is selected from (1-4C)alkyl, acyl, benzyl, ethoxycarbonyl, tert-butoxycarbonyl, pyridyl, cyclopropylmethyl, methoxyethyl, benzoyl, fluorobenzoyl, cyanobenzoyl, methylbenzoyl, pyridylcarbonyl, methylsulfonyl, methoxymethylcarbonyl, phenethylcarbonyl, anilino carbonyl, benzylaminocarbonyl, phenoxycarbonyl, benzyloxycarbonyl and aminocarbonyl,

25 In another aspect of this embodiment, in the compound of formula (IZC), R^1 is phenyl, optionally substituted as defined above.

In a further aspect of this embodiment, in the compound of formula (IZC), R^1 is phenyl, optionally substituted with 1, 2 or 3 substituents independently selected from

fluoro, chloro, trifluoromethyl, trifluoromethoxy, cyano, methyl and ethyl.

In particular, the present invention provides a compound of formula (IA) or a pharmaceutically-acceptable salt or prodrug thereof,



(IA)

wherein:


- R^{1a} is selected from $AR1^a$, $HET-1^a$ and $HET-2^a$; wherein R^{1a} is optionally substituted by 1, 2, 3 or 4 groups independently selected from R^3 ;
- R^3 is selected from halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $R^{8a}NHCO-$, $R^{8a}CONH$, $R^{8a}OCONH-$, $R^{8a}OCON(Me)-$, $R^{8a}(1-6C)alkyl-$, halo(1-6C)alkyl, halo(1-6C)alkoxy, (1-6C)alkylthio, di(1-6C)alkylaminocarbonyl, (3-8C)cycloalkyloxy(1-6C)alkyl, (3-8C)cycloalkyl(1-6C)alkoxy, (3-8C)cycloalkyl(1-6C)alkoxy(1-6C)alkyl, phenyl, $HET-1^b$, $AR1^b$ -(1-6C)alkoxy, $HET-1^b$ -(1-6C)alkoxy, $HET-3^b$ -(1-6C)alkoxy, $AR1^b$ -oxy, $HET-1^b$ -oxy, $HET-3^b$ -oxy, $AR1^b$ -oxy(1-6C)alkyl, $HET-1^b$ -oxy(1-6C)alkyl, $HET-3^b$ -oxy(1-6C)alkyl, $AR1^b$ -(1-6C)alkoxy(1-6C)alkyl, $HET-1^b$ -(1-6C)alkoxy(1-6C)alkyl, and $HET-3^b$ -(1-6C)alkoxy(1-6C)alkyl;
- wherein any phenyl, $AR1^b$, $HET-1^b$, $HET-3^b$ or cycloalkyl ring in any of the above values for R^3 may optionally be substituted by up to 4 substituents independently selected from R^{7a} ;
- and wherein any R^{8a} , (3-8C)cycloalkyl, or (1-6C)alkyl group in any of the above definitions for R^3 may optionally be substituted by 1, 2 or 3 substituents independently selected from halo, (1-6C)alkyl, (1-6C)alkoxy, (2-6C)alkenyl, (2-6C)alkynyl, hydroxy, cyano, carboxy, aminocarbonyl, halo C_{1-6} alkyl, amino, (1-6C)alkylamino, di(1-6C)alkylamino, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, aminosulfonyl, (1-6C)alkylaminosulfonyl, di(1-6C)alkylaminosulfonyl, (1-6C)alkylsulfonylamino, aminocarbonyl, (1-6C)alkylaminocarbonyl, di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino, (1-6C)alkoxycarbonylamino,

(1-6C)alkoxycarbonyl(N-methyl)amino and (1-6C)alkylaminocarbonyloxy;

Y is a direct bond, or a group $(CH_2)_s$ or $-O(CH_2)_t-$ where s is an integer of from 1 to 6 and t is an integer of from 2 to 6, provided that the oxygen atom of the group $-O(CH_2)_t-$ is attached to the R^{2a} group;


5 R^{2a} is:

a) a 6-membered ring selected from (i), (ii) or (iii)



10 wherein R^{36a} , R^{36b} , R^{36c} and R^{36d} are independently selected from hydrogen or a group R^{6a} as defined hereinafter; or

b) a 5-membered heteroaryl ring of sub-formula (iv)



15 wherein Z^5 , Z^6 and Z^7 are independently selected from a heteroatom (O, N or S) or CR^{37} , where each R^{37} is hydrogen or a group R^{6b} as defined hereinafter, and provided that at least one of Z^5 , Z^6 and Z^7 is a heteroatom and there are no O-O, S-S or O-S bonds); or

c) a group HET-2^b which is optionally substituted by 1 or 2 groups independently selected from R^4 or R^6 ;

20 d) an optionally substituted cycloalkyl group or an optionally substituted bicyclic aryl group (such as indanyl) wherein optional substituents are selected from groups R^4 and R^{6a} as hereinafter defined;

R^4 is selected from:

a) phenyl;

25 b) HET-3^c (wherein any available nitrogen atom is substituted by R^5 provided it is not thereby quaternised;

- c) HET-1^c;
- d) phenoxy, HET-1^c-oxy, R^{8b}O-, R^{8b}S(O)_q-, phenyl(1-6C)alkoxy and HET-1^c-(1-6C)alkoxy, where q is 0, 1 or 2;
- e) R^{8b}, R^{8b}O(1-6C)alkyl, amino(1-6C)alkyl, hydroxy(1-6C)alkyl, phenoxy(1-6C)alkyl,
- 5 HET-1^c-oxy(1-6C)alkyl; and
- f) phenylamino, phenyl(1-6C)alkylamino, HET-1^c-amino, HET-1^c-(1-6C)alkylamino, (1-6C)alkylamino, N-phenyl-N-(1-6C)alkylamino, N-phenyl(1-6C)alkyl-N-(1-6C)alkylamino, N-(HET-1^c)-N-(1-6C)alkylamino, N-(HET-1^c-(1-6C)alkyl)-N-(1-6C)alkylamino, and
- 10 di(1-6C)alkylamino;
- wherein any alkyl, R^{8b}, phenyl (including phenoxy), HET-1^c or HET-3^c ring in any of the definitions a) to f) for R⁴ may optionally be substituted by 1, 2, 3 or 4 substituents independently selected from R^{7b};
- R⁵ is selected from a group R^{8c}, phenyl, HET-1^d, R^{8c}CO-, R^{8c}OC(O)-,
- 15 phenylcarbonyl, HET-1^d-carbonyl, (1-6C)alkoxycarbonyl, H₂NCO, R^{8c}NHCO-, phenylNHCO-, (1-6C)alkoxysulfonyl, (1-6C)alkylaminosulfonyl, (1-6C)alkylsulfinyl and (1-6C)alkylsulfonyl;
- wherein any (1-6C)alkyl or (3-8C)cycloalkyl groups in any of the definitions of R⁵ including R^{8c} may optionally be substituted by a substituent selected from a group R¹⁵,
- 20 amino, (1-6C)alkylamino, hydroxy, carboxy, (2-6C)alkenyl, (2-6C)alkynyl, phenyl (optionally substituted with up to four groups selected from R^{15a} and HET-1^f;
- and wherein any phenyl or HET-1^d groups in any of the definitions of R⁵ are optionally substituted by up to four groups selected from R^{15b},
- where R¹⁵, R^{15a} and R^{15b} are independently selected from halo, cyano, C₁₋₆alkoxy,
- 25 haloC₁₋₆alkyl, haloC₁₋₆alkoxy, (1-6C)alkylS(O)p'- (wherein p' is 0, 1 or 2),
- R^{6a} and R^{6b} are independently selected from R^{8d}, R^{8d}O-, R^{8d}OCO-, R^{8d}COO-, halo, cyano, halo(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkoxy, (1-6C)alkylcarbonyl, (1-6C)alkylS(O)p- (wherein p is 0, 1 or 2), (1-6C)alkylaminocarbonyl, di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino,
- 30 (1-6C)alkylaminosulfonyl,
- di(1-6C)alkylaminosulfonyl, (1-6C)alkylsulfonylamino, AR1^c-(1-6C)alkoxy,

HET-1^e-(1-6C)alkoxy, HET-3^e-(1-6C)alkoxy, AR1^e-oxy(1-6C)alkyl,
HET-1^e-oxy(1-6C)alkyl, HET-3^e-oxy(1-6C)alkyl, AR1^e-(1-6C)alkoxy(1-6C)alkyl,
HET-1^e-(1-6C)alkoxy(1-6C)alkyl, and HET-3^e-(1-6C)alkoxy(1-6C)alkyl;

wherein any alkyl, R^{8d}, phenyl, AR1^e, HET-1^e, HET-3^e or cycloalkyl ring in any of the
5 above values for R^{6a} or R^{6b} may optionally be substituted by 1, 2 or 3 substituents
independently selected from R^{7c},

each R^{7a}, R^{7b} and R^{7c} is independently selected from R^{8e}, R^{8e}O(CR¹³R¹⁴)_{n'}-,
R^{8e}OCO(CR¹³R¹⁴)_{n'}-, R^{8e}COO(CR¹³R¹⁴)_{n'}-, HCOO(CR¹³R¹⁴)_{n'}-, R^{8e}CO(CR¹³R¹⁴)_{n'}-,
HCO(CR¹³R¹⁴)_{n'}-, R^{8e}NHCO(CR¹³R¹⁴)_{n'}-, H₂NCO(CR¹³R¹⁴)_{n'}-, (R¹⁰R¹¹)NCO(CH₂)_{n'},
10 R^{8e}CONH(CR¹³R¹⁴)_{n'}-, HCONH(CR¹³R¹⁴)_{n'}-, HCONR^{8e}(CR¹³R¹⁴)_{n'}-, R^{8e}CONR^{8f}(CR¹³R¹⁴)_{n'}-,
R^{8e}OCONH(CR¹³R¹⁴)_{n'}-, HCONR^{8e}(CR¹³R¹⁴)_{n'}-, R^{8e}OCON(R^{8e})(CR¹³R¹⁴)_{n'}-,
ROCON(H)(CR¹³R¹⁴)_{n'}-, HOCON(R^{8e})(CR¹³R¹⁴)_{n'}-, R^{8e}-(1-6C)alkyl-, halo, cyano, R¹²,
R¹²-(1-6C)alkyl, hydroxy, oxo, aminocarbonyl, halo(1-6C)alkyl, halo(1-6C)alkoxy,
R^{8e}S(O)_{p''}- (wherein p'' is 0, 1 or 2), amino, amino(1-6C)alkyl, R^{8e}NH(1-6C)alkyl, (R^{8e}R^{8f}
15)N(1-6C)alkyl, R^{8e}NHSO₂-, (R¹⁰R¹¹)NSO₂-, and R^{8e}SO₂NH-, where n' is 0 or an integer of
from 1 to 6;

R¹² is carboxy or a carboxylic acid mimic or bioisostere of carboxy;

R¹³ is selected from hydrogen, (1-6C)alkyl, hydroxy, halo, amino, cyano, ((1-
3C)alkyl)CONH-, carboxy, (1-6C)alkoxy, (1-6C)alkoxycarbonyl, carbamoyl, N-((1-
20 (6C)alkyl)carbamoyl, halo((1-6C)alkyl) (such as trifluoromethyl), (1-6C)alkylsulphonyl or
(1-6C)alkylsulphinyl;

R¹⁴ is hydrogen, (1-6C)alkyl or halo(1-6C)alkyl;

and wherein any alkyl group or R^{8e}, R^{8f}, R¹⁰ or R¹¹ group within a group R^{7a}, R^{7b}
or R^{7c} is optionally substituted by 1, 2 or 3 groups independently selected from R⁹;

25 each R^{8a}, R^{8b}, R^{8c}, R^{8d}, R^{8e} and R^{8f} is independently selected from (1-6C)alkyl,
(2-6C)alkenyl, (2-6C)alkynyl and (3-8C)cycloalkyl;

R¹⁰ and R¹¹ are independently selected from selected from (1-6C)alkyl,
(2-6C)alkenyl, (2-6C)alkynyl and (3-8C)cycloalkyl; or R¹⁰ and R¹¹ together with the
nitrogen atom to which they are attached form a 4-7 membered heterocyclic ring, which
30 optionally contains one or more further heteroatoms selected from O, N and S, and which
is optionally substituted by from 1, 2 or 3 groups selected from hydroxy, (1-6C)alkyl,
(1-6C)haloalkyl, carboxy or a carboxylic acid mimic or bioisostere thereof;

R^9 is selected from hydroxy, halo, amino, cyano, ((1-3C)alkyl)CONH-, carboxy, (1-6C)alkoxy, (1-6C)alkoxycarbonyl, carbamoyl, N-((1-6C)alkyl)carbamoyl, halo((1-6C)alkyl) (such as trifluoromethyl), (1-6C)alkylsulphonyl, (1-6C)alkylsulphinyl,

AR1^a, AR1^b and AR1^c are independently selected from phenyl or a 6,5-, or
 5 6,6-bicyclic fused ring system, wherein both rings are carbocyclic and wherein at least the ring attached to the linking nitrogen atom is fully aromatic;


HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e and HET-1^f are independently
 selected from a 5- or 6-membered heteroaryl ring containing 1, 2, or 3 heteroatoms
 independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds),
 10 wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein
 any carbon atom may optionally be oxidised to a carbonyl group;

HET-2^a and HET-2^b are independently selected from a 5,5-, 5,6- 6,5- or 6,6-fused
 bicyclic, aromatic or partially unsaturated heterocyclic ring system, containing 1, 2, 3 or 4
 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or
 15 O-S bonds), in which the ring directly attached to the linking nitrogen atom is aromatic,
 and either ring of the bicyclic ring system may be carbocyclic;

HET-3^b, HET-3^c, HET-3^c are independently selected from a partially or fully
 unsaturated 4-, 5-, 6- or 7 membered heterocyclic ring containing 1, 2, or 3 heteroatoms
 independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds),
 20 wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein
 any carbon atom may optionally be oxidised to a carbonyl group; except that HET-3^b and
 HET-3^c are not 3-oxomorpholin-4-yl.

In a particular embodiment, the present invention provides a compound of formula
 (IB) or a pharmaceutically-acceptable salt thereof,

25



wherein:

R^{1b} is selected from AR1^a, HET-1^a and HET-2^a; wherein R^{1b} is optionally
 30 substituted by 1, 2, 3 or 4 groups independently selected from R^{3a} ;

40

- R^{3a} is selected from halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $R^{8a}NHCO-$, $R^{8a}CONH$, $R^{8a}OCONH-$, $R^{8a}OCON(Me)-$, $R^{8a}(1-6C)alkyl-$, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), $-OC_6H_{(13-a)}F_a$ (wherein a is 1 to 13), (1-6C)alkylthio, di(1-6C)alkylaminocarbonyl, (3-8C)cycloalkyloxy(1-6C)alkyl, (3-8C)cycloalkyl(1-6C)alkoxy,
- 5 (3-8C)cycloalkyl(1-6C)alkoxy(1-6C)alkyl, phenyl, HET-1^b, AR1^b-(1-6C)alkoxy, HET-1^b-(1-6C)alkoxy, HET-3^b-(1-6C)alkoxy, AR1^b-oxy(1-6C)alkyl, HET-1^b-oxy(1-6C)alkyl, HET-3^b-oxy(1-6C)alkyl, AR1^b-(1-6C)alkoxy(1-6C)alkyl, HET-1^b-(1-6C)alkoxy(1-6C)alkyl, and HET-3^b-(1-6C)alkoxy(1-6C)alkyl; wherein any phenyl, AR1^b, HET-1^b, HET-3^b or cycloalkyl ring in any of the above values
- 10 for R^{3a} may optionally be substituted by 1 or 2 substituents independently selected from R^{7d} , and wherein any R^{8a} , (3-8C)cycloalkyl, or (1-6C)alkyl group in any of the above definitions for R^{3a} may optionally be substituted by 1 or 2 substituents independently selected from
- 15 (1-6C)alkyl, (1-6C)alkoxy, (2-6C)alkenyl, (2-6C)alkynyl, hydroxy, cyano, carboxy, aminocarbonyl, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), amino, (1-6C)alkylamino, di(1-6C)alkylamino, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, aminosulfonyl, (1-6C)alkylaminosulfonyl, di(1-6C)alkylaminosulfonyl, (1-6C)alkylsulfonylamino, aminocarbonyl, (1-6C)alkylaminocarbonyl,
- 20 di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino, (1-6C)alkoxycarbonylamino, (1-6C)alkoxycarbonyl(N-methyl)amino and (1-6C)alkylaminocarbonyloxy; and/or any R^{8a} , (3-8C)cycloalkyl, or (1-6C)alkyl group in any of the above definitions for R^{3a} may optionally be substituted by 1, 2 or 3 halo;

R^{2b} is:

- 25 a) a 6-membered ring selected from phenyl, 3-pyridyl and 3,5-pyrimidinyl; substituted in the position which is para to the linkage with the group $-Y-NH-$ by a group selected from R^{4a} and optionally additionally substituted on a carbon atom in one or both of the meta-positions to the linkage with the group $-Y-NH-$ (where possible) by 1 or 2 groups independently selected from R^{6c} ; or
- 30 b) a 5-membered heteroaryl ring containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds); substituted on a carbon atom in a position which is not adjacent the linkage to the group $-Y-NH-$, by a group

selected from R^{4a} and optionally additionally substituted on carbon (where available) by 1 or 2 groups independently selected from R^{6d} ;

R^{4a} is selected from:

- a) phenyl;
 - 5 b) HET-3^c (wherein any available nitrogen atom is substituted by R^{5a} provided it is not thereby quaternised);
 - c) HET-1^c;
 - d) phenoxy, HET-1^c-oxy, $R^{8b}O-$, phenyl(1-6C)alkoxy and HET-1^c-(1-6C)alkoxy;
 - e) R^{8b} , $R^{8b}O(1-6C)alkyl$, amino(1-6C)alkyl, hydroxy(1-6C)alkyl, phenoxy(1-6C)alkyl,
 - 10 HET-1^c-oxy(1-6C)alkyl; and
 - f) phenylamino, phenyl(1-6C)alkylamino, HET-1^c-amino, HET-1^c-(1-6C)alkylamino, (1-6C)alkylamino, N-phenyl-N-(1-6C)alkylamino, N-phenyl(1-6C)alkyl-N-(1-6C)alkylamino, N-(HET-1^c)-N-(1-6C)alkylamino, N-(HET-1^c-(1-6C)alkyl)-N-(1-6C)alkylamino, and
 - 15 di(1-6C)alkylamino;
- wherein any alkyl, R^{8b} , phenyl, HET-1^c or HET-3^c ring in any of the definitions a) to f) for R^{4a} may optionally be substituted by 1 or 2 substituents independently selected from R^{7e} ;

- R^{5a} is selected from (1-6C)alkyl, (3-8C)cycloalkyl, phenyl (optionally substituted with 1 or 2 halo), HET-1^d, $R^{8c}CO-$, phenylcarbonyl (wherein the phenyl is optionally substituted with 1 or 2 halo), HET-1^d-carbonyl, (1-6C)alkoxycarbonyl, $R^{8c}NHCO-$,
- 20 (1-6C)alkoxysulfonyl, (1-6C)alkylaminosulfonyl, (1-6C)alkylsulfinyl and (1-6C)alkylsulfonyl; wherein any (1-6C)alkyl group in any of the definitions of R^{5a} may optionally be substituted by a substituent selected from (1-6C)alkoxy, (1-6C)alkylS(O) p' (wherein p' is 0, 1 or 2), amino, (1-6C)alkylamino, hydroxy, carboxy, (2-6C)alkenyl,
- 25 (2-6C)alkynyl, phenyl (optionally substituted with 1 or 2 halo) and HET-1^f;

- R^{6c} and R^{6d} are independently selected from R^{8d} , $R^{8d}O-$, $R^{8d}OCO-$, $R^{8d}COO-$, halo, cyano, halo(1-6C)alkyl, dihalo(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkoxy, (1-6C)alkylcarbonyl, (1-6C)alkylS(O) p (wherein p is 0, 1 or 2), (1-6C)alkylaminocarbonyl, di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino,
- 30 (1-6C)alkylaminosulfonyl, di(1-6C)alkylaminosulfonyl, (1-6C)alkylsulfonylamino, $AR1^e$ -(1-6C)alkoxy, HET-1^e-(1-6C)alkoxy, HET-3^e-(1-6C)alkoxy, $AR1^e$ -oxy(1-6C)alkyl,

HET-1^e-oxy(1-6C)alkyl, HET-3^e-oxy(1-6C)alkyl, AR1^e-(1-6C)alkoxy(1-6C)alkyl, HET-1^e-(1-6C)alkoxy(1-6C)alkyl, and HET-3^e-(1-6C)alkoxy(1-6C)alkyl;

wherein any alkyl, R^{8d}, phenyl, AR1^e, HET-1^e, HET-3^e or cycloalkyl ring in any of the above values for R^{6e} or R^{6d} may optionally be substituted by 1, 2 or 3 substituents

5 independently selected from R^{7f};

R^{7d}, R^{7e} and R^{7f} are independently selected from R^{8e} (optionally substituted with 1, 2 or 3 halo or with hydroxy), R^{8e}O- (optionally substituted with 1, 2 or 3 halo), R^{8e}OCO-, R^{8e}COO-, R^{8e}CO-, R^{8e}NHCO-, (R^{8e})₂NCO, R^{8e}CONH, R^{8e}OCNH-, R^{8e}OCON(Me)-, R^{8e}(1-6C)alkyl-, halo, cyano, carboxy, carboxy(1-6C)alkyl, hydroxy, aminocarbonyl,

10 -C₆H_(13-a)F_a (wherein a is 1 to 13), -OC₆H_(13-a)F_a (wherein a is 1 to 13), R^{8e}S(O)p^{''}- (wherein p^{''} is 0, 1 or 2), amino(1-6C)alkyl, R^{8e}NH(1-6C)alkyl, (R^{8e})₂N(1-6C)alkyl, R^{8e}NHSO₂-, (R^{8e})₂NSO₂-, and R^{8e}SO₂NH-;

each R^{8a}, R^{8b}, R^{8c}, R^{8d}, R^{8e} and R^{8f} is selected from (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl and (3-8C)cycloalkyl;

15 AR1^a, AR1^b or AR1^c are independently selected from phenyl or a 6,5-, or 6,6-bicyclic fused ring system, wherein both rings are carbocyclic and wherein at least the ring attached to the linking nitrogen atom is fully aromatic;

HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e and HET-1^f are independently selected from a 5- or 6-membered heteroaryl ring containing 1, 2, or 3 heteroatoms

20 independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein any carbon atom may optionally be oxidised to a carbonyl group;


HET-2^a and HET-2^b are independently selected from a 5,5-, 5,6- 6,5- or 6,6-fused bicyclic, aromatic or partially unsaturated heterocyclic ring system, containing 1, 2, 3 or 4
25 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), in which the ring directly attached to the linking nitrogen atom is aromatic, and either ring of the bicyclic ring system may be carbocyclic;

HET-3^b, HET-3^c, HET-3^e are independently selected from a partially or fully unsaturated 4-, 5-, 6- or 7 membered heterocyclic ring containing 1, 2, or 3 heteroatoms


30 independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein

any carbon atom may optionally be oxidised to a carbonyl group; except that HET-3^b and HET-3^c are not 3-oxomorpholin-4-yl.


It will be understood that the definition of R^{2b} in relation to formula (IB) as a 6-membered ring selected from phenyl, 3-pyridyl and 3,5-pyrimidinyl; substituted in the position which is para to the linkage with the group -Y-NH- with a group R^{4a} and optionally additionally substituted on a carbon atom in one or both positions which are meta to the linkage with the group -Y-NH- (where possible) by 1 or 2 groups independently selected from R^{6c}; encompasses the following structures, for every definition of substituent R^{6c}:



Similarly, it will be understood that the definition of R^{2b} as a 5-membered heteroaryl ring containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds); substituted on a carbon atom in a position which is not directly adjacent the linkage to the group -Y-NH- by a group selected from R^{4a} and optionally additionally substituted on carbon (where available) by 1 or 2 groups independently selected from R^{6d}; encompasses the following structures:



For the avoidance of doubt it is to be understood that where in this specification a group is qualified by 'hereinbefore defined' or 'defined hereinbefore' the said group



encompasses the first occurring and broadest definition as well as each and all of the particular definitions for that group.

It is to be understood that where substituents contain two substituents on an alkyl chain, in which both are linked by a heteroatom (for example two alkoxy substituents), then these two substituents are not substituents on the same carbon atom of the alkyl chain.

Examples of AR1^a, AR1^b and AR1^c are phenyl, naphthyl, indenyl and dihydroindenyl.

Examples of AR1^a-(1-6C)alkyl, AR1^b-(1-6C)alkyl or AR1^c-(1-6C)alkyl are benzyl, phenethyl, naphthylmethyl and naphthylethyl.

Examples of AR1^a-oxy, AR1^b-oxy or AR1^c-oxy are phenoxy and naphthyloxy. Examples of AR1^a-oxy(1-6C)alkyl, AR1^b-oxy(1-6C)alkyl or AR1^c-oxy(1-6C)alkyl are phenoxymethyl and naphthyloxyethyl. Examples of AR1^a-(1-6C)alkoxy, AR1^b-(1-6C)alkoxy or AR1^c-(1-6C)alkoxy are any value of AR1^a, AR1^b or AR1^c bonded to any value of (1-6C)alkoxy, for example benzyloxy and naphthyloxy. The same convention applies wherever AR1^a, AR1^b or AR1^c is linked to another group.

Examples of HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e or HET-1^f are oxazolyl, oxadiazolyl, pyridyl, pyrimidinyl, imidazolyl, triazolyl, pyrazinyl, pyridazinyl, pyrrolyl, thienyl and furyl.

Further examples of HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e or HET-1^f include thiazolyl, tetrazolyl, isoxazolyl, pyrazolyl, triazinyl, thiadiazolyl, oxadiazolyl and isothiazolyl.

Examples of HET-1^a-oxy, HET-1^b-oxy, HET-1^c-oxy, HET-1^d-oxy, HET-1^e-oxy or HET-1^f-oxy are oxazolyl-2-oxy, 1,2,4-oxadiazolyl-3-oxy, pyridyl-2-oxy, and pyrimidinyl-4-oxy. The convention illustrated above for AR1^a, AR1^b and AR1^c applies to HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e or HET-1^f linked to any other group.

Examples of HET-2^a or HET-2^b are 1,3-benzodioxol-5-yl, quinazolinyl, purinyl, indolyl, benzofuryl, benzothienyl, chromanyl, isochromanyl, quinolinyl and isoquinolinyl. Further examples of HET-2^a or HET-2^b include quinoxalinyl, benzthiazolyl and benzoxazolyl.

Examples of HET-3^b, HET-3^c or HET-3^e are morpholinyl, thiomorpholinyl, piperazinyl, piperidinyl, 2-oxopiperidinyl, pyrrolidinyl, 2-oxopyrrolidinyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothienyl, azetidyl, homomorpholinyl,

diazepinyl and azepinyl.

Particular examples of HET-3^b, HET-3^c or HET-3^e are morpholinyl, thiomorpholinyl, piperazinyl and piperidinyl.

5 Examples of HET-3^b-oxy, HET-3^c-oxy or HET-3^e-oxy are morpholinylloxy, thiomorpholinylloxy, piperazinylloxy, piperidinylloxy, 2-oxopiperidinylloxy, pyrrolidinylloxy, 2-oxopyrrolidinylloxy, tetrahydropyranyloxy, dihydropyranyloxy, tetrahydrothienylloxy, azetidylloxy, homomorpholinylloxy, diazepinylloxy and azepinylloxy. The convention illustrated above for AR1^a, AR1^b and AR1^e applies to HET-3^b, HET-3^c or HET-3^e linked to any other group.

10 If not stated elsewhere, suitable optional substituents for a particular group are those as stated for similar groups herein.

A compound of formula (I) may form stable acid or basic salts, and in such cases administration of a compound as a salt may be appropriate, and pharmaceutically acceptable salts may be made by conventional methods such as those described following.

15 Suitable pharmaceutically-acceptable salts include acid addition salts such as methanesulfonate, tosylate, α -glycerophosphate, fumarate, hydrochloride, citrate, maleate, tartrate and (less preferably) hydrobromide. Also suitable are salts formed with phosphoric and sulfuric acid. In another aspect suitable salts are base salts such as an alkali metal salt for example sodium, an alkaline earth metal salt for example calcium or magnesium, an organic amine salt for example triethylamine, morpholine, *N*-methylpiperidine, *N*-ethylpiperidine, procaine, dibenzylamine, *N,N*-dibenzylethylamine, tris-(2-hydroxyethyl)amine, *N*-methyl d-glucamine and amino acids such as lysine. There may be more than one cation or anion depending on the number of charged functions and the valency of the cations or anions. A preferred pharmaceutically-acceptable salt is the sodium salt.

25 However, to facilitate isolation of the salt during preparation, salts which are less soluble in the chosen solvent may be preferred whether pharmaceutically-acceptable or not.

30 Within the present invention it is to be understood that a compound of the formula (I) or a salt thereof may exhibit the phenomenon of tautomerism and that the formulae drawings within this specification can represent only one of the possible tautomeric forms. It is to be understood that the invention encompasses any tautomeric form which inhibits

DGAT1 activity and is not to be limited merely to any one tautomeric form utilised within the formulae drawings.

Various forms of prodrugs are known in the art. For examples of such prodrug derivatives, see:

- 5 a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and
H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard
10 p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- d) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- e) N. Kakeya, *et al.*, Chem Pharm Bull, 32, 692 (1984).

Examples of such prodrugs are *in vivo* cleavable esters of a compound of the
15 invention. An *in vivo* cleavable ester of a compound of the invention containing a carboxy group is, for example, a pharmaceutically-acceptable ester which is cleaved in the human or animal body to produce the parent acid. Suitable pharmaceutically-acceptable esters for carboxy include (1-6C)alkyl esters, for example methyl or ethyl; (1-6C)alkoxymethyl esters, for example methoxymethyl; (1-
20 6C)alkanoyloxymethyl esters, for example pivaloyloxymethyl; phthalidyl esters; (3-8C)cycloalkoxycarbonyloxy(1-6C)alkyl esters, for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolan-2-ylmethyl esters, for example 5-methyl-1,3-dioxolan-2-ylmethyl; (1-6C)alkoxycarbonyloxyethyl esters, for example 1-methoxycarbonyloxyethyl; aminocarbonylmethyl esters and mono- or di- N-((1-
25 6C)alkyl) versions thereof, for example N,N-dimethylaminocarbonylmethyl esters and N-ethylaminocarbonylmethyl esters; and may be formed at any carboxy group in the compounds of this invention. An *in vivo* cleavable ester of a compound of the invention containing a hydroxy group is, for example, a pharmaceutically-acceptable ester which is cleaved in the human or animal body to produce the parent hydroxy group. Suitable
30 pharmaceutically acceptable esters for hydroxy include (1-6C)alkanoyl esters, for example acetyl esters; and benzoyl esters wherein the phenyl group may be substituted



with aminomethyl or N- substituted mono- or di- (1-6C)alkyl aminomethyl, for example 4-aminomethylbenzoyl esters and 4-N,N-diethylaminomethylbenzoyl esters.

It will be appreciated by those skilled in the art that certain compounds of formula (I) contain asymmetrically substituted carbon and/or sulfur atoms, and accordingly may exist in, and be isolated in, optically-active and racemic forms. Some compounds may exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic or stereoisomeric form, or mixtures thereof, which form possesses properties useful in the inhibition of DGAT1 activity, it being well known in the art how to prepare optically-active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, by enzymatic resolution, by biotransformation, or by chromatographic separation using a chiral stationary phase) and how to determine efficacy for the inhibition of DGAT1 activity by the standard tests described hereinafter.

It is also to be understood that certain compounds of the formula (I) and salts thereof can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which inhibit DGAT1 activity.

As stated before, we have discovered a range of compounds that have good DGAT1 inhibitory activity. They have good physical and/or pharmacokinetic properties in general. The following compounds possess preferred pharmaceutical and/or physical and/or pharmacokinetic properties.

Particular aspects of the invention comprise a compound of formula (I), or a pharmaceutically-acceptable salt thereof, wherein the substituents AR, HET, R¹ to R⁸ and other substituents mentioned above have values defined hereinbefore, or any of the following values (which may be used where appropriate with any of the definitions and embodiments disclosed hereinbefore or hereinafter):

In one embodiment of the invention there are provided compounds of formula (I), in an alternative embodiment there are provided pharmaceutically-acceptable salts of compounds of formula (I). In a further embodiment, there are provided pro-drugs, particularly in-vivo cleavable esters, of compounds of formula (I). In a further embodiment, there are provided pharmaceutically-acceptable salts of pro-drugs of compounds of formula (I). Reference herein to a compound of formula (I) should in

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO DIVISION DE NUEVAS CREACIONES				
PATENTE	☒	PCT/GB2005/004726	MODELO DE UTILIDAD ☐	DISEÑO INDUSTRIAL ☐
SOLICITANTE/S, RAZON SOCIAL O APELLIDOS Y NOMBRES <u>ASTRAZENECA AB</u>				
DOMICILIO <u>SÖDERTÄLJE, SUECIA</u>				
APODERADO <u>JUAN PABLO CADENA SARMIENTO</u>				
TITULO <u>DERIVADOS DE OXADIAZOL COMO INHIBIDORES DE DGAT</u>				
CLASIFICACION _____				
EXPEDIENTE No.	FECHA DE SOLICITUD _____		ROLLO MICROFILM _____	
CERTIFICADO No.	FECHA DE CONCESION _____		VIGENCIA DESDE _____ HASTA _____	
PRORROGA _____	CAMBIO DE NOMBRE _____			
TRASPASO A _____				
INVENTOR (ES) <u>ALAN MARTIN BIRCH, SUZANNE SAXON BOWKER, ROGER JOHN BUTLIN, CRAIG SAMUEL</u>				
OTROS <u>DONALD, WILLIAM MCCOULL, THORSTEN NOWAK, ALLEYN PLOWRIGHT.</u>				

OFICINA DE COMUNICACIONES

general be taken to apply also to compounds of formulae (IA), (IB), (IZA), (IZB) and (IZC).

Particular values of variable groups in compounds of formulae (I), (IA) and (IB) are as follows. Such values may be used where appropriate with any of the other values,

5 definitions, claims or embodiments defined hereinbefore or hereinafter.

- 1) R^1, R^{1a} or R^{1b} is $AR1^a$
- 2) R^1, R^{1a} or R^{1b} is HET-1^a
- 3) R^1, R^{1a} or R^{1b} is HET-2^a
- 4) R^1, R^{1a} or R^{1b} is unsubstituted
- 10 5) R^1 or R^{1a} is substituted by 1 group selected from R^3 or R^{1b} is substituted by 1 group selected from R^{3a}
- 6) R^1 or R^{1a} is substituted by 2 groups independently selected from R^3 or R^{1b} is substituted by 2 groups independently selected from R^{3a}
- 7) R^1 or R^{1a} is substituted by 3 groups independently selected from R^3 or R^{1b} is substituted by 3 groups independently selected from R^{3a}
- 15 8) R^1 or R^{1a} is substituted by 4 groups independently selected from R^3 or R^{1b} is substituted by 4 groups independently selected from R^{3a}
- 9) R^3 or R^{3a} is halo
- 10) R^3 or R^{3a} is selected from $R^{8a}, R^{8a}O-, R^{8a}OCO-, R^{8a}COO-$ and $R^{8a}CO-$
- 20 11) R^3 or R^{3a} is selected from $R^{8a}, R^{8a}O-$ and $R^{8a}OCO-$
- 12) R^3 or R^{3a} is selected from $R^{8a}NHCO-, R^{8a}CONH-, R^{8a}OCONH-$ and $R^{8a}OCON(Me)-$
- 13) R^3 or R^{3a} is selected from $R^{8a}(1-6C)alkyl-, haloC_{1-6}alkyl$ or $haloC_{1-6}alkoxy$
- 14) R^3 or R^{3a} is selected from $R^{8a}(1-6C)alkyl-, -C_6H_{(13-a)}F_a$ (wherein a is 1 to 14), $-OC_6H_{(13-a)}F_a$ (wherein a is 1 to 14)
- 25 15) R^3 or R^{3a} is selected from $(1-6C)alkylthio, alkylcarbonyl$ and $di(1-6C)alkylaminocarbonyl$
- 16) R^3 or R^{3a} is selected from $(3-8C)cycloalkyloxy(1-6C)alkyl, (3-8C)cycloalkyl(1-6C)alkoxy$, and $(3-8C)cycloalkyl(1-6C)alkoxy(1-6C)alkyl$
- 30 17) R^3 or R^{3a} is selected from halo, $R^{8a}, R^{8a}O-, R^{8a}OCO-, R^{8a}COO-, R^{8a}CO-, R^{8a}NHCO-, R^{8a}CONH-, R^{8a}OCONH-, R^{8a}OCON(Me)-, R^{8a}(1-6C)alkyl-, halo(1-6C)alkyl$ (such as $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), $halo(1-6C)alkoxy$ (such as $-OC_6H_{(13-a)}F_a$

- (wherein a is 1 to 13), (1-6C)alkylthio, di(1-6C)alkylaminocarbonyl, (3-8C)cycloalkyloxy(1-6C)alkyl, (3-8C)cycloalkyl(1-6C)alkoxy, (3-8C)cycloalkyl(1-6C)alkoxy(1-6C)alkyl, AR1^b-(1-6C)alkoxy, HET-1^b-(1-6C)alkoxy, HET-3^b-(1-6C)alkoxy, AR1^b-oxy(1-6C)alkyl, HET-1^b-oxy(1-6C)alkyl,
- 5 HET-3^b-oxy(1-6C)alkyl, AR1^b-(1-6C)alkoxy(1-6C)alkyl, HET-1^b-(1-6C)alkoxy(1-6C)alkyl, and HET-3^b-(1-6C)alkoxy(1-6C)alkyl
- 18) R³ or R^{3a} is selected from phenyl and HET-1^b
- 19) R³ or R^{3a} is phenyl
- 20) R³ or R^{3a} is selected from AR1^b-(1-6C)alkoxy, and HET-1^b-(1-6C)alkoxy,
- 10 HET-3^b-(1-6C)alkoxy
- 21) R³ or R^{3a} is selected from AR1^b-oxy(1-6C)alkyl, HET-1^b-oxy(1-6C)alkyl, HET-3^b-oxy(1-6C)alkyl, AR1^b-(1-6C)alkoxy(1-6C)alkyl, HET-1^b-(1-6C)alkoxy(1-6C)alkyl, and HET-3^b-(1-6C)alkoxy(1-6C)alkyl
- 22) a phenyl, AR1^b, HET-1^b, HET-3^b or cycloalkyl ring in any value for R³ is
- 15 substituted by 1 or 2 substituents independently selected from R^{7a}, and a phenyl, AR1^b, HET-1^b, HET-3^b or cycloalkyl ring in any value for R^{3a} is substituted by 1 or 2 substituents independently selected from R^{7d}
- 23) a phenyl, AR1^b, HET-1^b, HET-3^b or cycloalkyl ring in any value for R³ or R^{3a} is unsubstituted
- 20 24) an R^{8a}, (3-8C)cycloalkyl, or (1-6C)alkyl group in any definitions for R³ or R^{3a} is unsubstituted
- 25) an R^{8a}, (3-8C)cycloalkyl, or (1-6C)alkyl group in any definitions for R³ or R^{3a} is substituted by 1 or 2 substituents independently selected from (1-6C)alkyl, (1-6C)alkoxy, (2-6C)alkenyl, (2-6C)alkynyl, hydroxy, cyano, carboxy, aminocarbonyl, -C₆H_(14-a)F_a
- 25 (wherein a is 1 to 14), amino, (1-6C)alkylamino, di(1-6C)alkylamino, (1-6C)alkylthio, (1-6C)alkylsulfinyl, (1-6C)alkylsulfonyl, aminosulfonyl, (1-6C)alkylaminosulfonyl, di(1-6C)alkylaminosulfonyl, (1-6C)alkylsulfonylamino, aminocarbonyl, (1-6C)alkylaminocarbonyl, di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino, (1-6C)alkoxycarbonylamino, (1-6C)alkoxycarbonyl(N-methyl)amino and
- 30 (1-6C)alkylaminocarbonyloxy; and/or substituted by 1, 2 or 3 halo;
- 26) R² or R^{2a} is a 6-membered ring selected from phenyl, 3-pyridyl, 5-pyridyl and 3,5-pyrimidinyl; substituted in the 4-position by a group selected from R⁴ and optionally

additionally substituted on a carbon atom in one or both meta-positions relative to the linkage to the group $-Y-NH-$ (where possible) by 1 or 2 groups independently selected from R^{6a} ; or

- 27) R^{2b} is a 6-membered ring selected from phenyl, 3-pyridyl, 5-pyridyl and
 5 3,5-pyrimidinyl; substituted in the 4-position by a group selected from R^{4a} and optionally additionally substituted on a carbon atom in one or both meta-positions relative to the linkage to the group $-Y-NH-$ (where possible) by 1 or 2 groups independently selected from R^{6c} ;
- 28) R^2 , R^{2a} or R^{2b} is phenyl
- 10 29) R^2 , R^{2a} or R^{2b} is 3-pyridyl
- 30) R^2 , R^{2a} or R^{2b} is phenyl or 3-pyridyl
- 31) R^2 or R^{2a} is a six-membered ring substituted in the position which is para to the linkage with the group $-Y-NH-$ by a group selected from R^4 ; or a group R^{2b} is a six-membered ring substituted in the position which is para to the linkage with the group $-Y-NH-$ by a group selected from R^{4a}
- 15 32) R^2 or R^{2a} is a six-membered ring substituted in the position which is para to the linkage with the group $-Y-NH-$ by a group selected from R^4 and additionally substituted on a carbon atom in a 3-position by R^{6a} ; or R^{2b} is a six-membered ring substituted in the position which is para to the linkage with the group $-Y-NH-$ by a group selected from R^{4a} and additionally substituted on a carbon atom in a 3-position by R^{6c} ;
- 20 33) R^2 or R^{2a} is a group or sub-formula (iv) as defined above; or R^{2b} is a 5-membered heteroaryl ring containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds); substituted on a carbon atom in a position which is not adjacent the linkage to the group $-Y-NH-$ by a group selected from R^{4a} and optionally additionally substituted on carbon (where available) by 1 or 2 groups independently selected from R^{6d}
- 25 34) R^2 or R^{2a} is a group HET-2^b optionally substituted with methyl
- 35) R^4 or R^{4a} is phenyl
- 36) R^4 or R^{4a} is HET-3^c (wherein any available nitrogen atom is substituted by R^5 provided it is not thereby quaternised)
- 30 37) R^4 or R^{4a} is HET-1^c
- 38) R^4 or R^{4a} is selected from phenoxy, HET-1^c-oxy, $R^{8b}O-$, phenyl(1-6C)alkoxy and

HET-1^c-(1-6C)alkoxy

- 39) R^4 or R^{4a} is selected from R^{8b} , $R^{8b}O(1-6C)alkyl$, amino(1-6C)alkyl, hydroxy(1-6C)alkyl, phenoxy(1-6C)alkyl and HET-1^c-oxy(1-6C)alkyl
- 40) R^4 or R^{4a} is selected from R^{8b} and hydroxy(1-6C)alkyl
- 5 41) R^4 or R^{4a} is selected from phenylamino, phenyl(1-6C)alkylamino, HET-1^c-amino, HET-1^c-(1-6C)alkylamino, (1-6C)alkylamino, N-phenyl-N-(1-6C)alkylamino, N-phenyl(1-6C)alkyl-N-(1-6C)alkylamino, N-(HET-1^c)-N-(1-6C)alkylamino, N-(HET-1^c-(1-6C)alkyl)-N-(1-6C)alkylamino, and di(1-6C)alkylamino
- 42) R^4 or R^{4a} is selected from phenylamino, N-phenyl-N-(1-6C)alkylamino
- 10 43) R^4 or R^{4a} is selected from phenyl, HET-3^c, phenoxy, HET-1^c-oxy, $R^{8b}O$ -, phenyl(1-6C)alkoxy, HET-1^c-(1-6C)alkoxy, R^{8b} , hydroxy(1-6C)alkyl, phenylamino and N-phenyl-N-(1-6C)alkylamino
- 44) an alkyl, R^{8b} , phenyl, HET-1^c or HET-3^c ring in any of the definitions for R^4 is substituted by 1 or 2 substituents independently selected from R^{7b} ; or an alkyl, R^{8b} , phenyl, HET-1^c or HET-3^c ring in any of the definitions for R^{4a} is substituted by 1 or 2 substituents independently selected from R^{7c}
- 15 45) an alkyl, R^{8b} , phenyl, HET-1^c or HET-3^c ring in any of the definitions for R^4 or R^{4a} is unsubstituted
- 46) R^5 or R^{5a} is selected from (1-6C)alkyl and (3-8C)cycloalkyl
- 20 47) R^5 or R^{5a} is $R^{8c}CO$ -
- 48) R^5 or R^{5a} is phenyl (optionally substituted) or HET-1^d
- 49) R^5 or R^{5a} is phenylcarbonyl (optionally substituted) or HET-1^d-carbonyl
- 50) R^5 or R^{5a} is selected from carboxy, (1-6C)alkoxycarbonyl, $R^{8c}NHCO$ -, (1-6C)alkoxysulfonyl, (1-6C)alkylaminosulfonyl, (1-6C)alkylsulfinyl and
- 25 (1-6C)alkylsulfonyl
- 51) R^5 or R^{5a} is selected from (1-6C)alkoxycarbonyl and (1-6C)alkylsulfonyl
- 52) R^5 or R^{5a} is selected from (1-6C)alkyl, HET-1, $R^{8c}CO$ -, (1-6C)alkoxycarbonyl and (1-6C)alkylsulfonyl
- 53) any (1-6C)alkyl group in any of the definitions of R^5 or R^{5a} is substituted by a
- 30 substituent selected from (1-6C)alkoxy, (1-6C)alkylS(O)p- (wherein p is 0, 1 or 2), amino, (1-6C)alkylamino, hydroxy, carboxy, (2-6C)alkenyl, (2-6C)alkynyl, phenyl (optionally substituted) and HET-1^d

- 54) any (1-6C)alkyl group in any of the definitions of R^5 or R^{5a} is unsubstituted
- 55) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from R^{8d} , $R^{8d}O-$, $R^{8d}OCO-$ and $R^{8d}COO-$
- 56) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from halo and cyano
- 57) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from halo(1-6C)alkyl, dihalo(1-6C)alkyl
- 5 58) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from (1-6C)alkoxy(1-6C)alkyl and (1-6C)alkoxy(1-6C)alkoxy
- 59) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from (1-6C)alkylcarbonyl and (1-6C)alkylS(O)p- (wherein p is 0, 1 or 2)
- 60) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from (1-6C)alkylaminocarbonyl,
- 10 di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino, (1-6C)alkylaminosulfonyl, di(1-6C)alkylaminosulfonyl and (1-6C)alkylsulfonylamino
- 61) R^{6a} , R^{6b} , R^{6c} or R^{6d} is selected from $AR1^e$ -(1-6C)alkoxy, HET-1^e-(1-6C)alkoxy, HET-3^e-(1-6C)alkoxy, $AR1^e$ -oxy(1-6C)alkyl, HET-1^e-oxy(1-6C)alkyl, HET-3^e-oxy(1-6C)alkyl, $AR1^e$ -(1-6C)alkoxy(1-6C)alkyl,
- 15 HET-1^e-(1-6C)alkoxy(1-6C)alkyl, and HET-3^e-(1-6C)alkoxy(1-6C)alkyl
- 62) any alkyl, R^{8d} , phenyl, $AR1^e$, HET-1^e, HET-3^e or cycloalkyl ring in any value for R^{6a} , R^{6b} is substituted by 1, 2 or 3 substituents independently selected from R^{7c} , or in any value for R^{6c} or R^{6d} is substituted by 1, 2 or 3 substituents independently selected from R^{7f}
- 20 63) any alkyl, R^{8d} , phenyl, $AR1^e$, HET-1^e, HET-3^e or cycloalkyl ring in any value for R^{6a} , R^{6b} , R^{6c} or R^{6d} is unsubstituted
- 64) R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} or R^{7f} is R^{8e} (optionally substituted with 1, 2 or 3 halo or with hydroxy)
- 65) R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} or R^{7f} is $R^{8e}O-$ (optionally substituted with 1, 2 or 3 halo)
- 25 66) R^{7a} , R^{7b} or R^{7c} is selected from $R^{8e}OCO-$, $R^{8e}COO-$, $R^{8e}CO-$, $R^{8e}NHCO-$, $R^{10}R^{11}NCO$, $R^{8e}CONH$, $R^{8e}OCONH-$ and $R^{8e}OCON(Me)-$
- 67) R^{7d} , R^{7e} or R^{7f} is selected from $R^{8e}OCO-$, $R^{8e}COO-$, $R^{8e}CO-$, $R^{8e}NHCO-$, $(R^{8e})_2NCO$, $R^{8e}CONH$, $R^{8e}OCONH-$ and $R^{8e}OCON(Me)-$
- 68) R^{7a} , R^{7b} or R^{7c} is selected from R^{8e} (1-6C)alkyl-, halo, cyano, carboxy,
- 30 carboxy(1-6C)alkyl, hydroxy, aminocarbonyl, halo(1-6C)alkyl, halo(1-6C)alkoxy

- 69) R^{7d} , R^{7e} or R^{7f} is selected from R^{8a} (1-6C)alkyl-, halo, cyano, carboxy, carboxy(1-6C)alkyl, hydroxy, aminocarbonyl, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), $-OC_6H_{(14-a)}F_a$ (wherein a is 1 to 13)
- 70) R^{7a} , R^{7b} or R^{7c} is selected from $R^{8e}S(O)_p$ - (wherein p is 0, 1 or 2), amino(1-6C)alkyl, $R^{8c}NH(1-6C)alkyl$, $R^{11}R^{10}N(1-6C)alkyl$, $R^{8e}NHSO_2$ -, $R^{11}R^{10}NSO_2$ -, and $R^{8e}SO_2NH$ -
- 71) R^{7d} , R^{7e} or R^{7f} is selected from $R^{8e}S(O)_p$ - (wherein p is 0, 1 or 2), amino(1-6C)alkyl, $R^{8c}NH(1-6C)alkyl$, $(R^{8c})_2N(1-6C)alkyl$, $R^{8e}NHSO_2$ -, $(R^{8e})_2NSO_2$ -, and $R^{8e}SO_2NH$ -
- 72) R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} or R^{8f} is selected from (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (3-8C)cycloalkyl
- 73) R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} or R^{8f} is selected from (1-6C)alkyl and (3-8C)cycloalkyl
- 74) R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} or R^{8f} is selected from (1-6C)alkyl
- 75) $AR1^a$, $AR1^b$ or $AR1^c$ is phenyl
- 76) $AR1^a$, $AR1^b$ or $AR1^c$ is a 6,5-, or 6,6-bicyclic fused ring system,
- 77) $AR1^a$, $AR1^b$ or $AR1^c$ is naphthyl
- 78) $AR1^a$, $AR1^b$ or $AR1^c$ is phenyl, dihydroindenyl or naphthyl
- 79) $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ or $HET-1^f$ is 5- membered heteroaryl ring
- 80) $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ or $HET-1^f$ is a 6-membered heteroaryl ring
- 81) $HET-2^a$ or $HET-2^b$ is a 5,5- or 6,6-fused bicyclic heterocyclic ring system
- 82) $HET-2^a$ or $HET-2^b$ is a 5,6- 6,5- fused bicyclic heterocyclic ring system
- 83) $HET-3^b$, $HET-3^c$ or $HET-3^e$ is a 5-membered ring
- 84) $HET-3^b$, $HET-3^c$ or $HET-3^e$ is a 6-membered ring
- 85) $HET-3^b$, $HET-3^c$ or $HET-3^e$ is a 7-membered ring
- 86) $HET-3^b$, $HET-3^c$ or $HET-3^e$ is a 4-membered ring
- 87) s is suitably 0
- In yet further preferred compounds:
- 88) R^4 is a group R^{8b} where R^{8b} is (3-8C)cycloalkyl such as cyclohexyl
- 89) R^4 is a group R^{8b} where R^{8b} is substituted by a substituent R^{7b}
- 90) R^{7b} is $H_2NCO(CR^{13}R^{14})_n$ - or $(R^{10}R^{11})NCO(CH_2)_n$

- 91) R^{7b} is $HCONH(CR^{13}R^{14})_n$, $HCONR^{8c}(CR^{13}R^{14})_n$, $R^{8c}CONR^{9f}(CR^{13}R^{14})_n$
 92) R^{7b} is R^{12} or $R^{12}(1-6C)alkyl$,
 93) R^2 or R^{2a} is a cyclohexyl group.

In a further aspect of the invention there is provided a compound of the formula
 5 (IB') or a pharmaceutically-acceptable salt thereof, which is a compound of the formula (IB) as shown above, wherein:

R^{1b} is selected from $AR1^a$, $HET-1^a$ and $HET-2^a$; wherein R^{1b} is optionally substituted by 1, or 2 groups independently selected from R^{3a} ;

- R^{3a} is selected from halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $R^{8a}NHCO-$,
 10 $R^{8a}CONH$, $R^{8a}OCONH-$, $R^{8a}OCON(Me)-$, $R^{8a}(1-6C)alkyl-$, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), $-OC_6H_{(13-a)}F_a$ (wherein a is 1 to 13), (1-6C)alkylthio, di(1-6C)alkylaminocarbonyl, phenyl, $HET-1^b$, $AR1^b-(1-6C)alkoxy$, $HET-1^b-(1-6C)alkoxy$, $HET-3^b-(1-6C)alkoxy$, $AR1^b-oxy(1-6C)alkyl$, $HET-1^b-oxy(1-6C)alkyl$, $HET-3^b-oxy(1-6C)alkyl$, $AR1^b-(1-6C)alkoxy(1-6C)alkyl$, $HET-1^b-(1-6C)alkoxy(1-6C)alkyl$, and
 15 $HET-3^b-(1-6C)alkoxy(1-6C)alkyl$;

wherein any phenyl, $AR1^b$, $HET-1^b$, $HET-3^b$ or cycloalkyl ring in any of the above values for R^{3a} may optionally be substituted by 1 or 2 substituents independently selected from R^{7d} ;

- and wherein any R^{8a} , or (1-6C)alkyl group in any of the above definitions for R^{3a} may
 20 optionally be substituted by 1 or 2 substituents independently selected from (1-6C)alkyl, (1-6C)alkoxy, hydroxy, carboxy, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), (1-6C)alkylamino, di(1-6C)alkylamino, (1-6C)alkylsulfonyl, (1-6C)alkylaminosulfonyl, di(1-6C)alkylaminosulfonyl, (1-6C)alkylsulfonylamino, aminocarbonyl, (1-6C)alkylaminocarbonyl, di(1-6C)alkylaminocarbonyl, (1-6C)alkylcarbonylamino,
 25 (1-6C)alkoxycarbonylamino; and/or any R^{8a} , or (1-6C)alkyl group in any of the above definitions for R^{3a} may optionally be substituted by 1, 2 or 3 halo;

R^{2b} is:

- a) a 6-membered ring selected from phenyl, 3-pyridyl and 3,5-pyrimidinyl; substituted in the which is para to the linkage of the group $-Y-NH-$ by a group selected from R^{4a} and
 30 optionally additionally substituted on a carbon atom in one or both 3-positions (where possible) by 1 or 2 groups independently selected from R^{6c} ;

R^{4a} is selected from:

a) phenyl;

b) HET-3^c (wherein any available nitrogen atom is substituted by R^{5a} provided it is not thereby quaternised);

c) HET-1^c;

5 d) phenoxy, HET-1^c-oxy, R^{8b}O-, phenyl(1-6C)alkoxy and HET-1^c-(1-6C)alkoxy;

e) R^{8b}, R^{8b}O(1-6C)alkyl, amino(1-6C)alkyl, hydroxy(1-6C)alkyl, phenoxy(1-6C)alkyl, HET-1^c-oxy(1-6C)alkyl; and

f) phenylamino, phenyl(1-6C)alkylamino, HET-1^c-amino, HET-1^c-(1-6C)alkylamino,

(1-6C)alkylamino, N-phenyl-N-(1-6C)alkylamino, N-phenyl(1-6C)alkyl-N-(1-6C)alkylamin

10 o, N-(HET-1^c)-N-(1-6C)alkylamino, N-(HET-1^c-(1-6C)alkyl)-N-(1-6C)alkylamino, and di(1-6C)alkylamino;

wherein any alkyl, R^{8b}, phenyl, HET-1^c or HET-3^c ring in any of the definitions a) to f) for R^{4a} may optionally be substituted by 1 or 2 substituents independently selected from R^{7c};

R^{5a} is selected from (1-6C)alkyl, (3-8C)cycloalkyl, phenyl (optionally substituted
15 with 1 or 2 halo), HET-1^d, R^{8c}CO-, phenylcarbonyl (wherein the phenyl is optionally substituted with 1 or 2 halo), HET-1^d-carbonyl, (1-6C)alkoxycarbonyl, R^{8c}NHCO-, (1-6C)alkoxysulfonyl, (1-6C)alkylaminosulfonyl, (1-6C)alkylsulfinyl and (1-6C)alkylsulfonyl; wherein any (1-6C)alkyl group in any of the definitions of R^{5a} may optionally be substituted by a substituent selected from (1-6C)alkoxy, (1-6C)alkylS(O)p'-
20 (wherein p' is 0, 1 or 2), amino, (1-6C)alkylamino, hydroxy, carboxy, phenyl (optionally substituted with 1 or 2 halo) and HET-1^f;

R^{6c} and R^{6d} are independently selected from R^{8d}, R^{8d}O-, R^{8d}OCO-, R^{8d}COO-, halo, cyano, halo(1-6C)alkyl, dihalo(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkyl, (1-6C)alkoxy(1-6C)alkoxy, (1-6C)alkylcarbonyl and (1-6C)alkylS(O)p- (wherein p is 0, 1
25 or 2), wherein any alkyl or R^{8d} in any of the above values for R^{6c} or R^{6d} may optionally be substituted by 1, 2 or 3 substituents independently selected from R^{7f};

R^{7d}, R^{7e} and R^{7f} are independently selected from R^{8e} (optionally substituted with 1, 2 or 3 halo or with hydroxy), R^{8e}O- (optionally substituted with 1, 2 or 3 halo), halo, cyano, carboxy, carboxy(1-6C)alkyl, hydroxy, aminocarbonyl, -C₆H_(13-a)F_a (wherein a is 1
30 to 13), -OC₆H_(13-a)F_a (wherein a is 1 to 13), R^{8e}S(O)p''- (wherein p'' is 0, 1 or 2);

each R^{8a}, R^{8b}, R^{8c}, R^{8d}, R^{8e} or R^{8f} is selected from (1-6C)alkyl;

each $AR1^a$, $AR1^b$ or $AR1^c$ is phenyl or a 6,5-, or 6,6-bicyclic fused ring system, wherein both rings are carbocyclic and wherein at least the ring attached to the linking nitrogen atom is fully aromatic;

5 each $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ or $HET-1^f$ is 5- or 6-membered heteroaryl ring containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein any carbon atom may optionally be oxidised to a carbonyl group;

10 each $HET-2^a$ or $HET-2^b$ is a 5,5-, 5,6- 6,5- or 6,6-fused bicyclic, aromatic or partially saturated heterocyclic ring system, containing 1, 2, 3 or 4 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), in which the ring directly attached to the linking nitrogen atom is aromatic, and either ring of the bicyclic ring system may be carbocyclic;

15 $HET-3^b$, $HET-3^c$ or $HET-3^e$ is independently selected from a partially or fully saturated 4-, 5-, 6- or 7 membered heterocyclic ring containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein any carbon atom may optionally be oxidised to a carbonyl group; except that $HET-3^b$, $HET-3^c$ or $HET-3^e$ is not 3-oxomorpholin-4-yl.

20

In a further aspect of the invention there is provided a compound of the formula (IB'') or a pharmaceutically-acceptable salt thereof, which is a compound of the formula (IB) as shown above, wherein:

25 R^{1b} is selected from $AR1^a$, $HET-1^a$ and $HET-2^a$; wherein R^{1b} is optionally substituted by 1, or 2 groups independently selected from R^{3a} ;

R^{3a} is selected from halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13), $-OC_6H_{(13-a)}F_a$ (wherein a is 1 to 13), (1-6C)alkylthio, di(1-6C)alkylaminocarbonyl, phenyl, $HET-1^b$, $AR1^b$ -(1-6C)alkoxy, $HET-1^b$ -(1-6C)alkoxy, wherein any R^{8a} , or (1-6C)alkyl group in any of the above definitions for R^{3a} may
30 optionally be substituted by 1 or 2 substituents independently selected from (1-6C)alkoxy, hydroxy, carboxy, $-C_6H_{(13-a)}F_a$ (wherein a is 1 to 13); and/or by 1, 2 or 3 halo;

R^{2b} is phenyl or 3-pyridyl; substituted in the position which is para to the linkage with the group $-Y-NH-$ by a group selected from R^{4a} and optionally additionally substituted on a carbon atom in one or both meta positions with respect to the linkage with the group $-Y-NH-$ (where possible) by 1 or 2 groups independently selected from R^{6d} ;

5 R^{4a} is selected from:

- a) phenyl;
- b) HET-3^c (wherein any available nitrogen atom is substituted by R^{5a} provided it is not thereby quaternised);
- d) phenoxy, HET-1^c-oxy, $R^{8b}O-$, phenyl(1-6C)alkoxy and HET-1^c-(1-6C)alkoxy;
- 10 e) R^{8b} , $R^{8b}O(1-6C)alkyl$, hydroxy(1-6C)alkyl; and
- f) phenylamino, phenyl(1-6C)alkylamino, HET-1^c-amino, HET-1^c-(1-6C)alkylamino, (1-6C)alkylamino, N-phenyl-N-(1-6C)alkylamino and di(1-6C)alkylamino;

R^{5a} is selected from (1-6C)alkyl, phenyl (optionally substituted with 1 or 2 halo), HET-1^d, $R^{8c}CO-$, phenylcarbonyl (wherein the phenyl is optionally substituted with 1 or 2 halo), HET-1^d-carbonyl, (1-6C)alkoxycarbonyl, $R^{8c}NHCO-$, (1-6C)alkoxysulfonyl, 15 (1-6C)alkylaminosulfonyl, and (1-6C)alkylsulfonyl; wherein any (1-6C)alkyl group in any of the definitions of R^{5a} may optionally be substituted by a substituent selected from (1-6C)alkoxy, (1-6C)alkylS(O)p'- (wherein p' is 0, 1 or 2), amino, (1-6C)alkylamino, hydroxy, carboxy, phenyl (optionally substituted with 1 or 2 halo) and HET-1^f;

R^{6d} is selected from R^{8d} , halo and cyano;

R^{8a} , R^{8b} , R^{8c} or R^{8d} is (1-6C)alkyl;

25 ARI^a or ARI^b is phenyl or a 6,5-, or 6,6-bicyclic fused ring system, wherein both rings are carbocyclic and wherein at least the ring attached to the linking nitrogen atom is fully aromatic;

HET-1^a, HET-1^b, HET-1^c, HET-1^d or HET-1^f is 5- or 6-membered heteroaryl ring containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein any carbon atom may optionally be oxidised to a carbonyl group;

30 HET-2^a is a 5,5-, 5,6- 6,5- or 6,6-fused bicyclic, aromatic or partially saturated heterocyclic ring system, containing 1, 2, 3 or 4 heteroatoms independently selected from

O, N and S (provided there are no O-O, S-S or O-S bonds), in which the ring directly attached to the linking nitrogen atom is aromatic, and either ring of the bicyclic ring system may be carbocyclic;

HET-3^c is a partially or fully saturated 5- or 6- membered heterocyclic ring
 5 containing 1, 2, or 3 heteroatoms independently selected from O, N and S (provided there are no O-O, S-S or O-S bonds), wherein any sulfur atom may optionally be oxidised to an SO or SO₂ group, and wherein any carbon atom may optionally be oxidised to a carbonyl group; except that HET-3 is not 3-oxomorpholin-4-yl.

10 In a further aspect of the invention there is provided a compound of the formula (IB''') or a pharmaceutically-acceptable salt thereof, which is a compound of the formula (IB) as shown above, wherein:

R^{1b} is selected from AR1^a and HET-2^a; wherein R^{1b} is optionally substituted by 1 or 2 groups independently selected from R^{3a};

15 R^{3a} is selected from halo, R^{8a}, R^{8a}O-, R^{8a}OCO-, -C₆H_(13-a)F_a (wherein a is 1 to 13), and (1-6C)alkylthio;

R^{2b} is phenyl or 3-pyridyl; substituted in the position which is para to the linkage with the group -Y-NH- by a group selected from R^{4a} and optionally additionally substituted on a carbon atom in one or both of the positions which are meta to the the
 20 linkage with the group -Y-NH- (where possible) by 1 or 2 groups independently selected from R^{6c};

R^{4a} is selected from:

- a) phenyl;
- b) HET-3^c (wherein any available nitrogen atom is substituted by R^{5a} provided it is not
 25 thereby quaternised);
- d) phenoxy, HET-1^c-oxy, R^{8b}O-, phenyl(1-6C)alkoxy and HET-1^c-(1-6C)alkoxy;
- e) R^{8b} and hydroxy(1-6C)alkyl; and
- f) phenylamino and N-phenyl-N-(1-6C)alkylamino;

R^{5a} is selected from (1-6C)alkyl, benzyl, HET-1^d, R^{8c}CO-, phenylcarbonyl (wherein
 30 the phenyl is optionally substituted with 1 or 2 halo), HET-1^d-carbonyl, (1-6C)alkoxycarbonyl and (1-6C)alkylsulfonyl;

R^{6c} is selected from R^{8d}, halo and cyano;

R^{8a} , R^{8b} , R^{8c} and R^{8d} are independently selected from (1-6C)alkyl;

$AR1^a$ is selected from phenyl, naphthyl, indenyl and dihydroindenyl;

HET-1^c or HET-1^d is pyridyl, ;

HET-2^a is 1,3-benzodioxolyl;

5 HET-3^c is selected from morpholino, morpholinyl, piperazinyl and piperidinyl.

In a further aspect of the invention there is provided a compound of the formula (IB^{'''}), or a pharmaceutically-acceptable salt thereof, which is a compound of the formula (IB) as shown above, wherein

10 R^{1b} is selected from phenyl, 1-naphthyl and 2-naphthyl, optionally selected with 1, 2 or 3 fluoro;

R^{2b} is phenyl or pyridyl; substituted as described by a group R^{4a} and a group R^{6c}

R^{6c} is halo, preferably fluoro or chloro;

R^{4a} is morpholino, acylpiperazinyl or phenyl.

15 In a further aspect of the invention there is provided a compound of the formula (IB^{'''}), or a pharmaceutically-acceptable salt thereof, which is a compound of the formula (IB) as shown above, wherein

R^{1b} is selected from phenyl, 1-naphthyl and 2-naphthyl, optionally substituted with 1, 2 or 3 fluoro;

R^{2b} is phenyl or pyridyl; substituted as described by a group R^{4a} and a group R^{6c}

20 R^{6c} is halo, preferably fluoro or chloro;

R^{4a} is morpholino or acylpiperazinyl.

Preferred compounds of the invention are any one of the following, or their pharmaceutically acceptable salts and/or pro-drugs thereof:

25 N-[4-(4-morpholinyl)phenyl]-5-[[3-(phenylmethoxy)phenyl]amino]-1,3,4-oxadiazole-2-carboxamide;

5-[(4-chloro-3-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;

5-[(4-chlorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;

30 5-[(2,3-dihydro-1H-inden-5-yl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;

5-[(3-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;

- N-[4-(4-morpholinyl)phenyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
5-[(3,4-difluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
3-[[5-[[[4-(4-morpholinyl)phenyl]amino]carbonyl]-1,3,4-oxadiazol-2-yl]amino]-benzoic
5 acid, ethyl ester;
5-[(3-ethylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(3-chlorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
N-[4-(4-morpholinyl)phenyl]-5-(2-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
5-[(3-bromophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
10 5-[(3,4-dimethylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide;
5-[(3-chloro-4-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide,
5-[[3-(methylthio)phenyl]amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
15 carboxamide,
5-[(4-fluoro-3-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide,
5-[(3,5-dimethylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide,
20 5-[(2,5-difluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide,
5-[(2,3-difluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide,
5-[(2,4-difluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
25 carboxamide;
5-[[4-(1,1-dimethylethyl)phenyl]amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide;
5-[(4-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
carboxamide;
30 N-[4-(4-morpholinyl)phenyl]-5-(phenylamino)-1,3,4-oxadiazole-2-carboxamide;
N-[4-(4-morpholinyl)phenyl]-5-[[3-(trifluoromethyl)phenyl]amino]-1,3,4-oxadiazole-2-
carboxamide;

- 5-[(4-fluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-chlorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-chloro-4-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(2,3-dimethylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-fluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-methylphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 10 5-[(3-fluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[[2-(1-methylethyl)phenyl]amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(4-methoxyphenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 15 5-(1,3-benzodioxol-5-ylamino)-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-iodophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
N-[3-chloro-4-(4-morpholinyl)phenyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 20 N-[3-fluoro-4-(4-morpholinyl)phenyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
N-[3-chloro-4-(4-morpholinyl)phenyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
5-[(3,4-difluorophenyl)amino]-N-[3-methyl-4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-
- 25 carboxamide;
N-[3-methyl-4-(4-morpholinyl)phenyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
N-[3-fluoro-4-(4-morpholinyl)phenyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide,
- 30 N-[3-chloro-4-(4-morpholinyl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide,

- N-[3-fluoro-4-(4-morpholinyl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[3-methyl-4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5 N-[3-cyano-4-(4-morpholinyl)phenyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,5-difluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-morpholinyl)-3-pyridinyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
- 10 de;
- 5-[(2-fluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-fluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 15 5-[(3-methylphenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[[4-(1,1-dimethylethyl)phenyl]amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-chlorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 20 N-[6-(4-morpholinyl)-3-pyridinyl]-5-(2-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[5-chloro-6-(4-morpholinyl)-3-pyridinyl]-5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(4-chlorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[5-chloro-6-(4-morpholinyl)-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3,4-difluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 30 5-[(3-bromophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;

- 5-[(2,3-difluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[5-chloro-6-(4-morpholinyl)-3-pyridinyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(4-methylphenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(4-fluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-morpholinyl)-3-pyridinyl]-5-(phenylamino)-1,3,4-oxadiazole-2-carboxamide;
- 10 5-(1,3-benzodioxol-5-ylamino)-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-methoxyphenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,4-difluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 15 N-[4-(4-acetyl-1-piperazinyl)phenyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1-piperazinyl)phenyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
- 20 N-[4-(4-methyl-1-piperazinyl)phenyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1-piperazinyl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1-piperazinyl)-3-methylphenyl]-5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(2-fluorophenyl)amino]-N-[4-[4-(1-methylethyl)-1-piperazinyl]phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-methylphenyl)amino]-N-[4-(4-methyl-1-piperazinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 30 N-[4-(4-acetyl-1-piperazinyl)-3-methylphenyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;



- 4-[2-fluoro-4-[[[5-(1-naphthalenylamino)-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl]-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester;
- 5-[(2-fluorophenyl)amino]-N-[4-[4-(phenylmethyl)-1-piperazinyl]phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5 N-[4-(4-acetyl-1-piperazinyl)-3-methylphenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 4-[2-fluoro-4-[[[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl]-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester;
- 5-[(2-fluorophenyl)amino]-N-[4-(4-methyl-1-piperazinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 10 4-[4-[[[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl]-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester;
- 5-[(2-fluorophenyl)amino]-N-[4-[4-(4-pyridinyl)-1-piperazinyl]phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 15 5-(1-naphthalenylamino)-N-[4-(1-piperidinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-methylphenyl)amino]-N-[4-(1-piperidinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[4-(1-piperidinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-methyl-1-piperazinyl)-3-pyridinyl]-5-(1-naphthalenylamino)-1,3,4-oxadiazole-2-carboxamide;
- 20 5-[(3-methylphenyl)amino]-N-[6-(4-methyl-1-piperazinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-benzoyl-1-piperazinyl)-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-[4-(3-fluorobenzoyl)-1-piperazinyl]-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(2-fluorophenyl)amino]-N-[6-[4-(methylsulfonyl)-1-piperazinyl]-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-[4-(4-fluorobenzoyl)-1-piperazinyl]-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 30 5-[(2-fluorophenyl)amino]-N-[6-(4-methyl-1-piperazinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;



- 4-[5-[[[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]-2-pyridinyl]-1-piperazinecarboxylic acid, 1,1-dimethylethyl ester;
- N-[6-(4-acetyl-1-piperazinyl)-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(2-fluorophenyl)amino]-N-[6-[4-(4-pyridinylcarbonyl)-1-piperazinyl]-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-(6-phenoxy-3-pyridinyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[6-(phenylamino)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 10 N-(6-butoxy-3-pyridinyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[6-(3-pyridinyloxy)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[6-(4-hydroxybutyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide;
- 15 5-[(2-fluorophenyl)amino]-N-[4-(phenylmethoxy)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- N-(4-butylphenyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[4-(2-pyridinylmethoxy)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 20 5-[(2-fluorophenyl)amino]-N-[4-(methylphenylamino)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[1,1'-biphenyl]-4-yl-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide; and/or methyl {*trans*-4-[4-({[5-(pyridin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetate;
- 25 methyl (*trans*-4-[4-({[5-[(3-ethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl)acetate;
- methyl [*trans*-4-[4-({[5-[(3-benzyloxy)phenyl]amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl)acetate;
- methyl (*trans*-4-[4-({[5-[(4-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl)acetate;
- 30 methyl (*trans*-4-[4-({[5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl)acetate;



- methyl *trans*-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexanecarboxylate;
- methyl (*trans*-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetate;
- 5 {*trans*-4-{4-[(5-(pyridin-2-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl}acetic acid;
- (*trans*-4-{4-[(5-{(3-ethoxyphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
- {*trans*-4-{4-[(5-{(3-(benzyloxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl}acetic acid;
- 10 (*trans*-4-{4-[(5-{(4-chlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
- sodium (*trans*-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetate;
- 15 *trans*-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexanecarboxylic acid;
- (*trans*-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
- 5-{(2-fluorophenyl)amino}-N-{4-[4-(2-hydroxyethyl)cyclohexyl]phenyl}-1,3,4-oxadiazole-
- 20 2-carboxamide;
- N-{4-[*trans*-4-(2-amino-2-oxoethyl)cyclohexyl]phenyl}-5-{(3-ethoxyphenyl)amino}-1,3,4-oxadiazole-2-carboxamide;
- 5-{(2-fluorophenyl)amino}-N-(3-phenoxypropyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-{(3-methylphenyl)amino}-N-(3-phenoxypropyl)-1,3,4-oxadiazole-2-carboxamide;
- 25 5-{(3-(benzyloxy)phenyl)amino}-N-(3-phenoxypropyl)-1,3,4-oxadiazole-2-carboxamide;
- N-(4-cyclohexylphenyl)-5-{(2-fluorophenyl)amino}-1,3,4-oxadiazole-2-carboxamide;
- N-(4-cyclohexylphenyl)-5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazole-2-carboxamide;
- 5-{(3-(benzyloxy)phenyl)amino}-N-(4-cyclohexylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 30 N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-{(3-isopropoxyphenyl)amino}-1,3,4-oxadiazole-2-carboxamide;

- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-{{4-(methylthio)phenyl}amino}-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-{{4-(benzyloxy)phenyl}amino}-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(5-bromo-2,4-difluorophenyl)amino]-N-(4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-{{2,4-difluoro-5-(trifluoromethyl)phenyl}amino}-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- N-(6-morpholin-4-ylpyridin-3-yl)-5-{{3-(trifluoromethyl)phenyl}amino}-1,3,4-oxadiazole-2-carboxamide;
- 10 5-[(4-methylpyridin-2-yl)amino]-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-{{2-fluoro-5-(trifluoromethyl)phenyl}amino}-N-(4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 15 5-{{4-fluoro-3-(trifluoromethyl)phenyl}amino}-N-(4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-isopropoxyphenyl)amino]-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- N-(6-morpholin-4-ylpyridin-3-yl)-5-[(3-phenoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 20 N-(6-morpholin-4-ylpyridin-3-yl)-5-{{3-(trifluoromethoxy)phenyl}amino}-1,3,4-oxadiazole-2-carboxamide;
- N-(6-morpholin-4-ylpyridin-3-yl)-5-{{3-(pyridin-2-ylmethoxy)phenyl}amino}-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(3-ethoxyphenyl)amino]-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3-ethoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-(biphenyl-3-ylamino)-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 30 5-{{3-(benzyloxy)phenyl}amino}-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;



- ethyl 3-[(5-[(6-morpholin-4-ylpyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino]benzoate;
- tert-butyl 3-[(5-[(6-morpholin-4-ylpyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino]benzoate;
- 5 5-[(3-iodophenyl)amino]-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- N-(6-methoxypyridin-3-yl)-5-(1-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-(3-fluoro-4-morpholin-4-ylphenyl)-5-[(4-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 10 5-[(2,5-difluorophenyl)amino]-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-anilino-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- N-(3-fluoro-4-morpholin-4-ylphenyl)-5-[(3-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 15 N-(3-fluoro-4-morpholin-4-ylphenyl)-5-[(4-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,4-difluorophenyl)amino]-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-bromophenyl)amino]-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-
- 20 carboxamide;
- 5-[(4-chlorophenyl)amino]-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- N-(3-fluoro-4-morpholin-4-ylphenyl)-5-(2-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(3,4-difluorophenyl)amino]-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,3-difluorophenyl)amino]-N-(3-fluoro-4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- ethyl 3-[(5-[(3-fluoro-4-morpholin-4-ylphenyl)amino]carbonyl)-1,3,4-oxadiazol-2-
- 30 yl]amino]benzoate;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(4-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

- 85
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(2-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(3-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(2-methoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(4-methoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(2,5-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 10 5-anilino-N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-bromophenyl)amino]-N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(3-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 15 N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(4-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(2,6-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 20 N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(2-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(4-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-(2-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- 25 N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(3-methoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(4-tert-butylphenyl)amino]-N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 30 N-(5-chloro-6-morpholin-4-ylpyridin-3-yl)-5-[(2,3-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

- ethyl 3-[(5-[(5-chloro-6-morpholin-4-yl)pyridin-3-yl]amino)carbonyl]-1,3,4-oxadiazol-2-yl]amino]benzoate;
- N-(5-chloro-6-morpholin-4-yl)pyridin-3-yl)-5-(1-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- 5 N-(4-[cis-2,6-dimethylmorpholin-4-yl]phenyl)-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3,4-difluorophenyl)amino]-N-[4-[cis-2,6-dimethylmorpholin-4-yl]phenyl]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-
- 10 carboxamide;
- N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 15 N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-[(3-(benzyloxy)phenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-(1-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-[(2,4,5-trifluorophenyl)amino]-1,3,4-
- 20 oxadiazole-2-carboxamide;
- N-[4-(4-acetyl-1,4-diazepan-1-yl)phenyl]-5-[(2,5-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[2-(4-acetylpiperazin-1-yl)pyrimidin-5-yl]-5-[(3-(benzyloxy)phenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 N-(2-methyl-1,3-benzothiazol-5-yl)-5-(1-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(4-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(3-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 30 N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(2,5-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-anilino-1,3,4-oxadiazole-2-carboxamide;

- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(3-bromophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(3-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(4-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(2-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 10 N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(4-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-(2-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-(pyridin-2-ylamino)-1,3,4-oxadiazole-2-carboxamide;
- 15 ethyl 4-[[5-([6-(4-acetylpiperazin-1-yl)pyridin-3-yl]amino)carbonyl]-1,3,4-oxadiazol-2-yl]amino}benzoate;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-(1,3-benzodioxol-5-ylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(2,4-difluorophenyl)amino]-1,3,4-
- 20 oxadiazole-2-carboxamide;
- ethyl 3-[[5-([6-(4-acetylpiperazin-1-yl)pyridin-3-yl]amino)carbonyl]-1,3,4-oxadiazol-2-yl]amino}benzoate;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(4-tert-butylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(2,3-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-
- 30 carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[[3-(benzyloxy)phenyl]amino]-1,3,4-oxadiazole-2-carboxamide;

- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(3,5-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-(1-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- 5 N-[6-(4-acetylpiperazin-1-yl)pyridin-3-yl]-5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3-chlorophenyl)amino]-1,3,4-oxadiazole-2-
- 10 carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2,5-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-anilino-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3-bromophenyl)amino]-1,3,4-oxadiazole-2-
- 15 carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 20 N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-(2-naphthylamino)-1,3,4-oxadiazole-2-
- 25 carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-(1,3-benzodioxol-5-ylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 30 ethyl 3-{{5-([4-(4-acetylpiperazin-1-yl)phenyl]amino)carbonyl}-1,3,4-oxadiazol-2-yl}amino}benzoate;

- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-tert-butylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2,3-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[[3-(benzyloxy)phenyl]amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3,5-difluorophenyl)amino]-1,3,4-oxadiazole-2-
- 10 carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3-methoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 15 N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-methoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-anilino-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(4-chlorophenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-
- 20 oxadiazole-2-carboxamide;
- 5-[(2,3-difluorophenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[[3-(benzyloxy)phenyl]amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(2,4-difluorophenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,5-difluorophenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3,5-difluorophenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-
- 30 1,3,4-oxadiazole-2-carboxamide;
- N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-5-(1-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;

- N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-5-(2-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(4-tert-butylphenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- ethyl 3-[(5-[(6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino)benzoate;
- N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-5-(pyridin-2-ylamino)-1,3,4-oxadiazole-2-carboxamide;
- 10 5-[(2,6-difluorophenyl)amino]-N-{6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-anilino-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-methylphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-
- 15 carboxamide;
- 5-[(4-methylphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-fluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 20 5-[(4-fluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-methoxyphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-methoxyphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-
- 25 carboxamide;
- 5-[(4-methoxyphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-chlorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 30 5-[(3-chlorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;

- 5-[(4-chlorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,3-difluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(2,4-difluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,5-difluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 10 5-[(3,5-difluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-(1,3-benzodioxol-5-ylamino)-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-(1-naphthylamino)-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 15 5-(2-naphthylamino)-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- N-(6-thiomorpholin-4-ylpyridin-3-yl)-5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(4-tert-butylphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-
- 20 carboxamide;
- 5-[(3-bromophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-(pyridin-2-ylamino)-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[[3-(benzyloxy)phenyl]amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- ethyl 3-[(5-[[6-thiomorpholin-4-ylpyridin-3-yl)amino]carbonyl]-1,3,4-oxadiazol-2-yl)amino]benzoate;
- 5-[(2-fluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-
- 30 carboxamide;
- 5-[(3,4-difluorophenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;

- 5-[(3-methylphenyl)amino]-N-(6-thiomorpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(4-fluorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(3-chlorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-methoxyphenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,5-difluorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 10 5-anilino-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-bromophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-fluorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 15 5-[(4-methylphenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,6-difluorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 20 5-[(2-chlorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(4-chlorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-5-(2-naphthylamino)-1,3,4-oxadiazole-2-carboxamide;
- 25 N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-5-(pyridin-2-ylamino)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3-methoxyphenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 30 5-(1,3-benzodioxol-5-ylamino)-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;

- 5-[(2,4-difluorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,3-difluorophenyl)amino]-N-[6-(4-methylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;
- 5 N-(3-bromo-4-morpholin-4-ylphenyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(3-bromo-4-morpholin-4-ylphenyl)-5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(3-bromo-4-morpholin-4-ylphenyl)-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-
- 10 carboxamide;
- allyl 5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]-2-morpholin-4-ylbenzoate;
- allyl 5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]-2-morpholin-4-ylbenzoate;
- 15 5-[(2-fluorophenyl)amino]-N-{6-[4-(methoxyacetyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-{6-[4-(3-phenylpropanoyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- N-{6-[4-(2-fluorobenzoyl)piperazin-1-yl]pyridin-3-yl}-5-[(2-fluorophenyl)amino]-1,3,4-
- 20 oxadiazole-2-carboxamide;
- N-{6-[4-(4-cyanobenzoyl)piperazin-1-yl]pyridin-3-yl}-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-{6-[4-(4-methylbenzoyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(2-fluorophenyl)amino]-N-{6-[4-(2-methylbenzoyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-{6-[4-(3-methylbenzoyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-{6-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]pyridin-3-yl}-
- 30 1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[6-(4-glycoloylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide;

- 4-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperazine-1-carboxamide;
- 4-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}-N-phenylpiperazine-1-carboxamide;
- 5 N-benzyl-4-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperazine-1-carboxamide;
- phenyl 4-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperazine-1-carboxylate;
- ethyl 4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}piperazine-1-carboxylate;
- 10 benzyl 4-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperazine-1-carboxylate;
- ethyl 4-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperazine-1-carboxylate;
- 15 N-[4-(butylthio)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-(4-[(5-methoxy-1,3,4-thiadiazol-2-yl)amino]sulfonyl)phenyl)-1,3,4-oxadiazole-2-carboxamide;
- N-[4-(2,4-difluorophenoxy)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 20 N-[4-(1,4-dimethyl-6-oxo-1,4,5,6-tetrahydropyridazin-3-yl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-N-[4-(6-methyl-2-oxo-3,6-dihydro-2H-1,3,4-thiadiazin-5-yl)phenyl]-1,3,4-oxadiazole-2-carboxamide; and
- N-(4'-chlorobiphenyl-4-yl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 and/or
- N-cyclohexyl-5-[(3-(methylthio)phenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-cyclohexyl-5-[(4-ethoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- ethyl 4-[(5-[(3-bromophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}piperidine-1-carboxylate;
- 30 N-(1-benzylpiperidin-4-yl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-(2,3-dihydro-1H-inden-2-yl)-5-[(3-(2-oxo-1,3-oxazolidin-3-yl)phenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

- N-(2,3-dihydro-1H-inden-2-yl)-5-{{[2-(2-oxo-1,3-oxazolidin-3-yl)phenyl]amino}-1,3,4-oxadiazole-2-carboxamide;
ethyl 4-{{[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino]piperidine-1-carboxylate;
5 ethyl 4-{{[5-{{[4-(ethoxycarbonyl)phenyl]amino}-1,3,4-oxadiazol-2-yl]carbonyl}amino]piperidine-1-carboxylate;
ethyl 4-{{[5-(1-naphthylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino]piperidine-1-carboxylate;
tert-butyl 4-{{[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino]piperidine-1-carboxylate; and
10 N-(1-benzoylpiperidin-4-yl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
and/or
methyl (1-{5-[[5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]pyridin-2-yl}piperidin-4-yl)acetate;
15 (1-{5-[[5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]pyridin-2-yl}piperidin-4-yl)acetic acid;
methyl (trans-4-{4-[[5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl)acetate;
methyl (trans-4-{4-[[5-[(4-methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl)acetate;
20 methyl [trans-4-{4-[[5-[[3-(trifluoromethyl)phenyl]amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl]acetate;
methyl (trans-4-{4-[[5-[(3-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl)acetate;
25 methyl (trans-4-{4-[[5-[(2-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl)acetate;
methyl (trans-4-{4-[[5-[(2-isopropylphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl)acetate;
methyl [trans-4-{4-[[5-[[4-(benzyloxy)phenyl]amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl]acetate;
30 methyl (trans-4-{4-[[5-[(3-methylphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]phenyl}cyclohexyl)acetate;

- methyl (trans-4-{4-[(5-{(4-isopropylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl [trans-4-(4-{[(5-{[2-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- 5 methyl {trans-4-(4-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl (trans-4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl [trans-4-(4-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- 10 methyl [trans-4-(4-{[(5-{[2-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl (trans-4-{4-[(5-{(3-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- 15 methyl (trans-4-{4-[(5-{(2-methoxyphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl (trans-4-{4-[(5-{(3-anilinophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl [trans-4-(4-{[(5-{[3-(aminocarbonyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- 20 methyl (trans-4-{4-[(5-{(3-benzoylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl [trans-4-(4-{[(5-{[3-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- 25 methyl (trans-4-{4-[(5-{(3-phenoxyphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl [trans-4-(4-{[(5-{[2-(benzyloxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- methyl [trans-4-(4-{[(5-{[3-(benzoylamino)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
- 30 methyl {trans-4-[4-([5-(4-[(methylsulfonyl)oxy]phenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;

- methyl {trans-4-(4-{{5-{{3-(phenylsulfonyl)phenyl}amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl}acetate;
- methyl {trans-4-[4-{{5-{{3-[(methylsulfonyl)oxy]phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl}acetate;
- 5 methyl {trans-4-[4-{{5-{{2-[(methylsulfonyl)oxy]phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl}acetate;
- methyl {trans-4-[4-{{5-{{3-[(aminocarbonyl)amino]phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl}acetate;
- methyl {trans-4-[4-{{5-{{3-[(methoxycarbonyl)amino]phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl}acetate;
- 10 methyl (trans-4-{4-[[5-[(4-morpholin-4-yl)phenyl]amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl}acetate;
- methyl (trans-4-{4-[[5-[(2-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl}acetate;
- 15 methyl (trans-4-{4-[[5-[(4-anilinophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl}acetate;
- methyl [trans-4-(4-{{5-{{4-(phenylsulfonyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl]acetate;
- methyl [trans-4-(4-{{5-{{2-(phenylsulfonyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl]acetate;
- 20 methyl [trans-4-(4-{{5-{{4-(ethylsulfonyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl]acetate;
- (trans-4-{4-[[5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 25 (trans-4-{4-[[5-[(4-methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- [trans-4-(4-{{5-{{3-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino)phenyl}cyclohexyl]acetic acid;
- (trans-4-{4-[[5-[(3-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 30 (trans-4-{4-[[5-[(2-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;

- (trans-4-{4-[(5-[(2-isopropylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-[(4-(benzyloxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 5 (trans-4-{4-[(5-[(3-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(4-isopropylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-[(2-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 10 [trans-4-(4-{[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(4-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 15 [trans-4-(4-{[(5-[(4-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-[(2-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(3-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 20 [trans-4-(4-{[(5-[(3-(aminocarbonyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(3-benzoylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 25 [trans-4-(4-{[(5-[(3-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(3-phenoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 30 [trans-4-(4-{[(5-[(2-(benzyloxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-[(3-(benzoylamino)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

- {trans-4-[4-({[5-({4-[(methylsulfonyl)oxy]phenyl}amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 [trans-4-(4-({[5-({3-(phenylsulfonyl)phenyl}amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
- 5 {trans-4-[4-({[5-({3-[(methylsulfonyl)oxy]phenyl}amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 {trans-4-[4-({[5-({2-[(methylsulfonyl)oxy]phenyl}amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 (trans-4-{4-({[5-[(2-methoxyphenyl)amino]-1,3,4-oxadiazol-2-
- 10 yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 {trans-4-[4-({[5-({3-[(aminocarbonyl)amino]phenyl}amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 {trans-4-[4-({[5-({3-[(methoxycarbonyl)amino]phenyl}amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
- 15 (trans-4-{4-({[5-[(4-morpholin-4-yl)phenyl]amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 [trans-4-(4-({[5-({4-(phenylsulfonyl)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 [trans-4-(4-({[5-({2-(phenylsulfonyl)phenyl]amino)-1,3,4-oxadiazol-2-
- 20 yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 (trans-4-{4-({[5-[(2-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
 (trans-4-{4-({[5-[(4-anilinophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
- 25 (trans-4-{4-({[5-[(3-anilinophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid; and
 [trans-4-(4-({[5-({4-(ethylsulfonyl)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid; and/or
 methyl 1-[(trans-4-{4-({[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-
- 30 yl]carbonyl}amino)phenyl]cyclohexyl)acetyl]piperidine-4-carboxylate;
 methyl N-[(trans-4-{4-({[5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl)acetyl]-2-methylalaninate;

- methyl 1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-L-prolinate;
methyl 1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-D-prolinate;
- 5 methyl (4R)-1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-4-hydroxy-L-prolinate;
5-[(2-fluorophenyl)amino]-N-[4-(trans-4-{2-[(2-hydroxy-1,1-dimethylethyl)amino]-2-oxoethyl}cyclohexyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-fluorophenyl)amino]-N-[4-(trans-4-{2-oxo-2-[(3aR,6aS)-tetrahydro-5H-
- 10 [1,3]dioxolo[4,5-c]pyrrol-5-yl]ethyl)cyclohexyl]phenyl]-1,3,4-oxadiazole-2-carboxamide;
N-(4-{trans-4-[2-(dimethylamino)-2-oxoethyl]cyclohexyl}phenyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
5-[(2-fluorophenyl)amino]-N-[4-(trans-4-{2-[(2-hydroxyethyl)(methyl)amino]-2-oxoethyl}cyclohexyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 15 N-[4-(trans-4-{2-[(2,3-dihydroxypropyl)amino]-2-oxoethyl}cyclohexyl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]piperidine-4-carboxylic acid;
N-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-
- 20 yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-2-methylalanine;
1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-L-proline;
1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-D-proline;
- 25 (4R)-1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]-4-hydroxy-L-proline;
methyl (trans-4-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)cyclohexyl]acetate;
(trans-4-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-
- 30 yl)carbonyl]amino}pyridin-2-yl)cyclohexyl]acetic acid;
[trans-4-(4-{[(5-{[4-(2-hydroxyethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;

- (trans-4-{4-[(5-{(3-ethynylphenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
[trans-4-(4-{[(5-{[4-(cyanomethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
- 5 {trans-4-[4-[(5-(biphenyl-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
(trans-4-{4-[(5-{[4-(tert-butoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
(trans-4-{4-[(5-{[4-phenoxyphenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 10 [trans-4-(4-{[(5-{[4-(1H-1,2,4-triazol-1-yl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
(trans-4-{4-[(5-{[4-benzylphenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 15 [trans-4-(4-{[(5-{[3-(acetylamino)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
(trans-4-{4-[(5-{[3-chlorophenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
(trans-4-{4-[(5-{[4-ethynylphenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 20 (trans-4-{4-[(5-{[3-bromophenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
[trans-4-(4-{[(5-{[3-(difluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
- 25 [trans-4-(4-{[(5-{[3-(methoxymethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
[trans-4-(4-{[(5-{[3-(2-methylpyrimidin-4-yl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
(trans-4-{4-[(5-{[4-cyclohexylphenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 30 methyl {trans-4-[4-[(5-(pyrimidin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetate;

- methyl {trans-4-[4-({[5-[(1-methyl-1H-pyrazol-3-yl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetate;
methyl {trans-4-[4-({[5-(pyridin-3-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetate;
5 methyl {trans-4-[4-({[5-(pyridin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetate;
methyl {trans-4-[4-({[5-(pyrazin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetate;
methyl {trans-4-[4-({[5-(1,3-thiazol-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetate;
10 {trans-4-[4-({[5-(pyrimidin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
(trans-4-{4-[(5-[(1-methyl-1H-pyrazol-3-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
15 {trans-4-[4-({[5-(pyridin-3-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
{trans-4-[4-({[5-(pyridin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
{trans-4-[4-({[5-(pyrazin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
20 {trans-4-[4-({[5-(1,3-thiazol-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
(trans-4-{4-[(5-[(1-methyl-1H-pyrazol-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid;
25 {trans-4-[4-({[5-(pyrimidin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl}acetic acid;
trans-4-{4-[(5-[(2-methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexanecarboxylic acid;
trans-4-(4-{[(5-[(3-(hydroxymethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexanecarboxylic acid;
30 trans-4-(4-{[(5-[(2-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexanecarboxylic acid;

- trans-4-(4-((5-([2-(trifluoromethoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-((5-([3-(trifluoromethyl)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- 5 trans-4-(4-([5-([4-methoxyphenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-((5-([4-(trifluoromethoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-([5-([4-(trifluoromethyl)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- 10 trans-4-(4-([5-([2-(benzyloxy)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-([5-([3-(trifluoromethoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-([5-([3-(methoxymethyl)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- 15 trans-4-(4-([5-([3-(phenylsulfonyl)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-([5-([3-anilinophenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- 20 trans-4-(4-([5-([4-anilinophenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-([5-([2-([methylsulfonyl]oxy)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- 25 trans-4-(4-([5-([3-([methylsulfonyl]oxy)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- trans-4-(4-([5-([4-([methylsulfonyl]oxy)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl)cyclohexanecarboxylic acid;
- ethyl trans-4-(4-([5-([2,4,5-trifluorophenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenoxy)cyclohexanecarboxylate;
- 30 trans-4-(4-([5-([2,4,5-trifluorophenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenoxy)cyclohexanecarboxylic acid;

- methyl (trans-4-{4-[(5-{(3-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- methyl (trans-4-{4-[(5-{(4-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- 5 methyl (trans-4-{4-[(5-{(2,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- methyl (trans-4-{4-[(5-{(2,6-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- methyl (trans-4-{4-[(5-{(2,5-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- 10 methyl (trans-4-{4-[(5-{(2,3-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- methyl (trans-4-{4-[(5-{(2,4,6-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- 15 methyl (trans-4-{4-[(5-{(3,4-dimethoxyphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- methyl (trans-4-{4-[(5-{(3-{(1R)-2-methoxy-1-methylethoxy}phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- methyl (trans-4-{4-[(5-{(3-{(1S)-2-methoxy-1-methylethoxy}phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- 20 methyl (trans-4-{4-[(5-{(4-fluoro-3-methoxyphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate;
- (trans-4-{4-[(5-{(3-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 25 (trans-4-{4-[(5-{(4-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(2,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(2,6-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 30 (trans-4-{4-[(5-{(2,5-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid;

- (trans-4-{4-[(5-[(2,3-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(2,4,6-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
5 (trans-4-{4-[(5-[(3,4-dimethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
{trans-4-[4-[(5-[(3-[(1S)-2-methoxy-1-methylethoxy]phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
{trans-4-[4-[(5-[(3-[(1R)-2-methoxy-1-methylethoxy]phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
10 (trans-4-{4-[(5-[(4-fluoro-3-methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
1-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
15 1-{5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
1-{5-[(5-[(4-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
20 methyl 4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)butanoate;
methyl ({4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)thio)acetate;
25 methyl 3-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)adamantane-1-carboxylate;
methyl 3-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentanecarboxylate;
methyl 4'-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}biphenyl-4-carboxylate;
30 ethyl {4'-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}biphenyl-3-yl)acetate;

- methyl [(1,4-trans)-4'-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]-1,1'-bi(cyclohexyl)-4-yl]acetate;
ethyl 3-{2-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]-2,3-dihydro-1H-inden-5-yl}propanoate;
- 5 ethyl ((1R,3R)-3-{4-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetate;
ethyl (3-{4-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclopentyl)acetate;
methyl 4-{3-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]propoxy}benzoate;
- 10 methyl 3-{3-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]propoxy}benzoate;
methyl cis-3-{4-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclobutanecarboxylate;
- 15 ethyl cis-3-{4-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]benzyl}cyclobutanecarboxylate;
4-{4-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}butanoic acid;
3-{4-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}adamantane-1-carboxylic acid;
- 20 3-{4-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclopentanecarboxylic acid;
4'-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]biphenyl-4-carboxylic acid;
- 25 {4'-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]biphenyl-4-yl}acetic acid;
{4'-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]biphenyl-3-yl}acetic acid;
[(1,4-trans)-4'-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]-1,1'-bi(cyclohexyl)-4-yl]acetic acid;
- 30 3-{2-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]-2,3-dihydro-1H-inden-5-yl}propanoic acid;

- (3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)propyl)malonic acid;
- (4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}cyclohexyl)phenyl)acetic acid;
- 5 (3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- ((3R)-3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentyl)acetic acid;
- ((3S)-3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentyl)acetic acid;
- 10 4-{3-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}propoxy}benzoic acid;
- 3-{3-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}propoxy}benzoic acid;
- 15 3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclobutanecarboxylic acid;
- 3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}benzyl)cyclobutanecarboxylic acid;
- 5-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}pentanoic acid;
- 20 N-{4-[trans-4-(2-amino-2-oxoethyl)cyclohexyl]phenyl}-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N-{4-[trans-4-(cyanomethyl)cyclohexyl]phenyl}-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 25 5-[(2-fluorophenyl)amino]-N-[4-(propylsulfonyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- methyl 4'-{[5-({[4-(4-acetylpiperazin-1-yl)phenyl]amino}carbonyl)-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylate;
- N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-chloro-2-fluoro-5-hydroxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 30 ethyl (5-{[5-({[4-(4-acetylpiperazin-1-yl)phenyl]amino}carbonyl)-1,3,4-oxadiazol-2-yl]amino}-2-chloro-4-fluorophenoxy)acetate;

- methyl 3'-{[5-({[4-(4-acetylpiperazin-1-yl)phenyl]amino}carbonyl)-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylate;
- 3'-{[5-({[4-(4-acetylpiperazin-1-yl)phenyl]amino}carbonyl)-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylic acid;
- 5 2-({4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)sulfonyl)-2-methylpropanoic acid;
- (trans-4-{4-[(5-[(3-methylisoxazol-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- {trans-4-[4-[(5-(1,3,4-thiadiazol-2-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
- 10 (trans-4-{4-[(5-[(2,5-dimethylpyridin-4-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- (trans-4-{4-[(5-[(5,6-dimethylpyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- 15 (trans-4-{4-[(5-[(3-methylpyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- (trans-4-{4-[(5-[(5-bromopyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- (trans-4-{4-[(5-[(2-methoxypyrimidin-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- 20 (trans-4-{4-[(5-[(5-benzyl-1,3,4-oxadiazol-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- (trans-4-{4-[(5-[(3-methyl-1,2,4-thiadiazol-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- 25 (trans-4-{4-[(5-[(3-ethyl-1,2,4-thiadiazol-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- (trans-4-{4-[(5-[(2,6-dimethylpyrimidin-4-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- (trans-4-{4-[(5-[(1,3-dimethyl-1H-pyrazol-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;
- 30 (trans-4-{4-[(5-[(6-chloropyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexylacetic acid;

- (trans-4-{4-[(5-[(3,5-dimethylpyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-{4-(trifluoromethyl)-1,3-thiazol-2-yl}amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;
- 5 methyl (trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate;
(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
methyl (trans-4-{4-[(5-[(3-chloro-4-cyanophenyl)amino]-1,3,4-oxadiazol-2-
- 10 yl)carbonyl]amino}phenyl)cyclohexyl)acetate;
methyl [trans-4-(4-{[(5-{4-cyano-3-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
methyl (trans-4-{4-[(5-[(3,4-dicyanophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate;
- 15 methyl (trans-4-{4-[(5-[(4-cyano-3-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate;
methyl [trans-4-(4-{[(5-{4-cyano-3-(methylthio)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate;
(trans-4-{4-[(5-[(3-chloro-4-cyanophenyl)amino]-1,3,4-oxadiazol-2-
- 20 yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-{4-cyano-3-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;
(trans-4-{4-[(5-[(3,4-dicyanophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 25 (trans-4-{4-[(5-[(4-cyano-3-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-(4-{[(5-{4-cyano-3-(methylthio)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;
(trans-4-{4-[(5-[(3-chloro-4-fluorophenyl)amino]-1,3,4-oxadiazol-2-
- 30 yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(trans-4-{4-[(5-[(3,5-dimethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

- [trans-4-(4-{{(5-{{(4-fluoro-3-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- [trans-4-(4-{{(5-{{(3-fluoro-5-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- 5 (trans-4-(4-{{(5-{{(2-methyl-1,3-benzothiazol-6-yl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- (trans-4-(4-{{(5-{{(3-cyano-4-methylphenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- (trans-4-(4-{{(5-{{(3-cyano-4-fluorophenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- 10 (trans-4-(4-{{(5-{{(5-chloro-2-fluorophenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- [trans-4-(4-{{(5-{{(4-methoxy-3-(trifluoromethyl)phenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- 15 (trans-4-(4-{{(5-{{(4-chloro-3-cyanophenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- (trans-4-(4-{{(5-{{(4-chloro-3-methoxyphenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- [trans-4-(4-{{(5-{{(3-methoxy-5-(trifluoromethyl)phenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- 20 (trans-4-(4-{{(5-{{(4-chloro-3-fluorophenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- (trans-4-(4-{{(5-{{(4-cyano-2-fluorophenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- 25 (trans-4-(4-{{(5-{{(3-fluoro-4-methoxyphenyl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- (trans-4-(4-{{(5-{{(2-methyl-1,3-benzothiazol-5-yl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- (trans-4-(4-{{(5-{{(2-oxo-2,3-dihydro-1H-benzimidazol-5-yl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
- 30 (trans-4-(4-{{(5-{{(2-methyl-1H-benzimidazol-6-yl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;

- {trans-4-[4-({[5-(quinolin-3-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
[trans-4-(4-({[5-({[6-(4-chlorophenoxy)pyridin-3-yl]amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
5 (trans-4-{4-[(5-[(6-methoxypyridin-3-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
(trans-4-{4-[(5-[(4-methylpyridin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
(trans-4-{4-[(5-[(5-methylpyridin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
10 (trans-4-{4-[(5-[(2-methoxypyridin-3-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
(trans-4-{4-[(5-[(5-fluoropyridin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
15 [trans-4-(4-({[5-({[6-(4-methoxyphenoxy)pyridin-3-yl]amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl)cyclohexyl}acetic acid;
{trans-4-[4-({[5-({[6-(3-chlorobenzyl)oxy]pyridin-3-yl]amino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
{trans-4-[4-({[5-(1H-pyrrolo[2,3-b]pyridin-5-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid;
20 (trans-4-{4-[(5-[(6-cyanopyridin-3-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
(trans-4-{4-[(5-[(5-cyanopyridin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}acetic acid;
25 rac- ethyl 2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}propanoate;
rac- ethyl 2-(trans-4-{4-[(5-[(4-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}propanoate;
rac- ethyl 2-(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}propanoate;
30 rac- 2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl}propanoic acid;

- rac- 2-(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)propanoic acid;
- rac- 2-(trans-4-{4-[(5-[(4-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)propanoic acid;
- 5 (trans-4-{4-[(5-[(4-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- (trans-4-{4-[(5-[(4-bromophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- (trans-4-{4-[(5-[(4-ethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 10 (trans-4-{4-[(5-[(4-tert-butylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- (trans-4-{4-[(5-[(2-ethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 15 (trans-4-{4-[(5-[(2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- {trans-4-(4-{[(5-[(4-aminosulfonyl)phenyl]amino)-1,3,4-oxadiazol-2-yl}carbonyl]amino}phenyl}cyclohexyl)acetic acid;
- (trans-4-{4-[(5-[(2-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- 20 (trans-4-{4-[(5-[(3-methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid;
- {trans-4-[4-[(5-[(4-[(trifluoromethyl)thio]phenyl]amino)-1,3,4-oxadiazol-2-yl}carbonyl]amino]phenyl}cyclohexyl)acetic acid;
- 25 5-[(3,4-difluorophenyl)amino]-N-(4-phenoxyphenyl)-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3,4-difluorophenyl)amino]-N-[4-(2-methoxyphenoxy)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(3,4-difluorophenyl)amino]-N-[4-(4-methylphenoxy)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 30 5-[(3,4-difluorophenyl)amino]-N-[4-(3-methoxyphenoxy)phenyl]-1,3,4-oxadiazole-2-carboxamide;

- N*-(3,5-dichloro-4-morpholin-4-ylphenyl)-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N*-{6-[4-(cyclopropylmethyl)piperazin-1-yl]pyridin-3-yl}-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 5 5-[(2-fluorophenyl)amino]-*N*-{6-[4-(2-methoxyethyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-*N*-[4-(2-morpholin-4-ylethoxy)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2-fluorophenyl)amino]-*N*-[4-(3-hydroxypropoxy)phenyl]-1,3,4-oxadiazole-2-
- 10 carboxamide;
- N*-[4-(2-ethoxyethoxy)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- N*-{4-[2-(diethylamino)-1,1-dimethyl-2-oxoethoxy]phenyl}-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;
- 15 5-[(2-fluorophenyl)amino]-*N*-(4-pyrrolidin-1-ylphenyl)-1,3,4-oxadiazole-2-carboxamide;
- methyl trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl} cyclohexanecarboxylate;
- trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl} cyclohexanecarboxylic acid;
- 20 methyl 6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}piperidin-1-yl}nicotinate;
- 6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}piperidin-1-yl}nicotinic acid;
- 6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}piperidin-1-
- 25 yl}nicotinamide;
- (6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}piperidin-1-yl)pyridin-3-yl}acetic acid;
- 5-[(3-(anilinoacetyl)phenyl)amino]-*N*-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide;
- 30 *N*-{4-[trans-4-(aminomethyl)cyclohexyl]phenyl}-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

- tert-butyl [(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)methyl]carbamate;
methyl (1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetate;
5 (1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetic acid;
methyl [1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate;
[1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
10 (2R)-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)-propanoic acid;
(2S)-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoic acid;
15 (2R)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoic acid;
(2S)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoic acid;
(2R)-2-(trans-4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)-propanoic acid;
20 (2S)-2-(trans-4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)-propanoic acid; and/or
ethyl (2R)-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate;
25 ethyl (2S)-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate;
ethyl (2R)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate;
ethyl (2S)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate;
30 ethyl (2R)-2-(trans-4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate;

- ethyl (2S)-2-(trans-4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoate; and/or
- methyl 2-methyl-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoate;
- 5 2-methyl-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
- 1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
- 1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-
- 10 methylpiperidine-4-carboxylic acid;
- 4-methyl-1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 4-methyl-1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 15 4-methyl-1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 1-(5-{[(5-{[4-(4-chlorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
- methyl 1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-
- 20 yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylate;
- methyl 1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylate;
- methyl 4-methyl-1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate;
- 25 methyl 4-methyl-1-(5-{[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate;
- methyl 4-methyl-1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate;
- methyl 1-(5-{[(5-{[4-(4-chlorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-
- 30 yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylate;
- 1-(5-{[(5-{[4-(4-chlorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;

- methyl 1-(5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylate;
- methyl 1-(5-[(5-[(4-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylate;
- 5 methyl 1-(5-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylate;
- (*trans*-4-{4-[(5-[(2-chloro-4-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(4-methoxy-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 10 (*trans*-4-{4-[(5-[(4-fluoro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- [*trans*-4-{4-[(5-[(4-fluoro-2-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;
- 15 (*trans*-4-{4-[(5-[(2,4-dichlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(2-chloro-4-cyanophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(4-cyano-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 20 (*trans*-4-{4-[(5-[(4-chloro-2,5-dimethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(4-cyano-2-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 25 (*trans*-4-{4-[(5-[(2-chloro-4-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(4-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(2,3-dichlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 30 (*trans*-4-{4-[(5-[(2,5-dichlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

- methyl 1-(5-[(5-[(4-(4-chlorophenoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylate;
[1-(5-[(5-[(4-(4-chlorophenoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetic acid;
- 5 methyl [1-(5-[(5-[(4-(4-chlorophenoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetate;
rac-ethyl 2-(trans-4-(2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexyl)propanoate;
rac-ethyl 2-(trans-4-(3-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexyl)propanoate;
- 10 rac-2-(trans-4-(2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexyl)propanoic acid;
rac-2-(trans-4-(3-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexyl)propanoic acid;
- 15 ethyl cis-4-(4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)-phenoxy)cyclohexanecarboxylate;
cis-4-(4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)-phenoxy)cyclohexanecarboxylic acid;
(1-(5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl)acetic acid;
- 20 [1-(5-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetic acid;
methyl (1-(5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl)acetate;
- 25 methyl [1-(5-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetate;
1-(5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylic acid;
1-(5-[(5-[(4-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylic acid;
- 30 1-(5-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylic acid;

- (*trans*-4-{4-[(5-{(2-bromo-4-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[*trans*-4-(4-{[(5-{(2-chloro-5-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
5 [*trans*-4-(4-{[(5-{(2-bromo-4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[*trans*-4-(4-{[(5-{(2-chloro-4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(*trans*-4-{4-[(5-{(4-bromo-2-chlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
10 (*trans*-4-{4-[(5-{(3-chloro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(*trans*-4-{4-[(5-{(5-bromo-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
15 (*trans*-4-{4-[(5-{(2-chloro-5-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(*trans*-4-{4-[(5-{(2-chloro-5-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
(*trans*-4-{4-[(5-{(5-chloro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
20 ethyl 2-(3-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclobutyl)propanoate;
2-(3-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclobutyl)propanoic acid;
25 methyl 1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylate;
methyl 1-(3-methyl-5-{[(5-{(4-trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate;
methyl 1-(3-methyl-5-{[(5-{(4-trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate;
30 methyl 1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylate;



- methyl (1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]acetate;
- methyl [1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate;
- 5 methyl [1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate;
- methyl (1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]acetate;
- ethyl 2-(1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]propanoate;
- 10 ethyl 2-[1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]propanoate;
- ethyl 2-(1-{3-methyl-5-{[(5-{[2,4,5-trifluorophenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridine-2-yl)piperidin-4-yl]propanoate;
- 15 ethyl 2-(1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]propanoate;
- (1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]acetic acid;
- [1-(3-methyl-5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- 20 [1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- (1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]acetic acid;
- 25 1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylic acid;
- 1-(3-methyl-5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 30 1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylic acid;

- 2-(1-{3-methyl-5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)propanoic acid;
methyl (1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl)acetate;
- 5 1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
[1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- 10 methyl 1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate;
methyl [1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate;
- 15 methyl 1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylate;
(1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl)acetic acid;
(4-methyl-1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
- 20 [4-methyl-1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid; and/or
[1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl]acetic acid.
- 25 *In a further aspect of the invention, there is provided any one or more of the following compounds, or pharmaceutically-acceptable salts or pro-drugs thereof:*
N-[4-(4-morpholinyl)phenyl]-5-[[3-(phenylmethoxy)phenyl]amino]-1,3,4-oxadiazole-2-carboxamide;
- 30 5-[(3,4-difluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
- 5-[(2,4-difluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;

5-[(2-fluorophenyl)amino]-N-[4-(4-morpholinyl)phenyl]-1,3,4-oxadiazole-2-carboxamide;
and

5-[(2-fluorophenyl)amino]-N-[6-(4-morpholinyl)-3-pyridinyl]-1,3,4-oxadiazole-2-carboxamide.

5 In a further aspect of the invention, there is provided any one or more of the following compounds, or pharmaceutically-acceptable salts or pro-drugs thereof:

N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2,4-difluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

10 N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(3-chlorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

15 N-[4-(4-acetyl-1-piperazinyl)phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide;

N-[6-(4-acetyl-1-piperazinyl)-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide; and

20 5-[(2-fluorophenyl)amino]-N-{6-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]pyridin-3-yl}-1,3,4-oxadiazole-2-carboxamide.

In a further aspect of the invention, there is provided any one or more of the following compounds, or pharmaceutically-acceptable salts or pro-drugs thereof:

sodium (trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate;

25 {trans-4-[4-[(5-(pyridin-2-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

30 (trans-4-{4-[(5-[(2,4,6-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

[trans-4-(4-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

- (trans-4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
 [trans-4-(4-{[(5-{(4-(trifluoromethoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;
- 5 (trans-4-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)cyclohexyl)acetic acid;
 trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexanecarboxylic acid;
 (2R)-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)-propanoic acid;
- 10 (2S)-2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
 (2R)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
- 15 (2S)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
 trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenoxy)cyclohexanecarboxylic acid;
 1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-
- 20 yl)piperidine-4-carboxylic acid;
 (1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
 methyl (1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetate;
- 25 ((3S)-3-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentyl)acetic acid;
 ((3R)-3-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentyl)acetic acid;
- 3-4-{[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}benzyl)cyclobutanecarboxylic acid;
- 30 [trans-4-(4-{[(5-{(6-(4-chlorophenoxy)pyridin-3-yl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid;

- [trans-4-(4-{{(5-{{(6-(4-methoxyphenoxy)pyridin-3-yl}amino)}-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 {trans-4-[4-{{(5-(quinolin-3-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 5 (trans-4-[4-{{(5-{{(5-bromopyrazin-2-yl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 {trans-4-[4-{{(5-(pyridin-3-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 (trans-4-[4-{{(5-{{(6-methoxypyridin-3-yl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 10 {trans-4-[4-{{(5-{{(4-cyano-3-(trifluoromethyl)phenyl)amino)}-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid; and/or selected from
 2-methyl-2-(trans-4-[4-{{(5-{{(2,4,5-trifluorophenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}propanoic acid;
 15 1-(5-{{(5-{{(3,4-difluorophenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
 1-(5-{{(5-{{(4-ethylphenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
 4-methyl-1-(5-{{(5-{{(4-(trifluoromethoxy)phenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
 20 4-methyl-1-(5-{{(5-{{(2,4,5-trifluorophenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
 4-methyl-1-(5-{{(5-{{(4-(trifluoromethyl)phenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
 25 1-(5-{{(5-{{(4-(4-chlorophenoxy)phenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
 1-(5-{{(5-{{(4-(4-chlorophenoxy)phenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
 [1-(5-{{(5-{{(4-(4-chlorophenoxy)phenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
 30 rac-2-(trans-4-[2-chloro-4-{{(5-{{(2,4,5-trifluorophenyl)amino)}-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}propanoic acid;

- rac-2-(trans-4-{3-chloro-4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
- cis-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenoxy}cyclohexanecarboxylic acid;
- 5 1-(5-{[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
- [1-(5-{[(5-{(4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
- 10 1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 1-(5-{[(5-{(4-(trifluoromethoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 1-(5-{[(5-{(4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 15 (trans-4-{4-[(5-{(2-chloro-4-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(4-methoxy-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(4-fluoro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 20 [trans-4-{4-[(5-{(4-fluoro-2-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(2,4-dichlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 25 (trans-4-{4-[(5-{(2-chloro-4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(4-cyano-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (trans-4-{4-[(5-{(4-chloro-2,5-dimethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 30 (trans-4-{4-[(5-{(4-cyano-2-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

- (*trans*-4-{4-[(5-[(2-chloro-4-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(4-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 5 (*trans*-4-{4-[(5-[(2,3-dichlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(2,5-dichlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(2-bromo-4-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 10 [*trans*-4-(4-{[(5-[(2-chloro-5-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- [*trans*-4-(4-{[(5-[(2-bromo-4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 15 [*trans*-4-(4-{[(5-[(2-chloro-4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(4-bromo-2-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(3-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 20 (*trans*-4-{4-[(5-[(5-bromo-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(2-chloro-5-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 25 (*trans*-4-{4-[(5-[(2-chloro-5-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- (*trans*-4-{4-[(5-[(5-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
- 2-(3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclobutyl)propanoic acid;
- 30 (1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl)acetic acid;

- [1-(3-methyl-5-{{(5-{{(4-(trifluoromethoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- [1-(3-methyl-5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- 5 (1-{5-{{(5-{{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino)-3-methylpyridin-2-yl}piperidin-4-yl]acetic acid;
- 1-{5-{{(5-{{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino)-3-methylpyridin-2-yl}piperidine-4-carboxylic acid;
- 1-(3-methyl-5-{{(5-{{(4-(trifluoromethoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 10 1-(3-methyl-5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 1-{5-{{(5-{{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino)-3-methylpyridin-2-yl}piperidine-4-carboxylic acid;
- 15 2-(1-{3-methyl-5-{{(5-{{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl}propenoic acid;
- 1-{5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 1-(5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
- 20 [1-(5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- (1-{5-{{(5-{{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidin-4-yl]acetic acid;
- 25 (4-methyl-1-{5-{{(5-{{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
- [4-methyl-1-(5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)piperidin-4-yl]acetic acid; and/or
- [1-(5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidin-4-yl]acetic acid.
- 30

In a further aspect of the invention, there is provided any one or more of the following compounds, or pharmaceutically-acceptable salts or pro-drugs thereof:

- sodium (trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate;
(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
5 (trans-4-{4-[(5-[(2,4,6-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-{4-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
[trans-4-{4-[(5-[(4-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
10 (trans-4-{5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)cyclohexyl)acetic acid;
(2R)-2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
15 (2S)-2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
(2R)-2-(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
(2S)-2-(trans-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
20 ((3S)-3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentyl)acetic acid;
((3R)-3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclopentyl)acetic acid;
25 3-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}benzyl)cyclobutanecarboxylic acid;
[trans-4-{4-[(5-[(4-cyano-3-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;
2-methyl-2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;
30 rac-2-(trans-4-{2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;

rac-2-(trans-4-{3-chloro-4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid;

cis-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenoxy}cyclohexanecarboxylic acid;

5 (trans-4-{4-[(5-{(2-chloro-4-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(4-methoxy-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(4-fluoro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-

10 yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

[trans-4-{4-[(5-{(4-fluoro-2-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(2,4-dichlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

15 (trans-4-{4-[(5-{(2-chloro-4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(4-cyano-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(4-chloro-2,5-dimethylphenyl)amino}-1,3,4-oxadiazol-2-

20 yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(4-cyano-2-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(2-chloro-4-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

25 (trans-4-{4-[(5-{(4-chloro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(2,3-dichlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(2,5-dichlorophenyl)amino}-1,3,4-oxadiazol-2-

30 yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

(trans-4-{4-[(5-{(2-bromo-4-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid;

- [*trans*-4-(4-{{(5-{{(2-chloro-5-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 [*trans*-4-(4-{{(5-{{(2-bromo-4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 5 [*trans*-4-(4-{{(5-{{(2-chloro-4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 (*trans*-4-(4-{{(5-{{(4-bromo-2-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 (*trans*-4-(4-{{(5-{{(3-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 10 (*trans*-4-(4-{{(5-{{(5-bromo-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 (*trans*-4-(4-{{(5-{{(2-chloro-5-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 15 (*trans*-4-(4-{{(5-{{(2-chloro-5-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 (*trans*-4-(4-{{(5-{{(5-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid; and
 2-(3-(4-{{(5-{{(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclobutyl)propanoic acid.
 20

In a further aspect of the invention, there is provided any one or more of the following compounds, or pharmaceutically-acceptable salts or pro-drugs thereof:

- [*trans*-4-(4-{{(5-{{(6-(4-chlorophenoxy)pyridin-3-yl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 25 [*trans*-4-(4-{{(5-{{(6-(4-methoxyphenoxy)pyridin-3-yl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 {*trans*-4-(4-{{(5-{{(quinolin-3-yl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 (*trans*-4-(4-{{(5-{{(5-bromopyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;
 30 [*trans*-4-(4-{{(5-{{(pyridin-3-yl)amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid;

(trans-4-{4-[(5-{(6-methoxypyridin-3-yl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid.

In a further aspect of the invention, there is provided any one or more of the following compounds, or pharmaceutically-acceptable salts or pro-drugs thereof:

- 5 1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;
(1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}-4-methylpiperidine-4-carboxylic acid;
10 1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}-4-methylpiperidine-4-carboxylic acid;
4-methyl-1-{5-[(5-{(4-(trifluoromethoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;
15 4-methyl-1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;
4-methyl-1-{5-[(5-{(4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;
1-{5-[(5-{(4-(4-chlorophenoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}-4-methylpiperidine-4-carboxylic acid;
20 1-{5-[(5-{(4-(4-chlorophenoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;
[1-{5-[(5-{(4-(4-chlorophenoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
25 (1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
[1-{5-[(5-{(4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid;
1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;
30 1-{5-[(5-{(4-(trifluoromethoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid;

- 1-(5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
(1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl)acetic acid;
- 5 [1-(3-methyl-5-{{(5-{{(4-(trifluoromethoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
[1-(3-methyl-5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
(1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl)acetic acid;
- 10 1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylic acid;
1-(3-methyl-5-{{(5-{{(4-(trifluoromethoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
- 15 1-(3-methyl-5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylic acid;
2-(1-{3-methyl-5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)propanoic acid;
- 20 1-(5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidine-4-carboxylic acid;
1-(5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid;
- 25 [1-(5-{{(5-{{(4-(4-fluorophenoxy)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
(1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl)acetic acid;
(4-methyl-1-{5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid;
- 30 [4-methyl-1-(5-{{(5-{{(4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl)carbonyl}amino}pyridin-2-yl)piperidin-4-yl)acetic acid; and

[1-(5-{{{5-{{4-(4-fluorophenoxy)phenyl}amino}-1,3,4-oxadiazol-2-yl}carbonyl}amino}pyridin-2-yl)-4-methylpiperidin-4-yl]acetic acid.


Further preferred compounds of the invention are each of the Examples, each of which provides a further independent aspect of the invention. In further aspects, the present invention also comprises any two or more compounds of the Examples.

Process


A compound of formula (I) and its pharmaceutically-acceptable salts may be prepared by any process known to be applicable to the preparation of chemically related compounds. Such processes, when used to prepare a compound of the formula (I), or a pharmaceutically-acceptable salt thereof, are provided as a further feature of the invention.

In a further aspect the present invention also provides that the compounds of the formula (I) and pharmaceutically-acceptable salts or prodrugs thereof, can be prepared by a process a) to g) as follows (wherein all variables are as hereinbefore defined for a compound of formula (I) unless otherwise stated):


- a) reaction of a compound of formula (I) to form another compound of formula (I);
b) reaction of an amine of formula (2) with a carboxylate salt of formula (3);




- 20 c) cyclisation of a compound of formula (4) (where X is S or O):




- d) when R^2 is substituted by piperazinyl, by reaction of the piperazine nitrogen with R^5 -LG, wherein LG is a suitable leaving group such as halo, and R^5 is hydrocarbyl or a
25 suitable functional group such as acyl, for example:




e) when R^2 is aryl and is substituted by aryl, by transition metal catalysed aromatic substitution, for example:



f) when R^2 is substituted by piperazinyl, by reductive alkylation of the piperazine nitrogen with R^5 -CHO (wherein R^5 is for example hydrocarbyl), for example:



g) reaction of halogenated (for example with iodo) R^2 with an amide of formula (7) followed by subsequent removal of protecting group P^1 , wherein P^1 is for example benzyl or trimethylsilylethoxymethyl (SEM), for example;



wherein Hal is halogen;

and thereafter if necessary, removing any protecting groups, and/or forming a pharmaceutically-acceptable salt or prodrug thereof.

In the above schemes, R^1 , R^2 , Y and R^5 are as defined above. Suitably in the above schemes, R^1 is a group R^{1a} and R^2 is a group R^{2a} as defined above so that they can be used to prepare a compound of formula (IA). Furthermore, the group R^1 may be a group R^{1b} and R^2 may be a group R^{2b} in order to produce a compound of formula (IB). In the latter

case, Y is a direct bond. It will be understood that, where Y is a direct bond, processes a,b,d, and g apply to compounds of formula (1ZA), processes a, b, c and g apply to compounds of formula (1ZB) and processes a, b, c, d, f and g apply to compounds of formula (1ZC).

5 Process a)


Examples of conversions of a compound of formula (1) into another compound of Formula (1), well known to those skilled in the art, include functional group interconversions such as hydrolysis (in particular ester hydrolysis), oxidation or reduction (such as the reduction of an acid to an alcohol, or removal of an N protecting group),
10 and/or further functionalisation by standard reactions such as amide or metal-catalysed coupling, or nucleophilic displacement reactions.

Process b)

Compounds of formula (2) where Y is not a direct bond or where R^2 is not aromatic, may be made by application of standard synthetic methods well known in the art.
15 For example, reductive alkylation of ammonia (or a suitable amine such as a benzylamine or N,N-dibenzylamine) with a ketone or aldehyde $R^2Y=O$ (followed by deprotection as appropriate) provides R^2-Y-NH_2 . Alternatively, alkylation of an amine or amine equivalent (such as a Gabriel reagent or a guanidine) with a halide R^2-Y-X (where X is a halide) (followed by N-deprotection or hydrolysis as appropriate) provides the required
20 compounds of formula (2).


Compounds of formula (2) for other definitions of Y or R^2 may be made by metal catalysed couplings or nucleophilic displacement reactions among other methods. In particular, compounds of formula (2) may be prepared by reduction of a compound of formula (2A).

25




(2A)

Compounds of formula (2A) may be made by metal catalysed couplings or nucleophilic displacement reactions depending upon the nature of the R^2 group and Y. For example, production of a compound of formula (2A) may be represented as follows:




30


Examples of process (b) where Y is a direct bond are shown in Schemes 1 to 3 (wherein R^6 represents optional substituent on R^2 , including those defined hereinbefore for R^{6a-d}):



Scheme 1




Scheme 2




Scheme 3

Certain compounds of formula (2) may also have chiral centres or can exist in different isomeric forms such as cis/trans isomers, and may be prepared as individual isomers, as illustrated below in Schemes 4 and 5 and in the accompanying examples.




Scheme 4



Scheme 5



The process illustrated in Scheme 5 may also be used with cyclohexenone as a starting material. The opposite stereochemistry may be obtained by using known alternative chiral catalysts and/or chiral ligands. Elaboration of the bicyclic ketone intermediate may be carried out by processes known in the art, for example by Wittig or enolate/enol ether chemistry, optionally followed by functionalisation (such as alkylation) and functional group interconversion as desired to give the compound of formula (2) (wherein Ra and Rb may each for example be hydrogen or (optionally substituted) alkyl groups). Mixtures of diastereoisomers may be separated by standard procedures.

S_NAr chemistry may be used (under conditions well known in the art) to make certain compounds of formula (2), as illustrated in Scheme 6 (in which R is for example an alkyl group, X is for example Br or Cl, n is for example 0 to 4, group A may be a (hetero)aryl ring, a saturated ring or an alkyl chain).




Scheme 6

Compounds of formula (3) may be made by alkaline hydrolysis of ester (8a) as prepared using a published procedure (J. Het. Chem. 1977, 14, 1385-1388). Ester (8a) may be made by cyclisation of a compound of formula (8b) (where X is O or S) in a similar manner as described in process c) for compounds of formula (4).



An alternative method for making compounds of formula (8a) is illustrated below:



Compounds of formula (2) may be coupled with compounds of formula (3) under standard conditions for formation of amide bonds. For example using an appropriate coupling reaction, such as a carbodiimide coupling reaction performed with EDAC, optionally in the presence of DMAP, in a suitable solvent such as DCM, chloroform or DMF at room temperature.


For compounds of formula (2) other than when R^2 is aromatic and Y is a direct bond (ie other than compounds such as anilino compounds), an ester derivative of formula (8a) (or equivalent) may be used instead of the compound of formula (3) to couple with the compound of formula (2). Such a reaction may be carried out by any method known in the art such as by heating (thermally or by microwave) in a suitable solvent.

Process c)

Compounds of formula (4) and (8b) where X is S may be made by reaction of an aminocarbonyl acylhydrazine or ethoxycarbonyl acylhydrazine with a thioisocyanate or thioisocyanate equivalent such as aminothiocabonylimidazole in a suitable solvent such as DMF or MeCN at a temperature between 0 and 100 °C. The preparation of aminocarbonyl acylhydrazines from anilines and of ethoxycarbonyl acylhydrazines is well known in the art. For example reaction of an aniline with methyl chlorooxoacetate in the presence of pyridine in a suitable solvent such as DCM followed by reaction with hydrazine in a suitable solvent such as ethanol at a temperature between 0 and 100 °C.

The compound of formula (4) may then be cyclised using, for example agents such as carbonyldiimidazole, or tosyl chloride and a suitable base (such as triethylamine), under conditions known in the art.

An example of process c) is shown in Scheme 7 (wherein R^3 represents a substituent on R^1 as hereinbefore defined, R^6 represents an optional substituent on R^2 , including those defined hereinbefore as R^{6a-d});



Scheme 7

- 5 Iso(thio)cyanates R^1-NCX (where X is O or S) are commercially available or may be made by reaction of the acid chlorides R^1-NH_2 with for example (thio)phosgene or a (thio)phosgene equivalent followed by a suitable base (such as triethylamine).

Compounds of formula (4) may be made from compounds of formula (2) (wherein R^2 and Y are as defined for a compound of formula (1)) as illustrated above in Scheme 7.

Process d)

- 10 Compounds of formula (5) can be reacted with an acid chloride or sulfonyl chloride in the presence of a base such as triethylamine or pyridine in a suitable solvent such as DMF.

Process e)

- 15 Compounds of formula (6) can be reacted with aryl boronic acids in the presence of a suitable catalyst such as tetrakis(triphenyl phosphine)palladium(0) and a suitable base such as potassium phosphate in a suitable solvent such as DME-water (2:1) under microwave heating at 0 to 110 °C.

Process f)

Compounds of formula (5) can be reacted with aldehydes in the presence of a suitable acid such as acetic acid, and a reducing agent such as sodium borohydride in a suitable solvent such as DCM.

5 Process g)

Compounds of formula (7) can be reacted with arylbromides, aryliodides, heteroaryl bromides or heteroaryl iodides in the presence of a suitable catalyst such as copper(I) iodide, a suitable diamine ligand such as *trans*-*N,N'*-dimethyl-1,2-cyclohexyldiamine and a suitable base such as potassium phosphate in a suitable solvent such as DMF or dioxane heating at 80-110 °C.

It will be appreciated that certain of the various ring substituents in the compounds of the present invention, for example Z, Z_a, and/or R³, may be introduced by standard aromatic substitution reactions or generated by conventional functional group
15 modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions may convert one compound of the formula (I) into another compound of the formula (I). Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of
20 substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the
25 introduction of a halogen group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkanesulfinyl or alkanesulfonyl.

30 If not commercially available, the necessary starting materials for the procedures such as those described above may be made by procedures which are selected from standard organic chemical techniques, techniques which are analogous to the synthesis of

known, structurally similar compounds, techniques which are described or illustrated in the references given above, or techniques which are analogous to the above described procedure or the procedures described in the examples. The reader is further referred to Advanced Organic Chemistry, 5th Edition, by Jerry March and Michael Smith, published
5 by John Wiley & Sons 2001, for general guidance on reaction conditions and reagents.

It will be appreciated that some intermediates to compounds of the formula (I) are also novel and these are provided as separate independent aspects of the invention.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in compounds. The instances where
10 protection is necessary or desirable are known to those skilled in the art, as are suitable methods for such protection. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Greene, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991).

Protecting groups may be removed by any convenient method as described in the
15 literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Thus, if reactants include, for example, groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

20 Examples of a suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, a silyl group such as trimethylsilyl or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an
25 aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively a silyl group such as trimethylsilyl or SEM may be removed, for example, by fluoride or by aqueous acid; or an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation in the presence of a catalyst such as palladium-on-carbon.

30 A suitable protecting group for an amino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or tert-butoxycarbonyl group, an arylmethoxycarbonyl

group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base
5 such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulfuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or
10 by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine or 2-hydroxyethylamine, or with hydrazine.

A suitable protecting group for a carboxy group is, for example, an esterifying
15 group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

20 Resins may also be used as a protecting group.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art, or they may be removed during a later reaction step or work-up.

The skilled organic chemist will be able to use and adapt the information contained
25 and referenced within the above references, and accompanying Examples therein and also the examples herein, to obtain necessary starting materials, and products.

The removal of any protecting groups and the formation of a pharmaceutically-acceptable salt are within the skill of an ordinary organic chemist using standard techniques. Furthermore, details on these steps has been provided
30 hereinbefore.

When an optically active form of a compound of the invention is required, it may be obtained by carrying out one of the above procedures using an optically active starting

material (formed, for example, by asymmetric induction of a suitable reaction step), or by resolution of a racemic form of the compound or intermediate using a standard procedure, or by chromatographic separation of diastereoisomers (when produced). Enzymatic techniques may also be useful for the preparation of optically active compounds and/or intermediates.

Similarly, when a pure regioisomer of a compound of the invention is required, it may be obtained by carrying out one of the above procedures using a pure regioisomer as a starting material, or by resolution of a mixture of the regioisomers or intermediates using a standard procedure.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (1), (IA), (IB), (IZA), (IZB) and (IZC) as defined hereinbefore or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable excipient or carrier.

Certain intermediates used in the processes described above are novel, and these form a further aspect of the invention. In particular, compounds of formula (4) form a further aspect of the invention.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic

acid or talc; preservative agents such as ethyl or propyl *p*-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above.

5 Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these.

10 Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain
15 sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable
20 aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

25 Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

30 For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

According to a further aspect of the present invention there is provided a compound of formula (I) or a pharmaceutically acceptable salt thereof as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

Reference herein to a compound of formula (I) should be understood to refer equally to compounds of formulae (I), (IA), (IB) (and sub-formulae thereof), (IZA), (IZB) and/or (IZC).

We have found that compounds of the present invention inhibit DGAT1 activity and are therefore of interest for their blood glucose-lowering effects.

A further feature of the present invention is a compound of formula (I) or a pharmaceutically-acceptable salt thereof for use as a medicament.

Conveniently this is a compound of formula (I), or a pharmaceutically-acceptable salt thereof, for use as a medicament for producing an inhibition of DGAT1 activity in a warm-blooded animal such as a human being.

Particularly this is a compound of formula (I), or a pharmaceutically-acceptable salt thereof, for use as a medicament for treating diabetes mellitus and/or obesity in a warm-blooded animal such as a human being.

Thus according to a further aspect of the invention there is provided the use of a compound of formula (I), or a pharmaceutically-acceptable salt thereof in the manufacture of a medicament for use in the production of an inhibition of DGAT1 activity in a warm-blooded animal such as a human being.

Thus according to a further aspect of the invention there is provided the use of a compound of formula (I), or a pharmaceutically-acceptable salt thereof in the manufacture

of a medicament for use in the treatment of diabetes mellitus and/or obesity in a warm-blooded animal such as a human being.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I) as defined hereinbefore or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable excipient or carrier for use in producing an inhibition of DGAT1 activity in an warm-blooded animal, such as a human being.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I) as defined hereinbefore or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable excipient or carrier for use in the treatment of diabetes mellitus and/or obesity in an warm-blooded animal, such as a human being.

According to a further feature of the invention there is provided a method for producing an inhibition of DGAT1 activity in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) or a pharmaceutically-acceptable salt thereof as defined hereinbefore.

According to a further feature of the invention there is provided a method of treating diabetes mellitus and/or obesity in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) or a pharmaceutically-acceptable salt thereof as defined hereinbefore.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular disease state will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

As stated above compounds defined in the present invention are of interest for their ability to inhibit the activity of DGAT1. A compound of the invention may therefore be useful for the prevention, delay or treatment of a range of disease states including diabetes

14

mellitus, more specifically type 2 diabetes mellitus (T2DM) and complications arising there from (for example retinopathy, neuropathy and nephropathy), impaired glucose tolerance (IGT), conditions of impaired fasting glucose, metabolic acidosis, ketosis, dysmetabolic syndrome, arthritis, osteoporosis, obesity and obesity related disorders, (which include peripheral vascular disease, (including intermittent claudication), cardiac failure and certain cardiac myopathies, myocardial ischaemia, cerebral ischaemia and reperfusion, hyperlipidaemias, atherosclerosis, infertility and polycystic ovary syndrome); the compounds of the invention may also be useful for muscle weakness, diseases of the skin such as acne, Alzheimer's disease, various immunomodulatory diseases (such as psoriasis), HIV infection, inflammatory bowel syndrome and inflammatory bowel disease such as Crohn's disease and ulcerative colitis.

In particular, the compounds of the present invention are of interest for the prevention, delay or treatment of diabetes mellitus and/or obesity and/or obesity related disorders. In one aspect, the compounds of the invention are used for prevention, delay or treatment of diabetes mellitus. In another aspect, the compounds of the invention are used for prevention, delay or treatment of obesity. In a further aspect, the compounds of the invention are used for prevention, delay or treatment of obesity related disorders.

In a further aspect of the invention, compounds of the invention are used for the treatment of Alzheimer's disease.

The inhibition of DGAT1 activity described herein may be applied as a sole therapy or in combination with one or more other substances and/or treatments for the indication being treated. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. Simultaneous treatment may be in a single tablet or in separate tablets. For example such conjoint treatment may be beneficial in the treatment of metabolic syndrome [defined as abdominal obesity (as measured by waist circumference against ethnic and gender specific cut-points) plus any two of the following: hypertriglyceridemia (> 150 mg/dl; 1.7 mmol/l); low HDLc (< 40 mg/dl or < 1.03 mmol/l for men and < 50 mg/dl or 1.29 mmol/l for women) or on treatment for low HDL (high density lipoprotein); hypertension (SBP ≥ 130 mmHg DBP ≥ 85 mmHg) or on treatment for hypertension; and hyperglycemia (fasting plasma glucose ≥ 100 mg/dl or 5.6 mmol/l or impaired glucose

tolerance or pre-existing diabetes mellitus) - International Diabetes Federation & input from IAS/NCEP].

Such conjoint treatments may include the following main categories:

- 1) Anti-obesity therapies such as those that cause weight loss by effects on food intake, nutrient absorption or energy expenditure, such as orlistat, sibutramine and the like.
- 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide), prandial glucose regulators (for example repaglinide, nateglinide);
- 3) Agents that improve incretin action (for example dipeptidyl peptidase IV inhibitors, and GLP-1 agonists);
- 4) Insulin sensitising agents including PPARgamma agonists (for example pioglitazone and rosiglitazone), and agents with combined PPARalpha and gamma activity;
- 5) Agents that modulate hepatic glucose balance (for example metformin, fructose 1, 6 bisphosphatase inhibitors, glycogen phosphorylase inhibitors, glycogen synthase kinase inhibitors, glucokinase activators);
- 6) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 7) Agents that prevent the reabsorption of glucose by the kidney (SGLT inhibitors);
- 8) Agents designed to treat the complications of prolonged hyperglycaemia (for example aldose reductase inhibitors);
- 9) Anti-dyslipidaemia agents such as, HMG-CoA reductase inhibitors (eg statins); PPARalpha-agonists (fibrates, eg gemfibrozil); bile acid sequestrants (cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors); bile acid absorption inhibitors (IBATi) and nicotinic acid and analogues (niacin and slow release formulations);
- 10) Antihypertensive agents such as, beta-blockers (eg atenolol, inderal); ACE inhibitors (eg lisinopril); Calcium antagonists (eg. nifedipine); Angiotensin receptor antagonists (eg candesartan), alpha antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 11) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors; antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low molecular weight analogues, hirudin) and warfarin;
- 12) Agents which antagonise the actions of glucagon; and

13) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin) and steroidal anti-inflammatory agents (eg. cortisone).

In addition to their use in therapeutic medicine, compounds of formula (I) and their pharmaceutically-acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of DGAT1 activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

As indicated above, all of the compounds, and their corresponding pharmaceutically-acceptable salts, are useful in inhibiting DGAT1. The ability of the compounds of formula (I), and their corresponding pharmaceutically-acceptable acid addition salts, to inhibit DGAT1 may be demonstrated employing the following enzyme assay:

Human Enzyme Assay

The *in vitro* assay to identify DGAT1 inhibitors uses human DGAT1 expressed in insect cell membranes as the enzyme source (Proc. Natl. Acad. Sci. 1998, 95, 13018-13023). Briefly, sf9 cells were infected with recombinant baculovirus containing human DGAT1 coding sequences and harvested after 48 h. Cells were lysed by sonication and membranes isolated by centrifuging at 28000 rpm for 1 h at 4 °C on a 41% sucrose gradient. The membrane fraction at the interphase was collected, washed, and stored in liquid nitrogen.

DGAT1 activity was assayed by a modification of the method described by Coleman (Methods in Enzymology 1992, 209, 98-102). Compound at 1-10 μ M was incubated with 0.4 μ g membrane protein, 5 mM $MgCl_2$, and 10 0 μ M 1,2-dioleoyl-sn-glycerol in a total assay volume of 200 μ l in plastic tubes. The reaction was started by adding ^{14}C oleoyl coenzyme A (30 μ M final concentration) and incubated at room temperature for 30 minutes. The reaction was stopped by adding 1.5 mL 2-propanol:heptane:water (80:20:2). Radioactive triolein product was separated into the organic phase by adding 1mL heptane and 0.5 mL 0.1 M carbonate buffer pH 9.5. DGAT1 activity was quantified by counting aliquots of the upper heptane layer by liquid scintillography.

Using this assay the compounds generally show activity with $IC_{50} < 10mM$, preferably $< 1 \mu M$. Example 48 showed an $IC_{50} = 0.12 \mu M$.

The ability of the compounds of formula (I), and their corresponding pharmaceutically-acceptable acid salts, to inhibit DGAT1 may further be demonstrated employing the following whole cell assays 1) and 2):

1) Measurement of Triglyceride Synthesis in 3T3 Cells

- 5 Mouse adipocyte 3T3 cells were cultured to confluency in 6 well plates in new born calf serum containing media. Differentiation of the cells was induced by incubating in medium containing 10% foetal calf serum, 1 $\mu\text{g/mL}$ insulin, 0.25 μM dexamethasone and 0.5 mM isobutylmethyl xanthine. After 48 h the cells were maintained in medium containing 10% foetal calf serum and 1 $\mu\text{g/mL}$ insulin for a further 4-6 days. For the experiment, the
- 10 medium was changed to serum-free medium and the cells pre-incubated with compound solubilised in DMSO (final concentration 0.1%) for 30 minutes. De novo lipogenesis was measured by the addition of 0.25 mM sodium acetate plus 1 $\mu\text{Ci/mL}$ ^{14}C -sodium acetate to each well for a further 2 h (J. Biol. Chem., 1976, 251, 6462-6464). The cells were washed in phosphate buffered saline and solubilised in 1% sodium dodecyl sulfate. An aliquot was
- 15 removed for protein determination using a protein estimation kit (Perbio) based on the method of Lowry (J. Biol. Chem., 1951, 193, 265-275). The lipids were extracted into the organic phase using a heptane:propan-2-ol:water (80:20:2) mixture followed by aliquots of water and heptane according to the method of Coleman (Methods in Enzymology, 1992, 209, 98-104). The organic phase was collected and the solvent evaporated under a stream
- 20 of nitrogen. The extracts solubilised in iso-hexane:acetic acid (99:1) and lipids separated via normal phase high performance liquid chromatography (HPLC) using a Lichrospher diol-5, 4 $\times$ 250 mm column and a gradient solvent system of iso-hexane:acetic acid (99:1) and iso-hexane:propan-2-ol:acetic acid (85:15:1), flow rate of 1 mL/minute according to the method of Silversand and Haux (1997). Incorporation of radiolabel into the triglyceride
- 25 fraction was analysed using a Radiomatic Flo-one Detector (Packard) connected to the HPLC machine.

2) Measurement of Triglyceride Synthesis in MCF7 Cells

- Human mammary epithelial (MCF7) cells were cultured to confluency in 6 well plates in foetal calf serum containing media. For the experiment, the medium was changed to
- 30 serum-free medium and the cells pre-incubated with compound solubilised in DMSO (final concentration 0.1%) for 30 minutes. De novo lipogenesis was measured by the addition of

50 μ M sodium acetate plus 3 μ Ci/mL ^{14} C-sodium acetate to each well for a further 3 h (J. Biol. Chem., 1976, 251, 6462-6464). The cells were washed in phosphate buffered saline and solubilised in 1% sodium dodecyl sulfate. An aliquot was removed for protein determination using a protein estimation kit (Perbio) based on the method of Lowry (J. Biol. Chem., 1951, 193, 265-275). The lipids were extracted into the organic phase using a heptane:propan-2-ol:water (80:20:2) mixture followed by aliquots of water and heptane according to the method of Coleman (Methods in Enzymology, 1992, 209, 98-104). The organic phase was collected and the solvent evaporated under a stream of nitrogen. The extracts solubilised in iso-hexane:acetic acid (99:1) and lipids separated via normal phase high performance liquid chromatography (HPLC) using a Lichrospher diol-5, 4 $\times$ 250 mm column and a gradient solvent system of iso-hexane:acetic acid (99:1) and iso-hexane:propan-2-ol:acetic acid (85:15:1), flow rate of 1 mL/minute according to the method of Silversand and Haux (J. Chromat. B, 1997, 703, 7-14). Incorporation of radiolabel into the triglyceride fraction was analysed using a Radiomatic Flo-one Detector (Packard) connected to the HPLC machine.

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

Examples

The invention will now be illustrated by the following Examples in which, unless stated otherwise:

- (i) temperatures are given in degrees Celsius ($^{\circ}$ C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25 $^{\circ}$ C and under an atmosphere of an inert gas such as argon;
- (ii) organic solutions were dried over anhydrous magnesium sulfate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pa; 4.5-30 mmHg) with a bath temperature of up to 60 $^{\circ}$ C;
- (iii) chromatography means flash chromatography on silica gel; where a Biotage cartridge is referred to this means a cartridge containing KP-SILTM silica, 60 $\text{\AA}$, particle size 32-63 μ m, supplied by Biotage, a division of Dyax Corp., 1500 Avon Street Extended,

Charlottesville, VA 22902, USA;

(iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;

(v) yields are given for illustration only and are not necessarily those which can be

5 obtained by diligent process development; preparations were repeated if more material was required;

(vi) where given, NMR data (^1H) is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS), determined at 300 or 400 MHz (unless otherwise stated) using perdeuterio dimethyl sulfoxide

10 (DMSO- d_6) as solvent, unless otherwise stated; peak multiplicities are shown thus: s, singlet; d, doublet; dd, doublet of doublets; dt, doublet of triplets; dm, doublet of multiplets; t, triplet; q, quartet; m, multiplet; br, broad;

(vii) chemical symbols have their usual meanings; SI units and symbols are used;

(viii) solvent ratios are given in volume : volume (v/v) terms;

15 (ix) mass spectra (MS) (loop) were recorded on a Micromass Platform LC equipped with HP 1100 detector; unless otherwise stated the mass ion quoted is (MH^+);

(x) LCMS (liquid chromatography-mass spectrometry) were recorded on a system comprising Waters 2790 LC equipped with a Waters 996 Photodiode array detector and Micromass ZMD MS, using a Phenomenex® Gemini 5u C18 110A 50x2 mm column and
20 eluting with a flow rate of 1.1 ml/min with 5% (Water/Acetonitrile (1:1) + 1% formic acid) and a gradient increasing from 0-95% of acetonitrile over the first 4 minutes, the balance (95-0%) being water and where HPLC Retention Times are reported these are in minutes in this system unless otherwise stated; unless otherwise stated the mass ion quoted is (MH^+);

25 (xi) where phase separation cartridges are stated then ISOLUTE Phase Separator 70ml columns, supplied by Argonaut Technologies, New Road, Hengoed, Mid Glamorgan, CF82 8AU, United Kingdom, were used;

(xii) where a SiliCycle cartridge is referred to this means a cartridge containing Ultra Pure Silica Gel particle size 230-400 mesh, 40 -63 μm pore size, supplied by SiliCycle Chemical
30 Division, 1200 Ave St-Jean-Baptiste, Suite 114, Quebec City, Quebec, G2E 5E8, CANADA;


- (xiii) where an Isco Companion is referred to then a Combiflash companion chromatography instrument, supplied by ISOC Inc. Address Teledyne ISOC Inc, 4700 Superior Street, Lincoln, NE 68504, USA, was used;
- (xiv) where a microwave is referred to this means a Biotage Initiator sixty or Smith Creator microwave, supplied by Biotage, a division of Dyax Corp., 1500 Avon Street Extended, Charlottesville, VA 22902, USA;
- 5 (xv) where GCMS is referred to then a Gas Chromatography –Mass Spectrometry analysis was carried out on a QP-2010 GC-MS system fitted with an AOC 20i autosampler and controlled by 'GCMS solutions' software, version 2.0, supplied by Shimadzu, Milton
- 10 Keynes, MK12 5RE, UK; the GC column was a DB-5MS of length 25 m, 0.32 mm i.d. with a film thickness of 0.52 μ m supplied by J & W Scientific, Folsom, CA, USA;
- (xvi) where a centrifuge is referred to this means a Genevac EZ-2plus, supplied by Genevac Limited, The Sovereign Centre, Farthing Road, Ipswich, IP1 5AP, UK;
- (xvii) where chiral chromatography is referred to this is carried generally out using a 20 μ m
- 15 Merck 50mm Chiralpak AD column, (Chiral Stationary Phase supplied by Chiral Technologies Europe, Parc d'Innovation, Bd. Gonthier d'Andernach, 67404 Illkirch Cedex, France), using MeCN/2-propanol/AcOH (90/10/0.1) as eluent, flow rate 80 mL/min, wavelength 300nm, using a Gilson prep HPLC instrument (200ml heads);
- (xviii) melting points were determined using a Buchi 530 apparatus and are uncorrected;
- 20 (xix) The following abbreviations may be used below or in the process section hereinbefore:

	Et ₂ O or ether	diethyl ether
	DMF	dimethylformamide
	DCM	dichloromethane
25	DME	1,2-dimethoxyethane
	MeOH	methanol
	EtOH	ethanol
	H ₂ O	water
	TFA	trifluoroacetic acid
30	THF	tetrahydrofuran
	DMSO	dimethylsulfoxide
	HOBt	1-hydroxybenzotriazole

	EDCI (EDAC)	1-ethyl-3-(3-dimethylaminopropyl)carbodi-imide hydrochloride
	DIPEA	diisopropylethylamine
	DEAD	diethyl azodicarboxylate
5	EtOAc	ethyl acetate
	NaHCO ₃	sodium bicarbonate / sodium hydrogencarbonate
	K ₃ PO ₄	potassium phosphate
	PS	polymer supported.
	BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
10	Dppf	1,1'-bis(diphenylphosphino)ferrocene
	dba	dibenzylideneacetone
	PS-CDI	polymer supported carbonyldiimidazole
	CH ₃ CN or MeCN	acetonitrile
	h	hour
15	min	minute
	HA'TU	O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
	NaOH	sodium hydroxide
	AcOH	acetic acid
20	DMA	dimethyl acetamide
	nBuLi	n-butyl lithium
	MgSO ₄	magnesium sulfate
	Na ₂ SO ₄	sodium sulfate
	CDCl ₃	deutero chloroform
25	CD ₃ OD	per-deuterated methanol
	Boc	tert-butoxycarbonyl

All compound names were derived using ACD NAME computer package.


Example 1: 5-[(2-Fluorophenyl)amino]-N-(4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide










2-Fluorophenylisothiocyanate (92 μ L, 0.75 mmol) was added to a stirred suspension of 2-hydrazino-*N*-(4-morpholin-4-ylphenyl)-2-oxoacetamide (Intermediate 16, 160 mg, 0.6 mmol) in DMF (10 mL) and stirred at 50 °C until a clear solution was obtained. PS-Carbodiimide (909 mg, 1.2 mmol) was added and the reaction was heated at 80 °C for a further 4 h. The reaction was filtered and the resin washed with DMF (5 mL). The combined DMF solutions were concentrated *in vacuo* and the residue triturated with diethyl ether to give the title compound as a yellow powder (190 mg, 82%):
 $^1\text{H NMR}$ δ 10.83 (1H, s), 10.77 (1H, s), 8.05 (1H, t), 7.66 (2H, d), 7.30 (2H, m), 6.95 (2H, d), 3.80-3.65 (4H, m), 3.11-3.01 (4H, m); MS m/e MH^+ 384.








Examples 2-66








The following examples were prepared by the general procedure of Example 1, using Intermediates 23 to 28 and commercially available isothiocyanates.

















Example	R ⁶	R ¹	$^1\text{H NMR}$ δ	MS m/e MH^+
2	H		11.01 (1H, s), 10.86 (1H, s), 7.67 (2H, d), 7.51-7.45 (2H, m), 7.41 (1H, t), 7.37-7.25 (4H, m), 7.15 (1H, d), 6.95 (2H, d), 6.74 (1H, d), 5.15 (2H, s), 3.79-3.69 (4H, m), 3.15-3.03 (4H, m)	472








Example	R ⁶	R ⁷	¹ H NMR δ	MS m/e MII ⁺
3	H		11.10 (1H, s), 10.88 (1H, s), 7.68 (2H, d), 7.57-7.39 (3H, m), 6.96 (2H, d), 3.87-3.67 (4H, m), 3.20-3.03 (4H, m), 2.41 (3H, s)	414
4	H		11.10 (1H, s), 10.82 (1H, s), 7.62 (4H, dd), 7.39 (2H, d), 6.93 (2H, d), 3.81-3.65 (4H, m), 3.11-3.01 (4H, m)	400
5	H		10.82-10.76 (2H, m), 7.67 (2H, d), 7.51 (1H, s), 7.35 (1H, d), 7.22 (1H, d), 6.95 (2H, d), 3.79-3.70 (4H, m), 3.14-3.05 (4H, m), 2.93- 2.79 (4H, m), 2.09-1.98 (2H, m)	406
6	H		10.80 (2H, s), 7.7 (2H, d), 7.45 (1H, m), 7.4 (1H, s), 7.3 (1H, m), 6.96 (2H, d), 6.9 (1H, d), 3.80-3.64 (4H, m), 3.11-3.00 (4H, m)	380
7	H		10.80 (1H, s), 10.70 (1H, s), 8.30 (1H, d), 8.02 (1H, d), 7.95 (1H, d), 7.75 (1H, d), 7.70 (2H, d), 7.58 (3H, m), 6.95 (2H, d), 3.80- 3.65 (4H, m), 3.12-3.01 (4H, m)	416
8	H		11.27 (1H, s), 10.94 (1H, s), 7.49 (1H, q), 7.69 (3H, m), 7.35 (1H, m), 6.95 (2H, d), 3.80-3.66 (4H, m), 3.11-3.00 (4H, m)	402
9	H		10.89 (1H, s), 10.82 (1H, s), 7.67 (2H, d), 7.50-7.40 (2H, m), 7.30 (1H, t), 7.00-6.89 (3H, m), 3.80-3.70 (4H, m), 3.16-3.04 (4H, m), 2.62 (2H, q), 1.20 (3H, t)	394


Example	R ⁶	R ⁷	¹ H NMR δ	MS m/e MH ⁺
10	H		11.22 (1H, s), 10.87 (1H, s), 8.30 (1H, s), 7.87 (1H, d), 7.71-7.62 (3H, m), 7.56 (2H, t), 6.95 (2H, d), 4.35 (2H, q), 3.80-3.69 (4H, m), 3.15-3.04 (4H, m), 1.34 (3H, t)	438
11	H		11.20 (1H, s), 10.82 (1H, s), 7.72 (1H, s), 7.69 (2H, d), 7.50 (1H, d), 7.41 (1H, dd), 7.1 (1H, d), 6.95 (2H, d), 3.79-3.64 (4H, m), 3.10-3.01 (4H, m)	400
12	H		10.95 (1H, s), 10.64 (1H, s), 8.0 (1H, s), 7.73 (1H, s), 7.63 (2H, m), 7.44 (3H, m), 7.25 (1H, m), 7.16 (1H, m), 6.75 (2H, d), 3.60- 3.50 (4H, m), 2.93-2.84 (4H, m)	416
13	H		11.15 (1H, s), 10.81 (1H, s), 7.85 (1H, d), 7.70 (2H, d), 7.52 (1H, d), 7.31 (1H, t), 7.20 (1H, d), 6.93 (2H, d), 3.80-3.69(4H, m), 3.12-3.01 (4H, m)	446
14	H		10.81 (1H, s), 10.75 (1H, s), 7.67 (2H, d), 7.39-7.30 (2H, m), 7.14 (1H, d), 6.95 (2H, d), 3.79-3.69 (4H, m), 3.13-3.04 (4H, m), 2.27 (3H, s), 2.16 (3H, s)	394
15	H		11.12 (1H, s), 10.87 (1H, s), 7.73 (1H, d), 7.66 (2H, d), 7.45-7.33 (2H, m), 6.95 (2H, d), 3.80-3.70 (4H, m), 3.14-3.05 (4H, m), 2.32 (3H, s)	414
16	H		11.01 (1H, s), 10.85 (1H, s), 7.66 (2H, d), 7.59 (1H, s), 7.40-7.29 (2H, m), 7.00-6.90 (3H, m), 3.80-3.70 (4H, m), 3.28 (3H, s), 3.14-3.04 (4H, m)	412


Example	R ⁶	R ⁷	¹ H NMR δ	MS m/e MH ⁺
17	H		10.90 (1H, s), 10.82 (1H, s), 7.66 (2H, d), 7.51-7.40 (2H, m), 7.17 (1H, t), 6.95 (2H, d), 3.80-3.70 (4H, m), 3.15-3.01 (4H, m), 2.29 (3H, s)	398
18	H		10.77 (2H, d), 7.65 (2H, d), 7.25 (2H, s), 6.95 (2H, d), 6.73 (1H, s), 3.80-3.70 (4H, m), 3.13-3.04 (4H, m), 2.31 (6H, s)	394
19	H		11.07 (1H, s), 10.80 (1H, s), 7.98 (1H, m), 7.68 (2H, d), 7.30 (1H, m), 6.95 (3H, d), 3.79-3.62 (4H, m), 3.11-3.02 (4H, m)	402
20	H		11.05 (1H, s), 10.80 (1H, s), 7.91 (1H, t), 7.70 (2H, d), 7.23 (1H, m), 7.15 (1H, m), 6.95 (2H, d), 3.80-3.65 (4H, m), 3.11-3.02 (4H, m)	402
21	H		10.80 (1H, s), 10.69 (1H, m), 8.0 (2H, m), 7.60 (2H, d), 7.35 (1H, m), 7.16 (1H, m), 6.90 (2H, d), 3.79-3.63 (4H, m), 3.11-3.01 (4H, m)	402
22	H		10.80 (2H, s), 7.72 (2H, d), 7.55 (2H, d), 7.40 (2H, d), 6.95 (2H, d), 3.77-3.65 (4H, m), 3.15-3.03 (4H, d), 1.25 (9H, s)	422
23	H		10.78 (2H, s), 7.70 (2H, d), 7.45 (2H, d), 7.18 (2H, d), 6.91 (2H, d), 3.78-3.64 (4H, m), 3.11-3.02 (4H, m), 2.22 (3H, s)	380





Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
24	H		10.93 (1H, s), 10.83 (1H, s), 7.65 (2H, d), 7.59 (2H, d), 7.38 (1H, t), 7.06 (1H, t), 6.93 (2H, d), 3.77-3.69 (4H, m), 3.11-3.02 (4H, m)	366
25	H		11.38 (1H, s), 10.89 (1H, s), 8.04 (1H, s), 7.86 (1H, d), 7.71-7.59 (3H, m), 7.41 (1H, d), 6.95 (2H, d), 3.79-3.71 (4H, m), 3.14-3.04 (4H, m)	434
26	H		11.00 (1H, s), 10.84 (1H, s), 7.71-7.58 (4H, m), 7.25 (2H, t), 6.95 (2H, d), 3.79-3.70 (4H, m), 3.14-3.05 (4H, m)	384
27	H		10.83 (1H, s), 10.35 (1H, s), 7.95 (1H, m), 7.65 (2H, d), 7.56 (1H, d), 7.45 (1H, m), 7.20 (1H, m), 6.95 (2H, d), 3.80-3.66 (4H, m), 3.10-3.01 (4H, m)	400
28	H		10.78 (1H, s), 10.21 (1H, s), 7.78 (1H, d), 7.65 (2H, d), 7.36 (1H, s), 7.23 (1H, d), 6.95 (2H, d), 3.79-3.65 (4H, m), 3.11-3.02 (4H, m), 2.27 (3H, s)	414
29	H		10.78 (1H, s), 10.00 (1H, s), 7.64 (2H, d), 7.46 (1H, d), 7.18-7.09 (2H, m), 7.04 (1H, d), 6.94 (2H, d), 3.78-3.70 (4H, m), 3.14-3.04 (4H, m), 2.31 (3H, s), 2.16 (3H, s)	394
30	H		10.70 (1H, s), 9.90 (1H, s), 7.68 (1H, d), 7.58 (2H, d), 7.30 (2H, m), 7.05 (1H, m), 6.89 (2H, d), 3.69-3.59 (4H, m), 3.10-3.02 (4H, m), 2.21 (3H, s)	380








Example	R ⁶	R ⁷	¹ H NMR δ	MS m/e MH ⁺
31	H		11.25 (1H, s), 10.89 (1H, s), 7.66 (2H, d), 7.58-7.50 (1H, m), 7.47-7.32 (2H, m), 6.95 (2H, d), 6.92-6.85 (1H, m), 3.79-3.70 (4H, m), 3.13-3.05 (4H, m)	384
32	H		10.68 (1H, s), 9.95 (1H, s), 7.60 (2H, d), 7.51 (1H, d), 7.40 (1H, d), 7.15 (2H, m), 6.86 (2H, d), 3.79-3.65 (4H, m), 3.30-3.20 (1H, m), 3.00 (4H, m), 1.10 (6H, d)	408
33	H		10.80 (1H, s), 10.70 (1H, s), 7.70 (2H, d), 7.55 (2H, d), 7.00 (4H, dd), 3.82-3.67 (7H, m), 3.15-3.05 (4H, m)	396
34	H		10.80 (2H, s), 7.65 (2H, d), 7.23 (1H, s), 7.00 (1H, d), 6.93 (3H, m), 6.00 (2H, s), 3.80-3.65 (4H, m), 3.11-3.02 (4H, s)	410
35	H		10.75 (1H, s), 10.15 (1H, s), 7.97 (2H, d), 7.70 (3H, m), 7.50 (1H, m), 7.05 (1H, m), 6.95 (2H, d), 3.79-3.63 (4H, m), 3.15-3.03 (4H, m)	492
36	Cl		11.07 (1H, s), 10.89 (1H, s), 7.94 (1H, s), 7.68-7.76 (1H, m), 7.44 (1H, d), 7.38 (1H, s), 7.25 (1H, t), 7.18 (1H, d), 6.86 (1H, d), 3.70-3.78 (4H, m), 2.99-2.92 (4H, m), 2.32 (3H, s)	414
37	F		11.08 (1H, s), 10.80 (1H, s), 8.35-8.26 (1H, m), 8.08-7.95 (2H, m), 7.78 (1H, d), 7.69 (1H, dd), 7.53-7.64 (4H, m), 7.06 (1H, t), 3.75 (4H, t), 2.99 (4H, t)	434









Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
38	Cl		11.17 (1H, s), 10.86 (1H, s), 8.33-8.25 (1H, m), 8.08-7.94 (3H, m), 7.80-7.72 (2H, m), 7.62-7.55 (3H, m), 7.20 (1H, d), 3.79-3.70 (4H, m), 2.99-2.92 (4H, m)	450
39	Me		11.21 (1H, bs), 10.85 (1H, s), 7.74-7.39 (4H, m), 7.37-7.28 (1H, m), 7.01 (1H, d), 3.77-3.67 (4H, m), 2.85-2.75 (4H, m), 2.25 (3H, s)	416
40	Me		10.88-10.77 (2H, m), 7.60 (1H, s), 7.55 (1H, d), 7.42 (1H, d), 7.37 (1H, s), 7.25 (1H, t), 7.01 (1H, d), 6.86 (1H, d), 3.72 (4H, t), 2.80 (4H, t), 2.31 (3H, s), 2.31 (3H, s)	394
41	F		11.06 (1H, s), 10.89 (1H, s), 7.68 (1H, d), 7.57 (1H, d), 7.44 (1H, d), 7.39 (1H, s), 7.28 (1H, t), 7.05 (1H, t), 6.89 (1H, d), 3.75 (4H, t), 2.99 (4H, t), 2.33 (3H, s)	398
42	Cl		11.10 (1H, s), 10.79 (1H, s), 8.05 (1H, t), 7.96 (1H, s), 7.73 (1H, d), 7.38-7.25 (2H, m), 7.22-7.13 (2H, m), 3.80-3.72 (4H, m), 3.02-2.94 (4H, m)	418
43	F		11.08 (1H, s), 10.79 (1H, s), 8.10-7.99 (1H, m), 7.68 (1H, dd), 7.61-7.52 (1H, m), 7.36-7.23 (2H, m), 7.21-7.13 (1H, m), 7.05 (1H, t), 3.79-3.71 (4H, t), 3.02-2.94 (4H, t)	402
44	Me		10.82 (1H, s), 10.73 (1H, s), 8.07-7.98 (1H, m), 7.67-7.50 (2H, m), 7.34-7.20 (2H, m), 7.19-7.09 (1H, m), 7.01 (1H, d), 3.77-3.68 (4H, m), 2.80 (4H, t), 2.25 (3H, s)	398



Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
45	CN		11.22 (1H, s), 10.82 (1H, s), 8.32-8.26 (1H, m), 8.15 (1H, d), 8.05-7.96 (3H, m), 7.78 (1H, d), 7.63-7.54 (3H, m), 7.24 (1H, d), 3.78 (4H, t), 3.13 (4H, t)	441




Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
46	H		11.09 (1H, s), 11.00 (1H, s), 8.52 (1H, d), 8.03-7.93 (2H, m), 7.41-7.33 (1H, m), 7.01-6.94 (1H, m), 6.88 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	402
47	H		10.93 (1H, s), 10.75 (1H, s), 8.51 (1H, s), 8.31-8.22 (1H, m), 8.05-7.91 (3H, m), 7.78 (1H, d), 7.61-7.50 (3H, m), 6.87 (1H, d), 3.74-3.65 (4H, m), 3.44-3.35 (4H, m)	417
48	H		10.96 (1H, s), 10.75 (1H, s), 8.52 (1H, s), 8.04 (1H, t), 7.95 (1H, d), 7.35-7.23 (2H, m), 7.20-7.11 (1H, m), 6.88 (1H, d), 3.74-3.66 (4H, m), 3.45-3.37 (4H, m)	385
49	H		11.23 (1H, s), 10.98 (1H, s), 8.52 (1H, d), 7.96 (1H, dd), 7.53 (1H, dt), 7.43 (1H, dd), 7.38-7.34 (1H, m), 6.93-6.85 (2H, m), 3.71 (4H, t), 3.42 (4H, t)	385

Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
50	H		10.91 (1H, s), 10.85 (1H, s), 8.51 (1H, s), 7.98-7.90 (1H, m), 7.42 (1H, d), 7.38 (1H, s), 7.25 (1H, t), 6.90-6.83 (2H, m), 3.74-3.64 (4H, m), 3.45-3.37 (4H, m), 2.32 (3H, s).	381
51	H		10.92 (1H, s), 10.83 (1H, s), 8.52 (1H, d), 7.98-7.93 (1H, m), 7.54-7.50 (2H, m), 7.43- 7.39 (2H, m), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t), 2.51 (9H, s)	423
52	H		11.22 (1H, s), 10.98 (1H, s), 8.52 (1H, d), 7.98-7.94 (1H, m), 7.74 (1H, t), 7.54-7.50 (1H, m), 7.43 (1H, t), 7.14-7.10 (1H, m), 6.88 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	401
53	H		11.19 (1H, s), 10.96 (1H, s), 8.54 (1H, d), 8.21 (1H, s), 7.99-7.93 (2H, m), 7.90-7.85 (2H, m), 7.67-7.63 (1H, m), 7.51 (1H, t), 7.42 (1H, t), 6.88 (1H, d), 3.72 (4H, t), 3.43 (4H, t)	417
54	Cl		11.27 (2H, bs), 8.61 (1H, d), 8.25 (1H, d), 7.67 (1H, ddd), 7.47 (1H, dd), 7.37-7.29 (1H, m), 3.77-3.69 (4H, m), 3.24-3.17 (4H, m)	437
55	H		11.13 (1H, s), 11.01 (1H, s), 8.52 (1H, d), 7.96 (1H, dd), 7.63 (2H, d), 7.46 (2H, d), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	401
56	Cl		11.26 (1H, broad s), 10.79 (1H, broad s), 8.61 (1H, d), 8.24 (1H, d), 8.01 (1H, t), 7.34-7.11 (3H, m), 3.73 (4H, t), 3.21 (4H, t)	419

Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
57	H		11.23 (1H, s), 10.97 (1H, s), 8.52 (1H, d), 7.97-7.93 (1H, m), 7.70 (1H, ddd), 7.49 (1H, dd), 7.38-7.33 (1H, m), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	403
58	H		11.19 (1H, s), 10.97 (1H, s), 8.52 (1H, d), 7.98-7.94 (2H, m), 7.59-7.55 (1H, m), 7.36 (1H, t), 7.27-7.24 (1H, m), 6.88 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	446 and 447
59	H		11.07 (1H, broad s), 10.97 (1H, s), 8.52 (1H, d), 7.98-7.86 (2H, m), 7.32-7.14 (2H, m), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	403
60	Cl		11.24 (1H, s), 10.91 (1H, s), 8.62 (1H, d), 8.27 (1H, dd), 7.47-7.34 (2H, m), 7.26 (1H, d), 6.87 (1H, d), 3.73 (4H, t), 3.21 (4H, t), 2.31 (3H, s)	415
61	H		10.92 (1H, s), 10.83 (1H, s), 8.52 (1H, d), 7.95 (1H, dd), 7.49 (2H, d), 7.20 (2H, d), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t), 2.28 (3H, s)	381
62	H		10.99 (1H, s), 10.94 (1H, s), 8.52 (1H, d), 7.95 (1H, dd), 7.62 (2H, dd), 7.25 (2H, t), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	385
63	H		10.95 (2H, bs), 8.52 (1H, d), 7.96 (1H, dd), 7.61 (2H, d), 7.40 (2H, t), 7.07 (1H, t), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	367
64	H		10.92 (1H, s), 10.81 (1H, s), 8.52 (1H, d), 7.95 (1H, dd), 7.26-7.24 (1H, m), 7.02 (1H, dd), 6.94 (1H, d), 6.87 (1H, d), 6.02 (2H, s), 3.71 (4H, t), 3.42 (4H, t)	411

Example	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺
65	H		10.96 (1H, s), 10.94 (1H, s), 8.52 (1H, d), 7.96 (1H, dd), 7.33-7.25 (2H, m), 7.18-7.13 (1H, m), 6.87 (1H, d), 6.66 (1H, dd), 3.78 (3H, s), 3.71 (4H, t), 3.42 (4H, t)	397
66	H		10.94 (1H, s), 10.73 (1H, s), 8.51 (1H, d), 8.05-7.92 (2H, m), 7.40 (1H, ddd), 7.21-7.15 (1H, m), 6.87 (1H, d), 3.71 (4H, t), 3.42 (4H, t)	403


Example 72: 5-[(2-Fluorophenyl)amino]-N-[4-(4-isopropylpiperazin-1-yl)phenyl]-1,3,4-oxadiazole-2-carboxamide








- 5 Ethyl 5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxylate (Intermediate 40, 94 mg, 0.37 mmol) was stirred in water (1 mL). Sodium hydroxide (37 mg, 0.925 mmol) was dissolved in water (0.5 mL) and added to the reaction mixture. After stirring for 1 h the reaction was concentrated *in vacuo* to give a yellow solid which was dissolved in DMF (1 mL). [4-(4-Isopropylpiperazin-1-yl)phenyl]amine (WO02/057261) (89 mg, 0.41 mmol)
- 10 dissolved in DMF (1 mL), HOBt (100 mg, 0.74 mmol), and EDAC (142 mg, 0.74 mmol) were added. The reaction was allowed to stir for 3 h then water (20 mL) was added and the reaction mixture thoroughly shaken to give a precipitate. After filtration and drying under *in vacuo* the title compound was obtained as a cream solid (69 mg, 44%):
- ¹H NMR δ 10.78 (1H, s), 8.0 (1H, t), 7.6 (2H, d), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.9 (2H, d), 3.1 (4H, s), 2.6 (4H, s), 1.0 (6H, d);
- 15 MS m/e MH⁺ 425.38







Examples 67-71 and 73-84




The following examples were prepared by the general procedure of Example 1 (method A) using intermediates 19-22, 29-30 or the general procedure of Example 72 (method B) using intermediate 40 and 4-(4-benzylpiperazin-1-yl)aniline or 4-(4-pyridin-4-ylpiperazin-1-yl)aniline (J. Med. Chem., 2003; 46, 1803-1806).




Ex	X	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺	Method
67	NAc	H		10.84 (1H, s), 10.80 (1H, s), 7.65 (2H, d), 7.42 (1H, d), 7.38 (1H, s), 7.25 (1H, t), 6.95 (2H, d), 6.90 (1H, d) 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	421	A
68	NAc	H		10.84 (1H, s), 10.74 (1H, s), 8.22-8.35 (1H, m), 7.9-8.05 (2H, m), 7.80 (1H, d), 7.70 (2H, d), 7.50-7.60 (3H, m), 6.95 (2H, d), 3.60-3.90 (4H, m), 3.10 (4H, d), 2.00 (3H, s)	457	A
69	NMe	H		10.80 (1H, s), 10.72 (1H, s), 8.25-8.35 (1H, m), 8.01 (1H, d), 7.95-8.0 (1H, m), 7.75 (1H, d), 7.70 (2H, d), 7.55-7.65 (3H, m), 6.95 (2H, d), 3.10 (4H, s), 2.40 (4H, s), 2.21 (3H, s)	429	A

Ex	X	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺	Method
70	NAc	H		10.82 (1H, s), 10.72 (1H, s), 8.05 (1H, t), 7.90 (1H, s), 7.60 (2H, d), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.95 (2H, d), 3.6 (4H, s), 3.05 (4H, d), 2.00 (3H, s)	425	A
71	NAc	Me		11.21 (1H, s), 10.86 (1H, s), 7.68 (1H, ddd), 7.62 (1H, s), 7.58-7.50 (1H, m), 7.50-7.40 (1H, m), 7.37-7.29 (1H, m), 7.00 (1H, d), 3.61-3.51 (4H, m), 2.85-2.72 (4H, m), 2.27 (3H, s), 2.03 (3H, s)	457	A
73	NMe	H		10.84 (1H, s), 10.79 (1H, s), 7.70 (2H, d), 7.45 (2H, d), 7.40 (1H, s), 7.25 (1H, t), 6.85-6.95 (3H, m), 3.10 (4H, s), 2.45 (4H, s), 2.25 (3H, s)	393	A
74	NAc	Me		10.86 (1H, s), 10.81 (1H, s), 7.62 (1H, s), 7.54 (1H, d), 7.42 (1H, d), 7.37 (1H, s), 7.25 (1H, t), 7.00 (1H, d), 6.86 (1H, d), 3.62-3.50 (4H, m), 2.85-2.70 (4H, m), 2.31 (3H, s), 2.27 (3H, s), 2.03 (3H, s)	435	A
75	NBoc	F		11.09 (1H, s), 10.79 (1H, broad s), 8.32-8.26 (1H, m), 8.05-7.96 (2H, m), 7.78 (1H, d), 7.70 (1H, dd), 7.63-7.54 (4H, m), 7.07 (1H, t), 3.53-3.44 (4H, m), 2.98-2.91 (4H, m), 2.51 (9H, s)	531 (M-H) ⁺	A

Ex	X	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺	Method
76	NBn	H		10.78 (1H, s), 10.70 (1H, broad s), 8.00 (1H, t), 7.65 (2H, d), 7.1-7.3 (7H, m), 6.90 (2H, d), 3.30 (4H, s), 3.10 (4H, s)	473	B
77	NAc	Me		10.84 (1H, s), 10.74 (1H, s), 8.03 (1H, t), 7.61 (1H, s), 7.54 (1H, d), 7.35-7.21 (2H, m), 7.20-7.10 (1H, m), 7.00 (1H, d), 3.61-3.51 (4H, m), 2.85-2.71 (4H, m), 2.27 (3H, s), 2.03 (3H, s)	439	A
78	NBoc	F		11.07 (1H, s), 10.76 (1H, s), 8.02 (1H, t), 7.76-7.50 (2H, m), 7.34- 7.10 (3H, m), 7.05 (1H, t), 3.50- 3.40 (4H, m), 2.96-2.89 (4H, m), 2.49 (9H, s)	499 (M-H) ⁻	A
79	NMe	H		10.80 (1H, s), 8.05 (1H, t), 7.66 (2H, d), 7.20-7.35 (2H, m), 7.1-7.2 (1H, m), 6.95 (H, d), 3.10 (4H, s), 2.40 (4H, s), 2.30 (2H, s)	397	A
80	NBoc	H		10.82 (1H, s), 10.72 (1H, s), 8.05 (1H, t), 7.60 (2H, d), 7.20-7.30 (2H, m), 7.10-7.20 (1H, m), 6.95 (2H, d), 3.40-3.48 (4H, m), 3.05-3.10 (4H, m), 1.40 (9H, s)	427 [MH- C ₄ H ₈] ⁺	A
81		H			460	B

Ex	X	R ⁶	R ¹	¹ H NMR δ	MS m/e MH ⁺	Method
82	CH ₂	H		10.78 (1H, s), 10.72 (1H, s), 8.25-8.35 (1H, m), 8.05 (1H, d), 7.95-8.0 (1H, m), 7.80 (1H, d), 7.55-7.65 (5H, m), 6.95 (2H, d), 3.05-3.15 (4H, m), 1.45-1.65 (6H, m)	414	A
83	CH ₂	H		10.84 (1H, s), 10.75 (1H, s), 7.60 (1H, d), 7.45 (1H, d), 7.40 (1H, s), 7.25 (1H, t), 6.85-6.95 (3H, m), 3.05-3.15 (4H, m), 1.45-1.65 (6H, m)	378	A
84	CH ₂	H		10.78 (1H, s), 10.72 (1H, s), 8.05 (1H, t), 7.60 (2H, d), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.95 (2H, d), 3.05-3.15 (4H, m), 1.45-1.65 (6H, m)	382	A

Example 87: N-[6-(4-Benzoyl-1-piperazinyl)-3-pyridinyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide



5

5-[(2-Fluorophenyl)amino]-N-(6-piperazin-1-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (intermediate 41, 150 mg, 0.39 mmol), was stirred in DMF (4 mL) and pyridine (1 mL). Benzoyl chloride (60 mg, 0.43 mmol) was added and the reaction stirred for 18 h. Water (20 mL) was added and the reaction mixture thoroughly mixed. The resulting precipitate was filtered, washed with water (2 × 10 mL) then ether (2 × 10 mL) and dried *in vacuo* to give the title compound (130 mg) as a pale purple solid:


¹H NMR δ 10.94 (1H, s), 10.74 (1H, s), 8.50 (1H, s), 8.00 (1H, t), 7.95

10








(1H, d), 7.40 (4H, s), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.90 (2H, d), 3.42-3.83 (8H, m); MS m/e MH⁺ 488.





Examples 85, 86 and 88-94A

The following examples were prepared by the general procedure of Example 1 (method A) using Intermediates 31-33, or the general procedure of Example 72 (method B) using Intermediate 40 and the appropriate aniline, for example *N*²-phenylpyridine-2,5-diamine (J. Med. Chem. 1994, 37, 18-25), [4-(pyridin-2-ylmethoxy)phenyl]amine (DE 3607382), *N*-methyl-*N*-phenylbenzene-1,4-diamine (WO 03059905), 6-butoxypyridin-3-amine (J. Am. Chem. Soc., 1947, 69, 1204), or the general procedure of Example 87 (method C) using Intermediate 41 and the appropriate acid chloride.




Ex	X	R ¹	¹ H NMR (DMSO-d ₆) δ	MS m/e MH ⁺	Method
85	NMe		10.90 (1H, s), 10.76 (1H, s), 8.45-8.50 (1H, m), 8.25-8.35 (1H, m), 7.90-8.05 (3H, m), 7.80 (1H, d), 7.55-7.65 (3H, m), 6.80 (2H, d), 3.40-3.50 (4H, m), 2.35-2.45 (4H, m), 2.20 (3H, s)	430	A
86	NMe		10.88 (1H, s), 10.84 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.45 (1H, d), 7.40 (s, 1H), 7.30 (1H, t), 6.95 (2H, t), 3.40-3.45 (4H, m), 2.35-2.45 (4H, m), 2.20 (3H, s)	394	A

Ex	X	R ¹	¹ H NMR (DMSO-d ₆) δ	MS m/e MH ⁺	Method
88			10.94 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.00 (1H, t), 7.90 (1H, d), 7.50 (1H, q), 7.20-7.35 (5H, m), 7.10-7.20 (1H, m), 6.90 (1H, d), 3.47-3.81 (8H, bd)	506	C
89	NSO ₂ Me		10.95 (1H, s), 8.50 (1H, s), 8.00 (1H, t), 7.90 (1H, d), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.90 (1H, d), 3.55-3.65 (4H, m), 3.15-3.25 (4H, m), 2.90 (3H, s)	462	B
90			10.94 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.00 (1H, t), 7.90-8.00 (1H, m), 7.45-7.55 (2H, m), 7.20-7.35 (4H, m), 7.10-7.20 (1H, m), 6.90 (1H, d), 3.40-3.80 (8H, bs)	506	C
91	NMe		10.90 (1H, s), 10.70 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.90-7.95 (1H, m), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.95 (2H, d), 3.40-3.50 (4H, m), 2.35-2.45 (4H, m), 2.20 (3H, s)	398	A
92	NBoc		10.93 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.90-8.00 (1H, m), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.95 (2H, d), 3.40 (8H, d), 1.40 (9H, s)	484	A

Ex	X	R ¹	¹ H NMR (DMSO-d ₆) δ	MS m/e MH ⁺	Method
93	NAc		10.94 (1H, s), 10.74 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.90-8.00 (1H, m), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.95 (2H, d), 3.50 (6H, m), 3.40-3.50 (2H, m), 2.00 (3H, s)	424	A
94			10.95 (1H, s), 10.74 (1H, s), 8.70 (2H, s), 8.50 (1H, s), 8.00 (1H, t), 7.90-8.00 (1H, m), 7.50 (2H, d), 7.20-7.35 (2H, m), 7.10-7.20 (1H, m), 6.90 (1H, d), 3.70 (2H, s), 3.60 (2H, s), 3.50 (2H, s)	489	C
94 A	NMe			381	A



Examples 95-103

The following examples were prepared by the general procedure of Example 72 (method B) using Intermediate 40 and the appropriate aniline, for example Intermediate 42.




5

Example	R ¹	X	¹ H NMR δ	MS m/e MH ⁺
95	OPh	N	11.22 (1H, s), 10.78 (1H, s), 8.50 (1H, s), 8.20 (1H, d), 8.00 (1H, t), 7.40 (2H, t), 7.10-7.30 (7H, m)	392
96	NPh	N	10.98 (1H, s), 9.00 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.70 (2H, d), 7.10-7.30 (5H, m), 6.80-6.90 (2H, m)	391

Example	R ⁴	X	¹ H NMR δ	MS m/e MH ⁺
97	O-nBu	N		372
98		N	11.26 (1H, s), 8.70 (1H, s), 8.60-8.70 (2H, m), 8.30 (1H, d), 8.00 (1H, t), 7.60 (1H, d), 7.50 (1H, m), 7.10-7.30 (4H, m)	393
99	-(CH ₂) ₄ OH	N	11.18 (1H, s), 8.90 (1H, s), 8.00-8.10 (2H, m), 7.20-7.35 (4H, m), 7.10-7.20 (1H, m), 4.40 (1H, d), 3.40 (2H, s), 2.70 (2H, t), 1.60-1.70 (2H, m), 1.35-1.45 (2H, m)	372
100	OBn	C	10.80 (1H, s), 10.76 (1H, s), 8.00 (1H, t), 7.50 (1H, s), 7.20-7.40 (9H, m), 6.80 (1H, d), 5.10 (2H, s)	405
101	nBu	C	10.92 (1H, s), 10.75 (1H, s), 8.00 (1H, t), 7.70 (2H, d), 7.20-7.35 (2H, m), 7.10-7.20 (3H, d), 2.50 (2H, t), 1.50-1.60 (2H, m), 1.20-1.40 (2H, m), 0.90 (3H, t)	355
102		C	11.07 (1H, s), 8.60 (1H, d), 7.95-8.05 (2H, m), 7.90 (1H, t), 7.70 (1H, d), 7.50 (1H, d), 7.20-7.40 (4H, m), 7.10-7.20 (1H, m), 5.20 (2H, s)	440
103	N(Me)Ph	C		404

Example 104: N-Biphenyl-4-yl-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide




- 5 *N*-(4-Iodophenyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide
(Intermediate 35), 150 mg, 0.36 mmol), phenyl boronic acid (35 mg, 0.29 mmol),

potassium phosphate (230 mg, 1.44 mmol), and tetrakis(triphenyl phosphine)palladium(0) (22 mg, 0.02 mmol) were stirred in DME (1.2 mL) and water (0.6 mL), in a microwave tube, and heated at 110 °C for 3 h. The reaction mixture was partitioned between ethyl acetate (10 mL) and water (10 mL), and extracted with ethyl acetate (2 × 10 mL). The organics were combined, and dried by passing through a Varian Chem elut column. The filtrate was concentrated *in vacuo* and purified by preparative HPLC to afford the title compound as a cream solid (25 mg, 19%):

¹H NMR δ 11.20 (1H, s), 10.78 (1H, s), 8.10 (1H, t), 7.90 (2H, d), 7.65-7.75 (4H, m), 7.50 (2H, t), 7.20-7.35 (3H, m), 7.10-7.20 (1H, m); MS m/e MH⁺ 375.

Example 105: Methyl [(trans-4-[4-({[5-(pyridin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)phenyl]cyclohexyl]acetate



2-Isothiocyanatopyridine (163 mg) was added to a stirred suspension of methyl [(trans-4-(4-({[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl]acetate (Intermediate 43) (333 mg) in DMF (3 mL) and the mixture was heated to 40°C and stirred for 60 minutes. EDAC (230 mg) was added and the resulting mixture was heated to 85°C and stirred for 16 hours. Water (3 mL) was added. The precipitate was filtered off and washed with water, Et₂O then dried under vacuum overnight to give the title compound (333 mg) as a solid; ¹H NMR δ 11.60 (1H, bs), 10.93 (1H, s), 8.30 (1H, m), 7.88 (2H, m), 7.70 (2H, d), 7.23 (2H, d), 7.07 (1H, m), 3.61 (3H, s), 2.45 (1H+DMSO, m), 2.25 (2H, d), 1.80 (5H, m), 1.46 (2H, m), 1.13 (2H, m); MS m/e MH⁺ 436.

Examples 106-111

Using an analogous procedure to that described for Example 105 with the appropriate starting material (SM) and using the requisite isothiocyanate the following Examples were prepared:

Ex.	SM	Name	¹ H NMR	MS m/e M+H ⁺
106 ¹	Int. 43	methyl (<i>trans</i> -4-{4-[(5-[(3-ethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexyl)acetate	10.94 (1H, s), 10.92 (1H, s), 7.69 (2H, d), 7.30-7.19 (4H, m), 7.13 (1H, d), 6.63 (1H, d), 4.05 (2H, quartet), 3.61 (3H, s), 2.45 (1H+DMSO, m), 2.25 (2H, d), 1.79 (5H, m), 1.45 (2H, m), 1.35 (3H, t), 1.13 (2H, m).	479
107	Int. 43	methyl [<i>trans</i> -4-(4-{[(5-{[3-(benzyloxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetate	10.97 (1H, s), 10.93 (1H, s), 7.69 (2H, d), 7.48 (2H, d), 7.41 (2H, t), 7.36-7.28 (3H, m), 7.22 (2H, d), 7.16 (1H, d), 6.73 (1H, d), 5.12 (2H, s), 3.60 (3H, s), 2.45 (1H+DMSO, m), 2.25 (2H, d), 1.80 (5H, m), 1.47 (2H, m), 1.15 (2H, m)	541
108 ^{2,3}	Int. 43'	methyl (<i>trans</i> -4-{4-[(5-[(4-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexyl)acetate		469
109 ^{2,3}	Int. 43'	methyl (<i>trans</i> -4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexyl)acetate		489
110	Int. 44	methyl <i>trans</i> -4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexanecarboxylate	10.94 (1H, s), 10.75 (1H, s), 8.04 (1H, t), 7.69 (2H, d), 7.34-7.14 (5H, m), 3.62 (3H, s), 2.60-2.33 (2H+DMSO, m), 2.01 (2H, m), 1.84 (2H, m), 1.49 (4H, m)	439

Ex.	SM	Name	¹ H NMR	MS m/e M+H ⁺
111	Int. 43	methyl (<i>trans</i> -4-{4-[(5-[2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexyl)acetate	10.49 (1H, br.s), 8.02 (1H, t), 7.67 (2H, d), 7.28 (2H, d), 7.1-6.98 (2H, m), 6.79 (1H, br.s), 3.60 (3H, s), 2.42 (1H, m), 2.23 (2H, d), 1.87-1.67 (5H, m), 1.45 (2H, m), 1.12 (2H, m)	453

¹O-(pentafluorophenyl) (3-ethoxyphenyl)thiocarbamate (Intermediate 48) was used instead of the corresponding isothiocyanate.


²Prepared in similar manner to Example 105 except a mixture of methyl [*trans*-4-(4-[[hydrazino(oxo)acetyl]amino]phenyl)cyclohexyl]acetate and ethyl [*trans*-4-(4-[[hydrazino(oxo)acetyl]amino]phenyl)cyclohexyl]acetate used.

³Contains 17% ethyl (*trans*-4-{4-[(5-[(4-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexyl)acetate by LCMS; MS m/e M+H⁺ 483.

⁴Contains 26% ethyl (*trans*-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl)cyclohexyl)acetate by LCMS; MS m/e M+H⁺ 503.

10

Example 112: [*trans*-4-{4-[(5-(Pyridin-2-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl]cyclohexyl}acetic acid



Lithium hydroxide monohydrate (60 mg) in water (2 mL) was added to a stirred suspension of methyl [*trans*-4-{4-[(5-(pyridin-2-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl]cyclohexyl)acetate (Example 105, 309 mg) in THF (7 mL). After 16 hours the solution was heated to 40°C and stirred for a further 5 hours then allowed to cool to ambient temperature. The THF was removed *in vacuo* and the residue was acidified with 2M HCl. The precipitate was filtered off, washed with ether, warm DCM and warm ether then dried overnight. The resultant solid was purified by HPLC using a gradient of 5-25% MeCN in water with 1% ammonia as the eluent to give the title compound (119 mg) as a solid; ¹H NMR δ 10.80 (1H, bs), 10.91 (1H, s), 8.31 (1H, d), 7.86

(2H, m), 7.69 (2H, d), 7.23 (2H, d), 7.05 (1H, m), 2.45 (1H+DMSO, m), 2.14 (2H, d), 1.82 (5H, m), 1.47 (2H, m), 1.13 (2H, m); MS m/e MH^+ 422.

Examples 113-117

- 5 The following examples were prepared by the general procedure of Example 112 using the appropriate starting material selected from Examples 106-110.

Example	Name	^1H NMR	MS m/e $M+H^+$
113 ^{1,5}	(<i>trans</i> -4-{4-[(5-{(3-ethoxyphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	11.98 (1H, s), 10.94 (1H, s), 10.92 (1H, s), 7.69 (2H, d), 7.30-7.22 (4H, m), 7.13 (1H, d), 6.64 (1H, d), 4.04 (2H, quartet), 2.45 (1H+DMSO, m), 2.15 (2H, d), 1.82 (5H, m), 1.46 (2H, m), 1.35 (3H, t), 1.13 (2H, m).	464
114 ^{1,6}	[<i>trans</i> -4-(4-{[(5-{(3-benzyloxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid	11.98 (1H, bs), 10.98 (1H, s), 10.93 (1H, s), 7.69 (2H, d), 7.48 (2H, d), 7.41 (2H, m), 7.36-7.27 (3H, m), 7.22 (2H, d), 7.16 (1H, d), 6.74 (1H, dd), 5.12 (2H, s), 2.45 (1H+DMSO, m), 2.15 (2H, d), 1.81 (5H, m), 1.46 (2H, m), 1.15 (2H, m).	527
115 ^{2,4}	(<i>trans</i> -4-{4-[(5-{(4-chlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	11.98 (1H, s), 11.13 (1H, s), 10.94 (1H, s), 7.70 (2H, d), 7.63 (2H, d), 7.46 (2H, d), 7.23 (2H, d), 2.45 (1H+DMSO, m), 2.15 (2H, d), 1.80 (5H, m), 1.45 (2H, m), 1.13 (2H, m)	455

Example	Name	¹ H NMR	MSm/e M+H ⁺
116 ^{3,4}	Sodium (<i>trans</i> -4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetate	10.24 (1H, s), 8.16 (1H, m), 7.67 (2H, d), 7.15 (2H, d), 7.08 (1H, m), 2.43 (1H+DMSO, m), 1.90-1.60 (7H, m), 1.40 (2H, m), 1.00 (2H, m).	475
117	<i>trans</i> -4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexanecarboxylic acid	12.00 (1H, s), 10.94 (1H, s), 10.76 (1H, s), 8.04 (1H, t), 7.70 (2H, d), 7.35-7.13 (5H, m), 2.50 (1H+DMSO, m), 2.28 (1H, m), 2.00 (2H, m), 1.85 (2h, m), 1.48 (4H, m)	425

¹Final compound did not require HPLC purification

²HPLC Purification used an acidic modifier, 0.2% TFA, in place of 1% ammonia


³The crude material was partitioned between EtOAc and saturated aqueous Na₂CO₃ solution. A precipitate was filtered off, dried under vacuum to give title compound.

5 ⁴Prepared from a mixture of methyl and ethyl esters of appropriate starting material via Intermediate 43'.

⁵Reaction carried in EtOH at ambient temperature for 16 hours, then 5 hours at 40°C. Worked up in usual manner. Crude material was then hydrolysed further in THF for 16 hours.

10 ⁶Stirred for 64 hours at ambient temperature.


Example 118: (*trans*-4-{4-[(5-[(2-Fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}cyclohexyl)acetic acid



15 Lithium hydroxide monohydrate (45 mg, 1.06 mmol) was added to a solution of Example 111 (240 mg, 0.53 mmol) in MeOH/H₂O (1:1) (10 mL) and the reaction mixture was stirred for 18 hours. The mixture was then filtered, cooled in an ice bath and acidified to pH 5 with 2M HCl. The resulting precipitate was filtered off and dried under high vacuum to give the


title compound as a solid (127mg, 55%); ^1H NMR δ 11.90 (1H, br.s), 10.85 (s, 1H), 10.69 (1H, s), 7.97 (1H, t), 7.60 (2H, d), 7.30-7.03 (5H, m), 2.38 (1H, m), 2.09 (2H, d), 1.80-1.60 (5H, m), 1.38 (2H, m), 1.04 (2H, m); MS m/e MH^+ 439.

5 **Example 119: 5-[(2-Fluorophenyl)amino]-N-{4-[4-(2-hydroxyethyl)cyclohexyl]phenyl}-1,3,4-oxadiazole-2-carboxamide**



Borane (1 mL of a 1M solution in THF) was added to a stirred solution of (4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid (Intermediate 43, 79 mg) in THF (2 mL). The reaction mixture was stirred for 16 hours. The mixture was evaporated and the residue dissolved in DCM (10 mL) then washed with brine (10 mL), dried and concentrated *in vacuo* to give a solid. This solid was purified by HPLC eluting with a gradient of acetonitrile/water containing 0.2% TFA to give the title compound (35 mg) as a solid, a *ca.* 2:1 *trans:cis* isomeric mixture; ^1H NMR δ 10.90 (1H, s), 10.72 (1H, s), 8.01 (1H, t), 7.65 (2H, m), 7.33-7.12 (5H, m), 4.28 (1H, m), 3.46 (2H, bs), 2.58-2.40 (1H+DMSO, m), 1.80 (3H, bs), 1.68-1.50 (4H, m), 1.49-1.31 (3H, m), 1.05 (1H, m); MS m/e MH^+ 425.

20 **Example 120: N-{4-[trans-4-(2-Amino-2-oxoethyl)cyclohexyl]phenyl}-5-[(3-ethoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide**



Ammonia (0.06 mL of a 28-30% wt% solution in water) was added to a stirred mixture of (*trans*-4-{4-[(5-[(3-ethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid (Example 113, 65 mg), EDAC (32 mg), and HOBt (23 mg) in DMF (1 mL). The mixture was stirred for 16 h then was diluted with DMF (2 mL) and heated to 100°C for 30 minutes in an Emrys Optimizer™ microwave. The crude mixture was purified directly by HPLC using a gradient of 25-75% acetonitrile /

water with 1% ammonia as the eluent to give the title compound (10 mg) as a solid; ^1H NMR δ 10.82 (2H, bs), 7.68 (2H, d), 7.22 (5H, m), 7.08 (1H, 1H), 6.67 (1H, s), 6.56 (1H, m), 4.00 (2H, quartet), 2.50 (1H+DMSO, m), 1.96 (2H, d), 1.83-1.66 (5H, m), 1.42 (2H, m), 1.33 (3H, t), 1.08 (2H, m); MS m/e MH^+ 474.

5


Examples 121-126

The following examples were prepared by the general procedure of Example 105, using the indicated Intermediates as Starting Materials and commercially available isothiocyanates:

Ex.	SM	Name	^1H NMR	MS MH^+
121	Int. 46	[(2-fluorophenyl)amino]- <i>N</i> -(3-phenoxypropyl)- 1,3,4-oxadiazole-2- carboxamide	10.68 (1H, s), 9.14 (1H, m), 8.02 (1H, t), 7.30 (4H, m), 7.16 (1H, m), 6.95 (3H, m), 4.03 (2H, t), 3.43 (2H, m), 2.00 (2H, m)	357
122	Int. 46	5-[(3- methylphenyl)amino]- <i>N</i> - (3-phenoxypropyl)-1,3,4- oxadiazole-2-carboxamide	10.75 (1H, s), 9.10 (1H, m), 7.41 (2H, m), 7.28 (3H, m), 6.93 (3H, m), 6.86 (2H, d), 4.03 (2H, t), 3.44 (2H, m), 2.31 (3H, s), 1.99 (2H, m)	353
123	Int. 46	5-[(3-(benzyloxy)phenyl] amino]- <i>N</i> -(3- phenoxypropyl)-1,3,4- oxadiazole-2-carboxamide	10.88 (1H, s), 9.13 (1H, t), 7.53- 7.25 (9H, m), 7.13 (1H, d), 6.93 (3H, m), 6.70 (1H, d), 5.10 (2H, s), 4.02 (2H, t), 3.42 (2H, m), 2.00 (2H, m)	445
124	Int. 47	<i>N</i> -(4-cyclohexylphenyl)-5- [(2-fluorophenyl)amino]- 1,3,4-oxadiazole-2- carboxamide	10.92 (1H, s), 10.73 (1H, s), 8.05 (1H, t), 7.20 (2H, d), 7.37-7.10 (5H, m), 2.50 (1H+ DMSO, bs), 1.88- 1.67 (5H, m), 1.48-1.17 (5H, m)	381
125	Int. 47	<i>N</i> -(4-cyclohexylphenyl)-5- [(3,4-difluorophenyl) amino]-1,3,4-oxadiazole- 2-carboxamide	11.25 (1H, s), 10.96 (1H, s), 7.70 (3H, m), 7.49 (1H, m), 7.37 (1H, m), 7.23 (2H, d), 2.50	399

Ex.	SM	Name	¹ H NMR	MS MH ⁺
126	Int. 47	5-{[3-(benzyloxy)phenyl]amino}- -N-(4-cyclohexylphenyl)- 1,3,4-oxadiazole-2- carboxamide	10.96 (2H, d), 7.70 (2H, d), 7.50- 7.26 (7H, m), 7.20 (2H, d), 7.15 (1H, d), 6.23 (1H, d), 5.12 (2H, s), 2.50 (1H+DMSO, bs), 1.88-1.67 (5H, m), 1.48-1.20 (5H, m)	469

Example 127: N-[4-(4-Acetyl piperazin-1-yl)phenyl]-5-[(3-isopropoxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide




- 5 Thiophosgene (46 μ L, 0.60 mmol) was added dropwise to a stirred solution of 3-isopropoxy aniline (82 mg, 0.543 mmol) and calcium carbonate (272 mg, 2.72 mmol) in DCM/H₂O (1:1) (4 mL). This was stirred at room temperature for 1 h. The phases were separated and the organic layer was dried and concentrated *in vacuo*. The residue was dissolved in DMF (3 mL) and added to a stirred solution of N-[4-(4-acetyl piperazin-1-
- 10 yl)phenyl]-2-hydrazino-2-oxoacetamide (Intermediate 21, 100 mg, 0.33 mmol) in DMF at 85°C. This mixture was stirred at 85°C for 1 h then EDAC (105 mg, 0.54 mmol) was added. The mixture was stirred at 100°C for a further 1 h then allowed to cool and treated with H₂O (10 mL). The resulting precipitate was filtered and washed sequentially with DMSO/MeCN/H₂O (7:2:1) (5 mL) and ether (5 mL) to give the title compound as a white
- 15 solid (12 mg, 8%); ¹H NMR δ 10.90 (1H, s), 10.82 (s, 1H), 7.67 (2H, d), 7.25 (2H, d), 7.10 (1H, d), 6.97 (2H, d), 6.63 (1H, d), 4.59 (1H, m), 3.58 (4H, br. s), 3.15 (2H, m), 3.09 (2H, m), 2.04 (3H, s), 1.32 (3H, s), 1.30 (3H, s); MS m/e MH⁺ 465.

Examples 128-129


- 20 The following examples were prepared by the general procedure of Example 105 using Intermediate 21 and the appropriate isothiocyanate

Example 128: *N*-[4-(4-Acetylpiperazin-1-yl)phenyl]-5-[4-(methylthio)phenylamino]-1,3,4-oxadiazole-2-carboxamide




¹H NMR δ 10.95 (1H, s), 10.81 (1H, s), 7.66 (2H, d), 7.57 (2H, d), 7.34 (2H, d), 6.97 (2H, d), 3.58 (4H, br. s), 3.28 (3H, s), 3.14 (2H, m), 3.07 (2H, m), 2.05 (3H, s); MS m/e MH⁺ 453.

Example 129: *N*-[4-(4-Acetylpiperazin-1-yl)phenyl]-5-[4-(benzyloxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide



¹H NMR at 373K δ 10.25 (2H, s), 7.62 (2H, d), 7.48 (2H, d), 7.44 (2H, m), 7.39 (2H, t), 7.32 (1H, m), 7.04 (2H, d), 6.94 (2H, d), 5.10 (2H, s), 3.60 (4H, m), 3.15 (4H, m), 2.03 (3H, s); MS m/e MH⁺ 513.

Example 130: 5-[5-Bromo-2,4-difluorophenylamino]-*N*-(4-morpholin-4-ylphenyl)-1,3,4-oxadiazole-2-carboxamide




2,4-Difluoro-5-bromoaniline (208 mg, 1.04 mmol) was added to a solution of thiocarbonyldiimidazole (185 mg 1.04 mmol) in DMF (10 mL) and the mixture was stirred for 16 h. 2-Hydrazino-*N*-(4-morpholin-4-ylphenyl)-2 oxoacetamide (Intermediate 27, 211 mg, 0.8 mmol) was added and the mixture was stirred at 80°C until a clear solution was obtained. EDAC (306 mg, 1.6 mmol) was added and stirring was continued for a further 1 h. The mixture was cooled then diluted with water (25 mL) and the resulting precipitate was filtered and dried to give the title compound the title compound (135 mg) as a solid;






¹H NMR δ 11.09 (1H, s), 10.88 (1H, s), 8.45-8.33 (1H, m), 7.74-7.57 (3H, m), 6.95 (2H, d), 3.83-3.68 (5H, m), 3.15-2.99 (5H, m); MS m/e MH⁺ 482.

Examples 131-140


The following examples were prepared by the general procedure of Example 130 using either Intermediate 27 (for X = N) or Intermediate 18 (for X = CH) and the appropriate aniline.



Example	X	R	¹ H NMR	MS m/e M+H
131	N		11.15 (1H, s), 10.95 (1H, s), 8.58-8.48 (2H, m), 7.97-7.90 (1H, dd), 7.78 (1H, t), 6.87 (1H, d), 3.77-3.63 (4H, m), 3.48-3.38 (4H, m)	471
132	N		11.39 (1H, s), 11.00 (1H, s), 8.52 (1H, d), 8.03 (1H, s), 7.99-7.93 (1H, m), 7.86 (1H, d), 7.65 (1H, t), 7.42 (1H, d), 6.87 (1H, d), 3.77-3.64 (4H, m), 3.52-3.35 (4H, m)	435
133	N		11.90 (1H, s), 11.00 (1H, s), 8.51 (1H, d), 8.14 (1H, d), 7.95 (1H, d), 7.71 (1H, s), 6.92-6.84 (2H, m), 3.76-3.62 (4H, m), 3.50-3.38 (4H, m), 2.40 (3H, s)	382
134	CH		11.25 (1H, s), 10.89 (1H, s), 8.63-8.52 (1H, m), 7.67 (2H, d), 7.60-7.49 (2H, m), 6.95 (2H, d), 3.84-3.66 (4H, m), 3.13-3.04 (4H, m)	452
135	CH		11.37 (1H, s), 10.89 (1H, s), 8.08-7.98 (1H, m), 7.93-7.84 (1H, m), 7.68 (2H, d), 7.58 (1H, t), 6.95 (2H, d), 3.82-3.70 (4H, m), 3.13-3.04 (4H, m)	452


Example	X	R	¹ H NMR	MS m/e M+H
136	N		10.96 (1H, s), 10.88 (1H, s), 8.53 (1H, d), 7.99-7.91 (1H, m), 7.31-7.20 (2H, m), 7.13-7.07 (1H, m), 6.87 (1H, d), 6.66- 6.60 (1H, m), 4.65-4.54 (1H, m), 3.75- 3.66 (4H, m), 3.46-3.37 (4H, m), 1.30 (6H, d)	425
137	N		11.10 (1H, s), 10.96 (1H, s), 8.52 (1H, d), 7.98-7.92 (1H, m), 7.45-7.27 (6H, m), 7.18 (1H, t), 7.07 (2H, d), 6.87 (1H, d), 3.74-3.67 (4H, m), 3.45-3.38 (4H, m)	459
138	N		11.36 (1H, s), 11.09 (1H, s), 8.53 (1H, d), J = 10.1 Hz, 8.07-8.01 (1H, m), 7.72 (1H, s), 7.59-7.50 (m2H, m), 7.08-6.99 (2H, m), 3.75-3.69 (4H, m), 3.49-3.44 (4H, m)	451
139	N		11.05 (1H, s), 10.98 (1H, s), 8.66-8.63 (1H, m), 8.53 (1H, d), 8.05-8.01 (1H, m), 7.99-7.93 (1H, m), 7.62 (1H, d), 7.48- 7.44 (1H, m), 7.36 (1H, t), 7.32 (1H, t), 7.21-7.17 (1H, m), 6.99 (1H, d), 6.78- 6.73 (1H, m), 5.25 (2H, s), 3.75-3.70 (4H, m), 3.48-3.43 (4H, m)	474
140	N		10.98 (2H, s), 8.52 (1H, d), 7.98-7.93 (1H, m), 7.29 (1H, d), 7.27-7.23 (1H, m), 7.16-7.11 (1H, m), 6.87 (1H, d), 6.67- 6.61 (1H, m), 4.04 (2H, q), 3.75-3.67 (4H, m), 3.45-3.38 (4H, m), 1.35 (3H, t)	411

Example 141: *N*-[4-(4-Acetylpiperazin-1-yl)phenyl]-5-[3-ethoxyphenylamino]-1,3,4-oxadiazole-2-carboxamide



N-[4-(4-Acetylpiperazin-1-yl)phenyl]-2-hydrazino-2-oxoacetamide (Intermediate 21, 153 mg, 0.5 mmol) was added to a solution of *O*-(pentafluorophenyl) (3-ethoxyphenyl)thiocarbamate (Intermediate 48, 182 mg 0.5 mmol) in DMF (7 mL). The mixture was stirred at 60°C for 30 min then PS-CDI (500 mg 1.0 mmol) was added and stirring was continued for a further 2 hours. The mixture was allowed to cool then filtered to remove the resin, which was washed with a further 2 x 10 mL DMF. The combined
 10 filtrates were concentrated by evaporation to give a solid that was then triturated with 1:1 acetonitrile:water to give the title compound (220 mg) as a solid; ¹H NMR δ 10.95 (1H, s), 10.85 (1H, s), 7.67 (2H, d), 7.33-7.22 (2H, m), 7.12 (1H, d), 6.97 (2H, d), 6.64 (1H, d), 4.04 (2H, q), 3.65-3.51 (4H, m), 3.11 (4H, d), 2.07 (3H, s), 1.35 (3H, t); MS m/e MH⁺ 451.

Example 142: 5-(Biphenyl-3-ylanino)-*N*-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide




Anisole (0.14 mL, 1.30 mmol) and a mixture of TFA:water (9:1) (3 mL) was added to (5-(biphenyl-3-yl){2-(trimethylsilyl)ethoxy)methyl}amino)-*N*-(6-morpholin-4-ylpyridin-3-yl)-*N*-{2-(trimethylsilyl)ethoxy)methyl}-1,3,4-oxadiazole-2-carboxamide (Intermediate 49, 180 mg, 0.28 mmol). After 30 minutes the mixture was by diluted with DCM (40 mL) and washed with 1M aqueous NaHCO₃ (20 mL). The DCM layer was separated and MeOH (5 mL) was added. The aqueous was again extracted with MeOH:DCM (1:9) (4 x 10 mL). DMF (5 mL) was added to the aqueous and the aqueous was extracted with
 25 MeOH:DCM (1:9) (2 x 10 mL). All organics were combined, dried, filtered and evaporated to give a solid which was triturated with hot Et₂O (10 mL) and washed with Et₂O (2 x 5 mL) then hexane (2 x 5 mL). Chromatography through silica using 0-100% DCM/EtOAc then 10% MeOH/DCM as eluent gave the title compound (30 mg, 24%) as a


solid; $^1\text{H NMR}$ δ 11.10 (1H, s), 10.98 (1H, s), 8.53 (1H, d), 7.99-7.93 (1H, m), 7.91 (1H, s), 7.67-7.61 (3H, m), 7.54-7.47 (3H, m), 7.41 (1H, t), 7.35 (1H, d), 6.88 (1H, d), 3.74-3.69 (4H, m), 3.45-3.40 (4H, m); MS m/e MH^+ 443.


5 Examples 143-147

The following examples were prepared by the general procedure of Example 1 using the indicated Intermediates and commercially available isothiocyanates:



Ex.	R	Int.	$^1\text{H NMR}$	MS m/e -M+H
143		27	11.00 (2H, s), 8.53 (1H, d), 7.99-7.93 (1H, m), 7.51-7.26 (7H, m), 7.19-7.13 (1H, m), 6.87 (1H, d), 6.76-6.72 (1H, m), 5.13 (2H, s), 3.73-6.8 (4H, m), 3.44-3.40 (4H, m)	473
144		27	11.18 (1H, s), 10.95 (1H, s), 8.52 (1H, d), 8.31-8.26 (1H, m), 7.98-7.93 (1H, m), 7.89-7.85 (1H, m), 7.64 (1H, d), 7.55 (1H, t), 6.87 (1H, d), 4.33 (2H, c), 3.73-3.66 (4H, m), 3.44-3.39 (4H, m), 1.33 (3H, t)	439
145		27	11.16 (1H, s), 10.97 (1H, s), 8.53 (1H, d), 8.25-8.23 (1H, m), 7.98-7.93 (1H, m), 7.84-7.81 (1H, m), 7.60 (1H, d), 7.51 (1H, t), 6.87 (1H, d), 3.73-3.68 (4H, m), 3.44-3.39 (4H, m), 1.56 (9H, s)	467


Ex.	R	Int.	¹ H NMR	MS m/e M+H
146		27	11.14 (1H, s), 10.98 (1H, s), 8.52 (1H, d), 8.02 (1H, t), 7.98-7.93 (1H, m), 7.62-7.57 (1H, m), 7.42 (1H, d), 7.20 (1H, t), 6.87 (1H, d), 3.74-3.68 (4H, m), 3.45-3.38 (4H, m)	493




Example	SM	¹ H NMR	MS m/e M+H
147	Int. 58	11.07 (1H, s), 10.76 (1H, s), 8.59-8.57 (1H, m), 8.24-8.20 (1H, m), 8.10-7.90 (3H, m), 7.75 (1H, d), 7.60-7.50 (3H, m), 6.8 (1H, d), 3.8 (3H, s)	362

Examples 148-296








The following examples were prepared by the general procedure of Example 1, using the indicated Intermediate as starting material (SM) in conjunction with the appropriate commercially available isothiocyanate.








Starting material = Intermediate 24:

Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
148		11.07 (1H, s), 11.03 (1H, br s), 7.71-7.55 (4H, m), 7.28-7.21 (2H, m), 7.05 (1H, t), 3.77-3.72 (4H, m), 3.01-2.97 (4H, m)	402













Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
149		11.12 (2H, s), 8.03-7.97 (1H, m), 7.68 (1H, dd), 7.57 (1H, dd), 7.37 (1H, ddd), 7.06 (1H, t), 7.01-6.94 (1H, m), 3.77-3.73 (4H, m), 3.02-2.97 (4H, m)	420
150		11.07 (1H, s), 11.07 (1H, br s), 7.71-7.55 (4H, m), 7.43-7.37 (2H, m), 7.10-7.03 (2H, m), 3.77-3.73 (4H, m), 3.01-2.97 (4H, m)	384
151		11.25 (1H, br s), 11.10 (1H, s), 7.68 (1H, dd), 7.59-7.50 (2H, m), 7.47-7.40 (1H, m), 7.38-7.34 (1H, m), 7.06 (1H, t), 6.89 (1H, td), 3.77-3.72 (4H, m), 3.02-2.97 (4H, m)	402
152		11.03 (1H, s), 10.22 (1H, br s), 7.75-7.64 (2H, m), 7.56 (1H, dd), 7.28-7.23 (2H, m), 7.13-7.02 (2H, m), 3.77-3.72 (4H, m), 3.01-2.96 (4H, m), 2.30 (3H, s)	398
153		11.07 (1H, s), 11.07 (1H, br s), 8.06-7.95 (1H, m), 7.67 (1H, dd), 7.56 (1H, dd), 7.43-7.36 (1H, m), 7.22-7.15 (1H, m), 7.05 (1H, t), 3.77-3.72 (4H, m), 3.02-2.96 (4H, m)	420
154		11.10 (2H, br s), 7.89-7.86 (1H, m), 7.68 (1H, dd), 7.59-7.55 (2H, m), 7.36 (1H, t), 7.28-7.23 (1H, m), 7.06 (1H, t), 3.77-3.72 (4H, m), 3.02-2.96 (4H, m)	462 and 464
155		11.07 (2H, br s), 7.71-7.60 (3H, m), 7.57 (1H, dd), 7.49-7.44 (2H, m), 7.05 (1H, t), 3.77-3.72 (4H, m), 3.01-2.97 (4H, m)	418 and 420

Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
156		11.09 (2H, br s), 8.21 (1H, d), 7.95 (1H, d), 7.87 (2H, dd), 7.73-7.63 (2H, m), 7.58 (1H, dd), 7.54-7.48 (1H, m), 7.42 (1H, dd), 7.07 (1H, t), 3.78-3.73 (4H, m), 3.02-2.97 (4H, m)	434
157		11.09 (2H, s), 7.92-7.86 (1H, m), 7.68 (1H, dd), 7.56 (1H, dd), 7.31-7.24 (1H, m), 7.22-7.14 (1H, m), 7.06 (1H, t), 3.77-3.72 (4H, m), 3.01-2.97 (4H, m)	420
158		11.25 (1H, br s), 11.25 (1H, s), 7.73-7.65 (2H, m), 7.57 (1H, dd), 7.52-7.44 (1H, m), 7.37-7.32 (1H, m), 7.05 (1H, t), 3.77-3.72 (4H, m), 3.02-2.97 (4H, m)	420
159		11.21 (1H, br s), 11.09 (1H, s), 8.29-8.26 (1H, m), 7.87 (1H, ddd), 7.72-7.63 (2H, m), 7.59-7.53 (2H, m), 7.06 (1H, t), 4.35 (2H, q), 3.77-3.72 (4H, m), 3.02-2.96 (4H, m), 1.34 (3H, t)	456












Starting material = Intermediate 28:

Example	R ²	MS m/e MH ⁺
160		419


Example	R ²	MS m/e MH ⁺
161		415
162		435
163		431
164		431
165		437
166		401
167		479 and 481
168		419
169		415

- 169 -

~~12/0~~


Example	R ²	MS m/e MH ⁺
170		437
171		435
172		435
173		451
174		431
175		457
176		437
177		473
178		451

- 170 -








Starting material = Intermediate 59


Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
179		10.85 (1H, br s), 10.77 (1H, s), 7.63 (2H, d), 7.45-7.35 (2H, m), 7.25 (1H, t), 6.96-6.82 (3H, m), 3.74-3.61 (2H, m), 3.58-3.49 (2H, m), 2.31 (3H, s), 2.23 (2H, t), 1.14 (6H, d)	408
180		11.20 (1H, br s), 10.81 (1H, s), 7.73-7.59 (3H, m), 7.51-7.39 (1H, m), 7.36-7.28 (1H, m), 6.92 (2H, d), 3.74-3.62 (2H, m), 3.57-3.49 (2H, m), 2.23 (2H, t), 1.14 (6H, d)	430




5 Starting material = Intermediate 60:

Example	R ²	¹ H NMR δ	MS m/e MH ⁺
181		10.69 (2H, s), 8.02 (1H, t), 7.56 (2H, dd), 7.33-7.09 (3H, m), 6.72 (2H, dd), 3.68-3.45 (6H, m), 3.35-3.24 (2H, m), 1.98-1.72 (5H, m)	439
182		10.80 (1H, s), 10.66 (1H, s), 7.57 (2H, dd), 7.44-7.34 (2H, m), 7.23 (1H, t), 6.86 (1H, d), 6.72 (2H, dd), 3.68-3.45 (6H, m), 3.35-3.23 (2H, m), 2.30 (3H, s), 1.98-1.72 (5H, m)	435

Example	R ²	¹ H NMR δ	MS m/e MH ⁺
183		11.17 (1H, s), 11.17 (1H, s), 7.68 (1H, ddd), 7.57 (2H, dd), 7.52-7.39 (1H, m), 7.36-7.28 (1H, m), 6.72 (2H, dd), 3.68-3.44 (6H, m), 3.35-3.23 (2H, m), 1.98-1.72 (5H, m)	457
184		10.91 (1H, s), 10.68 (1H, s), 7.57 (2H, dd), 7.48-7.23 (7H, m), 7.16-7.10 (1H, m), 6.76-6.68 (3H, m), 5.09 (2H, s), 3.68-3.45 (6H, m), 3.35-3.24 (2H, m), 1.98-1.73 (5H, m)	527
185		10.71 (1H, s), 10.69 (1H, s), 8.30-8.24 (1H, m), 8.03-7.95 (2H, m), 7.75 (1H, d), 7.65-7.50 (5H, m), 6.73 (2H, d), 3.70-3.47 (6H, m), 3.37-3.25 (2H, m), 1.98-1.73 (5H, m)	471
186		10.99 (1H, s), 10.71 (1H, s), 8.21-8.09 (1H, m), 7.75-7.63 (1H, m), 7.65 (2H, dd), 6.72 (2H, dd), 3.68-3.46 (6H, m), 3.35-3.25 (2H, m), 1.98-1.74 (5H, m)	475
187		11.02 (1H, s), 10.73 (1H, s), 8.02-7.93 (1H, m), 7.57 (2H, dd), 7.35 (1H, ddd), 6.98-6.90 (1H, m), 6.74 (2H, dd), 3.69-3.44 (6H, m), 3.34-3.25 (2H, m), 1.98-1.75 (5H, m)	457




Example 188











Example 189








Example	SM	¹ H NMR δ	MS m/e MH ⁺
188	Int. 61	11.01 (1H, s), 10.96 (1H, s), 8.70 (2H, s), 7.48-7.23 (7H, m), 7.17-7.11 (1H, m), 6.72 (1H, dd), 5.09 (2H, s), 3.79-3.64 (4H, m), 3.55-3.47 (4H, m), 2.03 (3H, s)	515
189	Int. 62	11.15 (1H, s), 10.72 (1H, s), 8.38 (1H, s), 8.22 (1H, d), 7.99-7.90 (3H, m), 7.78 (1H, d), 7.70 (1H, d), 7.55-7.45 (3H, m), 2.72 (3H, s).	402








Starting material = Intermediate 32

Example	R ²	¹ H NMR δ	MS m/e MH ⁺
190		10.97 (1H, s), 10.92 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.65-7.55 (2H, m), 7.20 (2H, t), 6.95 (1H, d), 3.60 (6H, m), 3.30-3.20 (2H, m), 2.00 (3H, s)	426
191		11.20 (1H, s), 10.80 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.70 (1H, s), 7.50 (1H, d), 7.40 (1H, t), 7.10 (1H, d), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	442
192		11.07 (1H, s), 10.97 (1H, s), 8.5 (1H, s), 8.01-7.95 (2H, m), 7.40-7.30 (1H, m), 7.00-6.95 (1H, m), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	444
193		10.95-10.91 (2H, d), 8.5 (1H, s), 7.95 (1H, d), 7.60 (2H, d), 7.40 (1H, t), 7.05 (1H, t), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	408



Example	R ²	¹ H NMR δ	MS m/e MH ⁺
194		11.17 (1H, s), 10.97 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.90 (1H, s), 7.65 (1H, d), 7.35 (1H, t), 7.20 (1H, d), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	488
195		11.21 (1H, s), 10.97 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.55-7.30 (3H, m), 6.95 (2H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	426
196		10.90 (1H, s), 10.81 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.45 (2H, d), 7.20 (2H, d), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.20 (3H, s), 2.00 (3H, s)	422
197		10.92 (1H, d), 10.38 (1H, s), 8.5 (1H, s), 7.95-7.90 (2H, m), 7.55 (1H, d), 7.40 (1H, t), 7.20 (1H, t), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	442
198		11.10 (1H, s), 10.93 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.60 (2H, d), 7.40 (2H, d), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	442
199		11.18 (1H, s), 10.95 (1H, s), 8.5 (1H, s), 8.20 (1H, s), 8.0-7.90 (2H, m), 7.80 (2H, d), 7.65 (1H, d), 7.50 (1H, t), 7.40 (1H, t), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	458
200			409
201		11.40 (1H, s), 10.97 (1H, s), 8.5 (1H, s), 8.0-7.95 (3H, t), 7.70 (2H, d), 6.95 (1H, d), 4.30 (2H, q), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s), 1.30 (3H, t)	480









Example	R ²	¹ H NMR δ	MS m/e MH ⁺
202		10.90 (1H, s), 10.79 (1H, s), 8.5 (1H, s), 7.95 (1H, t), 7.20 (1H, s), 7.05- 6.95 (3H, m), 6.0 (2H, s), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	452
203		10.92 (1H, s), 10.71 (1H, s), 8.5 (1H, s), 8.01-7.90 (2H, m), 7.40 (1H, t), 7.20 (1H, t), 6.95 (1H, d), 3.60 (6H, m), 3.30-3.20 (2H, m), 2.00 (3H, s)	444
204		11.17 (1H, s), 10.94 (1H, s), 8.50 (1H, s), 8.25 (1H, s), 7.95 (1H, d), 7.85 (1H, d), 7.65 (1H, d), 7.50 (1H, t), 6.95 (1H, d), 4.30 (2H, q), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s), 1.30 (3H, t)	480
205		10.90 (1H, s), 10.81 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.45 (2H, d), 7.40 (2H, d), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s), 1.15 (9H, s)	464
206		11.20 (1H, s), 10.95 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.70-7.60 (1H, m), 7.45 (1H, q), 7.40-7.30 (1H, m), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	444
207		11.05 (1H, s), 10.95 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.90 (1H, d), 7.30 (1H, q), 7.20 (1H, q), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	444
208		10.90 (1H, s), 10.85 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.45 (1H, d), 7.39 (1H, s), 7.25 (1H, t), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.30 (3H, s), 2.00 (3H, s)	422









Example	R ²	¹ H NMR δ	MS m/e MH ⁺
209		10.95 (1H, s), 10.93 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.45-7.25 (7H, m), 7.15 (1H, d), 6.95 (1H, d), 6.85 (1H, d), 5.10 (2H, s), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	514
210		11.45 (1H, s), 10.98 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.30 (2H, d), 6.95 (2H, t), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	444
211		10.93 (1H, s), 10.75 (1H, s), 8.5 (1H, s), 8.30-8.20 (1H, m), 8.0-7.90 (3H, m), 7.80 (1H, d), 7.60-7.50 (3H, m), 6.95 (1H, d), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	458
212		11.04 (1H, s), 10.96 (1H, s), 8.5 (1H, s), 8.20-8.10 (1H, m), 7.95 (1H, d), 7.7 (1H, q), 6.95 (2H, t), 3.60 (6H, m), 3.50-3.40 (2H, m), 2.00 (3H, s)	462







Starting material = Intermediate 21

Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
213		10.95 (1H, s), 10.80 (1H, s), 8.7-8.5 (4H, m), 7.20 (2H, t), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	425
214		11.18 (1H, s), 10.84 (1H, s), 7.70 (1H, s), 7.60 (2H, d), 7.50 (1H, d), 7.40 (1H, t), 7.10 (1H, d), 6.95 (1H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	441




Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
215		10.85 (1H, s), 8.05-7.95 (1H, m), 7.65 (2H, d), 7.40-7.30 (1H, m), 6.95 (3H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	443
216		10.91 (2H, d), 10.80 (1H, s), 8.65- 8.55 (4H, m), 7.40 (2H, t), 7.05 (1H, t), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	407
217		11.17 (1H, s), 10.83 (1H, s), 7.85 (1H, s), 7.65 (2H, d), 7.55 (1H, d), 7.35 (1H, t), 7.20 (1H, d), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	487
218		11.21 (1H, s), 10.83 (1H, s), 7.65 (2H, d), 7.55-7.30 (3H, m), 6.95 (2H, d), 6.90-6.80 (1H, m), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	425
219		10.79 (1H, s), 10.77 (1H, s), 7.65 (2H, d), 7.45 (2H, d), 7.20 (2H, d), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.20 (3H, s), 2.00 (3H, s)	421
220		10.80 (1H, d), 10.34 (1H, s), 7.95 (1H, d), 7.65 (2H, d), 7.55 (1H, d), 7.40 (1H, t), 7.20 (1H, t), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	441
221		11.10 (1H, s), 10.84 (1H, s), 7.65-7.55 (4H, m), 7.40 (2H, d), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	441
222		11.15 (1H, s), 10.82 (1H, s), 8.20 (1H, s), 7.95 (1H, d), 7.85 (2H, d), 7.70-7.60 (3H, m), 7.50 (1H, t), 7.40 (1H, t), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	457








Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
223		10.77 (1H, s), 7.65 (2H, d), 7.20 (1H, s), 7.05-6.95 (4H, m), 6.0 (2H, s), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	451
224		10.80 (1H, s), 10.65 (1H, s), 8.01-7.95 (1H, m), 7.60 (2H, d), 7.40 (1H, t), 7.20 (1H, t), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	443
225		11.17 (1H, s), 10.94 (1H, s), 8.30 (1H, s), 7.90 (1H, d), 7.80 (3H, t), 7.50 (1H, t), 6.95 (2H, d), 4.30 (2H, q), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s), 1.30 (3H, t)	479
226		10.79 (1H, s), 10.77 (1H, s), 7.65 (2H, d), 7.50 (2H, d), 7.40 (2H, d), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s), 1.20 (9H, s)	463
227		11.20 (1H, s), 10.83 (1H, s), 7.70-7.60 (3H, m), 7.45 (1H, q), 7.40-7.30 (1H, m), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	443
228		10.83 (1H, s), 7.95 (1H, t), 7.60 (1H, d), 7.20 (2H, q), 7.10 (1H, q), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	443
229		10.93 (1H, s), 10.80 (1H, s), 7.60 (2H, d), 7.50-7.25 (7H, m), 7.15 (1H, d), 6.95 (2H, d), 6.85 (1H, d), 5.10 (2H, t), 3.60 (4H, s), 3.10 (4H, m), 2.00 (3H, s)	513
230		11.45 (1H, s), 10.85 (1H, s), 7.65 (2H, d), 7.30 (2H, d), 7.0-6.90 (3H, m), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	443





Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
231		11.00 (1H, s), 10.84 (1H, s), 8.20-8.10 (1H, m), 7.75-7.60 (3H, m), 6.95 (2H, d), 3.60 (4H, s), 3.10 (4H, m), 2.00 (3H, s)	461
232		10.90 (1H, s), 10.78 (1H, s), 7.65 (2H, d), 7.30-7.20 (2H, m), 7.10 (1H, t), 6.95 (2H, d), 6.60 (1H, d), 3.75 (3H, s), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	437
233		10.76 (1H, s), 10.68 (1H, s), 7.65 (2H, d), 7.45 (2H, d), 7.00-6.95 (4H, m), 3.70 (3H, s), 3.60 (4H, s), 3.10 (4H, d), 2.00 (3H, s)	437




Starting material = Intermediate 63



Example	R ²	¹ H NMR δ	MS m/e MH ⁺
234		10.97 (1H, s), 10.93 (1H, s), 8.50 (1H, s), 8.00 (1H, d), 7.60 (2H, d), 7.40 (2H, t), 7.05 (1H, t), 6.95 (1H, d), 3.60-3.50 (4H, m), 3.10-3.05 (4H, m), 2.80 (3H, s)	444
235		11.15 (1H, s), 10.95 (1H, s), 8.50 (1H, s), 8.00 (1H, d), 7.60 (2H, d), 7.40 (2H, t), 6.95 (1H, d), 3.60-3.55 (4H, m), 3.10-3.05 (4H, m), 2.85 (3H, s)	478
236		11.05 (1H, s), 10.95 (1H, s), 8.50 (1H, s), 8.00 (1H, d), 7.90 (1H, t), 7.30-7.10 (2H, m), 6.95 (1H, d), 3.60-3.55 (4H, m), 3.10-3.05 (4H, m), 2.85 (3H, s)	480









Example	R ²	¹ H NMR δ	MS m/e MH ⁺
237		10.94 (1H, s), 8.50 (1H, s), 7.95 (1H, d), 7.60 (1H, d), 7.50-7.30 (7H, m), 7.10 (1H, d), 6.95 (1H, d), 6.7 (1H, d), 3.60-3.55 (4H, m), 3.10-3.05 (4H, m), 2.85 (3H, s)	550
238		10.95 (1H, s), 10.70 (1H, s), 8.50 (1H, s), 8.00-7.90 (2H, m), 7.40-7.30 (1H, m), 7.20-7.10 (1H, m), 6.95 (1H, d), 3.60-3.55 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s)	480
239		11.07 (1H, s), 11.00 (1H, s), 8.50 (1H, s), 8.00-7.90 (2H, m), 7.40-7.30 (1H, m), 7.00-6.95 (2H, m), 3.60-3.55 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s)	480
240		11.45 (1H, s), 11.00 (1H, s), 8.50 (1H, s), 8.00 (1H, d), 7.30 (2H, d), 7.0-6.95 (2H, m), 3.60-3.55 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s)	480
241		10.97 (1H, s), 10.75 (1H, s), 8.50 (1H, s), 8.30-8.20 (1H, m), 8.00-7.90 (3H, m), 7.80 (2H, d), 7.60-7.50 (3H, m), 6.95 (1H, d), 3.60-3.50 (4H, m), 3.10-3.05 (4H, m), 2.85 (3H, s)	494
242		11.18 (1H, s), 10.98 (1H, s), 8.50 (1H, s), 8.20 (1H, s), 8.00 (1H, d), 7.95 (1H, d), 7.80 (2H, d), 7.60 (1H, d), 7.5 (1H, t), 7.40 (1H, t), 6.95 (1H, d), 3.60-3.50 (4H, m), 3.10-3.05 (4H, m), 2.85 (3H, s)	494
243		11.04 (1H, s), 10.98 (1H, s), 8.50 (1H, s), 8.10-8.00 (1H, m), 7.95 (1H, d), 7.70 (1H, q), 6.95 (1H, d), 3.60-3.55 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s)	498









Example	R ²	¹ H NMR δ	MS m/e MH ⁺
244		10.94 (1H, s), 10.81 (1H, s), 8.50 (1H, s), 8.00 (1H, d), 7.50 (2H, d), 7.40 (2H, t), 6.95 (1H, d), 3.60-3.50 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s), 1.20 (9H, m)	500
245		11.17 (1H, s), 10.97 (1H, s), 8.50 (1H, s), 8.25 (1H, s), 8.00 (1H, d), 7.85 (1H, d), 7.60 (1H, d), 7.50 (1H, t), 6.95 (1H, d), 4.30 (2H, q), 3.60-3.50 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s), 1.30 (3H, t)	516
246		No nmr data	445
247		10.91 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.55-7.45 (1H, m), 7.25 (1H, t), 6.95 (1H, d), 3.60-3.55 (4H, m), 3.20-3.10 (4H, m), 2.85 (3H, s)	480



Starting material = Intermediate 64








Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
248		10.93 (1H, s), 10.88 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.60 (2H, d), 7.40 (2H, t), 7.05 (1H, t), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	383
249		10.85 (1H, s), 9.95 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.70 (1H, d), 7.25-7.2 (2H, m), 7.05 (1H, t), 6.90 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m), 2.3 (3H, s)	397



Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
250		10.86 (1H, s), 10.80 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.50 (2H, d), 7.20 (2H, d), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m), 2.2 (3H, s)	397
251		11.20 (1H, s), 10.91 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.60 (1H, d), 7.50-7.30 (2H, m), 6.95-6.90 (2H, m), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	401
252		10.96 (1H, s), 10.88 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.65-7.55 (2H, m), 7.20 (2H, t), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	401
253		10.85 (1H, s), 10.02 (1H, s), 8.50 (1H, s), 7.95-7.90 (2H, m), 7.10-7.00 (2H, m), 7.05-6.95 (1H, m), 6.90 (1H, d), 3.90-3.80 (7H, m), 2.60-2.50 (4H, m)	413
254		10.92 (1H, s), 10.88 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.30-7.20 (2H, m), 7.10 (1H, d), 6.90 (1H, d), 6.80 (1H, d), 3.90-3.80 (4H, m), 3.70 (3H, s), 2.60-2.50 (4H, m)	413
255		10.85 (1H, s), 10.68 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.50 (2H, d), 6.95 (2H, d), 6.95 (1H, d), 3.90-3.80 (4H, m), 3.7 (3H, s), 2.60-2.50 (4H, m)	413
256		10.88 (1H, s), 10.36 (1H, s), 8.50 (1H, s), 8.00-7.85 (2H, m), 7.60 (1H, d), 7.40 (1H, t), 7.20 (1H, t), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	417
257		11.18 (1H, s), 10.91 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.70 (1H, s), 7.50 (1H, d), 7.40 (1H, t), 7.10 (1H, d), 6.90 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	417


Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
258		11.10 (1H, s), 10.90 (1H, s), 8.50 (1H, s), 7.95 (1H, d), 7.60 (2H, d), 7.40 (2H, d), 6.95 (1H, d), 3.90-3.80 (4H, m), 3.60-3.40 (4H, m)	417
259		10.91 (1H, s), 8.50 (1H, s), 7.90-7.80 (2H, m), 7.30-7.10 (2H, m), 6.95 (1H, d), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	419
260		10.88 (1H, s), 10.68 (1H, s), 8.50 (1H, s), 8.05-7.95 (1H, m), 7.90 (1H, d), 7.40-7.30 (1H, m), 7.20-7.10 (1H, m), 6.95 (1H, d), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	419
261		10.94 (1H, s), 8.50 (1H, s), 8.00-7.90 (2H, m), 7.40-7.30 (1H, m), 7.0-6.95 (1H, m), 6.90 (1H, d), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	419
262		11.45 (1H, s), 10.95 (1H, s), 8.50 (1H, s), 7.95-7.90 (1H, m), 7.30 (2H, d), 6.95-6.90 (2H, m), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	419
263		10.86 (1H, s), 10.78 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.30 (1H, s), 7.05-6.80 (3H, m), 6.00 (2H, s), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	427
264		10.90 (1H, s), 10.75 (1H, s), 8.50 (1H, s), 8.30-8.20 (1H, m), 8.00-7.90 (3H, m), 7.80 (1H, d), 7.60-7.50 (3H, m), 6.90 (1H, d), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	433
265		11.16 (1H, s), 10.91 (1H, s), 8.50 (1H, s), 8.20 (1H, s), 7.95-7.80 (2H, m), 7.80 (1H, d), 7.50 (1H, t), 7.40 (1H, t), 6.95 (1H, d), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	433

206







Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
266		10.92 (1H, s), 8.50 (1H, s), 8.20-8.10 (1H, m), 7.90 (1H, d), 7.70 (1H, q), 6.90 (1H, d), 3.90- 3.80 (4H, m), 3.60-3.50 (4H, m)	437
267		10.86 (1H, s), 10.80 (1H, s), 8.50 (1H, s), 7.95 (1H, d), 7.50 (2H, d), 7.40 (2H, d), 6.95 (1H, d), 3.90-3.80 (4H, m), 3.60-3.50 (4H, m)	439
268		11.15 (1H, s), 10.91 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.80 (1H, s), 7.55 (1H, d), 7.30 (1H, t), 7.25 (1H, d), 6.90 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	463
269		10.90 (1H, s), 8.50 (1H, s), 8.30 (1H, d), 7.95- 7.90 (3H, m), 7.05-7.00 (1H, m), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	384
270			489
271		11.16 (1H, s), 10.90 (1H, s), 8.50 (1H, s), 8.15 (1H, s), 7.9-7.80 (2H, m), 7.60 (1H, d), 7.50 (1H, t), 6.95 (1H, d), 4.30 (2H, q), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m), 1.3 (3H, t)	455
272		10.99 (1H, s), 10.75 (1H, s), 8.50 (1H, s), 8.05- 7.90 (2H, m), 7.30-7.20 (2H, m), 7.20-7.10 (1H, m), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	401

Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
273		11.20 (1H, s), 10.92 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.7-7.60 (1H, m), 7.50 (1H, q), 7.40-7.30 (1H, m), 6.95 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	419
274		10.94 (1H, s), 10.86 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.44 (2H, d), 7.35 (1H, s), 7.25 (1H, t), 6.90 (1H, dd), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m), 2.3 (3H, s)	397
















Starting material = Intermediate 31


Example	R ²	¹ H NMR δ	MS m/e MH ⁺
275			398
276			415
277			410
278			416

208


142



Example	R ²	¹ H NMR δ	MS m/e MH ⁺
279		10.93 (1H, s), 10.88 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.60 (2H, d), 7.40 (2H, t), 7.05 (1H, t), 6.95 (1H, d), 3.50-3.40 (4H, m), 2.40-2.30 (4H, m), 2.20 (3H, s)	380
280			459
281			398
282		10.90 (1H, s), 10.81 (1H, s), 8.5 (1H, s), 7.95 (1H, d), 7.50 (2H, d), 7.20 (2H, d), 6.90 (1H, d), 3.5-3.40 (4H, s), 2.50-2.40 (4H, m), 2.25 (3H, s), 2.20 (3H, s)	394
283			416
284			415
285		11.10 (1H, s), 10.90 (1H, s), 8.50 (1H, s), 7.95 (1H, d), 7.60 (2H, d), 7.40 (2H, d), 6.95 (1H, d), 3.50-3.40 (4H, m), 2.50-2.40 (4H, m)	415
286		11.18 (1H, s), 10.92 (1H, s), 8.50 (1H, s), 8.20 (1H, s), 7.95-7.90 (1H, m), 7.90 (1H, d), 7.65 (1H, d), 7.50 (1H, t), 7.4 (1H, t), 6.90 (1H, d), 3.50-3.40 (4H, m), 2.40-2.30 (4H, m), 2.20 (3H, s)	430


Example	R ²	¹ H NMR δ	MS m/e MH ⁺
287			381
288		10.92 (1H, s), 10.88 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.30-7.20 (2H, m), 7.10 (1H, d), 6.85 (1H, d), 6.60 (1H, d), 3.70 (3H, s), 3.50-3.40 (4H, m), 2.50-2.40 (4H, m), 2.2 (3H, s)	410
289		10.88 (1H, s), 10.78 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 7.25 (1H, s), 7.00 (1H, d), 6.95 (1H, d), 6.90 (1H, d), 6.00 (2H, s), 3.50-3.40 (4H, m), 3.40-3.30 (4H, m), 2.2 (3H, s)	424
290			416
291			416





Starting material = Intermediate 65

Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
292		11.08 (1H, s), 10.76 (1H, s), 8.10 (1H, s), 8.00 (1H, t), 7.75 (1H, d), 7.35-7.10 (4H, m), 3.80-3.70 (4H, m), 3.00-2.90 (4H, m)	464

293		11.25 (1H, s), 11.11 (1H, s), 8.10 (1H, s), 7.75 (1H, d), 7.70-7.60 (1H, m), 7.50 (1H, q), 7.40-7.30 (1H, m), 7.20 (1H, s), 3.80-3.70 (4H, m), 3.00-2.90 (4H, m)	482
294		11.06 (1H, s), 10.88 (1H, s), 8.10 (1H, s), 7.80 (1H, d), 7.40 (1H, d), 7.35 (1H, s), 7.25-7.20 (2H, m), 3.80-3.70 (4H, m), 3.00-2.90 (4H, m), 2.30 (3H, s)	460




Starting material = Intermediate 66

Example	R ²	¹ H NMR δ	MS m/e MH ⁺
295		11.08 (1H, s), 10.76 (1H, s), 8.10 (1H, s), 8.00 (1H, d), 7.80 (1H, d), 7.35-7.10 (4H, m), 6.10-5.95 (1H, m), 5.45 (1H, d), 5.30 (1H, d), 4.80 (2H, d), 3.80-3.70 (4H, m), 3.00-2.90 (4H, m)	468
296		11.24 (1H, s), 11.11 (1H, s), 8.10 (1H, s), 7.90 (1H, d), 7.70-7.60 (1H, m), 7.50 (1H, q), 7.40-7.30 (1H, m), 7.20 (1H, s), 6.10-5.95 (1H, m), 5.45 (1H, d), 5.30 (1H, d), 4.80 (2H, d), 3.80-3.70 (4H, m), 3.00-2.90 (4H, m)	486




5 Examples 297-304:

The following examples were prepared by the general procedure of Example 87, using Intermediate 41 and commercially available acid chlorides.


- 188 -



Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
297		10.93 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.35-7.10 (3H, m), 6.90 (1H, d), 4.10 (2H, s), 3.30 (8H, s)	456
298		10.98 (1H, s), 10.93 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.35-7.10 (8H, m), 6.90 (1H, d), 3.70-3.65 (1H, m), 3.60-3.50 (3H, m), 3.45-3.40 (3H, m), 3.20-3.10 (1H, m), 2.80 (2H, q), 2.70 (2H, q)	516
299		10.94 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.55-7.40 (2H, m), 7.35-7.20 (5H, m), 7.20-7.10 (1H, m), 6.90 (1H, d), 3.80-3.75 (2H, m), 3.60-3.50 (2H, m), 3.50-3.40 (2H, m), 3.30-3.20 (2H, m)	506
300		10.96 (1H, s), 10.74 (1H, s), 8.50 (1H, s), 8.05-7.90 (5H, m), 7.60 (2H, d) 7.30-7.10 (3H, m), 6.90 (1H, d), 3.80-3.75 (2H, m), 3.60-3.50 (2H, m), 3.50-3.40 (2H, m), 3.30- 3.20 (2H, m)	513
301		10.94 (1H, s), 10.74 (1H, s), 8.50 (1H, s), 8.05-7.90 (2H, m), 7.40-7.10 (7H, m), 6.90 (1H, d), 3.60-3.40 (8H, m), 2.30 (3H, s)	502

Example	R ¹	¹ H NMR δ	MS m/e MH ⁺
302		10.94 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.35-7.10 (8H, m), 6.90 (1H, d), 3.80-3.75 (2H, m), 3.60-3.50 (2H, m), 3.50-3.40 (2H, m), 3.30-3.20 (2H, m), 2.10 (3H, s)	502
303		10.94 (1H, s), 10.74 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.40-7.10 (8H, m), 6.90 (1H, d), 3.60-3.40 (8H, m), 2.30 (3H, s)	502
304		10.94 (1H, s), 10.74 (1H, s), 8.65 (2H, s), 8.50 (1H, s), 8.05-7.85 (3H, m), 7.55-7.45 (1H, m), 7.30-7.10 (3H, m), 6.90 (1H, d), 3.60-3.40 (8H, m)	489

Example 305: 5-[(2-Fluorophenyl)amino]-N-[6-(4-glycoloylpiperazin-1-yl)pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide




5 EDAC (60 mg, 0.31 mmol) was added to a stirred mixture of 5-[(2-fluorophenyl)amino]-N-(6-piperazin-1-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (Intermediate 41, 100 mg, 0.26 mmol), glycolic acid (20 mg (0.26 mmol) and HOBT (42 mg, 0.31 mmol) in 3 mL of DMF and stirring was continued for 16 h. Water (10 mL) was added and the resultant

10 precipitate was collected by filtration, washed with water then ether and dried *in vacuo* to give the title compound (75mg, 65%) as a solid; ¹H NMR δ 10.94 (1H, s), 10.70 (1H, s), 8.50 (1H, s), 8.05 (1H, t), 7.90 (1H, d), 7.35-7.20 (2H, m), 7.20-7.10 (1H, m), 6.90 (1H, d), 4.60 (1H, t), 4.10 (2H, d), 3.60-3.4 (8H, m); MS m/e MH⁺ 442.


~~11~~

Example 306: 4-{5-[(5-{(2-Fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperazine-1-carboxamide



Sodium cyanate 34mg (0.52 mmol) was added to a mixture of 5-[(2-fluorophenyl)amino]-N-(6-piperazin-1-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (Intermediate 41, 100mg, 0.26mmol) and glacial acetic acid (0.2 mL) in 0.6 mL of water (0.6 mL) and THF (0.3 mL). The mixture was stirred for 16 h then sodium cyanate (34 mg), glacial acetic acid (0.2 mL), dioxan (0.5 mL) and water (0.2 mL) were added. The mixture was stirred for another 2 h then directly purified by reverse phase HPLC, eluting with a gradient of acetonitrile / water containing 0.2% TFA, to give the title compound (90 mg, 82%) as a solid; ^1H NMR δ 11.02 (1H, s), 10.74 (1H, s), 8.50 (1H, s), 8.05-7.95 (2H, m), 7.35-7.20 (2H, m), 7.20-7.10 (1H, m), 6.90 (1H, d), 3.60-3.4 (8H, m); MS m/e MH^+ 427.


Example 307: 4-{5-[(5-{(2-Fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}-N-phenylpiperazine-1-carboxamide



Phenyl isocyanate (0.032 mL, 0.29 mmol) was added to a stirred mixture of 5-[(2-fluorophenyl)amino]-N-(6-piperazin-1-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (Intermediate 41, 100 mg, 0.26 mmol) and DMF (1.5 mL). After 2 h the crude reaction mixture was directly purified by reverse phase HPLC eluting with a gradient of acetonitrile / water containing 0.2% TFA to give the title compound (28 mg, 20%) as a solid; ^1H NMR δ 11.00 (1H, s), 10.74 (1H, s), 8.55 (1H, s), 8.50 (1H, s), 8.05-7.95 (2H, m), 7.45 (2H, d), 7.35-7.10 (4H, m), 7.00 (1H, d), 6.90 (1H, t), 3.80-3.60 (8H, m); MS m/e MH^+ 503.


Example 308

The following example was prepared by the general procedure of Example 307 using Intermediate 41 and the appropriate isocyanate.



^1H NMR δ 10.98 (1H, s), 10.74 (1H, s), 8.5 (1H, s), 8.05-7.95 (2H, m), 7.30-7.10 (8H, m), 6.95 (1H, d), 4.20 (2H, d), 3.50 (8H, s); MS m/e MH^+ 517.


5 **Example 309: Phenyl 4-((5-((2-fluorophenyl)amino)-1,3,4-oxadiazol-2-yl)carbamoyl)amino]pyridin-2-yl}piperazine-1-carboxylate**




Phenyl chloroformate (0.036 mL, 0.29 mmol) was added to a mixture of 5-[(2-fluorophenyl)amino]-*N*-(6-piperazin-1-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (Intermediate 41, 100 mg, 0.26 mmol) and triethylamine (0.07 mL, 0.52 mmol) in DMF (1.5 mL). The mixture was stirred for 2 h then directly purified by reverse phase HPLC eluting with a gradient of acetonitrile / water containing 0.2% TFA to give the title compound (40 mg, 31%) as a solid; ^1H NMR δ 10.95 (1H, s), 10.73 (1H, s), 8.50 (1H, s), 8.05-7.95 (2H, m), 7.45-7.10 (8H, m), 6.90 (1H, d), 3.80-3.60 (8H, m); MS m/e MH^+ 504.

15 **Examples 310-312**

The following Examples were prepared by the general procedure of Example 309 using Intermediate 73 in place of Intermediate 41 and using the appropriate commercially available chloroformate.




Example	R ¹	^1H NMR δ	MS m/e MH^+
310	Ethyl	10.95 (1H, s), 10.73 (1H, s), 8.5 (1H, s), 8.05-7.95 (2H, m), 7.30-7.10 (3H, m), 6.95 (1H, d),	456




		4.10 (3H, q), 3.50 (8H, s), 1.20 (2H, t)	
311		10.93 (1H, s), 10.72 (1H, s), 8.5 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.40-7.20 (7H, m), 7.20-7.10 (1H, m), 6.95 (1H, d), 5.10 (2H, s), 3.50 (8H, s)	518
312	Ethyl	10.81 (1H, s), 10.70 (1H, s), 8.0 (1H, t), 7.60 (2H, d), 7.30-7.10 (3H, m), 6.95 (2H, d), 4.10 (3H, q), 3.50-3.40 (4H, m), 3.10-3.00 (4H, m), 1.20 (2H, t)	455




Examples 313-318:

The following examples were prepared by the general procedure of Example 72 (method B) using Intermediate 40 and the appropriate aniline.



5

Example	R ⁴	¹ H NMR δ	MS m/e MH ⁺
313 ¹		11.02 (1H, s), 10.74 (1H, s), 8.0 (3H, t), 7.75 (2H, d), 7.35-7.20 (4H, m), 7.20-7.10 (1H, m), 2.90 (2H, t), 1.50 (2H, m), 1.40 (2H, m), 0.85 (3H, t)	387
314 ²			492
315 ³		11.10 (1H, s), 10.75 (1H, s), 8.10 (1H, t), 7.80-7.70 (2H, m), 7.50-7.40 (1H, m), 7.35-7.20 (3H, m), 7.20-7.10 (2H, m), 6.90 (1H, d)	427

Example	R ⁴	¹ H NMR δ	MS m/e MH ⁺
316 ⁴		11.20 (1H, s), 10.75 (1H, s), 8.10 (1H, t), 7.90 (2H, d), 7.85 (2H, d), 7.35-7.20 (2H, m), 7.20-7.10 (1H, m), 3.45 (1H, qn), 3.30 (3H, s), 2.75 (1H, dd), 2.35 (1H, d), 1.10 (3H, d)	423
317 ⁵		11.65 (1H, s), 11.20 (1H, s), 10.75 (1H, s), 8.10 (1H, t), 7.90 (2H, d), 7.85 (2H, d), 7.35-7.20 (2H, m), 7.20-7.10 (1H, m), 4.75 (1H, q), 1.10 (3H, d)	427
318 ⁶		11.12 (1H, s), 10.75 (1H, s), 8.05 (1H, t), 7.90 (1H, d), 7.70-7.60 (4H, m), 7.50 (2H, d), 7.10-7.30 (7H, m)	MH ⁺ 407

¹Starting material (SM) prepared as described in *Farmaco* 2004, 59(6), 443.

²SM prepared as described in *J. Chem. Soc. C (Organic)*, 1967, 24, 2700.


³SM prepared as described in WO 2001056990

⁴SM prepared as described in US 3746712

5 ⁵SM prepared as described in EP 52442.

⁶SM prepared as described in WO 2001055110

Example 319: Methyl [1-(5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]pyridin-2-yl)piperidin-4-yl]acetate



10


Methyl [1-(5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]pyridin-2-yl)piperidin-4-yl]acetate

(Intermediate 77, 215 mg, 0.64 mmol) was dissolved in DMA (4 mL) and treated with 2,4,5-trifluorophenylisothiocyanate (146mg, 0.77 mmol). The reaction mixture was heated at 45°C for 30 minutes then EDAC (148 mg, 0.77 mmol) was added and the mixture was heated to 85°C for 1 h. The mixture was cooled and diluted with water (4 mL). The resulting precipitate was filtered off, washed with water and dried under high vacuum to give the title compound (277mg, 89%) as a solid; ¹H NMR δ 1.04-1.20 (2H, m), 1.64 (2H, d), 1.78-1.95 (1H, m), 2.19 (2H, d), 2.63-2.78 (2H, m), 3.52 (3H, s), 4.16 (2H, d), 6.77

15

(1H, d), 7.57-7.69 (1H, m), 7.76-7.86 (1H, m), 8.01-8.17 (1H, m), 8.34-8.43 (1H, m), 10.85 (1H, s), 10.97 (1H, s); MS m/e MH⁺ 491.


Example 320: (1-{5-[(5-{(2,4,5-Trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetic acid hydrochloride



Methyl (1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)-amino]pyridin-2-yl}piperidin-4-yl)acetate (Example 319, 216 mg, 0.44 mmol) was suspended in a mixture of MeOH (4ml) and THF (8 mL). Lithium hydroxide (4.4 mL of a 1M aqueous solution) was added. The resulting clear solution was heated at 45°C for 15 minutes then cooled and acidified to pH ~ 2 with dropwise addition of concentrated aqueous HCl. Volatile material was removed by evaporation and the residue was washed with water then dried to give the title compound (182 mg, 87%) as a solid; ¹H NMR δ 1.16-1.45 (2H, m), 1.76-1.93 (2H, m), 1.98-2.13 (1H, m), 2.28 (2H, d), 3.12 (2H, t), 4.28 (2H, d), 7.25-7.41 (1H, m), 7.70-7.88 (1H, m), 8.13-8.32 (2H, m), 8.55 (1H, d), 11.16 (1H, s), 11.29 (1H, s); MS m/e MH⁺ 477.




Examples 321-337

The following Examples were prepared by the general procedure of Example 1, using Intermediate 43 and commercially available isothiocyanates.




Example	R	HPLC Retention Time (minutes)	MS m/e MH ⁺
321		2.98	463

322		2.68	465
323		2.99	503
324		2.71	460
325		2.86	463
326		2.94	477
327		3.10	541
328		2.84	449
329		3.10	477
330		2.80	503
331		3.00	503
332		2.69	460
333		3.04	519
334		2.96	519

335		2.97	463
336		2.83	465
337 ¹		2.52	460

¹ The commercially available isocyanate was used in place of isothiocyanate


Example 338: Methyl [*trans*-4-(4-[(5-[(3-(aminocarbonyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)phenyl)cyclohexyl]acetate

















3-Aminobenzamide (54 mg, 0.39 mmol) was added to a solution of thiocarbonyldiimidazole (71 mg, 0.39 mmol) in CH₃CN (5 mL) and the mixture was stirred for 16 h. Methyl [*trans*-4-(4-[(hydrazino(oxo)acetyl]amino)phenyl)cyclohexyl]acetate (Intermediate 43) (110 mg, 0.33 mmol) and DMF (5 mL) was added and the mixture was stirred at 50°C until a clear solution was obtained. PS-CDI (527 mg) was added and the temperature raised to 80°C for 16 h. The mixture was filtered whilst hot and the polymer washed with DMF (2 x 5 mL). The filtrates were concentrated by evaporation to give the title compound as a solid; MS m/e (M+Na)⁺ 500; HPLC retention time = 2.26 minutes.

Examples 339-352

The following examples were prepared by the general procedure of Example 338 using the appropriate aniline (commercially available unless indicated otherwise) in place of 3-aminobenzamide.



Example	R	HPLC Retention Time (min)	MS m/e MH ⁺
339		2.98	561
340		3.03	519
341		3.13	527
342		3.12	541
343 ¹		2.82	552 (M-H) ⁻
344 ²		2.65	529
345		2.86	575
346 ³		2.69	527 (M-H) ⁻
347 ⁴		2.71	529
348		2.32	515 (M+Na) ⁺
349 ⁵		2.64	508

350		2.90	520
351		3.08	526
352		3.14	526

The following footnotes refer to preparation of the aniline starting materials:

¹For preparative details: CAS Reg. No. 16091-26-2.

²Prepared as described in J. Am. Chem. Soc., 2003, 125(22), 6630.

³Prepared as described in US Patent Application No. US 3687998 (1972).


5 ⁴Prepared as described in Tetrahedron, 1988, 54(1/2), 45.




⁵For preparative details: CAS Reg. No. 6464-98-8

Examples 353-355

The following examples were prepared by the general procedure of Example 338 but using *O,O*-dipyridin-2-yl thiocarbonate in place of thiocarbonyldiimidazole and using the appropriate commercially available aniline.


10



Example	R	HPLC Retention Time (min)	MS m/e MH ⁺
353		2.87	575
354 ¹		3.11	575
355		2.64	527

¹Heated at 80°C for 24 hours when *O,O*-dipyridin-2-yl thiocarbonate and aniline mixed. DMF was also used in place of CH₃CN.


Example 356: (*trans*-4-{4-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid









A solution of lithium hydroxide monohydrate (110 mg, 2.64 mmol) in water (1.4mL) was added to a suspension of methyl (*trans*-4-{4-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetate (152 mg, 0.33mmol) in THF (1.6mL) and MeOH (2.6 mL). After 24 h, 1M aqueous HCl (5mL) was added followed by water (10mL). The solid was filtered off and washed with water (2 x 5mL), Et₂O (2 x 5mL) then hexane (2 x 5mL) to give the title compound as a solid (105 mg, 71%); ¹H NMR δ 1.04-1.19 (5H, m), 1.37-1.51 (2H, m), 1.65-1.85 (5H, m), 2.11-2.15 (2H, m), 2.39-2.47 (1H, m), 2.57 (2H, q), 7.21 (4H, m), 7.49 (2H, d), 7.68 (2H, d), 10.84 (1H, s), 10.91 (1H, s), 11.99 (1H, s); MS m/e MH⁺ 449.







Examples 357-390






The following examples were prepared by the general procedure of Example 356 using the appropriate ester as starting material selected from Examples 321-355.














Example	R	¹ H NMR δ	MS m/e MH ⁺
357		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.15 (2H, d), 2.42-2.48 (1H, m), 3.75 (3H, s), 6.97-6.99 (2H, m), 7.22 (2H, d), 7.50-7.54 (2H, m), 7.68-7.70 (2H, m), 10.71 (1H, s), 10.87 (1H, s), 11.96 (1H, s)	451






358		1.14-1.18 (2H, m), 1.42-1.51 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.39-2.47 (1H, m), 7.23 (2H, d), 7.42 (1H, d), 7.63-7.71 (3H, m), 7.85-7.87 (1H, m), 8.01 (1H, s), 10.94 (1H, s), 11.35 (1H, s), 11.97 (1H, s)	489
359		1.05-1.20 (2H, m), 1.40-1.55 (2H, m), 1.70-1.95 (5H, m), 2.14-2.16 (2H, m), 2.46-2.55 (1H, m), 7.23 (2H, d), 7.53 (1H, d), 7.63 (1H, t), 7.70 (2H, d), 7.87 (1H, d), 8.02 (1H, s), 10.97 (1H, s), 11.43 (1H, s), 11.97 (1H, s)	446
360		1.08-1.18 (5H, m), 1.41-1.51 (2H, m), 1.71-1.84 (5H, m), 2.14-2.16 (2H, m), 2.45 (1H, t), 2.70 (2H, q), 7.15-7.30 (5H, m), 7.67-7.70 (1H, m), 7.69-7.69 (2H, m), 10.00 (1H, s), 10.86 (1H, s), 11.97 (1H, s)	449
361		1.05-1.20 (8H, m), 1.41-1.50 (2H, m), 1.70-1.84 (5H, m), 2.15 (2H, d), 2.40-2.50 (1H, m), 3.25-3.35 (1H, m), 7.21-7.28 (4H, m), 7.36-7.40 (1H, m), 7.55-7.60 (1H, m), 7.66-7.69 (2H, m), 10.01 (1H, s), 10.83 (1H, s), 11.97 (1H, s)	463
362		1.11-1.17 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.14-2.16 (2H, m), 2.46-2.55 (1H, m), 5.10 (2H, s), 7.06 (2H, d), 7.22 (2H, d), 7.32-7.53 (7H, m), 7.68-7.71 (2H, m), 10.72 (1H, s), 10.89 (1H, s), 11.97 (1H, s)	527
363		1.08-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.33 (2H, s), 2.46-2.55 (1H, m), 6.89 (1H, d), 7.22-7.29 (3H, m), 7.39 (1H, s), 7.44 (1H, d), 7.68-7.71 (2H, m), 10.90 (1H, s), 11.97 (1H, s)	435

364		1.08-1.18 (2H, m), 1.21 (6H, d), 1.41-1.51 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.42-2.47 (1H, m), 2.84-2.91 (1H, m), 7.22-7.27 (4H, m), 7.50-7.53 (2H, m), 7.68-7.71 (2H, m), 10.82 (1H, s), 10.89 (1H, s), 11.96 (1H, s)	463
365		1.10-1.17 (2H, m), 1.41-1.50 (2H, m), 1.70-1.84 (5H, m), 2.15 (2H, d), 2.45 (1H, t), 7.22 (2H, d), 7.50 (1H, t), 7.66-7.68 (2H, m), 7.75-7.86 (3H, m), 10.33 (1H, s), 10.87 (1H, s), 11.97 (1H, s)	489
366		1.11-1.18 (2H, m), 1.42-1.51 (2H, m), 1.73-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.53 (1H, m), 7.23 (2H, d), 7.69-7.72 (2H, m), 7.76-7.82 (4H, m), 10.96 (1H, s), 11.49 (1H, s), 12.00 (1H, s)	489
367		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.73-1.85 (5H, m), 2.15 (2H, d), 2.42-2.53 (1H, m), 7.23 (2H, d), 7.69-7.72 (2H, m), 7.76-7.78 (2H, m), 7.86-7.89 (2H, m), 10.98 (1H, s), 11.57 (1H, s), 11.97 (1H, s)	446
368		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.54 (1H, m), 7.23 (2H, d), 7.42 (2H, d), 7.68-7.73 (4H, m), 10.94 (1H, s), 11.18 (1H, s), 11.97 (1H, s)	505
369		1.11-1.15 (2H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 7.22-7.26 (3H, m), 7.44-7.50 (2H, m), 7.68-7.71 (2H, m), 8.20-8.22 (1H, m), 10.87 (1H, s), 10.93 (1H, s), 11.94 (1H, s)	505


370		1.11-1.15 (2H, m), 1.20 (3H, q), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 2.63 (2H, q), 6.92 (1H, d), 7.23 (2H, d), 7.30 (1H, t), 7.43-7.48 (2H, m), 7.68-7.71 (2H, m), 10.88 (1H, s), 10.93 (1H, s), 11.98 (1H, s)	449
371		1.12-1.17 (2H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 7.23 (2H, d), 7.36 (1H, s), 7.47 (1H, t), 7.55 (1H, d), 7.69-7.71 (2H, m), 7.79-7.82 (1H, m), 7.94 (1H, s), 8.04 (1H, t), 10.92 (1H, s), 11.06 (1H, s), 11.97 (1H, s)	464
372		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 7.23 (2H, d), 7.41-7.43 (1H, m), 7.57-7.61 (3H, m), 7.68-7.73 (4H, m), 7.78-7.80 (2H, m), 7.88-7.91 (1H, m), 8.08 (1H, t), 10.92 (1H, s), 11.22 (1H, s), 11.99 (1H, s)	525
373		1.12-1.17 (2H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 7.04-7.06 (1H, m), 7.23 (2H, d), 7.53-7.57 (2H, m), 7.68-7.71 (3H, m), 10.94 (1H, s), 11.33 (1H, s), 11.91 (1H, s)	505
374		1.11-1.17 (2H, m), 1.44-1.48 (2H, m), 1.73-1.84 (5H, m), 2.14-2.16 (2H, m), 2.48-2.56 (1H, m), 6.67-6.70 (1H, m), 7.06-7.08 (2H, m), 7.16-7.23 (3H, m), 7.31-7.44 (5H, m), 7.67-7.70 (2H, m), 10.91 (1H, s), 11.06 (1H, s), 11.97 (1H, s)	513

375		1.11-1.18 (2H, m), 1.44-1.48 (2H, m), 1.73-1.85 (5H, m), 2.15 (2H, d), 2.48-2.56 (1H, m), 5.21 (2H, s), 6.99-7.03 (1H, m), 7.08-7.12 (1H, m), 7.15-7.17 (1H, m), 7.22 (2H, d), 7.29-7.39 (3H, m), 7.50 (2H, d), 7.67-7.70 (2H, m), 7.86-7.88 (1H, m), 10.06 (1H, s), 10.87 (1H, s), 11.97 (1H, s)	527
376		δ1.13 (2H, d), 1.44-1.48 (2H, m), 1.65-1.87 (5H, m), 2.14-2.16 (2H, m), 2.48-2.56 (1H, m), 7.23 (2H, d), 7.36 (1H, t), 7.43-7.46 (2H, m), 7.52-7.61 (3H, m), 7.69 (2H, d), 7.97-7.99 (2H, m), 8.07 (1H, t), 10.35 (1H, s), 10.89 (1H, s), 10.97 (1H, s), 11.98 (1H, s)	540
377		1.11-1.17 (2H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.15 (2H, d), 2.48-2.56 (1H, m), 3.35 (3H, s), 7.23 (2H, d), 7.39-7.41 (2H, m), 7.68-7.70 (4H, m), 10.93 (1H, s), 11.16 (1H, s), 11.98 (1H, s)	515
378		δ1.12-1.17 (2H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.15 (2H, d), 2.46-2.53 (1H, m), 7.22-7.25 (2H, m), 7.61-7.74 (7H, m), 7.82-7.84 (1H, m), 7.95-7.97 (2H, m), 8.28 (1H, d), 10.94 (1H, s), 11.39 (1H, s), 11.98 (1H, s)	561
379		1.08-1.18 (2H, m), 1.41-1.51 (2H, m), 1.72-1.85 (5H, m), 2.15 (2H, d), 2.42-2.48 (1H, m), 3.40 (3H, s), 7.05-7.08 (1H, m), 7.23 (2H, d), 7.51 (1H, t), 7.56-7.59 (1H, m), 7.66 (1H, t), 7.68 (1H, s), 7.71 (1H, s), 10.94 (1H, s), 11.27 (1H, s), 11.98 (1H, s)	515

380		1.07-1.17 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 3.50 (3H, s), 7.19-7.24 (3H, m), 7.42-7.48 (2H, m), 7.69 (2H, d), 8.18-8.21 (1H, m), 10.64 (1H, s), 10.95 (1H, s), 11.97 (1H, s)	515
381		1.08-1.17 (2H, m), 1.42-1.50 (2H, m), 1.73-1.85 (5H, m), 2.15 (2H, d), 2.42-2.48 (1H, m), 3.85 (3H, s), 6.99-7.03 (1H, m), 7.09-7.15 (2H, m), 7.22 (2H, d), 7.68 (2H, d), 7.92-7.94 (1H, m), 10.04 (1H, s), 10.88 (1H, s), 11.97 (1H, s)	451
382		1.08-1.17 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.15 (2H, d), 2.42-2.48 (1H, m), 5.82 (2H, s), 7.17-7.24 (5H, m), 7.61-7.61 (1H, m), 7.68-7.70 (2H, m), 8.61 (1H, s), 10.84 (1H, s), 10.88 (1H, s), 11.97 (1H, s)	479
383		1.08-1.14 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.42-2.48 (1H, m), 3.68 (3H, s), 7.14 (1H, d), 7.22-7.34 (4H, m), 7.69 (2H, d), 7.75 (1H, t), 9.70 (1H, s), 10.88-10.91 (2H, m), 11.97 (1H, s)	494
384		1.08-1.17 (2H, m), 1.41-1.50 (2H, m), 1.71-1.84 (5H, m), 2.15 (2H, d), 2.42-2.48 (1H, m), 3.53-5.30 (8H, m), 7.22 (4H, d), 7.54 (2H, d), 7.69 (2H, d), 10.82 (1H, s), 10.89 (1H, s)	506
385		1.11-1.17 (2H, m), 1.41-1.50 (2H, m), 1.79-1.82 (5H, m), 2.15 (2H, d), 2.42-2.50 (1H, m), 7.23 (2H, d), 7.60-7.70 (5H, m), 7.80 (2H, d), 7.93-7.95 (2H, m), 8.00 (2H, d), 10.96 (1H, s), 11.55 (1H, s), 11.97 (1H, s)	559

386		1.11-1.15 (2H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.15 (2H, d), 2.42 - 2.53 (1H, m), 7.23 (2H, d), 7.35 - 7.50 (1H, m), 7.53-7.57 (2H, m), 7.61 - 7.69 (3H, m), 7.70 - 7.80 (1H, m), 7.93-7.95 (3H, m), 8.12 (1H, d), 10.11 (1H, s), 10.84 (1H, s), 11.97 (1H, s)	561
387		1.13-1.18 (2H, m), 1.43-1.54 (2H, m), 1.74-1.85 (5H, m), 2.16 (2H, d), 2.43 - 2.51 (1H, m), 7.25 (2H, d), 7.45 (1H, d), 7.50 (1H, d), 7.76-7.78 (3H, m), 8.26-8.28 (1H, m), 10.67 (1H, s), 11.98 (1H, s), 12.47 (1H, s)	446
388		1.10-1.20 (2H, m), 1.40-1.52 (2H, m), 1.74-1.90 (5H, m), 2.15 (2H, d), 2.43-2.51 (1H, m), 6.77 (1H, t), 6.97-7.04 (2H, m), 7.13 (2H, d), 7.18-7.24 (4H, m), 7.33 (2H, d), 7.48-7.50 (2H, m), 7.69 (2H, d), 8.03 (1H, s), 10.72 (1H, s), 10.88 (1H, s), 11.97 (1H, s)	512
389		1.08-1.18 (2H, m), 1.41-1.51 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41 - 2.52 (1H, m), 6.74-6.76 (1H, m), 6.85 (1H, t), 7.02-7.04 (1H, m), 7.12-7.15 (2H, m), 7.19-7.27 (5H, m), 7.48 (1H, t), 7.69 (2H, d), 8.27 (1H, s), 10.82 (1H, s), 10.89 (1H, s), 11.97 (1H, s)	512
390		1.10-1.15 (5H, m), 1.44-1.48 (2H, m), 1.74-1.85 (5H, m), 2.15 (2H, d), 2.42 - 2.51 (1H, m), 3.25 (2H, q), 7.24 (2H, d), 7.70 (2H, d), 7.82-7.85 (2H, m), 7.90-7.92 (2H, m), 10.98 (1H, s), 11.56 (1H, s), 11.97 (1H, s)	513


Example 391: Methyl 1-[(trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl}cyclohexyl)acetyl]piperidine-4-carboxylate








To a solution of (trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl}cyclohexyl)acetic acid (Example 112, 300 mg, 0.68 mmol) in DMF (3 mL) was added HATU (286 mg, 0.75 mmol), diisopropylethylamine (0.14 mL, 0.75 mmol) followed by methyl isonipecotate (0.15 mL, 0.75 mmol). The reaction mixture was allowed to stir at ambient temperature overnight. The DMF was evaporated and the residue re-dissolved in DMSO/MeCN/H₂O (7:2:1); a suspension formed which was filtered, washed with ether (2 x 10 mL) and dried to leave the title compound (137 mg). The filtrate was purified on an acidic reverse phase HPLC system eluting 5-95% MeCN 0.2% TFA, isolated 112 mg of the title compound (249 mg, 65%) as a solid; ¹H NMR δ 1.07-1.16 (m, 2H), 1.36-1.53 (m, 4H), 1.75-1.88 (m, 7H), 2.24 (d, 2H), 2.41-2.47 (m, 1H), 2.59-2.65 (m, 2H), 3.11 (t, 1H), 3.62 (s, 3H), 3.85 (d, 1H), 4.26 (d, 1H), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H); MS m/e MH⁺ 564.



Examples 392-400

The following examples were prepared by the general procedure of Example 391 using the appropriate amine in place of methyl isonipecotate:



Example	R	¹ H NMR δ	MS m/e MH ⁺
392		1.04-1.13 (m, 2H), 1.34 (s, 6H), 1.39-1.48 (m, 2H), 1.72-1.81 (m, 5H), 1.99 (d, 2H), 2.41-2.46 (m, 1H), 3.55 (s, 3H), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 8.04 (t, 1H), 8.13 (s, 1H), 10.75 (s, 1H), 10.93 (s, 1H)	560 (M+Na) ⁺

393		1.07-1.17 (m, 2H), 1.40-1.49 (m, 2H), 1.77-1.93 (m, 8H), 2.12-2.28 (m, 3H), 2.41-2.46 (m, 1H), 3.53-3.58 (m, 2H), 3.61 (s, 3H), 4.28-4.31 (m, 1H), 7.13-7.34 (m, 5H), 7.69 (d, 2H), 8.06 (t, 1H), 10.75 (s, 1H), 10.99 (s, 1H)	550
394		1.07-1.17 (m, 2H), 1.40-1.49 (m, 2H), 1.76-1.93 (m, 8H), 2.12-2.28 (m, 3H), 2.44-2.46 (m, 1H), 3.53-3.58 (m, 2H), 3.61 (s, 3H), 4.28-4.30 (m, 1H), 7.13-7.35 (m, 5H), 7.69 (d, 2H), 8.06 (t, 1H), 10.75 (s, 1H), 11.00 (s, 1H)	572 (M+Na) ⁺
395		1.05-1.17 (m, 2H), 1.39-1.49 (m, 2H), 1.76-1.92 (m, 6H), 2.08-2.24 (m, 3H), 2.41-2.46 (m, 1H), 3.43-3.65 (m, 6H inc s, 3H 3.61), 4.30 (t, 1H), 4.34 (s, 1H), 7.14- 7.35 (m, 5H), 7.69 (d, 2H), 8.06 (t, 1H), 10.82 (s, 1H), 11.00 (s, 1H)	566
396		1.04-1.12 (m, 2H), 1.18 (s, 6H), 1.39-1.48 (m, 2H), 1.71-1.80 (m, 5H), 1.98 (d, 2H), 2.40-2.46 (m, 1H), 3.39 (s, 2H), 4.88 (s, 1H), 7.14-7.34 (m, 6H), 7.68 (d, 2H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H)	532 M+Na
397		1.08-1.17 (m, 2H), 1.24-1.27 (m, 1H), 1.40-1.49 (m, 2H), 1.78-1.84 (m, 5H), 2.11-2.23 (m, 2H), 2.41-2.47 (m, 1H), 3.53-3.57 (m, 1H), 3.70-3.76 (m, 2H), 4.64-4.67 (m, 1H), 4.70-4.73 (m, 1H), 4.79 (s, 1H), 4.99 (s, 1H), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H)	536

398		1.07-1.15 (m, 2H), 1.40-1.49 (m, 2H), 1.77-1.84 (m, 5H), 2.22 (d, 2H), 2.42-2.48 (m, 1H), 2.82 (s, 3H), 2.98 (s, 3H), 7.14- 7.34 (m, 5H), 7.68 (d, 2H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H)	466
399		1.11 (m, 2H), 1.45 (m, 2H), 1.81 (m, 5H), 2.24 (m, 2H), 2.43 (m, 1H), 3.02 (s, 1.5H, NCH3), 3.02 (s, 1.5H, NCH3), 3.33-3.38 (m, 2H), 3.45-3.55 (m, 2H), 4.60 (t, 0.5H, OH), 4.77 (t, 0.5H, OH), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H) (rotamers detected)	496
400		1.05-1.13 (m, 2H), 1.39-1.48 (m, 2H), 1.71-1.80 (m, 5H), 2.04 (d, 2H), 2.41-2.47 (m, 1H), 2.97-3.04 (m, 1H), 3.17-3.23 (m, 2H), 3.46-3.51 (m, 1H), 4.39-4.78 (m, 1H), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 7.77 (t, 1H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H)	533 (M+Na) ⁺


Examples 401-405

The following examples were prepared by the general procedure of Example 112 using the appropriate starting material selected from Examples 391-395.

Example	Name	¹ H NMR δ	MS m/e MH ⁺
401	1-[(<i>trans</i> -4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetyl]piperidine-4-carboxylic acid	1.10-1.16 (m, 2H), 1.34-1.49 (m, 4H), 1.74-1.83 (m, 7H), 2.24 (d, 2H), 2.68-2.75 (m, 1H), 3.10 (t, 3H), 3.84 (d, 1H), 4.25 (d, 1H), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H), 12.22 (s, 1H)	550

402	<i>N</i> -[(<i>trans</i> -4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetyl]-2-methylalanine	1.04-1.13 (m, 2H), 1.33 (s, 6H), 1.38-1.47 (m, 2H), 1.72-1.83 (m, 5H), 1.99 (d, 2H), 2.41-2.47 (m, 1H), 7.14-7.34 (m, 5H), 7.68 (d, 2H), 7.96 (s, 1H), 8.04 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H), 12.00 (s, 1H)	524
403	1-[(<i>trans</i> -4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetyl]-D-proline	1.05-1.16 (m, 2H), 1.39-1.49 (m, 2H), 1.76-1.93 (m, 8H), 2.12-2.26 (m, 3H), 2.41-2.46 (m, 1H), 3.51-3.56 (m, 2H), 4.21-4.24 (m, 1H), 7.13-7.34 (m, 5H), 7.69 (d, 2H), 8.06 (t, 1H), 10.82 (s, 1H), 11.00 (s, 1H). CO ₂ H not seen	536
404	1-[(<i>trans</i> -4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetyl]-L-proline	1.07-1.17 (m, 2H), 1.40-1.49 (m, 2H), 1.76-1.93 (m, 8H), 2.12-2.26 (m, 3H), 2.41-2.47 (m, 1H), 3.51-3.56 (m, 2H), 4.21-4.24 (m, 1H), 7.14-7.35 (m, 5H), 7.69 (d, 2H), 8.06 (t, 1H), 10.82 (s, 1H), 11.00 (s, 1H), 12.36 (s, 1H) CO ₂ H not seen	534 (M-H) ⁺
405	(4 <i>R</i>)-1-[(<i>trans</i> -4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetyl]-4-hydroxy-L-proline	1.04-1.18 (m, 2H), 1.39-1.48 (m, 2H), 1.76-1.92 (m, 6H), 2.07-2.19 (m, 3H), 2.41-2.47 (m, 1H), 3.60-3.64 (m, 2H), 4.23 (t, 1H), 4.99 (s, 1H), 4.99 (s, 1H), 5.15 (s, 1H), 7.13-7.34 (m, 5H), 7.69 (d, 2H), 8.06 (t, 1H), 11.00 (s, 1H) CO ₂ H not seen	552


Example 406: Methyl (*trans*-4-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}cyclohexyl)acetate



Following the procedure described in Example 105 except using methyl (*trans*-4-{5-
5 { [hydrazino(oxo)acetyl]amino}pyridin-2-yl}cyclohexyl)acetate (Intermediate 78) and
2,4,5-trifluorophenylisothiocyanate as starting materials, the title compound was isolated in
79% yield; $^1\text{H NMR}$ δ 1.10-1.19 (m, 2H), 1.48-1.58 (m, 2H), 1.72-1.89 (m, 5H), 2.25 (d,
2H), 2.57-2.66 (m, 1H), 3.61 (s, 3H), 7.29 (d, 1H), 7.67-7.74 (m, 1H), 8.08 (d, 1H), 8.12-
8.20 (m, 1H), 8.84 (d, 1H), 11.07 (s, 1H), 11.20 (s, 1H); MS m/e MH^+ 490.


10

Example 407: (*trans*-4-{5-[(5-{(2,4,5-Trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}cyclohexyl)acetic acid



To a solution of methyl (*trans*-4-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-
15 yl)carbonyl)amino]pyridin-2-yl}cyclohexyl)acetate (Example 406, 355 mg, 0.725 mmol)
in THF (5 mL) and MeOH (25 mL) was added a 2M solution of NaOH (1.8 mL). The
resulting yellow solution was allowed to stir at ambient temperature for 16 hr then a further
1.8mL of 2M NaOH was added and stirring was continued at ambient temperature for 16
hr. The volatile organics were removed by evaporation and residue adjusted to pH ~1 with
20 2M HCl and the suspension was filtered and dried to leave 410 mg of a solid. This was
suspended in acetic acid (4 mL) and heated to 120°C, giving an incomplete solution. This
was allowed to cool, filtered and washed with water then dried overnight, to give the title
compound (167 mg, 48%) as a solid; $^1\text{H NMR}$ (CDCl_3) δ 1.10-1.19 (m, 2H), 1.56-1.65 (m,
2H), 1.72-1.79 (m, 1H), 1.85-1.94 (m, 4H), 2.17 (d, 2H), 2.79-2.85 (m, 1H), 7.67-7.76 (m,
25 2H), 8.12-8.19 (m, 1H), 8.44 (d, 1H), 9.01 (s, 1H), 11.14 (s, 1H), 11.58 (s, 1H), CO_2H not
seen; MS m/c MH^+ 476.

Example 408: [trans-4-(4-{[(5-{[4-(2-Hydroxyethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]-amino}phenyl)cyclohexyl]acetic acid




Thiocarbonyl diimidazole (117 mg, 0.65 mmol) was added in one portion to a stirred solution of 4-aminophenethyl alcohol (89 mg, 0.65 mmol) in DMA (2.5 mL) and the mixture was stirred at room temperature under an argon atmosphere for 2 h. Methyl [*trans*-4-(4-{[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl]acetate (Intermediate 43, 167 mg, 0.50 mmol) was added and the mixture was stirred at 80°C for 10 minutes in a microwave. EDCI (125 mg, 0.65 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water, stirred for 30 minutes and then filtered to leave the crude oxadiazole methyl ester intermediate as a brown solid (230 mg) that was used with no further purification; MS *m/e* MH⁺ 479.







A solution of lithium hydroxide (170 mg, 4.0 mmol) in H₂O (3 mL) was added in one portion to a solution of the crude oxadiazole product (230 mg) in a mixture of THF:MeOH (1:1, 10 mL) and the reaction mixture was stirred for 20 h. Citric acid (15 mL) was added and the mixture was stirred for 30 minutes and then filtered. The solid was washed twice with water, twice with diethylether and once with isohexane to give the title compound (150 mg, 67%) as a solid; ¹H NMR δ 1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 2.71 (2H, t), 3.52-3.67 (2H, m), 4.6 (1H, t), 7.21 (2H, d), 7.22 (2H, d), 7.51 (2H, d), 7.7 (2H, d), 10.8 (1H, s), 10.9 (1H, s), 12.0 (1H, s); MS m/e MH⁺ 465.







Examples 409 – 423




25 The following Examples were prepared by the general procedure of Example 408, using methyl [*trans*-4-(4-{[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl]acetate (Intermediate 43) and the appropriate commercially available aniline, RNH₂.




- 212 -

Example	R	¹ H NMR δ	MS m/e MH ⁺
409		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 4.2 (1H, s), 7.18 (1H, dd), 7.23 (2H, d), 7.42 (1H, dd), 7.53 (1H, dd), 7.7 (1H, d), 7.77 (1H, s), 10.9 (1H, s), 11.15 (1H, s), 12.0 (1H, s)	445
410		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 4.0 (2H, s), 7.24 (2H, d), 7.4 (2H, d), 7.62 (2H, d), 7.7 (2H, d), 10.9 (1H, s), 11.1 (1H, s), 12.0 (1H, s)	460
411		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.2 (2H, d), 7.3-7.5 (8H, m), 7.65 (2H, d), 7.75 (1H, d), 10.1 (1H, s), 10.85 (1H, s), 12.05 (1H, s)	497
412		1.04-1.2 (2H, m), 1.29 (9H, s), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.01 (2H, d), 7.24 (2H, d), 7.52 (2H, d), 7.7 (2H, d), 10.8 (1H, s), 10.9 (1H, s), 12.0 (1H, s)	493
413		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.0 (2H, d), 7.1 (2H, d), 7.11 (1H, dd), 7.22 (2H, d), 7.36 (2H, dd), 7.61 (2H, d), 7.7 (2H, d), 10.9 (1H, s), 10.95 (1H, s), 12.0 (1H, s)	513
414		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.23 (2H, d), 7.7 (2H, d), 7.79 (2H, d), 7.9 (2H, d), 8.23 (1H, s), 9.2 (1H, s), 10.9 (1H, s), 11.2	488

		(1H, s)	
415		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 3.94 (2H, s), 7.25 (9H, m), 7.53 (2H, d), 7.7 (2H, d), 10.85 (1H, s), 10.89 (1H, s), 11.95 (1H, s)	511
416		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.05 (3H, s), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.22 (2H, d), 7.32 (3H, m), 7.7 (2H, d), 7.84 (1H, s), 10.0 (1H, s), 10.87 (1H, s), 10.92 (1H, s), 12.0 (1H, s)	478
417		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.13 (1H, dd), 7.22 (2H, d), 7.44 (1H, dd), 7.54 (1H, dd), 7.7 (2H, d), 7.75 (1H, d), 10.94 (1H, s), 11.2 (1H, s), 12.0 (1H, s)	455
418		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 4.11 (1H, s), 7.25 (2H, d), 7.54 (2H, d), 7.64 (2H, d), 7.7 (2H, d), 11.0 (1H, s), 11.28 (1H, s), 12.05 (1H, s)	445
419		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 7.22 (2H, d), 7.24 (1H, dd), 7.36 (2H, dd), 7.58 (2H, dd), 7.7 (2H, d), 7.98 (1H, d), 10.95 (1H, s), 11.19 (1H, s), 11.95 (1H, s)	499
420		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 6.85-6.95 (1H, m), 7.2 (1H, t), 7.22 (2H, d), 7.42 (2H, dd), 7.5 (1H, s), 7.7 (2H, d), 10.92 (1H, s), 11.19 (1H, s), 11.96 (1H, s)	487

421		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 3.3 (3H, s), 4.54 (2H, s), 7.21 (1H, dd), 7.23 (2H, d), 7.38 (1H, dd), 7.4 (1H, dd), 7.7 (2H, d), 7.81 (1H, dd), 9.88 (1H, s), 10.9 (1H, s), 12.0 (1H, s)	465
422		1.04-1.2 (2H, m), 1.38-1.55 (2H, m), 1.65-1.9 (5H, m), 2.15 (2H, d), 2.38-2.55 (1H, m), 2.72 (3H, s), 7.23 (2H, d), 7.58 (1H, dd), 7.7 (2H, d), 7.8-7.9 (3H, m), 8.47 (1H, s), 8.79 (1H, d), 10.9 (1H, s), 11.15 (1H, s), 12.0 (1H, s)	513
423		1.0-1.55 (10H, m), 1.62-1.9 (9H, m), 2.15 (2H, d), 2.36-2.6 (2H, m), 7.21 (2H, d), 7.23 (2H, d), 7.52 (2H, d), 7.7 (2H, d), 10.8 (1H, s), 10.9 (1H, s), 11.9 (1H, s)	503


Example 424: Methyl *trans*-4-[4-({[5-(pyrimidin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)-phenyl]cyclohexyl]acetate



Di-2-pyridyl thionocarbonate (367 mg, 1.58 mmol) was added in one portion to a stirred solution of 4-aminopyrimidine (150 mg, 1.58 mmol) in DCM (3 mL) and the reaction mixture was heated at 100°C for 20 minutes in the microwave and then cooled to room temperature. This solution was added to a suspension of methyl [*trans*-4-(4-
5 {[hydrazino(oxo)acetyl]amino}-phenyl)cyclohexyl]acetate (Intermediate 43, 421 mg, 1.26 mmol) in DCM (5 mL) and the mixture stirred at room temperature for 5 minutes. DMF (8
10 mL) was added and the DCM was removed under reduced pressure. The solution was heated at 85°C for 40 minutes in the microwave and then cooled to room temperature. EDCI (303 mg, 1.58 mmol) was added in one portion and the reaction mixture was heated at 85°C for 60 minutes in the microwave. The reaction was quenched by the addition of H₂O (200 mL) and EtOAc (250 mL) was added. The layers were separated and the


aqueous layer was extracted with EtOAc (2x150 mL). The combined organic layer was washed with brine (100 mL), dried (MgSO₄) and concentrated *in vacuo* to leave a dark brown gummy solid. MeOH (40 mL) was added and the mixture was heated to reflux for 2 minutes and then cooled to room temperature to leave a white suspension. The solid was
5 filtered to give the title compound (110 mg, 20%) as a solid; ¹H NMR δ 1.10-1.17 (2H, m), 1.43-1.47 (2H, m), 1.75-1.81 (5H, m), 2.24-2.26 (2H, d), 2.44 (1H, t), 3.61 (3H, s), 7.24 (2H, d), 7.69-7.73 (2H, m), 7.92 (1H, d), 8.69 (1H, d), 8.87 (1H, s), 11.02 (1H, s); MS m/e MH⁺ 437.

10 Example 425: Methyl (*trans*-4-{4-[(5-(1-methyl-1H-pyrazol-3-yl)amino)-1,3,4-oxadiazol-2-yl]-carbonyl}amino)phenyl)cyclohexyl)acetate



Di-2-pyridyl thionocarbonate (359 mg, 1.54 mmol) was added in one portion to a stirred solution of 3-amino-1-methylpyrazole (150 mg, 1.54 mmol) in CH₃CN (8 mL) and the reaction mixture was heated at 80°C for 10 minutes in the microwave and then cooled to
15 room temperature. Methyl [*trans*-4-(4-{[hydrazino(oxo)acetyl]amino}-phenyl)cyclohexyl)acetate (Intermediate 43, 412 mg, 1.24 mmol) in DCM (5 mL) was added in one portion and the mixture was heated at 85°C for 20 minutes and then cooled to room temperature. EDCI (297 mg, 1.54 mmol) was added in one portion and the reaction mixture was heated to 85°C for 30 minutes. The mixture was cooled to room temperature
20 and filtered to leave a solid. The solid was washed with CH₃CN (5 mL) to give the title compound (373 mg, 69%) as a solid; ¹H NMR δ 1.09-1.18 (2H, m), 1.41-1.50 (2H, m), 1.73-1.81 (5H, m), 2.24-2.26 (2H, d), 2.41-2.47 (1H, m), 3.61 (3H, s), 3.77 (3H, s), 6.41 (1H, d), 7.21-7.23 (2H, d), 7.64 (1H, d), 7.66-7.69 (2H, d), 10.84 (1H, s), 11.13 (1H, s); MS m/e MH⁺ 439.


25 Example 426: Methyl (*trans*-4-[4-[(5-(pyridin-3-ylamino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)-phenyl)cyclohexyl)acetate



3-Pyridyl isothiocyanate (41 mg, 0.3 mmol) was added in one portion to a suspension of methyl [*trans*-4-(4-{{[hydrazino(oxo)acetyl]amino}-phenyl}cyclohexyl]acetate (Intermediate 43, 100 mg, 0.3 mmol) in CH₃CN (5 mL) and the reaction mixture was heated at 80°C for 10 minutes in the microwave and then cooled to room temperature.


- 5 EDCI (58 mg, 0.3 mmol) was added in one portion and the reaction mixture was heated to 90°C for 20 minutes. The mixture was cooled to room temperature and filtered to leave a solid. The solid was washed with CH₃CN (5 mL) to give the title compound (90 mg, 69%) as a solid; ¹H NMR δ 1.12-1.18 (2H, m), 1.45 (2H, q), 1.80 (5H, m), 2.24-2.26 (2H, d), 2.42-2.48 (1H, m), 3.61 (3H, s), 7.23 (2H, d), 7.43-7.46 (1H, m), 7.69-7.71 (2H, m), 8.06-8.09 (1H, m), 8.28-8.29 (1H, m), 8.78 (1H, d), 10.95 (1H, s), 11.22 (1H, s); MS m/e MH⁺
- 10 436.

Example 427: Methyl {*trans*-4-[4-{{[5-(pyridin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl]amino}-phenyl]cyclohexyl}acetate




- 15 Thiocarbonyl diimidazole (190 mg, 1.06 mmol) was added in one portion to a solution of 4-aminopyridine (100 mg, 1.06 mmol) in DMA (8 mL) and the solution was stirred at room temperature for 3 h. Methyl [*trans*-4-(4-{{[hydrazino(oxo)acetyl]amino}phenyl}cyclohexyl]-acetate (Intermediate 43, 284 mg, 0.85 mmol) was added in one portion and the reaction mixture was stirred at room temperature for 30 minutes and then heated at
- 20 80°C for 10 minutes in the microwave. The solution was cooled to room temperature and EDCI (205 mg, 1.06 mmol) was added in one portion and the reaction mixture was heated to 90°C for 20 minutes in the microwave. The reaction mixture was cooled to room temperature and poured into water (200 mL). The mixture was filtered and the resulting solid was recrystallised (CH₃CN) to give the title compound (93 mg, 20%) as a solid; ¹H
- 25 NMR δ 1.15 (2H, t), 1.42-1.51 (2H, m), 1.74-1.82 (5H, m), 2.24-2.26 (2H, d), 3.61 (3H, s), 7.23 (2H, d), 7.54-7.55 (2H, m), 7.69-7.71 (2H, m), 8.43 (2H, s) MS m/e MH⁺ 436.

Example 428: Methyl {*trans*-4-[4-({[5-(pyrazin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)-phenyl]cyclohexyl}acetate



Thiocarbonyl diimidazole (179 mg, 1.00 mmol) was added in one portion to a solution of aminopyrazine (96 mg, 1.00 mmol) in DMA (5 mL) and the solution was stirred at room temperature for 6 h. Methyl [*trans*-4-(4-{{[hydrazino(oxo)acetyl]amino}phenyl})cyclohexyl]-acetate (Intermediate 43, 267 mg, 0.80 mmol) was added in one portion and the reaction mixture was stirred at room temperature for 20 h and then heated at 80°C for 10 minutes in the microwave. The solution was cooled to room temperature and EDCI (192 mg, 1.00 mmol) was added in one portion and the reaction mixture was heated to 90°C for 15 minutes in the microwave. The reaction mixture was cooled to room temperature and poured into water (200 mL). The mixture was filtered and the resulting solid was recrystallised (CH₃CN) to give the title compound as an off white solid (35 mg, 8%); ¹H NMR δ 1.15 (2H, m), 1.45-1.48 (2H, m), 1.79 (6H, m), 2.26 (2H, d), 2.45-2.49 (1H, m), 3.61 (3H, s), 7.23-7.25 (2H, d), 7.69-7.71 (2H, d), 8.36 (1H, d), 8.42 (1H, t), 9.19 (1H, s), 11.01 (1H, s); MS m/e MH⁺ 437.


Example 429: Methyl {*trans*-4-[4-({[5-(1,3-thiazol-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetate



Thiocarbonyl diimidazole (179 mg, 1.00 mmol) was added in one portion to a solution of 2-aminothiazole (100 mg, 1.00 mmol) in DMA (5 mL) and the solution was stirred at room temperature for 6 h. Methyl [*trans*-4-(4-{{[hydrazino(oxo)acetyl]amino}phenyl})cyclohexyl]-acetate (Intermediate 43, 267 mg, 0.80 mmol) was added in one portion and the reaction mixture was stirred at room temperature for 20 h and then heated at 80°C for 10 minutes in the microwave. The solution was cooled to room temperature and EDCI (192 mg, 1.00 mmol) was added in one portion and the reaction mixture was heated to

90°C for 15 minutes in the microwave. The reaction mixture was cooled to room temperature and poured into water (200 mL). The mixture was filtered and the resulting solid was recrystallised (CH₃CN) to give the title compound (130 mg, 29%) as a solid; ¹H NMR δ 1.09-1.18 (2H, m), 1.42-1.47 (2H, m), 1.79 (5H, m), 2.24-2.26 (2H, d), 2.45 (1H, t), 3.61 (3H, s), 6.99 (1H, d), 7.22 (2H, d), 7.38 (1H, d), 7.67-7.70 (2H, m), 10.78 (1H, s); MS m/e MH⁺ 442.

Example 430: {trans-4-[4-({[5-(Pyrimidin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)-phenyl]cyclohexyl}acetic acid




10

A solution of lithium hydroxide (101 mg, 2.41 mmol) in H₂O (1.5 mL) was added in one portion to a solution of methyl {trans-4-[4-({[5-(pyrimidin-4-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetate (Example 416, 105 mg, 0.24 mmol) in a mixture of THF (1.8 mL) and MeOH (3 mL) and the reaction mixture was stirred at 35°C for 2 h. Citric acid (50 mL) and EtOAc (50 mL) were added and the layers were separated and the aqueous layer was filtered to leave a solid. The solid was washed with water (10 mL) to give the title compound as a yellow brown solid (60 mg, 59%); ¹H NMR δ 1.07-1.18 (2H, m), 1.41-1.51 (2H, m), 1.69-1.85 (5H, m), 2.14-2.16 (2H, d), 7.24 (2H, d), 7.71 (2H, d), 7.92 (1H, d), 8.69 (1H, d), 8.87 (1H, s), 11.00 (1H, s), 12.06 (2H, s); MS m/e MH⁺ 423.






20

Examples 431-435


The following examples were prepared by the general procedure of Example 430 using the appropriate ester as starting material selected from Examples 425-429.



25

Example	R	¹ H NMR δ	MS m/e MH ⁺
431		1.07-1.17 (2H, m), 1.41-1.51 (2H, m), 1.70-1.84 (6H, m), 2.14-2.16 (2H, d), 3.77 (3H, s), 6.41-6.41 (1H, m), 7.21-7.23 (2H, d), 7.63-7.64 (1H, m), 7.66-7.69 (2H, d), 10.84 (1H, s), 11.13 (1H, s), 11.97 (1H, s)	425
432		1.07-1.18 (2H, m), 1.41-1.51 (2H, m), 1.70-1.85 (5H, m), 2.14-2.16 (2H, d), 7.23 (2H, d), 7.43-7.46 (1H, m), 7.69-7.71 (2H, d), 8.06-8.09 (1H, m), 8.28-8.29 (1H, m), 8.78 (1H, d), 10.94 (1H, s), 11.22 (1H, s), 11.97 (1H, s)	422
433		1.11-1.15 (2H, m), 1.45-1.49 (2H, m), 1.74-1.85 (5H, m), 2.15-2.16 (2H, d), 7.24 (2H, d), 7.54-7.55 (2H, m), 7.70 (2H, d), 8.42 (2H, s), 10.95 (1H, s)	422
434		1.12-1.15 (2H, m), 1.45-1.49 (2H, m), 1.82 (6H, m), 2.15-2.16 (2H, d), 7.24 (2H, d), 7.70 (2H, d), 8.36 (1H, d), 8.42-8.43 (1H, m), 9.19 (1H, s), 11.01 (1H, s)	423
435		1.07-1.17 (2H, m), 1.41-1.51 (2H, m), 1.69-1.84 (6H, m), 2.15 (2H, d), 2.42-2.53 (1H, m), 6.99 (1H, d), 7.22 (2H, d), 7.38 (1H, d), 7.68 (2H, d), 10.78 (1H, s)	428


Example 436: (trans-4-{4-[(5-[(1-Methyl-1H-pyrazol-5-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]-amino}phenyl)cyclohexyl)acetic acid



Thiocarbonyl diimidazole (179 mg, 1.00 mmol) was added in one portion to a solution of 5-amino-1-methyl pyrazole (98 mg, 1.00 mmol) in DMA (5 mL) and the solution was stirred at room temperature for 6 h. Methyl [*trans*-4-(4-{[hydrazino(oxo)acetyl]amino} phenyl)-cyclohexyl]acetate (Intermediate 43, 267 mg, 0.80 mmol) was added in one portion and the reaction mixture was stirred at room temperature for 20 h and then heated at 80°C for 10 minutes in the microwave. The solution was cooled to room temperature and EDCI (192 mg, 1.00 mmol) was added in one portion and the reaction mixture was heated to 90°C for 15 minutes in the microwave. The reaction mixture was cooled to room temperature and poured into water (200 mL). The mixture was concentrated *in vacuo* to leave a gum, which was recrystallised (CH₃CN) to leave a gummy solid. Filtration followed by concentration *in vacuo* of the filtrate gave the crude oxadiazole methyl ester intermediate that was used with no further purification; MS *m/e* MH⁺ 439.

Lithium hydroxide (420 mg, 10.0 mmol) was added in one portion to a solution of the crude oxadiazole methyl ester intermediate in a mixture of THF (4 mL), MeOH (8 mL) and water (4 mL) and the reaction mixture was stirred at 35°C for 4 h. Citric acid (30 mL) was added and the mixture was filtered to leave a solid that was recrystallised (MeOH) to give the title compound (87 mg, 26%) as a solid; ¹H NMR δ 1.07-1.17 (2H, m), 1.41-1.51 (2H, m), 1.70-1.84 (5H, m), 2.14-2.16 (2H, d), 3.75 (3H, s), 6.70 (1H, s), 7.21-7.24 (2H, m), 7.40 (1H, s), 7.67-7.74 (2H, m); MS *m/e* MH⁺ 425.

Example 437: [*trans*-4-[4-({[5-(Pyrimidin-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)-phenyl]cyclohexyl]acetic acid




A suspension of 2-aminopyrimidine (100 mg, 1.05 mmol) in DCM (2 mL) was added to a solution of thiophosgene (81 μL, 1.05 mmol) in DCM (3 mL) and then triethylamine (294 μL, 2.10 mmol) was added in one portion. The reaction mixture was stirred for 1 h 30 minutes and then methyl [*trans*-4-(4-{[hydrazino(oxo)acetyl]amino}phenyl) cyclohexyl]acetate (Intermediate 43, 300 mg, 0.90 mmol) was added in one portion, the reaction mixture was stirred at room temperature for 1 h and then heated at 80°C for 10 minutes in the microwave. The solution was cooled to room temperature and EDCI (150

mg, 0.80 mmol) was added in one portion and the reaction mixture was heated to 90°C for 20 minutes in the microwave. The reaction mixture was cooled to room temperature and filtered. The filtrate was concentrated *in vacuo* to leave a brown solid, which was purified by column chromatography, using 0-20% MeOH and EtOAc as eluent to give the crude oxadiazole methyl ester intermediate that was used with no further purification; MS *m/e* MH⁺ 437.

A solution of lithium hydroxide (35 mg, 0.8 mmol) in H₂O (4 mL) was added in one portion to a solution of the crude oxadiazole methyl ester intermediate in a mixture of THF (2 mL) and MeOH (2 mL) and the reaction mixture was stirred at 35°C for 2 h. Citric acid (15 mL) was added and the mixture was filtered to leave a solid that was recrystallised (MeOH) to give the title compound (5 mg, 1%) as a solid; ¹H NMR δ 1.12 (2H, m), 1.45-1.48 (2H, m), 1.80 (6H, m), 2.15-2.16 (2H, d), 7.17 (1H, t), 7.22 (2H, d), 7.70 (2H, d), 8.68 (2H, d), 11.00 (1H, s); MS *m/e* MH⁺ 423.

Example 438: trans-4-(4-[(5-[(2-Methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]aminophenyl)-cyclohexanecarboxylic acid




A solution of 2-methoxyphenyl isothiocyanate (59 mg, 0.36 mmol) in DMA (2 mL) was added to *tert*-butyl *trans*-4-(4-[[hydrazino(oxo)acetyl]amino}phenyl)cyclohexanecarboxylate (Intermediate 79, 108 mg, 0.30 mmol) and the mixture was stirred at 50°C for 2 h. EDCI (86 mg, 0.45 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water (20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl ester intermediate as a solid (145 mg) that was used with no further purification; MS *m/e* (M-H)⁺ 491.

The crude oxadiazole *tert*-butyl ester intermediate (145 mg) was dissolved in TFA (2.5 mL) and stirred for 6 h. The reaction mixture was concentrated *in vacuo* to leave a solid, which was purified by recrystallisation from a mixture of DMSO:CH₃CN:H₂O (7:2:1) to give the title compound (50 mg, 39%) as a solid; ¹H NMR δ 1.4-1.58 (4H, m), 1.8-1.9 (2H,

m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 3.85 (3H, s), 7.0 (1H, ddd), 7.14 (2H, m), 7.22 (2H, d), 7.7 (2H, d), 7.95 (1H, dd), 10.05 (1H, s), 10.9 (1H, s), 12.0 (1H, s); MS m/e (M-H)⁺ 435.

5 **Example 439: *trans*-4-(4-[[[5-[[3-(Hydroxymethyl)phenyl]amino]-1,3,4-oxadiazol-2-yl]carbonyl]-amino]phenyl)cyclohexanecarboxylic acid**




A solution of (3-hydroxymethyl)phenyl isothiocyanate (59 mg, 0.36 mmol) in DMA (2 mL) was added to *tert*-butyl *trans*-4-(4-[[[hydrazino(oxo)acetyl]amino]phenyl)-cyclohexanecarboxylate (Intermediate 79, 108 mg, 0.30 mmol) and the mixture was stirred at 50°C for 2 h. EDCI (86 mg, 0.45 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water (20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl ester intermediate as a solid (160 mg) that was used with no further purification; MS m/e (M-H)⁺ 491.

The crude oxadiazole *tert*-butyl ester intermediate (160 mg) was dissolved in TFA (2.5 mL) and stirred for 6 h. The reaction mixture was concentrated *in vacuo* to leave an oil, which was stirred with a mixture of MeOH:H₂O:triethylamine (4:1:1, 6 mL) for 10 minutes. Concentration *in vacuo* left a brown solid, which was purified by recrystallisation from a mixture of DMSO:CH₃CN:H₂O (7:2:1) to give the title compound (67 mg, 47%) as a solid; ¹H NMR δ 1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 4.5 (2H, d), 5.2 (1H, t), 7.0 (1H, d), 7.25 (2H, d), 7.35 (1H, dd), 7.55 (2H, m), 7.7 (2H, d), 10.9 (2H, s), 12.0 (1H, s); MS m/e (M-H)⁺ 435.


25 **Examples 440-445**

The following Examples were prepared by the general procedure of Example 431, using *tert*-butyl *trans*-4-(4-[[[hydrazino(oxo)acetyl]amino]phenyl)cyclohexane-carboxylate (Intermediate 79) and commercially available isothiocyanates, R-NCS.



Example	R	¹ H NMR δ	MS m/e M-H ⁺
440		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.2 (2H, d), 7.5 (1H, dd), 7.7 (2H, d), 7.7-7.9 (3H, m), 10.4 (1H, s), 10.9 (1H, s), 12.0 (1H, s)	473
441		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.2-7.3 (3H, m), 7.4-7.5 (2H, m), 7.7 (2H, d), 8.2 (1H, dd), 10.9 (1H, s), 11.0 (1H, s), 12.0 (1H, s)	489
442		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.2 (2H, d), 7.4 (1H, d), 7.65 (1H, dd), 7.7 (2H, d), 7.85 (1H, dd), 8.0 (1H, s), 10.9 (1H, s), 11.3-12.3 (2H, m)	473
443		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 3.75 (3H, s), 7.0 (2H, d), 7.2 (2H, d), 7.5 (2H, d), 7.7 (2H, d), 10.7 (1H, s), 10.9 (1H, s), 12.0 (1H, s)	435
444		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.23 (2H, d), 7.45 (1H, dd), 7.65-7.75 (4H, m), 10.9 (1H, s), 11.2 (1H, s), 12.0 (1H, s)	489
445		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.22 (2H, d), 7.5 (1H, dd), 7.7 (2H, d), 7.75-7.9 (4H, m), 10.95 (1H, s), 11.5 (1H, s), 12.0 (1H, s)	473

Example 446: *trans*-4-(4-{{(5-{{2-(Benzyloxy)phenyl}amino}-1,3,4-oxadiazol-2-yl)carbonyl}amino}-phenyl)cyclohexanecarboxylic acid




Thiocarbonyl diimidazole (64 mg, 0.36 mmol) was added in one portion to a stirred
 5 solution of 2-benzyloxyaniline (72 mg, 0.36 mmol) in DMA (2 mL) and the mixture was
 stirred at room temperature under an argon atmosphere for 2 h. *tert*-Butyl *trans*-4-(4-
 {{hydrazino-(oxo)acetyl}amino}phenyl)cyclohexanecarboxylate (Intermediate 79, 108 mg,
 0.30 mmol) was added and the mixture was stirred at 80°C for 10 minutes in a microwave.
 EDCI (69 mg, 0.36 mmol) was then added in one portion and the reaction mixture was
 10 heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water
 (20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl
 ester intermediate as a brown solid (130 mg) that was used with no further purification;
 MS *m/e* (M-H)⁺ 567.

The crude oxadiazole *tert*-butyl ester intermediate (130 mg) was dissolved in TFA (2 mL)
 15 and stirred for 2 h. The reaction mixture was concentrated *in vacuo* to leave a brown solid,
 which was purified by recrystallisation from a mixture of DMSO:CH₃CN:H₂O (7:2:1) to
 give the title compound (35 mg, 30%) as a solid; ¹H NMR δ 1.4-1.58 (4H, m), 1.8-1.9 (2H,
 m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 5.22 (2H, s), 7.02 (1H, dd), 7.1
 (1H, dd), 7.17 (1H, d), 7.22 (2H, d), 7.35 (3H, m), 7.5 (1H, d), 7.5 (1H, dd), 7.7 (2H, d),
 20 7.88 (1H, dd), 10.0 (1H, s), 10.9 (1H, s), 12.0 (1H, s); MS *m/e* (M-H)⁺ 511.


Examples 447 - 449

The following Examples were prepared by the general procedure of Example 446, using
 25 *tert*-butyl *trans*-4-(4-{{hydrazino(oxo)acetyl}amino}phenyl)cyclohexane-carboxylate
 (Intermediate 79) and the appropriate commercially available aniline, RNH₂.



Example	R	¹ H NMR δ	MS m/e M-H ⁺
447		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.05 (1H, dd), 7.22 (2H, d), 7.5-7.6 (2H, m), 7.7 (2H, d), 7.65-7.75 (1H, m), 10.95 (1H, s), 11.35 (1H, s), 12.0 (1H, s)	489
448		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 4.55 (2H, s), 7.15-7.25 (1H, m), 7.22 (2H, d), 7.38 (1H, dd), 7.4 (1H, dd), 7.5 (1H, dd), 7.7 (2H, d), 7.82 (1H, d), 9.95 (1H, s), 10.9 (1H, s), 12.0 (1H, s)	449
449		1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 7.24 (2H, d), 7.6-7.7 (7H, m), 7.8 (1H, dd), 7.95 (2H, dd), 8.3 (1H, s), 10.95 (1H, s), 11.4 (1H, s), 12.0 (1H, s)	545

Example 450: *trans*-4-{4-[(5-[(3-Anilinophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}-cyclohexanecarboxylic acid



- 5 Thiocarbonyl diimidazole (64 mg, 0.36 mmol) was added in one portion to a stirred solution of 3-aminodiphenylamine (66 mg, 0.36 mmol) in DMA (2 mL) and the mixture was stirred at room temperature under an argon atmosphere for 2 h. *tert*-Butyl *trans*-4-(4-{[hydrazino(oxo)-acetyl]amino}phenyl)cyclohexanecarboxylate (Intermediate 79, 108 mg, 0.30 mmol) was added and the mixture was stirred at 80°C for 10 minutes in a microwave.
- 10 EDCI (69 mg, 0.36 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water


247

(20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl ester intermediate as a brown solid (130 mg) that was used with no further purification; MS *m/e* (M-H)⁺ 552.

5 The crude oxadiazole *tert*-butyl ester intermediate (130 mg) was dissolved in TFA (2 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo* to leave a brown solid, which was triturated with MeOH and filtered to give the title compound (65 mg, 51%) as a solid; ¹H NMR δ 1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 6.75 (1H, dd), 6.85 (1H, dd), 7.05 (1H, dd), 7.1-7.35 (6H, m), 7.5 (1H, s), 7.7 (2H, d), 8.3 (1H, s), 10.8 (1H, s), 10.9 (1H, s), 12.0 (1H, s); MS *m/e* (M-H)⁺ 496.

10


Example 451: *trans*-4-[4-[(15-[(4-Anilinophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]aminophenyl]-cyclohexanecarboxylic acid



Example 451 was prepared using the procedure of Example 450, using *tert*-butyl *trans*-4-(4-[[hydrazino(oxo)acetyl]amino]phenyl)cyclohexanecarboxylate (Intermediate 79) and 4-aminodiphenylamine, to give the title compound (38 mg, 26%) as a solid; ¹H NMR δ 1.4-1.58 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 6.78 (1H, dd), 7.0 (2H, d), 7.15 (2H, d), 7.15-7.3 (3H, m), 7.5 (2H, d), 7.7 (2H, d), 8.0 (1H, s), 10.7 (1H, s), 10.9 (1H, s), 12.0 (1H, s); MS *m/e* (M-H)⁺ 496.

20

Example 452: *trans*-4-[4-[(15-[(2-[(Methylsulfonyl)oxy]phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]-amino]phenyl]cyclohexanecarboxylic acid




25 Thiocarbonyl diimidazole (64 mg, 0.36 mmol) was added in one portion to a stirred solution of 2-aminophenyl methanesulfonate (67 mg, 0.36 mmol) in DMA (2 mL) and the mixture was stirred at room temperature under an argon atmosphere for 2 h. *tert*-Butyl *trans*-4-(4-[[hydrazino(oxo)-acetyl]amino]phenyl)cyclohexanecarboxylate (Intermediate

79, 108 mg, 0.30 mmol) was added and the mixture was stirred at 80°C for 10 minutes in a microwave. EDCI (69 mg, 0.36 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water (20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl ester intermediate as a solid (210 mg) that was used with no further purification; MS *m/e* (M-H)⁺ 555.

The crude oxadiazole *tert*-butyl ester intermediate (210 mg) was dissolved in TFA (2 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo* to leave a brown solid, which was purified by recrystallisation from a mixture of DMSO:CH₃CN:H₂O (7:2:1) and then recrystallisation from AcOH to give the title compound (21 mg, 14%) as a solid; ¹H NMR δ 1.3-1.5 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 3.45 (3H, s), 7.1-7.2 (1H, m), 7.2 (2H, d), 7.4 (2H, dd), 7.65 (2H, d), 8.3 (1H, d), 10.7 (1H, s), 10.9 (1H, s), 12.0 (1H, s); MS *m/e* MH⁺ 501.

Example 453: *trans*-4-[4-({[5-({[3-({[Methylsulfonyl]oxy}phenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl})-amino]phenyl]cyclohexanecarboxylic acid



Thiocarbonyl diimidazole (64 mg, 0.36 mmol) was added in one portion to a stirred solution of 3-aminophenyl methanesulfonate (67 mg, 0.36 mmol) in DMA (2 mL) and the mixture was stirred at room temperature under an argon atmosphere for 2 h. *tert*-Butyl *trans*-4-(4-([hydrazino-(oxo)acetyl]amino)phenyl)cyclohexanecarboxylate (Intermediate 79, 108 mg, 0.30 mmol) was added and the mixture was stirred at 80°C for 10 minutes in a microwave. EDCI (69 mg, 0.36 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water (20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl ester intermediate as a solid (195 mg) that was used with no further purification; MS *m/e* (M-H)⁺ 555.


The crude oxadiazole *tert*-butyl ester intermediate (195 mg) was dissolved in TFA (2 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo* to leave a brown solid, which was purified by recrystallisation from a mixture of DMSO:CH₃CN:H₂O (7:2:1) and

237

then washed with AcOH to give the title compound (60 mg, 40%) as a solid; ^1H NMR δ 1.3-1.5 (4H, m), 1.8-1.9 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 3.35 (3H, s), 7.0 (1H, d), 7.15 (2H, d), 7.43 (1H, dd), 7.50 (1H, d), 7.58 (1H, s), 7.65 (2H, d), 10.9 (1H, s), 11.2 (1H, s), 12.0 (1H, s); MS m/e MH^+ 501.

5

Example 454: *trans*-4-[4-([5-([4-(Methylsulfonyl)oxy]phenyl)amino)-1,3,4-oxadiazol-2-yl]carbonyl]-amino)phenyl]cyclohexanecarboxylic acid




10 Thiocarbonyl diimidazole (64 mg, 0.36 mmol) was added in one portion to a stirred solution of 4-aminophenyl methanesulfonate (67 mg, 0.36 mmol) in DMA (2 mL) and the mixture was stirred at room temperature under an argon atmosphere for 2 h. *tert*-Butyl *trans*-4-[4-([hydrazino(oxo)-acetyl]amino)phenyl]cyclohexanecarboxylate (Intermediate 79, 108 mg, 0.30 mmol) was added and the mixture was stirred at 80°C for 10 minutes in a microwave. EDCI (69 mg, 0.36 mmol) was then added in one portion and the reaction mixture was heated at 90°C for 10 minutes in a microwave. The reaction mixture was poured onto water (20 mL), stirred for 10 minutes and then filtered to leave the crude oxadiazole *tert*-butyl ester intermediate as a brown solid (200 mg) that was used with no further purification; MS m/e (M-H) $^-$ 555.

20 The crude oxadiazole *tert*-butyl ester intermediate (200 mg) was dissolved in TFA (2 mL) and stirred for 2 h. The reaction mixture was concentrated *in vacuo* to leave a brown solid, which was purified by recrystallisation from a mixture of DMSO:CH₃CN:H₂O (7:2:1) and then washed with AcOH to give the title compound (80 mg, 53%) as a solid; ^1H NMR δ 1.3-1.5 (4H, m), 1.7-1.85 (2H, m), 1.94-2.1 (2H, m), 2.2-2.32 (1H, m), 2.43-2.6 (1H, m), 3.25 (3H, s), 7.18 (1H, d), 7.24 (2H, d), 7.6-7.75 (3H, m), 10.9 (1H, s), 11.2 (1H, s), 12.0 (1H, s); MS m/e MH^+ 501.


25

Example 455: Ethyl *trans*-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]-amino}phenoxy}cyclohexanecarboxylate



A solution of 2,4,5-trifluorophenyl isothiocyanate (70 mg, 0.37 mmol) in DMA (1 mL) was added in one portion to a stirred solution of ethyl *trans*-4-(4-
5 {[(hydrazino(oxo)acetyl)amino]-phenoxy}cyclohexanecarboxylate (Intermediate 80, 129 mg, 0.37 mmol) in DMA (3 mL) and the reaction mixture was stirred at room temperature for 1 h. EDCI (71 mg, 0.37 mmol) was added in one portion and the reaction mixture was heated to 85°C for 20 minutes in a microwave. The mixture was cooled to room
10 temperature and a further portion of EDCI (7.1 mg, 0.04 mmol) was added and the reaction mixture stirred at 85°C for 10 minutes in a microwave. The reaction mixture was concentrated *in vacuo* to leave a solid, which was recrystallised (CH₃CN) to give the title compound (125 mg, 69%) as a solid; ¹H NMR δ 1.33 (3H, t), 1.48-1.58 (2H, m), 1.68-1.72 (2H, m), 2.06-2.10 (2H, m), 2.18-2.22 (2H, m), 2.45-2.53 (1H, m), 4.21 (2H, q), 4.39-4.46
15 (1H, m), 7.07-7.11 (2H, d), 7.81-7.84 (2H, d), 7.80-7.91 (1H, m), 8.29-8.36 (1H, m), 11.12-11.25 (1H, m); MS *m/e* MH⁺ 505.

Example 456: *trans*-4-{4-[(5-[(2,4,5-Trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-phenoxy}cyclohexanecarboxylic acid




20 Lithium hydroxide monohydrate (104 mg, 2.48 mmol) was added in one portion to a solution of methyl *trans*-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]-amino}phenoxy}cyclohexanecarboxylate (Example 449, 125 mg, 0.25 mmol) in a mixture of THF (2 mL), H₂O (2 mL) and MeOH (4 mL) and the reaction mixture was stirred at 35°C for 1 h. The reaction mixture was concentrated *in vacuo* and citric acid (15
25 mL) was added. The resulting mixture was filtered, washed with H₂O and the white solid

was then triturated with hot MeOH and filtered to give the title compound (93 mg, 79%) as a solid. The compound may be recrystallised from a 4:6:1 mixture of EtOH:MeOH:H₂O; melting point 263-265°C.

¹H NMR δ 1.34-1.44 (2H, m), 1.46-1.56 (2H, m), 1.93-1.97 (2H, m), 2.05-2.09 (2H, m),
 5 2.23-2.30 (1H, m), 4.24-4.31 (1H, m), 6.94-6.96 (2H, d), 7.67-7.69 (2H, d), 7.71 (1H, m),
 8.13-8.20 (1H, m), 10.90 (1H, s), 11.03 (1H, s), 12.06 (1H, s); MS m/e (M-H)⁺ 475.

Examples 457-463

The following Examples were prepared by the general procedure of Example 105 except
 10 that DMA was used as solvent in place of DMF, and using the appropriate commercially
 available arylisothiocyanates, R-NCS.




Example	Name	¹ H NMR δ	MS m/e MH ⁺
457	Methyl (<i>trans</i> -4-{4-[(5-[(3-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)acetate	1.01-1.23 (m, 2H), 1.36-1.56 (m, 2H), 1.66-1.87 (m, 5H), 2.25 (d, 2H), 2.37-2.46 (m, 1H), 3.60 (s, 3H), 6.89 (t, 1H), 7.21 (d, 2H), 7.36 (d, 1H), 7.43 (q, 1H), 7.51-7.57 (m, 1H), 7.70 (d, 2H), 11.01 (s, 1H), 11.28 (s, 1H)	453
458	Methyl (<i>trans</i> -4-{4-[(5-[(4-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)acetate	1.01-1.22 (m, 2H), 1.35-1.56 (m, 2H), 1.66-1.88 (m, 5H), 2.25 (d, 2H), 2.39-2.47 (m, 1H), 3.60 (s, 3H), 7.17-7.31 (m, 4H), 7.56-7.66 (m, 2H), 7.66-7.75 (m, 2H), 10.98 (s, 1H), 11.04 (s, 1H)	453
459	Methyl (<i>trans</i> -4-{4-[(5-[(2,4-difluorophenyl)amino]-	1.00-1.23 (m, 2H), 1.34-1.59 (m, 2H), 1.66-1.89 (m, 5H), 2.25 (d, 2H), 2.37-2.46 (m, 1H), 3.60 (s,	471

	1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl }cyclohexyl)acetate	3H), 7.11-7.31 (m, 3H), 7.30-7.49 (m, 1H), 7.68 (d, 2H), 7.95-8.13 (m, 1H), 10.77 (s, 1H), 10.98 (s, 1H)	
460	Methyl (<i>trans</i> -4-{4-[(5-[(2,6-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl }cyclohexyl)acetate	1.01-1.21 (m, 2H), 1.36-1.56 (m, 2H), 1.69-1.86 (m, 5H), 2.25 (d, 2H), 2.38-2.47 (m, 1H), 3.60 (s, 3H), 7.21 (d, 2H), 7.28 (t, 2H), 7.36-7.50 (m, 1H), 7.66 (d, 2H), 10.55 (s, 1H), 10.93 (s, 1H)	471
461	Methyl (<i>trans</i> -4-{4-[(5-[(2,5-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl }cyclohexyl)acetate	1.01-1.24 (m, 2H), 1.37-1.54 (m, 2H), 1.67-1.87 (m, 5H), 2.25 (d, 2H), 2.39-2.48 (m, 1H), 3.60 (s, 3H), 6.92-7.04 (m, 1H), 7.23 (d, 2H), 7.32-7.43 (m, 1H), 7.70 (d, 2H), 7.95-8.10 (m, 1H), 11.03 (s, 1H), 11.12 (s, 1H)	471
462	Methyl (<i>trans</i> -4-{4-[(5-[(2,3-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl }cyclohexyl)acetate	1.01-1.22 (m, 2H), 1.37-1.56 (m, 2H), 1.66-1.88 (m, 5H), 2.25 (d, 2H), 2.37-2.48 (m, 1H), 3.60 (s, 3H), 7.09-7.40 (m, 4H), 7.69 (d, 2H), 7.90 (s, 1H), 11.01 (s, 1H), 11.10 (s, 1H)	471
463	Methyl (<i>trans</i> -4-{4-[(5-[(2,4,6-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl }cyclohexyl)acetate	1.03-1.19 (m, 2H), 1.37-1.54 (m, 2H), 1.66-1.88 (m, 5H), 2.25 (d, 2H), 2.38-2.48 (m, 1H), 3.60 (s, 3H), 7.21 (d, 2H), 7.43 (t, 2H), 7.60-7.72 (m, 2H), 10.50 (s, 1H), 10.93 (s, 1H)	489

234
255


Example 464: Methyl (*trans*-4-{4-[(5-{[3,4-dimethoxyphenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate



Di-2-pyridyl thionocarbonate (233 mg, 1.00 mmol) was added to 3,4-dimethoxyaniline
 5 (154 mg, 1.00 mmol) in DCM (3 mL) and the mixture was stirred for 1 h. Methyl [*trans*-4-
 (4-{[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl)acetate (Intermediate 43) (267mg;
 0.80mmol) was added, followed by DMA. The resultant solution was heated at 85°C for 30
 minutes (DCM was allowed to boil off) then EDAC (192 mg, 1.00 mmol) was added. After
 a further 30 minutes the mixture was allowed to cool with stirring and was then diluted
 10 with water (7 mL). Stirring was continued and the resultant solid was collected by
 filtration, washed with water and dried at 60°C under vacuum to give the title compound
 (288 mg, 73%); ¹H NMR δ 1.04-1.20 (m, 2H), 1.38-1.54 (m, 2H), 1.69-1.89 (m, 5H), 2.25
 (d, 2H), 2.38-2.48 (m, 1H), 3.61 (s, 3H), 3.66-3.73 (m, 1H), 3.74 (s, 3H), 3.78 (s, 3H), 6.99
 (d, 1H), 7.11-7.17 (m, 1H), 7.19-7.28 (m, 3H), 7.65-7.74 (m, 2H), 10.72 (s, 1H), 10.87 (s,
 15 1H); MS m/e MH⁺ 495.

Examples 465-467

The following Examples were prepared by the general procedure of Example 464 using the appropriate aniline as starting material in place of 3,4-dimethoxyaniline.



Example	Name	¹ H NMR δ	MS m/e MH ⁺
465 ¹	Methyl (<i>trans</i> -4-{4-[(5-{[3-[(1R)-2-methoxy-1-methylethyl]oxy}phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate	1.05-1.20 (m, 2H), 1.25 (d, 3H), 1.38-1.54 (m, 2H), 1.70-1.87 (m, 5H), 2.25 (d, 2H), 2.37-2.47 (m, 1H), 3.22-3.33 (m, 3H), 3.42-3.55 (m, 2H), 3.61 (s, 3H), 4.51-4.68 (m, 1H), 6.67 (d, 1H),	523

	l)cyclohexyl)acetate	7.11 (d, 1H), 7.17-7.34 (m, 4H), 7.63-7.77 (m, 2H), 10.93 (s, 2H)	
466 ²	Methyl (<i>trans</i> -4-{4-[(5-[(3-[(1 <i>S</i>)-2-methoxy-1-methylethyl]oxy}phenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl)cyclohexyl)acetate	1.05-1.20 (m, 2H), 1.25 (d, 3H), 1.38-1.54 (m, 2H), 1.70-1.87 (m, 5H), 2.25 (d, 2H), 2.37-2.47 (m, 1H), 3.22-3.33 (m, 3H), 3.42-3.55 (m, 2H), 3.61 (s, 3H), 4.51-4.68 (m, 1H), 6.67 (d, 1H), 7.11 (d, 1H), 7.17-7.34 (m, 4H), 7.63-7.77 (m, 2H), 10.93 (s, 2H)	523
467	Methyl (<i>trans</i> -4-{4-[(5-[(4-fluoro-3-methoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl)cyclohexyl)acetate	1.04-1.22 (m, 2H), 1.36-1.55 (m, 2H), 1.70-1.87 (m, 5H), 2.25 (d, 2H), 2.38-2.48 (m, 1H), 3.60 (s, 3H), 3.85 (s, 3H), 7.12-7.20 (m, 1H), 7.20-7.31 (m, 3H), 7.41 (d, 1H), 7.69 (d, 2H), 10.96 (s, 1H), 11.01 (s, 1H)	483

¹Preparation of the aniline starting material: Intermediate 81²Preparation of the aniline starting material: Intermediate 82**Examples 468-478**

- 5 The following Examples were prepared by the general procedure of Example 357, using the appropriate ester selected from Examples 457-467 as starting material, except that the crude product was recrystallised from acetic acid.


Example	Name	¹ H NMR δ	MS m/e MH ⁺
468	(<i>trans</i> -4-{4-[(5-[(3-Fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl)cyclohexyl)acetic acid	1.12 (q, 2H), 1.46 (q, 2H), 1.67-1.88 (m, 5H), 2.15 (d, 2H), 2.44 (t, 1H), 6.82-6.98 (m, 1H), 7.23 (d, 2H), 7.31-7.39 (m, 1H), 7.39-7.50 (m, 1H), 7.49-7.59 (m, 1H), 7.70 (d, 2H), 11.01 (s, 1H), 11.28 (s, 1H), 12.03 (s, 1H)	453
469	(<i>trans</i> -4-{4-[(5-[(4-	1.11 (q, 2H), 1.45 (q, 2H), 1.64-	439

	Fluorophenyl)amino]- 1,3,4-oxadiazol-2- yl}carbonyl)amino]phe nyl}cyclohexyl)acetic acid	1.88 (m, 5H), 2.15 (d, 2H), 2.44 (t, 1H), 7.17-7.31 (m, 4H), 7.58-7.66 (m, 2H), 7.66-7.77 (m, 2H), 10.98 (s, 1H), 11.04 (s, 1H), 12.03 (s, 1H)	
470	(<i>trans</i> -4-{4-[(5-[(2,4- Difluorophenyl)amino]- 1,3,4-oxadiazol-2- yl}carbonyl)amino]phe nyl}cyclohexyl)acetic acid	1.11 (q, 2H), 1.45 (q, 2H), 1.66- 1.89 (m, 5H), 2.15 (d, 2H), 2.37- 2.45 (m, 1H), 7.10-7.29 (m, 3H), 7.36-7.48 (m, 1H), 7.69 (d, 2H), 7.95-8.11 (m, 1H), 10.78 (s, 1H), 10.98 (s, 1H), 12.03 (s, 1H)	457
471	(<i>trans</i> -4-{4-[(5-[(2,6- Difluorophenyl)amino]- 1,3,4-oxadiazol-2- yl}carbonyl)amino]phe nyl}cyclohexyl)acetic acid	1.11 (q, 2H), 1.45 (q, 2H), 1.64- 1.88 (m, 5H), 2.14 (d, 2H), 2.36- 2.48 (m, 1H), 7.22 (d, 2H), 7.28 (t, 2H), 7.37-7.49 (m, 1H), 7.67 (d, 2H), 10.56 (s, 1H), 10.93 (s, 1H), 12.03 (s, 1H)	457
472	(<i>trans</i> -4-{4-[(5-[(2,5- Difluorophenyl)amino]- 1,3,4-oxadiazol-2- yl}carbonyl)amino]phe nyl}cyclohexyl)acetic acid	1.12 (q, 2H), 1.46 (q, 2H), 1.66- 1.89 (m, 5H), 2.15 (d, 2H), 2.44 (t, 1H), 6.92-7.03 (m, 1H), 7.23 (d, 2H), 7.32-7.45 (m, 1H), 7.70 (d, 2H), 7.95-8.09 (m, 1H), 11.03 (s, 1H), 11.14 (s, 1H), 12.03 (s, 1H)	457
473	(<i>trans</i> -4-{4-[(5-[(2,3- Difluorophenyl)amino]- 1,3,4-oxadiazol-2- yl}carbonyl)amino]phe nyl}cyclohexyl)acetic acid	1.11 (q, 2H), 1.46 (q, 2H), 1.66- 1.87 (m, 5H), 2.15 (d, 2H), 2.38- 2.47 (m, 1H), 7.12-7.35 (m, 4H), 7.69 (d, 2H), 7.92 (t, 1H), 11.01 (s, 1H), 11.11 (s, 1H), 12.04 (s, 1H)	457
474	(<i>trans</i> -4-{4-[(5- [(2,4,6-	1.02-1.21 (m, 2H), 1.37-1.54 (m, 2H), 1.64-1.89 (m, 5H), 2.14 (d,	475

	Trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetate	2H), 2.37-2.49 (m, 1H), 7.22 (d, 2H), 7.37-7.50 (m, 2H), 7.66 (d, 2H), 10.51 (s, 1H), 10.93 (s, 1H), 12.04 (s, 1H)	
475	(<i>trans</i> -4-{4-[(5-{(3,4-Dimethoxyphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.04-1.19 (m, 2H), 1.38-1.54 (m, 2H), 1.66-1.90 (m, 5H), 2.15 (d, 2H), 2.38-2.48 (m, 1H), 3.74 (s, 3H), 3.78 (s, 3H), 6.99 (d, 1H), 7.11-7.18 (m, 1H), 7.20-7.27 (m, 3H), 7.65-7.74 (m, 2H), 10.72 (s, 1H), 10.87 (s, 1H), 11.98 (s, 1H)	481
476	(<i>trans</i> -4-{4-[(5-{(3-{[(1R)-2-methoxy-1-methylethyl]oxy}phenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.07-1.26 (m, 2H), 1.30 (d, 3H), 1.42-1.59 (m, 2H), 1.72-1.97 (m, 5H), 2.21 (d, 2H), 2.44-2.54 (m, 1H), 3.36 (s, 3H), 3.46-3.63 (m, 2H), 4.58-4.71 (m, 1H), 6.72 (d, 1H), 7.12-7.20 (m, 1H), 7.23-7.39 (m, 4H), 7.75 (d, 2H), 11.04 (s, 2H), 12.10 (s, 1H)	509
477	(<i>trans</i> -4-{4-[(5-{(3-{[(1S)-2-methoxy-1-methylethyl]oxy}phenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.07-1.26 (m, 2H), 1.30 (d, 3H), 1.42-1.59 (m, 2H), 1.72-1.97 (m, 5H), 2.21 (d, 2H), 2.44-2.54 (m, 1H), 3.36 (s, 3H), 3.46-3.63 (m, 2H), 4.58-4.71 (m, 1H), 6.72 (d, 1H), 7.12-7.20 (m, 1H), 7.23-7.39 (m, 4H), 7.75 (d, 2H), 11.04 (s, 2H), 12.10 (s, 1H)	509
478	(<i>trans</i> -4-{4-[(5-{(4-fluoro-3-methoxyphenyl)amino]-1,3,4-oxadiazol-2-	1.03-1.21 (m, 2H), 1.38-1.55 (m, 2H), 1.64-1.88 (m, 5H), 2.15 (d, 2H), 2.38-2.49 (m, 1H), 3.85 (s, 3H), 7.13-7.19 (m, 1H), 7.20-7.29	469

	yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	(m, 3H), 7.38-7.45 (m, 1H), 7.66-7.73 (m, 2H), 10.96 (s, 1H), 11.01 (s, 1H), 12.04 (s, 1H)	
--	---	--	--


Example 479: 1-{5-[(5-[(2-Fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidine-4-carboxylic acid hydrochloride



- i) Following the procedure of Example 319 except that Intermediate 83 was used in place of Intermediate 77 and 2-fluorophenylisothiocyanate in place of 2,4,5-trifluorophenylisothiocyanate, there was obtained (in quantitative yield) *tert*-butyl 1-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidine-4-carboxylate as a solid; ^1H NMR δ 1.41 (s, 9H), 1.44-1.60 (m, 2H), 1.83 (d, 2H), 2.92 (t, 1H), 3.33 (d, 2H), 4.09-4.20 (m, 2H), 6.88 (d, 1H), 7.12-7.22 (m, 1H), 7.27 (q, 1H), 7.33 (d, 1H), 7.86-7.96 (m, 1H), 8.06 (t, 1H), 8.44-8.52 (m, 1H), 10.80 (s, 1H), 10.96 (s, 1H); MS m/e MH^+ 483.
- ii) TFA (0.307 mL; 4.0 mmol) was added to a suspension of *tert*-Butyl 1-{5-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidine-4-carboxylate (483 mg; 1.0 mmol) in DCM (5 mL). After 3 days more TFA was added in aliquots (0.31 mL) over 2 days then HCl (1 mL of a 4M solution in 1,4-dioxan) was added. After 24 h the mixture was filtered and the solid was washed with ether and dried under vacuum at 60°C to give the title compound (193 mg, 42%) as a solid; ^1H NMR δ 1.55-1.74 (m, 2H), 1.88-2.06 (m, 2H), 2.56-2.69 (m, 1H), 2.69-2.81 (m, 1H), 3.18-3.35 (m, 2H), 4.17 (d, 2H), 7.13-7.22 (m, 1H), 7.24-7.31 (m, 1H), 7.31-7.37 (m, 1H), 7.44 (d, 1H), 7.98-8.09 (m, 1H), 8.19-8.28 (m, 1H), 8.51-8.59 (m, 1H), 10.87 (s, 1H), 11.37 (s, 1H); MS m/e MH^+ 427.


Example 480-482

- The following Examples were prepared by the general procedure of Example 479, using the appropriate isothiocyanate in place of 2-fluorophenylisothiocyanate



Example	Name	¹ H NMR δ	MS m/e MH ⁺
480	1-{5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid hydrochloride	1.54-1.69 (m, 2H), 1.87-2.05 (m, 2H), 2.41-2.50 (m, 1H), 2.53-2.66 (m, 1H), 3.10-3.30 (m, 2H), 4.15 (d, 2H), 7.18-7.41 (m, 1H), 7.66-7.81 (m, 1H), 8.08-8.25 (m, 2H), 8.51 (s, 1H), 11.15 (s, 1H), 11.30 (s, 1H)	463
481	1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid hydrochloride	1.55-1.70 (m, 2H), 1.87-2.02 (m, 2H), 2.57-2.71 (m, 1H), 3.22 (t, 2H), 4.16 (d, 2H), 7.23-7.43 (m, 3H), 7.50 (q, 1H), 7.65-7.78 (m, 1H), 8.17 (d, 1H), 8.53 (d, 1H), 11.32 (s, 1H), 11.37 (s, 1H)	445
482	1-{5-[(5-[(4-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid hydrochloride	1.50-1.71 (m, 2H), 1.86-2.02 (m, 2H), 2.54-2.67 (m, 1H), 3.07-3.30 (m, 2H), 4.14 (d, 2H), 7.19-7.40 (m, 3H), 7.56-7.70 (m, 2H), 8.15 (d, 1H), 8.51 (s, 1H), 11.10 (s, 1H), 11.26 (s, 1H)	427


Example 483: Methyl 4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl}butanoate













- Methyl 4-(4-aminophenyl)butanoate (Journal of Medicinal Chemistry (1990), 33(11), 3014-19) (193 mg, 1 mmol) was stirred in dry DCM (9 mL) and PS-di-
isopropylethylamine (750 mg, 3 mmol) was added. Methyl chloro(oxo)acetate (101 μ L, 1.1 mmol) was added to the reaction mixture which was then stirred at room temperature for 2
5 h. The polymer supported base (HCl salt) was filtered off and washed with DCM (3 mL). Hydrazine monohydrate (53 μ L, 1.1 mmol) was added to the stirred filtrate and a white precipitate formed after 2 h. The reaction mixture was stirred at room temperature overnight then DMF (5 mL) was added to dissolve the solid. 2-Fluorophenylisothiocyanate (0.135 mL, 1.1 mmol) was added via a pipette and the mixture stirred at room temperature
10 for 2.5 h before adding EDAC (230 mg, 1.2 mmol). After stirring at room temperature for a further 2.5 h the mixture was heated to 50°C and the DCM was allowed to evaporate. After 2h the reaction had gone to completion and the mixture was concentrated in vacuo. The orange solid obtained was crystallised from water and then filtered. The solid was then washed with Et₂O, EtOAc and then MeCN (1 mL) to give the title compound (174 mg,
15 44%) as a solid; ¹H NMR δ 1.79-1.87 (2H, m), 2.31 (2H, t), 2.58 (2H, t), 3.59 (3H, s), 7.15-7.20 (3H, m), 7.25-7.34 (2H, m), 7.71 (2H, d), 8.02-8.07 (1H, m), 10.76 (1H, s), 10.95 (1H, s); MS m/e MH⁺ 399.









Examples 484-497



- The following examples were prepared by the general procedure of Example 483, using the
20 appropriate amine R¹NH₂ and isothiocyanate R²-NCS.



Ex	SM	R ¹	R ²	¹ H NMR δ	MS m/e MH ⁺
484 ¹	Lit. ²			3.63 (3H, s), 3.86 (2H, s), 7.16-7.20 (1H, m), 7.26-7.28 (1H, m), 7.29-7.34 (1H, m), 7.37-7.41 (2H, m), 7.76-7.80 (2H, m), 8.02-8.06 (1H, m), 10.78 (1H, s), 11.09 (1H, s)	403
485	Int 84			1.70 (2H, s), 1.85 (9H, s), 1.93 (2H, s), 2.18 (2H, s), 3.61 (3H, s), 7.16-7.19 (1H, m), 7.25-7.38 (4H, m), 7.71-7.74 (2H, m), 8.02-8.07 (1H, m), 10.76 (1H, s), 10.95 (1H, s)	491
486	Int 85			1.56-1.66 (1H, m), 1.74-1.89 (2H, m), 1.94-2.03 (1H, m), 2.07-2.11 (1H, m), 2.23-2.33 (1H, m), 3.01-3.14 (2H, m), 3.63 (3H, d), 7.14-7.19 (1H, m), 7.25-7.34 (3H, m), 7.71 (2H, d), 8.02-8.07 (1H, m), 10.75 (1H, s), 10.95 (1H, s)	425
487	Lit. ³			3.89 (3H, s), 7.17-7.20 (1H, m), 7.26-7.35 (2H, m), 7.79 (2H, d), 7.84-7.86 (2H, m), 7.96 (2H, d), 8.05 (3H, t), 10.80 (1H, s), 11.19 (1H, s)	433

488	Int 86			1.21 (3H, t), 3.75 (2H, s), 4.11 (2H, q), 7.18 (1H, s), 7.25-7.33 (3H, m), 7.41 (1H, d), 7.58 (2H, s), 7.68 (2H, d), 7.93 (2H, d), 8.08 (1H, s), 10.85 (1H, s), 11.20 (1H, s)	461
489	Int 87			0.82-1.30 (m, 6H), 1.34-1.85 (m, 12H), 2.14-2.23 (m, 2H), 3.58 (s, 3H), 3.93 (s, 1H), 4.38 (s, 1H), 7.11-7.18 (m, 1H), 7.21-7.34 (m, 2H), 8.03 (t, 1H), 8.72 (d, 1H), 8.96 (d, 1H)	459
490	Int 88				457
491	Int 89				485
492	Int 90			1.19 (t, 3H), 1.22-2.44 (m, 9H), 2.97-3.17 (m, 1H), 4.06 (q, 2H), 7.20-7.27 (m, 2H), 7.32-7.39 (m, 1H), 7.45-7.55 (m, 1H), 7.66-7.76 (m, 3H), 11.02 (s, 1H), 11.02 (s, 1H)	471

493	Int 91			1.19 (t, 3H), 1.22-2.44 (m, 9H), 2.97-3.17 (m, 1H), 4.06 (q, 2H), 7.20-7.27 (m, 2H), 7.32-7.39 (m, 1H), 7.45-7.55 (m, 1H), 7.66-7.76 (m, 3H), 11.02 (s, 1H), 11.02 (s, 1H)	471
494	Lit. ⁴			2.04 (2H, t), 3.46 (2H, q), 3.84 (3H, s), 4.14 (2H, t), 7.06-7.08 (2H, m), 7.36 (1H, t), 7.47-7.54 (1H, m), 7.68-7.73 (1H, m), 7.92-7.96 (2H, m), 9.26 (1H, t), 11.21 (1H, s)	433
495	Int 92			1.97-2.04 (2H, m), 3.44 (2H, q), 3.84 (3H, s), 4.09 (2H, t), 7.21-7.24 (1H, m), 7.31-7.34 (1H, m), 7.44 (1H, s), 7.42-7.49 (2H, m), 7.53-7.56 (1H, m), 7.65-7.70 (1H, m), 9.23 (1H, t), 11.18 (1H, s)	433
496	Lit. ⁵			2.19-2.27 (1H, m), 2.34-2.41 (1H, m), 2.54-2.60 (2H, m), 3.13-3.22 (1H, m), 3.42 (1H, m), 3.59 (1.5H, s), 3.67 (1.5H, s), 7.22-7.32 (2H, m), 7.36 (1H, d), 7.48 (1H, d), 7.67-7.75 (3H, m), 10.98 (1H, s), 11.23 (1H, s)	429

497	Int 94			1.10-1.23 (3H, m), 1.83-2.03 (2H, m), 2.18-2.24 (2H, m), 2.56 (1H, m), 2.63 (1H, d), 2.72 (1H, d), 2.90-3.16 (1H, m), 4.03-4.07 (2H, m), 7.17 (2H, t), 7.35 (1H, d), 7.45-7.52 (1H, m), 7.64-7.73 (3H, m), 10.95 (1H, s), 11.22 (1H, s)	457
-----	--------	---	---	---	-----

¹Purified by preparative HPLC eluting with a gradient of water-MeCN containing formic acid modifier


²Aniline prepared according to J. Pharm. Sci., 1974, 63(8), 1333

³Aniline prepared according to J. Combinatorial Chem., 2003, 5(3), 205

5 ⁴Amine prepared according to Helv. Chim. Acta, 1983, 66(2), 489

⁵Aniline prepared according to German Patent application DE 2626287 (1976)

Example 498: 4-((5-((2-Fluorophenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl]butanoic acid



10

Methyl 4-((5-((2-fluorophenyl)amino)-1,3,4-oxadiazol-2-

yl)carbonyl)amino]phenyl]butanoate (Example 483) (174 mg, 0.44mmol) was stirred in

THF: MeOH (8 mL of a 1:1 mixture) and NaOH (2.5 mL of 2M aqueous solution) was

added. The mixture was stirred for 8.5 h then acidified (2M HCl), allowed to stand

15 overnight, then concentrated *in vacuo*. The precipitate was filtered and washed with water


and then Et₂O to give the title compound (94 mg, 56%) as a solid; ¹H NMR δ 1.77-1.84

(2H, m), 2.22 (2H, t), 2.59 (2H, t), 7.14-7.21 (3H, m), 7.25-7.34 (2H, m), 7.70-7.72 (2H,









m), 8.02-8.07 (1H, m), 10.74 (1H, s), 10.94 (1H, s), 11.98 (1H, s); MS m/e MH⁺ 385.









20 **Examples 499-514**









The following examples were prepared by the general procedure of Example 498, using the appropriate ester as starting material (SM).



Example	SM	R ¹	R ²	¹ H NMR δ	MS m/e MH ⁺
499	Ex 485			1.70 (2H, s), 1.83 (8H, d), 1.91 (2H, s), 2.17 (2H, s), 7.16-7.20 (1H, m), 7.26-7.38 (4H, m), 7.71-7.74 (2H, m), 8.02-8.06 (1H, m), 10.74 (1H, s), 10.94 (1H, s), 12.04 (1H, s)	477
500	Ex 486			1.28 (0H, s), 1.52-2.38 (5H, q), 2.85-3.11 (2H, m), 3.59-3.64 (1H, m), 7.14-7.22 (1H, m), 7.25-7.34 (4H, m), 7.70 (2H, d), 8.04 (1H, t), 10.74 (1H, s), 10.94 (1H, s), 12.02 (1H, s)	411
501	Ex 487			7.17-7.21 (1H, m), 7.26-7.35 (2H, m), 7.77-7.83 (4H, m), 7.95 (2H, d), 8.01-8.06 (3H, m), 10.78 (1H, s), 11.16 (1H, s)	419
502	Int 94			3.59-3.63 (2H, s), 7.15-7.20 (1H, m), 7.26-7.36 (4H, m), 7.62-7.64 (2H, m), 7.67-7.70 (2H, m), 7.90-7.92 (2H, m), 8.03-8.08 (1H, m), 10.79 (1H, s), 11.13 (1H, s), 12.32 (1H, s)	433


503	Ex 488			3.66 (2H, s), 7.18 (1H, t), 7.24-7.35 (3H, m), 7.41 (1H, t), 7.56 (2H, d), 7.68 (2H, d), 7.92 (2H, d), 8.07 (1H, t), 10.84 (1H, s), 11.18 (1H, s), 12.38 (1H, s)	433
504 ¹	Ex 489			0.81-1.08 (m, 2H), 1.09-1.28 (m, 4H), 1.32-1.47 (m, 2H), 1.48-1.63 (m, 3H), 1.65-1.86 (m, 6H), 2.08 (t, 2H), 2.51-2.55 (m, 1H), 3.60-3.74 (m, 1H), 3.87-3.99 (m, 1H), 7.10-7.18 (m, 1H), 7.22-7.34 (m, 2H), 7.99-8.06 (m, 1H), 8.67-8.97 (m, 1H), 10.68 (s, 1H), 10.68 (s, 1H)	445
505 ²	Ex 490			2.76 (2H, t), 3.07 (2H, t), 3.14-3.19 (2H, m), 3.49-3.50 (2H, m), 4.95 (1H, m), 7.22 (1H, d), 7.32 (2H, m), 7.42-7.45 (2H, m), 7.79-7.84 (1H, m)	429
506 ²	Int 95			1.58 (2H, m), 1.73 (2H, m), 2.57 (2H, m), 3.24 (1H, m), 7.18 (2H, d), 7.34-7.37 (1H, m), 7.48 (1H, d), 7.68-7.73 (3H, m), 10.96 (1H, s), 11.23 (1H, s), 12.60 (2H, s)	461

507 ¹³	Int 96			(d ₄ MeOH) 1.41-1.62 (m, 4H), 1.71-2.05 (m, 4H), 2.14-2.21 (m, 1H), 2.42-2.54 (m, 1H), 3.46 (s, 1H), 3.57 (s, 1H), 6.97-7.28 (m, 6H), 7.92-8.02 (m, 2H)	439
508	Ex 491			(d ₄ MeOH) 0.87-1.91 (m, 8H), 2.13 (d, 1H), 2.24-2.32 (m, 0.5H), 2.38 (d, 1H), 2.45-2.54 (m, 1H), 2.67-2.78 (m, 0.5H), 7.09-7.24 (m, 4H), 7.48-7.62 (m, 3H)	457
509	Ex 492			δ 1.24-1.43 (2H, m), 1.56-1.82 (2H, m), 1.98-2.06 (2H, m), 2.17 (0.5H, m), 2.30-2.35 (2H, m), 2.44-2.45 (0.5H, m), 3.03-3.14 (1H, m), 7.23-7.26 (2H, m), 7.35-7.37 (1H, m), 7.46-7.52 (1H, m), 7.68-7.73 (3H, m), 10.97 (1H, s), 11.24 (1H, s), 11.98 (1H, s)	443
510	Ex 493			1.24-1.43 (2H, m), 1.56-1.82 (2H, m), 1.98-2.06 (2H, m), 2.17 (0.5H, m), 2.30-2.35 (2H, m), 2.44-2.45 (0.5H, m), 3.03-3.14 (1H, m), 7.23-7.26 (2H, m), 7.35-7.37 (1H, m), 7.46-7.52 (1H, m), 7.68-7.73 (3H, m), 10.97 (1H, s), 11.24 (1H, s), 11.98 (1H, s)	443

511	Ex 494			1.99-2.05 (2H, m), 3.40-3.46 (2H, m), 4.12 (2H, t), 7.00-7.03 (2H, m), 7.32-7.36 (1H, m), 7.42-7.47 (1H, m), 7.65-7.71 (1H, m), 7.87-7.91 (2H, m), 9.16 (1H, t), 11.15 (1H, s)	419
512	Ex 495			1.98-2.04 (2H, m), 3.44 (2H, q), 4.09 (2H, t), 7.17-7.20 (1H, m), 7.32-7.35 (1H, m), 7.39-7.47 (3H, m), 7.52-7.54 (1H, m), 7.65-7.70 (1H, m), 9.16 (1H, t), 11.13 (1H, s)	419
513	Ex 496			2.17-2.25 (1H, m), 2.30-2.38 (1H, m), 2.53-2.58 (2H, m), 3.03-3.09 (1H, m), 3.39-3.44 (0.5H, m), 3.62 (0.5H, m), 7.22-7.37 (3H, m), 7.45-7.52 (1H, m), 7.67-7.75 (3H, m), 10.98 (1H, s), 12.00 (1H, s)	415
514	Ex 497			1.77-1.96 (2H, m), 2.15-2.24 (2H, m), 2.43-2.46 (0.5H, m), 2.54 (0.5H, m), 2.63 (1H, d), 2.70-2.72 (1H, d), 2.87-3.04 (1H, m), 7.17 (2H, t), 7.35-7.37 (1H, m), 7.45-7.52 (1H, m), 7.68-7.73 (3H, m), 10.95 (1H, s), 11.23 (1H, s), 11.96 (1H, s)	429


¹20 equivalents of 2M NaOH was used²10 equivalents of 2M NaOH was used³Purified by preparative HPLC eluting with a gradient of water-MeCN containing formic acid modifier

Example 515: 5-{4-[(5-{(3,4-Difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}propyl)malonic acid




A solution of (3-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}propyl)malonic acid (Example 506) (40 mg, 0.087 mmol) in AcOH (4 mL) was heated under reflux with stirring for 4 h. The reaction mixture was concentrated in vacuo and the solid filtered and washed with water then Et₂O to give the title compound (20 mg, 55%) as a solid; ¹H NMR δ 1.50-1.60 (4H, m), 2.24 (2H, t), 2.55 (2H, t), 7.15-7.20 (2H, m), 7.35-7.37 (1H, m), 7.48-7.52 (1H, m), 7.68-7.73 (3H, m), 10.97 (1H, s); MS m/e MH⁺ 417.

Example 516: N-{4-[trans-4-(2-Amino-2-oxoethyl)cyclohexyl]phenyl}-5-(2-fluorophenyl)-amino]-1,3,4-oxadiazole-2-carboxamide




Ethyl chloroformate (0.056 mL, 0.58 mmol) was added dropwise to a stirred solution of (trans-4-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid (Example 118) (170 mg, 0.39 mmol) and N-methyl morpholine (0.064 mL, 0.58 mmol) in DCM (5 mL) at 0°C. The mixture was stirred for 40 minutes then ammonia (2 mL of a 7M solution in MeOH) was added and stirring was continued for 16 h whilst warming to ambient temperature. Volatile material was removed by evaporation and the residue was triturated with water and washed with 10% ammonium hydroxide solution (10 mL) followed by ether (10 mL) and dried under reduced pressure to give the title compound (121 mg, 71%) as a solid; ¹H NMR δ 0.98-1.15 (m, 2H), 1.33-1.50 (m, 2H), 1.63-1.85 (m, 5H), 1.95 (d, 2H), 2.37-2.44 (m, 1H), 6.75 (s, 1H), 7.10-7.35 (m, 5H), 7.66 (d, 2H), 8.03 (t, 1H), 10.78 (s, 1H), 10.99 (s, 1H); MS m/e MH⁺ 438.

Example 517: N-[4-[trans-4-(Cyanomethyl)cyclohexyl]phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide



Trifluoroacetic anhydride 0.042 mL, 0.05 mmol) was added to a stirred solution of N-[4-
5 [trans-4-(2-amino-2-oxoethyl)cyclohexyl]phenyl]-5-[(2-fluorophenyl)amino]-1,3,4-
oxadiazole-2-carboxamide (Example 516) (20 mg, 0.05 mmol) and pyridine (90 μ L, 0.11
mmol) in DCM (2 mL). After 15 minutes DCM (10 mL) was added and the reaction
mixture was washed sequentially with 1M HCl (5 mL), water (10 mL) and 10%
ammonium hydroxide solution (5 mL) then dried and concentrated under reduced pressure
10 to give the title compound (18mg, 95%) as a solid; ^1H NMR δ 1.13-1.30 (m, 3H), 1.40-
1.54 (m, 2H), 1.60-1.75 (m, 1H), 1.76-1.92 (m, 5H), 2.40-2.50 (m, 1H), 7.10-7.19 (m, 1H),
7.20-7.35 (m, 4H), 7.68 (d, 2H), 8.03 (t, 1H), 10.75 (s, 1H), 10.93 (s, 1H); MS m/e MH^+
420.


15 **Example 518: 5-[(2-Fluorophenyl)amino]-N-[4-(propylsulfonyl)phenyl]-1,3,4-oxadiazole-2-carboxamide**



Following the general procedure of Example 105 but using 2-hydrazino-2-oxo-N-[4-
(propylsulfonyl)phenyl]acetamide (Intermediate 97) as starting material in place of methyl
20 [trans-4-(4-[[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl]acetate and using 2-
fluorophenylisothiocyanate in place of 2-isothiocyanatopyridine the title compound was
obtained as a solid; ^1H NMR δ 0.90 (t, 3H), 1.46-1.62 (m, 2H), 3.18-3.27 (m, 2H), 7.12-
7.20 (m, 1H), 7.22-7.35 (m, 2H), 7.87 (d, 2H), 7.99-8.10 (m, 3H), 10.90 (s, 1H), 11.50 (s,
1H); MS m/e MH^+ 405.

Examples 519-522


The following examples were prepared by the general procedure of Example 127 using Intermediate 21 and the appropriate commercially available aniline as starting materials.



Example	Name	¹ H NMR δ	MS m/e MH ⁺
519	Methyl 4'-{[5-({[4-(4-acetylpiperazin-1-yl)phenyl]amino}carbonyl)-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylate	2.07 (s, 3H), 3.05-3.11 (m, 2H), 3.12-3.19 (m, 2H), 3.54-3.63 (m, 4H), 3.93 (s, 3H), 6.97 (d, 2H), 7.64-7.88 (m, 8H), 8.03 (d, 2H), 10.90 (s, 1H), 11.20 (s, 1H)	541
520	N-[4-(4-acetylpiperazin-1-yl)phenyl]-5-[(4-chloro-2-fluoro-5-hydroxyphenyl)amino]-1,3,4-oxadiazole-2-carboxamide	2.08 (s, 3H), 3.02-3.20 (m, 4H), 3.53-3.62 (m, 4H), 6.97 (d, 2H), 7.39 (d, 1H), 7.65 (d, 2H), 7.84 (d, 1H), 10.35 (s, 1H), 10.80 (s, 1H), 10.84 (s, 1H)	475
521	Ethyl {5-({[5-({[4-(4-acetylpiperazin-1-yl)phenyl]amino}carbonyl)-1,3,4-oxadiazol-2-yl]amino}-2-chloro-4-fluorophenoxy)acetate	1.22 (t, 3H), 2.08 (s, 3H), 3.04-3.19 (m, 4H), 3.53-3.63 (m, 4H), 4.20 (q, 2H), 4.92 (s, 2H), 6.97 (d, 2H), 7.56 (d, 1H), 7.65 (d, 2H), 7.85 (d, 1H), 10.85 (s, 1H), 11.00 (s, 1H)	561


522	Methyl 3'-{[5-([4-(4-acetylpiperazin-1-yl)phenyl]amino)carbonyl]-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylate	2.07 (s, 3H), 3.04-3.20 (m, 4H), 3.53-3.64 (m, 4H), 3.94 (s, 3H), 6.97 (d, 2H), 7.43 (d, 1H), 7.54 (t, 1H), 7.67 (d, 3H), 7.81 (d, 2H), 7.96-8.00 (m, 1H), 8.09 (d, 2H), 10.89 (s, 1H), 11.16 (s, 1H)	541
-----	--	---	-----

Example 523: 3'-{[5-([4-(4-Acetylpiperazin-1-yl)phenyl]amino)carbonyl]-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylic acid




- 5 Following the procedure of Example 357 but using methyl 3'-{[5-([4-(4-acetylpiperazin-1-yl)phenyl]amino)carbonyl]-1,3,4-oxadiazol-2-yl]amino}biphenyl-4-carboxylate as starting material in place of methyl (*trans*-4-{4-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate the title compound was obtained as a solid; ¹H NMR δ 2.09 (s, 3H), 3.01-3.20 (m, 4H), 3.52-3.67 (m, 4H), 6.98 (d, 2H), 7.41 (d, 1H), 7.53 (t, 1H), 7.61-7.71 (m, 3H), 7.78 (d, 2H), 8.00 (s, 1H), 8.07 (d, 2H), 10.87 (s, 1H), 11.18 (s, 1H), 13.37 (s, 1H); MS m/e MH⁺ 527.

Example 524: 2-([4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl)sulfonyl)-2-methylpropanoic acid



- 15 Following the general procedure described in Example 357 except using ethyl 2-([4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl)sulfonyl)-2-methylpropanoate (Intermediate 98) as the ester starting material the title compound was obtained as a solid; ¹H NMR δ 1.50 (s, 6H), 7.12-7.20 (m, 1H), 7.23-7.36 (m, 2H), 7.79 (d, 2H), 7.98-8.12 (m, 3H), 10.90 (s, 1H), 11.56 (s, 1H); MS m/e MH⁺ 449.

Example 525: (trans-4-{4-[(5-{(3-Methylisoxazol-5-yl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid



- Sodium hydride (50 mg of a 60% dispersion in mineral oil, 1.25 mmol) was added to a stirred solution of 5-amino-3-methylisoxazole in THF. After 5 minutes di-2-pyridyl thionocarbonate (290 mg, 1.25 mmol) was added. Stirring was continued for a further 10 minutes then methyl [trans-4-(4-{[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl]acetate (Intermediate 43) (333 mg, 1.00 mmol) was added followed by DMF (5 mL). The mixture heated to 85°C for 15 minutes then EDAC (240 mg, 1.25 mmol) was added.
- 10 After a further 1 h at 85°C volatile material was removed by evaporation and the residue was redissolved in MeOH (5 mL), THF (2.5 mL) and NaOH (3 mL of a 2M aqueous solution) and stirred for 16h. The reaction mixture was concentrated under reduced pressure then redissolved in water (10 mL) and acidified with HCl (2M aqueous solution) to ca pH 5. The resulting precipitate was centrifuged, washed with water, dried under
- 15 reduced pressure and triturated with ether to give the title compound (83 mg, 20%) as a solid; ¹H NMR δ 1.02-1.20 (m, 2H), 1.36-1.54 (m, 2H), 1.65-1.88 (m, 5H), 2.13 (d, 2H), 2.24 (s, 3H), 2.39-2.48 (m, 1H), 6.10 (s, 1H), 7.24 (d, 2H), 7.69 (d, 2H), 11.00 (s, 1H), 12.11 (s, 1H); MS m/e MH⁺ 426.

Examples 526-539


- 20 The following examples were prepared by the general procedure of Example 525 using the appropriate commercially available amino-heterocycle in place of 5-amino-3-methylisoxazole.

Example	Name	¹ H NMR δ	MS m/e MH ⁺
526	{ <i>trans</i> -4-[4-({[5-(1,3,4-thiadiazol-2-ylamino)-1,3,4-oxadiazol-2-yl]carbonyl}amino)phenyl]cyclohexyl}acetic acid	1.03-1.19 (m, 2H), 1.34-1.54 (m, 2H), 1.65-1.91 (m, 5H), 2.13 (d, 2H), 2.38-2.48 (m, 1H), 7.20 (d, 2H), 7.68 (d, 2H), 8.90 (s, 1H), 10.86 (s, 1H)	428
527	(<i>trans</i> -4-{4-[(5-[(2,5-dimethylpyridin-4-yl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.21 (m, 2H), 1.36-1.53 (m, 2H), 1.67-1.89 (m, 5H), 2.14 (d, 2H), 2.22 (s, 3H), 2.41-2.48 (m, 4H), 7.21 (d, 2H), 7.70 (d, 2H), 7.95-8.06 (m, 2H)	
528	(<i>trans</i> -4-{4-[(5-[(5,6-dimethylpyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.02-1.20 (m, 2H), 1.37-1.53 (m, 2H), 1.65-1.88 (m, 5H), 2.36-2.47 (m, 7H), 7.20 (d, 2H), 7.66 (d, 2H), 8.91 (s, 1H), 10.94 (s, 1H)	451
529	(<i>trans</i> -4-{4-[(5-[(3-methylpyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)acetic acid	0.99-1.22 (m, 2H), 1.30-1.53 (m, 2H), 1.64-1.89 (m, 5H), 2.07-2.18 (m, 2H), 2.38-2.47 (m, 1H), 2.58 (s, 3H), 6.67 (d, 1H), 6.98 (d, 1H), 7.20 (d, 2H), 7.65 (d, 2H)	437
530	(<i>trans</i> -4-{4-[(5-[(5-bromopyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.02-1.20 (m, 2H), 1.36-1.54 (m, 2H), 1.63-1.90 (m, 5H), 2.14 (d, 2H), 2.38-2.48 (m, 1H), 7.22 (d, 2H), 7.68 (d, 2H), 8.48 (s, 1H), 8.91 (s, 1H), 10.85 (s, 1H)	502

531	(<i>trans</i> -4-{4-[(5-{(2-methoxypyrimidin-5-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	0.93-1.12 (m, 2H), 1.30-1.46 (m, 2H), 1.55-1.81 (m, 5H), 2.00-2.11 (m, 2H), 2.32-2.41 (m, 1H), 3.86 (s, 3H), 7.12 (d, 2H), 7.60 (d, 2H), 8.70 (s, 2H)	453
532	(<i>trans</i> -4-{4-[(5-{(5-benzyl-1,3,4-oxadiazol-2-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	7.67 (d, 2H), 7.10-7.44 (m, 7H), 4.02 (s, 2H), 2.38-2.48 (m, 1H), 2.14 (d, 2H), 1.61-1.93 (m, 5H), 1.34-1.54 (m, 2H), 1.02-1.27 (m, 2H)	503
533	(<i>trans</i> -4-{4-[(5-{(3-methyl-1,2,4-thiadiazol-5-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	0.99-1.20 (m, 2H), 1.35-1.55 (m, 2H), 1.63-1.92 (m, 5H), 2.14 (d, 2H), 2.35-2.47 (m, 4H), 7.22 (d, 2H), 7.67 (d, 2H), 10.87 (s, 1H)	443
534	(<i>trans</i> -4-{4-[(5-{(3-ethyl-1,2,4-thiadiazol-5-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.04-1.19 (m, 2H), 1.23 (t, 3H), 1.37-1.55 (m, 2H), 1.65-1.94 (m, 5H), 2.15 (d, 2H), 2.37-2.49 (m, 1H), 2.67 (q, 2H), 7.21 (d, 2H), 7.68 (d, 2H), 10.72 (s, 1H)	457
535	(<i>trans</i> -4-{4-[(5-{(2,4-dimethylpyrimidin-5-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.20 (m, 2H), 1.36-1.53 (m, 2H), 1.65-1.89 (m, 5H), 2.14 (d, 2H), 2.36-2.53 (m, 7H), 7.24 (d, 2H), 7.56-7.76 (m, 3H), 10.98 (s, 1H), 11.97 (s, 1H)	451
536	(<i>trans</i> -4-{4-[(5-{(1,3-dimethyl-1 <i>H</i> -pyrazol-5-yl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.20 (m, 2H), 1.38-1.53 (m, 2H), 1.67-1.88 (m, 5H), 2.09-2.17 (m, 5H), 2.36-2.48 (m, 1H), 3.68 (s, 3H), 6.14 (s, 1H), 7.22 (d, 2H), 7.67 (d, 2H), 10.91 (s, 1H)	439

537	(<i>trans</i> -4-{4-[(5-[(6-chloropyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid	1.03-1.20 (m, 2H), 1.36-1.54 (m, 2H), 1.65-1.91 (m, 5H), 2.15 (d, 2H), 2.40-2.49 (m, 1H), 7.25 (d, 2H), 7.69 (d, 2H), 8.48 (s, 1H), 9.20 (s, 1H), 11.05 (s, 1H), 11.86 (s, 1H)	457
538	(<i>trans</i> -4-{4-[(5-[(3,5-dimethylpyrazin-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid	1.23-1.41 (m, 2H), 1.56-1.75 (m, 2H), 1.85-2.10 (m, 5H), 2.34 (d, 2H), 2.56-2.74 (m, 7H), 7.43 (d, 2H), 7.87 (d, 2H), 8.35 (s, 1H), 11.12 (s, 1H), 12.14 (s, 1H)	451
539	[<i>trans</i> -4-(4-{[(5-[(4-(trifluoromethyl)-1,3-thiazol-2-yl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid	1.02-1.20 (m, 2H), 1.36-1.55 (m, 2H), 1.66-1.93 (m, 5H), 2.14 (d, 2H), 2.38-2.49 (m, 1H), 7.21 (d, 2H), 7.68 (d, 2H), 7.89 (s, 1H), 10.93 (s, 1H), 11.98 (s, 1H)	496


Example 540: Methyl (*trans*-4-{4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetate



- 5 Following the general procedure of Example 457, except that 3,4-difluorophenyl-isothiocyanate was used in place of 3-fluorophenylisothiocyanate, the title compound was obtained in 90% yield as a solid; ^1H NMR δ 1.04-1.21 (m, 2H), 1.37-1.54 (m, 2H), 1.69-1.87 (m, 5H), 2.25 (d, 2H), 2.38-2.48 (m, 1H), 3.61 (s, 3H), 7.22 (d, 2H), 7.31-7.40 (m, 1H), 7.48 (q, 1H), 7.65-7.76 (m, 3H), 10.96 (s, 1H), 11.24 (s, 1H); MS m/e ($M-H$)⁺ 469.

278

Example 541: (trans-4-{4-[(5-[(3,4-Difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid




Following the procedure of Example 468 except the starting material was methyl (trans-4-
 5 {4-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)
 cyclohexyl)acetate, the title compound was obtained in 56% yield as a solid; ^1H NMR δ
 1.12 (q, 2H), 1.46 (q, 2H), 1.65-1.75 (m, 1H), 1.79 (t, 4H), 2.15 (d, 2H), 2.37-2.48 (m, 1H),
 7.23 (d, 2H), 7.31-7.39 (m, 1H), 7.49 (q, 1H), 7.70 (d, 2H), 7.71-7.75 (m, 1H); MS m/e
 457.

10

Examples 542-546

The following Examples were prepared by the procedure of Example 525 with
 Intermediate 43 and the appropriate commercially available aniline RNH_2 .




Example	R	HPLC Retention Time (minutes) (in LCMS system)	MS m/e MH^+
542		2.92	494
543		2.99	567
544		2.78	483



545		2.85	474
546		2.92	506

Examples 547-551

The following Examples were prepared by the procedure of Example 357 starting with the appropriate ester selected from Examples 542-546 to give the illustrated acid.





Example	R	¹ H NMR δ	MS m/e MH ⁺
547		1.11-1.14 (2H, m), 1.44-1.48 (2H, m), 1.73-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 7.23 (2H, d), 7.70 (3H, d), 7.93 (1H, d), 7.99 (1H, d), 11.01 (1H, s), 11.81 (1H, s), 11.97 (1H, s)	480
548		1.11-1.18 (2H, m), 1.42-1.51 (2H, m), 1.73-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 7.24 (2H, d), 7.70 (2H, d), 8.01-8.04 (1H, m), 8.17-8.20 (2H, m), 11.01 (1H, s), 11.97 (2H, s)	514
549		1.08-1.18 (2H, m), 1.44-1.48 (2H, m), 1.72-1.33 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 7.24 (2H, d), 7.70 (2H, d), 8.00-8.03 (1H, m), 8.15-8.17 (2H, m), 11.03 (1H, s), 12.02 (1H, s)	471

550		1.11-1.18 (1H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 7.23 (2H, d), 7.60-7.64 (2H, m), 7.69 (2H, d), 7.80 (1H, d), 10.96 (1H, s), 11.47 (1H, s), 11.97 (1H, s)	460
551		1.08-1.12 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 2.60 (3H, s), 7.23 (2H, d), 7.48-7.50 (1H, m), 7.70 (2H, d), 7.74 (1H, d), 7.81 (1H, d), 10.97 (1H, s), 11.60 (1H, s), 11.97 (1H, s)	492







Examples 552-569







The following Examples were prepared in two steps: (i) oxadiazole ring synthesis by the procedure of Example 354 with Intermediate 43 and the appropriate commercially available aniline RNH₂ then (ii) ester hydrolysis by the procedure of Example 357 to give the illustrated acid.








Example	R	¹ H NMR δ	MS m/e MH ⁺
552		1.08-1.17 (2H, m), 1.39-1.53 (2H, m), 1.74-1.85 (5H, m), 2.15 (2H, d), 2.42-2.51 (1H, m), 7.23 (2H, d), 7.47 (1H, t), 7.52-7.56 (1H, m), 7.69 (2H, d), 7.82-7.85 (1H, m), 10.94 (1H, s), 11.20 (1H, s), 11.96 (1H, s)	473

281


553		1.08-1.14 (2H, m), 1.44-1.48 (2H, m), 1.74-1.82 (5H, m), 2.15 (2H, d), 2.28 (6H, s), 2.41-2.50 (1H, m), 6.71 (1H, s), 7.22-7.26 (4H, m), 7.68-7.70 (2H, m), 10.77 (1H, s), 10.88 (1H, s), 11.97 (1H, s)	449
554		1.08-1.14 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 7.23 (2H, d), 7.59 (1H, t), 7.69 (2H, d), 7.88-7.92 (1H, m), 8.01-8.03 (1H, m), 10.94 (1H, s), 11.34 (1H, s), 11.97 (1H, s)	507
555		1.08-1.17 (2H, m), 1.42-1.51 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 7.24 (2H, d), 7.35 (1H, d), 7.70 (2H, d), 7.76-7.79 (2H, m), 10.98 (1H, s), 11.62 (1H, s), 11.97 (1H, s)	507
556		1.08-1.17 (2H, m), 1.42-1.51 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 2.78 (3H, s), 7.23 (2H, d), 7.60-7.62 (1H, m), 7.70 (2H, d), 7.92 (1H, d), 8.31 (1H, d), 10.93 (1H, s), 11.19 (1H, s), 11.97 (1H, s)	492
557		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 2.46 (3H, s), 7.23 (2H, d), 7.50 (1H, d), 7.69 (2H, d), 7.73-7.76 (1H, m), 7.96 (1H, d), 10.95 (1H, s), 11.27 (1H, s), 11.97 (1H, s)	458
558		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 2.46 (3H, s), 7.23 (2H, d), 7.50 (1H, d), 7.69 (2H, d), 7.73-7.76 (1H, m), 7.96 (1H, d), 10.95 (1H, s), 11.27 (1H, s), 11.97 (1H, s)	464

559		1.08-1.17 (2H, m), 1.41-1.48 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 7.18-7.24 (3H, m), 7.35-7.40 (1H, m), 7.64-7.70 (2H, m), 8.20-8.22 (1H, m), 10.96 (1H, s), 11.07 (1H, s), 11.97 (1H, s)	473
560		1.11-1.17 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 3.88 (3H, s), 7.23 (2H, d), 7.35 (1H, d), 7.69 (2H, d), 7.82-7.84 (1H, m), 7.90 (1H, d), 10.90 (1H, s), 11.01 (1H, s), 11.97 (1H, s)	519
561		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.73-1.85 (5H, m), 2.13 (2H, d), 2.41-2.50 (1H, m), 7.23 (2H, d), 7.70 (2H, d), 7.78 (1H, d), 7.86-7.89 (1H, m), 8.09 (1H, d), 10.98 (1H, s), 11.55 (1H, s), 11.97 (1H, s)	478
562		1.08-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.41-2.50 (1H, m), 3.88 (3H, s), 7.20-7.24 (3H, m), 7.42-7.44 (2H, m), 7.69 (2H, d), 10.91 (1H, s), 11.11 (1H, s), 11.97 (1H, s)	485
563		1.11-1.17 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 3.86 (3H, s), 6.95 (1H, s), 7.23 (2H, d), 7.52 (2H, d), 7.69 (2H, d), 10.94 (1H, s), 11.33 (1H, s), 11.97 (1H, s)	519
564		1.07-1.17 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (5H, m), 2.14-2.16 (2H, m), 2.42-2.48 (1H, m), 7.23 (2H, d), 7.38-7.41 (1H, m), 7.61 (1H, t), 7.68-7.72 (3H, m), 10.96 (1H, s), 11.38 (1H, s), 11.97 (1H, s)	473

565		1.08-1.18 (2H, m), 1.41-1.50 (2H, m), 1.71-1.85 (5H, m), 2.15 (2H, d), 2.42-2.53 (1H, m), 7.24 (2H, d), 7.67 (2H, d), 7.79-7.81 (1H, m), 7.92-7.95 (1H, m), 8.38 (1H, t), 11.01 (1H, s), 11.44 (1H, s), 11.97 (1H, s)	464
566		1.30-1.39 (2H, m), 1.63-1.72 (2H, m), 1.93-2.06 (5H, m), 2.36 (2H, d), 2.41-2.50 (1H, m), 4.04 (3H, s), 7.43 (3H, t), 7.51-7.54 (1H, m), 7.72-7.76 (1H, m), 7.90 (2H, d), 11.12-11.15 (2H, m), 12.18 (1H, s)	469
567		1.11-1.18 (2H, m), 1.42-1.51 (2H, m), 1.72-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 2.81 (3H, s), 7.23 (2H, d), 7.53-7.56 (1H, m), 7.71 (2H, d), 8.01 (1H, d), 8.31 (1H, d), 10.95 (1H, s), 11.17 (1H, s), 11.97 (1H, s)	492
568		1.11-1.18 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 6.92 (1H, d), 7.08-7.11 (1H, m), 7.22 (2H, d), 7.37 (1H, d), 7.69 (2H, d), 10.49 (1H, s), 10.58 (1H, s), 10.73 (1H, s), 10.87 (1H, s), 11.97 (1H, s)	477
569		1.13 (2H, d), 1.44-1.48 (2H, m), 1.82 (5H, m), 2.14-2.16 (2H, m), 2.41-2.50 (1H, m), 2.55 (3H, s), 7.24 (2H, d), 7.50-7.53 (1H, m), 7.69-7.75 (3H, m), 8.14 (1H, d), 10.94 (1H, s), 11.26 (1H, s), 11.97 (1H, m)	475

284

Example 570: (trans-4-[4-({5-(Quinolin-3-ylamino)-1,3,4-oxadiazol-2-yl}carbonyl)amino)-phenyl]cyclohexyl)acetic acid









Thiocarbonyl diimidazole (178 mg, 1.00 mmol) was added in one portion to a solution of
 5 3-aminoquinoline (144 mg, 1.00 mmol) in DMA (5 mL) and the solution was stirred at
 room temperature for 40 h. Methyl (trans-4-[4-({[hydrazino(oxo)acetyl]amino)phenyl]
 cyclohexyl)acetate (Intermediate 43, 333 mg, 1.00 mmol) was added in one portion and the
 reaction mixture was stirred at room temperature for 20 h and then heated at 80°C for 15
 mins in the microwave. The solution was cooled to room temperature and EDCI (192 mg,
 10 1.00 mmol) was added in one portion and the reaction mixture was heated to 90°C for 20
 mins in the microwave. The reaction mixture was concentrated *in vacuo* to give a gum,
 which was recrystallised (CH₃CN) to give the crude oxadiazole methyl ester intermediate
 as a gummy solid that was used with no further purification.






Lithium hydroxide (420 mg, 10.0 mmol) was added in one portion to a solution of the
 15 crude oxadiazole methyl ester intermediate in a mixture of THF (4 mL), MeOH (8 mL) and
 H₂O (4 mL) and the reaction mixture was stirred at 35°C for 3 h. Citric acid (30 mL) was
 added and the mixture was filtered to leave a solid that was triturated (MeOH). The
 resulting solid was recrystallised using DMSO:H₂O:CH₃CN (7:1:2) to give the title
 compound (216 mg, 46%) as a solid; ¹H NMR δ 1.08-1.18 (2H, m), 1.42-1.52 (2H, m),
 20 1.71-1.85 (6H, m), 2.15 (2H, d), 7.24 (2H, d), 7.59-7.68 (2H, m), 7.70-7.73 (2H, m), 7.99
 (2H, t), 8.62 (1H, d), 8.99 (1H, d), 10.96 (1H, s); MS *m/e* (M-H)⁺ 471.


Examples 571-581

The following examples were prepared by the general procedure of Example 570 using the
 25 appropriate commercially available amino-heterocycle as starting material except where
 stated otherwise.



Example	R	¹ H NMR δ	MS m/e (M-H) ⁻
571		1.07-1.17 (2H, m), 1.41-1.51 (2H, m), 1.69-1.84 (6H, m), 2.14-2.16 (2H, d), 7.14-7.18 (3H, m), 7.22 (2H, d), 7.44- 7.47 (2H, m), 7.68-7.71 (2H, m), 8.10- 8.13 (1H, m), 8.40 (1H, d), 10.92 (1H, s)	547
572		1.07-1.17 (2H, m), 1.41-1.50 (2H, m), 1.70-1.75 (1H, m), 1.78-1.85 (5H, m), 2.13 (2H, d), 3.61 (2H, s), 3.84 (3H, s), 6.88 (1H, d), 7.22 (2H, d), 7.69 (2H, d), 7.92-7.95 (1H, m), 8.39 (1H, d), 10.85 (1H, s)	451
573		1.07-1.18 (2H, m), 1.42-1.50 (2H, m), 1.71-1.85 (7H, m), 2.15 (2H, d), 2.37 (3H, s), 6.89 (1H, d), 7.21-7.24 (2H, m), 7.68-7.73 (3H, m), 8.14 (1H, d), 10.87 (1H, s), 11.84 (2H, s)	435
574		1.08-1.18 (2H, m), 1.41-1.50 (2H, m), 1.71-1.85 (6H, m), 2.14-2.16 (2H, d), 2.27 (3H, s), 7.21-7.24 (2H, m), 7.68-7.72 (3H, m), 7.82 (1H, d), 8.15 (1H, s)	435
575		1.07-1.17 (2H, m), 1.41-1.50 (2H, m), 1.71-1.84 (6H, m), 2.14-2.16 (2H, d), 3.96 (3H, s), 7.06-7.09 (1H, m), 7.23 (2H, d), 7.67-7.70 (2H, d), 7.90-7.91 (1H, m), 8.30-8.33 (1H, m), 10.40 (1H, s), 10.91 (1H, s), 11.97 (1H, s)	451


576		1.07-1.17 (2H, m), 1.41-1.51 (2H, m), 1.69-1.84 (6H, m), 2.15 (2H, d), 2.45- 2.48 (1H, m), 7.23 (2H, d), 7.69 (2H, d), 7.82-7.90 (1H, m), 7.96-8.00 (1H, m), 8.36 (1H, d), 10.95 (1H, s)	439
577		1.11-1.17 (2H, m), 1.41-1.51 (2H, m), 1.71-1.84 (6H, m), 2.15 (2H, d), 3.77 (3H, s), 6.94-6.98 (2H, m), 7.04-7.08 (3H, m), 7.22 (2H, d), 7.68-7.70 (2H, m), 8.04- 8.07 (1H, m), 8.36 (1H, d), 10.91 (1H, s)	543
578 ¹		1.06-1.17 (2H, m), 1.41-1.51 (2H, m), 1.69-1.84 (5H, m), 2.15 (2H, d), 5.36 (2H, d), 6.99-7.01 (1H, m), 7.22 (2H, d), 7.37-7.44 (4H, m), 7.51 (1H, s), 7.68-7.70 (2H, d), 7.96-7.99 (1H, m), 8.40-8.41 (1H, m), 10.90 (1H, s)	561
579		1.08-1.17 (2H, m), 1.42-1.50 (2H, m), 1.72-1.85 (6H, m), 2.14-2.16 (2H, d), 6.48 (1H, d), 7.23 (2H, d), 7.49 (1H, d), 7.70 (2H, d), 8.27 (1H, d), 8.36 (1H, d), 10.84 (1H, s), 10.89 (1H, s), 11.61 (1H, s)	460
580 ²		1.14 (2H, m), 1.46 (2H, m), 1.82 (5H, m), 2.14-2.16 (2H, d), 7.24 (2H, d), 7.69-7.72 (2H, d), 8.08 (1H, d), 8.26-8.29 (1H, m), 8.84 (1H, d), 11.97 (1H, s)	447

581 ²			447
------------------	---	--	-----

¹The amino-heterocycle was synthesised as described as in WO2001021597

²Purified by reverse phase preparative HPLC, eluting with a gradient of CH₃CN and H₂O containing 0.2% TFA

5 **Example 582: *rac-trans*-Ethyl 2-[4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]-amino}phenyl)cyclohexyl]propanoate**



2,4,5-Trifluorophenyl isothiocyanate (252 mg, 1.33 mmol) was added in one portion to a suspension of *trans*-ethyl 2-[4-(4-[[hydrazino(oxo)acetyl]amino}phenyl)cyclo-


10 hexyl]propanoate (Intermediate 99, 400 mg, 1.11 mmol) in DMA (6 mL) and the reaction mixture was stirred at room temperature for 2 h. EDCI (319 mg, 1.66 mmol) was added in one portion and the reaction mixture was heated to 90°C for 10 mins in the microwave.



The mixture was then cooled to room temperature and H₂O (20 mL) was added. The mixture was filtered to leave a solid, which was washed with H₂O (10 mL) to give the title

15 compound (550 mg, 95%) as a solid; ¹H NMR δ 1.1 (3H, d), 1.12-1.28 (2H, m), 1.2 (3H, t), 1.35-1.51 (2H, m), 1.52-1.64 (1H, m), 1.65-1.75 (1H, m), 1.77-1.9 (3H, m), 2.22-2.32 (1H, m), 2.38-2.51 (1H, m), 4.01-4.16 (2H, m), 7.2 (2H, d), 7.65-7.78 (1H, m), 7.69 (2H, d), 8.1-8.25 (1H, m), 10.95 (1H, s), 11.05 (1H, s); MS m/e MH⁺ 517.


20 **Examples 583-584**

The following examples were prepared by the general procedure of Example 582 using the appropriate commercially available isothiocyanate as starting material.



Example	R	¹ H NMR δ	MS m/e (M-H) ⁻
583		1.08 (3H, d), 1.1-1.28 (2H, m), 1.2 (3H, t), 1.35-1.5 (2H, m), 1.51-1.64 (1H, m), 1.65-1.74 (1H, m), 1.75-1.9 (3H, m), 2.22-2.34 (1H, m), 2.39-2.5 (1H, m), 4.01-4.16 (2H, m), 7.21 (2H, d), 7.7 (2H, d), 7.78 (2H, d), 7.89 (2H, d), 11.0 (1H, s), 11.55 (1H, s)	488
584		1.1 (3H, d), 1.12-1.28 (2H, m), 1.2 (3H, t), 1.37-1.51 (2H, m), 1.52-1.64 (1H, m), 1.65- 1.75 (1H, m), 1.77-1.9 (3H, m), 2.22-2.34 (1H, m), 2.39-2.5 (1H, m), 4.03-4.17 (2H, m), 7.22 (2H, d), 7.32-7.4 (1H, m), 7.44-7.55 (1H, m), 7.63-7.75 (1H, m), 7.7 (2H, d), 10.95 (1H, s), 11.25 (1H, s)	499

Example 585: rac- 2-(trans-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid



- 5 A solution of lithium hydroxide (17 mg, 0.39 mmol) in H₂O (1.0 mL) was added in one portion to a solution of *trans*-ethyl 2-(4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]-amino}phenyl)cyclohexyl)propanoate (Example 582, 25 mg, 0.05 mmol) in a mixture of THF (1.5 mL) and MeOH (1.5 mL) and the reaction mixture was stirred at room temperature for 24 h. A 1M aqueous solution of citric acid (10 mL)
- 10 was added and the mixture was filtered to leave a solid. The solid was washed with H₂O (2 mL) and purified by reverse phase preparative HPLC, eluting with a gradient of CH₃CN and H₂O containing 0.2 % TFA to give the title compound (10 mg, 42 %) as a solid. The individual isomers may be separated by chiral chromatography under standard conditions and purified by recrystallisation from EtOH to give:


(2R)-2-(trans-4-{4-[(5-{(2,4,5-Trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid

¹H NMR δ 1.01-1.1 (5H, m), 1.1-1.28 (2H, m), 1.37-1.5 (2H, m), 1.51-1.63 (1H, m), 1.71-1.9 (4H, m), 2.14-2.22 (1H, m), 2.4-2.5 (1H, m), 3.45 (2H, dq), 4.32 (1H, t), 7.21 (2H, d),
5 7.62-7.75 (m, 1H), 7.7 (2H, d), 8.1-8.22 (m, 1H), 10.95 (1H, s), 11.03 (1H, s), 12.0 (1H, s);
MS m/e MH⁺ 489.

(2S)-2-(trans-4-{4-[(5-{(2,4,5-Trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid

¹H NMR δ 1.01-1.1 (5H, m), 1.1-1.28 (2H, m), 1.37-1.5 (2H, m), 1.51-1.63 (1H, m), 1.71-1.9 (4H, m), 2.14-2.22 (1H, m), 2.4-2.5 (1H, m), 3.45 (2H, dq), 4.32 (1H, t), 7.21 (2H, d),
10 7.62-7.75 (m, 1H), 7.7 (2H, d), 8.1-8.22 (m, 1H), 10.95 (1H, s), 11.03 (1H, s), 12.0 (1H, s);
MS m/e MH⁺ 489.

Example 586: rac-trans-2-(4-{4-[(5-{(3,4-Difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-phenyl)cyclohexyl)propanoic acid



The following example was prepared using the method described for Example 585 but using Example 584 as starting material to give the title compound (9.5 mg, 40 %) as a solid. The individual isomers may be separated by chiral chromatography under standard
20 conditions and purified by recrystallisation from EtOH to give:

(2R)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid


¹H NMR δ 1.05 (3H, d), 1.1-1.26 (2H, m), 1.35-1.5 (2H, m), 1.51-1.62 (1H, m), 1.7-1.9 (4H, m), 2.12-2.23 (1H, m), 2.38-2.5 (1H, m), 7.21 (2H, d), 7.29-7.38 (1H, m), 7.41-7.51 (1H, m), 7.64-7.76 (1H, m), 7.69 (2H, d), 11.0 (1H, s); MS m/e MH⁺ 471.
25

(2S)-2-(trans-4-{4-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-phenyl)cyclohexyl)propanoic acid

¹H NMR δ ¹H NMR δ: 1.05 (3H, d), 1.1-1.26 (2H, m), 1.35-1.5 (2H, m), 1.51-1.62 (1H, m), 1.7-1.9 (4H, m), 2.12-2.23 (1H, m), 2.38-2.5 (1H, m), 7.21 (2H, d), 7.29-7.38 (1H, m),

7.41-7.51 (1H, m), 7.64-7.76 (1H, m), 7.69 (2H, d), 11.0 (1H, s), 11.7 (1H, s); MS m/e MH⁺ 471.


Example 587: *rac-trans*-2-(4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]-phenyl}cyclohexyl)propanoic acid








Potassium trimethylsilanolate (65 mg, 0.51 mmol) was added in one portion to a solution of *trans*-ethyl 2-(4-{4-[(5-{(4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]-phenyl}cyclohexyl)propanoate (Example 583, 25 mg, 0.051 mmol) in THF (2 mL) and the reaction mixture was stirred at room temperature for 48 h. A 1M aqueous solution of citric acid (10 mL) was added and the mixture was filtered to leave a solid. The solid was washed with H₂O (2 mL) and purified by reverse phase preparative HPLC, eluting with a gradient of CH₃CN and H₂O containing 0.2% TFA, to give the title compound (11 mg, 47 %) as a solid; ¹H NMR δ 1.05 (3H, t), 1.1-1.3 (2H, m), 1.38-1.51 (2H, m), 1.52-1.63 (1H, m), 1.7-1.9 (4H, m), 2.13-2.25 (1H, m), 2.4-2.5 (1H, m), 7.22 (2H, d), 7.7 (2H, d), 7.76 (2H, d), 7.85 (2H, d), 11.0 (1H, s); MS m/e MH⁺ 460.





Examples 588-597

The following Examples were prepared in two steps: (i) oxadiazole ring synthesis by the procedure of Example 1 with Intermediate 43 and the appropriate commercially available isothiocyanate R-NCS then (ii) hydrolysis of the crude ester product by the procedure of Example 357 to give the indicated acid.



Example	R	¹ H NMR δ	MS m/e MH ⁺
588		1.07 - 1.17 (2H, m), 1.41 - 1.51 (2H, m), 1.70 - 1.85 (6H, m), 2.15 (2H, d), 2.28 (3H, s), 7.19 - 7.24 (4H, m), 7.48 - 7.50 (2H, m),	433 (M-H) ⁺


		7.69 (2H, d), 10.83 - 10.89 (2H, m), 11.97 (1H, s)	
589		1.11 - 1.14 (2H, m), 1.44 - 1.48 (2H, m), 1.82 (6H t), 2.14 - 2.16 (2H, m), 7.23 (2H d), 7.58 (4H, s), 7.69 (2H d), 10.94 (1H, s), 11.13 (1H, s), 11.98 (1H, s)	499
590		1.14 (2H, d), 1.33 (3H, t), 1.44 - 1.48 (2H, m), 1.82 (6H, t), 2.15 (2H d), 4.01 (2H, q), 6.95 - 6.98 (2H, m), 7.22 (2H, d), 7.49 - 7.51 (2H, m), 7.69 (2H, d), 10.71 (1H, s), 10.88 (1H, s), 11.97 (1H, s)	465
591		1.08 - 1.18 (2H, m), 1.29 (9H, s), 1.42 - 1.50 (2H, m), 1.70 - 1.85 (6H, m), 2.15 (2H, d), 7.23 (2H, d), 7.39 - 7.43 (2H, m), 7.51 - 7.53 (2H, m), 7.69 (2H, d), 10.83 (1H, s), 10.89 (1H, s), 11.98 - 11.99 (1H, m)	477
592		1.13 (2H, d), 1.37 (3H, t), 1.44 - 1.48 (2H, m), 1.82 (6H, m), 2.15 (2H, d), 4.11 (2H, q), 6.97 - 7.02 (1H, m), 7.08 - 7.10 (2H, m), 7.23 (2H, d), 7.68 (2H, d), 7.92 - 7.94 (1H, m), 9.96 (1H, s), 10.89 (1H, s), 11.97 (1H, s)	465
593		1.07 - 1.17 (2H, m), 1.40 - 1.51 (2H, m), 1.68 - 1.84 (6H, m), 2.16 (2H, m), 2.30 (3H, s), 7.08 - 7.15 (1H, m), 7.17 - 7.37 (4H, m), 7.67 - 7.74 (3H, m), 10.02 (1H, s), 10.89 (1H, s), 12.00 (1H, s)	435

594		1.07 - 1.17 (2H, m), 1.41 - 1.51 (2H, m), 1.70 - 1.84 (6H, m), 2.15 (2H, d), 7.22 - 7.26 (4H, m), 7.68 - 7.71 (2H, m), 7.73 - 7.75 (2H, m), 7.84 - 7.86 (2H, m), 10.96 (1H, s), 11.41 (1H, s), 11.99 (1H, s)	498 (M-H) ⁻
595		1.10 - 1.14 (2H, m), 1.44 - 1.48 (2H, m), 1.73 - 1.84 (6H, m), 2.14 - 2.16 (2H, m), 7.20 - 7.24 (3H, m), 7.41 - 7.45 (1H, m), 7.54 - 7.57 (1H, m), 7.68 (2H, d), 7.95 - 7.98 (1H, m), 10.41 (1H, s), 10.93 (1H, s), 12.00 (1H, s)	453 (M-H) ⁻
596		1.08 - 1.15 (2H, m), 1.44 - 1.48 (2H, m), 1.74 - 1.85 (6H, m), 2.14 - 2.16 (2H, m), 3.74 - 3.78 (3H, m), 6.65 - 6.67 (1H, m), 7.14 - 7.32 (5H, m), 7.68 - 7.71 (2H, m), 10.91 - 10.94 (2H, m), 11.97 (1H, s)	451
597		1.10 - 1.14 (2H, m), 1.44 - 1.48 (2H, m), 1.82 (6H, t), 2.15 (2H, q), 7.23 (2H, d), 7.70 (2H, d), 7.76 (4H, s), 10.96 (1H, s), 11.40 (1H, s), 12.00 (1H, s)	519 (M-H) ⁻





Examples 598-601

The following examples were prepared by the general procedure of Example 483, using the appropriate aniline (commercially available unless indicated otherwise) and 3,4-

5 difluorophenylisothiocyanate.




- 270 -

Example	R	¹ H NMR δ	MS m/e MII ⁺
598		6.99-7.07 (4H, m), 7.12-7.16 (1H, m), 7.34-7.42 (3H, m), 7.45-7.52 (1H, m), 7.68-7.73 (1H, m), 7.81-7.83 (2H, m), 11.07 (1H, s) 11.23 (1H, s)	407 (M-H) ⁺
599 ¹		3.76 (3H, s), 6.85-6.87 (2H, m), 6.96-6.99 (1H, m), 7.00-7.04 (1H, m), 7.17-7.21 (2H, m), 7.35 (1H, t), 7.45-7.52 (1H, m), 7.67-7.73 (3H, m), 10.98 (1H, s), 11.22 (1H, s)	439
600 ²		2.30 (3H, s), 6.90-6.93 (2H, m), 6.99-7.01 (2H, m), 7.19-7.21 (2H, m), 7.35 (1H, d), 7.44-7.49 (1H, m), 7.67-7.73 (1H, m), 7.78-7.80 (2H, m), 11.04 (2H, s)	423
601		3.75 (3H, s), 6.53-6.58 (2H, m), 6.70-6.73 (1H, m), 7.04-7.08 (2H, m), 7.28 (1H, t), 7.34-7.38 (1H, m), 7.45-7.52 (1H, m), 7.68-7.73 (1H, m), 7.80-7.84 (2H, m), 11.08 (1H, s), 11.24 (1H, s)	439

¹Preparation of aniline starting material described in Patent Application WO 2001056990²Preparation of aniline starting material described in Patent Application WO 9521815

Example 602: *N*-(3,5-Dichloro-4-morpholin-4-ylphenyl)-5-[(3-methylphenyl)amino]-1,3,4-oxadiazole-2-carboxamide




Following the general procedure of Example 1, using Intermediate 100 and 3-methylphenylisothiocyanate as starting materials, the title compound was obtained; ¹H

294


NMR δ 2.30 (3H, s), 3.11 (4H, t), 3.67 - 3.69 (4H, m), 6.87 (1H, d), 7.25 (1H, t), 7.36 (1H, s), 7.42 (1H, d), 7.91 (2H, s), 10.94 (1H, broad s), 11.17 (1H, broad s); MS m/e MH^+ 448.

Example 603: N-[6-[4-(Cyclopropylmethyl)piperazin-1-yl]pyridin-3-yl]-5-[(2-Fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide



5
10
15
5-[(2-Fluorophenyl)amino]-N-[6-piperazin-1-yl]pyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (Intermediate 41) (120 mg, 0.31 mmol) in DMA (2 mL) was placed in a microwave tube with cyclopropylmethyl bromide (88 mg, 0.62 mmol), and triethylamine (64 mg, 0.62 mmol). The mixture was heated with stirring in the microwave at 110°C for 60 minutes then transferred directly onto a reverse phase HPLC column and eluted with a gradient of acetonitrile/water containing 0.2%/TFA to give the title compound (125 mg, 91%) as a solid; ^1H NMR δ 10.65 (1H, s), 10.40 (1H, bs), 8.20 (1H, s), 7.65 (2H, m), 6.90 (2H, m), 6.80 (1H, m), 6.65 (1H, d), 4.00 (2H, d), 3.25 (2H, d), 2.80 (6H, m), 0.75 (1H, m), 0.30 (2H, m), 0.00 (2H, m); MS m/e MH^+ 438.


Example 604: 5-[(2-Fluorophenyl)amino]-N-[6-[4-(2-methoxyethyl)piperazin-1-yl]pyridin-3-yl]-1,3,4-oxadiazole-2-carboxamide



20
Following the procedure of Example 603 except that 1-bromo-2-methoxyethane was used in place of cyclopropylmethyl bromide, the title compound was obtained; ^1H NMR δ 10.98 (1H, s), 10.73 (1H, bs), 8.55 (1H, s), 8.00 (2H, m), 7.30 (2H, m), 7.15 (1H, m), 6.95 (1H, d), 4.30 (2H, bd), 3.70 (2H, m), 3.55 (2H, m), 3.35 (5H, m), 3.15 (4H, m); MS m/e MH^+ 442.

25 **Examples 605-609**

The following examples were prepared by the general procedure of Example 72 using Intermediate 40 and the appropriate aniline.



Example	R	¹ H NMR δ	MS m/e MH ⁺
605		10.90 (1H, s), 10.75 (1H, s), 8.05 (1H, t), 7.70 (1H, d), 7.30 (2H, m), 7.15 (1H, m), 7.00 (2H, d), 4.20 (2H, m), 3.70 (4H, bs), 3.25 (4H, bs), 2.70 (2H, m)	428
606		10.90 (1H, s), 10.85 (1H, s), 8.05 (1H, t), 7.70 (2H, d), 7.30 (2H, d), 7.20 (1H, m), 6.95 (2H, d), 4.00 (2H, t), 3.55 (2H, t), 1.85 (2H, q)	394 (M+Na) ⁺
607		10.85 (1H, s), 10.75 (1H, s), 8.05 (1H, t), 7.70 (2H, d), 7.30 (2H, m), 7.15 (1H, m), 6.95 (2H, d), 4.05 (2H, t), 3.70 (2H, t), 3.50 (2H, q), 1.15 (3H, t)	385 (M-H) ⁻
608		10.95 (1H, s), 10.70 (1H, bs), 8.05 (1H, t), 7.65 (2H, d), 7.30 (2H, m), 7.15 (1H, m), 6.80 (2H, d), 3.65 (2H, q), 3.06 (2H, q), 1.55 (6H, s), 1.05 (3H, t), 0.90 (3H, t)	456
609			368

The aniline starting materials are described in the following references:

¹PCT Patent Application WO2004005282 (4-(2-morpholin-4-ylethoxy)aniline)


²PCT Patent Application WO9515952 (3-(4-aminophenoxy)propan-1-ol)

5 ³German Patent Application DE2732384 (4-(2-ethoxyethoxy)aniline)

⁴Indian Journal of chemistry, Section B, 1986,25B(12),1277 (2-(4-aminophenoxy)-N,N-diethyl-2-methylpropanamide)

⁵PCT Patent Application WO2004073634A2 (4-pyrrolidin-1-ylaniline)


Example 610: Methyl *trans*-4-(4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl)cyclohexanecarboxylate



Following the procedure of Example 105 except that


- 5 methyl *trans*-4-(4-[(hydrazino(oxo) acetyl]amino)phenyl)cyclohexanecarboxylate (Intermediate 44) and 2,4,5-trifluorophenylisothiocyanate were used as starting materials the title compound was obtained as solid; ^1H NMR δ 1.47 (1H, s), 1.49 (2H, d), 1.52 (1H, s), 1.85 (2H, d), 2.01 (2H, d), 2.45 (1H+DMSO, m), 2.53 - 2.56 (1H, m), 3.62 (3H, s), 7.22 - 7.24 (2H, m), 7.68 (1H, s), 7.71 (2H, q), 8.15 - 8.18 (1H, m), 10.96 (1H, s), 11.04
- 10 (1H, s) MS m/e MH^+ 475.

Example 611: *trans*-4-{4-[(5-[(2,4,5-Trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]-amino]phenyl)cyclohexanecarboxylic acid




- Following the procedure of Example 112 except that the starting material was methyl
- 15 *trans*-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl)cyclohexanecarboxylate (Example 610) and purification was by trituration with MeCN the title compound was obtained as a solid; ^1H NMR δ 1.44 (1H, s), 1.48 (3H, d), 1.52 (1H, s), 1.84 (2H, d), 2.00 (2H, d), 2.25 (2H, d), 2.45 (1H+DMSO, m), 2.53-2.56 (1H, m), 7.23 (2H, d), 7.68 (1H, s), 7.69 (2H, d), 8.15 (1H, s); MS m/e MH^+
- 20 461.

Example 612: Methyl 6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]-piperidin-1-yl}nicotinate




A solution of 5-[(2-fluorophenyl)amino]-N-piperidin-4-yl-1,3,4-oxadiazole-2-carboxamide (Intermediate 101) (100 mg, 0.33 mmol) and methyl 6-chloronicotinate (56 mg, 0.33 mmol) in dry DMF (3 mL) was heated in the microwave at 120°C for 1 h. The reaction mixture was filtered and the filtrate was purified by preparative HPLC to give the title compound (17 mg, 12%) as a solid; ¹H NMR δ 1.66-1.53 (m, 2H), 1.92-1.82 (m, 2H), 3.12-2.99 (m, 2H), 3.80 (s, 3H), 4.16-4.04 (m, 1H), 4.54-4.42 (m, 2H), 6.92 (d, 1H), 7.18-7.10 (m, 1H), 7.33-7.21 (m, 2H), 7.97-7.92 (m, 1H), 8.04-7.98 (m, 1H), 8.65 (d, 1H), 9.03 (d, 1H), 10.62 (s, 1H); MS m/e MH⁺ 441.

10 **Example 613: 6-{4-[(5-[(2-Fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-piperidin-1-yl}nicotinic acid**



A mixture of 2M NaOH (2 mL, 4 mmol) and methyl 6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}piperidin-1-yl}nicotinate (17 mg, 0.04 mmol) (Example 612) in MeOH (10 mL) was stirred for 16 h then taken to pH 7 by addition of 2M HCl. The solution was concentrated by evaporation to give a cloudy aqueous mixture that was cooled on an ice bath. The precipitate was collected by filtration, washed with water and Et₂O and dried to give the title compound (13 mg, 88%) as a solid; ¹H NMR (CD₃OD) δ 8.61 (s, 1H), 8.02-7.88 (m, 2H), 7.16-7.05 (m, 2H), 7.05-6.97 (m, 1H), 6.75 (d, 1H), 4.47-4.37 (m, 2H), 4.17-4.06 (m, 1H), 3.08-2.97 (m, 2H), 1.99-1.89 (m, 2H), 1.63-1.50 (m, 2H); MS m/e MH⁺ 427.

Example 614: 6-{4-[(5-[(2-Fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}-piperidin-1-yl}nicotinamide




A solution of 5-[(2-fluorophenyl)amino]-N-piperidin-4-yl-1,3,4-oxadiazole-2-carboxamide (Intermediate 101) (200 mg, 0.66 mmol) and 6-chloronicotinamide (103 mg, 0.66 mmol) in dry DMF (4 mL) was heated in the microwave at 125°C for 20 minutes and then for a

- 275 -


further 10 minutes at 135°C. The mixture was filtered and purified by preparative HPLC to give the title compound (7 mg, 2.5%) as a solid; ¹H NMR δ 1.55-1.65 (2H, m), 1.83-1.86 (2H, m), 2.99 (2H, t), 4.04-4.12 (1H, m), 4.42-4.46 (2H, m), 6.87 (1H, d), 7.11 (2H, m), 7.21-7.30 (2H, m), 7.72 (1H, s), 7.94-8.03 (2H, m), 8.62 (1H, d), 8.99 (1H, d), 10.61 (1H, s); MS m/e MH⁺ 426.

Example 615: (6-{4-[(5-{(2-Fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]piperidin-1-yl}pyridin-3-yl)acetic acid



- 10 A mixture of 2M NaOH (0.09 mL, 0.18 mmol) and methyl (6-{4-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]piperidin-1-yl}pyridin-3-yl)acetate (Intermediate 102) (41 mg, 0.09 mmol) in MeOH (1 mL) was stirred for 16 h then acidified with 2M HCl and concentrated by evaporation. The residue purified by preparative HPLC to give the title compound (1 mg) as a gum; ¹H NMR (CD₃OD) δ 1.78-1.63 (m, 3H), 2.11-2.00 (m, 2H), 3.35-3.25 (m, 2H), 3.55 (s, 2H), 4.23-4.07 (m, 3H), 7.07-6.99 (m, 1H), 7.15-7.07 (m, 2H), 7.30-7.23 (m, 1H), 7.79 (s, 1H), 7.89-7.83 (m, 1H), 7.99-7.91 (m, 1H); MS m/e MH⁺ 441.


- 20 **Example 616: 5-{[3-(Anilino)carbonyl]phenyl}amino-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide**



- EDCI (77 mg, 0.40 mmol) was added to a suspension of 3-[(5-{[3-(Anilino)carbonyl]phenyl}amino)-1,3,4-oxadiazole-2-carboxamide]benzoic acid (Intermediate 103) (150 mg, 0.37 mmol), aniline (0.043 mL, 0.44 mmol) and HOBt (54 mg, 0.40 mmol) in DMF (4.5 mL). After 72 hours the mixture was diluted with H₂O (20 mL) and the solid filtered. Purification by reverse phase preparative HPLC (2 injections of 55 mg in 2 mL DMF) gave the title compound (as a trifluoroacetate salt) (6.8 mg, 3%) as a solid; ¹H NMR δ 3.45 (4H, t), 3.72 (4H, t), 6.96 (1H, d), 7.12 (1H, t), 7.37 (2H, t), 7.56 (1H, t), 7.65 (1H, d), 7.77 (1H,


s), 7.79 - 7.79 (1H, m), 7.84 - 7.86 (1H, m), 8.00 - 8.03 (1H, m), 8.11 (1H, d), 8.53 (1H, d), 10.29 (1H, s), 11.03 (1H, s), 11.17 (1H, s); MS m/e (MH)⁺ 486.

Example 617: *N*-(4-[*trans*-4-(Aminomethyl)cyclohexyl]phenyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide




4M HCl in 1,4-dioxane (1 mL) was added to *tert*-butyl [(*trans*-4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl}cyclohexyl)-methyl]carbamate (Example 618) (50 mg, 0.10 mmol) and the suspension stirred for 18 hours. All volatiles were removed *in vacuo* and the residue triturated with Et₂O (3 mL), the solid filtered, washed with Et₂O (2 x 5 mL), isohexane (2 x 5 mL) to give the title compound (as a hydrochloride salt) (44 mg, 99%) as a white solid; ¹H NMR δ 1.07 - 1.16 (2H, m), 1.39 - 1.47 (2H, m), 1.57 - 1.69 (1H, m), 1.87 (4H, t), 2.41 - 2.52 (1H, m), 2.71 (2H, d), 7.14 - 7.20 (1H, m), 7.23 (1H, s), 7.25 - 7.27 (2H, m), 7.29 - 7.34 (1H, m), 7.70 (2H, d), 7.87 (3H, s), 8.01 - 8.06 (1H, m), 10.76 (1H, s), 10.94 (1H, s); MS m/e (MH)⁺ 410.

Example 618 : *tert*-Butyl [(*trans*-4-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]phenyl}cyclohexyl)methyl]carbamate




The following example was prepared according to the procedure of Example 141 using Intermediate 104 and 2-fluorophenyl isothiocyanate to give the title compound as a solid; ¹H NMR δ 1.00 - 1.06 (2H, m), 1.35 - 1.39 (12H, m), 1.79 - 1.81 (4H, m), 2.44 (1H, t), 2.82 (2H, t), 6.81 (1H, t), 7.13 - 7.34 (5H, m), 7.68 (2H, d), 8.02 - 8.07 (1H, m), 10.75 (1H, s), 10.92 (1H, s); MS m/e (MH)⁺ 510.

Example 619: Methyl (1-{5-[(5-{[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetate



To a suspension methyl [1-(5-{[hydrazino(oxo)acetyl]amino}pyridin-2-yl)piperidin-4-yl]acetate (Intermediate 77) (250 mg, 0.745 mmol) in anhydrous DMF (3 mL) was added 3,4-difluoroisothiocyanate (155 mg, 0.894 mmol) in anhydrous DMF (1 mL) and the suspension heated to 45 °C. A clear solution quickly formed and heating and stirring continued for a further ~1hr. EDAC.HCl (175 mg, 0.913 mmol) was added and the reaction mixture heated to 85 °C for 2.5 hrs, and allowed to cool to ambient temperature overnight. Water was added to the reaction mixture and a yellow suspension formed which was filtered and dried. This material was suspended in absolute EtOH and heated to 80 °C, after ~5 mins the suspension was filtered and dried to provide the title compound as a yellow solid (189 mg, 0.727 mmol; 79%); ¹H NMR δ 1.14 - 1.24 (m, 2H), 1.70 - 1.73 (m, 2H), 1.89 - 1.98 (m, 1H), 2.28 (d, 2H), 2.76 - 2.83 (m, 2H), 3.61 (s, 3H), 4.22 - 4.25 (m, 2H), 6.85 (d, 1H), 7.34 - 7.37 (m, 1H), 7.45 - 7.52 (m, 1H), 7.67 - 7.73 (m, 1H), 7.89 (d, 1H), 8.46 (d, 1H), 10.90 (s, 1H), 11.21 (s, 1H); MS m/e MH⁺ 473.


Example 620: (1-{5-[(5-{[(3,4-Difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetic acid



To a solution of methyl (1-{5-[(5-{[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetate (Example 619, 189 mg, 0.40 mmol) in MeOH (10 mL) was added a 2M solution of NaOH (1 mL). The resulting yellow solution was allowed to stir at ambient temperature overnight. The volatile organics were removed by evaporation and residue adjusted to pH ~1 with 2M HCl and the suspension


was filtered and dried to leave a solid which was dissolved in DMSO/MeCN/water(7:2:1) and purified on a reverse phase preparative HPLC column eluting 5-95% MeCN 0.2% TFA. The title compound was isolated as a yellow solid (108 mg, 0.235 mmol; 58.9%); ¹H NMR δ 1.16 - 1.27 (m, 2H), 1.76 (d, 2H), 1.91 - 1.99 (m, 1H), 2.19 (d, 2H), 2.90 (t, 2H), 4.21 (d, 2H), 7.00 (br d, 1H), 7.34 - 7.37 (m, 1H), 7.49 (q, 1H), 7.67 - 7.73 (m, 1H), 7.98 (d, 1H), 8.46 (s, 1H), 11.01 (s, 1H), 11.23 (s, 1H); CO₂H not seen; MS m/e MH⁺ 459.

Example 621: Methyl [1-(5-[(5-[(4-(trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]pyridin-2-yl)piperidin-4-yl]acetate



Prepared as described for Example 619 except 4-(trifluoromethoxy)phenylisothiocyanate (197mg, 0.89 mmol) was used as starting material. The title compound was obtained as a yellow solid (270 mg, 0.519 mmol; 69%); ¹H NMR δ 1.14 - 1.24 (m, 2H), 1.71 (d, 2H), 1.89 - 1.98 (m, 1H), 2.28 (d, 2H), 2.79 (t, 2H), 3.60 (s, 3H), 4.23 (d, 2H), 6.84 (d, 1H), 7.42 (d, 2H), 7.71 (d, 2H), 7.89 (d, 1H), 8.46 (s, 1H), 10.89 (s, 1H), 11.17 (s, 1H); MS m/e MH⁺ 521.


Example 622: [1-(5-[(5-[(4-(Trifluoromethoxy)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]pyridin-2-yl)piperidin-4-yl]acetic acid



Prepared as described for Example 620 except Example 621 was used as starting material. The title compound was isolated as a yellow solid, (172 mg, 0.361 mmol; 69.4%).


¹H NMR δ 1.16 - 1.26 (m, 2H), 1.76 (d, 2H), 1.89 - 1.98 (m, 1H), 2.19 (d, 2H), 2.90 (t, 2H), 4.21 (d, 2H), 7.01 (d, 1H), 7.42 (d, 2H), 7.71 (d, 2H), 7.98 (d, 1H), 8.46 (s, 1H), 11.00 (s, 1H), 11.19 (s, 1H); CO₂H not seen; MS m/e MH⁺ 507.

5 **Example 623: Methyl 2-methyl-2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)propanoate**



A solution of 2,4,5-trifluorophenyl isothiocyanate (183 mg, 0.96 mmol) in DMA (1 mL) was added in one portion to a suspension of methyl 2-[trans-4-(4-
10 {[hydrazino(oxo)acetyl]amino}-phenyl)cyclohexyl]-2-methylpropanoate (Intermediate 105, 290 mg, 0.80 mmol) in DMA (3 mL) and the reaction mixture was stirred at room temperature for 1 h 30 mins. EDCI (230 mg, 1.20 mmol) was added in one portion and the reaction mixture was heated to 90°C for 10 mins in the microwave. The mixture was then cooled to room temperature, poured into H₂O (20 mL) and stirred for 5 mins. The mixture
15 was centrifuged and the aqueous layer decanted off to leave a solid residue. The solid residue was taken up in H₂O (5 mL), stirred for 5 mins and then centrifuged and the aqueous layer decanted off to leave a solid residue. H₂O (5 mL) was added and the mixture was filtered to give the title compound (398 mg, 96%) as a solid; ¹H NMR δ 1.1 (6H, s), 1.12-1.26 (2H, m), 1.36-1.5 (2H, m), 1.57-1.7 (3H, m), 1.78-1.89 (2H, m), 2.39-
20 2.49 (1H, m), 3.6 (3H, s), 7.2 (2H, d), 7.65-7.78 (1H, m), 7.68 (2H, d), 8.12-8.24 (1H, m), 11.0 (1H, s), 11.1 (1H, s); MS m/e MH⁺ 517.


Example 624: 2-Methyl-2-(trans-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)propanoic acid



- 280 -

Potassium trimethylsilanolate (970 mg, 7.6 mmol) was added in one portion in a stirred solution of methyl 2-methyl-2-(*trans*-4-{4-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate (Example 623, 390 mg, 0.76 mmol) in THF (10 mL) and the reaction mixture was heated in the microwave at 90°C for 20 min, then 100°C for 1 h and finally at 120°C for 30 min. The reaction mixture was concentrated *in vacuo* to leave a yellow solid and then a 1M aqueous solution of citric acid (40 mL) was added and the mixture was stirred for 30 mins. The mixture was filtered to leave a solid. The solid was washed with H₂O (20 mL) and purified by recrystallisation, using ethanol as solvent, to give the title compound (219 mg, 57 %) as a white solid; ¹H NMR δ 1.06 (6H, s), 1.12-1.27 (2H, m), 1.37-1.5 (2H, m), 1.58-1.75 (3H, m), 1.8-1.9 (2H, m), 2.39-2.5 (1H, m), 7.22 (2H, m), 7.65 (2H, m), 7.68-7.78 (1H, m), 8.14-8.24 (1H, m), 11.0 (1H, s), 11.13 (1H, s), 12.1 (1H, s); MS m/e MH⁺ 502.

Example 625: 1-{5-[(5-{(3,4-Difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}-4-methylpiperidine-4-carboxylic acid




To a solution of methyl 1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}-4-methylpiperidine-4-carboxylate (Example 631; 282 mg, 0.60 mmol) in MeOH (10 mL) was added a 2M solution of NaOH (1.5 mL). The resulting yellow solution was allowed to stir at ambient temperature overnight. The reaction mixture was then heated to 50 °C for 8 hours and allowed to stir at ambient temperature overnight. The volatile organics were removed by evaporation, the aqueous residue was washed with ether (50 mL) and the aqueous phase adjusted to pH ~1 with 2M HCl. The resultant suspension was filtered and dried under vacuum at ambient temperature to give the title compound (230 mg, 0.502 mmol, 84%); ¹H NMR δ 1.21 (s, 3H), 1.47 - 1.54 (m, 2H), 2.03 - 2.08 (m, 2H), 3.29 - 3.36 (m, 2H), 3.95 - 4.00 (m, 2H), 7.30 (d, 1H), 7.37 - 7.41 (m, 1H), 7.48 (q, 1H), 7.69 - 7.75 (m, 1H), 8.15 (d, 1H), 8.54 (s, 1H), 11.25 (s, 1H), 11.37 (s, 1H); CO₂H not seen; MS m/e MH⁺ 459.

The following examples were prepared by the general procedure described above using the appropriate starting esters (Examples 632-636).

Example	Name	¹ H NMR δ	MS m/e M+H ⁺
626	1-{5-[(5-{[4-ethylphenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}-4-methylpiperidine-4-carboxylic acid	1.16 - 1.20 (m, 6H), 1.42 - 1.49 (m, 2H), 2.00 - 2.05 (m, 2H), 2.58 (q, 2H), 3.18 - 3.24 (m, 2H), 3.86 - 3.89 (m, 2H), 7.03 - 7.07 (m, 1H), 7.23 (d, 2H), 7.51 (d, 2H), 8.01 (d, 1H), 8.47 (s, 1H), 10.83 (s, 1H), 11.00 (s, 1H); CO ₂ H not seen.	451
627	4-methyl-1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid	1.41 (s, 3H), 1.66 (t, 2H), 2.22 - 2.26 (m, 2H), 3.41 (t, 2H), 4.06 - 4.11 (m, 2H), 7.21 (d, 1H), 7.64 (d, 2H), 7.93 (d, 2H), 8.20 (d, 1H), 8.69 (s, 1H), 11.21 (s, 1H), 11.41 (s, 1H); CO ₂ H not seen.	507
628	4-methyl-1-(5-{[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid	1.20 (s, 3H), 1.46 - 1.53 (m, 2H), 2.02 - 2.08 (m, 2H), 3.25 - 3.33 (m, 2H), 3.91 - 3.94 (m, 2H), 7.23 (d, 1H), 7.65 - 7.75 (m, 1H), 8.09 - 8.20 (m, 2H), 8.50 (s, 1H), 11.08 (s, 1H), 11.19 (s, 1H); CO ₂ H not seen.	477
629	4-methyl-1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-	1.19 (s, 3H), 1.40 - 1.47 (m, 2H), 2.00 - 2.04 (m, 2H), 3.15 - 3.22 (m, 2H), 3.83 - 3.88 (m, 2H), 6.98 (d, 1H), 7.76 - 7.82	491

	yl)piperidine-4-carboxylic acid	(m, 4H), 7.98 (d, 1H), 8.48 (s, 1H), 11.01 (s, 1H), 11.43 (s, 1H); CO ₂ H not seen.	
630	1-(5-(((5-([4-(4-chlorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid	1.25 (s, 3H), 1.49 - 1.55 (m, 2H), 2.06 - 2.11 (m, 2H), 3.27 - 3.33 (m, 2H), 3.93 - 3.98 (m, 2H), 7.05 (d, 2H), 7.17 - 7.21 (m, 3H), 7.47 (d, 2H), 7.70 (d, 2H), 8.10 (d, 1H), 8.55 (s, 1H), 11.08 (s, 1H), 11.13 (s, 1H); CO ₂ H not seen.	549


Example 631: Methyl 1-(5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]aminopyridin-2-yl)-4-methylpiperidine-4-carboxylate



- 5 To a suspension of methyl 1-(5-[[hydrazino(oxo)acetyl]amino]pyridin-2-yl)-4-methylpiperidine-4-carboxylate (Intermediate 109; 250 mg, 0.75 mmol) in anhydrous DMF (3 mL) was added 3,4-difluorophenylisothiocyanate (154 mg, 0.89 mmol) in anhydrous DMF (1 mL) and the resulting suspension heated to 45 °C for 1 hr. EDAC.HCl (172 mg, 0.89 mmol) was added and the reaction mixture heated to 85 °C for 2 hrs and then
- 10 allowed to cool to ambient temperature. Water (15 mL) was added and the suspension was filtered, washed with water (2 x 10 mL) and ether (15 mL) then dried to give the title product as a yellow solid (282 mg, 0.597 mmol, 79.6%);
- ¹H NMR δ 1.12 (s, 3H), 1.34 - 1.42 (m, 2H), 1.92 - 1.97 (m, 2H), 3.02 - 3.08 (m, 2H), 3.59 (s, 3H), 3.76 - 3.81 (m, 2H), 6.79 (d, 1H), 7.27 - 7.30 (m, 1H), 7.40 (q, 1H), 7.60 -
- 15 7.66 (m, 1H), 7.83 (d, 1H), 8.40 (s, 1H), 10.84 (s, 1H), 11.13 (s, 1H); MS m/e MH⁺ 473.

The following examples were prepared by the general procedure described above using the appropriate isothiocyanates, which were commercially available except 1-chloro-4-(4-isothiocyanatophenoxy)benzene which was made from 4-(4-chlorophenoxy)aniline and


1,1'-carbonylbis(1*H*-imidazole) as described in Barclay, Tristin K.; Santillan, Alejandro, Jr.; Tang, Liu Y.; Venkatesan, Hariharan; Wolin, Ronald L. PCT Int. Appl. (2005) WO 2005044810.



Example	R	¹ H NMR δ	MS m/e MH ⁺
632	R ¹ , R ³ =H, R ² =Et, methyl 1-{5-[(5-{[4- ethylphenyl]amino}-1,3,4- oxadiazol-2-yl)carbonyl] amino}pyridin-2-yl}-4- methylpiperidine-4- carboxylate	1.16 - 1.20 (m, 6H), 1.42 - 1.49 (m, 2H), 1.99 - 2.04 (m, 2H), 2.59 (q, 2H), 3.10 - 3.16 (m, 2H), 3.67 (s, 3H), 3.82 - 3.88 (m, 2H), 6.86 (d, 1H), 7.23 (d, 2H), 7.51 (d, 2H), 7.90 (d, 1H), 8.47 (s, 1H), 10.81 (s, 1H), 10.85 (s, 1H)	465
633	R ¹ , R ³ =H, R ² =OCF ₃ , methyl 4-methyl-1-(5-{[(5- {[4-(trifluoromethoxy) phenyl]amino}-1,3,4- oxadiazol-2-yl)carbonyl] amino}pyridin-2- yl)piperidine-4-carboxylate	1.19 (s, 3H), 1.42 - 1.48 (m, 2H), 2.00 - 2.04 (m, 2H), 3.10 - 3.17 (m, 2H), 3.66 (s, 3H), 3.83 - 3.88 (m, 2H), 6.86 (d, 1H), 7.42 (d, 2H), 7.71 (d, 2H), 7.90 (d, 1H), 8.47 (s, 1H), 10.90 (s, 1H), 11.17 (s, 1H)	521
634	R ¹ , R ² , R ³ =F, methyl 4-methyl-1-{5-[(5- [(2,4,5-trifluorophenyl) amino]-1,3,4-oxadiazol-2- yl)carbonyl]amino}pyridin-2- yl)piperidine-4-carboxylate	1.19 (s, 3H), 1.42 - 1.49 (m, 2H), 1.99 - 2.04 (m, 2H), 3.10 - 3.16 (m, 2H), 3.66 (s, 3H), 3.83 - 3.88 (m, 2H), 6.87 (d, 1H), 7.70 (q, 1H), 7.89 (d, 1H), 8.13 - 8.19 (m, 1H), 8.47 (s, 1H), 10.91 (s, 1H),	491

		11.02 (s, 1H)	
635	$R^1, R^2=H, R^3=CF_3$, methyl 4-methyl-1-(5-{{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate	1.20 (s, 3H), 1.41 - 1.49 (m, 2H), 2.00 - 2.05 (m, 2H), 3.09 - 3.17 (m, 2H), 3.66 (s, 3H), 3.82 - 3.88 (m, 2H), 6.86 (d, 1H), 7.76 - 7.82 (m, 4H), 7.90 (d, 1H), 8.47 (s, 1H), 10.93 (s, 1H), 11.41 (s, 1H)	505
636	$R^1, R^2=H, R^3=4$ -chlorophenoxy, methyl 1-(5-{{[5-{{[4-(4-chlorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl]amino}pyridin-2-yl}-4-methyl)piperidine-4-carboxylate	1.20 (s, 3H), 1.41 - 1.49 (m, 2H), 1.99 - 2.05 (m, 2H), 3.10 - 3.16 (m, 2H), 3.66 (s, 3H), 3.82 - 3.88 (m, 2H), 6.86 (d, 1H), 7.00 (d, 2H), 7.13 (d, 2H), 7.41 (d, 2H), 7.64 (d, 2H), 7.90 (d, 1H), 8.47 (s, 1H), 10.88 (s, 1H), 10.97 (s, 1H)	563

Example 637: 1-(5-{{[5-{{[4-(4-Chlorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylic acid




5

To a stirred solution of methyl 1-(5-{{[5-{{[4-(4-chlorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl}carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylate (Example 638; 340 mg, 0.62 mmol) in MeOH (5 mL) was added 2M NaOH (1.5 mL, 3.10 mmol) and the reaction mixture was allowed to stir at ambient temperature overnight. The solvent was evaporated and the aqueous residue adjusted to pH ~1-2 with 2M HCl, the solid was filtered and dried to give the title compound (284 mg, 0.531 mmol, 86%); ^1H NMR δ 1.79 - 1.90 (m, 2H), 2.14 - 2.19 (m, 2H), 2.78 - 2.84 (m, 1H), 3.33 - 3.40 (m, 2H), 4.35 - 4.39 (m, 2H), 7.23 (d, 2H), 7.37 (d, 2H), 7.43 (d, 1H), 7.64 (d, 2H), 7.88 (d, 2H), 8.33 (d, 1H), 8.73 (s, 1H), 11.26 (s, 1H), 11.36 (s, 1H); CO_2H not seen; MS m/e MH^+ 535.

10

Example 638: Methyl 1-(5-((5-([4-(4-chlorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylate




Prepared using the procedures described above for Example 631 except starting from

- 5 methyl 1-(5-([hydrazino(oxo)acetyl]amino)pyridin-2-yl)piperidine-4-carboxylate (Intermediate 83) and 1-chloro-4-(4-isothiocyanatophenoxy)benzene. ^1H NMR δ 1.51 - 1.61 (m, 2H), 1.86 - 1.91 (m, 2H), 2.59 - 2.66 (m, 1H), 2.91 - 2.98 (m, 2H), 3.63 (s, 3H), 4.14 - 4.19 (m, 2H), 6.87 (d, 1H), 7.00 (d, 2H), 7.13 (d, 2H), 7.41 (d, 2H), 7.64 (d, 2H), 7.90 (d, 1H), 8.47 (s, 1H), 10.88 (s, 1H), 10.97 (s, 1H); MS m/e MH^+ 549.

10

Example 639: [1-(5-((5-([4-(4-Chlorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetic acid




Prepared by the procedure described for 1-(5-((5-([4-(4-Chlorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidine-4-carboxylic acid (Example 637) except starting with methyl [1-(5-((5-([4-(4-chlorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetate (Example 640).

- 15 ^1H NMR δ 1.44 - 1.54 (m, 2H), 2.03 (d, 2H), 2.18 - 2.28 (m, 1H), 2.44 (d, 2H), 3.31 (t, 2H), 4.47 (d, 2H), 7.22 (d, 2H), 7.36 (d, 2H), 7.49 - 7.52 (m, 1H), 7.64 (d, 2H), 7.88 (d, 2H), 8.37 (d, 1H), 8.74 (s, 1H), 11.28 (s, 1H), 11.42 (s, 1H); CO_2H not seen; MS m/e MH^+ 549.

20

Example 640: Methyl [1-(5-((5-([4-(4-chlorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl)carbonyl]amino)pyridin-2-yl)piperidin-4-yl]acetate



25


- 286 -

207

Prepared using the procedures described above for Example 631 except using methyl [1-(5-{{[hydrazino(oxo)acetyl]amino}pyridin-2-yl]piperidin-4-yl}acetate (Intermediate 77) and 1-chloro-4-(4-isothiocyanatophenoxy)benzene.

¹H NMR δ 1.20 - 1.30 (m, 2H), 1.77 (d, 2H), 1.97 - 2.05 (m, 1H), 2.32 (d, 2H), 2.98 (t, 2H), 3.63 (s, 3H), 4.20 (d, 2H), 7.01 (d, 2H), 7.11 - 7.16 (m, 3H), 7.43 (d, 2H), 7.66 (d, 2H), 8.06 (d, 1H), 8.48 (s, 1H), 11.01 (s, 1H), 11.07 (s, 1H); MS m/e MH⁺ 563.


Example 641: rac-Ethyl 2-(trans-4-{2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)propanoate



10

Following the general procedure of Example 623, except that ethyl 2-{trans-4-(2-chloro-4-{{[hydrazino(oxo)acetyl]amino}phenyl}-cyclohexyl)propanoate (Intermediate 106) was used in place of methyl 2-[trans-4-(4-{{[hydrazino(oxo)acetyl]amino}-phenyl}cyclohexyl]-2-methylpropanoate (Intermediate 105), the title compound was obtained in 88% yield as a solid; ¹H NMR δ 1.05 (3H, d), 1.1-1.27 (2H, m), 1.18 (3H, t), 1.38-1.51 (2H, m), 1.52-1.65 (1H, m), 1.66-1.87 (4H, m), 2.22-2.35 (1H, m), 2.78-2.9 (1H, m), 4.09 (2H, dq), 7.39 (1H, d), 7.68-7.79 (1H, m), 7.72 (1H, dd), 7.93 (1H, d), 8.13-8.27 (1H, m), 11.18 (1H, s), 11.24 (1H, s); MS m/e MH⁺ 551.

Example 642: rac-Ethyl 2-(trans-4-{3-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)propanoate




Following the general procedure of Example 623, except that ethyl 2-[trans-4-(3-chloro-4-{{[hydrazino(oxo)acetyl]amino}phenyl}-cyclohexyl)propanoate (Intermediate 107) was used in place of methyl 2-[trans-4-(4-{{[hydrazino(oxo)acetyl]amino}-phenyl}cyclohexyl]-2-methylpropanoate (Intermediate 105), the title compound was obtained in 92% yield as a solid; ¹H NMR δ 1.05 (3H, d), 1.1-1.27 (2H, m), 1.18 (3H, t), 1.38-1.51 (2H, m), 1.52-1.65

25

- 287 -

(1H, m), 1.66-1.87 (4H, m), 2.22-2.35 (1H, m), 2.78-2.9 (1H, m), 4.09 (2H, dq), 7.2 (1H, dd), 7.38 (1H, d), 7.47 (1H, d), 7.62-7.75 (1H, m), 8.07-8.19 (1H, m), 11.2 (1H, s), 11.25 (1H, s); MS m/e MH⁺ 551.

5 Example 643: *rac*-2-(*trans*-4-{2-Chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid



Potassium trimethylsilanolate (1.29 g, 9.98 mmol) was added in one portion to a stirred solution of *rac*-ethyl 2-(*trans*-4-{2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoate (Example 641, 550 mg, 1.0 mmol) in THF (10 mL) and the reaction mixture was heated in the microwave at 90°C for 20 min. The reaction mixture was concentrated *in vacuo* to leave a yellow solid and then a 1M aqueous solution of citric acid (40 mL) was added and the mixture was stirred for 30 mins. The mixture was filtered to leave a solid, which was washed with H₂O (20 mL) to give the title compound (570 mg, 97 %) as an off-white solid. The individual isomers may be separated by chiral chromatography under standard conditions and recrystallised from EtOH/water to give

(2*R*)-2-(*trans*-4-{2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)-propanoic acid:


20 ¹H NMR δ 1.07 (3H, d), 1.1-1.31 (2H, m), 1.37-1.53 (2H, m), 1.54-1.67 (1H, m), 1.71-1.9 (4H, m), 2.15-2.27 (1H, m), 2.79-2.91 (1H, m), 7.39 (1H, d), 7.64-7.76 (1H, m), 7.69 (1H, dd), 7.94 (1H, d), 8.13-8.24 (1H, m), 11.5 (1H, s), 11.2 (1H, s), 12.1 (1H, s); MS m/e MH⁺ 523;

and (2*S*)-2-(*trans*-4-{2-chloro-4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)propanoic acid:

25 ¹H NMR δ 1.07 (3H, d), 1.1-1.31 (2H, m), 1.37-1.53 (2H, m), 1.54-1.67 (1H, m), 1.71-1.9 (4H, m), 2.15-2.27 (1H, m), 2.79-2.91 (1H, m), 7.39 (1H, d), 7.64-7.76 (1H, m), 7.69 (1H, dd), 7.94 (1H, d), 8.13-8.24 (1H, m), 11.5 (1H, s), 11.2 (1H, s), 12.1 (1H, s); MS m/e MH⁺ 523.


205

Example 644: *rac*-2-(*trans*-4-{3-Chloro-4-[(5-(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoic acid



Following the general procedure of Example 643, except that *rac*-ethyl 2-(*trans*-4-{3-chloro-4-[(5-(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)propanoate (Example 642) was used in place of *rac*-ethyl 2-(*trans*-4-{2-chloro-4-[(5-(2,4,5-trifluorophenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)-propanoate (Example 641), the title compound was obtained in 95% yield as an off-white solid; ¹H NMR δ 1.0 (3H, d), 1.02-1.22 (2H, m), 1.32-1.6 (3H, m), 1.63-1.84 (4H, m), 2.06-2.18 (1H, m), 2.4-2.53 (1H, m), 7.21 (1H, dd), 7.38 (1H, d), 7.47 (1H, d), 7.62-7.75 (1H, m), 8.07-8.19 (1H, m), 10.58 (1H, s), 11.1 (1H, s), 12.02 (1H, s); MS m/e MH⁺ 523. The individual isomers may be separated by chiral chromatography under standard conditions to give (2*R*)-2-(*trans*-4-{3-chloro-4-[(5-(2,4,5-trifluorophenyl)amino)-1,3,4-oxadiazol-2-yl]-carbonyl)amino]phenyl}cyclohexyl)propanoic acid and (2*S*)-2-(*trans*-4-{3-chloro-4-[(5-(2,4,5-trifluorophenyl)amino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenyl}cyclohexyl)-propanoic acid.

Example 645: Ethyl *cis*-4-{4-[(5-(2,4,5-trifluorophenyl)amino)-1,3,4-oxadiazol-2-yl]carbonyl)amino]phenoxy}cyclohexanecarboxylate




Hydrazine monohydrate (32 μL, 0.65 mmol) was added in one portion to a stirred solution of ethyl *cis*-4-(4-{[methoxy(oxo)acetyl]amino}phenoxy)cyclohexanecarboxylate (Intermediate 108) (188 mg, 0.54 mmol) in EtOH (3 mL) and the reaction mixture was stirred at room temperature for 10 mins. EtOH (40 mL) and hydrazine monohydrate (9.0 μL, 0.18 mmol) were added and the reaction mixture was stirred for a further 20 mins.

- 289 -

Diethylether (50 mL) was added and the mixture was filtered to leave a solid. The filtrate was concentrated *in vacuo* to leave a gum. The solid and the gum were combined to give the crude hydrazide that was used with no further purification; MS *m/e* (M-H)⁺ 348.


A solution of 2,4,5-trifluorophenyl isothiocyanate (103 mg, 0.54 mmol) in DMA (1 mL) was added in one portion to a stirred solution of the crude hydrazide (189 mg, 0.54 mmol) in DMA (4 mL) and the reaction mixture was stirred at room temperature for 1 h. EDCI (125 mg, 0.65 mmol) was added in one portion and the reaction mixture was heated to 60°C for 2 mins and then stirred at room temperature for 60 h. The reaction mixture was concentrated *in vacuo* to leave an oil. Hot water was added and the residue was stirred for 5 min to give a precipitate, which was filtered to give the title compound as a light brown solid (221 mg, 81%); ¹H NMR δ 1.20 (3H, t), 1.65 - 1.71 (4H, m), 1.76 - 1.85 (4H, m), 2.52 (1H, m), 4.09 (2H, q), 4.51 (1H, s), 6.96 (2H, d), 7.67 - 7.74 (3H, m), 8.15 - 8.18 (1H, m), 10.95 (1H, s), 11.05 (1H, s); MS *m/e* (M-H)⁺ 503.

Example 646: *cis*-4-{4-[(5-[(2,4,5-Trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]-phenoxy}cyclohexanecarboxylic acid




Lithium hydroxide monohydrate (184 mg, 4.38 mmol) was added in one portion to a solution of ethyl *cis*-4-{4-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]-phenoxy}cyclohexanecarboxylate (Example 645, 221 mg, 0.44 mmol) in a mixture of THF (4 mL), H₂O (4 mL) and MeOH (8 mL) and the reaction mixture was stirred at 35°C for 3 h. The reaction mixture was concentrated *in vacuo* and citric acid (40 mL) was added. The resulting mixture was filtered, washed with H₂O and recrystallised from ethanol/H₂O to give the title compound as a pale yellow solid (118 mg, 56%); ¹H NMR δ 1.66 - 1.70 (4H, m), 1.75 - 1.83 (4H, m), 2.33 - 2.40 (1H, m), 4.50 (1H, s), 6.96 (2H, d), 7.67 - 7.74 (3H, m), 8.17 (1H, q), 10.95 (1H, s), 11.10 (1H, s), 12.05 (1H, s); MS *m/e* (M-H)⁺ 475. The compound may be further recrystallised from EtOH, melting point 251-253°C.

The following examples were prepared by the general procedure as described for (1-{5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid (Example 619) using the appropriate starting material selected from Examples 649 and 650.



5


Example	R	Name	¹ H NMR δ	MS m/e M+H ⁺
647	Et	(1-{5-[(5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetic acid	1.17 - 1.27 (m, 5H), 1.77 (d, H), 1.90 - 2.00 (m, 1H), 2.20 (d, 2H), 2.56 - 2.62 (m, 2H), 2.92 (t, 2H), 4.20 (d, 2H), 7.04 (d, 1H), 7.23 (d, 2H), 7.51 (d, 2H), 8.00 (d, 1H), 8.46 (s, 1H), 10.83 (s, 1H), 10.99 (s, 1H); CO ₂ H not seen.	451
648	CF ₃	[1-(5-[(5-[(4-(trifluoromethyl)phenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid	1.17 - 1.26 (m, 2H), 1.77 (d, 2H), 1.90 - 2.00 (m, 1H), 2.19 (d, 2H), 2.91 (t, 2H), 4.21 (d, 2H), 7.03 (d, 1H), 7.76 - 7.83 (m, 4H), 8.00 (d, 1H), 8.46 (s, 1H), 11.04 (s, 1H), 11.43 (s, 1H); CO ₂ H not seen	491



The following examples were prepared by the general procedure as described for methyl (1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetate (Example 620) using the appropriate isothiocyanate:

Example	R	Name	¹ H NMR δ	MS m/e M+H ⁺
649	Et	methyl (1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetate	1.16 - 1.21 (m, 5H), 1.69 - 1.73 (m, 2H), 1.89 - 1.98 (m, 1H), 2.28 (d, 2H), 2.76 - 2.83 (m, 2H), 3.61 (s, 3H), 4.22 - 4.25 (m, 2H), 4.31 (t, 2H), 6.84 (d, 1H), 7.23 (d, 2H), 7.51 (d, 2H), 7.89 (d, 1H), 8.46 (d, 1H), 10.81 (s, 1H), 10.85 (s, 1H)	465
650	CF ₃	methyl [1-(5-{[(5-{(4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl)acetate	1.14 - 1.25 (m, 2H), 1.71 (d, 2H), 1.89 - 1.99 (m, 1H), 2.28 (d, 2H), 2.77 - 2.84 (m, 2H), 3.61 (s, 3H), 4.24 (d, 2H), 6.85 (d, 1H), 7.76 - 7.82 (m, 4H), 7.89 (d, 1H), 8.46 (s, 1H), 10.92 (s, 1H), 11.42 (s, 1H)	505

- 5 The following examples were prepared by the general procedure as described previously as in Example 480:




Example	R	Name	¹ H NMR δ	MS m/e M+H ⁺
651	Et	1-{5-[(5-{(4-	1.39 (t, 3H), 1.74 - 1.84 (m,	437

- 292 -

		ethylphenylamino]-1,3,4-oxadiazol-2-yl}carbonyl]amino]pyridin-2-yl]piperidine-4-carboxylic acid	2H), 2.09 - 2.13 (m, 2H), 2.75 - 2.82 (m, 3H), 3.25 (t, 2H), 4.31 - 4.36 (m, 2H), 7.25 (d, 1H), 7.43 (d, 2H), 7.72 (d, 2H), 8.22 (d, 1H), 8.68 (d, 1H), 11.04 (s, 1H), 11.20 (s, 1H); CO ₂ H not seen.	
652	OCF ₃	1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid	1.61 - 1.70 (m, 2H), 1.96 - 2.00 (m, 2H), 2.62 - 2.64 (m, 1H), 3.11 (t, 2H), 4.18 - 4.23 (m, 2H), 7.10 (d, 1H), 7.50 (d, 2H), 7.79 (d, 2H), 8.08 (d, 1H), 8.56 (d, 1H), 11.10 (s, 1H), 11.26 (s, 1H); CO ₂ H not seen.	493
653	CF ₃	1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid	1.52 - 1.62 (m, 2H), 1.88 - 1.92 (m, 2H), 2.54 - 2.56 (m, 1H), 2.98 - 3.05 (m, 2H), 4.11 - 4.16 (m, 2H), 6.99 (d, 1H), 7.76 - 7.82 (m, 4H), 7.98 (d, 1H), 8.48 (d, 1H), 11.02 (s, 1H), 11.43 (s, 1H); CO ₂ H not seen	477

The following examples were prepared by the general procedure as described previously, as in Example 620, using the appropriate isothiocyanate:



Example	R	Name	¹ H NMR δ	MS m/e M+H ⁺
654	Et	methyl 1-{5-[(5-{[4-ethylphenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl}piperidine-4-carboxylate	1.18 (t, 3H), 1.51 - 1.61 (m, 2H), 1.86 - 1.91 (m, 2H), 2.56 - 2.62 (m, 3H), 2.91 - 2.98 (m, 2H), 3.63 (s, 3H), 4.14 - 4.18 (m, 2H), 6.87 (d, 1H), 7.23 (d, 2H), 7.51 (d, 2H), 7.91 (dd, 1H), 8.48 (d, 1H), 10.81 (s, 1H), 10.86 (s, 1H)	451
655	OCF ₃	methyl 1-(5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate	1.51 - 1.61 (m, 2H), 1.86 - 1.91 (m, 2H), 2.59 - 2.65 (m, 1H), 2.91 - 2.98 (m, 2H), 3.63 (s, 3H), 4.14 - 4.18 (m, 2H), 6.88 (d, 1H), 7.42 (d, 2H), 7.71 (d, 2H), 7.91 (dd, 1H), 8.48 (d, 1H), 10.91 (s, 1H), 11.17 (s, 1H)	507
656	CF ₃	methyl 1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate	1.51 - 1.61 (m, 2H), 1.87 - 1.91 (m, 2H), 2.59 - 2.66 (m, 1H), 2.91 - 2.98 (m, 2H), 3.63 (s, 3H), 4.14 - 4.18 (m, 2H), 6.88 (d, 1H), 7.76 - 7.82 (m, 4H), 7.91 (dd, 1H), 8.49 (d, 1H), 10.94 (s, 1H), 11.41 (s, 1H)	491

Examples 657-679

The following examples were prepared by the general procedure of Example 525 using the appropriate commercially available aniline in place of 5-amino-3-methylisoxazole.

Example	Name	¹ H NMR δ	MS m/e MH ⁺
657 ¹	(<i>trans</i> -4-{4-[(5-{(2-chloro-4-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid	0.97-1.12 (m, 2H), 1.29-1.46 (m, 2H), 1.60-1.80 (m, 5H), 2.07 (d, 2H), 2.24 (s, 3H), 2.31-2.41 (m, 1H), 7.15 (m, 3H), 7.31 (s, 1H), 7.60 (d, 2H), 7.69 (d, 1H), 10.20 (s, 1H), 11.51 (s, 1H), 11.82 (s, 1H)	469
658 ¹	(<i>trans</i> -4-{4-[(5-{(4-methoxy-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid	1.04-1.19 (m, 2H), 1.39-1.52 (m, 2H), 1.66-1.87 (m, 5H), 2.14 (d, 2H), 2.25 (s, 3H), 2.38-2.48 (m, 1H), 3.76 (s, 3H), 6.80-6.84 (m, 1H), 6.85-6.88 (m, 1H), 7.22 (d, 2H), 7.47 (d, 1H), 7.67 (d, 2H), 9.82 (s, 1H), 11.61 (s, 1H), 11.95 (s, 1H)	465
659 ¹	(<i>trans</i> -4-{4-[(5-{(4-fluoro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid	0.97-1.11 (m, 2H), 1.30-1.44 (m, 2H), 1.60-1.79 (m, 5H), 2.07 (d, 2H), 2.22 (s, 3H), 2.31-2.41 (m, 1H), 6.97-7.11 (m, 2H), 7.15 (d, 2H), 7.56-7.65 (m, 3H), 9.93 (s, 1H), 10.78 (s, 1H), 11.88 (s, 1H)	453
660 ¹	[<i>trans</i> -4-(4-{[(5-{[4-fluoro-2-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl]acetic acid	1.04-1.20 (m, 2H), 1.38-1.54 (m, 2H), 1.66-1.88 (m, 5H), 2.14 (d, 2H), 2.38-2.49 (m, 1H), 7.21 (d, 2H), 7.61-7.75 (m, 4H), 7.84-7.94 (m, 1H), 10.35 (s, 1H), 10.87 (s, 1H), 11.95 (s, 1H)	507

661 ¹	(<i>trans</i> -4-{4-[(5-{(2,4-dichlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	0.97-1.12 (m, 2H), 1.30-1.44 (m, 2H), 1.59-1.81 (m, 5H), 2.07 (d, 2H), 2.31-2.41 (m, 1H), 7.15 (d, 2H), 7.44 (d, 1H), 7.57-7.66 (m, 3H), 7.95 (d, 1H), 10.45 (s, 1H), 10.85 (s, 1H), 11.89 (s, 1H)	489
662 ¹	(<i>trans</i> -4-{4-[(5-{(2-chloro-4-cyanophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.04-1.21 (m, 2H), 1.37-1.53 (m, 2H), 1.65-1.89 (m, 5H), 2.15 (d, 2H), 2.40-2.49 (m, 1H), 7.23 (d, 2H), 7.69 (d, 2H), 7.89-7.95 (m, 1H), 8.12 (d, 1H), 8.36 (d, 1H), 10.89 (s, 1H), 11.00 (s, 1H), 11.94 (s, 1H)	480
663 ¹	(<i>trans</i> -4-{4-[(5-{(4-cyano-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.02-1.20 (m, 2H), 1.37-1.54 (m, 2H), 1.66-1.88 (m, 5H), 2.13 (d, 2H), 2.31-2.44 (m, 1H), 7.23 (d, 2H), 7.64-7.81 (m, 4H), 8.13 (d, 1H), 10.46 (s, 1H), 10.99 (s, 1H), 11.99 (s, 1H)	460
664 ¹	(<i>trans</i> -4-{4-[(5-{(4-chloro-2,5-dimethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	0.96-1.12 (m, 2H), 1.30-1.45 (m, 2H), 1.59-1.80 (m, 5H), 2.06 (d, 2H), 2.18 (s, 3H), 2.24 (s, 3H), 2.32-2.40 (m, 1H), 7.14 (d, 2H), 7.25 (s, 1H), 7.57-7.65 (m, 3H), 9.97 (s, 1H), 10.78 (s, 1H), 11.87 (s, 1H)	483
665 ¹	(<i>trans</i> -4-{4-[(5-{(4-cyano-2-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.24 (m, 5H), 1.37-1.54 (m, 2H), 1.66-1.89 (m, 5H), 2.15 (d, 2H), 2.39-2.48 (m, 1H), 2.80 (q, 2H), 7.23 (d, 2H), 7.65-7.79 (m, 4H), 8.18 (d, 1H), 10.45 (s, 1H), 10.97 (s, 1H), 11.96 (s, 1H)	474

207

666 ¹	(<i>trans</i> -4-{4-[(5-{(2-chloro-4-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.01-1.22 (m, 2H), 1.34-1.55 (m, 2H), 1.63-1.89 (m, 5H), 2.14 (d, 2H), 2.38-2.48 (m, 1H), 7.22 (d, 2H), 7.30-7.41 (m, 1H), 7.54-7.63 (m, 1H), 7.70 (d, 2H), 7.90-8.01 (m, 1H), 10.47 (s, 1H), 10.97 (s, 1H), 12.04 (s, 1H)	473
667 ¹	(<i>trans</i> -4-{4-[(5-{(4-chloro-2-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	0.97-1.13 (m, 2H), 1.31-1.47 (m, 2H), 1.60-1.81 (m, 5H), 2.08 (d, 2H), 2.23 (s, 3H), 2.32-2.41 (m, 1H), 7.16 (d, 2H), 7.23-7.36 (m, 2H), 7.62 (d, 2H), 7.76 (d, 1H), 10.09 (s, 1H), 10.90 (s, 1H), 11.97 (s, 1H)	469
668	(<i>trans</i> -4-{4-[(5-{(2,3-dichlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.02-1.19 (m, 2H), 1.35-1.54 (m, 2H), 1.65-1.88 (m, 5H), 2.14 (d, 2H), 2.38-2.48 (m, 1H), 7.22 (d, 2H), 7.40-7.50 (m, 2H), 7.69 (d, 2H), 7.97-8.04 (m, 1H), 10.68 (s, 1H), 11.01 (s, 1H), 12.05 (s, 1H)	489
669	(<i>trans</i> -4-{4-[(5-{(2,5-dichlorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.18 (m, 2H), 1.38-1.53 (m, 2H), 1.65-1.87 (m, 5H), 2.14 (d, 2H), 2.38-2.48 (m, 1H), 7.19-7.30 (m, 3H), 7.58 (d, 1H), 7.69 (d, 2H), 8.16-8.21 (m, 1H), 10.69 (s, 1H), 11.02 (s, 1H), 12.05 (s, 1H)	489
670	(<i>trans</i> -4-{4-[(5-{(2-bromo-4-methylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.18 (m, 2H), 1.38-1.51 (m, 2H), 1.66-1.87 (m, 5H), 2.14 (d, 2H), 2.32 (s, 3H), 2.38-2.48 (m, 1H), 7.21 (d, 2H), 7.28 (d, 1H), 7.56 (s, 1H), 7.63-7.72 (m, 3H), 10.24 (s, 1H), 10.94 (s, 1H), 12.04 (s, 1H)	515

204


671	[<i>trans</i> -4-(4-{{(5-{{(2-chloro-5-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid	1.02-1.19 (m, 2H), 1.37-1.53 (m, 2H), 1.66-1.88 (m, 5H), 2.14 (d, 2H), 2.38-2.49 (m, 1H), 7.24 (d, 2H), 7.55 (d, 1H), 7.70 (d, 2H), 7.81 (d, 1H), 8.51 (s, 1H), 10.84 (s, 1H), 11.01 (s, 1H), 12.04 (s, 1H)	523
672	[<i>trans</i> -4-(4-{{(5-{{(2-bromo-4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid	0.96-1.11 (m, 2H), 1.29-1.46 (m, 2H), 1.57-1.81 (m, 5H), 2.07 (d, 2H), 2.30-2.41 (m, 1H), 7.16 (d, 2H), 7.63 (d, 2H), 7.80 (d, 1H), 8.02 (s, 1H), 8.17 (d, 1H), 10.96 (s, 1H), 10.96 (s, 1H), 11.97 (s, 1H)	567
673	[<i>trans</i> -4-(4-{{(5-{{(2-chloro-4-(trifluoromethyl)phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid	1.01-1.20 (m, 2H), 1.34-1.53 (m, 2H), 1.64-1.89 (m, 5H), 2.14 (d, 2H), 2.36-2.48 (m, 1H), 7.23 (d, 2H), 7.70 (d, 2H), 7.84 (d, 1H), 7.96 (s, 1H), 8.37 (d, 1H), 10.81 (s, 1H), 11.05 (s, 1H), 12.04 (s, 1H)	523
674	(<i>trans</i> -4-{4-[(5-[(4-bromo-2-chlorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid	1.03-1.20 (m, 2H), 1.37-1.53 (m, 2H), 1.65-1.88 (m, 5H), 2.14 (d, 2H), 2.37-2.48 (m, 1H), 7.23 (d, 2H), 7.62-7.73 (m, 3H), 7.83 (s, 1H), 7.99 (d, 1H), 10.58 (s, 1H), 11.00 (s, 1H), 12.04 (s, 1H)	535
675	(<i>trans</i> -4-{4-[(5-[(3-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl}amino}phenyl)cyclohexyl}acetic acid	1.03-1.18 (m, 2H), 1.37-1.52 (m, 2H), 1.66-1.86 (m, 5H), 2.14 (d, 2H), 2.34 (s, 3H), 2.39-2.49 (m, 1H), 7.11-7.25 (m, 3H), 7.27-7.34 (m, 1H), 7.65-7.75 (m, 3H), 10.31 (s, 1H), 10.96 (s, 1H), 12.03 (s, 1H)	469

676	(<i>trans</i> -4-{4-[(5-{(5-bromo-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.19 (m, 2H), 1.37-1.51 (m, 2H), 1.65-1.88 (m, 5H), 2.14 (d, 2H), 2.28 (s, 3H), 2.37-2.48 (m, 1H), 7.17-7.30 (m, 4H), 7.69 (d, 2H), 8.08 (s, 1H), 10.24 (s, 1H), 10.98 (s, 1H), 12.02 (s, 1H)	513
677	(<i>trans</i> -4-{4-[(5-{(2-chloro-5-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.20 (m, 2H), 1.38-1.53 (m, 2H), 1.64-1.87 (m, 5H), 2.15 (d, 2H), 2.37-2.49 (m, 1H), 7.03-7.10 (m, 1H), 7.23 (d, 2H), 7.56-7.63 (m, 1H), 7.69 (d, 2H), 7.97-8.05 (m, 1H), 10.69 (s, 1H), 11.03 (s, 1H), 12.07 (s, 1H)	473
678 ¹	(<i>trans</i> -4-{4-[(5-{(2-chloro-5-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.04-1.20 (m, 2H), 1.38-1.52 (m, 2H), 1.67-1.87 (m, 5H), 2.14 (d, 2H), 2.34 (s, 3H), 2.39-2.49 (m, 1H), 7.05 (d, 1H), 7.23 (d, 2H), 7.42 (d, 1H), 7.68 (d, 2H), 7.75 (s, 1H), 10.37 (s, 1H), 10.95 (s, 1H), 12.02 (s, 1H)	469
679 ¹	(<i>trans</i> -4-{4-[(5-{(5-chloro-2-methylphenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]phenyl}cyclohexyl)acetic acid	1.03-1.19 (m, 2H), 1.37-1.52 (m, 2H), 1.67-1.87 (m, 5H), 2.14 (d, 2H), 2.29 (s, 3H), 2.38-2.48 (m, 1H), 7.13 (d, 1H), 7.22 (d, 2H), 7.28 (d, 1H), 7.69 (d, 2H), 7.97 (s, 1H), 10.25 (s, 1H), 10.99 (s, 1H), 12.02 (s, 1H)	469



¹The product was recrystallised from acetic acid

Example 680

The following example was prepared by the general procedure of Example 483 using the appropriate aniline R¹NH₂ and isothiocyanate R²-NCS.




20/0

Example	SM	R ¹	R ²	¹ H NMR δ	MS m/e MH ⁺
680	Int 110			1.02-1.10 (3H, m), 1.17-1.22 (3H, m), 2.16-2.22 (3H, m), 2.33-2.68 (3H, m), 3.50 (1H, m), 4.06-4.11 (2H, m), 7.21-7.28 (2H, m), 7.36 (1H, d), 7.47-7.50 (1H, m), 7.68-7.75 (3H, m), 10.95-10.97 (1H, m), 11.23 (1H, s)	471

Example 681


The following example was prepared by the general procedure of Example 498, using Example 680 as starting material (SM).



Example	¹ H NMR δ	MS m/e MH ⁺
681	0.98-1.12 (3H, m), 1.71-1.92 (1H, m), 2.13-2.24 (3H, m), 2.28-2.44 (2H, m), 3.42-3.53 (1H, m), 7.21-7.28 (2H, m), 7.35-7.40 (1H, m), 7.45-7.50 (1H, m), 7.65-7.75 (3H, m), 10.96 (1H, d), 11.23 (1H, s), 12.01 (1H, s)	443

5

Example 682: Methyl 1-(5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino)-3-methylpyridin-2-yl)piperidine-4-carboxylate



To a stirred suspension of methyl 1-(5-[(hydrazino(oxo)acetyl]amino)-3-methylpyridin-2-yl)piperidine-4-carboxylate (Intermediate 111) (336 mg; 1.00 mmol) in anhydrous DMA (5 mL) was added 3,4-difluorophenylisothiocyanate (206 mg; 1.20 mmol) and the clear solution, obtained on warming, was heated to 45°C for 1hr. EDAC (231 mg) was added


10

- 300 -

24X

and the reaction heated at 85 °C for 1 hr. The reaction mixture was cooled and water (~5 mL) was added. The precipitated solid was filtered off and dried under vacuum at 60 °C to leave the title compounds as a yellow powder (381mg, 81% yield). ¹H NMR δ 1.62 - 1.81 (m, 2H), 1.83 - 2.00 (m, 2H), 2.16 (s, 3H), 2.51 - 2.59 (m, 1H), 2.65 - 2.84 (m, 2H), 3.23 - 3.41 (m, 2H), 3.70 (s, 3H), 7.28 - 7.40 (m, 1H), 7.39 - 7.57 (m, 1H), 7.59 - 7.76 (m, 1H), 7.90 (d, 1H), 8.44 (d, 1H), 10.97 (s, 1H), 11.16 (s, 1H). MS m/e MH⁺ = 473.

In a similar manner, from the appropriate hydrazides (Intermediates 111 to 113) and the commercially available isothiocyanates, the following example were synthesised.




10

Example	Name	¹ H NMR δ	MS m/e M+H ⁺
683 R = 4-OCF ₃	Methyl 1-(3-methyl-5-{{(5- [[4-trifluoromethoxy]phenyl]amino -1,3,4-oxadiazol-2- yl)carbonyl]amino}pyridin-2- yl)piperidine-4-carboxylate	1.62 - 1.79 (m, 2H), 1.83 - 1.97 (m, 2H), 2.25 (s, 3H), 2.44 - 2.52 (m, 1H), 2.66 - 2.81 (m, 2H), 3.19 - 3.37 (m, 2H), 3.66 (s, 3H), 7.40 (d, 2H), 7.69 (d, 2H), 7.90 (d, 1H), 8.43 (d, 1H), 10.98 (s, 1H), 11.16 (s, 1H)	521
684 R = 4-CF ₃	Methyl 1-(3-methyl-5-{{(5- [[4-trifluoromethyl]phenyl]amino} -1,3,4-oxadiazol-2- yl)carbonyl]amino}pyridin-2- yl)piperidine-4-carboxylate	1.61 - 1.82 (m, 2H), 1.85 - 1.99 (m, 2H), 2.18 (s, 3H), 2.52 - 2.58 (m, 1H), 2.68 - 2.83 (m, 2H), 3.24 - 3.42 (m, 2H), 3.58 (s, 3H), 7.62 - 7.88 (m, 4H), 7.92 (d, 1H), 8.45 (d, 1H), 11.00 (s, 1H), 11.39 (s, 1H)	505

- 301 -


685 R = 4-Et	Methyl 1-{5-[(5-{(4-ethylphenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]-3-methylpyridin-2-yl}piperidine-4-carboxylate	1.16 (t, 3H), 1.59 - 1.80 (m, 2H), 1.86 - 1.99 (m, 2H), 2.19 (s, 3H), 2.51 - 2.64 (m, 3H), 2.68 - 2.83 (m, 2H), 3.31 (d, 2H), 3.58 (s, 3H), 7.22 (d, 2H), 7.50 (d, 2H), 7.90 (d, 1H), 8.44 (d, 1H), 10.83 (s, 1H), 10.95 (s, 1H)	465
-----------------	---	--	-----



Example	Name	¹ H NMR δ	MS m/e M+H ⁺
686 R = 3,4-diF	Methyl [1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]-3-methylpyridin-2-yl}piperidin-4-yl]acetate	1.24 - 1.45 (m, 2H), 1.66 - 1.78 (m, 2H), 1.78 - 1.92 (m, 1H), 2.18 (s, 3H), 2.31 (d, 2H), 2.60 - 2.75 (m, 2H), 3.25 - 3.39 (m, 2H), 3.56 (s, 3H), 7.28 - 7.39 (m, 1H), 7.39 - 7.53 (m, 1H), 7.62 - 7.74 (m, 1H), 7.89 (d, 1H), 8.43 (d, 1H), 11.06 (s, 1H), 11.30 (s, 1H)	487
687 R = 4-OCF ₃	Methyl [1-(3-methyl-5-{[(5-{(4-trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate	1.24 - 1.43 (m, 2H), 1.63 - 1.79 (m, 2H), 1.79 - 1.92 (m, 1H), 2.25 (s, 3H), 2.32 (d, 2H), 2.61 - 2.75 (m, 2H), 3.22 - 3.39 (m, 2H), 3.65 (s, 3H), 7.40 (d, 2H), 7.69 (d, 2H), 7.90 (d, 1H), 8.43 (d, 1H), 10.98 (s, 1H), 11.23 (s,	535

- 302 -

		1H).	
688 R = 4-CF ₃	Methyl [1-(3-methyl-5-({(5-{[4-trifluoromethyl]phenyl}amino)-1,3,4-oxadiazol-2-yl}carbonyl)amino)pyridin-2-yl)piperidin-4-yl]acetate	1.23 - 1.48 (m, 2H), 1.66 - 1.79 (m, 2H), 1.78 - 1.93 (m, 1H), 2.17 (s, 3H), 2.32 (d, 2H), 2.59 - 2.75 (m, 2H), 3.24 - 3.41 (m, 2H), 3.70 (s, 3H), 7.61 - 7.84 (m, 4H), 7.90 (d, 1H), 8.43 (d, 1H), 10.98 (s, 1H), 11.48 (s, 1H)	519
689 R = 4-Et	Methyl (1-{5-[(5-{[4-ethylphenyl]amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]-3-methylpyridin-2-yl}piperidin-4-yl)acetate	1.17 (t, 3H), 1.24 - 1.44 (m, 2H), 1.67 - 1.78 (m, 2H), 1.79 - 1.90 (m, 1H), 2.16 (s, 3H), 2.31 (d, 2H), 2.57 (q, 2H), 2.62 - 2.75 (m, 2H), 3.22 - 3.39 (m, 2H), 3.67 (s, 3H), 7.21 (d, 2H), 7.49 (d, 2H), 7.89 (d, 1H), 8.43 (d, 1H), 10.81 (s, 1H), 11.02 (s, 1H)	479




Example	Name	¹ H NMR δ	MS m/e M+H ⁺
690 R = 3,4-diF	Ethyl 2-(1-{5-[(5-{[3,4-difluorophenyl]amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino]-3-methylpyridin-2-yl}piperidin-4-yl)propanoate	1.08 (d, 3H), 1.19 (t, 3H), 1.29 - 1.45 (m, 2H), 1.55 - 1.67 (m, 2H), 1.67 - 1.79 (m, 1H), 1.92 (s, 3H), 2.10 - 2.23 (m, 2H), 2.24 - 2.34 (m, 1H), 2.56 - 2.73 (m, 2H), 3.21 - 3.46 (m, 3H), 4.02 - 4.13 (m, 2H), 7.27	515

- 303 -

		- 7.38 (m, 1H), 7.38 - 7.53 (m, 1H), 7.61 - 7.75 (m, 1H), 7.89 (d, 1H), 8.43 (d, 1H), 11.37 (s, 1H).	
691 R = 4-OCF ₃	Ethyl 2-[1-(3-methyl-5-[[5- [[4-trifluoromethyl]phenyl]amino]- 1,3,4-oxadiazol-2- yl]carbonyl]amino}pyridin-2- yl)piperidin-4-yl]propanoate	1.08 (d, 3H), 1.19 (t, 3H), 1.29 - 1.47 (m, 2H), 1.51 - 1.68 (m, 2H), 1.67 - 1.79 (m, 1H), 2.24 (s, 3H), 2.23 - 2.35 (m, 1H), 2.55 - 2.76 (m, 2H), 3.22 - 3.44 (m, 3H), 4.07 (q, 2H), 7.39 (d, 2H), 7.67 (d, 2H), 7.87 (d, 1H), 8.40 (d, 1H), 11.05 (s, 1H)	563
692 R = 2,4,5-triF	Ethyl 2-[1-(3-methyl-5-[[5- [(2,4,5-trifluorophenyl)amino]- 1,3,4-oxadiazol-2- yl]carbonyl]amino]pyridine-2- yl)piperidin-4-yl]propanoate	1.06 (d, 3H), 1.20 (q, 3H), 1.25 - 1.48 (m, 2H), 1.51 - 1.67 (m, 2H), 1.68 - 1.78 (m, 1H), 2.14 (s, 3H), 2.24 - 2.38 (m, 1H), 2.53 - 2.73 (m, 2H), 3.20 - 3.45 (m, 3H), 4.08 (q, 2H), 7.67 (q, 1H), 7.88 (d, 1H), 8.15 (q, 1H), 8.43 (d, 1H), 10.98 (s, 1H)	533
693 R = 4-Et	Ethyl 2-[1-(5-[[5-[(4-ethylphenyl)amino]-1,3,4-oxadiazol-2-yl]carbonyl]amino]-3-methylpyridin-2-yl)piperidin-4-yl]propanoate	1.08 (d, 3H), 1.19 (q, 6H), 1.35 (m, 2H), 1.60 (m, 2H), 1.73 (m, 1H), 2.20 (s, 3H), 2.30 (m, 1H), 2.58 (q, 2H), 2.60 (m, 2H), 3.37 (m, 2H), 4.07 (q, 2H), 7.20 (d, 2H), 7.48 (d, 2H), 7.85 (d, 1H), 8.42 (d, 2H), 10.84 (s, 1H), 10.97 (s, 1H)	507

- 304 -

The following examples were prepared from the appropriate esters (Examples 682-689) by hydrolysis following the method of Example 320.




Example	Name	¹ H NMR δ	MS m/e M+H ⁺
694 R = 3,4-diF	(1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]-3-methylpyridin-2-yl}piperidin-4-yl)acetic acid	1.18 - 1.41 (m, 2H), 1.61 - 1.77 (m, 2H), 1.75 - 1.89 (m, 1H), 2.13 (d, 2H), 2.28 (s, 3H), 2.69 - 2.91 (m, 2H), 3.31 - 3.48 (m, 2H), 7.22 - 7.32 (m, 1H), 7.32 - 7.47 (m, 1H), 7.56 - 7.68 (m, 1H), 7.97 (s, 1H), 8.43 (d, 1H), 11.13 (s, 1H), 11.31 (s, 1H)	473
695 R = 4-OCF ₃	[1-(3-methyl-5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid	1.27 - 1.48 (m, 2H), 1.70 - 1.85 (m, 2H), 1.84 - 1.99 (m, 1H), 2.21 (d, 2H), 2.35 (s, 3H), 2.79 - 3.02 (m, 2H), 3.42 - 3.54 (m, 2H), 7.41 (d, 2H), 7.70 (d, 2H), 8.05 (s, 1H), 8.44 (s, 1H), 11.14 (s, 2H)	521
696 R = CF ₃	[1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid	1.27 - 1.49 (m, 2H), 1.71 - 1.85 (m, 2H), 1.85 - 1.98 (m, 1H), 2.22 (d, 2H), 2.34 (s, 3H), 2.81 - 3.00 (m, 2H), 3.43 - 3.54 (m, 2H), 7.78 (d, 4H), 8.06 (s, 1H), 8.51 (d, 1H), 11.23 (s, 1H), 11.55 (s, 1H)	505

- 305 -

312


697 R = 4-Et	(1-{5-[(5-{[4-ethylphenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl)acetic acid	1.16 (t, 3H), 1.27 - 1.46 (m, 2H), 1.72 - 1.85 (m, 2H), 1.85 - 1.94 (m, 1H), 2.21 (d, 2H), 2.35 (s, 3H), 2.57 (q, 2H), 2.80 - 2.96 (m, 2H), 3.43 - 3.54 (m, 2H), 7.21 (d, 2H), 7.51 (d, 2H), 8.03 (s, 1H), 8.50 (d, 1H), 10.86 (s, 1H), 11.26 (s, 1H)	465
-----------------	---	---	-----



Example	Name	¹ H NMR δ	MS m/e M+H ⁺
698 R = 3,4-diF	1-{5-[(5-{[3,4-difluorophenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylic acid	1.55 - 1.73 (m, 2H), 1.78 - 1.92 (m, 2H), 2.25 (s, 3H), 2.30 - 2.39 (m, 1H), 2.73 - 2.92 (m, 2H), 3.29 - 3.43 (m, 2H), 7.19 - 7.32 (m, 1H), 7.32 - 7.45 (m, 1H), 7.54 - 7.71 (m, 1H), 8.01 (s, 1H), 8.41 (d, 1H), 11.08 (s, 1H), 11.18 (s, 1H), 11.30 (s, 1H)	459
699 R = OCF ₃	1-(3-methyl-5-{[(5-{[4-(trifluoromethoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid	1.64 - 1.82 (m, 2H), 1.86 - 2.01 (m, 2H), 2.32 (s, 3H), 2.38 - 2.45 (m, 1H), 2.77 - 2.97 (m, 2H), 3.32 - 3.48 (m, 2H), 7.38 (d, 2H), 7.65 - 7.77 (m, 2H), 8.09 (s, 1H), 8.49 (d, 1H), 11.14 (s, 1H), 11.26 (s, 1H)	507

		1H), 11.31 (s, 1H)	
700 R = CF ₃	1-(3-methyl-5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid	1.55 - 1.74 (m, 2H), 1.80 - 1.93 (m, 2H), 2.16 (s, 3H), 2.33 - 2.39 (m, 1H), 2.76 - 2.92 (m, 2H), 3.28 - 3.43 (m, 2H), 7.62 - 7.79 (m, 4H), 7.95 (s, 1H), 8.44 (d, 1H), 11.12 (s, 1H), 11.38 (s, 1H), 11.51 (s, 1H)	491
701 R = 4-Et	1-{5-[(5-{[4-ethylphenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylic acid	1.17 (t, 3H), 1.59 - 1.82 (m, 2H), 1.87 - 1.99 (m, 2H), 2.23 - 2.33 (m, 3H), 2.37 - 2.47 (m, 1H), 2.59 (q, 2H), 2.79 - 2.93 (m, 2H), 3.34 - 3.47 (m, 2H), 7.21 (d, 2H), 7.45 - 7.55 (m, 2H), 7.97 (s, 1H), 8.44 (s, 1H), 10.84 (s, 1H), 10.94 (s, 1H), 11.16 (s, 1H)	451

Example 702: 1-(5-{[(5-{[4-(4-Fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylic acid



- 5 To a stirred solution of methyl 1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidine-4-carboxylate (Example 705, 415 mg, 0.78 mmol) in MeOH (5 mL) was added 2M NaOH (2 mL, 4 mmol) and the reaction mixture was allowed to stir at ambient temperature overnight. The solvent was evaporated and the aqueous residue adjusted to pH ~1-2 with 2M HCl, the solid was filtered and dried
- 10 to leave a cream solid (284 mg). Glacial AcOH (~40 mL) was added and the suspension heated to 150 °C, a small amount of material still didn't dissolve so the suspension was


- 307 -

filtered and allowed to cool. The resulting solid was filtered and dried under vacuum at 40 °C to leave a white solid, 106 mg.

¹H NMR δ 1.57 - 1.66 (m, 2H), 1.91 - 1.95 (m, 2H), 2.56 - 2.62 (m, 1H), 3.14 - 3.20 (m, 2H), 4.14 (d, 2H), 7.01 - 7.04 (m, 2H), 7.11 (d, 2H), 7.22 (t, 2H), 7.26 - 7.31 (m, 1H),
 5 7.63 (d, 2H), 8.14 (d, 1H), 8.51 (s, 1H), 11.06 (s, 1H), 11.24 (s, 1H); MS m/e MH⁺ 519.


Examples 703 and 704 were prepared using the procedure described above, from Examples 707 and 706 respectively.

10 Example 703: 1-(5-((5-((4-(4-Fluorophenoxy)phenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl)amino)pyridin-2-yl)-4-methylpiperidine-4-carboxylic acid




¹H NMR δ 1.17 (s, 3H), 1.37 - 1.43 (m, 2H), 1.97 - 2.00 (m, 2H), 3.11 (t, 2H), 3.86 - 3.89 (m, 2H), 6.87 (d, 1H), 7.01 - 7.04 (m, 2H), 7.09 (d, 2H), 7.22 (t, 2H), 7.62 (d, 2H), 7.90
 15 (d, 1H), 8.47 (s, 1H), 10.94 (s, 1H), 11.00 (s, 1H), 12.41 (s, 1H); MS m/e MH⁺ 533.

Example 704: 1-(5-((5-((4-(4-Fluorophenoxy)phenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl)amino)pyridin-2-yl)piperidin-4-yl]acetic acid



20 ¹H NMR δ 1.15 - 1.24 (m, 2H), 1.73 - 1.76 (m, 2H), 1.90 - 1.94 (m, 1H), 2.19 (d, 2H), 2.83 - 2.89 (m, 2H), 4.20 - 4.24 (m, 2H), 6.93 - 7.04 (m, 3H), 7.11 (d, 2H), 7.19 - 7.24 (m, 2H), 7.62 (d, 2H), 7.96 (d, 1H), 8.46 (s, 1H), 11.01 (s, 2H); MS m/e MH⁺ 533.

25 Example 705: Methyl 1-(5-((5-((4-(4-fluorophenoxy)phenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl)amino)pyridin-2-yl)piperidine-4-carboxylate




- 308 -

To a stirred solution of 4-(4-fluorophenoxy)aniline (190 mg, 0.93 mmol) in anhydrous DMA (4 mL) was added 1,1'-carbonyldiimidazole (167 mg, 0.93 mmol) and the mixture stirred at ambient temperature for 1 hr. Methyl 1-(5-
 5 {[(hydrazino(oxo)acetyl)amino]pyridin-2-yl)piperidine-4-carboxylate (250 mg, 0.78 mmol) was added and the reaction mixture was heated to 50 °C for 1 hr. EDCl.HCl (180 mg, 0.93 mmol) was added and the reaction mixture heated to 85 °C overnight and then allowed to cool to ambient temperature. Water (~20 mL) was added and the suspension filtered, washed with ether (~15 mL) and dried to leave the title compound as a pale yellow solid (415 mg, 0.780 mmol, 100%). ¹H NMR δ 1.50 - 1.58 (m, 2H), 1.87 - 1.90 (m, 2H), 2.61 - 2.67 (m, 1H), 2.90 - 2.94 (m, 2H), 3.62 (s, 3H), 4.17 (d, 2H), 6.87 - 6.89 (m, 1H), 7.00 - 7.10 (m, 4H), 7.19 - 7.23 (m, 2H), 7.60 - 7.63 (m, 2H), 7.90 - 7.92 (m, 1H), 8.48 (s, 1H), 10.95 (s, 1H), 10.99 (s, 1H); MS m/e MH⁺ 533.


Examples 706 and 707 were prepared using the procedure described above.

Example 706: Methyl [1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate




Methyl [1-(5-{[(hydrazino(oxo)acetyl)amino]pyridin-2-yl)piperidin-4-yl]acetate (262 mg, 0.78 mmol) was used instead of ethyl 1-(5-{[(hydrazino(oxo)acetyl)amino]pyridin-2-yl)piperidine-4-carboxylate. The title compound was isolated as a pale yellow solid (379 mg, 0.694 mmol, 89%). ¹H NMR δ 1.14 - 1.20 (m, 2H), 1.68 - 1.73 (m, 2H), 1.91 - 1.95 (m, 1H), 2.28 (d, 2H), 2.76 - 2.82 (m, 2H), 3.60 (s, 3H), 4.22 - 4.25 (m, 2H), 6.84 - 6.86 (m, 1H), 7.00 - 7.10 (m, 4H), 7.19 - 7.23 (m, 2H), 7.60 - 7.63 (m, 2H), 7.88 - 7.90 (m, 1H), 8.46 (s, 1H), 10.93 (s, 1H), 10.99 (s, 1H); MS m/e MH⁺ 548.

Example 707: Methyl 1-(5-([5-([4-(4-fluorophenoxy)phenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)pyridin-2-yl)-4-methylpiperidine-4-carboxylate




- 5 Methyl 1-(5-([hydrazino(oxo)acetyl]amino)pyridin-2-yl)-4-methylpiperidine-4-carboxylate (262 mg, 0.78 mmol) was used instead of ethyl 1-(5-([hydrazino(oxo)acetyl]amino)pyridin-2-yl)piperidine-4-carboxylate. The title compound was isolated as a pale brown solid (410 mg, 0.75 mmol, 96%). ^1H NMR δ 1.18 (s, 3H), 1.41 - 1.47 (m, 2H), 1.96 - 2.02 (m, 2H), 3.08 - 3.13 (m, 2H), 3.65 (s, 3H), 3.84 - 3.88 (m, 2H), 6.86 - 6.88 (m, 1H), 7.00 - 7.10 (m, 4H), 7.19 - 7.23 (m, 2H), 7.60 - 7.63 (m, 2H), 7.89 - 7.91 (m, 1H), 8.47 (s, 1H), 10.94 (s, 1H), 10.99 (s, 1H); MS m/e MH^+ 548.

Example 708: (1-(5-([5-([3,4-difluorophenyl]amino)-1,3,4-oxadiazol-2-yl]carbonyl]amino)pyridin-2-yl)-4-methylpiperidin-4-yl)acetic acid



- 15 Prepared from Intermediate 114 by hydrolysis using 5 equivalents of 2M NaOH in MeOH followed by purification on a reverse phase system eluting 5-95% water (0.2% TFA) - MeCN (0.2% TFA). ^1H NMR δ 1.07 (s, 3H), 1.42 - 1.46 (m, 2H), 1.56 - 1.61 (m, 2H), 2.24 (s, 2H), 3.40 - 3.47 (m, 2H), 3.58 - 3.62 (m, 2H), 6.88 (d, 1H), 7.34 - 7.36 (m, 1H), 7.50 (q, 1H), 7.68 - 7.73 (m, 1H), 7.91 (d, 1H), 8.46 (s, 1H), 10.98 (s, 1H), 11.27 (s, 1H), 12.04 (s, 1H); MS m/e MH^+ 473.

The following compounds were prepared from the appropriate esters (Intermediates 115-117) in the same manner as described above:




- 310 -

317

Example	Name	¹ H NMR (400 MHz, DMSO) δ	MS m/e M+H ⁺
709 R ¹ =R ² =R ³ =F	(4-methyl-1-{5-[(5-{(2,4,5-trifluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl}piperidin-4-yl)acetic acid	1.02 (s, 3H), 1.37 - 1.41 (m, 2H), 1.51 - 1.56 (m, 2H), 2.19 (s, 2H), 3.34 - 3.39 (m, 2H), 3.53 - 3.56 (m, 2H), 6.82 (d, 1H), 7.65 - 7.72 (m, 1H), 7.84 (d, 1H), 8.10 - 8.17 (m, 1H), 8.41 (s, 1H), 10.94 (s, 1H), 11.03 (s, 1H), 11.99 (s, 1H)	491
710 R ¹ =R ³ =H, R ² =CF ₃	[4-methyl-1-{5-[(5-{(4-(trifluoromethyl)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetic acid	11.50 (s, 1H), 11.19 (s, 1H), 8.48 (s, 1H), 8.06 (d, 1H), 7.82 - 7.77 (m, 4H), 7.16 - 7.11 (m, 1H), 3.66 - 3.62 (m, 2H), 3.51 - 3.47 (m, 2H), 2.27 (s, 2H), 1.66 - 1.62 (m, 2H), 1.52 - 1.46 (m, 2H), 1.09 (s, 3H); CO ₂ H not seen.	505
711 R ¹ =R ³ =H, R ² =O(4-F-C ₆ H ₄)	[1-{5-[(5-{(4-(4-fluorophenoxy)phenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl]acetic acid	1.07 (s, 3H), 1.41 - 1.46 (m, 2H), 1.56 - 1.62 (m, 2H), 2.24 (s, 2H), 3.41 - 3.45 (m, 2H), 3.57 - 3.63 (m, 2H), 6.84 (d, 1H), 7.00 - 7.04 (m, 2H), 7.09 (d, 2H), 7.21 (t, 2H), 7.62 (d, 2H), 7.91 (d, 1H), 8.46 (s, 1H), 10.95 (s, 1H), 11.00 (s, 1H), 12.05 (s, 1H)	547

Example 712: 2-(1-(3-Methyl-5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridin-2-yl)piperidin-4-yl)propanoic acid




Ethyl 2-(1-(3-methyl-5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl)amino]pyridine-2-yl)piperidin-4-yl)propanoate (Example 692) (70 mg; 0.13 mmol) was dissolved in dry THF (2.0 mL), treated with potassium trimethylsilanolate (167 mg; 1.30 mmol) and placed in a Biotage 'Initiator' microwave. The reaction was heated at 90°C for 35 mins., monitoring by LCMS, as a yellow precipitate formed. The reaction was incomplete (~9% ester remained). A further portion of potassium trimethylsilanolate (17 mg) was added and the reaction was returned to the microwave for 15 mins. Very little more reaction was seen (ie ~5% of starting ester remained). The resulting yellow suspension was treated with water (3 mL) and acidified, to about pH 2, with dropwise addition of aq. 1M HCl. With vigorous stirring for ~15mins, a white precipitate formed which was filtered, washed with water and dried under vacuum at 60 °C to give the title compound (45 mg; 79% yield).

¹H NMR δ 1.08 (d, 3H), 1.29 - 1.53 (m, 2H), 1.58 - 1.83 (m, 3H), 2.16 - 2.28 (m, 1H), 2.34 (s, 3H), 2.74 - 2.94 (m, 2H), 3.43 - 3.56 (m, 2H), 7.64 - 7.78 (m, 1H), 8.01 (s, 1H), 8.08 - 8.21 (m, 1H), 8.44 (s, 1H), 11.02 (s, 1H), 11.26 (s, 1H). MH⁺ = 505.53.

Preparation of Starting Materials

Intermediate 1: Methyl [(4-morpholin-4-yl phenyl)amino]oxoacetate




Methyl chlorooxoacetate (4.64 mL, 50 mmol) was added dropwise to an ice cooled solution of 4-morpholinoaniline (8.91 g, 50 mmol) and ethyldiisopropylamine (9.4 mL, 55 mmol) in DCM (125 mL). The reaction was stirred for 2 h at room temperature then quenched with water (100 mL). The organic layer was removed, dried (MgSO₄), filtered and evaporated to

- 312 -

give the title compound (11.7 g, 89%): ^1H NMR δ 10.63 (1H, s), 7.61 (2H, d), 6.92 (2H, d), 3.88 (3H, s), 3.78-3.68 (4H, m), 3.15-3.04 (4H, m); MS m/e MH^+ 265.

The following intermediates were prepared by the general procedure of intermediate 1 (method D1) using intermediates 36 and 38, known anilines [*tert*-butyl 4-(4-amino-2-fluorophenyl)piperazine-1-carboxylate (WO 2004018439), 5-amino-2-morpholin-4-ylbenzonitrile (WO 2000047558), (3-methyl-4-morpholin-4-ylphenyl)amine (DE 10220048), (3-fluoro-4-morpholin-4-ylphenyl)amine (J. Med. Chem. 1996, 39, 673-679)] and commercially available anilines. In method D2 trichloramine was used as the base instead of ethyldiisopropylamine and in Method D3 pyridine was used:



Inter	X	Y	R ⁶	^1H NMR δ	MS m/e MH^+	Method
2	CH ₂	CH	H		263	D2
3	NMe	CH	H		278	D2
4	NAc	CH	H		306	D2
5	NBoc	CH	H	10.58 (1H, s), 7.80 (1H, d), 6.95 (1H, d), 3.80 (3H, s), 3.40-3.45 (4H, m), 3.00-3.10 (4H, m), 1.40 (9H, s)	308	D2
6	O	CCl	H	10.85 (1H, s), 7.89 (1H, d), 7.68 (1H, d), 7.17 (1H, d), 3.85 (3H, s), 3.76-3.70 (4H, m), 2.98-2.90 (4H, m)	299	D1
7	O	CF	H	10.74 (1H, s), 7.65-6.97 (3H, m,), 3.86-3.69 (7H, m), 3.31- 2.46 (4H, m)	283	D3

- 313 -



330

Inter	X	Y	R ⁶	¹ H NMR δ	MS m/e MH ⁺	Method
8	O	CCN	H	10.91 (1H, s), 8.07-7.64 (2H, m), 7.21 (1H, t), 3.86-3.71 (7H, m), 3.28-3.06 (4H, m)	290	D3
9	O	CMe	H	10.53 (1H, s), 7.55-7.46 (2H, m), 7.00 (1H, d), 3.83 (3H, s), 3.75-3.68 (4H, m), 2.82-2.76 (4H, m), 2.23 (3H, s)	279	D3
10	O	N	H	10.74 (1H, s), 8.48 (1H, d), 7.91 (1H, dd), 6.88 (1H, d), 3.85 (3H, s), 3.74-3.66 (4H, m), 3.45-3.37 (4H, m)	266	D1
11	O	N	Cl	11.01 (1H, s), 8.57 (1H, d), 8.19 (1H, d), 3.85 (3H, s), 3.72 (4H, t), 3.23-3.17 (4H, m)	300	D3
12	NMe	N	H		279	D2
13	NAc	CMe	H	10.59 (1H, s), 7.56-7.46 (2H, m), 6.98 (1H, d), 3.83 (3H, s), 3.60-3.50 (4H, m), 2.84-2.69 (4H, m), 2.25 (3H, s), 2.02 (3H, s)	320	D3
14	NBoc	CF	H	10.82 (1H, s), 7.61 (1H, dd), 7.48 (1H, d), 7.03 (1H, t), 3.83 (3H, s), 3.50-3.41 (4H, m), 2.96-2.87 (4H, m), 1.41 (9H, s)	326 [MH-C ₄ H ₉] ⁺	D3
15	NAc	N	H		307	

Inter	X	Y	R ⁶	¹ H NMR δ	MS m/e MH ⁺	Method
16	NBoc	N	H	10.69 (1H, s), 8.45 (1H, s), 7.85 (1H, d), 5.75 (2H, s), 3.80 (3H, s), 3.40 (8H, s), 1.40 (9H, s)	365	

Intermediate 17: Methyl [(4-iodophenyl)amino](oxo)acetate


The title compound was prepared by the general procedure of intermediate 1 using triethylamine as base and 4-iodo-aniline. This yielded the title compound as a pale brown solid; MS m/e MH⁺ 306.

**Intermediate 18: 2-Hydrazino-N-(4-morpholin-4-ylphenyl)-2-oxoacetamide**

Hydrazine hydrate (1.25 mL, 25 mmol) was added to a stirred suspension of methyl [(4-morpholin-4-ylphenyl)amino](oxo)acetate (intermediate 1, 6.6 g, 25 mmol) in methanol (150 mL). The reaction was heated to 75 °C for 2 h during which time the precipitate thickened. After cooling the precipitate was filtered and washed with diethyl ether (50 mL) and dried to give the title compound (6.32 g, 94%):

¹H NMR δ 10.38 (1H, s), 10.15 (1H, s), 7.69 (2H, d), 6.91 (2H, d), 3.75 (4H, m), 3.08 (4H, m); MS m/e MH⁺ 265.

Intermediates 19-33 were prepared by the general procedure of intermediate 18 using intermediates 2-16 and either methanol or ethanol as solvent.




Intermediate	X	Y	R ⁶	¹ H NMR (DMSO-d ₆) δ	MS m/e MH ⁺
19	CH ₂	CH	H		263
20	NMe	CH	H		278
21	NAc	CH	H		306
22	NBoc	CH	H		308
23	O	CCl	H	10.68 (1H, s), 10.22 (1H, bs), 7.97 (1H, s), 7.75 (1H, dd), 7.17 (1H, d), 4.60 (2H, bs), 3.78-3.71 (4H, m), 2.98-2.91 (4H, m)	299
24	O	CF	H	7.67 (1H, dd), 7.60-7.54 (1H, m), 7.00 (1H, t), 4.78-4.24 (3H, m), 3.71 (4H, t), 2.95 (4H, t)	283
25	O	CC N	H	10.78 (1H, s), 10.27 (1H, s), 8.15-8.11 (1H, m), 8.01 (1H, dd), 7.18 (1H, d), 4.55 (2H, bs), 3.74 (4H, t), 3.09 (4H, t)	290
26	O	CMe	H	10.34 (1H, bs), 7.65-7.53 (2H, m), 6.98 (1H, d), 4.57 (2H, bs), 3.71 (4H, t), 2.78 (4H, t), 2.22 (3H, s)	279
27	O	N	H	10.54 (1H, s), 10.20 (1H, bs), 8.54 (1H, s), 7.95 (1H, d), 6.84 (1H, d), 4.60 (2H, bs), 3.73-3.66 (4H, m), 3.44-3.36 (4H, m)	266
28	O	N	Cl	10.85 (1H, bs), 10.29 (1H, bs), 8.65 (1H, d), 8.25 (1H, d), 4.62 (2H, bs), 3.71 (4H, t), 3.18 (4H, t)	300

Intermediate	X	Y	R ⁶	¹ H NMR (DMSO-d ₆) δ	MS m/e MH ⁺
29	NAc	CMc	H	10.45 (1H, s), 7.65-7.53 (2H, m), 6.97 (1H, d), 4.57 (2H, bs), 3.59- 3.50 (4H, m), 2.82-2.69 (4H, m), 2.24 (3H, s), 2.02 (3H, s)	320
30	NBoc	CF	H	10.62 (1H, s), 10.22 (1H, s), 7.68 (1H, dd), 7.56 (1H, d), 7.01 (1H, t), 4.59 (2H, bs), 3.49-3.41 (4H, m), 2.94-2.86 (4H, m), 1.41 (9H, s)	326 [MH- C ₄ H ₈] ⁺
31	NMe	N	H		279
32	NAc	N	H	10.45 (1H, s), 10.16 (1H, s), 8.50 (1H, d), 7.90 (1H, dd), 6.90 (1H, d), 4.60 (2H, bs), 3.50 (6H, m), 3.40 (2H, m), 2.00 (3H, s)	307
33	NBOC	N	H	10.45 (1H, s), 10.16 (1H, s), 8.50 (1H, d), 7.90 (1H, dd), 6.90 (1H, d), 4.60 (2H, bs), 3.00 (8H, s), 2.00 (9H, s)	309 [MH- C ₄ H ₈] ⁺

Intermediate 34 : *N*-(4-Iodophenyl)-2-hydrazino-2-oxoacetamide

The title compound was prepared by the general procedure of intermediate 18 using Intermediate 17.



5

MS m/e MH⁺ 306.


Intermediate 35: *N*-(4-Iodophenyl)-5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxamide

The title compound was prepared by the general procedure of Example 1 using methyl [(4-iodophenyl)amino](oxo)acetate.

10

- 317 -

34




MS m/e MH^+ 425.

This compound is a compound of formula (I) and so forms part of the invention.

Intermediate 36: 1-Acetyl-4-(4-amino-2-methylphenyl)-piperazine

5




Palladium on carbon catalyst (2.0 g) was added to a solution of intermediate 37 (19.5 g, 74.1 mmol) in EtOAc/EtOH (1:1) (250 mL). The reaction was allowed to stir under a balloon atmosphere of hydrogen for 18 h. The reaction was filtered through a pad of celite® and the resulting clear solution evaporated under reduced pressure. The resulting white solid was triturated with ether and dried *in vacuo* at 60 °C; ^1H NMR δ 6.74 (1H, d), 6.41 (1H, d), 6.36 (1H, dd), 4.67 (2H, broad s), 3.56-3.48 (4H, m), 2.72-2.60 (4H, m), 2.15 (3H, s), 2.02 (3H, s); MS m/e MH^+ 234.

10

Intermediate 37: 1-Acetyl-4-(2-methyl-4-nitrophenyl)piperazine

15




1-Butyl-3-methylimidazolium hexafluorophosphate (1.75 g, 7.7 mmol) and acetyl piperazine (39.7 g, 309.4 mmol) was added to a stirred solution of 2-fluoro-5-nitrotoluene (12 g, 77.4 mmol) in MeCN (3 mL) and heated to 85 °C for 2 h. The reaction was cooled and concentrated *in vacuo*. Water (100 mL) was added to the resulting residue and the reaction extracted with EtOAc (3 $\times$ 100 mL). The solvent was dried ($MgSO_4$), filtered, and concentrated *in vacuo* to yield the product as a yellow solid which was triturated with isohexane and dried *in vacuo* at 60 °C; ^1H NMR δ 8.05 (2H, m), 7.14 (1H, d), 3.64-3.58 (4H, m), 3.05-2.94 (4H, m), 2.37 (3H, s), 2.37 (3H, s); MS m/e MH^+ 264.


20

Intermediate 38: 5-Chloro-6-morpholin-4-ylpyridin-3-amine


The title compound was prepared by the general procedure of Intermediate 36 using 4-(3-chloro-5-nitro-pyridin-2-yl)-morpholine.



¹H NMR (300 MHz, DMSO-d₆) δ 7.64-7.60 (1H, m), 7.06-7.03 (1H, m), 5.15 (2H, broad s), 3.68 (4H, t, J = 4.5 Hz), 2.98-2.92 (4H, m); MS m/e MH⁺ 214.

Intermediate 39: 4-(3-Chloro-5-nitropyridin-2-yl)morpholine

2,3-Dichloro-5-nitropyridine (Synthesis, 1990, 6, 499-501) (20 g, 104 mmol) was added portionwise to morpholine (40 mL) at 0 °C and heated at 120 °C for 1 h. The reaction was cooled to room temperature, water (100 mL) added and the reaction extracted with DCM (3 × 150 mL). The organics were dried (MgSO₄), filtered, and the solvent removed under reduced pressure to give the product as a yellow solid, which was triturated with isohexane and allowed to dry under vacuum at 60 °C. ¹H NMR δ 8.98 (1H, d), 8.47 (1H, d), 3.74-3.69 (4H, m), 3.67-3.62 (4H, m); MS m/e MH⁺ 244.

Intermediate 40: Ethyl 5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxylate

Hydrazine monohydrate (834 mg, 16.6 mmol) was in DCM (70 mL), and cooled in a dry ice/acetone bath to -60 °C. 2-Fluoroisocyanate (2.28 g, 16.6 mmol) was dissolved in DCM (5 mL) and added dropwise to the hydrazine solution, which was then stirred at -60 °C for 2 h. The reaction mixture was allowed to warm to room temperature and stirred for 18 h. The solid formed was filtered and washed with DCM (2 × 20 mL). The filtrate was


concentrated *in vacuo* to quantitatively afford *N*-(2-fluorophenyl)hydrazinecarboxamide as a white solid (MS *m/e* MH^+ 170) which was used without further purification. This solid was stirred in ethanol-water (1:1, 50 mL). Ethyl glycoylate (50% solution in toluene, 2.55 g, 24.9 mmol), was added and the reaction allowed to stir for 18 h. The ethanol was

5 evaporated off from the reaction mixture *in vacuo*, and the resulting precipitate was filtered and washed with water (2×20 mL), then dried *in vacuo* to afford ethyl (2*E*)-({[(2-fluorophenyl)amino]carbonyl}hydrazono)acetate as a white solid (2.93 g, 70%, MS *m/e* MH^+ 254). This solid (3.55 g, 14.0 mmol), and sodium acetate (4.58 g, 14.7 mmol), were mixed together and suspended with stirring in glacial acetic acid (20 mL). A solution of

10 bromine (2.37 g, 14.7 mmol) in glacial acetic acid (6 mL), was added dropwise to the reaction and left to stir at room temperature for 30 min. The reaction was then heated in an oil bath at 130 °C for 1 h and poured into water (50 mL). The resulting solid was filtered and washed with water (2×20 mL), dissolved in DCM (50 mL), washed with water (2×20 mL), then the organics were dried ($MgSO_4$), filtered and concentrated *in vacuo* to afford

15 the title compound as an off white solid (2.92 g, 82%); ^1H NMR ($CDCl_3$) δ 8.18(1H, t), 7.72 (1H, s), 7.05-7.26(3H, m), 4.48 (2H, q), 1.45 (3H, t); MS *m/e* MH^+ 252.

Intermediate 41: 5-[(2-fluorophenyl)amino]-*N*-(6-piperazin-1-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide




20

Example 78 (9.8 g, 20.3 mmol) was stirred in DCM (20 mL), 4N HCl in Dioxan (30 mL) was added and the reaction stirred for 2 h. The reaction was concentrated *in vacuo*, EtOAc (30 mL), water (30 mL) and triethylamine (10 mL) added, and stirred for 5 min. The resulting solid was filtered and washed with water (10 mL) then EtOAc (10 mL), then

25 dried *in vacuo* to afford the title compound as a pale pink solid (7.4 g, 95%); ^1H NMR δ 8.50 (1H, s), 8.05 (1H, t), 7.95 (1H, d), 7.2-7.35(2H, m), 7.10-7.20 (1H, m), 3.45 (4H, s), 2.95 (4H, s); MS *m/e* MH^+ 384.



This compound is a compound of formula (I) and so forms part of the invention.

32X

Intermediate 42: 4-(5-Aminopyridin-2-yl)butan-1-ol

A mixture of 2-bromo-5-nitro pyridine (2.03 g, 10 mmol), 3-butyn-1-ol (1.05 g, 15 mmol), copper(I) iodide (19 mg, 0.1 mmol), dichlorobis(triphenylphosphine)palladium (II) (140 mg, 0.02 mmol), triethylamine (5.06 g, 50 mmol) in acetonitrile (30 mL) was stirred for 4 h. The reaction mixture was evaporated to dryness and loaded onto a Biotage KPSil60A cartridge, and purified by flash chromatography using Et₂O-DCM (1:9) as eluent to afford 4-(5-nitropyridin-2-yl)but-3-yn-1-ol as an oil (1.42 g, MS m/e MH⁺ 193). This oil (1 g, 5.21 mmol) was stirred in methanol (50 mL) and 10% palladium on carbon (0.2 g) was added and the reaction stirred under an atmosphere of hydrogen. The reaction was filtered through celite[®] and concentrated *in vacuo* to afford the title compound as an oil (0.87g).
¹H NMR δ 7.99 (1H, s), 6.93 (2H, s), 3.86 (2H, t), 3.55 (2H, s), 2.73 (2H, t), 1.70-1.74(2H, m), 1.60-1.62 (2H, m); MS m/e MH⁺ 167.


**Intermediate 43: Methyl [*trans*-4-(4-
 {[hydrazino(oxo)acetyl]amino}phenyl)cyclohexyl]acetate**

**i) Methyl (*trans*-4-phenylcyclohexyl)acetate**

10% Pd/C (4520 mg) was added to a solution of methyl [*trans*-4-(4-
 {[(trifluoromethyl)sulfonyl]oxy}phenyl)cyclohexyl]acetate (prepared as described in Patent Application WO2004/047755) (8100 mg) in MeOH (150 mL). The resulting suspension was stirred for 16 h under an atmosphere of Hydrogen. The suspension was filtered through diatomaceous earth and concentrated *in vacuo* to give a slurry. This was extracted into EtOAc (300 mL). The organic extract was washed with saturated aqueous sodium hydrogen carbonate solution (75 mL) and then brine (75 mL). The organic layer was dried and concentrated *in vacuo* to give the title compound (4720 mg) as an oil; ¹H


NMR δ 7.28-7.11 (5H, m), 3.58 (3H, s), 2.43 (1H+DMSO, m), 2.22 (2H, d), 1.83-1.67 (5H, m), 1.44 (2H, m), 1.13 (2H, m); MS m/e MH^+ 233.

ii) Methyl [*trans*-4-(4-aminophenyl)cyclohexyl]acetate



5 A mixture of 65% nitric acid (3.95 mL) and 95% sulphuric acid (4.97 mL) was added dropwise to a stirred solution of methyl (*trans*-4-phenylcyclohexyl)acetate (4710 mg) in CCl_4 (20 mL) at 5°C and the solution was allowed to warm to ambient temperature and stirred for 16 hours. Ice/water (50 mL) was added and the mixture was extracted with
10 DCM (2 x 40 mL). The organic extracts were combined, washed with brine (50 mL), dried, and concentrated *in vacuo* to give an oil. This oil was purified by flash chromatography on a 80 g Biotage™ silica column, using a gradient of 0-20% EtOAc in hexane as the eluent to give crude methyl [*trans*-4-(4-nitrophenyl)cyclohexyl]acetate which was dissolved in ethyl acetate (30 mL). 10% palladium on carbon (0.400 g) was added and the resulting
15 suspension was stirred at ambient temperature for 16 hours under an atmosphere of hydrogen. The suspension was filtered through diatomaceous earth and concentrated *in vacuo* to give a solid. This was purified by flash chromatography on a 40 g Biotage™ silica column using a gradient of 20-45% EtOAc / hexane as eluent to give the title compound (1740 mg) as a solid; ^1H NMR δ 6.83 (2H, d), 6.46 (2H, d), 4.72 (2H, s), 3.59
20 (3H, s), 2.23 (3H, m), 1.72 (5H, m), 1.35 (2H, m), 1.09 (2H, m); MS m/e MH^+ 248.

iii) Methyl ({4-[*trans*-4-(2-methoxy-2-oxoethyl)cyclohexyl]phenyl}amino)(oxo)acetate



Methyl chloro(oxo)acetate (0.842 mL) was added to a stirred solution of methyl [*trans*-4-(4-aminophenyl)cyclohexyl]acetate (1740 mg) and pyridine (0.689 mL) in DCM (50 mL)
25 at 0°C. After the addition was complete the mixture was allowed to warm to ambient temperature and stirred for 64 hours. The solution was diluted with DCM (100 mL), washed with water (50 mL) and brine (50 mL), then dried and concentrated *in vacuo* to give the title compound (2267 mg) as a solid; ^1H NMR δ 7.60 (2H, d), 7.18 (2H, d), 3.83


- 322 -

(3H, s), 3.58 (3H, s), 2.58-3.5 (1H+DMSO, m), 2.21 (2H, d), 1.75 (5H, m), 1.43 (2H, m), 1.13 (2H, m); MS m/e MH⁺ 332.


iv) Hydrazine hydrate (0.361 mL) was added to a stirred solution of methyl ({4-[*trans*-4-(2-methoxy-2-oxoethyl)cyclohexyl]phenyl}amino)(oxo)acetate (2260 mg) in EtOH (50 mL). The mixture was stirred for 1 hour. The precipitate was filtered off, washed with Et₂O, and dried under vacuum overnight to give the title compound (Intermediate 43, 1845 mg) as a solid; ¹H NMR δ 10.44 (1H, s), 10.20 (1H, s), 7.70 (2H, d), 7.21 (2H, d), 4.60 (2H, s), 3.60 (3H, s), 2.42 (1H, m), 1.79 (5H, m), 1.45 (2H, m), 1.11 (2H, m); MS m/e MH⁺ 334.

Intermediate 43': This refers to material prepared by exactly the same procedure as for Intermediate 43 except that ethanol was used as solvent in step (i), giving rise to a small proportion of ethyl ester in the final product.


Intermediate 44: Methyl *trans*-4-(4-{[hydrazino(oxo)acetyl]amino}phenyl)cyclohexanecarboxylate



i) Methyl *trans*-4-phenylcyclohexanecarboxylate



10% Palladium on carbon (1800 mg) was added to a solution of methyl *trans*-4-(4-chlorophenyl)cyclohexanecarboxylate (8670 mg) in EtOH (100 mL). The resulting suspension was stirred for 16 h under an atmosphere of hydrogen at 25 bar in a high pressure cell then was filtered through diatomaceous earth and concentrated *in vacuo* to give the title compound (6940 mg) as a solid; ¹H NMR δ 7.30-7.11 (5H, m), 3.59 (3H, s), 2.53-2.30 (2H+DMSO, m), 1.98 (2H, m), 1.82 (2H, m), 1.47 (4H, m); MS m/e M⁺ 218.

~~230~~ii) Methyl *trans*-4-(4-{{methoxy(oxo)acetyl}amino}phenyl)cyclohexanecarboxylate

A mixture of 65% nitric acid (6.2 mL) and 95% sulphuric acid (7.8 mL) in carbon tetrachloride (10 mL) was added dropwise to a solution of methyl *trans*-4-phenylcyclohexanecarboxylate (6940 mg) keeping the internal temperature below 5°C. The mixture was allowed to warm and stirred for 16 h. Ice/water (50 mL) was added and the mixture was extracted with DCM (2 x 75 mL). The extracts were combined, washed with brine (50 mL), dried and concentrated *in vacuo* to give an oil. This oil was chromatographed on an 80 g Biotage™ silica column, using a gradient of 0-20% EtOAc / hexane as the eluent to give crude methyl *trans*-4-(4-nitrophenyl)cyclohexanecarboxylate (4445 mg); MS *m/e* MH⁺ 264. 10% Pd/C (450 mg) was added to a solution of this material (4440 mg) in EtOAc (50 mL) and the suspension was stirred for 16 hours under an atmosphere of hydrogen then filtered through diatomaceous earth and concentrated *in vacuo*. The residue was purified by chromatography on an 80 g Biotage™ silica column using a gradient of 0-50% EtOAc / hexane as eluent to give crude methyl *trans*-4-(4-aminophenyl)cyclohexanecarboxylate (2900 mg). A solution of this material (2890 mg) and pyridine (1.67 mL) in DCM (50 mL) was cooled to 5°C. Methyl chloro(oxo)acetate (1.48 mL) was added and the solution was allowed to warm up with stirring for 16 h. Water (75 mL) was added and the mixture was extracted with DCM (2 x 75 mL). The extracts were combined, dried, and concentrated *in vacuo* to give a solid. This solid was purified by flash chromatography on 80g Biotage™ silica column eluting with a gradient of 0-50 % EtOAc / hexane to give the title compound (919 mg) as a solid; ¹H NMR δ 10.66 (1H, s), 7.60 (2H, d), 7.20 (2H, d), 3.83 (3H, s), 3.60 (3H, s), 2.56-2.30 (2H+DMSO, m), 2.00 (2H, m), 1.80 (2H, m), 1.46 (4H, m); MS *m/e* MH⁺ 320.

25


iii) Hydrazine hydrate (0.152 mL) was added to a stirred solution of methyl *trans*-4-(4-{{methoxy(oxo)acetyl}amino}phenyl)cyclohexanecarboxylate (913 mg) in EtOH (25 mL). The mixture was stirred for 2 hours. The precipitate was filtered off, washed with EtOH, and dried under vacuum to give the title compound (intermediate 44, 277 mg) as a solid;

331


¹H NMR δ 10.43 (1H, s), 7.67 (2H, d), 7.17 (2H, d), 4.52 (2H, s), 3.59 (3H, s), 2.53-2.33 (2H+DMSO, m), 1.97 (2H, m), 1.81 (2H, m), 1.46 (4H, m); MS m/e MH⁺ 320.

Intermediate 45: (4-{4-[(5-{(2-Fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid

The starting material (4-{4-[(5-{(2-Fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}phenyl)cyclohexyl)acetic acid for Example 119 was a *ca.* 2:1 *trans:cis* isomeric mixture and it was prepared in an analogous manner to that described for the pure *trans* isomer (Example 118) except that the starting material was ethyl (4-phenylcyclohexyl)acetate and it was a mixture of *cis/trans* isomers, prepared as described in Patent Application WO2004/047755.



Intermediate 46: 2-Hydrazino-2-oxo-N-(3-phenoxypropyl)acetamide




- 15 i) Methyl chloro(oxo)acetate (1.77 mL) was added to a stirred solution of (3-phenoxypropyl)amine hydrochloride (3000 mg) and pyridine (2.98 mL) in DCM (100 mL) at 0°C. After the addition was complete the solution was allowed to warm to ambient temperature and stirred for 16 hours. Water (50 mL) was added and the mixture was extracted with DCM (2 x 75 mL). The organic extracts were combined, washed with brine
- 20 (50 mL), dried and concentrated *in vacuo* to give an oil. This was purified by flash chromatography on a 90g Biotage™ silica column using a gradient of 10-60% EtOAc in hexane as the eluent to give methyl oxo[(3-phenoxypropyl)amino]acetate (3212 mg) as a solid; ¹H NMR δ (CDCl₃) 7.60 (1H, bs), 7.28 (2H+CHCl₃, m), 6.94 (3H, m), 4.07 (2H, t), 3.90 (3H, s), 3.58 (2H, quartet), 2.08 (2H, m), + 0.1 mole of EtOAc; MS m/e MH⁺ 238.
- 25 ii) Hydrazine monohydrate (6.3 mL) was added to a stirred suspension of methyl oxo[(3-phenoxypropyl)amino]acetate (3080 mg) in ethanol (50 mL). The reaction mixture was heated to 85°C for 3 hours then allowed to cool. The precipitate was filtered off, washed

- 325 -

332


with Et₂O and dried under vacuum, to give the title compound (Intermediate 46, 2.269 g) as a solid; ¹H NMR δ 9.95 (1H, s), 8.77 (1H, s), 7.28 (2H, m), 6.92 (3H, m), 4.50 (2H, s), 3.99 (2H, t), 3.30 (2H+H₂O, m), 1.94 (2H, t); MS m/e MNa⁺ 260.

5 **Intermediate 47: N-(4-Cyclohexylphenyl)-2-hydrazino-2-oxoacetamide**



- i) Methyl chloro(oxo)acetate (1.88 mL) was added to a stirred solution of 4-cyclohexylaniline (2980 mg) and pyridine (1.80 mL) in DCM (100 mL) at 0°C. After the addition was complete allowed to warm to ambient temperature and stirred for 16 hours.
- 10 Water (50 mL) added. Extracted with DCM (2 x 75 mL). The organics were combined washed with brine (50 mL), dried over MgSO₄ and concentrated *in vacuo* to give methyl [(4-cyclohexylphenyl)amino](oxo)acetate (4133 mg) as a solid; ¹H NMR δ 10.66 (1H, s), 7.60 (2H, d), 7.19 (2H, d), 3.84 (3H, s), 2.50 (1H+ DMSO, m), 1.84-2.65 (5H, m), 1.46-1.14 (5H, m); MS m/e M-H⁺ 260.
- 15 ii) Hydrazine monohydrate (7.40 mL) was added to a stirred suspension of methyl [(4-cyclohexylphenyl)amino](oxo)acetate (3990 mg) in ethanol (50 mL) at ambient temperature. The reaction was heated to 85°C and allowed to stir for a further 3 hours. Allowed to cool to ambient temperature overnight. The precipitate was filtered off, washed
- 20 with Et₂O and dried under vacuum, to give the title compound (Intermediate 47, 3750 mg) as a solid; ¹H NMR δ 10.42 (1H, s), 10.17 (1H, s), 7.68 (2H, d), 7.16 (2H, d), 4.58 (1H, s), 2.48 (1H+ DMSO, m), 1.89-1.65 (5H, m), 1.46-1.13 (5H, m); MS m/e M-H⁺ 260.

Intermediate 48: O-(Pentafluorophenyl) (3-ethoxyphenyl)thiocarbamate




25

A solution of pentafluorophenylchlorothioformate (4.0 mL 24.9 mmol) in dichloromethane (10 mL) was added dropwise to a stirred, ice-cooled solution of 3-ethoxyaniline (3.02 mL,


337

22.65 mmol) and pyridine (1.33 mL, 37.4 mmol) in dichloromethane (250 mL). The mixture was allowed to reach room temperature and stirred for a further 14 h. The mixture was washed with 1M aqueous citric acid (100 mL), saturated aqueous NaHCO₃ (100 mL) and water (100 mL) then dried. Volatile material was removed by evaporation to give an oil that was purified by flash chromatography on SiO₂ eluting with isohexane to give the title compound (6.2 g) as an oil; ¹H NMR (CDCl₃) δ 7.28-7.17 (1H, m), 6.84-6.76 (2H, m), 6.75-6.69 (1H, m), 4.01 (2H, q), 1.41 (3H, t).

Intermediate 49: 5-(Biphenyl-3-yl){2-(trimethylsilyl)ethoxy)methyl}amino)-N-(6-morpholin-4-ylpyridin-3-yl)-N-{[2-(trimethylsilyl)ethoxy)methyl]-1,3,4-oxadiazole-2-carboxamide



(i) 5-((3-Iodophenyl){2-(trimethylsilyl)ethoxy)methyl}amino)-N-(6-morpholin-4-ylpyridin-3-yl)-N-{[2-(trimethylsilyl)ethoxy)methyl]-1,3,4-oxadiazole-2-carboxamide



15

5-[(3-Iodophenyl)amino]-N-(6-morpholin-4-ylpyridin-3-yl)-1,3,4-oxadiazole-2-carboxamide (Example 146) (610 mg, 1.24 mmol) was added portionwise to an ice-cooled suspension of sodium hydride (60% dispersion in oil) (119 mg, 2.97 mmol) in anhydrous DMF (5 mL) in dry glassware. The mixture was stirred at 0°C for 10 minutes and ambient temperature for 30 minutes then recooled in an ice bath and [2-(chloromethoxy)ethyl](trimethyl)silane (0.482 mL, 2.73 mmol) was added dropwise. The mixture was allowed to warm up and stirred for 16 h. ether (60 mL) was added, and the mixture was washed with water (3 x 20 mL) and the aqueous extracts were re-extracted with ether (20 mL). The organic extracts were combined, dried and concentrated by

20

334

evaporation and the residue was purified by chromatography, eluting with 0-20% ether/DCM to give the title compound (610 mg, 65%) as gum; $^1\text{H NMR}$ δ 8.09 (1H, s), 7.94 (1H, s), 7.74 (1H, d), 7.59-7.40 (2H, m), 7.28 (1H, t), 6.86 (1H, d), 5.24-5.11 (1H, m), 3.80-3.48 (12H, m), 0.98-0.84 (4H, m), 0.10-0.05 (18H, m).

5


(ii) Potassium phosphate (214 mg, 1.01 mmol) was added to 5-((3-iodophenyl){[2-(trimethylsilyl)ethoxy]methyl}amino)-*N*-(6-morpholin-4-ylpyridin-3-yl)-*N*-{[2-(trimethylsilyl)ethoxy]methyl}-1,3,4-oxadiazole-2-carboxamide (190 mg, 0.25 mmol) and phenylboronic acid (37mg, 0.30mmol) in DMF (3 mL) and the mixture was degassed.

- 10 Tetrakis(triphenylphosphine)palladium (29 mg, 0.03 mmol) was added and the mixture was heated to 80°C for 90 minutes. Phenylboronic acid (8 mg, 0.06 mmol) and tetrakis(triphenylphosphine)palladium (10 mg, 0.01 mmol) was added, the mixture degassed and heated to 80°C for 240 minutes. The mixture was cooled, diluted with ether (60 mL) and washed with water (3 x 20 mL). The aqueous layers were re-extracted with
- 15 ether (20 mL) and the organic layers were combined, dried and concentrated by evaporation. The residue was purified by chromatography, eluting with 0-20% ether/DCM to give the title compound (Intermediate 49, 149 mg, 84%) as a gum; $^1\text{H NMR}$ δ 8.09 (1H, s), 7.83-7.66 (4H, m), 7.66-7.37 (6H, m), 6.86 (1H, d), 5.37 (2H, s), 5.19 (2H, s), 3.79-3.62 (8H, m), 3.55-3.49 (4H, m), 0.93 (4H, t), 0.06 (9H, s), 0.01 (9H, s).


20

Intermediates 50-57








The following intermediates were prepared by the general procedure of intermediate 1 (Method D3) using intermediates as indicated in the table, anilines known in the literature as referenced or commercially available anilines.



25

Int.	Het/Ar	$^1\text{H NMR}$ δ	MS m/e MH ⁺	SM
50		8.48 (1H, d), 8.0 (1H, dd), 6.8 (1H, d), 3.81 (3H, s), 3.83 (3H, s).	211	Purchased


331

Int.	Het/Ar	¹ H NMR δ	MS m/e MH ⁺	SM
51		10.55 (1H, s), 7.57 (2H, d), 6.90 (2H, d), 3.82 (3H, s), 3.72-3.60 (2H, m), 3.56-3.48 (2H, m), 2.20 (2H, t), 1.13 (6H, d)	293	Int. 67
52		10.45 (1H, s), 7.51 (2H, dd), 6.75-6.66 (2H, m), 3.81 (3H, s), 3.67-3.42 (6H, m), 3.36-3.23 (2H, m), 1.98-1.71 (5H, m)	320	Int. 69
53			MH ⁺ - C ₄ H ₈ : 365	Int. 72
54			251	Purchased
55		10.68 (1H, s), 8.40 (1H, s), 7.80 (1H, d), 6.80 (1H, d), 3.85 (3H, s), 3.90-3.80 (4H, m), 2.60-2.50 (4H, m)	282	Lit. ¹
56		10.08 (1H, s), 8.10 (1H, s), 7.70 (1H, d), 7.20 (1H, d), 3.85 (3H, s), 3.80-3.70 (4H, m), 2.95-2.85 (4H, m)	345	Int. 75
57		10.82 (1H, s), 8.10 (1H, s), 7.80 (1H, d), 7.10 (1H, d), 6.10-5.95 (1H, m), 5.40 (1H, d), 5.30 (1H, d), 4.70 (2H, d), 3.85 (3H, s), 3.80-3.70 (4H, m), 2.95-2.85 (4H, m)	349	Int. 76





¹6-Thiomorpholin-4-ylpyridin-3-amine, prepared as described in Patent Application WO02081453.


Intermediates 58-66

- 5 The following intermediates were prepared by the general procedure of Intermediate 18 using Intermediates 50-57 or 74 and either methanol or ethanol as solvent.




Int.	Het/Ar	¹ H NMR δ	MS m/e MH ⁺	SM Int.
58		10.48 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 6.90 (1H, d), 3.90-3.80 (4H, m), 2.60-2.50 (1H, m)	211	50
59		10.32 (1H, s), 7.63 (2H, d), 6.89 (2H, d), 4.65-4.39 (2H, m), 3.72-3.60 (3H, m), 3.52 (2H, d), 2.19 (2H, t), 1.13 (6H, d)	293	51
60		10.21 (1H, s), 10.07 (1H, s), 7.62-7.54 (2H, m), 6.69 (2H, dd), 4.54 (2H, s), 3.66-3.42 (6H, m), 3.34-3.24 (2H, m), 1.97-1.71 (5H, m)	320	52
61		8.71 (2H, s), 3.76-3.62 (4H, m), 3.53-3.46 (4H, m), 2.02 (3H, s)	308	71
62			251	54

Int.	Het/Ar	¹ H NMR δ	MS m/e MH ⁺	SM Int.
63			343	74
64		10.48 (1H, s), 8.50 (1H, s), 7.90 (1H, d), 6.90 (1H, d), 3.90-3.80 (4H, m), 2.60- 2.50 (1H, m)	282	55
65		10.65 (1H, s), 10.22 (1H, s), 8.10 (1H, s), 7.80 (1H, d), 7.20 (1H, d), 4.6 (2H, s), 3.80-3.70 (4H, m), 2.95-2.85 (4H, m)	345	56
66		10.65 (1H, s), 10.20 (1H, s), 8.20 (1H, s), 7.85 (1H, d), 7.10 (1H, d), 6.10-5.95 (1H, m), 5.40 (1H, d), 5.30 (1H, d), 4.70 (2H, d), 4.60-4.55 (2H, m), 3.80-3.70 (4H, m), 2.95-2.85 (4H, m)	349	57


Intermediate 67: 4-[cis-2,6-dimethylmorpholin-4-yl]aniline

- 5 A mixture of *cis*-2,6-dimethyl-4-(4-nitrophenyl)morpholine (Intermediate 68, 8.0 g) and 10% palladium on carbon (800 mg) in ethanol (100 mL) was stirred under an atmosphere of hydrogen at 50°C for 6 h. The mixture was filtered and the filtrate was concentrated by evaporation to give the title compound (4.3 g, 100%) as an oil which subsequently solidified; ¹H NMR δ (CDCl₃): 6.79 (2H, d), 6.65 (2H, d), 3.81 (2H, m), 3.25 (2H, d), 2.33 (2H, t), 1.23 (6H, d); MS m/e MH⁺: 207.

Intermediate 68: *cis*-2,6-Dimethyl-4-(4-nitrophenyl)morpholine


1-Fluoro-4-nitrobenzene (4.9 g, 34.75 mmol) was added to a stirred mixture of *cis*-2,6-dimethylmorpholine (44.4 g, 38.26 mmol) and anhydrous potassium carbonate (2.5 g, 18.12 mmol) in acetonitrile (50 mL). The mixture was heated under reflux for 19 h, cooled, filtered and the solid washed with acetonitrile. The filtrate was concentrated by evaporation to give the title compound (8.1 g, 99%) as a solid; ¹H NMR δ (CDCl₃): 8.13 (2H, d), 6.82 (2H, d), 3.76 (2H, m), 3.66 (2H, dt), 2.61 (2H, t), 1.28 (6H, d); MS m/e MH⁺: 237.

10

Intermediate 69: [4-(4-acetyl-1,4-diazepan-1-yl)phenyl]amine

Following the general procedure of Intermediate 36 but using 1-acetyl-4-(4-nitrophenyl)-1,4-diazepane (Intermediate 70) as starting material there was thus obtained the title compound in 97% yield; ¹H NMR δ: 6.55-6.46 (4H, m), 4.30 (2H, br s), 3.59-3.48 (3H, m), 3.41-3.26 (5H, m), 1.99-1.72 (5H, m); MS m/e MH⁺: 234.

15


Intermediate 70: 1-Acetyl-4-(4-nitrophenyl)-1,4-diazepane

N-Acetylhomopiperazine (7.2 mL, 54.6 mmol) and DIPEA (12.3 mL, 74.4 mmol) were added to a stirred solution of 4-fluoronitrobenzene (7.0 g, 49.6 mmol) in acetonitrile (100 mL) and the mixture was heated to 80°C under nitrogen with stirring for 16 h. Volatile material was removed by evaporation and the residue was dissolved in EtOAc (150 mL) and water (150 mL) was added. The organic layer was separated and the aqueous layer was extracted again with EtOAc (2 x 150 mL). The organic layers were combined and dried, concentrated by evaporation then the residue was triturated with isohexane to give the title

25


compound (5.6 g, 53%) as a solid; ^1H NMR δ 8.01 (2H, qd), 6.92-6.82 (2H, m), 3.80 (1H, t), 3.70-3.56 (5H, m), 3.35 (2H, dt), 1.97-1.71 (5H, m); MS m/e MH^+ : 264.

Intermediate 71: Methyl {[2-(4-acetyl)piperazin-1-yl]pyrimidin-5-yl}amino}{oxo}acetate



A solution of *tert*-Butyl 4-(5-{[methoxy(oxo)acetyl]amino}pyrimidin-2-yl)piperazine-1-carboxylate (Intermediate 53; 4.14 g, 11.33 mmol) in a 4M solution of HCl in dioxane (40 mL) was stirred for 16 h. The precipitate was collected by filtration and washed with ether to give *N*-(5-{[methoxy(oxo)acetyl]amino}pyrimidin-2-yl)piperazine hydrochloride as a solid. Pyridine (9.16 mL, 113.30 mmol), followed by acetyl chloride (0.97 mL, 13.60 mmol) were added to an ice-cooled suspension of this solid (3.4 g) in DCM (100 mL) and the mixture was allowed to warm to ambient temperature and stirred for 1 h. Water (100 mL) was added and the mixture was extracted with DCM (3 x 100 mL). The organic extracts were combined, dried and concentrated by evaporation. The residue was triturated with isohexane to give the title compound (2.45 g, 70%) as a solid; ^1H NMR δ : 10.79 (1H, s), 8.65 (2H, s), 3.84 (3H, s), 3.75-3.65 (4H, m), 3.52-3.45 (4H, m), 2.04 (3H, s); MS m/e MH^+ : 308.


Intermediate 72: *tert*-Butyl 4-(5-aminopyrimidin-2-yl)piperazine-1-carboxylate



Bis(dibenzylideneacetone)palladium (0.17 g, 0.29 mmol) was added to a mixture of *tert*-butyl 4-(5-bromopyrimidin-2-yl)piperazine-1-carboxylate (6.15 g, 17.92 mmol, prepared as described in Patent Application WO 2001087870), sodium *tert*-butoxide (2.41 g, 25.09 mmol), benzophenone imine (3.61 g, 21.50 mmol), and BINAP (0.34 g, 21.5 mmol) in anhydrous toluene (150 mL). The mixture was heated to 80°C and stirred for 16 h under an atmosphere of nitrogen. The mixture was diluted with EtOAc (100 mL) and filtered through diatomaceous earth then concentrated by evaporation to give an oil. This

intermediate was stirred together with THF (150 mL) and 2N HCl (100 mL) for 2 h. The mixture was then neutralised by the addition of saturated aqueous sodium hydrogen carbonate solution and the mixture was extracted with EtOAc (3 x 100 mL). The organic extracts were dried, filtered, concentrated by evaporation and triturated with hexane to give the title compound as a solid (4.6 g, 92%); ^1H NMR δ : 7.90 (2H, s), 4.61 (2H, s), 3.51-3.45 (4H, m), 3.40-3.30 (4H, m), 1.20 (9H, s); MS m/e MH^+ : 280.

Intermediate 73: 5-[(2-Fluorophenyl)amino]-N-(4-piperazin-1-ylphenyl)-1,3,4-oxadiazole-2-carboxamide




10

Following the general procedure of Intermediate 41 but using Example 80 as starting material in place of Example 78 there was thus obtained the title compound in 69% yield; ^1H NMR δ 10.88 (1H, s), 9.40 (1H, s), 8.00 (1H, t), 7.60 (2H, d), 7.35-7.10 (3H, m), 7.00 (2H, d), 3.40-3.30 (4H, m), 3.20-3.10 (4H, m); MS m/e MH^+ 383.


15 This compound is a compound of formula (I) and so forms part of the invention.

Intermediate 74: Methyl (6-[4-(methylsulfonyl)piperazin-1-yl]pyridin-3-yl)amino(oxo)acetate




20 A solution of *tert*-butyl 4-(5-([methoxy(oxo)acetyl]amino)pyridin-2-yl)piperazine-1-carboxylate (Intermediate 16, 4 g, 11 mmol) in 4N HCl/dioxan (25 mL) was stood for 16 h then concentrated by evaporation. The residue was dissolved in a mixture of DMF (20 mL) and pyridine (7 mL) then methanesulphonyl chloride (1.28 mL, 16.48 mmol) was added. The mixture was stirred for 3 hours then was diluted with water (50 mL) and extracted with 25 dichloromethane (2 x 100 mL). The organic extracts were dried and concentrated by evaporation to give the title compound as a solid that was used without further purification; MS m/e MH^+ 343.

- 334 -


~~341~~Intermediate 75: (3-bromo-4-morpholin-4-ylphenyl)amine

i) 4-(2-bromo-4-nitrophenyl)morpholine:




- 5 Morpholine (7.92 mL, 90.9 mmol) was added to a solution of 2-bromo-1-fluoro-4-nitrobenzene (10 g, 45.5 mmol) in acetonitrile (100 mL) and the mixture was heated under reflux for 6 h. Volatile material was removed by evaporation and the residue was diluted with water (100 mL) then extracted with ethyl acetate (2 x 100 mL). The organic extracts were combined, dried and concentrated to give the title compound (12.9 g, 99%) as a solid;
- 10 ¹H NMR δ 8.40 (1H, s), 8.20 (1H, d), 7.10 (1H, d), 3.80-3.70 (4H, m), 3.20-3.10 (4H, m); MS m/e MH⁺ 287.

- ii) Following the general procedure of Intermediate 36 but using 4-(2-bromo-4-nitrophenyl)morpholine as starting material and allowing 5 h for hydrogenation and
- 15 subsequently purifying by chromatography eluting with EtOAc / isohexane there was thus obtained the title compound (Intermediate 75) in 24% yield; ¹H NMR δ 6.90 (1H, d), 6.80 (1H, s), 6.50 (1H, dd), 5.00 (2H, s), 5.40 (1H, d), 3.70-3.60 (4H, m), 2.80-2.70 (4H, m); MS m/e MH⁺ 258.

20 Intermediate 76: Allyl 5-amino-2-morpholin-4-ylbenzoate


i) Allyl 2-morpholin-4-yl-5-nitrobenzoate




Morpholine (7.78 mL, 88.9 mmol) was added to a solution of 2-fluoro-5-nitrobenzoic acid allyl ester (prepared as described in WO 9315078; 10 g, 44.4 mmol) in acetonitrile (100 mL) and the mixture was heated under reflux for 6 h. Volatile material was removed by evaporation and the residue was diluted with water (100 mL) and extracted with EtOAc (2 x 100 mL). The organic layers were dried and concentrated by evaporation to give the title compound (12.87 g, 99%) as a viscous liquid; ^1H NMR δ : 8.40(1H, s), 8.20 (1H, d), 7.10 (1H, d), 6.10-5.95 (1H, m), 5.40(1H, d), 5.30 (1H, d), 4.80 (2H, s), 3.70-3.60 (4H, m), 3.20-3.10 (4H, m); MS m/e MH^+ 293.

ii) Tin(II) chloride dihydrate (17.37 g, 77 mmol) was added to a solution of allyl 2-morpholin-4-yl-5-nitrobenzoate (11.24 g, 38.5 mmol) in EtOAc (250 mL) and the mixture was stirred and heated under reflux for 4 h. Ethanol (150 mL) was added, followed by another portion of Tin(II) chloride dihydrate (17.4 g) and the mixture was heated for another 2 h. The mixture was cooled and filtered then the filtrate was washed repeatedly with aqueous ammonium hydroxide and re-filtered. The organic layer was dried and concentrated by evaporation to give the title compound (Intermediate 76, 5.42 g, 54%) as a solid; ^1H NMR δ 6.90 (1H, d), 6.80 (1H, s), 6.70 (1H, d), 6.10-5.95 (1H, m), 5.40 (1H, d), 5.30 (1H, d), 5.00 (2H, s), 4.80 (2H, s), 3.70-3.60 (4H, m), 3.20-3.10 (4H, m); MS m/e MH^+ 263.

Intermediate 77: Methyl [1-(5-{{hydrazine(oxo)acetyl}amino}pyridin-2-yl)piperidin-4-yl]acetate




i) Methyl [1-(5-nitropyridin-2-yl)piperidin-4-yl]acetate



A mixture of 2-Chloro-5-nitropyridine (1.59 g, 10 mmol) and methyl piperidin-4-ylacetate hydrochloride (1.8 g, 10 mmol) (prepared as described in Patent Application WO 2000029407) in DMA (50 mL) was treated with 2-tert-butylimino-2-diethylamino-1,3-dimethyl-perhydro-1,3,2-diazaphosphorine (BEMP) (15 g of a ~2.2 mmol/g loading on polystyrene support, ~33 mmol). The resulting mixture was stirred for 2.5 h then filtered through diatomaceous earth. Volatile material was removed by evaporation to give the title compound (740 mg, 21%) as a solid; ¹H NMR δ 1.08-1.28 (2H, m), 1.77 (2H, d), 1.99-2.12 (1H, m), 2.29 (2H, d), 3.07 (2H, t), 3.60 (3H, s), 4.51 (2H, d), 6.94 (1H, d), 8.13-8.25 (1H, m), 8.95 (d, 1H); MS m/e MH⁺ 280.

ii) Methyl [1-(5-aminopyridin-2-yl)piperidine-4-yl]acetate




A solution of methyl [1-(5-nitropyridin-2-yl)piperidin-4-yl]acetate (710 mg, 2.60 mmol) in EtOH (26 mL) was treated with 10% Pd on charcoal (71 mg), and stirred vigorously under an atmosphere of hydrogen for 4h. The reaction mixture was filtered through diatomaceous earth and the filtrate was evaporated to dryness to give the title compound (556mg, 70%) as an oil; ¹H NMR δ 1.12-1.27 (2H, m), 1.67 (2H, d), 1.75-1.89 (1H, m), 2.25 (2H, d), 2.56 (2H, t), 3.33 (2H, s), 3.58 (3H, s), 4.52 (2H, s), 6.61 (1H, d), 6.86-6.93 (1H, m), 7.58 (1H, d); MS m/e MH⁺ 250.


iii) Following the general procedure described for Intermediate 47 there was thus obtained, from the first step following chromatography eluting with 0-50% EtOAc / isohexane, ({6-[4-(2-Methoxy-2-oxoethyl)piperidin-1-yl]pyridin-3-yl} amino)(oxo)acetate (in 97% yield) as a solid; ¹H NMR δ 1.09-1.26 (2H, m), 1.69 (2H, d), 1.84-2.00 (1H, m), 2.29 (2H, d), 2.71-2.86 (2H, m), 3.58 (3H, s), 3.80 (3H, s), 4.14-4.28 (2H, m), 6.84 (1H, d), 7.79-7.91 (1H, m), 8.42 (1H, d), 10.71 (1H, s); MS m/e MH⁺ 336; and from the second step, the title

compound (Intermediate 77, in 33% yield) as a solid; ^1H NMR δ 1.11-1.23 (2H, m), 1.69 (2H, d), 1.84-2.00 (1H, m), 2.28 (2H, d), 2.70-2.85 (2H, m), 3.60 (3H, s), 4.22 (2H, d), 4.60 (2H, s), 6.82 (1H, d), 7.85-7.94 (1H, m), 8.49 (1H, d), 10.22 (1H, s), 10.51 (1H, s); MS m/e MH^+ 336.

Intermediate 78: Methyl [(trans-4-(5-{[hydrazino(oxo)acetyl]amino}pyridin-2-yl)cyclohexyl]acetate




i) 1,4-Dioxaspiro[4.5]dec-7-en-8-yl trifluoromethanesulphonate



To a solution of diisopropylamine (10 mL, 67.61 mmol) in anhydrous THF (100 mL) cooled to -78°C was added $n\text{BuLi}$ (34 mL, 67.6 mmol, 2M solution in cyclohexane) and the reaction mixture was allowed to stir at -78°C for 10 minutes. 1,4-Dioxaspiro[4.5]decan-8-one (9.60 g, 61.46 mmol) in anhydrous THF (100 mL) was added and the reaction mixture allowed to stir for 90 minutes at -78°C . A solution of *N*-phenyltrifluoromethanesulphonimide (25 g, 76.83 mmol) in anhydrous THF (100 mL) was added and the reaction mixture was allowed to warm slowly to ambient temperature overnight. The solvent was concentrated under reduced pressure and the residue was redissolved in ether (500 mL) and washed with 1M HCl (2 x 200 mL), 1M NaOH (2 x 200 mL) brine (200 mL). The organic phase was separated, dried (MgSO_4) and concentrated to leave crude product. The residue was columned on a 300 g silica Biotage cartridge loading in isohexane and eluting a gradient of isohexane – DCM 0-50% to leave a clear oil, 11.86 g (41.14 mmol, 67%): ^1H NMR (CDCl_3) δ 1.89-1.91 (m, 2H), 1.90 (t, 2H), 2.52-2.56 (m, 2H), 3.99 (s, 4H), 5.66 (s, 1H)


ii) 8-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-1,4-dioxaspiro[4.5]dec-7-ene



- 338 -

To a stirred solution of 1,4-dioxaspiro[4.5]dec-7-en-8-yl trifluoromethanesulphonate prepared as above (11.18 g, 38.79 mmol) in dry dioxane (200 mL) was added bis(pinacolato)diboron (10.8 g, 42.67 mmol), Pd(dppf)₂Cl₂DCM ([1,1'-bis(diphenylphosphino(ferrocene)]dichloropalladium(II), complex with DCM (1:1) (951 mg, 1.16 mmol), bis(diphenylphosphino)ferrocene (646 mg, 1.16 mmol) and potassium acetate (11.4 g, 0.116 mol). The reaction mixture was degassed by bubbling nitrogen through it for 15 minutes then heated to 80 °C for 3 hours. The reaction mixture was allowed to cool, the solvent removed and the residue was partitioned between EtOAc (200 mL) and water (200 mL), the organic phase was washed with brine (200 mL) and aqueous extracts were combined and reextracted with EtOAc (200 mL). The organic washings were combined, dried (MgSO₄) and concentrated under reduced pressure to leave an oil. This was dissolved in DCM-isohexane (1:1, 20 mL) and the filtrate purified on a 120 g SiliCycle cartridge, eluting with 50%-100% DCM/isohexane to give the title compound (8.4 g, 81%); ¹H NMR (CDCl₃) δ 1.25 (s, 12H), 1.71-1.75 (m, 2H), 2.34-2.37 (m, 4H), 3.98 (s, 4H), 6.48 (s, 1H); GCMS EI 266 M⁺.

lii) 2-(1,4-Dioxaspiro[4.5]dec-7-en-8-yl)-5-nitropyridine



A stirred solution of 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,4-dioxaspiro[4.5]dec-7-ene prepared as above (8.40 g, 31.56 mmol), 2-bromo-5-nitropyridine (6.0 g, 29.55 mmol) in DME (200 mL), degassed 2M potassium carbonate (40 mL) was added palladium tetrakis(triphenylphosphine) (Pd(PPh₃)₄; 1.8 g, 1.58 mmol). The reaction mixture was evacuated with nitrogen (3 cycles) and allowed to stir under vacuum for 5 minutes, then heated to 80°C overnight. The reaction mixture was allowed to cool to ambient temperature, EtOAc (200 mL) was added followed by water (200 mL), the suspension was filtered and the organic phase was separated and the aqueous layer re-extracted into EtOAc (200 mL). The organic extracts were combined, washed with brine (200 mL), dried (MgSO₄) and concentrated to leave crude product. DCM was added and the suspension was filtered and dried to leave the title compound (1.64 g). The filtrate was concentrated after the addition of DCM a further 1.51 g of product was obtained, the


- 339 -

filtrate was purified on a 120 g SiliCycle cartridge eluting 20-50-70% EtOAc in isohexane to provide 1.72 g of product. Total yield is 4.85 g (18.51 mmol, 58%);

^1H NMR (CDCl_3) δ 1.96 (t, 2H), 2.57-2.58 (m, 2H), 2.79-2.83 (m, 2H), 4.03 (s, 4H), 6.85-6.88 (m, 1H), 7.53 (d, 1H), 8.39 (d, 1H), 9.35 (s, 1H); GCMS 263 MH^+ .

5

iv) 6-(1,4-Dioxaspiro[4.5]dec-8-yl)pyridin-3-amine




Wet 50% palladium on carbon (750 mg) was added to a solution of 2-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-5-nitropyridine (3.96g, 15.11 mmol), prepared as above, in MeOH (500 mL). The reaction mixture was deoxygenated by evacuation and placed under an atmosphere of hydrogen (5 cycles) then stirred for 16 h. The mixture was filtered, and the filtrate concentrated to give the title compound (3.10 g, 87%) as a solid;

10

^1H NMR (CDCl_3) δ 1.68-1.96 (m, 8H), 2.63-2.70 (m, 1H), 3.97 (s, 4H), 3.97 (s, 2H), 6.93-7.00 (m, 2H), 8.03 (s, 1H); MS m/e MH^+ 235.

15

v) Benzyl [6-(1,4-dioxaspiro[4.5]dec-8-yl)pyridin-3-yl]carbamate



To an ice water cooled solution of 6-(1,4-dioxaspiro[4.5]dec-8-yl)pyridin-3-amine (3.08g, 13.15 mmol), prepared as above, in THF (50 mL) was added pyridine (3.19 mL, 39.44 mmol) followed by benzyl chloroformate (1.9 mL, 13.15 mmol). The resulting suspension was allowed to stir at 0 °C, then after 2 hrs a further 0.5 mL of benzyl chloroformate was added and reaction mixture was allowed to stir at ambient temperature overnight. It was partitioned between EtOAc (100 mL) and water (50 mL), the organic phase was separated and the aqueous phase re extracted into EtOAc (200 mL). The organic extracts were combined, dried and concentrated to leave crude product. This was purified on a 120 g silica SiliCycle cartridge, eluting with 30-50% EtOAc/isohexane using an Isco Companion, to provide the title compound (4.94 g, 100%); ^1H NMR (CDCl_3) δ 1.66-1.97 (m, 8H), 2.70-2.77 (m, 1H), 3.97 (s, 4H), 5.21 (s, 2H), 6.71 (s, 1H), 7.16 (d, 1H), 7.35-7.41 (m, 5H), 7.91 (s, 1H), 8.37 (s, 1H); MS m/e MH^+ 369.

20


25

30

- 340 -


~~341~~

vi) Benzyl [6-(4-oxocyclohexyl)pyridin-3-yl]carbamate



To an ice water cooled solution of benzyl [6-(1,4-dioxaspiro[4.5]dec-8-yl)pyridin-3-yl]carbamate, prepared as above, (4.94 g, 13.41 mmol) was added TFA (20 mL) followed by water (1.1 mL). The reaction mixture was allowed to stir at ambient temperature for 2 hours. It was treated with 2M NaOH to adjust the pH to ~10 and then the mixture was extracted into EtOAc (3 x 200 mL), the organic phase was separated, dried (MgSO₄) and concentrated to leave a gum, 4.05 g (12.26 mmol, 93%) which was used directly in the next stage; ¹H NMR (CDCl₃) δ 1.97-2.10 (m, 2H), 2.24-2.29 (m, 2H), 2.49-2.53 (m, 4H), 3.13-3.21 (m, 1H), 5.22 (s, 2H), 6.82 (s, 1H), 7.18 (d, 1H), 7.36-7.40 (m, 5H), 7.96 (s, 1H), 8.41 (s, 1H); MS m/e MH⁺ 325.


vii) Methyl [4-(5-(((benzyloxy)carbonyl)amino)pyridin-2-yl)cyclohexylidene]acetate



To an ice water cooled suspension of a 60% mineral oil dispersion of sodium hydride (585 mg, 24.35 mmol) in anhydrous THF (60 mL) was added trimethylphosphonoacetate (3.1 mL, 18.73 mmol). After 10 minutes a solution of benzyl [6-(4-oxocyclohexyl)pyridin-3-yl]carbamate, prepared as above, (4.05 g, 12.49 mmol) in THF (40 mL) was added and the reaction mixture allowed to warm to ambient temperature over the weekend. Water (100 mL) was added and the volatile material was removed under reduced pressure and the mixture extracted into EtOAc (4 x 200 mL) the organic extracts were combined, washed with brine (50 mL), dried (MgSO₄) and concentrated under reduced pressure to leave crude product. This was purified on a 120 g silica SiliCycle cartridge, loading in and eluting 30-50% EtOAc/ isohexane to provide the title compound (3.17 g, 66.7%); ¹H NMR (CDCl₃) δ 1.63-1.76 (m, 2H), 2.03-2.15 (m, 3H), 2.31-2.44 (m, 2H), 2.92-2.97 (m, 1H), 3.69 (s, 3H), 3.92-3.96 (m, 1H), 5.21 (s, 2H), 5.68 (s, 1H), 6.71 (s, 1H), 7.12 (d, 1H), 7.35-7.40 (m, 5H), 8.37 (s, 1H), 8.37 (s, 1H); MS m/e MH⁺ 381.

viii) Methyl [*trans*-4-(5-aminopyridin-2-yl)cyclohexyl]acetate hydrochloride salt

To a stirred solution of methyl [4-(5-{{[(benzyloxy)carbonyl]amino}}pyridin-2-yl)cyclohexylidene]acetate, prepared as above, (3.17 g, 8.33 mmol) in methanol (200 mL) was added 50% wet Palladium on carbon (1.0 g), deoxygenated by evacuation and placed under an atmosphere of hydrogen (5 cycles) then stirred for 16 h. The mixture was filtered and the filtrate concentrated to give a gum (2.05 g). The residue was dissolved in DCM/ether (40 mL of a 1:1 mixture) and HCl (8.5 mL of a 2M solution in ether) was added with stirring. The suspension was allowed to stir at ambient temperature and concentrated to provide the salt as a white solid. This was recrystallised from absolute ethanol-ether to provide the title compound (1.10 g, 54%) as a solid; ¹H NMR δ 1.07-1.21 (m, 2H), 1.52-1.62 (m, 2H), 1.72-1.88 (m, 5H), 2.27 (d, 2H), 2.74-2.81 (m, 1H), 3.61 (s, 3H), 7.59-7.59 (m, 2H), 7.87 (d, 1H); MS m/e MH⁺ 249.


ix) Methyl ({6-[*trans*-4-(2-methoxy-2-oxoethyl)cyclohexyl]pyridin-3-yl}amino)(oxo)acetate

To an ice water cooled solution of methyl [*trans*-4-(5-aminopyridin-2-yl)cyclohexyl]acetate hydrochloride, prepared as above, (831 mg, 2.59 mmol) in DCM (50 mL) was added triethylamine (1.3 mL, 9.31 mmol) followed by methyl chlorooxoacetate (0.31 mL, 3.36 mmol) and the reaction mixture was allowed to stir at ~0°C for 2 hrs. The reaction mixture was diluted with DCM (100 mL) and washed with water (80 mL) then brine (80 mL) and the aqueous layer was re extracted with DCM (50 mL). The organic washings were combined, dried and concentrated to give the title compound, 865 mg (2.59 mmol, 100%) which was taken through to the next stage; ¹H NMR (CDCl₃) δ 1.13-1.27 (m, 1H), 1.57-1.64 (m, 3H), 1.84-2.00 (m, 5H), 2.26 (d, 2H), 2.62-2.70 (m, 1H), 3.68 (s, 3H), 3.98 (s, 3H), 7.18 (d, 1H), 8.13 (d, 1H), 8.60 (s, 1H), 8.82 (brs, 1H); MS m/e MH⁺ 335.


x) To a suspension of methyl ((6-{4-(2-methoxy-2-oxoethyl)cyclohexyl}pyridin-3-yl)amino)(oxo)acetate, prepared as above, (865 mg, 2.59 mmol) in absolute ethanol (50 mL) was added hydrazine monohydrate (0.15 mL, 3.09 mmol). A thick white suspension formed which was then allowed to stir at ambient temperature for ~1.5 hrs, filtered, washed with ether (2 x 20 mL) and then dried to give the title compound (Intermediate 78), 610 mg (1.82 mmol, 70%) as a solid; ^1H NMR δ 1.09-1.21 (m, 2H), 1.47-1.56 (m, 2H), 1.71-1.87 (m, 5H), 2.25 (d, 2H), 2.54-2.63 (m, 1H), 3.60 (s, 3H), 4.62 (s, 2H), 7.24 (d, 1H), 8.07 (d, 1H), 8.85 (s, 1H), 10.27 (s, 1H) (not all NH protons seen); MS m/e MH^+ 335.

Intermediate 79: *tert*-Butyl *trans*-4-(4-

***hydrazino(oxo)acetyl*(amino)phenyl)cyclohexanecarboxylate**




i) *trans*-4-Phenylcyclohexanecarboxylic acid



Palladium (10 wt. %) on carbon (50 g) and triethylamine (90 mL, 0.65 mol) were added sequentially, each in one portion, to a stirred solution of 4-(4-chlorophenyl)cyclohexanecarboxylic acid (150 g, 0.63 mol) in MeOH (1.5 L). The reaction mixture was stirred under a hydrogen atmosphere for 4 h at 50°C at a pressure of 5 bar then cooled, filtered and then concentrated *in vacuo* to give the title compound (160 g, 83%) as a solid, which was contaminated with triethylamine hydrochloride (0.65 mol) and used with no further purification; ^1H NMR δ 1.23 (9H, t), 1.44-1.53 (4H, m), 1.8-1.9 (2H, m), 1.96-2.05 (2H, m), 2.22-2.32 (1H, m), 2.45-2.58 (1H, m), 3.05 (6H, q), 7.1-7.35 (5H, m).


ii) *trans*-4-(4-Nitrophenyl)cyclohexanecarboxylic acid



A mixture of concentrated sulphuric acid (1.5 mL) and concentrated nitric acid (0.9 mL) was added dropwise to a vigorously stirred solution of *trans*-4-phenylcyclohexanecarboxylic acid (1.7 g, 8.3 mmol) in nitrobenzene (4 mL) at 5°C. The reaction mixture was warmed to room temperature and stirred for 2.5 h and then poured


onto ice. The aqueous mixture was extracted with DCM and the organic layer was concentrated *in vacuo* to leave a yellow oil, which crystallised to a white solid upon standing. The solid was triturated with toluene and filtered to give the title compound (600 mg, 60%) as a solid; ^1H NMR δ 1.4-1.6 (4H, m), 1.75-1.9 (2H, m), 1.95-2.1 (2H, m), 2.2-2.38 (1H, m), 2.64-2.77 (1H, m), 7.52 (2H, d), 8.15 (2H, d); MS m/e (M-H) $^+$ 248.

iii) *tert*-Butyl *trans*-4-(4-nitrophenyl)cyclohexanecarboxylate



4-(Dimethylamino)pyridine (9.7 g, 80.2 mmol) and a solution of di-*tert*-butyl dicarbonate (17.5 g, 80.2 mmol) in DCM (5 mL) were added sequentially, each in one portion, to a mixture of *trans*-4-(4-nitrophenyl)cyclohexanecarboxylic acid (10 g, 40.1 mmol) in DCM (45 mL) at room temperature. The reaction was stirred for 16 h and then evaporated to leave a yellow oil. The oil was triturated with a mixture of isohexane:EtOAc (4:1) and filtered. The filtrate was concentrated *in vacuo* and purified by chromatography, eluting with 20-33% EtOAc in isohexane, to give the title compound as a waxy solid (4.3 g, 35%); ^1H NMR δ 1.4 (9H, s), 1.4-1.6 (4H, m), 1.8-1.9 (2H, m), 1.95-2.05 (2H, m), 2.2-2.35 (1H, m), 2.65-2.75 (1H, m), 7.55 (2H, d), 8.18 (2H, d).


iv) *tert*-Butyl *trans*-4-(4-aminophenyl)cyclohexanecarboxylate



10% Palladium on charcoal (500 mg) was added in one portion to a stirred solution of *tert*-butyl *trans*-4-(4-nitrophenyl)cyclohexanecarboxylate (4.3 g, 14.1 mmol) in EtOAc (75 mL) and the mixture was stirred under a hydrogen atmosphere for 20 h. The mixture was filtered through a pad of diatomaceous earth and the filtrate was concentrated *in vacuo* to give the title compound (3.8 g, 98%) as a waxy solid; ^1H NMR δ 1.4 (9H, s), 1.3-1.5 (4H, m), 1.7-1.8 (2H, m), 1.9-2.0 (2H, m), 2.15-2.25 (1H, m), 2.25-2.35 (1H, m), 4.75 (2H, s), 6.5 (2H, d), 6.85 (2H, d); MS m/e MH $^+$ 276.

251


v) *tert*-Butyl *trans*-4-(4-([methoxy(oxo)acetyl]amino)phenyl)cyclohexanecarboxylate




Methyl chlorooxoacetate (2.5 g, 20.7 mmol) was added dropwise to a mixture of *tert*-butyl *trans*-4-(4-aminophenyl)cyclohexanecarboxylate (3.8 g, 13.8 mmol) and pyridine (2.2 mL, 27.6 mmol) in DCM (40 mL) at 10°C. The reaction mixture was warmed to room temperature and stirred for 16 h and then quenched with water. The layers were separated and the organic layer was washed with brine. The organic layer was dried and concentrated *in vacuo* and the residue was purified by chromatography, eluting with 10-50% ethyl acetate in isohexane, to give the title compound (4.0 g, 80%) as an oil; ¹H NMR δ 1.4 (9H, s), 1.4-1.55 (4H, m), 1.8-1.9 (2H, m), 1.9-2.05 (2H, m), 2.2-2.3 (1H, m), 2.4-2.6 (1H, m), 3.75 (3H, s), 7.2 (2H, d), 7.65 (2H, d); MS *m/e* (M-H)⁺ 360.

vi) Hydrazine hydrate (0.59 mL, 12.2 mmol) was added in one portion to a stirred mixture of *tert*-butyl *trans*-4-(4-([methoxy(oxo)acetyl]amino)phenyl)cyclohexanecarboxylate (4.0 g, 11.1 mmol) in EtOH (50 mL) and the reaction mixture was stirred at room temperature for 3 h. EtOH (25 mL) was added and the mixture was filtered to give the title compound (Intermediate 79) (3.7 g, 92%) as a solid; ¹H NMR δ 1.4 (9H, s), 1.4-1.5 (4H, m), 1.8-1.9 (2H, m), 1.9-2.0 (2H, m), 2.15-2.3 (1H, m), 2.4-2.55 (1H, m), 4.5 (1H, s), 4.6 (2H, s), 7.2 (2H, d), 7.7 (2H, d), 10.45 (1H, s); MS *m/e* (M-H)⁺ 360.

20 Intermediate 80: Ethyl *trans*-4-(4-([hydrazino(oxo)acetyl]amino)phenoxy)cyclohexanecarboxylate



i) Ethyl *trans*-4-(4-aminophenoxy)cyclohexanecarboxylate



- 345 -

352

A solution of ethyl 4-hydroxycyclohexanecarboxylate (2.0 g, 11.6 mmol) in DMF (3mL) was added in one portion to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 490 mg, 12.2 mmol) in DMF (15 mL) at 0°C under an argon atmosphere. The mixture was stirred at 0°C for 10 minutes and then warmed to room temperature and

5 stirred for 20 minutes and then a solution of 1-fluoro-4-nitrobenzene (1.24 mL, 11.6 mmol) in DMF (2mL) was added in one portion. The reaction mixture was stirred at room temperature for 2 h and then heated at 110°C for 24 h. The solution was concentrated *in vacuo* to leave a residue that was taken up in EtOAc (250 mL) and water (150 mL) was added. The layers were separated and the organic layer was washed with H₂O (2x100 mL)

10 and brine (50 mL) and then dried (MgSO₄) and concentrated *in vacuo* to leave an oil. The oil was taken up in a mixture of hot EtOAc and isohexane (1:1) and left for 20 h. Filtration, followed by concentration *in vacuo* of the filtrate left an oil, which was purified by column chromatography, using 10-40% EtOAc in isohexane as eluent, to give crude ethyl 4-(4-nitrophenoxy)cyclohexanecarboxylate as an off white solid that was used


15 without further purification.

Palladium (10 wt. %) on carbon (50 mg) was added in one portion to a solution of crude ethyl 4-(4-nitrophenoxy)cyclohexanecarboxylate in a mixture of EtOH (15 mL) and THF (10 mL). The reaction mixture was stirred under a hydrogen atmosphere at room temperature for 16 h then filtered and concentrated *in vacuo* to give a solid that was

20 purified by chromatography, using 30-60% EtOAc in isohexane as eluent, to give the title compound (127 mg, 4% over 2 steps) as a solid; ¹H NMR (CDCl₃) δ 1.16-1.20 (3H, t), 1.31-1.42 (2H, m), 1.43-1.52 (2H, m), 1.96-2.01 (2H, m), 2.05-2.10 (2H, m), 2.20-2.28 (1H, m), 3.35 (2H, br s), 3.89-3.96 (1H, m), 4.05 (2H, q), 6.53-6.57 (2H, d), 6.65-6.69 (2H, d); MS m/e MH⁺ 264.

25

ii) Ethyl *trans*-4-(4-([methoxy(oxo)acetyl]amino)phenoxy)cyclohexanecarboxylate



Methyl chlorooxoacetate (45 µL, 0.48 mmol) was added in one portion to a stirred solution of ethyl 4-(4-aminophenoxy)cyclohexanecarboxylate (127 mg, 0.48 mmol) and pyridine


30 (47 µL, 0.58 mmol) in DCM (5 mL) and the reaction mixture was stirred for 30 minutes.

- 346 -


Volatile material was removed by evaporation and the residue was taken up in EtOAc (40 mL) and an aqueous solution of hydrochloric acid (1M, 20 mL) was added. The layers were separated and the organic layer was washed with brine (10 mL), dried and concentrated *in vacuo* to give the title compound (152 mg, 91%) as a solid; ¹H NMR (CDCl₃) δ 1.19 (3H, t), 1.39-1.45 (2H, m), 1.48-1.56 (2H, m), 1.97-2.03 (2H, m), 2.08-2.12 (2H, m), 2.23-2.31 (1H, m), 3.89 (3H, s), 4.07 (2H, q), 4.11-4.14 (1H, m), 6.80-6.84 (2H, d), 7.44-7.48 (2H, d), 8.66 (1H, s); MS m/e MH⁺ 350.

iii) Hydrazine monohydrate (0.15 mL, 3.34 mmol) was added in one portion to a stirred solution of ethyl *trans*-4-(4-
10 {[methoxy(oxo)acetyl]amino}phenoxy)cyclohexanecarboxylate (150 mg, 3.04 mmol) in EtOH (3 mL) and the reaction mixture was stirred for 20 minutes. Ether (10 mL) was added and the mixture was cooled to 5°C. The mixture was filtered to give the title compound (Intermediate 80) (129 mg, 85%) as a solid that was used without further
15 purification; MS m/e MH⁺ 350.

Intermediate 81: 3-[[[(1R)-2-Methoxy-1-methylethyl]oxy]aniline



i) 1-[[[(1R)-2-Methoxy-1-methylethyl]oxy]-3-nitrobenzene



DEAD (1.97 mL, 10.8 mmol) was added dropwise to a stirred mixture of 3-nitrophenol (2.50g, 18 mmol), (2S)-1-methoxypropan-2-ol (648 mg, 7.2 mmol) and triphenylphosphine (2.84 g, 10.8 mmol) in DCM (10ml). The reaction mixture was stirred for 16 h then treated with saturated aqueous sodium hydrogen carbonate solution. The resultant mixture was partitioned and the organics were washed with 1M NaOH, dried and concentrated by
20 evaporation. The residue was triturated with 1:1 EtOAc/hexanes, and filtered. The filtrate was concentrated by evaporation and the residue was purified by chromatography eluting with 25% EtOAc/hexane to give the title compound (1 g) as an oil; ¹H NMR (400


- 347 -

351
 MHz, DMSO) δ 1.24 (d, 3H), 3.24 (s, 3H), 3.44-3.60 (m, 2H), 4.71-4.89 (m, 1H), 7.38-7.48 (m, 1H), 7.57 (t, 1H), 7.70-7.76 (m, 1H), 7.76-7.86 (m, 1H).


- ii) 1-{{{1*R*}-2-methoxy-1-methylethyl}oxy}-3-nitrobenzene (1.00 g; 4.74 mmol) was dissolved in MeOH (20ml) and treated with 10% palladium on carbon then placed under an atmosphere of hydrogen and stirred for 24 h. The reaction mixture was filtered then concentrated by evaporation to give the title compound (Intermediate 81) (0.88 g) as an oil; $^1\text{H NMR } \delta$ 1.18 (d, 3H), 3.24 (s, 3H), 3.34-3.42 (m, 1H), 3.42-3.52 (m, 1H), 4.37-4.49 (m, 1H), 4.90 (s, 2H), 6.04-6.11 (m, 1H), 6.10-6.18 (m, 2H), 6.83-6.91 (m, 1H); MH^+ 182.

10

Intermediate 82: 3-[[*(1S)*-2-Methoxy-1-methylethyl]oxy]aniline




i) 1-{{{1*S*}-2-Methoxy-1-methylethyl}oxy}-3-nitrobenzene




- Following the procedure described for Intermediate 81 part (i) except that (2*R*)-1-methoxypropan-2-ol was used in place of (2*S*)-1-methoxypropan-2-ol, the title compound was obtained as an oil; $^1\text{H NMR } \delta$ 1.25 (d, 3H), 3.26 (s, 3H), 3.43-3.61 (m, 2H), 4.69-4.88 (m, 1H), 7.40-7.47 (m, 1H), 7.57 (t, 1H), 7.69-7.75 (m, 1H), 7.76-7.83 (m, 1H).
- ii) Reduction of 1-{{{1*S*}-2-Methoxy-1-methylethyl}oxy}-3-nitrobenzene following the procedure described for Intermediate 81 part (ii) gave the title compound (Intermediate 82) as an oil; $^1\text{H NMR } \delta$ 1.17 (d, 3H), 3.26 (s, 3H), 3.35-3.42 (m, 1H), 3.42-3.51 (m, 1H), 4.37-4.49 (m, 1H), 4.90 (s, 2H), 6.03-6.11 (m, 1H), 6.10-6.19 (m, 2H), 6.81-6.94 (m, 1H); MS $m/e \text{ MH}^+$ 182.

25

Intermediate 83: *tert*-Butyl 1-(5-[[hydrazino(oxo)acetyl]amino]pyridin-2-yl)piperidine-4-carboxylate




i) *tert*-Butyl 1-(5-nitropyridin-2-yl)piperidine-4-carboxylate



Following the procedure described for Intermediate 77 part (i), but using *tert*-butylpiperidine-4-carboxylate hydrochloride in place of methyl piperidin-4-ylacetate

- 5 hydrochloride, the title compound was obtained in 91% yield; ^1H NMR δ 1.35 (s, 9H), 1.51 (q, 2H), 1.90 (d, 2H), 2.55-2.69 (m, 1H), 3.20 (t, 2H), 4.39 (d, 2H), 6.96 (d, 1H), 8.20 (d, 1H), 8.95 (d, 1H); MS m/e MH^+ 252.

ii) *tert*-Butyl 1-(5-aminopyridin-2-yl)piperidine-4-carboxylate



10


Following the procedure described for Intermediate 77 part (ii), the title compound was obtained in 95% yield as a solid; ^1H NMR (CDCl_3) δ 1.41 (s, 9H), 1.68-1.85 (m, 2H), 1.89-2.02 (m, 2H), 2.28-2.44 (m, 1H), 2.74-2.87 (m, 2H), 3.91-4.06 (m, 2H), 6.64 (d, 1H), 6.98 (d, 1H), 7.79 (d, 1H); MS m/e MH^+ 252.


15

iii) Following the procedure described for Intermediate 77 part (iii), the first intermediate, *tert*-Butyl 1-(5-([methoxy(oxo)acetyl]amino)pyridin-2-yl)piperidine-4-carboxylate, was obtained in 97% yield as a solid; ^1H NMR (CDCl_3) δ 1.39 (s, 9H), 1.64-1.82 (m, 2H), 1.96 (d, 2H), 2.36-2.52 (m, 1H), 2.98 (t, 2H), 3.95 (s, 3H), 4.10-4.26 (m, 2H), 6.68 (d, 1H), 7.94 (d, 1H), 8.29 (d, 1H), 8.69 (s, 1H); MS m/e MH^+ 364; then condensation with hydrazine gave the title compound (Intermediate 83) in 72% yield as a solid; ^1H NMR δ 1.35 (s, 9H), 1.44-1.61 (m, 2H), 1.75-1.88 (m, 2H), 2.38-2.49 (m, 1H), 2.90 (t, 2H), 4.12 (d, 2H), 4.62 (s, 2H), 6.84 (d, 1H), 7.91 (d, 1H), 8.51 (d, 1H), 10.23 (s, 1H), 10.53 (s, 1H); MS m/e MH^+ 364.

25



Intermediate 84: Methyl 3-(4-aminophenyl)cyclopentanecarboxylate



i) Methyl 3-(4-nitrophenyl)cyclopentanecarboxylate

A mixture of 65% nitric acid (0.95 mL) and 95% sulphuric acid (1.2 mL) in carbon tetrachloride (2.5 mL) was added dropwise to a solution of methyl 3-phenylcyclopentanecarboxylate (Journal of Organic Chemistry, 1988, 53(17), 4021) (1 g, 4.9 mmol) and the mixture was stirred at 5-10°C for 2 h. Ice/water (30 mL) was added and the mixture was extracted with DCM (2 x 50 mL). The extracts were combined, washed with water and dried. The residue was adsorbed onto silica and purified by chromatography eluting with 0-30% EtOAc in hexane to give the title compound (782 mg, 64%) as an oil; ¹H NMR δ 3.90 (3H, s), 7.94-7.97 (2H, m), 8.03-8.06 (2H, m), 8.09-8.12 (2H, m), 8.33-8.36 (2H, m).

ii) 10% palladium on carbon (78 mg) was added to a stirred solution of methyl 3-(4-nitrophenyl)cyclopentanecarboxylate (782 mg, 3.14 mmol) in EtOH (40 mL). The mixture was stirred under an atmosphere of hydrogen for 16 h then filtered through diatomaceous earth and the filter washed with EtOAc. Volatile material was removed by evaporation to give the title compound (Intermediate 84) (608 mg, 88%) as an oil; ¹H NMR δ 2.24-1.45 (m, 6H), 3.03-2.83 (m, 2H), 3.62 (s, 3H), 4.80 (s, 2H), 6.53-6.46 (m, 2H), 6.92-6.86 (m, 2H); MS m/e MH⁺ 220.

Intermediate 85: Methyl 3-(4-aminophenyl)adamantane-1-carboxylate**i) Methyl 3-(4-Nitrophenyl)adamantane-1-carboxylate**

- 350 -


3-(4-Nitrophenyl)adamantane-1-carboxylic acid (500 mg, 1.66 mmol) was stirred in MeOH (8 mL) and concentrated sulphuric acid (0.1 mL) was added. After stirring at room temperature overnight the reaction mixture was basified with aqueous potassium carbonate (30 mL of a saturated solution) and then concentrated in vacuo. The residue was extracted with EtOAc (2 x 250 mL) and washed with water and brine. The organics were dried and solvent removed in vacuo to give a white solid (554 mg, 100%); ^1H NMR δ 8.18 (d, 2H), 7.68 (d, 2H), 3.61 (s, 3H), 2.23-2.17 (m, 2H), 2.01-1.97 (m, 2H), 1.91-1.84 (m, 8H), 1.74-1.69 (m, 2H); MS m/e M+MeCN 357.

ii) Methyl 3-(4-Nitrophenyl)adamantane-1-carboxylate was reduced by the method of Intermediate 84, step (ii), to give the title compound (Intermediate 85); ^1H NMR δ 6.99 (d, 2H), 6.51 (d, 2H), 4.80 (s, 2H), 3.59 (s, 3H), 2.17-2.10 (m, 2H), 1.87-1.79 (m, 6H), 1.77-1.72 (m, 4H), 1.69-1.63 (m, 2H); MS m/e M+MeCN 327.

15 **Intermediate 86: Ethyl 4'-aminobiphenyl-3-yl]acetate**



i) **Ethyl [4'-(dibenzylamino)biphenyl-3-yl]acetate**



20 [4-(Dibenzylamino)phenyl]boronic acid (UK Patent Application No. GB 2276161 (1994)) (400 mL, 1.26mmol) was stirred in 5 mL dry toluene and potassium carbonate was added followed by ethyl (3-bromophenyl)acetate (338mg, 1.39mmol) as a solution in toluene (1 mL). The reaction vessel was evacuated and refilled with nitrogen (5 times). Pd(PPh₃)₄ (73
25 mg, 0.06 mmol) was added as a solution in toluene (1 mL) and the vessel was deoxygenated as before. The reaction mixture was heated at 90°C for 16 h then cooled and filtered through diatomaceous earth. Volatile material was removed by evaporation, the residue was adsorbed onto silica and purified by chromatography eluting with 10-50%


- 351 -

~~238~~

EtOAc/isohexane to give the title compound (175mg, 32%) as a gum; ^1H NMR δ 1.18 (t, 3H), 3.67 (s, 2H), 4.07 (q, 2H), 4.76 (s, 4H), 6.76 (d, 2H), 7.11 (d, 2H), 7.44-7.20 (m, 14H); MS m/e MH^+ 436.


- 5 ii) Nitrogen was bubbled through a solution of ethyl [4'-(dibenzylamino)biphenyl-3-yl]acetate (170 mg, 0.39 mmol) in EtOH (15 mL) and EtOAc (15 mL) then 10% palladium on carbon (30 mg) was added and the mixture was stirred under an atmosphere of hydrogen for 16 h. The mixture was filtered through diatomaceous earth and concentrated by evaporation to give the title compound (Intermediate 86) (98 mg) as a gum; ^1H NMR δ
- 10 1.19 (t, 3H), 3.69 (s, 2H), 4.09 (q, 2H), 5.24 (s, 2H), 6.64 (d, 2H), 7.11 (d, 1H), 7.37-7.30 (m, 3H), 7.42 (d, 2H); MS m/e $\text{M}+\text{MeCN}$ 297.

Intermediate 87: Methyl [(1*R*,4*R*)-4'-amino-1,1'-bi(cyclohexyl)-4-yl]acetate




- A mixture of methyl [*trans*-4-(4-aminophenyl)cyclohexyl]acetate (from Intermediate 43, step (ii)) (200 mg, 0.81 mmol) and Rhodium (5% on alumina; 15 mg) in EtOH/water (6
- 15 mL of a 1:2 mixture) was stirred under an atmosphere of hydrogen at 48 bar, 65°C for 3h then filtered. The filtrate was concentrated by evaporation to give the title compound as a gum.

- 20 **Intermediate 88: Ethyl 3-(2-amino-2,3-dihydro-1*H*-inden-5-yl)propanoate**




i) Benzyl (5-bromo-2,3-dihydro-1*H*-inden-2-yl)carbamate



- Benzyl chloroformate (0.804 mL, 5.63 mmol) was added slowly with stirring to a mixture
- 25 of 5-bromoindan-2-amine hydrobromide (Advanced Synth. & Catalysis 343(5), 461-472, 2001) (1.5g, 5.12 mmol) and triethylamine (1.57 mL, 11.3 mmol) in DCM (20 mL) at 0°C.



The reaction mixture was allowed to warm to room temperature and stirred for 16 h. DCM (150 mL) was added and the solution was washed with 0.5M aqueous HCl (100 mL) then saturated aqueous sodium hydrogencarbonate solution (100 mL). MeOH (10 mL) was added as cosolvent and the solution was dried and concentrated by evaporation. The residue was purified by chromatography to give the title compound (804 mg, 45%) as a solid; ^1H NMR δ 2.59-2.87 (m, 2H), 3.03-3.23 (m, 2H), 4.23-4.39 (m, 1H), 4.50 (d, 1H), 5.04 (s, 2H), 7.14-7.44 (m, 8H); MS m/e MH^+ 214.

ii) Ethyl (2*E*)-3-(2-((benzyloxy)carbonyl)amino)-2,3-dihydro-1*H*-inden-5-yl)acrylate




A mixture of palladium(II) acetate (3 mg, 0.01 mmol), Tri-*o*-tolylphosphine (7 mg, 0.02 mmol) and DIPEA (0.088 mL, 0.5 mmol) was stirred in MeCN (2 mL) at 60°C under a nitrogen atmosphere for 20 minutes in a 5 mL microwave vial. Benzyloxycarbonyl(5-bromo-2,3-dihydro-1*H*-inden-2-yl)carbamate (174 mg, 0.5 mmol) and ethyl acrylate (28 μL , 0.25 mmol) were added and the mixture was heated in the microwave at 180°C for 0.5 h. The mixture was adsorbed onto silica and purified by chromatography with 0-40% EtOAc/isohexane as eluent to give the title compound (105mg, 57%) as a gum; ^1H NMR (CDCl_3) δ 1.32 (t, 3H), 2.62-2.88 (m, 2H), 3.12-3.36 (m, 2H), 4.18-4.29 (m, 2H), 4.46-4.62 (m, 1H), 4.76 (d, 1H), 5.09 (s, 2H), 6.32-6.41 (m, 1H), 7.10-7.44 (m, 8H), 7.57-7.68 (m, 1H); MS m/e MH^+ 366.

iii) Ethyl (2*E*)-3-(2-((benzyloxy)carbonyl)amino)-2,3-dihydro-1*H*-inden-5-yl)acrylate was reduced by the general procedure of Intermediate 84, step (ii), to give the title compound (Intermediate 88); ^1H NMR (CD_3OD) δ 1.01-1.18 (m, 3H), 2.44-2.70 (m, 2H), 2.73-2.96 (m, 2H), 3.02-3.35 (m, 4H), 3.92-4.05 (m, 2H), 4.30-4.42 (m, 1H), 6.84-7.23 (m, 3H); MS m/e MH^+ 234.


Intermediate 89: Ethyl [3-(4-aminophenyl)cyclohexyl]acetate**i) 3-[4-(Benzyloxy)phenyl]cyclohexanone**

- 5 [4-(Benzyloxy)phenyl]boronic acid (2 g, 8.77 mmol) was dissolved in dioxane/water (50 mL / 5 mL) and Rh(acac)(C₂H₄)₂ (68 mg, 0.26 mmol) and racemic BINAP (164 mg, 0.26 mmol) were added. The mixture was degassed 3 times and put under a nitrogen atmosphere. Cyclohexenone (1.02 mL, 10.52 mmol) was added to the yellow solution and the mixture heated at reflux overnight. The reaction mixture was cooled, saturated
- 10 aqueous sodium hydrogen carbonate solution (50 mL) added, then extracted with EtOAc (2 x 100 mL). The organic extracts were washed with water (100 mL) and dried. Volatile material was removed by evaporation and purification by chromatography gave the title compound (565 mg, 23%) as a solid; ¹H NMR 8.1.65-1.69 (1H, m), 1.78-1.93 (2H, m), 1.99-2.04 (1H, m), 2.22-2.35 (2H, m), 2.39-2.43 (1H, m), 2.65 (1H, m), 2.90-2.96 (1H, m), 5.08 (2H, s), 6.96 (2H, d), 7.20 (2H, d), 7.31-7.45 (5H, m); MS m/e MH⁺ 281.
- 15

ii) Ethyl [3-[4-(benzyloxy)phenyl]cyclohexylidene]acetate


- Triethylphosphonoacetate (0.48 mL, 2.42 mmol) was added to a stirred suspension of
- 20 sodium hydride (97 mg of a 60% dispersion in oil, 2.42 mmol) in THF (10 mL) at 0°C. The mixture was allowed to warm to ambient temperature and stirred for 15 minutes then 3-[4-(benzyloxy)phenyl]cyclohexanone (565 mg, 2.02 mmol) was added as a solution in THF (10 mL). After 3 h saturated aqueous ammonium chloride solution (50 mL) was added and the mixture was extracted with EtOAc (2 x 50 mL). The organic layers were
- 25 washed with water (50 mL) then dried and concentrated by evaporation to give the title compound as an oil that was used directly in the next step; MS m/e MH⁺ 351.

iii) Ethyl 3-(4-(4-hydroxyphenyl)cyclohexyl)acetate




Ethyl 3-[4-(benzyloxy)phenyl]cyclohexylidene} acetate was reduced following the general procedure described for Intermediate 84, part (ii), to give the title compound; ¹H NMR δ 1.12-1.19 (3H, m), 0.93-1.82 (8H, m), 2.18 (1H, d), 2.45 (1H, d), 2.65 (1H, m), 2.99 (1H, m), 4.04 (2H, q), 6.65-6.67 (2H, m), 7.00 (2H, t), 9.12 (1H, s); MS m/e (M-H)⁺ 261.

iv) Ethyl 3-(4-((trifluoromethyl)sulphonyl)oxy)phenyl)cyclohexyl)acetate



10 Ethyl 3-(4-(4-hydroxyphenyl)cyclohexyl)acetate (329 mg, 1.25 mmol) was stirred in dry DCM (15 mL) under nitrogen. The mixture was cooled with an ice bath and trifluoromethanesulphonic anhydride (0.264 mL, 1.57 mmol) was added. Triethylamine (0.263 mL, 1.88 mmol) was then added dropwise over 5 minutes, keeping the temperature below 10°C and the mixture was stirred for 2 h. Saturated aqueous sodium hydrogen carbonate solution (10 mL) was added, the layers were separated and the aqueous phase was extracted with DCM (20 mL). The organic layers were dried and concentrated by evaporation to give an oil that was purified by chromatography eluting with 0-40% EtOAc/isohexane to give the title compound (345 mg, 70%) as an oil; ¹H NMR δ 0.94-2.00 (9H, s), 1.16 (3H, m), 2.23 (1H, d), 2.47 (1H, d), 2.62-2.85 (1H, m), 4.01-4.09 (2H, m), 7.41-7.44 (4H, m); MS m/e (M-H)⁺ 393.

v) Ethyl 3-(4-((diphenylmethylene)amino)phenyl)cyclohexyl)acetate



25 Ethyl 3-(4-((trifluoromethyl)sulphonyl)oxy)phenyl)cyclohexyl)acetate (322 mg, 0.85 mmol) was stirred in dry THF (8 mL) under a nitrogen atmosphere and caesium carbonate

~~26~~

(387 mg, 1.19 mmol), palladium(II) acetate (12 mg, 0.05 mmol) and BINAP (32 mg, 0.05 mmol) were added followed by benzophenone imine (0.214 mL, 1.27 mmol). The reaction vessel was degassed 6 times and stirred under reflux for 16 h. The mixture was extracted with ether (2 x 100 mL) and washed with water (100 mL). The organic extracts were dried and concentrated by evaporation to give an oil that was purified by chromatography eluting with 0-30% EtOAc/isohexane to give the title compound (137mg, 38%) as an oil; ¹H NMR δ 0.81-2.48 (m, 9H), 1.12-1.21 (m, 3H), 2.58-2.72 (m, 1H), 3.97-4.12 (m, 2H), 6.57-6.65 (m, 1H), 6.94-7.04 (m, 1H), 7.10-7.18 (m, 1H), 7.29-7.36 (m, 1H), 7.37-7.78 (m, 10H); MS m/e MH⁺ 426.

10

vi) Aqueous HCl (1 mL of a 1M solution) was added to ethyl (3-{4-[(diphenylmethylene)amino]phenyl}cyclohexyl)acetate (137 mg, 0.32 mmol) in EtOH (10 mL) with stirring. After 1 h the reaction mixture was basified with aqueous potassium carbonate (10 mL of a saturated solution) and the mixture was extracted with Et₂O (2 x 15 mL), washed with water (15 mL) and dried. Volatile material was removed by evaporation and the residue was purified by chromatography eluting with 0-35% EtOAc/isohexane to give the title compound (Intermediate 89) (40mg, 48%) as a gum; MS m/e MH⁺ 262.


15

Intermediate 90: Ethyl [(3R)-3-(4-aminophenyl)cyclopentyl]acetate

20



i) (3R)-3-(4-Bromophenyl)cyclopentanone




25

4-Bromophenylboronic acid (4 g, 19.92 mmol), Rh(acac)(C₂H₄)₂ (acetylacetonatobis(ethylene)Rhodium (I); 103 mg, 0.4 mmol) and (R)-BINAP (249 mg, 0.4 mmol) were placed in a 100 mL 3-necked round bottomed flask and 1,4-dioxane (40 mL) was added. The reaction vessel was degassed several times before putting under a nitrogen atmosphere. Water (4 mL) was added followed by cyclopent-2-en-1-one (1.185 mL, 14.14 mmol). The reaction mixture was then heated at 100°C overnight before cooling and washing with saturated aqueous sodium hydrogen carbonate solution (50 mL).

39


The mixture was extracted with EtOAc (2 x 100 mL) and the organics dried then purified by chromatography eluting with 0-40% EtOAc/isohexane) to give the title compound (1.62 g, 48%) as an oil that crystallised upon standing; $^1\text{H NMR}$ δ 1.86-1.93 (1H, m), 2.23-2.36 (3H, m), 2.53-2.57 (1H, m), 3.34-3.43 (1H, m), 7.29-7.32 (2H, m), 7.50-7.53 (2H, m); $\alpha_D = +20^\circ$ (MeOH, $c=1$, $L=1$)

ii) Ethyl [(3*R*)-3-(4-bromophenyl)cyclopentylidene]acetate



Following the general procedure described for ethyl [3-(4-(benzyloxy)phenyl)cyclohexylidene]acetate (Intermediate 89 (i)), replacing 3-(4-(benzyloxy)phenyl)cyclohexanone with (3*R*)-3-(4-bromophenyl)cyclopentanone the title compound was obtained; $^1\text{H NMR}$ δ 1.18-1.23 (3H, m), 1.61-1.78 (1H, m), 2.06-2.20 (1H, m), 2.53-2.67 (3H, m), 2.84-3.04 (1H, m), 3.08-3.24 (1H, m), 4.01-4.10 (2H, m), 5.82-5.85 (1H, m), 7.24-7.28 (2H, m), 7.47-7.51 (2H, m); MS m/e MH^+ 310.

iii) Ethyl [(3*R*)-3-{4-[(diphenylmethylene)amino]phenyl}cyclopentylidene]acetate



Following the procedure described for ethyl [3-(4-[(diphenylmethylene)amino]phenyl)cyclohexylidene]acetate (Intermediate 89(iii)), replacing ethyl [3-(4-[(trifluoromethyl)sulphonyloxy]phenyl)cyclohexylidene]acetate with ethyl [(3*R*)-3-(4-bromophenyl)cyclopentylidene]acetate, the title compound was obtained; MS m/e MH^+ 410.

iv) Ethyl [(3*R*)-3-(4-aminophenyl)cyclopentylidene]acetate



Following the procedure described for ethyl [3-(4-aminophenyl)cyclohexylidene]acetate (Intermediate 89iv), replacing ethyl [3-(4-

- 357 -

364


[(diphenylmethylene)amino]phenyl)cyclohexyl]acetate with ethyl ((3*R*)-3-{4-[(diphenylmethylene)amino]phenyl}cyclopentylidene)acetate the title compound was obtained; ¹H NMR δ 1.17-1.22 (3H, m), 1.57-1.68 (1H, m), 2.27-2.48 (2H, m), 2.53 (0H, s), 2.56-2.68 (1H, m), 2.74-2.76 (1H, m), 2.93-3.00 (1H, m), 3.15-3.24 (0.5H, m), 3.69-3.71 (0.5H, m), 4.01-4.12 (2H, m), 4.79-4.83 (2H, m), 5.49-5.53 (1H, m), 5.78-5.82 (1H, m), 6.47-6.52 (2H, m), 6.80-6.93 (2H, m); MS m/e M+MeCN 287.07

v) Ethyl [(3*R*)-3-(4-aminophenyl)cyclopentyl]acetate




Following the procedure described for methyl 3-(4-aminophenyl)cyclopentanecarboxylate, replacing methyl 3-(4-nitrophenyl)cyclopentanecarboxylate (Intermediate 84(ii)) with ethyl [(3*R*)-3-(4-aminophenyl)cyclopentylidene]acetate, the title compound was obtained; ¹H NMR δ 1.18 (t, 3H), 1.21-1.76 (m, 2H), 1.80-2.12 (m, 2H), 2.23-2.48 (m, 2H), 2.80-3.01 (m, 1H), 3.16-3.30 (m, 2H), 4.06 (q, 2H), 4.79 (s, 2H), 6.45-6.51 (m, 2H), 6.84-6.91 (m, 2H); MS m/e M+MeCN 289.

Intermediate 91: Ethyl [(3*S*)-3-(4-aminophenyl)cyclopentyl]acetate




i) (3*S*)-3-(4-Bromophenyl)cyclopentanone



Following the procedure described for (3*S*)-3-(4-Bromophenyl)cyclopentanone (Intermediate 90(i)), replacing (*R*)-BINAP with (*S*)-BINAP, the title compound was obtained; ¹H NMR δ 1.86-1.93 (1H, m), 2.23-2.36 (3H, m), 2.53-2.57 (1H, m), 3.34-3.43 (1H, m), 7.29-7.32 (2H, m), 7.50-7.53 (2H, m); α_D = -18° (MeOH, c=1, L=1)


ii) Ethyl [(3*S*)-3-(4-bromophenyl)cyclopentylidene]acetate



- 358 -


Following the procedure described for ethyl {3-[4-(benzyloxy)phenyl]cyclohexylidene}acetate (Intermediate 89(ii)), replacing 3-[4-(benzyloxy)phenyl]cyclohexanone with (3*S*)-3-(4-bromophenyl)cyclopentanone the title compound was obtained; ¹H NMR δ 1.18-1.23 (3H, m), 1.61-1.78 (1H, m), 2.06-2.20 (1H, m), 2.53-2.67 (3H, m), 2.84-3.04 (1H, m), 3.08-3.24 (1H, m), 4.01-4.10 (2H, m), 5.82-5.85 (1H, m), 7.24-7.28 (2H, m), 7.47-7.51 (2H, m); MS m/e MH⁺ 309.

iii) Ethyl ((3*S*)-3-{4-[(diphenylmethylene)amino]phenyl}cyclopentylidene)acetate




Following the procedure described for ethyl {3-[4-[(diphenylmethylene)amino]phenyl]cyclohexyl}acetate (Intermediate 89(iii)), replacing ethyl {3-[4-[(trifluoromethyl)sulphonyloxy]phenyl]cyclohexyl}acetate with ethyl [(3*S*)-3-(4-bromophenyl)cyclopentylidene]acetate the title compound was obtained; MS m/e MH⁺ 410.

iv) Ethyl ((3*S*)-3-(4-aminophenyl)cyclopentylidene)acetate




Following the procedure described for ethyl {3-(4-aminophenyl)cyclohexyl}acetate (Intermediate 89(iv)), replacing ethyl {3-[4-[(diphenylmethylene)amino]phenyl]cyclohexyl}acetate with ethyl ((3*S*)-3-{4-[(diphenylmethylene)amino]phenyl}cyclopentylidene)acetate the title compound was obtained; ¹H NMR δ 1.17-1.22 (3H, m), 1.57-1.68 (1H, m), 2.27-2.48 (2H, m), 2.53 (0H, s), 2.56-2.68 (1H, m), 2.74-2.76 (1H, m), 2.93-3.00 (1H, m), 3.15-3.24 (0.5H, m), 3.69-3.71 (0.5H, m), 4.01-4.12 (2H, m), 4.79-4.83 (2H, m), 5.49-5.53 (1H, m), 5.78-5.82 (1H, m), 6.47-6.52 (2H, m), 6.80-6.93 (2H, m); MS m/e M+MeCN 287.

v) Ethyl {(3*S*)-3-(4-aminophenyl)cyclopentyl}acetate




Following the procedure described for methyl 3-(4-aminophenyl)cyclopentanecarboxylate, replacing methyl 3-(4-nitrophenyl)cyclopentanecarboxylate (Intermediate 84(ii)) with ethyl
5 {(3*S*)-3-(4-aminophenyl)cyclopentylidene}acetate the title compound was obtained; ¹H NMR δ 1.18 (t, 3H), 1.21-1.76 (m, 2H), 1.80-2.12 (m, 2H), 2.23-2.48 (m, 2H), 2.80-3.01 (m, 1H), 3.16-3.30 (m, 2H), 4.06 (q, 2H), 4.79 (s, 2H), 6.45-6.51 (m, 2H), 6.84-6.91 (m, 2H); MS m/e M+MeCN 289.

10 Intermediate 92: Methyl 3-(3-aminopropoxy)benzoate



i) Methyl 3-(3-(((benzyloxy)carbonyl)amino)propoxy)benzoate




A mixture of methyl 3-hydroxybenzoate (609 mg, 4 mmol), benzyl *N*-(3-
15 hydroxypropyl)carbamate (754mg, 3.6mmol) and polymer supported triphenylphosphine (PS-PPh₃) (4 g, 12 mmol) in THF (40 mL) was stirred and diethyl azodicarboxylate (948 mg, 5.44 mmol) was added as a solution in THF (4 mL). The mixture was stirred for 16 h then was filtered and washed with DCM and concentrated by evaporation. The residue was purified by chromatography eluting with DCM to give the title compound (481mg, 35%)
20 as an oil; ¹H NMR δ 1.85-1.92 (2H, m), 3.19 (2H, q), 3.86 (3H, s), 4.05 (2H, t), 5.02 (2H, s), 7.20-7.22 (1H, m), 7.29-7.38 (5H, m), 7.42-7.45 (2H, m), 7.53-7.56 (1H, m).

ii) Following the procedure described for methyl 3-(4-aminophenyl)cyclopentanecarboxylate, replacing methyl 3-(4-nitrophenyl)cyclopentanecarboxylate (Intermediate 84 (i)) with methyl 3-(3-
25 (((benzyloxy)carbonyl)amino)propoxy)benzoate and using EtOAc as solvent instead of


- 360 -

EtOH the title compound was obtained; ^1H NMR δ 1.77-1.83 (2H, m), 2.70 (2H, t), 3.86 (3H, s), 4.09 (2H, t), 7.21-7.23 (1H, m), 7.43 (2H, t), 7.54 (1H, d); MS m/e MH^+ 210.

Intermediate 93: Ethyl 3-(4-aminobenzyl)cyclobutanecarboxylate




i) Ethyl 3-(4-nitrobenzyl)cyclobutanecarboxylate




- A mixture of concentrated nitric acid (0.5 mL) and concentrated sulphuric acid (0.8 mL) was added slowly with stirring to an ice-cooled solution of ethyl 3-benzylcyclobutanecarboxylate (German Patent DE 2626287) (876 mmol, 4.02 mmol) in nitrobenzene (2 mL). The reaction mixture was warmed to ambient temperature and stirred for 1.5 h then poured onto ice/water. DCM (5 mL) was added and the mixture was stirred before passing through a phase separation cartridge. Solvent was removed by evaporation to give the title compound that was used directly in the next step.
- ii) Ethyl 3-(4-nitrobenzyl)cyclobutanecarboxylate (4.02 mmol) was stirred in EtOAc (20 mL) at room temperature and nitrogen was bubbled through. 10% Palladium on carbon (160 mg) was added and the reaction vessel was put under an atmosphere of hydrogen. The mixture was stirred overnight then filtered through diatomaceous earth and concentrated by evaporation under reduced pressure. The residue was purified by chromatography eluting with 0-55% EtOAc/isohexane to give the title compound (466 mg, 50%) as an oil; ^1H NMR δ 1.15-1.20 (3H, m), 1.77-1.94 (2H, m), 2.13-2.20 (2H, m), 2.33-2.37 (2H, m), 2.43-2.46 (1H, m), 2.53-2.56 (1H, m), 2.92-3.07 (1H, m), 4.01-4.08 (2H, m), 4.77 (2H, d), 6.46-6.49 (2H, m), 6.78-6.82 (2H, m); MS m/e $\text{M}+\text{MeCN}$ 275.

Intermediate 94: Diethyl {4'-[(5-{(2-fluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}biphenyl-4-yl}malonate




i) Diethyl (4'-nitrobiphenyl-4-yl)malonate



5 A mixture of 4-(4-bromophenyl)nitrobenzene (Journal of Organic Chemistry, 2005, 70(9), 3730) (800 mg, 2.88 mmol), 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (Xphos) (69 mg) and caesium carbonate (2.81 g, 8.63 mmol) in toluene (40 mL) was deoxygenated then palladium(II) acetate (13 mg, 0.06 mmol) and diethyl malonate (0.480 mL, 3.2 mmol) were added. The reaction mixture was heated under reflux for 2 h, cooled and then filtered through diatomaceous earth and the filter washed with EtOAc. The filtrate was concentrated by evaporation and adsorbed onto silica then purified by chromatography, eluting with 0-25% EtOAc in isohexane, to give the title compound (1.047 g) as a solid.

15

ii) Diethyl (4'-aminobiphenyl-4-yl)malonate




Diethyl (4'-nitrobiphenyl-4-yl)malonate was reduced by the method of Intermediate 84, step (ii), to give the title compound; ^1H NMR δ 1.17-1.22 (6H, m), 4.10-4.22 (4H, m), 4.92 (1H, s), 5.22 (2H, s), 6.63-6.66 (2H, m), 7.35-7.38 (4H, m), 7.52-7.54 (2H, m); MS m/e MH^+ 328.

20


iii) Following the procedure of Example 483 except that diethyl (4'-aminobiphenyl-4-yl)malonate was used as starting material in place of methyl 4-(4-aminophenyl)butanoate, the title compound (Intermediate 94) was obtained as a solid; ^1H NMR δ 1.20 (6H, t), 4.15-4.22 (4H, m), 5.00 (1H, s), 7.17-7.20 (1H, m), 7.26-7.35 (2H, m), 7.47 (2H, d), 7.68-7.73 (4H, m), 7.92 (2H, d), 8.03-8.08 (1H, m), 10.79 (1H, s), 11.14 (1H, s); MS m/e MH^+ 533.

25

Intermediate 95: Diethyl (3-{4'-[({5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino}biphenyl-4-yl}propyl)malonate



i) Diethyl (3-(4-aminophenyl)propyl)malonate




Following the general procedure of Intermediate 84 except that diethyl (3-phenylpropyl)malonate (J. Med. Chem., 2004, 47(12), 3282) was used in place of methyl 3-phenylcyclopentanecarboxylate, nitration and reduction (using EtOAc as solvent in place of EtOH) gave the title compound (purified by chromatography, eluting with 10-50% EtOAc/Isohexane); $^1\text{H NMR}$ δ 1.17 (t, 6H), 1.42-1.53 (m, 2H), 1.69-1.78 (m, 2H), 2.40 (t, 2H), 3.45 (t, 1H), 4.07-4.16 (m, 4H), 4.81 (s, 2H), 6.48 (d, 2H), 6.81 (d, 2H); MS m/e MH^+ 294.

15


ii) Following the procedure of Example 483 except that diethyl (3-(4-aminophenyl)propyl)malonate was used as starting material in place of methyl 4-(4-aminophenyl)butanoate, the title compound (Intermediate 95) was obtained; MS m/e MH^+ 517.

20

Intermediate 96: Diisopropyl (4-{4'-[({5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl}carbonyl)amino}cyclohexyl}phenyl)malonate




i) 4-(4-Oxocyclohexyl)phenyl 4-methylbenzenesulphonate



25


4-(4-Hydroxyphenyl)cyclohexanone (2 g, 10.51 mmol) was stirred in 10 mL acetone with potassium carbonate (1.6g, 11.56 mmol) at room temperature. *p*-Toluenesulphonyl chloride (2.2 g, 11.56 mmol) was added (as a solution in 5 mL acetone) and the mixture was heated at reflux overnight and the reaction mixture was cooled and then quenched with water (20 mL). The mixture was extracted with EtOAc (2x30 mL) and the organics separated and dried (MgSO₄). The solvent was concentrated in vacuo to leave a colourless oil which yielded a white solid upon trituration with Et₂O and isohexane. The solid was filtered off and washed with isohexane. The solid was dissolved in EtOAc and passed through an Isolute NH₂ column. The product was isolated as a white crystalline solid (2.49g, 69%); ¹H NMR δ 1.78-1.88 (2H, m), 1.99-2.09 (2H, m), 2.23-2.28 (2H, m), 2.43 (3H, s), 2.53-2.58 (2H, m), 3.02-3.10 (1H, m), 6.94-6.98 (2H, m), 7.30-7.33 (2H, m), 7.48 (2H, d), 7.74-7.76 (2H, m); MS m/e M+Na⁺ 367.

ii) 4-(1,4-Dioxaspiro[4.5]dec-8-yl)phenyl 4-methylbenzenesulphonate



Boron trifluoride diethyletherate (3.3 mL, 26.0 mmol) was added dropwise via a syringe to a stirred solution of 4-(4-oxocyclohexyl)phenyl 4-methylbenzenesulphonate (2.49 g, 7.23 mmol) in dry DCM/ethylene glycol (15 mL/14.11 mL) cooled with an isopropanol/ice bath and stirred for 40 minutes. The mixture was poured onto water (50 mL) and extracted with DCM (2 x 100 mL). The organic phase was washed with 10% sodium hydrogen carbonate solution (100 mL) then dried and concentrated by evaporation to give the title compound (2.685 g, 96%) as a solid; ¹H NMR δ 1.54-1.66 (4H, m), 1.68-1.75 (4H, m), 2.43 (3H, s), 2.54-2.60 (1H, m), 3.86-3.89 (4H, s), 6.91-6.95 (2H, m), 7.21-7.25 (2H, m), 7.46-7.48 (2H, m), 7.73-7.75 (2H, m); MS m/e M+Na⁺ 411.

iii) Diethyl 4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl malonate




4-(1,4-Dioxaspiro[4.5]dec-8-yl)phenyl 4-methylbenzenesulphonate (2.05 g, 5.27 mmol) was stirred with 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (Xphos) (502 mg,

- 364 -

1.05 mmol) and caesium carbonate (5.15 g, 15.81 mmol) in toluene (40 mL) at room temperature under a nitrogen atmosphere. The reaction mixture was degassed several times then palladium(II) acetate (95 mg, 0.42 mmol) was added, followed by diethyl malonate (0.96 mL, 6.324 mmol). The mixture was heated under reflux for 40 h then was
5 filtered through diatomaceous earth and concentrated by evaporation. The residue was adsorbed onto silica and purified by chromatography eluting with 0-40% EtOAc/isohexane to give the title compound (294 mg, 15%) as an oil; ^1H NMR δ 1.18 (6H, t), 1.58-1.71 (4H, m), 1.75 (4H, t), 2.56 (1H, t), 3.89 (4H, q), 4.08-4.20 (4H, m), 4.88 (1H, s), 7.15-7.29 (4H, m); MS m/e MH^+ 377.


10

iv) Diethyl [4-(4-oxocyclohexyl)phenyl]malonate



Aqueous HCl (1.15 mL of a 2M solution) was added to a stirred solution of diethyl [4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl]malonate (290 mg, 0.77 mmol) in 4 mL THF. The mixture
15 was stirred for 64 h then was partitioned between water and EtOAc. The organic phase was separated and the aqueous further extracted with EtOAc. The organics were combined and dried then concentrated by evaporation to give the title compound (249 mg, 97%) as a gum; ^1H NMR δ 1.18 (6H, t), 1.82-1.93 (2H, m), 2.06-2.11 (2H, m), 2.26-2.29 (2H, m), 2.54-2.63 (2H, m), 3.04-3.10 (1H, m), 4.08-4.21 (4H, m), 4.88 (1H, s), 7.31 (4H, s); MS
20 m/e $\text{M}+\text{Na}^+$ 355.

v) Diisopropyl [4-(4-aminocyclohexyl)phenyl]malonate



Titanium tetra(isopropoxide) (426 mg, 1.5 mmol) was added to a solution of diethyl [4-(4-oxocyclohexyl)phenyl]malonate (249 mg, 0.75 mmol) in a 2M solution of ammonia in
25 isopropanol (1.88 mL, 3.75 mmol) and the reaction mixture was stirred for 64 h. Sodium borohydride (43 mg, 1.125 mmol) was added and the mixture was stirred for 4 h then aqueous ammonia (1 mL concentrated ammonia in 4 mL water) was added. The mixture was filtered and the filtrate was separated and the aqueous phase further extracted with

- 365 -


EtOAc (2 x 20 mL). The organic phase was dried and solvent was removed by evaporation to give the title compound (116 mg, 46%) as a gum; $^1\text{H NMR}$ δ 1.14-1.23 (m, 12H), 1.35-2.65 (m, 10H), 4.73-4.79 (m, 1H), 4.87-5.01 (m, 2H), 7.08-7.38 (m, 4H); MS m/e MH^+ 362.

5


vi) Following the general procedure of Example 72 except using diisopropyl [4-(4-aminocyclohexyl)phenyl]malonate as the amine in place of [4-(4-isopropylpiperazin-1-yl)phenyl]amine, the title compound (Intermediate 96) was obtained as a solid; MS m/e MH^+ 567.

10

Intermediate 97: 2-Hydrazino-2-oxo-N-[4-(propylsulphonyl)phenyl]acetamide



i) Methyl oxo{[4-(propylsulphonyl)phenyl]amino}acetate




15 The procedure described for Intermediate 43 (part iii) was followed except that the aniline used was 4-(propylsulphonyl)aniline (prepared as described in *Helv. Chim. Acta* 1983, 66(4), 1046), to give the title compound; $^1\text{H NMR}$ δ 11.27 (s, 1H), 8.02 (d, 2H), 7.87 (d, 2H), 3.93 (s, 3H), 3.19-3.27 (m, 2H), 1.48-1.63 (m, 2H), 0.85-0.97 (m, 3H); MS m/e MH^+ 286.

20

ii) The procedure described for Intermediate 43 (part iv) was followed starting with methyl oxo{[4-(propylsulphonyl)phenyl]amino}acetate to give the title compound (Intermediate 97); $^1\text{H NMR}$ δ 0.85-0.97 (m, 3H), 1.46-1.65 (m, 2H), 3.18-3.27 (m, 2H), 4.77 (s, 2H), 7.85 (d, 2H), 8.10 (d, 2H), 10.44 (s, 1H), 11.09 (s, 1H);

25 MS m/e MH^+ 286.

Intermediate 98: Methyl 2-((4-((5-((2-fluorophenyl)amino)-1,3,4-oxadiazol-2-yl)carbonyl)amino)phenyl)sulphonyl)-2-methylpropanoate




i) Methyl 2-((4-aminophenyl)sulphonyl)-2-methylpropanoate




The general procedure described for Intermediate 77 (part ii) was followed, except the starting material was methyl 2-methyl-2-((4-nitrophenyl)sulphonyl)propanoate (CAS Reg. No. 364044-08-6), to give the title compound; ^1H NMR δ 1.44 (s, 6H), 3.60 (s, 3H), 6.27 (s, 2H), 6.63 (d, 2H), 7.33 (d, 2H); MS m/e ($M+\text{Na}$) $^+$ 281.

ii) Methyl 2-((4-((methoxy(oxo)acetyl)amino)phenyl)sulphonyl)-2-methylpropanoate



The procedure described for Intermediate 43 (part iii) was followed except that the aniline used was methyl 2-((4-aminophenyl)sulphonyl)-2-methylpropanoate, to give the title compound; ^1H NMR δ 1.57 (s, 6H), 3.67 (s, 3H), 3.94 (s, 3H), 7.80 (d, 2H), 8.06 (d, 2H), 11.32 (s, 1H); MS m/e $M\text{H}^+$ 344.


iii) Methyl 2-((4-((hydrazino(oxo)acetyl)amino)phenyl)sulphonyl)-2-methylpropanoate




The procedure described for Intermediate 43 (part iv) was followed starting with methyl 2-((4-((methoxy(oxo)acetyl)amino)phenyl)sulphonyl)-2-methylpropanoate to give the title compound; ^1H NMR δ 1.56 (s, 6H), 3.67 (s, 3H), 4.77 (s, 2H), 7.77 (d, 2H), 8.11 (d, 2H); MS m/e ($M+\text{Na}$) $^+$ 366.

iv) The procedure described for Example 105 was followed starting with methyl 2-[(4-
{[hydrazino(oxo)acetyl]amino}phenyl)sulphonyl]-2-methylpropanoate to give the title
compound (Intermediate 98); $^1\text{H NMR}$ δ 1.54 (s, 6H), 3.65 (s, 3H), 7.12-7.20 (m, 1H),
7.23-7.35 (m, 2H), 7.78 (d, 2H), 8.03 (t, 1H), 8.10 (d, 2H), 10.89 (s, 1H), 11.57 (s, 1H);
5 MS m/e MH^+ 463.


Intermediate 99: *trans*-Ethyl 2-[4-(4-{[hydrazino(oxo)acetyl]amino}phenyl)
cyclohexyl]propanoate




10 i) Ethyl 2-[4-(4-hydroxyphenyl)cyclohexylidene]propanoate



A solution of triethylphosphonopropionate (12.5 g, 52.6 mmol) in THF (10 mL) was added
in one portion to a stirred mixture of sodium hydride (60 % in mineral oil, 2.3 g, 57.6
mmol) in THF (75 mL) and the reaction mixture was stirred at 10°C under an argon
15 atmosphere for 1 h. In a separate flask tetramethylguanidine (6.61 g, 57.4 mmol) was
added in one portion to a solution of 4-(4-hydroxyphenyl)cyclohexanone (9.1 g, 47.8
mmol) in THF (40 mL) at 10°C under an argon atmosphere and the stirred mixture was
warmed to room temperature over 1 h. This mixture was added in one portion to the
triethylphosphonopropionate mixture at 10 °C and the combined reaction mixture was
20 warmed to ambient temperature and stirred for 18 h under an argon atmosphere. A
saturated aqueous solution of ammonium chloride (75 mL) and EtOAc (100 mL) were
added. The layers were separated and the aqueous layer was extracted with EtOAc (100
mL). The combined organic layers were dried and concentrated *in vacuo* to leave a
residue. The residue was triturated with a mixture of ether and isohexane (2:1), filtered
25 and the filtrate was concentrated *in vacuo* to give the title compound (6.9 g, 53 %) as a
solid; $^1\text{H NMR}$ δ 1.21 (3H, t), 1.33-1.5 (2H, m), 1.78-2.1 (4H, m), 1.84 (3H, s), 2.6-2.78
(2H, m), 2.99-3.09 (1H, m), 4.12 (2H, q), 6.67 (2H, c), 7.02 (2H, d), 9.08 (1H, s); MS m/e
 MH^+ 275.

~~39/~~ii) *trans*-Ethyl 2-[4-(4-hydroxyphenyl)cyclohexyl]propanoate


10% Palladium on carbon (250 mg) was added in one portion to a stirred solution of ethyl 2-[4-(4-hydroxyphenyl)cyclohexylidene]propanoate (6.9 g, 25.3 mmol) in EtOAc (50 mL) and the reaction mixture was stirred under a hydrogen atmosphere for 24 h. The reaction mixture was filtered and concentrated *in vacuo* to give a solid that was recrystallised from a mixture of isohexane and EtOAc (10:1) to give the title compound (2.2 g, 32 %) as a solid; ¹H NMR δ (CDCl₃) 1.0-1.3 (2H, m), 1.08 (3H, d), 1.2 (3H, t), 1.3-1.45 (2H, m), 1.45-1.64 (2H, m), 1.65-1.75 (1H, m), 1.77-1.9 (3H, m), 2.15-2.26 (1H, m), 2.28-2.39 (1H, m), 4.1 (2H, q), 6.68 (2H, d), 7.0 (2H, d).

iii) *trans*-Ethyl 2-[4-(4-[[[(trifluoromethyl)sulfonyl]oxy]phenyl]cyclohexyl]propanoate

Ethyl-diisopropylamine (1.6 mL) and *N*-phenyltrifluoromethanesulfonimide (4.12 g, 11.5 mmol) was added sequentially, each in portions over 1 minute, to a stirred solution of *trans*-ethyl 2-[4-(4-hydroxyphenyl)cyclohexyl]propanoate (2.6 g, 9.23 mmol) in DCM (25 mL) at 5 °C. The reaction mixture was warmed to room temperature and stirred for 18 h and then an aqueous solution of sodium hydroxide (2M, 30 mL) was added. The layers were separated and the organic layer was washed with H₂O (30 mL), dried and concentrated *in vacuo* to leave a residue. The residue was purified by column chromatography, eluting with 0-20% EtOAc in isohexane as eluent, to give the title compound (3.6 g, 96 %) as an oil; ¹H NMR δ (CDCl₃) 1.0-1.25 (2H, m), 1.1 (3H, d), 1.2 (3H, t), 1.3-1.45 (2H, m), 1.5-1.64 (2H, m), 1.7-1.79 (1H, m), 1.8-1.95 (3H, m), 2.17-2.28 (1H, m), 2.38-2.48 (1H, m), 4.1 (2H, q), 7.1 (2H, d), 7.2 (2H, d).

22/10


iv) *trans*-Ethyl 2-(4-{4-[(diphenylmethylene)amino]phenyl}cyclohexyl)propanoate



Caesium carbonate (4.0 g, 12.4 mmol), *R*-BINAP (330 mg, 0.53 mmol), palladium acetate (119 mg, 0.53 mmol) and benzophenone imine (2.2 mL, 13.2 mmol) were added

- 5 sequentially, each in one portion, to a stirred solution of *trans*-ethyl 2-[4-(4-
 {(trifluoromethyl)-sulfonyl}oxy)phenyl]cyclohexyl]propanoate in THF (50 mL) and the
 reaction mixture was heated at 65 °C for 70 h under an argon atmosphere. The reaction
 mixture was cooled to room temperature and concentrated *in vacuo* to leave a residue. The
 residue was taken up in EtOAc (75 mL) and H₂O (25 mL) was added. The layers were
 10 separated and the organic layer was extracted with brine (25 mL), dried and concentrated
in vacuo to leave a residue. The residue was purified by chromatography, eluting with 10-
 50% EtOAc in isohexane, to give the title compound (2.3g, 59 %) as an oil; ¹H NMR δ
 (CDCl₃) 1.07-1.25 (2H, m), 1.15 (3H, d), 1.3 (3H, t), 1.35-1.5 (2H, m), 1.55-1.7 (2H, m),
 1.72-1.8 (1H, m), 1.83-1.95 (2H, m), 2.24-2.43 (2H, m), 4.15 (2H, q), 6.7 (2H, d), 7.0 (2H,
 15 d), 7.15 (2H, d), 7.23-7.35 (1H, m), 7.4 (2H, dd), 7.47-7.55 (2H, m), 7.78 (2H, d), 7.84
 (1H, d); MS *m/e* MH⁺ 440.

v) *trans*-Ethyl 2-[4-(4-aminophenyl)cyclohexyl]propanoate




- 20 10% Palladium on carbon (500 mg) was added in one portion to a stirred solution of *trans*-
 ethyl 2-(4-{4-[(diphenylmethylene)amino]phenyl}cyclohexyl)propanoate (2.3 g, 5.23
 mmol) in ethanol (50 mL) and the reaction mixture was stirred under a hydrogen
 atmosphere for 6 h. The reaction mixture was filtered and concentrated *in vacuo* to leave
 an oil. The oil was purified by chromatography, using 20-50% EtOAc in isohexane as
 25 eluent, to give the title compound (1.0 g, 71 %) as an oil; ¹H NMR δ (CDCl₃) 0.9-1.1 (2H,
 m), 0.95 (3H, d), 1.1 (3H, t), 1.18-1.32 (2H, m), 1.38-1.5 (1H, m), 1.54-1.64 (1H, m), 1.65-

- 370 -

1.8 (3H, m), 2.04-2.15 (1H, m), 2.18-2.3 (1H, m), 3.98 (2H, q), 6.8 (2H, d), 6.92 (2H, d);
MS m/e MH⁺ 276.

vi) *trans*-Ethyl 2-[4-(4-{[methoxy(oxo)acetyl]amino}phenyl)cyclohexyl]propanoate




A solution of methyl chlorooxoacetate (670 mg, 5.45 mmol) in DCM (3 mL) was added in one portion to a solution of *trans*-ethyl 2-[4-(4-aminophenyl)cyclohexyl]propanoate (1.0 g, 3.63 mmol) and pyridine (0.59 mL, 7.26 mmol) in DCM (22 mL) at 5 °C and the mixture was warmed to room temperature and stirred for 2 h. The reaction mixture was


concentrated *in vacuo* and EtOAc (30 mL) was added. The layers were separated and the organic layer was washed with water (25 mL) and brine (25 mL). The organic layer was dried and concentrated *in vacuo* to give the title compound (1.3 g, 99 %) as a solid; ¹H NMR δ (CDCl₃) 1.0-1.1.25 (2H, m), 1.1 (3H, d), 1.2 (3H, t), 1.3-1.47 (2H, m), 1.48-1.55 (2H, m), 1.68-1.79 (1H, m), 1.8-1.95 (2H, m), 2.18-2.28 (1H, m), 2.35-2.5 (1H, m), 3.9 (3H, t), 4.08 (2H, q), 7.12 (2H, d), 7.47 (2H, d); MS m/e (M-H)⁺ 360.

vii) Hydrazine hydrate (229 mg, 4.57 mmol) was added in one portion to a stirred solution of *trans*-ethyl 2-[4-(4-{[methoxy(oxo)acetyl]amino}phenyl)cyclohexyl]propanoate (1.3 g, 3.6 mmol) in ethanol (25 mL) and the reaction mixture was stirred at room temperature for 18 h. The mixture was filtered to leave a solid, which was washed with ethanol (10 mL) to give the title compound (Intermediate 99) (1.2 g, 92 %) as a solid; ¹H NMR δ 1.07 (3H, d), 1.1-1.3 (2H, m), 1.2 (3H, t), 1.36-1.5 (2H, m), 1.5-1.63 (1H, m), 1.65-1.72 (1H, m), 1.75-1.88 (3H, m), 2.2-2.32 (1H, m), 2.35-2.49 (1H, m), 4.0-4.17 (2H, m), 4.6 (2H, s), 7.19 (2H, d), 7.7 (2H, d), 10.44 (1H, s); MS m/e MH⁺ 362.

Intermediate 100 : *N*-(3,5-dichloro-4-morpholin-4-ylphenyl)-2-hydrazino-2-oxoacetamide




i) (3,5-Dichloro-4-morpholin-4-ylphenyl)amine



Following the general procedure of Intermediate 36 using 4-(2,6-dichloro-4-nitrophenyl)morpholine (prepared as described in J. Med. Chem., 1980, 23(10), 1083) the title compound was obtained; ^1H NMR δ 3.01 (4H, t), 3.64 (4H, t), 5.50 (1H, broad s), 6.57 (2H, s); MS m/e MH^+ 247.


ii) Methyl [(3,5-dichloro-4-morpholin-4-ylphenyl)amino](oxo)acetate



Following the general procedure of Intermediate 1, using pyridine as base and 4-(2,6-dichloro-4-nitrophenyl)morpholine, the title compound was obtained; ^1H NMR δ 3.09 (4H, t), 3.67 (4H, t), 3.84 (3H, s), 7.85 (2H, s), 10.98 (1H, s); MS m/e MH^+ 333.


Following the general procedure of Intermediate 18, using methyl [(3,5-dichloro-4-morpholin-4-ylphenyl)amino](oxo)acetate as starting material, the title compound (Intermediate 100) was obtained; ^1H NMR δ 3.08 (4H, t), 3.67 (4H, t), 4.62 (2H, s), 7.93 (2H, s), 10.31 (1H, broad s), 10.83 (1H, broad s); MS m/e MH^+ 333.

Intermediate 101: S-[(2-Fluorophenyl)amino]-N-piperidin-4-yl-1,3,4-oxadiazole-2-carboxamide



- 372 -

i) *tert*-Butyl 4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]piperidine-1-carboxylate



5

A suspension of ethyl 5-[(2-fluorophenyl)amino]-1,3,4-oxadiazole-2-carboxylate (intermediate 40) (102 mg, 0.41 mmol) and *tert*-butyl 4-aminopiperidine-1-carboxylate (81 mg, 0.41 mmol) was heated to 85 °C for 18 hours. Upon cooling the solid was filtered and washed with EtOH (10 mL), Et₂O (10 mL) then hexane (10 mL) to leave the title compound (123 mg, 74%) as a white solid; ¹H NMR δ 1.41 (9H, s), 1.46 - 1.54 (2H, m), 1.73 - 1.76 (2H, m), 2.81 (2H, s), 3.93 - 3.96 (3H, m), 7.13 - 7.17 (1H, m), 7.23 - 7.32 (2H, m), 7.99 - 8.04 (1H, m), 9.02 (1H, d), 10.63 (1H, s); MS *m/e* (M-H)⁺ 404.


10

ii) *tert*-Butyl 4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]piperidine-1-carboxylate (5.66 g, 13.96 mmol) was suspended in 4M HCl in 1,4-dioxane (50 mL) and stirred for 18 hours. After removal of the solvent *in vacuo* the residue was partitioned between MeOH:DCM (1:9) (500 mL) and 2M aqueous potassium carbonate (500 mL). The aqueous was extracted with MeOH:DCM (1:9) (7 x 50 mL) and the combined organics dried (Na₂SO₄), filtered and concentrated *in vacuo*. The solid was triturated with hot Et₂O (20 mL), filtered then washed with Et₂O (20 mL), hexane (20 mL) to leave the title product (Intermediate 101, 2.80 g, 66%) as a beige powder; ¹H NMR δ 1.44 - 1.57 (2H, m), 1.66 - 1.74 (2H, m), 2.47 - 2.59 (2H, m), 2.94 - 3.02 (2H, m), 3.74 - 3.86 (1H, m), 7.04 - 7.11 (1H, m), 7.17 - 7.28 (2H, m), 7.96 - 8.04 (1H, m), 8.92 (1H, d); MS *m/e* (MH)⁺ 306.


15

20

25 Intermediate 102: Methyl (6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]piperidin-1-yl}pyridin-3-yl)acetate




i) Methyl (6-chloro-1-oxidopyridin-3-yl)acetate



Hydrogen peroxide (0.409 mL of a 30% aqueous solution, 3.6 mmol) was added slowly to a stirred solution of trifluoroacetic anhydride (0.915 mL, 6.55 mmol) in DCM (5 mL) with heating under reflux. After 30min, methyl (6-chloropyridin-3-yl)acetate (Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry (2004), 43B(4), 864) (608 mg, 3.28 mmol) was added as a solution in DCM (5 mL). The reaction mixture was stirred under reflux for 1 h and then allowed to cool with stirring for 16 h. Solid potassium carbonate was added until there was no more effervescence then the mixture was filtered, concentrated by evaporation and the resultant oil was triturated with Et₂O to give the title compound (291 mg, 44%) as a solid; ¹H NMR δ 3.64 (s, 3H), 3.79 (s, 2H), 7.30 (d, 1H), 7.75 (d, 1H), 8.45 (s, 1H); MS m/e MH⁺ 202.

ii) Methyl (6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]piperidin-1-yl}-1-oxidopyridin-3-yl)acetate




5-[(2-Fluorophenyl)amino]-N-piperidin-4-yl-1,3,4-oxadiazole-2-carboxamide (Intermediate 101) (100 mg, 0.33 mmol) and methyl (6-chloro-1-oxidopyridin-3-yl)acetate (67 mg, 0.33 mmol) were heated with sodium hydrogen carbonate (56 mg, 0.66 mmol) in 2-methyl-2-butanol (4 mL) at 100°C for 22 h. The reaction mixture was concentrated by evaporation and the residue was dissolved in DMSO then filtered and purified by preparative HPLC to give the title compound (50 mg, 32%) as a gum; ¹H NMR δ 1.94-1.74 (m, 4H), 2.58-2.43 (m, 2H), 2.95-2.84 (m, 2H), 3.65 (s, 3H), 4.07-3.93 (m, 3H), 7.21-7.12 (m, 2H), 7.35-7.23 (m, 2H), 7.45-7.40 (m, 1H), 8.06-7.98 (m, 1H), 8.25-8.20 (m, 1H), 9.13 (d, 1H), 10.64 (s, 1H); MS m/e MH⁺ 471.

iii) Methyl (6-{4-[(5-[(2-fluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]piperidin-1-yl}-1-oxidopyridin-3-yl)acetate (40 mg, 0.09 mmol) was


stirred in THF (1.3 mL) at room temperature and water (1.3 mL) was added. Ammonium chloride (334 mg, 6.24 mmol) was added to the mixture followed by Zn dust (26 mg, 0.38 mmol). The reaction mixture was stirred overnight before diluting with 2 mL DMSO and filtering. The filtrate was purified by preparative HPLC to yield the title compound

5 (Intermediate 102) as a gum; MS m/e MH^+ 455.

Intermediate 103: 3-[(5-[(6-morpholin-4-ylpyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino]benzoic acid



10 i) *tert*-butyl 3-[(5-[(6-morpholin-4-ylpyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino]benzoate



Prepared from *tert*-butyl 3-isothiocyanatobenzoate and Intermediate 27 by the method of Example 1. ^1H NMR δ 1.57 (9H, s), 3.42 (4H, t), 3.71 (4H, t), 6.86 - 6.88 (1H, m), 7.52
15 (1H, t), 7.58 - 7.61 (1H, m), 7.82 - 7.85 (1H, m), 7.95 - 7.98 (1H, m), 8.25 (1H, t), 8.53 (1H, d), 10.97 (1H, s), 11.17 (1H, s); MS m/e $(MH)^+$ 467


ii) 3-[(5-[(6-morpholin-4-ylpyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino]benzoic acid

20 4M HCl in 1,4-dioxane (70 mL) was added to *tert*-butyl 3-[(5-[(6-morpholin-4-ylpyridin-3-yl)amino]carbonyl)-1,3,4-oxadiazol-2-yl]amino]benzoate (3.22 g, 6.90 mmol) and the suspension stirred for 18 hours. All volatiles were removed *in vacuo* and the residue triturated with Et₂O (20 mL), the solid filtered, washed with Et₂O (2 x 10 mL), hexane (2 x 10 mL) to give the title compound (2.80 g, 99%) as a cream powder; MS m/e $(MH)^+$ 411


25

~~238~~

Intermediate 104 : tert-butyl {[trans-4-(4-
[[hydrazino(oxo)acetyl]amino]phenyl)cyclohexyl]methyl}carbamate



5 i) Methyl [trans-4-(4-[[[(benzyloxy)carbonyl]amino]phenyl)cyclohexyl]acetate




To a solution of Intermediate 43 (3.31 g, 13.40 mmol) in dry pyridine (18 mL) under an inert atmosphere at 0 °C was added benzyl chloroformate (2.30 mL, 16.08 mmol) then allowed to warm to ambient temperature and stirred for 18 hours. After concentrating *in vacuo* the residue was partitioned between Et₂O (150 mL) and 1M aqueous HCl (20 mL). The organic phase was dried (MgSO₄), filtered and evaporated to a yellow solid.

10 Purification by chromatography through silica gave the title compound (3.50 g, 68%) as an off white solid; ¹H NMR (CDCl₃) δ 1.03 - 1.12 (2H, m), 1.35 - 1.46 (2H, m), 1.74 - 1.82 (5H, m), 2.17 - 2.18 (2H, m), 2.33 - 2.39 (1H, m), 3.61 (3H, s), 5.12 (2H, s), 6.50 (1H, s), 7.06 (2H, d), 7.19 - 7.34 (7H, m)

15

ii) [trans-4-(4-[[[(Benzyloxy)carbonyl]amino]phenyl)cyclohexyl]acetic acid



To a solution of methyl [trans-4-(4-[[[(benzyloxy)carbonyl]amino]phenyl)cyclohexyl]acetate (3.34 g, 8.76 mmol) in THF (42 mL) and MeOH (42 mL) was added 0.6M aqueous lithium hydroxide (43.6 mL, 26.27 mmol) and the mixture stirred at ambient temperature for 18 hours. The mixture was partitioned between EtOAc (200 mL) and 1M HCl (100 mL). The organic phase was concentrated *in vacuo* and the residue partitioned between EtOAc (300 mL) and brine (20 mL). The organic phase was dried (MgSO₄),

20 filtered and evaporated to give the title compound (3.19 g, 99%) as a white solid; ¹H NMR


25

- 376 -

387


(CDCl₃) δ 1.05 - 1.15 (2H, m), 1.36 - 1.46 (2H, m), 1.74 - 1.87 (5H, m), 2.22 (2H, d), 2.34 - 2.41 (1H, m), 5.13 (2H, s), 6.58 (1H, s), 7.07 (2H, d), 7.21 (3H, t), 7.28 - 7.35 (4H, m);
MS m/e (MH)⁺ 368

5 iii) Benzyl [4-(*trans*-4-[(*tert*-butoxycarbonyl)amino]methyl)cyclohexyl]phenyl]carbamate



To a suspension of [*trans*-4-(4-[(benzyloxy)carbonyl]amino)phenyl]cyclohexyl]acetic acid (516 mg, 1.40 mmol) in *tert*-butanol (3 mL) under an inert atmosphere was added triethylamine (0.45 mL, 3.23 mmol) and diphenyl phosphorylazide (0.70 mL, 3.23 mmol) and the mixture heated to 80 °C for 24 hours. Upon cooling the mixture was partitioned between EtOAc (100 mL) and H₂O (20 mL). The aqueous phase was further extracted with DCM (2 x 20 mL) and the combined organics dried (MgSO₄), filtered and evaporated. Purification by chromatography through silica gave the title compound (525 mg, 86%) as a white solid. ¹H NMR (CDCl₃) δ 0.96 - 1.07 (2H, m), 1.30 - 1.44 (12H, m), 1.78 - 1.85 (4H, m), 2.33 - 2.39 (1H, m), 2.94 (2H, t), 4.52 (1H, s), 5.12 (2H, s), 6.51 (1H, s), 7.04 - 7.07 (2H, m), 7.19 - 7.34 (7H, m)

iv) *tert*-Butyl {[*trans*-4-(4-aminophenyl)cyclohexyl]methyl}carbamate




20 Benzyl [4-(*trans*-4-[(*tert*-butoxycarbonyl)amino]methyl)cyclohexyl]phenyl]carbamate (0.50 g, 1.14 mmol) in EtOH (10 mL) and EtOAc (10 mL) was hydrogenated over 10% Pd/C (100 mg) at ambient pressure and 40 °C. After 2 hours the catalyst was removed by filtration and the solution concentrated *in vacuo* to leave the title compound (320 mg, 86%) as a white solid; ¹H NMR δ 0.95 - 1.02 (2H, m), 1.25 - 1.39 (12H, m), 1.73 - 1.75 (4H, m), 2.22 - 2.25 (1H, m), 2.80 (2H, t), 4.75 (2H, s), 6.46 - 6.49 (2H, m), 6.78 (1H, t), 6.85 (2H, d)

- 377 -

321

v) Methyl {[4-(*trans*-4-[(*tert*-butoxycarbonyl)amino]methyl)cyclohexyl]phenyl}amino}(oxo)acetate



Prepared from *tert*-butyl {[*trans*-4-(4-aminophenyl)cyclohexyl]methyl} carbamate and methyl chlorooxoacetate by method described for Intermediate 1, method D3.


¹H NMR (CDCl₃) δ 1.08 - 1.18 (2H, m), 1.41 - 1.52 (12H, m), 1.93 (4H, t), 2.46 - 2.53 (1H, m), 3.05 (2H, t), 4.00 (3H, s), 4.63 (1H, s), 7.23 (2H, d), 7.57 (2H, d), 8.81 (1H, s)

vi) *tert*-Butyl {[*trans*-4-(4-[[hydrazino(oxo)acetyl]amino]phenyl)cyclohexyl]methyl}carbamate


Prepared using hydrazine hydrate in EtOH from methyl {[4-(*trans*-4-[(*tert*-butoxycarbonyl)amino]methyl)cyclohexyl]phenyl}amino}(oxo)acetate as described for Intermediate 18. ¹H NMR δ 0.99 - 1.06 (2H, m), 1.37 - 1.42 (12H, m), 1.77 - 1.80 (4H, m), 2.39 - 2.49 (1H, m), 2.81 (2H, t), 4.60 (2H, s), 6.80 (1H, t), 7.19 (2H, d), 7.68 (2H, d),

10.21 (1H, s), 10.45 (1H, s)

Intermediate 105: Methyl 2-[*trans*-4-(4-[[hydrazino(oxo)acetyl]amino]phenyl)-cyclohexyl]-2-methylpropanoate



i) Methyl 2-[*trans*-4-(4-[(*tert*-butyl(dimethyl)silyl]oxy)phenyl)cyclohexyl]propanoate




A solution of *tert*-butyl dimethylsilylchloride (7.29 g, 48.3 mmol) in DMA (80 mL) was added dropwise to a stirred solution of methyl [*trans*-4-(4-hydroxyphenyl)cyclohexyl]acetate (prepared as described in Patent Application

- 378 -

WO2004/047755) (10.0 g, 40.3 mmol) and imidazole (6.86 g, 100.7 mmol) and the reaction mixture was stirred at room temperature for 20 h. Ethyl acetate (250 mL) was added and the mixture was washed with H₂O (6x30 mL) and brine. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave the silyl-protected phenol as a colourless oil, which was used with no further purification.

- 5 A solution of lithium diisopropylamide (14.9 mmol) in THF (10 mL) was added dropwise to a stirred solution of the crude silyl-protected phenol (5.43 g, 14.9 mmol) in THF (50 mL) at -78°C under an argon atmosphere and the mixture was warmed to 0°C. The mixture was cooled to -78°C and methyl iodide (0.98 mL, 15.7 mmol) was added dropwise. The reaction mixture was warmed to room temperature and stirred for 20 h. Saturated aqueous ammonium chloride (50 mL) and ethyl acetate were added sequentially, both in one portion. The organic layer was separated and washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave an oil. The oil was purified by column chromatography, using a gradient of 10-25% ethyl acetate:isohexane as eluent, to give the title compound (5.0 g, 89 %) as a colourless oil; ¹H NMR δ 0.0 (6H, s), 0.8 (9H, s), 0.9-1.08 (2H, m), 0.97 (3H, d), 1.18-1.32 (2H, m), 1.37-1.5 (2H, m), 1.52-1.62 (1H, m), 1.66-1.8 (2H, m), 2.08-2.28 (2H, m), 3.5 (3H, s), 6.57 (2H, d), 6.86 (2H, d).
- 10
- 15

- 20 **ii) Methyl 2-[(*trans*-4-(4-[[*tert*-butyl(dimethyl)silyl]oxy}phenyl)cyclohexyl]-2-methylpropanoate**




- A solution of lithium diisopropylamide (12.9 mmol) in THF (20 mL) was added dropwise to a stirred solution of methyl 2-[(*trans*-4-(4-[[*tert*-butyl(dimethyl)silyl]oxy}phenyl)cyclohexyl]-propanoate (4.42 g, 11.7 mmol) in THF (40 mL) at -78°C under an argon atmosphere and the mixture was warmed to room temperature. The mixture was cooled to -78°C and methyl iodide (0.92 mL, 14.7 mmol) was added dropwise. The reaction mixture was warmed to room temperature and stirred for 20 h. Saturated aqueous ammonium chloride (100 mL) and ethyl acetate were added
- 25

- 379 -

sequentially, both in one portion. The organic layer was separated and washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave an oil. The oil was purified by column chromatography, using a gradient of 0-15% ethyl acetate/isohexane as eluent, to give the title compound (4.0 g, 87 %) as a colourless oil; ¹H NMR δ 0.0 (6H, s), 0.8 (9H, s), 0.98 (6H, s), 1.0-1.1 (2H, m), 1.18-1.31 (2H, m), 1.45-1.56 (3H, m), 1.7-1.8 (2H, m), 2.15-2.28 (1H, m), 3.5 (3H, s), 6.55 (2H, d), 6.85 (2H, d).

iii) Methyl 2-methyl-2-[*trans*-4-(4-[[trifluoromethyl]sulfonyl]oxy]phenyl)cyclohexyl]-propanoate



10

A solution of tetrabutylammonium fluoride (1M in THF, 11.3 mL, 11.3 mmol) was added in one portion to a stirred solution of methyl 2-[*trans*-4-(4-[[*tert*-butyl(dimethyl)silyl]oxy]-phenyl)cyclohexyl]-2-methylpropanoate (4.0 g, 10.2 mmol) in THF (40 mL) and the reaction mixture was stirred at room temperature for 18 h. Ethyl acetate (75 mL) was added and the mixture was washed with H₂O (2x50 mL) and brine. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave the crude phenol as an oil, which was used with no further purification.


15

N-Phenyltrifluoromethanesulfinamide (6.47 g, 18.1 mmol) was added portionwise over 1 min to a stirred solution of the crude phenol (4.0 g, 14.5 mmol) and DIPEA (2.52 mL, 14.5 mmol) in DCM (70 mL) at 10 °C. The reaction mixture was stirred at room temperature for 18 h and then a 2N aqueous solution of NaOH (100 mL) was added. The layers were separated and the organic layer was washed with H₂O and then passed through a phase separation cartridge and concentrated *in vacuo* to leave an oil. The oil was purified by column chromatography, eluting with 0-20% ethyl acetate in isohexane, to give the title compound (4.57 g, 77 %) as a colourless oil; ¹H NMR δ 1.15 (6H, s), 1.17-1.3 (2H, m), 1.38-1.52 (2H, m), 1.63-1.39 (3H, m), 1.9-2.0 (2H, m), 2.43-2.56 (1H, m), 3.68 (3H, s), 7.18 (2H, d), 7.27 (2H, d).

20

25

(iv) Methyl 2-[*trans*-4-(4-aminophenyl)cyclohexyl]-2-methylpropanoate



- Caesium carbonate (5.11 g, 15.7 mmol), *R*-BINAP (420 mg, 0.67 mmol), palladium acetate (151 mg, 0.67 mmol) and benzophenone imine (2.82 mL, 16.8 mmol) were added sequentially, each in one portion, to a stirred solution of methyl 2-methyl-2-[*trans*-4-4-
5 {[(trifluoromethyl)sulfonyl]oxy}phenyl)cyclohexyl]-propanoate (4.57 g, 11.2 mmol) in THF (60 mL) and the reaction mixture was heated at 65 °C for 18 h under an argon atmosphere. The reaction mixture was cooled to room temperature and concentrated *in vacuo* to leave a residue. The residue was taken up in EtOAc and H₂O was added. The
10 layers were separated and the organic layer was washed with brine, dried (MgSO₄) and concentrated *in vacuo* to leave a residue. The residue was purified by column chromatography, eluting with 10-50% EtOAc in isohexane, to give the crude benzophenone imine product (1.7 g) as an oil, which was used with no further purification; MS *m/e* MH⁺ 440.
- 15 10% Palladium on carbon (500 mg) was added in one portion to a stirred solution of the crude benzophenone imine (1.7 g) in a mixture of MeOH (25 mL) and THF (25 mL) and the reaction mixture was stirred under a hydrogen atmosphere for 20 h. The reaction mixture was filtered and concentrated *in vacuo* to leave an oil. The oil was purified by column chromatography, using a gradient of 11-50% EtOAc:isohexane as eluent, to give
20 the title compound (250 mg, 8 %) as a solid; ¹H NMR δ 1.14 (6H, s), 1.15-1.28 (2H, m), 1.36-1.48 (2H, m), 1.61-1.74 (3H, m), 1.86-1.95 (2H, m), 2.3-2.4 (1H, m), 3.52 (2H, s), 3.67 (3H, s), 6.61 (2H, d), 6.99 (2H, d); MS *m/e* MH⁺ 276.


- (v) A solution of methyl chlorooxoacetate (167 mg, 1.36 mmol) in DCM (2 mL) was
25 added in one portion to a solution of methyl 2-[*trans*-4-(4-aminophenyl)cyclohexyl]-2-methyl-propanoate (250 mg, 0.91 mmol) and pyridine (0.22 mL, 2.72 mmol) in DCM (10 mL) at 10 °C and the mixture was stirred at 10 °C for 30 mins and then warmed to room temperature and stirred for 2 h. The reaction mixture was concentrated *in vacuo* and

381


EtOAc was added. The mixture was washed with water and brine, dried (MgSO_4) and concentrated *in vacuo* to give the crude methyl ester (0.35 g) as a solid, which was used with no further purification; $^1\text{H NMR}$ δ 1.16 (6H, s), 1.18-1.3 (2H, m), 1.38-1.52 (2H, m), 1.64-1.78 (3H, m), 1.9-2.0 (2H, m), 2.4-2.51 (1H, m), 3.68 (3H, s), 3.97 (3H, m), 7.2 (2H, d), 7.53 (2H, d), 8.76 (1H, s); MS m/e ($\text{M}-\text{H}^-$) 360.

A solution of hydrazine hydrate (54 mg, 1.08 mmol) in ethanol (1 mL) was added dropwise to a stirred solution of the crude methyl ester (0.35 g, 0.91 mmol) in ethanol (9 mL) and the reaction mixture was stirred at room temperature for 18 h. The mixture was filtered to leave a solid, which was washed with ethanol to give the title compound (Intermediate 105) (0.30 g, 91 %) as a white solid; $^1\text{H NMR}$ δ 1.13 (6H, s), 1.16-1.3 (2H, m), 1.4-1.54 (2H, m), 1.63-1.74 (3H, m), 1.82-1.93 (2H, m), 1.9-2.54 (1H, m), 3.68 (3H, s), 4.69 (2H, s), 7.25 (2H, d), 7.75 (2H, d), 10.33 (1H, s), 10.57 (1H, s); MS m/e MH^+ 362.

**Intermediate 106: Ethyl 2-*trans*-4-(2-chloro-4-
hydrazino(oxo)acetyl)amino}phenyl)-cyclohexylpropanoate**



i) 3-Chloro-4-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)aniline




A 2N aqueous solution of potassium carbonate (26.5 mL) and tetrakis(triphenylphosphine) palladium(0) (1.23 g, 1.06 mmol) were added sequentially, each in one portion, to a stirred solution of 8-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,4-dioxaspiro[4.5]dec-7-ene (Intermediate 78ii, 5.65 g, 21.2 mmol) and 4-bromo-3-chloro aniline (4.39 g, 21.2 mmol) in dimethoxyethane (80 mL) at room temperature under an argon atmosphere. The reaction mixture was then heated at 80 °C for 24 h under an argon atmosphere. The reaction mixture was cooled to room temperature and ethyl acetate was added and then the mixture was washed twice with H_2O and then brine. The organic layer was dried (MgSO_4) and concentrated *in vacuo* to leave an oil, which was purified by column chromatography, using a gradient of 20-50% ethyl acetate:isohexane as eluent, to give the title compound

- 382 -

~~381~~

(3.5 g, 62%) as a solid; ^1H NMR δ 1.7-1.8 (2H, m), 2.26-2.43 (4H, m), 3.91 (4H, s), 5.28 (2H, s), 5.4-5.47 (1H, m), 6.45 (1H, dd), 6.59 (1H, d), 6.86 (1H, d).

ii) Benzyl [3-chloro-4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl]carbamate



5

Platinum oxide (250 mg) was added in one portion to a stirred solution of 3-chloro-4-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)aniline (3.4 g, 12.8 mmol) in ethyl acetate (50 mL) and the mixture was stirred at room temperature under a hydrogen atmosphere for 9 h. A further portion of platinum oxide (250 mg) was added in one portion and the mixture was stirred for 8 h at room temperature under a hydrogen atmosphere. The reaction mixture was filtered and concentrated *in vacuo* to leave the cyclohexane-aniline product (2.1 g) as a solid, which was used with no further purification.


10

Pyridine (1.91 mL, 23.5 mmol) and benzyl chloroformate (1.34g, 7.84 mmol) were added sequentially, each in one portion, to a stirred solution of the crude cyclohexane-aniline (2.1 g, 7.84 mmol) in THF (30 mL) at 5°C and the reaction mixture was stirred at 10°C for 3 h. Ethyl acetate (75 mL) was added and the mixture was washed with H₂O (2x30 mL) and brine. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave a solid, which was purified by column chromatography, using a gradient of 20-50% ethyl acetate:isohexane as eluent, to give the title compound (2.3 g, 74%) as a white solid; ^1H NMR δ 1.58-1.96 (8H, m), 4.0 (4H, s), 5.22 (2H, s), 6.63 (1H, s), 7.2 (1H, dd), 7.25 (1H, d), 7.33-7.45 (5H, m), 7.5 (1H, d); MS *m/e* MH⁺ 402.

15

20

iii) Ethyl 2-[4-(4-[(benzyloxy)carbonyl]amino)-2-chlorophenyl]cyclohexylidene]propanoate




25

Water (1 mL) was added in one portion to a stirred solution of benzyl [3-chloro-4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl]carbamate (2.30 g, 5.72 mmol) in TFA (20 mL) and the reaction mixture was stirred at room temperature for 2 h. The mixture was cooled to 10°C and a 2N aqueous solution of sodium hydroxide was added until the pH of the mixture was 10. Ethyl acetate (50 mL) was added and the layers were separated and the aqueous layer was extracted with ethyl acetate (50 mL). The combined organic layers were washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave the crude cyclohexanone product (2.0 g), which was used with no further purification.

Triethylphosphonopropionate (1.60 g, 6.71 mmol) was added dropwise to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 0.29 g, 7.27 mmol) in THF (30 mL) at 10°C under an argon atmosphere and the mixture was warmed to room temperature. The reaction mixture was cooled to 10°C and then a solution of the crude cyclohexanone (2.0 g) in THF (10 mL) was added dropwise. The reaction mixture was warmed to room temperature and stirred for 4 h. A saturated aqueous solution of ammonium chloride (50 mL) and ethyl acetate (70 mL) were added. The layers were separated and the organic layer was washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave an oil. Purification by column chromatography, using a gradient of 11-66% ethyl acetate:isohexane as eluent, gave the title compound (1.50 g, 60%) as a solid; ¹H NMR δ 1.3 (3H, t), 1.4-1.6 (2H, m), 1.9 (3H, s), 1.94-2.12 (4H, m), 2.72-2.85 (1H, m), 3.1-3.25 (2H, m), 4.2 (2H, q), 5.2 (2H, s), 6.59 (1H, s), 7.13 (1H, dd), 7.19 (1H, dd), 7.3-7.43 (5H, m), 7.5 (1H, d); MS m/e (M-H)⁺ 440.

iv) Ethyl 2-[4-(2-chloro-4-{[methoxy(oxo)acetyl]amino}phenyl)cyclohexyl]propanoate




Platinum oxide (300 mg) was added in one portion to a stirred solution of ethyl 2-[4-(4-[(benzyloxy)carbonyl]amino)-2-chlorophenyl)cyclohexylidene]propanoate (1.5 g, 3.39

mmol) in ethyl acetate (30 mL) and the mixture was stirred at room temperature under a hydrogen atmosphere for 7 h. A further portion of platinum oxide (1.0 g) was added in one portion and the mixture was stirred for 1 h at room temperature under a hydrogen atmosphere. The reaction mixture was filtered and concentrated *in vacuo* to leave a residue, which was purified by column chromatography, using a gradient of 11-50% ethyl acetate:isohexane as eluent, to give the aniline product as a mixture of *cis*- and *trans*-cyclohexane isomers, that was used with no further purification.


A solution of methyl chlorooxoacetate (266 mg, 2.17 mmol) in DCM (2 mL) was added in one portion to a solution of the crude aniline (500 mg, 1.44 mmol) and pyridine (0.35 mL, 4.33 mmol) in DCM (8 mL) at 10 °C and the mixture was stirred at 10 °C for 30 mins and then warmed to room temperature and stirred for 2 h. The reaction mixture was concentrated *in vacuo* and EtOAc was added. The mixture was washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave the title compound as an oil (0.58 g, 43% over 2 steps); ¹H NMR (for major *trans*-isomer) δ 1.15 (3H, d), 1.19-1.33 (2H, m), 1.28 (3H, t), 1.35-1.5 (2H, m), 1.6-1.75 (1H, m), 1.76-1.85 (1H, m), 1.86-2.0 (3H, m), 2.25-2.35 (1H, m), 2.89-3.01 (1H, m), 3.99 (3H, s), 4.16 (2H, dq), 7.24 (1H, d), 7.48 (1H, dd), 7.7 (1H, d), 8.75 (1H, s); MS m/e (M-H)⁺ 394.

v) A solution of hydrazine hydrate (86 mg, 1.73 mmol) in ethanol (1 mL) was added dropwise to a stirred solution of ethyl 2-[4-(2-chloro-4-([methoxy(oxo)acetyl]amino)phenyl)cyclo-hexyl]propanoate (0.58 g, 1.44 mmol) in ethanol (14 mL) and the reaction mixture was stirred at room temperature for 18 h. The mixture was filtered to leave a solid, which was washed with ethanol to give the title compound (Intermediate 106) (0.45 g, 81 %) as a white solid; ¹H NMR δ 1.14 (3H, d), 1.16-1.28 (2H, m), 1.27 (3H, t), 1.4-1.58 (2H, m), 1.6-1.7 (1H, m), 1.71-1.93 (4H, m), 2.29-2.39 (1H, m), 2.82-2.94 (1H, m), 4.15 (2H, dq), 4.7 (2H, s), 7.4 (1H, d), 7.8 (1H, dd), 8.01 (1H, d), 10.4 (1H, s), 10.8 (1H, s); MS m/e MH⁺ 396.

**Intermediate 107: Ethyl 2-[(trans-4-(3-chloro-4-
hydrazino(oxo)acetyl)amino)phenyl]-cyclohexyl]propanoate**




i) 2-Chloro-4-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)aniline




Following the general procedure of Intermediate 106, except that 4-bromo-2-chloroaniline was used in place of 4-bromo-3-chloroaniline, the title compound was obtained in 68% yield as a solid; ¹H NMR δ 1.75-1.82 (2H, m), 2.3-2.36 (2H, m), 2.41-2.5 (2H, m), 3.91 (4H, s), 5.3 (2H, s), 5.81-5.88 (1H, m), 6.74 (1H, d), 7.11 (1H, dd), 7.21 (1H, d).

ii) 2-Chloro-4-(1,4-dioxaspiro[4.5]dec-8-yl)aniline



Platinum oxide (200 mg) was added in one portion to a stirred solution of 2-chloro-4-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)aniline (3.8 g, 14.3 mmol) in ethyl acetate (50 mL) and the mixture was stirred at room temperature under a hydrogen atmosphere for 8 h. The reaction mixture was filtered and concentrated *in vacuo* to leave the title compound (2.4 g, 63%) as a solid; ¹H NMR δ 1.5-1.65 (4H, m), 1.67-1.8 (4H, m), 2.38-2.48 (1H, m), 3.88 (4H, s), 5.1 (2H, s), 6.72 (1H, d), 6.9 (1H, dd), 7.0 (1H, d); MS m/e MH⁺ 268.


iii) Benzyl [2-chloro-4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl]carbamate



Pyridine (2.18 mL, 26.9 mmol) and benzyl chloroformate (1.53 g, 8.96 mmol) were added sequentially, each in one portion, to a stirred solution of 2-chloro-4-(1,4-


dioxaspiro[4.5]dec-8-yl)aniline (2.40 g, 8.96 mmol) in THF (30 mL) at 5°C and the reaction mixture was stirred at 10°C for 3 h. Ethyl acetate (75 mL) was added and the mixture was washed with H₂O (2x30 mL) and brine. The organic layer was dried (MgSO₄) and concentrated *in vacuo* to leave a solid, which was purified by column chromatography, using a gradient of 20-50% ethyl acetate:isohexane as eluent, to give the title compound (2.6 g, 72%) as a white solid; ¹H NMR δ 1.65-1.93 (8H, m), 2.49-2.6 (1H, m), 4.0 (4H, s), 5.25 (2H, s), 7.12 (1H, s), 7.15 (1H, dd), 7.25 (1H, d), 7.37-7.48 (5H, m), 8.08 (1H, d); MS m/e (M-H)⁺ 400.

10 iv) Benzyl [2-chloro-4-(4-oxocyclohexyl)phenyl]carbamate



Water (1 mL) was added in one portion to a stirred solution of benzyl [2-chloro-4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl]carbamate (2.60 g, 6.47 mmol) in TFA (20 mL) and the reaction mixture was stirred at room temperature for 2 h. The mixture was cooled to 10°C and a 2N aqueous solution of sodium hydroxide was added until the pH of the mixture was 10. Ethyl acetate (50 mL) was added and the layers were separated and the aqueous layer was extracted with ethyl acetate (50 mL). The combined organic layers were washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave the title compound (2.3 g, 99%) as a solid; ¹H NMR δ 1.82-1.99 (2H, m), 2.17-2.28 (2H, m), 2.45-2.57 (4H, m), 2.94-3.07 (1H, m), 5.24 (2H, s), 7.17 (1H, dd), 7.25 (1H, d), 7.32-7.5 (5H, m), 8.13 (1H, d); MS m/e (M-H)⁺ 356.


v) Ethyl 2-[4-(4-((benzyloxy)carbonyl)amino)-3-chlorophenyl)cyclohexylidene]-propanoate



- 387 -

A solution of triethylphosphonopropionate (1.54 g, 6.46 mmol) in THF (10 mL) was added dropwise to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 0.28 g, 7.05 mmol) in THF (30 mL) at 10°C under an argon atmosphere and the mixture was warmed to room temperature. The reaction mixture was cooled to 10°C and then a solution of benzyl (2-chloro-4-(4-oxocyclohexyl)phenyl)carbamate (2.36 g, 5.87 mmol) in THF (10 mL) was added dropwise. The reaction mixture was warmed to room temperature and stirred for 4 h. A saturated aqueous solution of ammonium chloride (50 mL) and ethyl acetate (70 mL) were added. The layers were separated and the organic layer was washed with H₂O and brine, dried (MgSO₄) and concentrated *in vacuo* to leave an oil. Purification by column chromatography, using a gradient of 11-66% ethyl acetate:isohexane as eluent, gave the title compound (1.80 g, 69%) as a solid; ¹H NMR δ 1.34 (3H, t), 1.46-1.53 (2H, m), 1.93 (3H, s), 1.95-2.1 (4H, m), 2.67-2.87 (2H, m), 3.13-3.24 (1H, m), 4.22 (2H, q), 5.24 (2H, s), 7.1-7.18 (2H, m), 7.21 (1H, d), 7.33-7.5 (5H, m), 8.08 (1H, d); MS m/e (M-H)⁺ 440.

vi) Ethyl 2-[*trans*-4-(4-amino-3-chlorophenyl)cyclohexyl]propanoate




Platinum oxide (300 mg) was added in one portion to a stirred solution of ethyl 2-[4-({[(benzyloxy)carbonyl]amino}-3-chlorophenyl)cyclohexylidene]propanoate (1.7 g, 3.85 mmol) in ethyl acetate (50 mL) and the mixture was stirred at room temperature under a hydrogen atmosphere for 9 h. The reaction mixture was filtered and concentrated *in vacuo* to leave a residue, which was purified by column chromatography, using a gradient of 11-66% ethyl acetate:isohexane as eluent, to give the aniline product as a mixture of *cis*- and *trans*-cyclohexane isomers. Recrystallisation from a mixture of ethanol and diethyl ether gave the title compound (0.37 g, 28%) as a white solid; ¹H NMR δ 1.08 (3H, d), 1.2-1.3 (1H, m), 1.27 (3H, t), 1.31-1.49 (1H, m), 1.52-1.7 (1H, m), 1.71-1.81 (1H, m), 1.81-1.97 (3H, m), 2.2-2.32 (1H, m), 2.33-2.48 (1H, m), 4.15 (2H, dq), 7.02 (1H, dd), 7.2 (1H, d), 7.22 (1H, d); MS m/e MH⁺ 310.


vii) A solution of methyl chlorooxoacetate (197 mg, 1.60 mmol) in DCM (2 mL) was added in one portion to a solution of ethyl 2-[*trans*-4-(4-amino-3-chlorophenyl)cyclohexyl]-propanoate (370 mg, 1.07 mmol) and pyridine (0.26 mL, 3.21 mmol) in DCM (8 mL) at 10 °C and the mixture was stirred at 10 °C for 30 mins and then warmed to room temperature and stirred for 2 h. The reaction mixture was concentrated *in vacuo* and EtOAc was added. The mixture was washed with water and brine, dried (MgSO₄) and concentrated *in vacuo* to give the crude methyl ester (0.44 g) as a solid, which was used with no further purification; ¹H NMR δ 1.15 (3H, d), 1.18-1.3 (1H, m), 1.27 (3H, t), 1.36-1.5 (2H, m), 1.58-1.7 (1H, m), 1.73-1.83 (1H, m), 1.85-1.98 (3H, m), 2.23-2.34 (1H, m), 2.39-2.51 (1H, m), 4.0 (3H, s), 4.15 (2H, dq), 7.15 (1H, dd), 7.26 (1H, d), 8.3 (1H, d), 9.35 (1H, s); MS m/e MH⁺ 396.

Hydrazine hydrate (64 mg, 1.28 mmol) was added to a stirred solution of the crude methyl ester (0.44 g) in ethanol (5 mL) and the reaction mixture was stirred at room temperature for 18 h. The mixture was filtered to leave a solid, which was washed with ethanol to give the title compound (Intermediate 107) (0.35 g, 83 %) as a white solid; ¹H NMR δ 1.11 (3H, d), 1.15-1.3 (2H, m), 1.24 (3H, t), 1.42-1.58 (2H, m), 1.59-1.7 (1H, m), 1.7-1.8 (1H, m), 1.81-1.95 (3H, m), 2.28-2.38 (1H, m), 2.49-2.6 (1H, m), 4.08 (2H, dq), 4.72 (2H, s), 7.32 (1H, dd), 7.47 (1H, d), 7.9 (1H, d), 10.05 (1H, s), 10.45 (1H, s); MS m/e MH⁺ 396.

20 Intermediate 108: Ethyl *cis*-4-(4-{[methoxy(oxo)acetyl]amino}phenoxy)cyclohexanecarboxylate




i) Ethyl *cis*-4-(4-nitrophenoxy)cyclohexanecarboxylate



25 A solution of ethyl 4-hydroxycyclohexanecarboxylate (2.0 g, 11.6 mmol) in DMF (3 mL) was added in one portion to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 490 mg, 12.2 mmol) in DMF (15 mL) at 0 °C under an argon atmosphere. The mixture was stirred at 0 °C for 10 mins and then warmed to room temperature and stirred


for 20 mins and then a solution of 1-fluoro-4-nitrobenzene (1.24 mL, 11.6 mmol) in DMF (2 mL) was added in one portion. The reaction mixture was stirred at room temperature for 2 h and then heated at 110°C for 24 h. The solution was concentrated *in vacuo* to leave a residue that was taken up in EtOAc (250 mL) and H₂O (150 mL) was added. The layers were separated and the organic layer was washed with H₂O (2x100 mL) and brine (50 mL) and then dried (MgSO₄) and concentrated *in vacuo* to leave an oil. The oil was taken up in a mixture of hot EtOAc and isohexane (1:1) and left for 20 h. Filtration, followed by concentration *in vacuo* of the filtrate left an oil, which was purified by column chromatography, using a gradient of 10-40% EtOAc:isohexane as eluent, to give the title compound (0.27 g, 16%) as an oil; ¹H NMR (CDCl₃) δ 1.24 (3H, q), 1.66 - 1.72 (2H, m), 1.74 - 1.79 (2H, m), 1.90 - 1.95 (2H, m), 1.98 - 2.04 (2H, m), 2.38 - 2.44 (1H, m), 4.11 - 4.16 (2H, q), 4.57 - 4.60 (1H, m), 6.90 - 6.93 (2H, d), 8.15 - 8.18 (2H, d).

ii) Ethyl *cis*-4-(4-aminophenoxy)cyclohexanecarboxylate




Palladium (10 wt. %) on carbon (50 mg) was added in one portion to a solution of ethyl *cis*-4-(4-nitrophenoxy)cyclohexanecarboxylate (0.23 g, 0.78 mmol) in a mixture of EtOH (15 mL) and THF (10 mL) and the reaction mixture was stirred under a hydrogen atmosphere at room temperature for 16 h. The reaction mixture was filtered and then concentrated *in vacuo* to leave the title compound (0.20 g, 100%) as an oil; ¹H NMR (CDCl₃) δ 1.23 - 1.28 (3H, t), 1.55 - 1.63 (2H, m), 1.68 - 1.74 (2H, m), 1.93 - 2.03 (4H, m), 2.34 - 2.40 (1H, m), 4.10 - 4.17 (2H, q), 4.26 - 4.30 (1H, m), 6.60 - 6.63 (2H, d), 6.73 - 6.78 (2H, d) MS *m/e* MH⁺ 264.

iii) Ethyl *cis*-4-(4-([methoxy(oxo)acetyl]amino)phenoxy)cyclohexanecarboxylate




Methyl chlorooxoacetate (72 μ L, 0.77 mmol) was added in one portion to a stirred solution of ethyl *cis*-4-(4-aminophenoxy)cyclohexanecarboxylate (204 mg, 0.77 mmol) and pyridine (76 μ L, 0.93 mmol) in DCM (5 mL) and the reaction mixture was stirred at room temperature for 1 h. Concentration *in vacuo* left a residue which was taken up in EtOAc (50 mL) and an aqueous solution of hydrochloric acid (1M, 30 mL) was added. The layers were separated and the organic layer was washed with brine (20 mL), dried (MgSO_4) and concentrated *in vacuo* to leave an oil. The oil was purified by column chromatography, using a gradient of 10-70% ethyl acetate:isohexane as eluent, to give the title compound (Intermediate 108) (188 mg, 70%) as an oil; ^1H NMR (CDCl_3) δ 1.23 - 1.28 (3H, t), 1.57 - 1.68 (2H, m), 1.71 - 1.78 (2H, m), 1.92 - 2.04 (4H, m), 2.36 - 2.43 (1H, m), 3.96 (3H, s), 4.14 (2H, q), 4.43 - 4.45 (1H, m), 6.89 - 6.93 (2H, d), 7.51 - 7.55 (2H, d), 8.76 (1H, s); MS m/e (M-H) $^+$ 348.

Intermediate 109: Methyl 1-(5-[[hydrazino(oxo)acetyl]amino]pyridin-2-yl)piperidine-4-carboxylate



i) 1-*tert*-Butyl 4-methyl 4-methylpiperidine-1,4-dicarboxylate




To a solution of 1-(*tert*-butoxycarbonyl)-4-methylpiperidine-4-carboxylic acid (5.0 g, 20.55 mmol) in methanol (60 mL) and toluene (100 mL) was added a 2M solution of trimethylsilyldiazomethane (20 mL) in hexanes (the addition was accompanied by effervescence and a small exotherm). The resulting pale yellow solution was allowed to stir at ambient temperature for 1 hr. The solvent was removed under reduce pressure to give the title compound as a pale yellow oil (5.28 g 20.55 mmol, 100%); ^1H NMR (CDCl_3) δ 1.20 (s, 3H), 1.34 - 1.39 (m, 2H), 1.44 (s, 9H), 2.03 - 2.07 (m, 2H), 2.96 - 3.03 (m, 2H), 3.70 (s, 3H), 3.72 - 3.77 (m, 2H); MS m/e MH^+ 257

ii) Methyl 4-methylpiperidine-4-carboxylate hydrochloride




4M HCl in dioxane (25 mL) was added to 1-*tert*-butyl 4-methyl 4-methylpiperidine-1,4-dicarboxylate (5.28 g, 20.55 mmol) at ambient temperature. The reaction mixture was allowed to stir at ambient temperature for 5 hrs. The solvent was removed under reduced pressure and ether was added to the residue and the suspension was filtered and dried to give the title compound as a white solid (3.82 g, 19.72 mmol, 96%); ^1H NMR δ 1.12 (s, 3H), 1.53 - 1.60 (m, 2H), 1.99 - 2.04 (m, 2H), 2.76 - 2.82 (m, 2H), 3.07 - 3.12 (m, 2H), 3.61 (s, 3H), 8.72 (s, 1H); MS m/e MH^+ 157

iii) Methyl 4-methyl-1-(5-nitropyridin-2-yl)piperidine-4-carboxylate



To a solution of 2-chloro-5-nitropyridine (2.0 g, 12.62 mmol) in DMSO (20 mL) was added sodium carbonate (6.70 g, 63.08 mmol) followed by methyl 4-methylpiperidine-4-carboxylate hydrochloride (2.44 g, 12.62 mmol). The reaction mixture was heated to 60 °C for 2 hours. The reaction mixture was poured onto water (100 mL) and the suspension filtered, washed with water (4 x 150 mL) and dried to leave the title compound as a yellow solid (3.32 g, 11.89 mmol, 94%); ^1H NMR (CDCl_3) δ 1.26 (s, 3H), 1.46 - 1.52 (m, 2H), 2.21 - 2.25 (m, 2H), 3.26 - 3.32 (m, 2H), 3.74 (s, 3H), 4.16 - 4.20 (m, 2H), 6.56 (d, 1H), 8.18 (d, 1H), 9.03 (s, 1H); MS m/e MH^+ 280.

iv) Methyl 1-(5-aminopyridin-2-yl)-4-methylpiperidine-4-carboxylate




To a solution of methyl 4-methyl-1-(5-nitropyridin-2-yl)piperidine-4-carboxylate (3.32 g, 11.89 mmol) in abs EtOH (200 mL) and THF (200 mL) was added 10% Pd/C (400 mg) and then the reaction mixture was evacuated with hydrogen (5 cycles) and allowed to stir at ambient temperature overnight. The reaction mixture was filtered and then concentrated and the black residue purified on a 120 g silica Crawford silicycle cartridge loading in 40%

- 392 -

EtOAc/isohexane with some DCM and eluting a gradient of 40-80% EtOAc using an Isco companion to provide the title compound (2.96 g, 11.89 mmol, 100%).



¹H NMR (CDCl₃) δ 1.23 (s, 3H), 1.51 - 1.58 (m, 2H), 2.16 - 2.20 (m, 2H), 2.98 - 3.05 (m, 2H), 3.25 (s, 2H), 3.67 - 3.70 (m, 2H), 3.70 (s, 3H), 6.57 (d, 1H), 6.97 (d, 1H), 7.78 (d, 1H); MS m/e MH⁺ 250.

v) Methyl 1-(5-([methoxy(oxo)acetyl]amino)pyridin-2-yl)-4-methylpiperidine-4-carboxylate




To a solution of methyl 1-(5-aminopyridin-2-yl)-4-methylpiperidine-4-carboxylate (2.96 g, 11.89 mmol) in DCM (100 mL) was added triethylamine (2.2 mL, 15.84 mmol) followed by methyl chlorooxoacetate (1.6 mL, 17.15 mmol) and the reaction mixture was allowed to stir at ambient temperature for 1.5 hrs. The reaction mixture was washed with brine (2 x 100 mL), the aqueous layer was re extracted with DCM (50 mL) and the organic washings were combined, dried and concentrated to leave the title compound as a brown gum (3.98 g, 11.89 mmol, 100%). ¹H NMR (CDCl₃) δ 1.72 - 1.82 (m, 2H), 1.96 - 2.02 (m, 2H), 2.52 - 2.59 (m, 1H), 2.95 - 3.03 (m, 2H), 3.70 (s, 3H), 3.97 (s, 3H), 4.16 - 4.22 (m, 2H), 6.67 (d, 1H), 7.93 (d, 1H), 8.28 (s, 1H), 8.66 (s, 1H); MS m/e MH⁺ 336.


vi) Methyl 1-(5-([hydrazino(oxo)acetyl]amino)pyridin-2-yl)piperidine-4-carboxylate
To a vigorously stirred suspension of methyl 1-(5-([methoxy(oxo)acetyl]amino)pyridin-2-yl)piperidine-4-carboxylate (3.12 g, 9.71 mmol) in absolute ethanol was added hydrazine monohydrate (0.52 mL, 10.76 mmol) and the suspension was allowed to stir at ambient temperature for 2 hours. The suspension was filtered, washed with ether (4 x 100 mL) and dried under vacuum at ambient temperature to provide the title compound (2.62 g, 7.81 mmol, 80%). ¹H NMR δ 1.18 (s, 3H), 1.40 - 1.47 (m, 2H), 1.98 - 2.03 (m, 2H), 3.07 - 3.14 (m, 3H), 3.66 (s, 3H), 3.80 - 3.86 (m, 2H), 4.58 (s, 1H), 6.83 (d, 1H), 7.90 (d, 1H), 8.49 (s, 1H), 10.17 (s, 1H), 10.46 (s, 1H); MS m/e MH⁺ 336.

Intermediate 110: Ethyl 2-[3-(4-aminophenyl)cyclobutyl]propanoate**i) Ethyl 2-(3-phenylcyclobutylidene)propanoate**

- 5 Following the general procedure described for ethyl {3-[4-(benzyloxy)phenyl]cyclohexylidene}acetate (Intermediate 89ii), replacing 3-[4-(benzyloxy)phenyl]cyclohexanone with 3-phenylcyclobutanone (*J. Org. Chem.*, 2005, 70(11), 4549-4552) and triethyl phosphonoacetate with triethyl phosphonopropionate the title compound was obtained; $^1\text{H NMR}$ δ (CDCl_3): 1.24-1.32 (3H, m), 1.75-1.76 (3H, m), 2.87-2.91 (1H, m), 3.19-3.23 (2H, m), 3.54-3.59 (2H, m), 4.09-4.21 (2H, m), 7.19-7.36 (5H, m); MS m/e MH^+ 231.
- 10

ii) Ethyl 2-(3-phenylcyclobutyl)propanoate

- 15 Following the general procedure described for ethyl 3-(4-aminobenzyl)cyclobutanecarboxylate (Intermediate 93ii), replacing ethyl 3-(4-nitrobenzyl)cyclobutanecarboxylate with ethyl 2-(3-phenylcyclobutylidene)propanoate the title compound was obtained without chromatography; $^1\text{H NMR}$ δ (CDCl_3): 1.08-1.10 (0.75H, d), 1.12-1.15 (3.25H, d), 1.22-1.30 (3H, m), 2.17-2.64 (6H, m), 3.34 (0.25H, m), 3.48-3.54 (0.75H, m), 4.09-4.18 (2H, m), 7.15-7.36 (5H, m); MS m/e MH^+ 233.
- 20

iii) Ethyl 2-[3-(4-nitrophenyl)cyclobutyl]propanoate

- Following the general procedure described for ethyl 3-(4-nitrobenzyl)cyclobutanecarboxylate (Intermediate 93i), replacing ethyl 3-
- 25


benzylcyclobutanecarboxylate with ethyl 2-(3-phenylcyclobutyl)propanoate the title compound was obtained which was used directly in the next step.

iv) Following the general procedure described for ethyl 3-(4-


- 5 aminobenzyl)cyclobutanecarboxylate (Intermediate 93ii), replacing ethyl 3-(4-nitrobenzyl)cyclobutanecarboxylate with ethyl 2-[3-(4-nitrophenyl)cyclobutyl]propanoate the title compound (Intermediate 110) was obtained; $^1\text{H NMR}$ δ (CDCl_3): 1.07-1.16 (3H, m), 1.22-1.27 (3H, m), 1.65-1.90 (1H, m), 2.11-2.62 (5H, m), 3.22-3.43 (1H, m), 4.09-4.16 (2H, m), 6.65 (2H, t), 6.98-7.07 (2H, m); MS m/e MH^+ 248.

10

Intermediate 111: Methyl 1-(5-{[hydrazino(oxo)acetyl]amino}-3-methylpyridin-2-yl)piperidine-4-carboxylate




i) Methyl 1-(3-methyl-5-nitropyridin-2-yl)piperidine-4-carboxylate



15


- To a solution of 2-chloro-3-methyl-5-nitropyridine (1.73 g, 10.00 mmol) in DMSO (20 mL) was added sodium carbonate (5.30 g, 50.00 mmol) followed by methyl isonipecotate (1.35 mL, 10.00 mmol). The reaction mixture was heated to 60°C for 110 minutes. The reaction mixture was left to stand at room temperature overnight. A further aliquot of methyl isonipecotate (3.33 mmol, 476 mg, 450 μL) was added and heating resumed at 60°C for 100mins. The reaction mixture was poured onto ice water and the suspension was filtered, washed with water (4 x 75 mL) and dried under vacuum to leave a yellow powder; (2.43 g, 87% yield). This was used in the next stage without further purification. $^1\text{H NMR}$ δ 1.61 - 1.74 (m, 2H), 1.91 - 2.10 (m, 2H), 2.31 (s, 3H), 2.61 - 2.73 (m, 1H), 2.94 - 3.09 (m, 2H), 3.61 (s, 3H), 3.83 - 3.98 (m, 2H), 8.19 (d, 1H), 8.86 (d, 1H). MS m/e MH^+ 280.18.
- 20
- 25

ii) Methyl 1-(5-amino-3-methylpyridin-2-yl)piperidine-4-carboxylate



Methyl 1-(3-methyl-5-nitropyridin-2-yl)piperidine-4-carboxylate (2.40 g) was dissolved in EtOH (86 mL), treated with 10% Pd/C (860 mg) and placed under an atmosphere of hydrogen (balloon). LCMS indicated complete reaction at ~100 mins. The catalyst was filtered off and solvent removed under vacuum to leave a colourless oil (2.16 g). This was used in the next step without further purification. ¹H NMR δ 1.58 - 1.75 (m, 2H), 1.81 - 1.91 (m, 2H), 2.10 (s, 3H), 2.37 - 2.50 (m, 1H), 2.55 - 2.75 (m, 2H), 2.99 - 3.09 (m, 2H), 3.61 (s, 3H), 6.76 (d, 1H), 7.48 (d, 1H). MS m/e MH⁺ 250.27.

iii) Methyl 1-(5-([methoxy(oxo)acetyl]amino)-3-methylpyridin-2-yl)piperidine-4-carboxylate




To an ice-water cooled solution of methyl 1-(5-amino-3-methylpyridin-2-yl)piperidine-4-carboxylate (2.14 g; 8.59 mmol) in DCM (43 mL) was added methyl chloro(oxo)acetate (1.03 mL; 11.17 mmol) followed by pyridine (834 μL; 10.31 mmol) and the reaction mixture was allowed to stir at ~0 °C for 30 mins, then allowed to warm to room temperature. The reaction solution was washed with water (40 mL) then saturated brine (40 mL). The organics were dried over MgSO₄, filtered and evaporated under vacuum to yield a pale tan solid (2.01 g, 70% yield). This was used for the next stage without further purification. ¹H NMR δ 1.60 - 1.80 (m, 2H), 1.85 - 1.99 (m, 2H), 2.18 (s, 3H), 2.54 - 2.62 (m, 1H), 2.69 - 2.88 (m, 2H), 3.26 - 3.40 (m, 2H), 3.56 (s, 3H), 3.82 (s, 3H), 7.87 (d, 1H), 8.40 (d, 1H), 10.75 (s, 1H). MS m/e MH⁺ 336.20.


iv) Methyl 1-(5-([hydrazino(oxo)acetyl]amino)-3-methylpyridin-2-yl)piperidine-4-carboxylate

Methyl 1-(5-([methoxy(oxo)acetyl]amino)-3-methylpyridin-2-yl)piperidine-4-carboxylate (2.01 g; 5.99 mmol) was dissolved in EtOH (140 mL) and treated with dropwise addition of hydrazine hydrate (320 μ L; 6.59 mmol). Within a few minutes a flocculent precipitate formed. Vigorous stirring was continued for 17 hrs. A further 32 μ L (0.65 mmol) hydrazine hydrate was added and stirring continued at ambient temperature for 24 hrs. The precipitated solid was filtered off, washed with ether on the filter and dried under high vacuum at 60°C to give the title compound (1.76g, 88% yield). This was used in the next stage without further purification. ^1H NMR δ 1.59 - 1.80 (m, 2H), 1.83 - 1.98 (m, 2H), 2.16 (s, 3H), 2.52 - 2.60 (m, 1H), 2.66 - 2.84 (m, 2H), 3.22 - 3.37 (m, 2H), 3.58 (s, 3H), 4.54 (d, 2H), 7.89 (d, 1H), 8.47 (d, 1H), 10.14 (s, 1H), 10.48 (s, 1H). MS m/e MH^+ 336.24.

Intermediate 112: Methyl [1-(5-([hydrazino(oxo)acetyl]amino)-3-methylpyridin-2-yl)piperidin-4-yl]acetate




i) Methyl [1-(3-methyl-5-nitropyridin-2-yl)piperidin-4-yl]acetate



Prepared as in Intermediate 111 part i), using methyl piperidin-4-ylacetate as starting material. ^1H NMR δ 1.20 - 1.41 (m, 2H), 1.68 - 1.83 (m, 2H), 1.87 - 2.06 (m, 1H), 2.26 - 2.35 (m, 5H), 2.85 - 3.04 (m, 2H), 3.55 (s, 3H), 3.88 - 3.99 (m, 2H), 8.17 (d, 1H), 8.86 (d, 1H). MS m/e MH^+ 294.24.


- 397 -

ii) Methyl [1-(5-amino-3-methylpyridin-2-yl)piperidin-4-yl]acetate



Prepared following the method of Intermediate 111 part ii). ^1H NMR δ 1.31 (m, 2H), 1.69 (m, 2H), 1.74 (m, 1H), 2.09 (s, 3H), 2.28 (d, 2H), 2.57 (m, 2H), 3.02 (m, 2H), 3.59 (s, 3H), 4.73 (s, 2H), 6.75 (d, 1H), 7.28 (d, 1H). MS m/e MH^+ 264.23.

iii) Methyl ((6-[4-(2-methoxy-2-oxoethyl)piperidin-1-yl]-5-methylpyridin-3-yl)amino)(oxo)acetate




Prepared following the method of Intermediate 111 part iii). ^1H NMR δ 1.24 - 1.44 (m, 2H), 1.67 - 1.79 (m, 2H), 1.79 - 1.93 (m, 1H), 2.17 (s, 3H), 2.30 (d, 2H), 2.61 - 2.78 (m, 2H), 3.26 - 3.41 (m, 2H), 3.57 (s, 3H), 3.89 (s, 3H), 7.85 (d, 1H), 8.39 (d, 1H), 10.72 (s, 1H). MS m/e MH^+ 350.23.


iv) Methyl [1-(5-([hydrazino(oxo)acetyl]amino)-3-methylpyridin-2-yl)piperidin-4-yl]acetate

Prepared following the method of Intermediate 111 part iv). ^1H NMR δ 1.23 - 1.44 (m, 2H), 1.66 - 1.79 (m, 2H), 1.77 - 1.93 (m, 1H), 2.16 (s, 3H), 2.30 (d, 2H), 2.59 - 2.76 (m, 2H), 3.22 - 3.40 (m, 2H), 3.56 (s, 3H), 4.52 (s, 2H), 7.88 (d, 1H), 8.46 (d, 1H), 10.14 (s, 1H), 10.51 (s, 1H). MS m/e MH^+ 350.21.

Intermediate 113: Ethyl 2-[1-(5-[[hydrazino(oxo)acetyl]amino]-3-methylpyridin-2-yl)piperidin-4-yl]propanoate




i) *tert*-Butyl 4-(2-methoxy-1-methyl-2-oxoethylidene)piperidine-1-carboxylate



To a suspension of 60% NaH (2.005 g; 50.00 mmol) in anhydrous THF (100 mL), under an atmosphere of argon, was added triethyl 2-phosphonopropionate (10.73 mL; 50.00 mmol), at such a rate as to keep the reaction temperature below 30 °C. The resulting mixture was stirred at room temperature for a further 1hr. A solution of *N*-Boc-4-piperidone (9.965 g; 50.00 mmol), in anhydrous THF (50 mL), was added dropwise over five minutes. The resulting clear solution was stirred at ambient temperature for ~1hr. The reaction was quenched by the addition of water (100 mL). The resulting mixture was partitioned, the aq layer extracted once with EtOAc and the combined organics washed with saturated brine. The organics were dried over MgSO₄, filtered and evaporated to a colourless oil; 14.76 g. (~quantitative yield). This was used in the next stage without further purification. ¹H NMR δ 1.19 (t, 3H), 1.36 (s, 9H), 1.77 (s, 3H), 2.25 - 2.37 (m, 2H), 2.51 - 2.56 (m, 2H), 3.22 - 3.44 (m, 4H), 4.11 (q, 2H). MS m/e [MH - (CO₂, C₄H₈)]⁺ = 184.31.

ii) Ethyl 2-piperidin-4-ylidenepropanoate hydrochloride




tert-Butyl 4-(2-methoxy-1-methyl-2-oxoethylidene)piperidine-1-carboxylate (2.83 g; 10 mmol) was dissolved in EtOH (50 mL) and treated with 4M HCl / 1,4-dioxane solution (20 mL, 80 mmol). The resulting solution was stirred at ambient temperature for 6 hrs. The

- 399 -

solvents were removed under vacuum to leave a syrup. This was dried under vacuum to give the title product (3.199 g, ~quantitative yield). This was used in the next stage without further purification. ^1H NMR δ 1.22 (t, 3H), 1.78 (s, 3H), 2.44 - 2.57 (m, 2H), 2.65 - 2.77 (m, 2H), 2.98 - 3.15 (m, 4H), 4.14 (q, 2H), 9.52 (s, 2H). MS m/e MH^+ = 184.32.

5

iii) Ethyl 2-[1-(3-methyl-5-nitropyridin-2-yl)piperidin-4-ylidene]propanoate




Prepared as in Intermediate 111 part i) using ethyl 2-piperidin-4-ylidenepropanoate.

^1H NMR δ 1.23 (t, 3H), 1.89 (s, 3H), 2.39 (s, 3H), 2.49 - 2.56 (m, 2H), 2.67 - 2.78 (m, 2H), 3.47 - 3.66 (m, 4H), 4.14 (q, 2H), 8.18 (d, 1H), 8.86 (d, 1H). MS m/e MH^+ 320.21.

10

iv) Ethyl 2-[1-(5-amino-3-methylpyridin-2-yl)piperidin-4-yl]propanoate




Ethyl 2-[1-(3-methyl-5-nitropyridin-2-yl)piperidin-4-ylidene]propanoate (2.30 g; 7.20 mmol) was suspended in EtOH (100 mL) and treated with 10% Pd/C (230 mg). The resulting mixture was placed under an atmosphere of hydrogen (balloon) at ambient temperature and stirred vigorously at ambient temperature for 2 hrs. LCMS indicated no reduction of alkene but reduction of the nitro-group was essentially complete. The catalyst was filtered off and the filtrate was evaporated to a gum (2.42 g). The reaction was repeated under 20 bar of hydrogen, at 30 °C, for 3 hrs with 241 mg 10% Pd/C catalyst. The catalyst was filtered off (Celite) and the solvent removed under vacuum to leave a dark blue gum (2.12 g). This was used in the next step without further purification. ^1H NMR δ 1.06 (d, 3H), 1.18 (t, 3H), 1.24 - 1.43 (m, 2H), 1.45 - 1.63 (m, 2H), 1.61 - 1.74 (m, 1H), 2.06 (s, 3H), 2.21 - 2.35 (m, 1H), 2.50 - 2.63 (m, 1H), 2.99 - 3.13 (m, 2H), 3.36 - 3.50 (m, 1H), 3.98 - 4.14 (m, 2H), 4.65 (s, 2H), 6.76 (d, 1H), 7.47 (d, 1H). MS m/e MH^+ 292.27.

15

20

25

v) Ethyl 2-[1-(5-{[methoxy(oxo)acetyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]propanoate



Prepared following the method described for Intermediate 111 part iii).


5 ¹H NMR δ 1.07 (d, 3H), 1.18 (t, 3H), 1.28 - 1.45 (m, 2H), 1.54 - 1.67 (m, 2H), 1.67 - 1.80 (m, 1H), 2.17 (s, 3H), 2.24 - 2.36 (m, 1H), 2.59 - 2.78 (m, 2H), 3.31 - 3.47 (m, 2H), 3.82 (s, 3H), 4.01 - 4.15 (m, 2H), 7.86 (d, 1H), 8.39 (d, 1H), 10.76 (s, 1H). MS m/e MH⁺ 378.17.

10 vi) Ethyl 2-[1-(5-{[hydrazine(oxo)acetyl]amino}-3-methylpyridin-2-yl)piperidin-4-yl]propanoate

Prepared following the method described for Intermediate 111 part iv)


15 ¹H NMR δ 1.06 (d, 3H), 1.19 (t, 3H), 1.24 - 1.46 (m, 2H), 1.52 - 1.67 (m, 2H), 1.66 - 1.81 (m, 1H), 2.15 (s, 3H), 2.24 - 2.36 (m, 1H), 2.53 - 2.75 (m, 2H), 3.30 - 3.40 (m, 2H), 4.07 (q, 2H), 4.36 (s, 1H), 4.70 (s, 1H), 7.87 (d, 1H), 8.46 (d, 1H), 9.80 (s, 1H), 10.51 (s, 1H). MS m/e MH⁺ 378.22.

Intermediate 114: Methyl (1-{5-[(5-{(3,4-difluorophenyl)amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl)acetate



20 MS m/e MH⁺ 487. This intermediate is also a compound of formula (I) and is a further feature of the invention.


i) Methyl [4-methyl-1-(5-nitropyridin-2-yl)piperidin-4-yl]acetate



- 401 -


To a solution of 2-chloro-5-nitropyridine (2.97 g, 18.77 mmol) in DMSO (30 mL) was added sodium carbonate (6.4 g, 60.38 mmol) followed by methyl (4-methylpiperidin-4-yl)acetate (3.21g, 18.77 mmol) (prepared according to the procedures described by Janet Ralbovsky, Joseph Lisko and Wei He, *Synthetic Communications*, 2005, 35 (12), 1613-1625 and Gerhard Hund, and Woldemar Schneider, *Chem Ber* 1980, 113 (2), 401-7). The reaction mixture was heated to 60 °C for 1.5 hours, then allowed to cool and water (100 mL) was added. The suspension filtered, and dried to leave a yellow solid (3.70 g, 12.62mmol, 67%). ¹H NMR δ 1.08 (s, 3H), 1.45 - 1.48 (m, 2H), 1.55 - 1.61 (m, 2H), 2.38 (s, 2H), 3.59 (s, 3H), 3.64 - 3.70 (m, 2H), 3.85 - 3.91 (m, 2H), 6.95 (d, 1H), 8.19 (d, 1H), 8.95 (s, 1H); MS m/e MH⁺ 294.

ii) Methyl [1-(5-aminopyridin-2-yl)-4-methylpiperidin-4-yl]acetate



Prepared from methyl [4-methyl-1-(5-nitropyridin-2-yl)piperidin-4-yl]acetate following the method for Intermediate 77 stage ii). ¹H NMR δ 1.01 (s, 3H), 1.41 - 1.45 (m, 2H), 1.52 - 1.58 (m, 2H), 2.30 (s, 2H), 3.12 - 3.19 (m, 2H), 3.34 - 3.38 (m, 2H), 3.58 (s, 3H), 4.52 (s, 2H), 6.62 (d, 1H), 6.89 (d, 1H), 7.58 (s, 1H); MS m/e MH⁺ 264.


iii) Methyl ({6-[4-(2-methoxy-2-oxoethyl)-4-methylpiperidin-1-yl]pyridin-3-yl}amino)(oxo)acetate



To a solution of methyl [1-(5-aminopyridin-2-yl)-4-methylpiperidin-4-yl]acetate (3.29 g, 13.20 mmol) in DCM (100 mL) was added triethylamine (2.2 mL, 15.84 mmol) followed by methyl chlorooxoacetate (1.6 mL, 17.15 mmol) and the reaction mixture was allowed to stir at ambient temperature for 1.5 hrs. The reaction mixture was washed brine (2 x 100 mL), the aqueous layer was re extracted with DCM (50 mL) the organic washings were combined, dried and concentrated to leave the title compound as a brown gum (4.4 g, 13.2 mmol, 100%). ¹H NMR δ 1.06 (s, 3H), 1.40 - 1.46 (m, 2H), 1.53 - 1.57 (m, 2H), 2.36 (s,

3H), 3.46 - 3.52 (m, 2H), 3.59 (s, 3H), 3.67 - 3.73 (m, 2H), 3.78 (s, 3H), 6.90 (d, 1H), 7.56 (d, 1H), 8.06 (s, 1H); MS m/e MH⁺ 350.

iv) Methyl [1-(5-[[hydrazino(oxo)acetyl]amino]pyridin-2-yl)-4-methylpiperidin-4-yl]acetate




Prepared from the product of the previous reaction and hydrazine hydrate following the method of Intermediate 47 part ii).

¹H NMR δ 1.04 (s, 3H), 1.40 - 1.45 (m, 2H), 1.51 - 1.58 (m, 2H), 2.33 (s, 2H), 3.36 - 3.42 (m, 2H), 3.57 - 3.62 (m, 5H, inc s, 3H), 4.48 (s, 1H), 4.60 (s, 1H), 6.83 (d, 1H), 7.90 (d, 1H), 8.50 (s, 1H), 9.96 (s, 1H), 10.23 (s, 1H); MS m/e MH⁺ 350.

v) Methyl (1-[5-[(5-[(3,4-difluorophenyl)amino]-1,3,4-oxadiazol-2-yl)carbonyl]amino]pyridin-2-yl)-4-methylpiperidin-4-yl)acetate

Prepared from the product of the previous reaction and 3,4-difluorophenylisothiocyanate, following the method of Example 319, but the product was taken through directly to the hydrolysis stage; MS m/e MH⁺ 487.

The following intermediates were prepared from the hydrazide product of Intermediate A part iv) and the appropriate isothiocyanate following the method of Example 319. They were taken through crude to the hydrolysis stage without further purification or characterisation:




Intermediate	Name	MS m/e M+H ⁺
115 R ¹ =R ² =R ³ =F	Methyl (4-methyl-1-[5-[(5-[(2,4,5-trifluorophenyl)amino]-1,3,4-oxadiazol-2-	505

	yl}carbonyl)amino]pyridin-2-yl}piperidin-4-yl]acetate	
116 $R^1=R^3=H, R^2=CF_3$	Methyl [4-methyl-1-(5-{[(5-{[4-(trifluoromethyl)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)piperidin-4-yl]acetate	519
117 $R^1=R^3=H, R^2=O(4-F-C_6H_4)$	Methyl {1-(5-{[(5-{[4-(4-fluorophenoxy)phenyl]amino}-1,3,4-oxadiazol-2-yl)carbonyl]amino}pyridin-2-yl)-4-methylpiperidin-4-yl]acetate	561

Claims

1. A compound of formula (I)



(I)

or a pharmaceutically-acceptable salt or prodrug thereof, wherein:

R^1 is an optionally substituted aryl or optionally substituted heteroaryl group, wherein the optional substituents are one or more groups selected from a group $-Z^a$, a group $-X^2-(CR^{52}R^{53})_w-Z^a$, a group $-X^2-(CR^{52}R^{53})_n-X^3-Z^a$, a group $-(CR^{52}R^{53})_aX^3-Z^a$ or a functional group (which is other than a group $-X^2-(CR^{52}R^{53})_w-Z^a$ or a group $-X^2-(CR^{52}R^{53})_n-X^3-Z^a$);

Y is a direct bond, or a group $(CR^{40}R^{41})_s$ or $-X^6(CR^{40}R^{41})_t-$ where each R^{40} and R^{41} is independently selected from hydrogen, (1-4C)alkyl, hydroxy, halo, halo(1-4C)alkyl, amino, cyano, (1-4C)alkoxy, (1-4C)haloalkoxy or $((1-3C)alkyl)CONH-$, s is an integer of from 1 to 6 and t is an integer of from 1 to 6, provided that the X^6 atom of the group $-X^6(CR^{40}R^{41})_t-$ is attached to the R^2 group and that a single sp^3 hybridised carbon atom does not carry two or more bonds to a heteroatom unless the heteroatom is a halo;

R^2 is an optionally substituted aryl, an optionally substituted cycloalkyl or an optionally substituted heterocyclic group, wherein optional substituents are one or more groups selected from a group $-Z$, a group $-X-(CR^{42}R^{43})_u-Z$, a group $-X-(CR^{42}R^{43})_v-X^1-Z$ or a group $-(CR^{42}R^{43})_vX^1-Z$ or a functional group (which is other than a group $-X-(CR^{42}R^{43})_v-Z$ or a group $-X-(CR^{42}R^{43})_v-X^1-Z$);

wherein Z and Z^a are independently selected from a hydrocarbyl group or a heterocyclic group or a combination thereof, wherein the group Z and Z^a is optionally substituted on any available atom by one or more functional groups, or by a group $-X^7-(CR^{62}R^{63})_bR^{64}$;

X , X^1 , X^2 , X^3 , X^6 and X^7 are linking groups independently selected from $-C(O)_x-$, $-O-$, $-S(O)_y-$, $-NR^{44}-$, $-C(O)NR^{44}-$, $-OC(O)NR^{44}-$, $-CH=NO-$, $-NR^{44}C(O)_x-$, $-NR^{44}CONR^{45}-$, $-S(O)_2NR^{44}-$ or $-NR^{44}S(O)_2-$ where x is an integer of 1 or 2, y is 0, 1 or 2, and R^{44} and R^{45} are independently selected from hydrogen or C_{1-6} alkyl,

u and w are independently selected from 0 or an integer of from 1 to 6;


- 405 -

- v, a and b are independently selected from an integer of from 1 to 6;
each R^{42} , R^{43} , R^{52} , R^{53} , R^{62} and R^{63} is independently selected from hydrogen, (1-4C)alkyl, hydroxy, halo, halo(1-4C)alkyl, amino, cyano, (1-4C)alkoxy, (1-4C)haloalkoxy, ((1-3C)-alkyl)CONH-, carboxy or a carboxylic acid mimic or bioisostere thereof, and
- 5 R^{64} is a functional group.
2. A compound according to claim 1 or a pharmaceutically-acceptable salt or prodrug thereof, wherein R^1 is an optionally substituted aryl group.
- 10 3. A compound according to claim 1 or a pharmaceutically-acceptable salt or prodrug thereof, wherein R^1 is an optionally substituted monocyclic heteroaryl group.
4. A compound according to any one of the preceding claims, or a pharmaceutically-acceptable salt or prodrug thereof, wherein R^1 is unsubstituted or is
- 15 substituted by 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino.
- 20 5. A compound according to claim 1 or a pharmaceutically-acceptable salt or prodrug thereof, wherein R^1 is selected from phenyl, naphthyl, indanyl, pyrimidinyl, pyridyl, pyrazolyl, pyrazinyl, thiazolyl, oxadiazolyl, isoxazolyl, thiadiazolyl, pyrrolopyridyl, 1,3-benzodioxan-5-yl, benzthiazolyl, benzimidazolyl and quinolyl; optionally substituted with 1, 2 or 3 substituents independently selected from
- 25 halo, (1-4C)alkyl, ethynyl, (1-4C)alkoxy, hydroxy, (1-4C)alkoxy(1-4C)alkoxy, methoxymethyl, cyanomethyl, hydroxy(1-4C)alkyl, trifluoromethyl, difluoromethoxy, trifluoromethoxy, trifluoromethylthio, cyano, methylthio, methylsulfonyloxy, methylsulfonyl, ethylsulfonyl, aminocarbonylamino, methoxycarbonylamino, methylcarbonylamino, (1-4C)alkoxycarbonyl, methoxycarbonylmethoxy, benzyloxy,
- 30 pyridylmethoxy, phenoxy (optionally substituted by methoxy or halo), phenyl (optionally substituted by methoxycarbonyl or carboxy), benzyl, anilino, anilino-carbonyl,

aminocarbonyl, benzoyl, benzoylamino, phenylsulfonyl, aminosulfonyl, cyclohexyl, methylpyrimidinyl, triazolyl and morpholino.

6. A compound according to any one of the preceding claims wherein Y is a direct
5 bond.

7. A compound according to any one of the preceding claims wherein R² is a 5- or
6-membered aromatic ring of sub-structure (a)



(a)

Z¹, Z², Z³ and Z⁴ are independently selected from -CH-, -CR⁶- or a heteroatom
selected from O, S, N(R⁵⁰)_r, where r is 0 or 1 depending upon the requirements of the
aromatic ring, and R⁵⁰ is hydrogen or C₁₋₆alkyl, and Z⁴ may additionally be a direct bond,

15 R⁴ is a group -Z, a group -X-(CR⁴²R⁴³)_u-Z, a group -X-(CR⁴²R⁴³)_v-X¹-Z or a group -
(CR⁴²R⁴³)_v-X¹-Z, wherein Z, X, X¹, R⁴², R⁴³, u and v are as defined in claim 1,


each R⁶ is independently selected from halo, cyano, nitro, amino, hydroxy,
haloC₁₋₆alkyl, a group -Z, a group -X-(CR⁴²R⁴³)_u-Z, a group -X-(CR⁴²R⁴³)_v-X¹-Z or a group
-(CR⁴²R⁴³)_v-X¹-Z, wherein Z, X, X¹, R⁴², R⁴³, u and v are as defined in claim 1.

20

8. A compound according to any one of the preceding claims wherein R² is substituted
by a group Z.

9. A compound according to claim 6 or claim 7 wherein Z is a group of sub formula
25 (x), (y) or (z):

- 407 -



wherein each ring A or A' is independently selected from an optionally substituted heterocyclic ring, an optionally substituted cycloalkyl ring or an optionally substituted aryl ring, each R⁶⁰ is an optionally substituted (1-6C)alkyl, an optionally substituted (2-6C)alkenyl or an optionally substituted (2-6C)alkynyl, and R⁶¹ is an optionally substituted (1-6C)alkylene, an optionally substituted (2-6C)alkenylene or an optionally substituted (2-6C)alkynylene.


10. A compound according to claim 1 or claim 5, or a pharmaceutically-acceptable salt or prodrug thereof, wherein R²-Y is selected from:
- 4-(piperazino)phenyl (substituted on the available piperazine nitrogen by a substituent selected from (1-4C)alkyl, acyl, benzyl, ethoxycarbonyl, tert-butoxycarbonyl, pyridyl, cyclopropylmethyl and methoxyethyl),
- 4-(piperazino)-3-methyl-phenyl (substituted on the available piperazine nitrogen by a substituent selected from acyl and tert-butoxycarbonyl),
- 4-(piperazino)pyridyl (substituted on the available piperazine nitrogen by a substituent selected from (1-4C)alkyl, acyl, ethoxycarbonyl, tert-butoxycarbonyl, benzoyl, fluorobenzoyl, cyanobenzoyl, methylbenzoyl, pyridylcarbonyl, methylsulfonyl, methoxymethylcarbonyl, phenethylcarbonyl, anilino carbonyl, benzylaminocarbonyl, phenoxycarbonyl, benzyloxycarbonyl and aminocarbonyl),
- 4-piperidinophenyl, benzyloxyphenyl, (1-4C)alkylphenyl, pyridylmethoxyphenyl, N-methylanilinophenyl, biphenyl, (chloro)biphenyl, (methoxycarbonyl)biphenyl, (ethoxycarbonylmethyl)biphenyl, (carboxy)biphenyl, morpholinophenyl, dimethylmorpholinophenyl, (morpholino)(chloro)phenyl, (morpholino)(dichloro)phenyl

- (morpholino)(bromo)phenyl, (morpholino)(methyl)phenyl, (morpholino)(fluoro)phenyl, (morpholino)(cyano)phenyl, (morpholino)(allyloxycarbonyl)phenyl, (morpholinoocthoxy)phenyl, morpholinopyridyl, (morpholino)(chloro)pyridyl, (morpholino)(fluoro)pyridyl, phenoxypyridyl, anilinopyridyl, methoxypyridyl, 5 butoxypyridyl, pyridyloxypyridyl, hydroxybutylpyridyl, thiomorpholinopyridyl, phenoxypropyl, (fluoro)(trifluoromethyl)phenyl, (N-acyl)homopiperazinophenyl, (N-acyl)piperazinopyrimidinyl, methylbenzthiazolyl, butylthiophenyl, (methoxythiadiazolyl)aminosulfonylphenyl, (difluorophenoxy)phenyl, (dimethyl-oxo-pyridazinyl)phenyl, (methyl-oxo-thiadiazinyl)phenyl, methoxycarbonylpropylphenyl, 10 (methoxycarbonyl)methylthiophenyl, (methoxycarbonyl)adamantyl, carboxyadamantyl, (methoxycarbonylmethyl)bi-cyclohexyl, (carboxy)bi-cyclohexyl, (ethoxycarbonylethyl)indanyl, (carboxyethyl)indanyl, (methoxycarbonyl)phenoxypropyl, (carboxy)phenoxypropyl, (carboxypropyl)phenyl, dicarboxybutylphenyl, carboxybutylphenyl, propylsulfonylphenyl, (carboxy)isopropylsulfonylphenyl, 15 phenoxyphenyl, (methoxyphenoxy)phenyl, methylphenoxyphenyl, hydroxypropoxyphenyl, ethoxyethoxyphenyl, (dimethylaminocarbonyl)isopropoxyphenyl, (carboxymethyl)piperidinophenyl, (methoxycarbonylmethyl)piperidinopyridyl, (carboxymethyl)piperidinopyridyl, (carboxy)piperidinopyridyl, (methoxycarbonyl)piperidinopyridyl, (carboxy)(methyl)piperidinopyridyl, 20 (methoxycarbonyl)(methyl)piperidinopyridyl, (carboxymethyl)(methyl)piperidinopyridyl, (methoxycarbonyl)piperidyl[(methyl)pyridyl], (carboxy)piperidyl[(methyl)pyridyl], (methoxycarbonylmethyl)piperidyl[(methyl)pyridyl], (carboxymethyl)piperidyl[(methyl)pyridyl], (carboxyethyl)piperidyl[(methyl)pyridyl], (ethoxycarbonylethyl)piperidyl[(methyl)pyridyl], 25 [(methoxycarbonyl)piperidinocarbonylmethyl]cyclohexylphenyl, [(carboxy)piperidinocarbonylmethyl]cyclohexylphenyl, [(methoxycarbonyl)isopropylaminocarbonylmethyl]cyclohexylphenyl, [(carboxy)isopropylaminocarbonylmethyl]cyclohexylphenyl, [(methoxycarbonyl)pyrrolidinylcarbonylmethyl]cyclohexylphenyl, 30 [(carboxy)pyrrolidinylcarbonylmethyl]cyclohexylphenyl, [(methoxycarbonyl)(hydroxy)pyrrolidinylcarbonylmethyl]cyclohexylphenyl, [(carboxy)(hydroxy)pyrrolidinylcarbonylmethyl]cyclohexylphenyl,

- 409 -

- (hydroxyisobutylaminocarbonylmethyl)cyclohexylphenyl,
 [(tetrahydrodioxolopyrrolyl)carbonyl]methylcyclohexylphenyl,
 (methoxycarbonylpyridyl)piperidino, (carboxypyridyl)piperidino,
 (aminocarbonylpyridyl)piperidino, (carboxymethylpyridyl)piperidino,
 5 (ethoxycarbonylmethyl)cyclohexylphenyl, (methoxycarbonylmethyl)cyclohexylphenyl,
 cyclohexylphenyl, (carboxymethyl)cyclohexylphenyl, (hydroxyethyl)cyclohexylphenyl,
 (aminocarbonylmethyl)cyclohexylphenyl, (cyanomethyl)cyclohexylphenyl,
 carboxycyclohexylphenyl, (dimethylaminocarbonylmethyl)cyclohexylphenyl, [(N-
 hydroxyethyl-N-methylaminocarbonylmethyl)cyclohexylphenyl, [N-
 10 (dihydroxypropyl)aminocarbonylmethyl]cyclohexylphenyl,
 (aminomethyl)cyclohexylphenyl, (tertbutoxycarbonylaminomethyl)cyclohexylphenyl,
 (methoxycarbonyl)cyclohexyloxyphenyl, (ethoxycarbonyl)cyclohexyloxyphenyl,
 carboxycyclohexyloxyphenyl, (methoxycarbonyl)cyclopentylphenyl,
 (ethoxycarbonylmethyl)cyclopentylphenyl, (carboxy)cyclopentylphenyl
 15 (carboxymethyl)cyclopentylphenyl, (methoxycarbonyl)cyclobutylphenyl,
 (ethoxycarbonyl)cyclobutylmethylphenyl, (carboxy)cyclobutylphenyl,
 (carboxy)cyclobutylmethylphenyl, (carboxyethyl)cyclobutylphenyl,
 (ethoxycarbonyl)ethylcyclobutylphenyl, (carboxymethyl)phenylcyclohexyl, 1-
 (ethoxycarbonyl)ethylcyclohexylphenyl, 1-carboxyethylcyclohexylphenyl, (2-
 20 methoxycarbonyl)isopropylcyclohexyl, (2-carboxy)isopropylcyclohexyl,
 pyrrolidinylphenyl, (methoxycarbonylmethyl)cyclohexylpyridyl and
 (carboxymethyl)cyclohexylpyridyl.


11. A compound of formula (I) as claimed in claim 1, which is a compound of formula
 25 (IZA), or a pharmaceutically-acceptable salt or pro-drug thereof,



(IZA)

- 410 -

- wherein R^1 is selected from phenyl (optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino), 2-pyridyl (optionally substituted by chlorophenoxy, chlorobenzyloxy or methoxyphenoxy, and/or substituted with a substituent selected from halo, trifluoromethyl, (1-4C)alkyl, (1-4C)alkoxy and cyano), 3-pyridyl (optionally substituted as for 2-pyridyl), halopyrimidinyl and trifluoromethylthiazolyl;
- Z^2 is N or CH;
- 10 R^{ZA1} and R^{ZA2} are each independently hydrogen or methyl;
- R^{ZA3} is hydrogen or methyl;
- R^{6ZA} is hydrogen, fluoro, chloro or methyl;
- A is N or CH;
- X^{ZA} is a direct bond, $-CH_2-$ or $-O-$ (except when A is N);
- 15 m is 0, 1 or 2;
- n is 0 or 1;
- provided that $m + n = 0, 1$ or 2;
- p is 0 or 1.
- 20 12. A compound of formula (I) as claimed in claim 1, which is a compound of formula (IZB), or a pharmaceutically-acceptable salt or pro-drug thereof,



(IZB)

- 25 wherein R^1 is selected from phenyl (optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino), 2-pyridyl

- 411 -


(optionally substituted by chlorophenoxy, chlorobenzoyloxy or methoxyphenoxy, and/or substituted with a substituent selected from halo, trifluoromethyl, (1-4C)alkyl, (1-4C)alkoxy and cyano), 3-pyridyl (optionally substituted as for 2-pyridyl), halopyrimidinyl and trifluoromethylthiazolyl;

5 Z^2 is N or CH;

R^{6ZB} is hydrogen, fluoro, chloro or methyl;

X^{ZB} is O or S, particularly O.

13. A compound of formula (I) as claimed in claim 1, which is a compound of formula
10 (IZC), or a pharmaceutically-acceptable salt or pro-drug thereof,




(IZC)

wherein R^1 is selected from phenyl (optionally substituted with 1, 2 or 3 substituents independently selected from fluoro, chloro, bromo, trifluoromethyl, methoxy,
15 difluoromethoxy, trifluoromethoxy, cyano, methyl, ethyl, ethynyl, benzyloxy, 3-chlorobenzoyloxy, phenoxy, 4-chlorophenoxy, phenyl, benzoyl and anilino), 2-pyridyl (optionally substituted by chlorophenoxy, chlorobenzoyloxy or methoxyphenoxy, and/or substituted with a substituent selected from halo, trifluoromethyl, (1-4C)alkyl, (1-4C)alkoxy and cyano), 3-pyridyl (optionally substituted as for 2-pyridyl), halopyrimidinyl and
20 trifluoromethylthiazolyl;
 Z^2 is N or CH;
 R^{6ZC} is hydrogen, fluoro, chloro or methyl;
 R^{ZC} is selected from (1-4C)alkyl, acyl, benzyl, ethoxycarbonyl, tert-butoxycarbonyl, pyridyl, cyclopropylmethyl, methoxyethyl, benzoyl, fluorobenzoyl, cyanobenzoyl, methylbenzoyl,
25 pyridylcarbonyl, methylsulfonyl, methoxymethylcarbonyl, phenethylcarbonyl, anilino carbonyl, benzylaminocarbonyl, phenoxycarbonyl, benzyloxycarbonyl and aminocarbonyl.


14. A compound as claimed in Claim 1, or a pharmaceutically-acceptable salt or pro-drug thereof, which is any one of Examples 1 to 712 herein.
15. A compound according to any one of the preceding claims or a
5 pharmaceutically-acceptable salt or prodrug thereof for use as a medicament.
16. A method for producing an inhibition of DGAT1 activity in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) as claimed in any one of
10 claims 1 to 14 or a pharmaceutically-acceptable salt thereof.
17. A method of treating diabetes mellitus and/or obesity in a warm-blooded animal, such as a human being, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I) as claimed in any one of claims 1
15 to 14 or a pharmaceutically-acceptable salt thereof.
18. The use of a compound according to any one of claims 1 to 14 or a pharmaceutically-acceptable salt or prodrug thereof in the manufacture of a medicament for use in the production of an inhibition of DGAT1 activity in a warm-blooded animal such as
20 a human being.
19. The use as claimed in Claim 18 wherein the medicament is for use in the treatment of diabetes mellitus and/or obesity in a warm-blooded animal such as a human being.
- 25 20. A pharmaceutical composition which comprises a compound of formula (I) as claimed in any one of claims 1 to 14 or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable excipient or carrier.
21. A process for preparing a compound according to claim 1 which comprises one of
30 the following steps (wherein all variables are as hereinbefore defined for a compound of formula (I) unless otherwise stated):
- a) reaction of a compound of formula (I) to form another compound of formula (I);

- 413 -


b) reaction of an amine of formula (2) with a carboxylate salt of formula (3);




c) cyclisation of a compound of formula (4) (where X is S or O);




d) when R² is substituted by piperazinyl, by reaction of the piperazine nitrogen with R⁵-LG wherein LG is a suitable leaving group such as halo, and R⁵ is hydrocarbyl or a suitable functional group such as acyl, for example:



e) when R² is aryl and is substituted by aryl, by transition metal catalysed aromatic substitution, for example:




f) when R² is substituted by piperazinyl, by reductive alkylation of the piperazine nitrogen with R⁵-CHO (wherein R⁵ is for example hydrocarbyl), for example:



- 414 -

g) reaction of halogenated (for example with iodo) R^2 with an amide of formula (7) followed by subsequent removal of protecting group P^1 , wherein P^1 is for example benzyl or trimethylsilylethoxymethyl (SEM), for example;



5

wherein Hal is halogen;

and thereafter if necessary, removing any protecting groups, and/or forming a pharmaceutically-acceptable salt or prodrug thereof.

10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2005/004726

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D271/10 C07D413/12 A61K31/4245 A61P3/04 A61P3/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/20484 A (ASTRAZENECA AB; SANGANEE, HITEH; SPRINGTHORPE, BRIAN) 14 March 2002 (2002-03-14) page 2, line 13 - page 4, line 1 page 25, line 24; example 31	1, 2, 4, 5, 15-21
A	WO 2004/100881 A (BAYER PHARMACEUTICALS CORPORATION; SMITH, ROGER; CAMPBELL, ANN-MARIE;) 25 November 2004 (2004-11-25) cited in the application paragraph [0006]; claim 1	1-21

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earliest document but published on or after the international filing date

L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

17 March 2006

Date of mailing of the international search report

24/03/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 6819 Patentkan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seymour, L

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2005/004726

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 16 and 17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2005/004726

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0220484	A	14-03-2002	AU 6458401 A	22-03-2002
			EP 1322611 A1	02-07-2003
			JP 2004508355 T	18-03-2004
			US 2004102432 A1	27-05-2004
WO 2004100881	A	25-11-2004	AU 2004238258 A1	25-11-2004
			CA 2524470 A1	25-11-2004
			EP 1633722 A2	15-03-2006

(12) SOLICITUD INTERNACIONAL PUBLICADA SEGÚN EL TRATADO DE COOPERACIÓN INTERNACIONAL EN MATERIA DE PATENTES (PCT)

(19) Organización Mundial de Propiedad Intelectual
Oficina Internacional



(43) Fecha de Publicación Internacional
22 de junio de 2006 (22.06.2006)

(10) Número de Publicación Internacional
WO 2006/064189 A1

PCT

(51) Clasificación Internacional de Patentes:
C07D 271/10 (2006.01) A61P 3/04 (2006.01)
C07D 413/12 (2006.01) A61P 3/10 (2006.01)
A61K 31/4245 (2006.01)

Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
MCCOULL, William [GB/GB]; AstraZeneca R & D
Alderley, Alderley Park, Macclesfield, Cheshire SK10
4TG (GB). NOWAK, Thorsten [DE/GB]; AstraZeneca
R & D Alderley, Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB). PLOWRIGHT, Alleyn [GB/GB];
AstraZeneca R & D Alderley, Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB).

(21) No. Solicitud Internacional: PCT/GB2005/004726

(22) Fecha de Presentación Internacional:
9 de diciembre de 2005 (09.12.2005)

(74) Agente: GLOBAL INTELLECTUAL PROPERTY:
AstraZeneca AB, S-151 85 Södertälje (SE).

(25) Idioma de Presentación: Inglés

(26) Idioma de Publicación: Inglés

(30) Datos relativos a la Prioridad:
0427328.0 14 de diciembre de 2004 (14.12.2004) GB
0507403.4 13 de abril de 2005 (13.04.2005) GB
0520383.1 7 de octubre de 2005 (07.10.2005) GB

(81) Países designados (a menos que se indique lo
contrario para cada clase de protección disponible):
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW,
BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK,
MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH,
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,
ZA, ZM, ZW

(71) Solicitante (para todos los países destinatarios con
excepción de MG, US): ASTRAZENECA AB [SE/SE];
S-151 85 Södertälje (SE).

(84) Países designados (a menos que se indique lo
contrario para cada clase de protección disponible):
ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL,
SZ, TZ, UG, ZM, ZW), Eurasiática (AM, AZ, BY, KG,
KZ, MD, RU, TJ, TM), Europea (AT, BE, BG, CH, CY,
CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT,
LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR),
OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML,
MR, NE, SN, TD, TG).

(71) Solicitante (para MG únicamente): ASTRAZENECA UK
LIMITED [GB/GB], 15 Stanhope Gate, London, Greater,
London W1K 1LN (GB).


Publicado:
— con informe internacional de búsqueda

(72) Inventores, e

(75) Inventores/Solicitantes (para US únicamente): BIRCH,
Alan, Martin [GB/GB]; AstraZeneca R & D Alderley,
Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
BOWKER, Suzanne, Saxon [GB/GB]; AstraZeneca R &
D Alderley, Alderley Park, Macclesfield, Cheshire SK10
4TG (GB). BUTLIN, Roger, John [GB/GB];
AstraZeneca R & D Alderley, Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). DONALD,
Craig, Samuel [GB/GB]; AstraZeneca R & D Alderley.

Para códigos de dos letras y otras abreviaturas, remítase a
las "Notas Guía sobre Códigos y Abreviaturas" que aparece
al comienzo de cada edición regular del Boletín Oficial PCT.

(54) Título: DERIVADOS DE OXADIAZOL COMO INHIBIDORES DE DGAT



(I)

(57) Extracto: Se describen compuestos de fórmula (I), y sus sales y profármacos, en donde por ejemplo R¹ es
arilo o heteroarilo opcionalmente sustituido; Y es un grupo enlazante seleccionado por ejemplo de un enlace
directo, y una cadena alquilo (sustituido); R² es arilo opcionalmente sustituido, un cicloalquilo opcionalmente
sustituido o un grupo heterocíclico opcionalmente sustituido. Se describen asimismo procesos para elaborar
tales compuestos y su empleo como inhibidores de DGAT, por ejemplo, en el tratamiento de obesidad.

DERIVADOS DE OXADIAZOL COMO INHIBIDORES DE DGAT

La presente invención se refiere a compuestos que inhiben la actividad de la acetil CoA(acetil acetilcoenzima A):diacilglicerol aciltransferasa (DGAT1), a procesos para su preparación, a composiciones farmacéuticas que los contienen como ingrediente activo, a métodos para el tratamiento de estados patológicos asociados con actividad DGAT1, a su uso como medicamentos y a su empleo en la fabricación de medicamentos para uso en la inhibición de DGAT1 en animales de sangre caliente tales como humanos. En particular esta invención se refiere a compuestos útiles para el tratamiento de diabetes tipo II, resistencia a la insulina, tolerancia anormal a la glucosa y obesidad en animales de sangre caliente tales como humanos, más particularmente al empleo de estos compuestos en la fabricación de medicamentos para uso en el tratamiento de diabetes tipo II, resistencia a la insulina, tolerancia anormal a la glucosa y obesidad en animales de sangre caliente tales como humanos.

La acil CoA:diacilglicerol aciltransferasa (DGAT) se encuentra en la fracción microsómica de las células. Cataliza la reacción final en la vía del glicerol fosfato, considerada la vía principal de síntesis de triglicéridos en las células facilitando el enlace de un diacilglicerol con una acil CoA grasa, resultando en la formación de triglicérido. Aunque no es claro si la DGAT limita la tasa de síntesis de triglicéridos, cataliza el único paso en la vía que se dedica a producir este tipo de molécula [Lehner & Kuksis (1996) Biosynthesis of triacylglycerols. Prog. Lipid Res. 35: 169-201].

Se han clonado y caracterizado dos genes DGAT. Dos de las proteínas codificadas catalizan la misma reacción, aunque no comparten homología de secuencia. El gen DGAT1 se identificó a partir de búsquedas de bases de datos de secuencias, debido a su similitud con los genes de la acil CoA:colesterol aciltransferasa (ACAT). [Cases *et al* (1998) Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis. Proc. Natl. Acad.

Sci. USA 95: 13018-13023]. La actividad DGAT1 se ha detectado en muchos tejidos mamíferos, incluidos adipocitos.

Debido a la falta previa de sondas moleculares, poco se sabe sobre la regulación de DGAT1. DGAT1 es conocida por ser significativamente regulada por aumento durante la diferenciación adipocítica.

Estudios en ratones con genes inactivados ("*in gene knockout mice*") han indicado que moduladores de la actividad de DGAT1 serían útiles en el tratamiento de diabetes tipo II y obesidad. Ratones con el gen DGAT1 inactivado (*Dgat1*^{-/-}), son viables y capaces de sintetizar triglicéridos, como se evidenció por niveles normales de triglicéridos séricos en ayunas y composición normal del tejido adiposo. Los ratones *Dgat1*^{-/-} tienen menos tejido adiposo que los ratones naturales en la línea de referencia y son resistentes a obesidad inducida por la dieta. La tasa metabólica es ~20% más elevada en ratones *Dgat1*^{-/-} que en ratones naturales tanto con dietas regulares como con alto contenido de grasa [Smith *et al* (2000) Obesity resistance and multiple mechanisms of triglyceride synthesis in mice lacking DGAT. Nature Genetics 25:87-90]. Una actividad física creciente en ratones *Dgat1*^{-/-} explica parcialmente el mayor gasto de energía. En igual forma los ratones *Dgat1*^{-/-} exhiben un aumento de sensibilidad a la insulina y un incremento del 20% en la tasa de acumulación de glucosa. Los niveles de leptina se encuentran disminuidos en un 50% en los ratones *Dgat1*^{-/-} en línea con una reducción del 50% en la masa de grasa.


Si ratones *Dgat1*^{-/-} se cruzan con ratones *ob/ob*, estos ratones exhiben el fenotipo *ob/ob* [Chen *et al* (2002) Increased insulin and leptin sensitivity in mice lacking acyl CoA:diacylglycerol acyltransferase J. Clin. Invest. 109: 1049-1055] indicando que el fenotipo *Dgat1*^{-/-} requiere una vía leptina íntacta. Si ratones *Dgat1*^{-/-} se cruzan con ratones *Agouti* se observa una disminución en el peso corporal con niveles normales de glucosa y niveles de insulina reducidos en un 70% comparados con ratones *Dgat1*^{-/-} *ob/ob* o *agouti*, naturales.

El trasplante de tejido adiposo de ratones *Dgat1*^{-/-} a ratones naturales confiere resistencia a la obesidad, inducida por la dieta, y un mejor

metabolismo de la glucosa en estos ratones [Chen *et al* (2003) Obesity resistance and enhanced glucose metabolism in mice transplanted with white adipose tissue lacking acyl CoA:diacylglycerol acyltransferase J. Clin. Invest. 111: 1715-1722].

Las Solicitudes de Patente Internacional WO2004/047755 (Tularik and Japan Tobacco) y WO2005/013907 (Japan Tobacco and Amgen) describen heterociclos con contenido de nitrógeno, bicíclicos, fusionados, que son inhibidores de DGAT-1. JP2004-67635 (Otsuka Pharmaceuticals) describe compuestos fenilo tiazoleamido sustituido, los cuales se sustituyen además con alquilfosfonatos y que inhiben DGAT-1. WO2004/100881 (Bayer) describe compuestos bifenilamino sustituidos con imidazol, oxazol o tiazol, que inhiben DGAT-1.

Por consiguiente, la presente invención provee un compuesto de fórmula (I)



(I)

o su sal o profármaco farmacéuticamente aceptable, en donde:

R¹ es un grupo arilo opcionalmente sustituido o heteroarilo opcionalmente sustituido, en donde los sustituyentes opcionales son uno o más grupos seleccionados de un grupo -Z^a, un grupo -X²-(CR⁵²R⁵³)_w-Z^a, un grupo -X²-(CR⁵²R⁵³)_a-X³-Z^a, un grupo -(CR⁵²R⁵³)_a-X³-Z^a o un grupo funcional (el cual es distinto de un grupo -X²-(CR⁵²R⁵³)_w-Z^a o un grupo -X²-(CR⁵²R⁵³)_a-X³-Z^a);

Y es un enlace directo, o un grupo (CR⁴⁰R⁴¹)_s o -X⁶(CR⁴⁰R⁴¹)_t - donde cada R⁴⁰ y R⁴¹ se selecciona independientemente de hidrógeno, (1-4C)alquilo, hidroxí, halo, halo(1-4C)alquilo, amino, ciano, (1-4C)alcoxí, (1-4C)haloalcoxí o ((1-3C)alquil)CONH-, s es un entero de 1 a 6 y t es un entero de 1 a 6, siempre que el átomo X⁶ del grupo -X⁶(CR⁴⁰R⁴¹)_t- se una al grupo R² y que un átomo de carbono simple hibridado a sp³ no lleve dos o más enlaces a un heteroátomo a menos que el heteroátomo sea un halo;

R^2 es un grupo arilo opcionalmente sustituido, cicloalquilo opcionalmente sustituido o heterocíclico opcionalmente sustituido, en donde sustituyentes opcionales son uno o más grupos seleccionados de un grupo $-Z$, un grupo $-X-(CR^{42}R^{43})_u-Z$, un grupo $-X-(CR^{42}R^{43})_vX^1-Z$ o un grupo $-(CR^{42}R^{43})_vX^1-Z$ o un grupo funcional (el cual es distinto de un grupo $-X-(CR^{42}R^{43})_u-Z$ o un grupo $-X-(CR^{42}R^{43})_vX^1-Z$);

en donde Z y Z^a se seleccionan independientemente de un grupo hidrocarbilo o un grupo heterocíclico o una combinación de los mismos, en donde el grupo Z y Z^a se sustituye opcionalmente en cualquier átomo disponible por uno o más grupos funcionales, o por un grupo $-X^7-(CR^{62}R^{63})_bR^{64}$;

X , X^1 , X^2 , X^3 , X^6 y X^7 son grupos enlazantes seleccionados independientemente de $-C(O)_x-$, $-O-$, $-S(O)_y-$, $-NR^{44}-$, $-C(O)NR^{44}-$, $-OC(O)NR^{44}-$, $-CH=NO-$, $-NR^{44}C(O)_x-$, $-NR^{44}CONR^{45}-$, $-S(O)_2NR^{44}-$ o $-NR^{44}S(O)_2-$ donde x es un entero de 1 ó 2, y es 0, 1 ó 2, y R^{44} y R^{45} se seleccionan independientemente de hidrógeno o (1-6C)alquilo,

u y w se seleccionan independientemente de 0 o un entero de desde 1 a 6;

v , a y b se seleccionan independientemente de un entero de desde 1 a 6;

cada R^{42} , R^{43} , R^{52} , R^{53} , R^{62} y R^{63} se selecciona independientemente de hidrógeno, (1-4C)alquilo, hidroxilo, halo, halo(1-4C)alquilo, amino, ciano, (1-4C)alcoxi, (1-4C)haloalcoxi, ((1-3C)-alquil)CONH-, carboxi o réplica de ácido carboxílico o bioisótero del mismo, y R^{64} es un grupo funcional.

Tal como se emplea en esta solicitud, el término "grupo funcional" incluye halo, halo(1-6C)alquilo, ciano, nitro, $-C(O)_nR^{20}$, una réplica de ácido carboxílico o bioisótero del mismo, $-OR^{20}$, $-S(O)_mR^{20}$, $-OS(O)_2R^{20}$, $-NR^{21}R^{22}$, $-C(O)NR^{21}R^{22}$, $-OC(O)NR^{21}R^{22}$, $-CH=NOR^{20}$, $-NR^{21}C(O)_nR^{20}$, $-NR^{20}CONR^{21}R^{22}$, $-N=CR^{21}R^{22}$, $-S(O)_2NR^{21}R^{22}$ o $-NR^{21}S(O)_2R^{22}$ donde R^{20} , R^{21} y R^{22} se seleccionan independientemente de hidrógeno o hidrocarbilo opcionalmente sustituido o heterocíclico opcionalmente sustituido, o R^{21} y R^{22} , junto con el átomo de nitrógeno al cual se unen, forman un anillo opcionalmente sustituido con desde 3 a 10 átomos, el cual contiene

opcionalmente heteroátomos adicionales tales como $S(O)_m$ oxígeno y nitrógeno, n es un entero de 1 ó 2, m es 0 o un entero de 1-2.

Sustituyentes opcionales apropiados para grupos hidrocarbilo o grupos heterocíclicos R^{20} , R^{21} y R^{22} (inclusive para anillos formados por $NR^{21}R^{22}$) incluyen halo, halo(1-4C)alquilo (tales como trifluorometilo, difluorometilo o fluorometilo), mercapto, hidroxí, alcoxi, oxo, heteroariloxi, alqueniíloxi, alquiniíloxi, alcoxialcoxi, ariloxi (donde el grupo arilo puede sustituirse por halo, ciano, nitro, hidroxí(1-4C)alquilo, halo(1-4C)alquilo, amino, (1-4C)alcoxi, (1-4C)haloalcoxi, ((1-3C)alquil)CONH-, carboxi o una réplica de ácido carboxílico o bioisótero del mismo), ciano, nitro, amino, mono- o di-alquilamino, alquilamido, oximino (por ejemplo hidroxíimino o alquiloimino), carbamoilo, carboxi o una réplica de ácido carboxílico o bioisótero del mismo, o $-S(O)_mR^{23}$ donde m es como ya se definió y R^{23} es alquilo (opcionalmente sustituido por uno o más grupos seleccionados de hidroxí, halo, amino, ciano, ((1-3C)alquil)CONH-, carboxi o una réplica de ácido carboxílico o bioisótero del mismo), (1-6C)alcoxi, (1-6C)alcocarbonilo, carbamoilo, N-((1-6C)alquil)carbamoilo, halo(1-6C)alquilo (tal como trifluorometilo), (1-6C)alquilsulfonilo, (1-6C)alquilsulfinilo. Los grupos heterocíclicos R^{20} , R^{21} y R^{22} pueden sustituirse opcionalmente también por uno o más grupos hidrocarbilo tales como (1-4C)alquilo.

En esta especificación el término "alquilo" incluye los dos grupos alquilo de cadena recta y ramificada, pero las referencias a grupos alquilo individuales tales como "propilo" son específicos para la versión de cadena recta únicamente. Una convención análoga se aplica a otros términos genéricos. A menos que se indique lo contrario, el término "alquilo" se refiere ventajosamente a cadenas con 1-10 átomos de carbono, apropiadamente de 1-6 átomos de carbono, preferiblemente 1-4 átomos de carbono.

En esta especificación el término "alcoxi" significa un grupo alquilo como se definió anteriormente enlazado a un átomo de oxígeno.

Hay que entender que sustituyentes opcionales en cualquier grupo pueden unirse a cualquier átomo disponible, según sea apropiado, a menos que se indique lo contrario, incluidos heteroátomos, siempre que con ello no se cuaternicen.

En esta especificación el término "heteroátomo" se refiere a átomos diferentes de carbono tal como átomos de oxígeno, nitrógeno o azufre. Además, donde el heteroátomo puede tener una sola valencia, éste puede comprender un halo. Los términos "alquenilo" y "alquinilo" se refieren a estructuras rectas o ramificadas insaturadas las cuales, a menos que se especifique lo contrario, contienen por ejemplo desde 2 a 10, preferiblemente desde 2 a 6 átomos de carbono. Fracciones cíclicas tales como cicloalquilo y cicloalquenilo son similares en naturaleza pero tienen por lo menos 3 átomos de carbono. Ejemplos de grupos alquilo, alquenilo y cicloalquilo se dan a continuación, tales como ejemplos de (1-6C)alquilo, (3-8C)cicloalquilo, etc.

Referencias a grupos arilo incluyen grupos carbocíclicos aromáticos tales como fenilo y naftilo, así como grupos parcialmente aromáticos tales como indenilo e indanilo. El término "aralquilo" se refiere a grupos alquilo arilo sustituidos tales como bencilo.

El término "heterociclilo" o "heterocíclico" incluye anillos saturados o insaturados, los cuales pueden ser anillos aromáticos, no aromáticos o parcialmente aromáticos, por ejemplo que contengan desde 3 a 20, convenientemente desde 4 a 10 átomos de anillo, por lo menos uno de los cuales es un heteroátomo tal como oxígeno, azufre o nitrógeno. Pueden ser sistemas de anillo mono- o bicíclico, en donde uno o ambos anillos pueden ser saturados o insaturados, por ejemplo pueden ser aromáticos. En particular, los sistemas de anillo bicíclico comprenderán anillo fusionados de 5,6 miembros o de 6,6 miembros.

"Heteroarilo" se refiere a aquellos grupos heterocíclicos ya descritos que tienen un carácter aromático. Donde "heteroaril" es un sistema de anillo bicíclico, entonces por lo menos un anillo es aromático y uno o ambos anillos contienen heteroátomos de anillo.

En general, ejemplos de anillos heterociclilo incluyen furilo, tienilo, pirrolilo, pirrolidinilo, imidazolilo, triazolilo, tiazolilo, tetrazolilo, oxazolilo, isoxazolilo, pirazolilo, piridilo, pirimidinilo, pirazinilo, picridazinilo, triazinilo, indolilo, quinolinilo, isoquinolinilo, quinoxalinilo, benzotiazolilo, benzoxazolilo, benzotienilo y benzofurilo. Otros ejemplos incluyen tiadiazolilo, oxadiazolilo e isotiazolilo, morfolino, piperazinilo y piperidinilo. Otros ejemplos de anillos heteroarilo monocíclicos son los enumerados a continuación como ejemplos apropiados de HET-1^{a-f}. Otros ejemplos de anillos heterocíclicos bicíclicos son los enumerados a continuación como ejemplos apropiados de HET-2^{a-b}. Otros ejemplos de anillos heterocíclicos monocíclicos (no heteroarilo) son los enumerados a continuación como ejemplos apropiados de HET-3^{b,c,e}. Ejemplos apropiados de anillos heteroarilo bicíclicos incluyen indolilo, quinolinilo, isoquinolinilo, quinoxalinilo, benzotiazolilo, benzoxazolilo, benzotienilo, benzofurilo, benzimidazolilo, benzodioxolanilo, pirrolpiridilo, quinazolinilo y naftiridinilo. Se entenderá que estructuras tales como 2-oxo-2,3-dihidro-1H-benzimidazolilo y oxotiadiazolilo, que caen dentro de la definición del término heteroarilo, conservan sus características aromáticas en ambos anillos en virtud del tautomerismo.

Otras expresiones utilizadas en la especificación incluyen "hidrocarbilo", el cual se refiere a cualquier estructura que comprende átomos de carbono e hidrógeno. Éstos pueden disponerse en anillos o cadenas o combinaciones en las cuales los anillos se unen a cadenas o a otros anillos, o se fusionan a otros anillos. Por lo general, los grupos hidrocarbilo contendrán desde 1 a 20, por ejemplo desde 1-12 átomos de carbono. Éstos pueden ser alquilo, alquenilo, alquinilo, arilo, aralquilo, aralquenilo, aralquinilo, cicloalquilo o cicloalquenilo, en donde cualquier fracción cíclica tal como arilo, aralquilo, cicloalquilo o cicloalquenilo se sustituyen opcionalmente con alquilo, alquenilo, alquinilo y/o con otras fracciones cíclicas, y en donde cualesquiera grupos alquilo, alquenilo o alquinilo se sustituyen opcionalmente con cicloalquilo, o cicloalquenilo. El

término cicloalquilo incluye asimismo anillos bi- y tri-cicloalquilo, tales como adamantilo y biciclo[2.2.2]octanilo.

Combinaciones apropiadas de anillos y cadenas, que se abarcan por el término hidrocarbilo, incluyen

a) cicloalquilo enlazado a un grupo (1-6C)alquilo (en particular ciclohexilmetilo, ciclopentilmetilo, ciclobutilmetilo, ciclohexiletilo), o a dos grupos (1-6C)alquilo (por ejemplo metilciclobutilmetilo);

b) ciclohexilo enlazado a un segundo grupo ciclohexilo o ciclopentilo por un enlace directo, o con un grupo enlazante (1-6C)alquilo;

c) un grupo fenilo enlazado a un segundo grupo fenilo por un enlace directo, o con un grupo enlazante (1-6C)alquilo;

d) un grupo (3-8C)cicloalquilo (tal como ciclohexilo o ciclopentilo) enlazado a un grupo fenilo por un enlace directo o con un enlazante (1-6C)alquilo;

e) un grupo bencilo o metilfenilo (tal como toluilo).

Referencias a una "combinación" de grupos hidrocarbilo y heterocíclicos designan fracciones que contienen uno o más grupos heterocíclicos unidos entre sí, o uno o más grupos heterocíclicos unidos a uno o más grupos hidrocarbilo.

Combinaciones apropiadas de grupos hidrocarbilo y heterocíclicos incluyen un grupo heterociclilo (tal como pirídilo, morfolino, tiomorfolino, piperazinilo o piperidinilo) enlazado a (o sustituido por) un grupo hidrocarbilo (tal como un grupo (1-6C)alquilo y/o un grupo (3-8C)cicloalquilo; en particular un grupo (1-6C)alquilo. Por ejemplo metilpirídilo (en donde el metilo puede sustituirse adicionalmente por un grupo funcional tal como carboxi), bencilpiperazina, (metil)oxopiridazina, (metil)oxotiadiazol, (opcionalmente carboxi sustituido)metilpiperidilo, (opcionalmente carboxi sustituido)metilpiperidilmetilo, (opcionalmente carboxi sustituido)dimetilpiperidilo, (opcionalmente carboxi sustituido)etilpiperidilo y (ciclopropilmetil)piperazinilo.

A menos que se especifique lo contrario, la expresión "haloalquilo" se refiere a grupos alquilo que llevan por lo menos un sustituyente halo.

Esto incluye grupos perhalo donde todos los átomos de hidrógeno se reemplazan por halo, tal como fluoro.

Hay que entender que sustituyentes opcionales en cualquier grupo pueden unirse a cualquier átomo disponible, según sea apropiado, a menos que se especifique lo contrario, incluidos heteroátomos siempre que con ello no se cuaternicen.

Dentro de esta especificación, términos compuestos se utilizan para describir grupos que comprenden más de una funcionalidad tal como $-(1-6C)alquilNHSO_2(1-6C)alquilo$. Tales términos tienen que interpretarse de acuerdo con el significado que se entiende por una persona experta en el arte para cada parte componente. Por ejemplo $-(1-6)alquilNHSO_2(1-6C)alquilo$ incluye $-metilaminosulfonilmetilo$, $-metilaminosulfoniletilo$, $-etilaminosulfonilmetilo$, y $-propilaminosulfonilbutilo$.

Donde se seleccionan sustituyentes opcionales de "0, 1, 2 ó 3" grupos, hay que entender que esta definición incluye todos los sustituyentes que se escogen de uno de los grupos especificados o que los sustituyentes se escogen de dos o más de los grupos especificados. Una convención análoga se aplica a los sustituyentes seleccionados de "0, 1 ó 2" grupos y "1 ó 2" y cualesquiera otros grupos análogos.

Sustituyentes pueden estar presentes en cualquier posición adecuada en, por ejemplo, un grupo alquilo. Por lo tanto, hidroxí sustituido $(1-6C)alquilo$ incluye hidroxímetilo, 1-hidroxietilo, 2-hidroxietilo y 3-hidroxipropilo.

Ejemplos de $(1-4C)alquilo$ incluyen metilo, etilo, propilo e isopropilo; ejemplos de $(1-6C)alquilo$ incluyen metilo, etil, propilo, isopropilo, t-butilo, pentilo, iso-pentilo, 1-2-dimetilpropilo y hexilo; ejemplos de $(2-6C)alqueno$ incluyen etenilo, propenilo, isopropenilo, 2-pentenilo, 3-pentenilo, 4-pentenilo, 2-metilpropenilo y hexenilo; ejemplos de $(2-6C)alquinilo$ incluyen etinilo, propinilo, 2-pentinilo, 3-pentinilo, 4-pentinilo y hexinilo; ejemplos de $(1-4C)alcoxi$ incluyen metoxi, etoxi, propoxi, isopropoxi y *terc*-butoxi; ejemplos de $(1-6C)alcoxi$ incluyen metoxi, etoxi, propoxi, isopropoxi, *terc*-butoxi y pentoxi; ejemplos de $(1-$

6C)alcoxi(1-6C)alquilo incluyen metoximetilo, etoximetilo, metoxietilo, propoximetilo, isopropoximetilo, pentoxietilo, metoxihexilo y *terc*-butoxibutilo; ejemplos de **(3-8C)cicloalquil(3-6C)cicloalquilo** incluyen (ciclopropilo, ciclobutilo, ciclopentilo, ciclohexilo), cicloheptilo y ciclooctilo; ejemplos de **(3-8C)cicloalcoxi** incluyen ciclopropoxi, ciclobutoxi, ciclopentoxi, ciclohexiloxi, ciclopentiloxi y ciclooctiloxi; ejemplos de **(3-8C)cicloalquil(1-6C)alquilo** incluyen ciclopropilmetilo, ciclopropiletilo, ciclopropilbutilo, ciclobutilmetilo, ciclopentilmetilo, ciclohexilmetilo, ciclopentiletilo y ciclooctilpropilo; ejemplos de **(3-8C)cicloalcoxi(1-6C)alquilo** incluyen ciclopropoximetilo, ciclopropoxietilo, ciclopropoxibutilo, ciclobutoximetilo, ciclopentoximetilo, ciclohexiloximetilo, ciclopentoxietilo y ciclooctiloxipropilo; ejemplos de **(3-8C)cicloalcoxi(1-6C)alcoxi** incluyen ciclopropoximetoxi, ciclopropoxietoxi, ciclopropoxibutoxi, ciclobutoximetoxi, ciclopentoximetoxi, ciclohexiloximetoxi, ciclopentoxietoxi y ciclooctiloxipropoxi; ejemplos de **(3-8C)cicloalcoxi(1-6C)alcoxi(1-6C)alquilo** incluyen ciclopropoximetoximetilo, ciclopropoxietoximetilo, ciclopropoxibutoximetilo, ciclobutoximetoxietilo, ciclopentoximetoxipropilo, ciclohexiloximetoximetilo, ciclopentoxietoximetilo y ciclooctiloxipropoximetilo; ejemplos de **halo** son cloro, bromo, yodo y fluoro; ejemplos de **halo(1-6C)alquilo** incluyen **halo(1-4C)alquilo** tal como clorometilo, fluoroetilo, fluorometilo, fluoropropilo, fluorobutilo, diclorometilo, difluorometilo, 1,2-difluoroetilo y 1,1-difluoroetilo, así como perhalo(1-6C)alquilo (incluido perhalo(1-4C)alquilo) tal como trifluorometilo, pentafluoroetilo, y heptafluoropropilo; ejemplos de **halo(1-6C)alcoxi** incluyen **halo(1-4C)alcoxi** tal como clorometoxi, fluoroetoxi y fluorometoxi, difluorometoxi, así como perhaloalcoxi tal como pentafluoroetoxi, trifluorometoxi y heptafluoropropoxi; ejemplos de **hidroxi(1-6C)alquil** incluyen hidroxi(1-4C)alquilo tal como hidroxi metilo, 1-hidroxietilo, 2-hidroxietilo y 3-hidroxibutilo; ejemplos de **carboxi(1-6C)alquilo** incluyen carboxi(1-4C)alquilo, tal como carboximetilo, carboxietilo, carboxipropilo y

carboxibutilo; ejemplos de **amino(1-6C)alquilo** incluyen aminometilo, aminoetilo, 2-aminopropilo, 3-aminopropilo, 2-aminoiso-propilo, aminobutilo y 2-amino *terc*-butilo; ejemplos de **(1-6C)alquilamino** incluyen (1-4C)alquilamino tal como metilamino, etilamino y propilamino; ejemplos de **di-((1-6C)alquil)amino** incluyen di-(1-4C)alquilamino tal como dimetilamino, *N*-etil-*N*-metilamino, dietilamino, *N*-metil-*N*-propilamino y di-isopropilamino; ejemplos de **(1-6C)alquilcarbonilo** incluyen (1-4C)alquilcarbonilo tal como metilcarbonilo, etilcarbonilo, propilcarbonilo, iso-propilcarbonilo y *terc*-butilcarbonilo; ejemplos de **(1-6C)alquilcarboniloxi** incluyen (1-4C)alquilcarboniloxi tal como metilcarboniloxi, etilcarboniloxi, propilcarboniloxi, iso-propilcarboniloxi y *terc*-butilcarboniloxi; ejemplos de **(1-6C)alcoxycarbonil(N-(1-6C)alquilcarbamoilo)** incluyen (1-4C)alcoxycarbonilo tal como metoxycarbonilo, etoxycarbonilo, propoxycarbonilo, iso-propoxycarbonilo y *terc*-butoxycarbonilo; ejemplos de **(1-6C)alcoxycarbonilamino** incluyen (1-4C)alcoxycarbonilamino tal como metoxycarbonilamino, etoxycarbonilamino, propoxycarbonilamino, iso-propoxycarbonilamino y *terc*-butoxycarbonilamino; ejemplos de **(1-6C)alcoxycarbonil(N-metil)amino** incluyen (1-4C)alcoxycarbonil(N-metil)amino tal como metoxycarbonil(N-metil)amino, etoxycarbonil(N-metil)amino, propoxycarbonil (N-metil)amino, iso-propoxycarbonil(N-metil)amino y *terc*-butoxycarbonil(N-metil)amino; ejemplos de **(1-6C)alquiltio** incluyen metiltio, etiltio, propiltio, isopropiltio y butiltio; ejemplos de **(1-6C)alquilsulfinilo** incluyen metilsulfinilo, etilsulfinilo, propilsulfinilo, isopropilsulfinilo y butilsulfinilo; ejemplos de **(1-6C)alquilsulfonilo** incluyen metilsulfonilo, etilsulfonilo, propilsulfonilo, isopropilsulfonilo y butilsulfonilo; ejemplos de **(1-6C)alcoxisulfonilo** incluyen metoxisulfonilo, etoxisulfonilo, propoxisulfonilo, isopropoxisulfonilo y butoxisulfonilo; ejemplos de **(1-6C)alquilcarbonilamino** incluyen (1-4C)alquilcarbonilamino tal como (1-3C)alquilCONH (metilcarbonilamino, etilcarbonilamino, propilcarbonilamino, iso-propilcarbonilamino) y *terc*-butilcarbonilamino; ejemplos de **(1-6C)alquilaminocarbonilo** incluyen (1-

4C)alquilaminocarbonilo tal como metilaminocarbonilo, etilaminocarbonilo, propilaminocarbonilo, iso-propilaminocarbonilo y *terc*-butilaminocarbonilo; ejemplos de **di(1-6C)alquilaminocarbonilo** incluyen di(1-4C)alquilaminocarbonilo tal como dimetilaminocarbonilo, *N*-metil-*N*-etilaminocarbonilo, dietilaminocarbonilo, *N*-metil-*N*-propilaminocarbonilo y di-isopropilaminocarbonilo; ejemplos de **(1-6C)alquilaminocarboniloxi** incluyen (1-4C)alquilaminocarboniloxi tal como metilaminocarboniloxi, etilaminocarboniloxi, propilaminocarboniloxi, iso-propilaminocarboniloxi y *terc*-butilaminocarboniloxi; ejemplos de **-S(O)_p(1-4C)alquilo** (en donde *p* es 0, 1 ó 2) incluyen (1-6C)alquiltio, (1-6C)alquilsulfinilo y (1-6C)alquilsulfonilo; ejemplos de **(1-6C)alquilaminosulfonilo** incluyen -SO₂NH(1-4C)alquilo tal como metilaminosulfonilo, etilaminosulfonilo, propilaminosulfonilo, *iso*-propilaminosulfonilo y *terc*-butilaminosulfonilo; ejemplos de **di(1-6C)alquilaminosulfonilo** incluyen di(1-4C)alquilaminosulfonilo tal como dimetilaminosulfonilo, *N*-metil-*N*-etilaminosulfonilo, dietilaminosulfonilo, *N*-metil-*N*-propilaminosulfonilo y di-isopropilaminosulfonilo; ejemplos de **(1-6C)alquilsulfonilamino** incluyen (1-4C)alquilsulfonilamino tal como metilsulfonilamino, etilsulfonilamino, propilsulfonilamino, *iso*-propilsulfonilamino y *terc*-butilsulfonilamino.

Valores particulares de grupos variables en compuestos de fórmula (I) son los siguientes. Tales valores puede utilizarse donde se considere apropiado con cualquiera de los demás valores, definiciones, reivindicaciones o incorporaciones definidas anteriormente o más adelante.

En una incorporación particular, R¹ es un grupo arilo opcionalmente sustituido tal como fenilo o naftilo opcionalmente sustituido. R¹ como un grupo arilo opcionalmente sustituido puede ser también indanilo. Se entenderá que cuando R¹ es un grupo arilo parcialmente saturado, tal como indanilo, es la porción del anillo aromático de R¹ la que se une directamente al átomo de nitrógeno enlazante.

Alternativamente, R¹ es un grupo heteroarilo opcionalmente sustituido y, en particular, es un grupo heteroarilo monocíclico

opcionalmente sustituido tal como piridilo. Valores apropiados para R^1 , como un anillo heteroarilo, incluyen pirimidinilo, piridilo, pirazolilo, pirazinilo, tiazolilo, oxadiazolilo, isoxazolilo y tiadiazolilo.

Se entenderá que cuando R^1 es un grupo heteroarilo bicíclico parcialmente saturado, tal como benzodioxolanilo, es la porción del anillo aromático de R^1 la que se une directamente al átomo de nitrógeno enlazante.

Valores apropiados para R^1 , como un anillo heteroarilo bicíclico, incluyen pirrolopiridilo, benzodioxolanilo, benzotiazolilo, benzimidazolilo y quinolilo.

Valores más apropiados para R^1 incluyen fenilo, naftilo, indanilo, pirimidinilo, piridilo, pirazolilo, pirazinilo, tiazolilo, oxadiazolilo, isoxazolilo, tiadiazolilo, pirrolopiridilo, 1,3-benzodioxan-5-ilo, benzotiazolilo, benzimidazolilo y quinolilo.

En una incorporación, R^1 no puede ser pirrolo(1,2-b)piridazina.

Sustituyentes opcionales preferidos para R^1 incluyen grupos funcionales o grupos (1-6C)alquilo tal como metilo. Grupos funcionales particulares para sustituyentes en R^1 incluyen halo, $-C(O)_nR^{20}$ o $-OR^{20}$, donde R^{20} es como ya se definió y, en particular, es un grupo arilo o aralquilo.

Grupos funcionales apropiados como sustituyentes en R^1 incluyen halo, $-OR^{20}$ (en donde R^{20} es hidrógeno, fenilo o (1-4C)alquilo, sustituido opcionalmente por uno o más halo, de modo que por ejemplo R^{20} es difluorometilo o trifluorometilo, o sustituido opcionalmente por (1-4C)alcoxi), ciano, halo(1-4C)alquilo, $-S(O)_mR^{20}$ (en donde R^{20} es fenilo o (1-4C)alquilo, particularmente metilo o etilo, m es 0, 1 ó 2, particularmente 0 ó 2), trifluorometiltio, $-NR^{20}CONR^{21}R^{22}$ (en donde R^{20} , R^{21} y R^{22} son todos convenientemente hidrógeno), $-C(O)_nR^{20}$ (en donde n es 1 ó 2, particularmente 2 y R^{20} es (1-4C)alquilo o fenilo), $-OSO_2R^{20}$ (en donde R^{20} es convenientemente (1-4C)alquilo), $-SO_2NR^{21}R^{22}$ (en donde R^{21} y R^{22} son convenientemente ambos hidrógeno), $-NR^{21}C(O)_nR^{20}$ (en donde n es 1 ó 2, particularmente 1, R^{21} es convenientemente hidrógeno y R^{20} es

convenientemente fenilo o (1-4C)alquilo), y $-\text{CONR}^{21}\text{R}^{22}$ (en donde R^{21} y R^{22} son convenientemente hidrógeno).

Valores apropiados para Z^a incluyen fenilo (sustituido opcionalmente por un grupo funcional como se definió anteriormente, por ejemplo por $-\text{CO}_2\text{Me}$, o carboxi), bencilo, ciclohexilo, piridilo, pirimidinilo (sustituido opcionalmente por (1-4C)alquilo), triazolilo, morfolino, (2-4C)alquinilo (por ejemplo etinilo) y (1-4C)alquilo (sustituido opcionalmente por un sustituyente seleccionado de $-\text{CO}_2\text{Me}$, carboxi, metoxi, hidroxilo y ciano).

Donde R^1 se sustituye por un grupo $-\text{X}^2-(\text{CR}^{52}\text{R}^{53})_w-\text{Z}^a$, convenientemente w es 0 ó 1; Z^a se selecciona de los valores apropiados ya dados, particularmente hidrocarbilo (tal como alquilo opcionalmente sustituido, fenilo o bencilo) o piridilo, y es más convenientemente fenilo opcionalmente sustituido; X^2 es convenientemente $-\text{SO}_2-$, $-\text{CO}-$, $\text{NHCO}-$, $-\text{NH}-$, $-\text{O}-$, y R^{52} y R^{53} son convenientemente ambos hidrógeno.

En otro aspecto, sustituyentes opcionales preferidos en R^1 son 1, 2 ó 3 sustituyentes independientemente seleccionados de alquilo (por ejemplo (1-6C)alquilo tal como metilo o etilo, halo, haloalquilo (tal como halo(1-6C)alquilo, tal como halometilo, por ejemplo trifluorometilo), haloalcoxi (tal como halo(1-6C)alcoxi, tal como halometoxi, por ejemplo trifluorometoxi) y ciano.

En otro aspecto, sustituyentes opcionales preferidos en R^1 son 1, 2 ó 3 sustituyentes independientemente seleccionados de halo, halo(1-4C)alquilo, (1-4C)alcoxi, halo(1-4C)alcoxi, ciano, (1-4C)alquilo, (2-4C)alquinilo, $\text{Ph}(\text{CH}_2)_{0-1}\text{O}-$ (en donde el grupo fenilo se sustituye opcionalmente por halo), fenilo, benzoilo y anilino.

En otro aspecto, sustituyentes opcionales preferidos en R^1 son 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, bromo, trifluorometilo, metoxi, difluorometoxi, trifluorometoxi, ciano, metilo, etilo, etinilo, benciloxi, 3-clorobenciloxi, fenoxi, 4-clorofenoxi, fenilo, benzoilo y anilino.

En un aspecto, R^1 es mono-sustituido en la posición 3 en relación con el enlace a NH ; en otro aspecto R^1 es mono-sustituido en la posición 4. En

un aspecto adicional R^1 es 2,4-di-sustituido, 2,6-di-sustituido, 3,4-di-sustituido, 2,4-di-sustituido, o 2,5-di-sustituido por cualquiera de los posibles sustituyentes incluidos anteriormente o a partir de ahora, pero particularmente aquellos sustituyentes opcionales preferidos ya indicados y, más particularmente, di-halo, por ejemplo di-fluoro. En un aspecto adicional R^1 es tri-sustituido, por ejemplo 2,4,5-tri-sustituido, tal como 2,4,5-trihalo (por ejemplo 2,4,5-trifluoro).

Cuando R^1 es di- o tri-sustituido, convenientemente los sustituyentes se seleccionan independientemente de un grupo funcional, Z^a y $-X^2-(CR^{52}R^{53})_w-Z^a$, por ejemplo cualquiera de aquellos valores dados en esta solicitud para estos grupos. Por ejemplo, R^1 puede sustituirse por di-halo (tal como difluoro, dicloro, mono-fluoro mono-cloro y mono-cloro mono-bromo), tri-halo (tal como trifluoro), mono-halo mono-alquilo (tal como mono-metilo, mono-cloro), mono-halo (tal como fluoro o cloro) mono-trifluorometilo, mono-alquilo (tal como metilo) mono-ciano, di-metoxi, mono-cloro mono-metoxi, di-halo mono-hidroxilo (tal como 2-F, 4-Cl, 5-OH), o puede ser por ejemplo di-halo mono $-O-Z^a$ (tal como $-OCH_2CO_2Me$). Cuando R^1 es di-sustituido, en un aspecto por lo menos uno de los sustituyentes se selecciona de halo, (1-4C)alquilo, (1-4C)alcoxi, trifluorometilo y ciano. Cuando R^1 es tri-sustituido, en un aspecto por lo menos uno, por ejemplo por lo menos dos, de los sustituyentes se seleccionan de halo, (1-4C)alquilo, (1-4C)alcoxi, trifluorometilo y ciano.

Donde R^1 se sustituye por un grupo $-X^2-(CR^{52}R^{53})_w-Z^a$, un grupo $-X^2-(CR^{52}R^{53})_a-X^3-Z^a$ o un grupo $-(CR^{52}R^{53})_vX^3-Z^a$, R^{52} y R^{53} son convenientemente hidrógeno.

Cuando R^1 se sustituye por Z^a , en donde Z^a es un anillo heterociclilo, tal como un anillo morfolino, preferiblemente Z^a no se une al átomo de carbono de R^1 que está en una posición *orto* al enlace al átomo de nitrógeno enlazante.

En una incorporación, R^1 se selecciona de fenilo, naftilo, indanilo, pirimidinilo, piridilo, pirazolilo, pirazinilo, tiazolilo, oxadiazolilo, isoxazolilo, tiadiazolilo, pirrolopiridilo, 1,3-benzodioxan-5-ilo,

benztiazolilo, benzimidazolilo y quinolilo; opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de halo, (1-4C)alquilo, etinilo, (1-4C)alcoxi, hidroxilo, (1-4C)alcoxi(1-4C)alcoxi, metoximetilo, cianometilo, hidroxilo(1-4C)alquilo, trifluorometilo, difluorometoxi, trifluorometoxi, trifluorometiltio, ciano, metiltio, metilsulfonilo, metilsulfonilo, etilsulfonilo, aminocarbonilamino, metoxycarbonilamino, metilcarbonilamino, (1-4C)alcoxycarbonilo, metoxycarbonilmetoxi, benciloxi, piridilmetoxi, fenoxi (sustituido opcionalmente por metoxi o halo), fenilo (sustituido opcionalmente por metoxycarbonilo o carboxi), bencilo, anilino, anilincarbonilo, aminocarbonilo, benzoilo, benzoilamino, fenilsulfonilo, aminosulfonilo, ciclohexilo, metilpirimidinilo, triazolilo y morfolino.

En una incorporación preferida, Y es un enlace directo.

Donde Y es un grupo $-X^6(CR^{40}R^{41})_t$, X^6 es convenientemente oxígeno y t es preferiblemente un entero de desde 2 a 6.

Alternativamente, Y es un grupo $(CH_2)_s$ o más preferiblemente $-O(CH_2)_t$ - donde s es un entero de 1 a 6 y t es un entero de 2 a 6, y, en particular, s o t son 3.

Cuando R^2 es arilo no sustituido o cicloalquilo no sustituido, Y es preferiblemente distinto de un enlace directo.


R^2 es convenientemente un grupo fenilo sustituido o heteroarilo sustituido (por ejemplo cualquiera de aquellos grupos heteroarilo enumerados anteriormente). Ejemplos apropiados de R^2 incluyen fenilo, piridilo, pirimidinilo, indanilo, ciclohexilo, piperidinilo y benztiazolilo.

Cuando R^2 es un grupo cicloalquilo opcionalmente sustituido, éste es preferiblemente un grupo monocíclico tal como (3-8C)cicloalquilo o (3-6C)cicloalquilo.

Cuando R^2 es un grupo sustituido, éste es convenientemente sustituido por, como mínimo, uno y, opcionalmente, más de un grupo sustituyente -Z, un grupo $-X-(CR^{42}R^{43})_n-Z$, un grupo $-X-(CR^{42}R^{43})_n-X^1-Z$ o un grupo $-(CR^{42}R^{43})_n-X^1-Z$, donde uno o más sustituyentes adicionales pueden seleccionarse de halo, ciano, nitro, amino, hidroxilo o halo(1-

6C)alquilo. Preferiblemente R^2 se sustituye por 1 ó 2 grupos independientemente seleccionados de los definidos anteriormente o a partir de ahora, más preferiblemente por 1 grupo. Cuando R^2 se sustituye por 2 grupos, preferiblemente uno es un grupo funcional como se definió anteriormente, tal como halo, $-CO_2R^{20}$ (en donde R^{20} es hidrógeno, (1-4C)alquilo o alilo) o ciano, o un sustituyente es (1-4C)alquilo.


Ejemplos particulares de grupos Z o Z^a incluyen grupos de sub-fórmula (x), (y) o (z).




en donde cada anillo A o A' se selecciona independientemente de un anillo heterocíclico opcionalmente sustituido, un anillo cicloalquilo opcionalmente sustituido o un anillo arilo opcionalmente sustituido, cada R^{60} es un (1-6C)alquilo opcionalmente sustituido, un (2-6C)alquenilo opcionalmente sustituido o un (2-6C)alquinilo opcionalmente sustituido, y R^{61} es un (1-6C)alquilenio opcionalmente sustituido, un (2-6C)alquenileno opcionalmente sustituido o un (2-6C)alquinileno opcionalmente sustituido.

Convenientemente sustituyentes opcionales para grupos A, A', R^{60} y R^{61} son grupos funcionales.

Un ejemplo particular adicional de grupos Z o Z^a incluye grupos de sub-fórmula (zz), en donde A, R^{60} y R^{61} , y sustituyentes opcionales apropiados en ella son con se definen anteriormente para las sub-fórmulas (x), (y) y (z).



En una incorporación particularmente preferida, Z es un grupo de la subfórmula (x) ya indicada. En una incorporación, R² es un anillo aromático de 5 ó 6 miembros de la sub-estructura (a):



Z¹, Z², Z³ y Z⁴ se seleccionan independientemente de -CH-, -CR⁶- o un heteroátomo seleccionado de -O-, -S-, -N(R⁵⁰)_r-, donde r es 0 ó 1 dependiendo de los requerimientos del anillo aromático, y R⁵⁰ es hidrógeno o (1-6C)alquilo, y Z⁴ puede ser adicionalmente un enlace directo;

R⁴ es un grupo -Z, un grupo -X-(CR⁴²R⁴³)_u-Z, un grupo -X-(CR⁴²R⁴³)_v-X¹-Z o un grupo -(CR⁴²R⁴³)_v-X¹-Z, en donde Z, X, X¹, R⁴², R⁴³, u y v son como ya se definieron;

cada R⁶ se selecciona independientemente de halo, ciano, nitro, amino, hidroxí, haloC₁₋₆alquilo, un grupo -Z, un grupo -X-(CR⁴²R⁴³)_u-Z, un grupo -X-(CR⁴²R⁴³)_v-X¹-Z o un grupo -(CR⁴²R⁴³)_v-X¹-Z, en donde Z, X, X¹, R⁴², R⁴³, u y v son como ya se definieron.

Convenientemente, cuando Z es un enlace directo, uno de Z¹ o Z² es un heteroátomo, en particular oxígeno o azufre.

Preferiblemente Z⁴ es distinto de un enlace directo.

Convenientemente en este caso, Z² y Z³ se seleccionan independientemente de -CH-, -CR^{2a}- o un átomo de nitrógeno.

Convenientemente Z¹ es un grupo -CH-.


Grupos preferidos R⁶ se enumeran a continuación como R^{6a}.

Convenientemente, Z¹, Z², Z³ y Z⁴ son -CH-.

Convenientemente R⁴² y R⁴³ son hidrógeno

Donde uno de Z¹ a Z⁴ es N(R⁵⁰)_r, preferiblemente es Z² o Z³. Donde uno de Z¹ a Z⁴ es -CR⁶, preferiblemente es Z² o Z³.

En una incorporación alternativa, R² es un grupo cicloalquilo tal como ciclohexilo de la sub-fórmula (b)



donde R^4 es como ya se definió, y R^a , R^b , R^c y R^d se seleccionan independientemente de hidrógeno o un grupo R^6 como ya se definió.

En todavía otra incorporación, R^2 es un anillo bicíclico, el cual puede ser un anillo bicíclico o un anillo heterocíclico bicíclico. Por ejemplo, R^2 comprende anillos de 6,6 miembros, fusionados, o anillo de 5,6 miembros, fusionados, uno o los dos de dichos anillos pueden ser insaturados. Ejemplos de tales anillos incluyen benzimidazol (preferiblemente enlazado al grupo $-Y-NH-$ por medio del anillo benceno), indanilo, indenilo. Particularmente anillos bicíclicos apropiados son parcialmente insaturados, de modo que el anillo enlazado al grupo $-Y-NH-$ es saturado y éste se fusiona a un anillo aromático. Ejemplos particulares de tales anillos son anillos indanilo, tal como 2-indanilo. En una incorporación, R^2 no puede ser pirrolo(1,2-b)piridazina.

En particular, R^4 es un grupo Z.

Convenientemente Z es un grupo arilo, heterociclilo o cicloalquilo, cualquiera de los cuales se sustituye opcionalmente por un grupo funcional o un grupo (1-6C)alquilo, (2-6C)alquenilo o (2-6C)alquinilo.

Preferiblemente Z se sustituye por un grupo funcional o por un grupo (1-6C)alquilo, el cual se sustituye por un grupo funcional. Ejemplos particulares de tales grupos funcionales incluyen $-C(O)_2R^{20}$ o una réplica de ácido carboxílico o bioisótero del mismo, $-C(O)NR^{21}R^{22}$ y $-NR^{21}C(O)_nR^{20}$, donde R^{20} , R^{21} y R^{22} son como ya se definieron.

En otra incorporación, R^2 se sustituye por Z y Z es un grupo heterociclilo opcionalmente sustituido. Ejemplos apropiados de Z como un grupo heterociclilo opcionalmente sustituido incluyen cualquiera de los valores apropiados para heterociclilo dados anteriormente y, en particular, incluyen pírrolidinilo, piperazinilo, piperidinilo, piridilo, morfolino,

tiomorfolino, homopiperazinilo, tiadiazolilo, (oxo)piridazinilo y (oxo)tiadiazolilo.

En otra incorporación, R^2 se sustituye por Z y Z es un grupo hidrocarbilo opcionalmente sustituido. Ejemplos apropiados de Z como un grupo hidrocarbilo opcionalmente sustituido incluyen (todos opcionalmente sustituidos) (1-6C)alquilo (tal como (1-4C)alquilo), fenilo, cicloalquilo (tal como adamantilo, ciclobutilo, ciclopentilo y ciclohexilo), cicloalquilo combinado con (1-4C)alquilo (tal como metilciclohexilo, etilciclohexilo, isopropilciclohexilo, ciclohexilmetilo, etilciclobutilo, ciclobutilmetilo y metilciclopentilo) y fenilo combinado con (1-4C)alquilo (tal como bencilo y metilfenilo (tal como toliilo)).

En otra incorporación, R^2 se sustituye por Z y Z es una combinación opcionalmente sustituida de grupos hidrocarbilo y heterociclilo. Ejemplos apropiados de Z como una combinación opcionalmente sustituida de grupos hidrocarbilo y heterociclilo incluyen heterociclos no aromáticos tal como piperazinilo o piperidilo sustituido por (1-4C)alquilo (por ejemplo metilo, etilo e isopropilo), bencilo o cicloalquil(1-4C)alquilo (por ejemplo ciclopropilmetilo); heterociclos oxidados tal como oxopiridazina u oxotiadiazina sustituida por uno o dos (1-4C)alquilo (tal como metilo); heterociclos aromáticos (tal como piridilo) sustituido por uno o dos (1-4C)alquilo (tal como metilo). Por ejemplo, piridilmetilo (en donde el metilo puede sustituirse adicionalmente por un grupo funcional tal como carboxi), bencilpiperazinilo, (metil)oxopiridazinilo, (metil)oxotiadiazolilo, (opcionalmente carboxi sustituido)metilpiperidilo, (opcionalmente carboxi sustituido)metilpiperidilmetilo, (opcionalmente carboxi sustituido) dimetilpiperidilo, (opcionalmente carboxi sustituido)etilpiperidilo y (ciclopropilmetil)piperazinilo.

En otra incorporación R^2 se sustituye por Z y Z es una combinación opcionalmente sustituida de dos grupos heterociclilo, por ejemplo piridilo en combinación con piperazinilo.

Sustituyentes apropiados en un grupo Z incluyen halo, hidroxilo, carboxi, $-CO_nR^{20}$ [en donde R^{20} es hidrógeno, hidrocarbilo opcionalmente

sustituido (tal como (1-4C)alquilo, bencilo, fenilo, metilfenilo, fenetilo) o heterociclilo opcionalmente sustituido (tal como piridilo) y en donde n es 1 ó 2], $-\text{CONH}_2$, $-\text{CONHR}^{21}$ (en donde R^{21} se selecciona de hidrógeno, alquilo y bencilo), ciano, amino, $-\text{NHCO}_2(1-4\text{C})\text{alquilo}$, y $-\text{CONR}^{21}\text{R}^{22}$ (en donde $\text{NR}^{21}\text{R}^{22}$ forma un anillo heterociclilo opcionalmente sustituido).

Convenientemente un anillo formado por $\text{NR}^{21}\text{R}^{22}$ contiene 0 ó 1 heteroátomo adicional seleccionado de O, N y S y puede ser, por ejemplo, piperidinilo, piperazinilo, pirrolidinilo o morfolino. Un anillo formado por $\text{NR}^{21}\text{R}^{22}$ puede fusionarse también a otro anillo, por ejemplo comprender por ello un anillo pirrolidinilo fusionado con dioxolano.

En general, convenientemente R^{20} es hidrógeno o se selecciona de (todos opcionalmente sustituidos) (1-4C)alquilo, fenilo, piridilo, bencilo, fenetilo, metilfenilo y alilo.

En general, R^{21} y R^{22} son convenientemente cada uno independientemente hidrógeno o se seleccionan de (opcionalmente sustituido) fenilo, (1-4C)alquilo, y bencilo.

Convenientemente R^{20} , R^{21} y R^{22} (y anillos formados por $\text{NR}^{21}\text{R}^{22}$) son no sustituidos o se sustituyen por 1 ó 2 sustituyentes. Sustituyentes opcionales adecuados para R^{20} , R^{21} y R^{22} incluyen halo, ciano, hidroxí, (1-4C)alcoxi, carboxi y $-\text{CO}_2(1-4\text{C})\text{alquilo}$. Un sustituyente particular para R^{21} y R^{22} es hidroxí. Sustituyentes particulares para anillos formados por $\text{NR}^{21}\text{R}^{22}$ son hidroxí, carboxi y $-\text{CO}_2(1-4\text{C})\text{alquilo}$.

En otra incorporación R^2 se sustituye por $-\text{X}-(\text{CR}^{42}\text{R}^{43})_u\text{Z}$, en donde X es preferiblemente O, $-\text{NH}-$, $-\text{NMe}-$, o $-\text{SO}_2\text{NH}-$, u es 0, 1 ó 2, R^{42} y R^{43} son cada uno hidrógeno y Z se selecciona de cualquiera de los valores mencionados anteriormente, particularmente morfolino o fenilo opcionalmente sustituido (tal como metoxifenilo) o metilfenilo.

En otra incorporación, R^2 se sustituye únicamente por un grupo funcional como se definió anteriormente. En particular, el grupo funcional puede seleccionarse de (1-4C)alcoxi, (1-4C)alquiltio y (1-4C)alquilsulfonilo, en donde los grupos (1-4C)alquilo ya mencionados pueden sustituirse opcionalmente por carboxi o (1-4C)alcoxycarbonilo.

En un aspecto, R^2 -Y se selecciona de:


4-(piperazino)fenilo (sustituido en el nitrógeno de la piperazina disponible por un sustituyente seleccionado de (1-4C)alquilo, acilo, bencilo, etoxycarbonilo, *tert*-butoxicarbonilo, piridilo, ciclopropilmetilo y metoxietilo), 4-(piperazino)-3-metil-fenilo (sustituido en el nitrógeno de la piperazina disponible por un sustituyente seleccionado de acilo y *tert*-butoxicarbonilo),

4-(piperazino)piridilo (sustituido en el nitrógeno de la piperazina disponible por un sustituyente seleccionado de (1-4C)alquilo, acilo, etoxycarbonilo, *tert*-butoxicarbonilo, benzoilo, fluorobenzoilo, cianobenzoilo, metilbenzoilo, piridilcarbonilo, metilsulfonilo, metoximetilcarbonilo, fenetilcarbonilo, anilino-carbonilo, bencilaminocarbonilo, fenoxicarbonilo, benciloxicarbonilo y aminocarbonilo), 4-piperidinofenilo, benciloxifenilo, (1-4C)alquilfenilo, piridilmetoxifenilo, N-metilanilino-fenilo, bifenilo, (cloro)bifenilo, (metoxycarbonil)bifenilo, (etoxycarbonilmetil)bifenilo, (carboxi) bifenilo, morfolinofenilo, dimetilmorfolinofenilo, (morfolino) (cloro)fenilo, (morfolino)(dicloro)fenil(morfolino)(bromo)fenilo, (morfolino)(metil)fenilo, (morfolino)(fluoro)fenilo, (morfolino) (ciano)fenilo, (morfolino)(aliloxicarbonil)fenilo, (morfolinoetoxi)fenilo, morfolinopiridilo, (morfolino)(cloro)piridilo, (morfolino)(fluoro)piridilo, fenoxipiridilo, anilino-piridilo, metoxipiridilo, butoxipiridilo, piridiloxipiridilo, hidroxibutilpiridilo, tiomorfolinopiridilo, fenoxipropilo, (fluoro)(trifluorometil)fenilo, (N-acil)homopiperazinofenilo, (N-acil)piperazinopirimidinilo, metilbenzotiazolilo, butiltiofenilo, (metoxitiadiazolil)aminosulfonilfenilo, (difluorofenoxi)fenilo, (dimetil-oxo-piridazinil)fenilo, (metil-oxo-tiadiazinil)fenilo, metoxycarbonilpropilfenilo, (metoxycarbonil)metiltiofenilo, (metoxycarbonil)adamantilo, carboxiadamantilo, (metoxycarbonilmetil)bi-ciclohexilo, (carboxi)bi-ciclohexilo, (etoxycarboniletal)indanilo, (carboxietal)indanilo, (metoxycarbonil) fenoxipropilo, (carboxi)fenoxipropilo, (carboxipropil)fenilo, dicarboxibutilfenilo, carboxibutilfenilo, propilsulfonilfenilo, (carboxi)isopropilsulfonilfenilo, fenoxifenilo,


(metoxifenoxi)fenilo, metilfenoxifenilo, hidroxipropoxifenilo, etoxietoxifenilo, (dimetilaminocarbonil)isopropoxifenilo, (carboximetil)piperidinofenilo, (metoxicarbonilmetil)piperidinopiridilo, (carboximetil)piperidinopiridilo, (carboxi) piperidinopiridilo, (metoxicarbonil)piperidinopiridilo, (carboxi)(metil)piperidinopiridilo, (metoxicarbonil)(metil)piperidinopiridilo, (carboximetil)(metil)piperidinopiridilo, (metoxicarbonil)piperidil [(metil)piridilo], (carboxi)piperidil[(metil)piridilo], (metoxicarbonilmetil)piperidil [(metil)piridilo], (carboximetil)piperidil [(metil)piridilo], (carboxietil)piperidil[(metil)piridilo], (etoxi carboniletal)piperidil[(metil)piridilo], [(metoxicarbonil)piperidinocarbonilmetil]ciclohexilfenilo, [(carboxi)piperidinocarbonilmetil]ciclohexilfenilo, [(metoxicarbonil)isopropilaminocarbonilmetil]ciclohexilfenilo, [(carboxi)isopropilaminocarbonilmetil]ciclohexilfenilo, [(metoxicarbonil)pirrolidinilcarbonilmetil]ciclohexilfenilo, [(carboxi)pirrolidinilcarbonilmetil]ciclohexilfenilo, [(metoxicarbonil)(hidroxi)pirrolidinilcarbonilmetil]ciclohexilfenilo, [(carboxi)(hidroxi)pirrolidinilcarbonilmetil]ciclohexilfenilo, (hidroxiisobutilaminocarbonilmetil)ciclohexilfenilo, [(tetrahidrodioxolopirrolil)carbonil]metilciclohexilfenilo, (metoxicarbonilpiridil)piperidino, (carboxipiridil)piperidino, (aminocarbonilpiridil)piperidino, (carboximetilpiridil)piperidino, (etoxicarbonilmetil)ciclohexilfenilo, (metoxicarbonilmetil)ciclohexilfenilo, ciclohexilfenilo, (carboximetil)ciclohexilfenilo, (hidroxietil)ciclohexilfenilo, (aminocarbonilmetil)ciclohexilfenilo, (cianometil)ciclohexilfenilo, carboxiciclohexilfenilo, (dimetilaminocarbonilmetil)ciclohexilfenilo, [(N-hidroxietil-N-metilaminocarbonilmetil]ciclohexilfenilo, [N-(dihidroxipropil) aminocarbonilmetil]ciclohexilfenilo, (aminometil)ciclohexilfenilo, (terc-butoxicarbonilaminometil)ciclohexilfenilo, (metoxicarbonil) ciclohexiloxifenilo, (etoxicarbonil)ciclohexiloxifenilo,

carboxiciclohexiloxifenilo, (metoxycarbonil)ciclopentilfenilo,
 (etoxycarbonilmetil)ciclopentilfenilo,
 (carboxi)ciclopentilfenil(carboximetil)ciclopentilfenilo,
 (metoxycarbonil)ciclobutilfenilo, (etoxycarbonil)ciclobutilmetilfenilo,
 (carboxi)ciclobutilfenilo, (carboxi)ciclobutilmetilfenilo,
 (carboxietil)ciclobutilfenilo, (etoxycarbonietil)ciclobutilfenilo,
 (carboximetil)fenilciclohexilo, 1-(etoxycarbonil)etilciclohexilfenilo, 1-
 carboxietilciclohexilfenilo, (2-metoxycarbonil)isopropi-2-ilciclohexilo, (2-
 carboxi)isoprop-2-ilciclohexilo, pirrolidinilfenilo,
 (metoxycarbonilmetil)ciclohexilpiridilo y (carboximetil)ciclohexilpiridilo.


Tal como se emplea en esta solicitud, la referencia a una réplica de ácido carboxílico o bioisótero incluye grupos como se definen en The Practice of Medicinal Chemistry, Wermuth CG. Ed.: Academic Press: New York, 1996, p203. Ejemplos particulares de tales grupos incluyen $-SO_3H$, $S(O)_2NHR^{13}$, $S(O)_2NHC(O)R^{13}$, $-CH_2S(O)_2R^{13}$, $-C(O)NHS(O)_2R^{13}$, $-C(O)NHOH$, $-C(O)NHCN$, $-CH(CF_3)OH$, $C(CF_3)_2OH$, $-P(O)(OH)_2$ y grupos de la sub-fórmula (a)-(i') como se expone a continuación:




(a)




(b)




(c)




(d)




(e)




(f)




(g)




(h)




(i)




(j)




(k)




(l)




(m)




(n)




(o)




(p)




(q)




(r)




(s)




(t)




(u)




(v)




(w)




(x)




(y)




(z)




(a')




(b')




(c')



(d')



(e)




donde p es 1 ó 2, R^{27} y R^{28} se seleccionan independientemente de hidrógeno, hidroxilo, (1-6C)alcoxi, tior, (1-6C)alquiltio, $-C(O)R^{29}$, $-S(O)R^{30}$, $-SO_2R^{31}$, $-NR^{32}R^{33}$, $-NHCN$, halógeno y trihalometilo, donde R^{29} , R^{30} y R^{31} son $-OR^{34}$, (1-6C)alquilo, $-NR^{32}R^{33}$ o trihalometilo,

R^{32} y R^{33} se seleccionan independientemente de hidrógeno, (1-6C)alquilo, $-SO_2R^{34}$ y $-COR^{35}$, donde R^{35} es (1-6C)alquilo o trihalometilo, y R^{34} es hidrógeno, (1-6C)alquilo o trihalometilo.

Ejemplos particulares de R^{27} o R^{28} son hidroxilo.

En un aspecto adicional de la invención, se provee un compuesto de fórmula (IZA), o su sal o profármaco farmacéuticamente aceptable,



(IZA)

en donde R^1 se selecciona de fenilo (opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, bromo, trifluorometilo, metoxi, difluorometoxi, trifluorometoxi, ciano, metilo, etilo, etinilo, benciloxi, 3-clorobenciloxi, fenoxi, 4-clorofenoxi, fenilo, benzoilo y anilino), 2-piridilo (sustituido opcionalmente por clorofenoxi, clorobenciloxi o metoxifenoxi, y/o sustituido con un sustituyente seleccionado de halo, trifluorometilo, (1-4C)alquilo, (1-4C)alcoxi y ciano), 3-piridilo (opcionalmente sustituido como para 2-piridilo), halopirimidinilo y trifluorometiltiazolilo;

Z^2 es N o CH;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno o metilo;

R^{6ZA} es hidrógeno, fluoro, cloro o metilo;

A es N o CH;

X^{ZA} es un enlace directo, $-CH_2-$ o $-O-$ (excepto cuando A es N);

m es 0, 1 ó 2;

n es 0 ó 1;

siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En otro aspecto de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido como ya se definió.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es CH;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es CH;

X^{ZA} es un enlace directo o $-CH_2-$;

m es 0, 1 ó 2;

n es 0 ó 1;

siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es N;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es CH;

X^{ZA} es un enlace directo o $-CH_2-$;

m es 0, 1 ó 2;

n es 0 ó 1;

siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es CH;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es N;

X^{ZA} es un enlace directo o $-CH_2-$;

m es 0, 1 ó 2;

n es 0 ó 1;

siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es N;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es N;

X^{ZA} es un enlace directo o $-CH_2-$;

m es 0; 1 ó 2;

n es 0 ó 1;

siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es N;

R^{ZA3} es metilo;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es N;

X^{ZA} es un enlace directo o $-CH_2-$;

m es 1;

n es 1;

p es 0.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es CH;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es CH;

X^{ZA} es $-O-$;

m es 0, 1 ó 2;

n es 0 ó 1;

siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZA), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo;

Z^2 es N;

R^{ZA1} y R^{ZA2} son cada uno independientemente hidrógeno o metilo;

R^{ZA3} es hidrógeno;

R^{6ZA} es hidrógeno, cloro, fluoro o metilo, particularmente hidrógeno;

A es CH;

X^{ZA} es -O-;

m es 0, 1 ó 2;

n es 0 ó 1;


siempre que $m + n = 0, 1$ ó 2;

p es 0 ó 1.

En un aspecto, para un compuesto de fórmula (IZA), en donde A es -CH-, los sustituyentes en el anillo que contiene A (es decir el grupo X^{ZA} -piridilo/fenilo y el grupo carboxi(alquilo)) son *trans* en relación unos con otros.

En un aspecto, para un compuesto de fórmula (IZA), en donde A es -CH-, los sustituyentes en el anillo que contiene A (es decir, el grupo X^{ZA} -piridilo/fenilo y el grupo carboxi(alquilo)) son *cis* en relación unos con otros.

En un aspecto adicional de la invención, se provee un compuesto de fórmula (IZB), o su sal o profármaco farmacéuticamente aceptable,



(IZB)

en donde R^1 se selecciona de fenilo (opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, bromo, trifluorometilo, metoxi, difluorometoxi, trifluorometoxi, ciano, metilo, etilo, etinilo, benciloxi, 3-clorobenciloxi, fenoxi, 4-clorofenoxi, fenilo, benzoilo y anilino), 2-piridilo (sustituido opcionalmente por clorofenoxi, clorobenciloxi o metoxifenoxi, y/o sustituido con un sustituyente seleccionado de halo, trifluorometilo, (1-4C)alquilo, (1-4C)alcoxi y ciano), 3-piridilo (opcionalmente sustituido como para 2-piridilo), halopirimidinilo y trifluorometiltiazolilo;

Z^2 es N o CH;


R^{6ZB} es hidrógeno, fluoro, cloro o metilo;

X^{ZB} es O o S, particularmente O.

En otro aspecto de esta incorporación, en el compuesto de fórmula (IZB), R^1 es fenilo, opcionalmente sustituido como ya se definió.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZB), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo.

En un aspecto adicional de la invención, se provee un compuesto de fórmula (IZC), o su sal o profármaco farmacéuticamente aceptable,



(IZC)

en donde R^1 se selecciona de fenilo (opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, bromo, trifluorometilo, metoxi, difluorometoxi, trifluorometoxi, ciano, metilo, etilo, etinilo, benciloxi, 3-clorobenciloxi, fenoxi, 4-clorofenoxi, fenilo, benzoilo y anilino), 2-piridilo (sustituido opcionalmente por clorofenoxi, clorobenciloxi o metoxifenoxi, y/o sustituido con un sustituyente

seleccionado de halo, trifluorometilo, (1-4C)alquilo, (1-4C)alcoxi y ciano), 3-piridilo (opcionalmente sustituido como para 2-piridilo), halopirimidinilo y trifluorometiltiazolilo;

Z^2 es N o CH;


R^{6Zc} es hidrógeno, fluoro, cloro o metilo;

R^{Zc} se selecciona de (1-4C)alquilo, acilo, bencilo, etoxycarbonilo, terc-butoxycarbonilo, piridilo, ciclopropilmetilo, metoxietilo, benzoilo, fluorobenzoilo, cianobenzoilo, metilbenzoilo, piridilcarbonilo, metilsulfonilo, metoximetilcarbonilo, fenetilcarbonilo, anilincarbonilo, bencilaminocarbonilo, fenoxycarbonilo, benciloxycarbonilo y aminocarbonilo.

En otro aspecto de esta incorporación, en el compuesto de fórmula (IZC), R^1 es fenilo, opcionalmente sustituido como ya se definió.

En un aspecto adicional de esta incorporación, en el compuesto de fórmula (IZC), R^1 es fenilo, opcionalmente sustituido con 1, 2 ó 3 sustituyentes independientemente seleccionados de fluoro, cloro, trifluorometilo, trifluorometoxi, ciano, metilo y etilo.

En particular, la presente invención provee un compuesto de fórmula (IA) o su sal o profármaco farmacéuticamente aceptable,



en donde:

R^{1a} se selecciona de AR^{1a} , HET-1^a y HET-2^a; en donde R^{1a} se sustituye opcionalmente por 1, 2, 3 ó 4 grupos independientemente seleccionados de R^3 ;

R^3 se selecciona de halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $R^{8a}NHCO-$, $R^{8a}CONH-$, $R^{8a}OCONH-$, $R^{8a}OCON(Me)-$, $R^{8a}(1-6C)$ alquil-, halo(1-6C)alquilo, halo(1-6C)alcoxi, (1-6C)alquiltio, di(1-6C)alquilaminocarbonilo, (3-8C)cicloalquiloxi(1-6C)alquilo, (3-

8C)cicloalquil(1-6C)alcoxi, (3-8C)cicloalquil(1-6C)alcoxi(1-6C)alquilo, fenilo, HET-1^b, AR1^b-(1-6C)alcoxi, HET-1^b-(1-6C)alcoxi, HET-3^b-(1-6C)alcoxi, AR1^b-oxi, HET-1^b-oxi, HET-3^b-oxi, AR1^b-oxi(1-6C)alquilo, HET-1^b-oxi(1-6C)alquilo, HET-3^b-oxi(1-6C)alquilo, AR1^b-(1-6C)alcoxi(1-6C)alquilo, HET-1^b-(1-6C)alcoxi(1-6C)alquilo, y HET-3^b-(1-6C)alcoxi(1-6C)alquilo;


en donde cualquier fenilo, AR1^b, HET-1^b, HET-3^b o anillo cicloalquilo en cualquiera de los valores anteriores para R³ puede sustituirse opcionalmente por hasta 4 sustituyentes independientemente seleccionados de R^{7a};

y en donde cualquier R^{8a}, grupo (3-8C)cicloalquilo, o (1-6C)alquilo en cualquiera de las definiciones anteriores para R³ puede sustituirse opcionalmente por 1, 2 ó 3 sustituyentes independientemente seleccionados de halo, (1-6C)alquilo, (1-6C)alcoxi, (2-6C)alquenilo, (2-6C)alquinilo, hidroxilo, ciano, carboxi, aminocarbonilo, haloC₁₋₆alquilo, amino, (1-6C)alquilamino, di(1-6C)alquilamino, (1-6C)alquiltio, (1-6C)alquilsulfinilo, (1-6C)alquilsulfonilo, aminosulfonilo, (1-6C)alquilaminosulfonilo, di(1-6C)alquilaminosulfonilo, (1-6C)alquilsulfonilamino, aminocarbonilo, (1-6C)alquilaminocarbonilo, di(1-6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alcoxycarbonilamino, (1-6C)alcoxycarbonil(N-metil)amino y (1-6C)alquilaminocarboniloxi;

Y es un enlace directo, o un grupo (CH₂)_s o -O(CH₂)_t - donde s es un entero de 1 a 6 y t es un entero de 2 a 6, siempre que el átomo de oxígeno del grupo -O(CH₂)_t se una al grupo R²;


R^{2a} es:

a) un anillo de 6 miembros seleccionado de (i), (ii) o (iii)



en donde R^{36a}, R^{36b}, R^{36c} y R^{36d} se seleccionan independientemente de hidrógeno o un grupo R^{6a} como se define a partir de ahora; o

b) un anillo heteroarilo de 5 miembros de la sub-fórmula (iv)



en donde Z^5 , Z^6 y Z^7 se seleccionan independientemente de un heteroátomo (O, N o S) o CR^{37} , donde cada R^{37} es hidrógeno o un grupo R^{6b} como se define a partir de ahora, y siempre que por lo menos uno de Z^5 , Z^6 y Z^7 sea un heteroátomo y no haya enlaces O-O, S-S u O-S); o

c) un grupo HET-2^b el cual se sustituye opcionalmente por 1 ó 2 grupos independientemente seleccionados de R^4 o R^6 ;

d) un grupo cicloalquilo opcionalmente sustituido o grupo arilo bicíclico opcionalmente sustituido (tal como indanilo) en donde sustituyentes opcionales se seleccionan de grupos R^4 y R^{6a} como se definen a partir de ahora;

R^4 se selecciona de:

a) fenilo;

b) HET-3^c (en donde cualquier átomo de nitrógeno disponible se sustituye por R^5 siempre que, con ello, no se cuaternice;

c) HET-1^c;

d) fenoxi, HET-1^c-oxi, $R^{8b}O-$, $R^{8b}S(O)_q-$, fenil(1-6C)alcoxi y HET-1^c-(1-6C)alcoxi, donde q es 0, 1 ó 2;

e) R^{8b} , $R^{8b}O(1-6C)$ alquilo, amino(1-6C)alquilo, hidroxí(1-6C)alquilo, fenoxí(1-6C)alquilo, HET-1^c-oxi(1-6C)alquil; y

f) fenilamino, fenil(1-6C)alquilamino, HET-1^c-amino, HET-1^c-(1-6C)alquilamino, (1-6C)alquilamino, N-fenil-N-(1-6C)alquilamino, N-fenil(1-6C)alquil-N-(1-6C)alquilamino, N-(HET-1^c)-N-(1-6C)alquilamino, N-(HET-1^c-(1-6C)alquil)-N-(1-6C)alquilamino, y di(1-6C)alquilamino;

en donde cualquier alquilo, R^{8b} , fenilo (incluido fenoxi), anillo HET-1^c o HET-3^c en cualquiera de las definiciones a) a f) para R^4 puede sustituirse

opcionalmente por 1, 2, 3 ó 4 sustituyentes independientemente seleccionados de R^{7b} ;

R^5 se selecciona de un grupo R^{8c} , fenilo, HET-1^d, $R^{8c}CO-$, $R^{8c}OC(O)-$, fenilcarbonilo, HET-1^d-carbonilo, (1-6C)alcoxicarbonilo, H_2NCO , $R^{8b}NHCO-$, fenilNHCO-, (1-6C)alcoxisulfonilo, (1-6C)alquilaminosulfonilo, (1-6C)alquilsulfinilo y (1-6C)alquilsulfonilo;

en donde cualesquiera grupos (1-6C)alquilo o (3-8C)cicloalquilo en cualquiera de las definiciones de R^5 , incluido R^{8c} , puede sustituirse opcionalmente por un sustituyente seleccionado de un grupo R^{15} , amino, (1-6C)alquilamino, hidroxilo, carboxi, (2-6C)alquenilo, (2-6C)alquinilo, fenilo (opcionalmente sustituido con hasta cuatro grupos seleccionados de R^{15a} y HET-1^f;

y en donde cualesquiera grupos fenilo o HET-1^d en cualquiera de las definiciones de R^5 se sustituyen opcionalmente por hasta cuatro grupos seleccionados de R^{15b} ,

donde R^{15} , R^{15a} y R^{15b} se seleccionan independientemente de halo, ciano, C_{1-6} alcoxi, halo C_{1-6} alquilo, halo C_{1-6} alcoxi, (1-6C)alquilS(O)p' (en donde p' es 0, 1 ó 2),

R^{6a} y R^{6b} se seleccionan independientemente de R^{8d} , $R^{8d}O-$, $R^{8d}OCO-$, $R^{8d}COO-$, halo, ciano, halo(1-6C)alquilo, (1-6C)alcoxi(1-6C)alquilo, (1-6C)alcoxi(1-6C)alcoxi, (1-6C)alquilcarbonilo, (1-6C)alquilS(O)p (en donde p es 0, 1 ó 2), (1-6C)alquilaminocarbonilo, di(1-6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alquilaminosulfonilo, di(1-6C)alquilaminosulfonilo, (1-6C)alquilsulfonilamino, $AR1^c$ -(1-6C)alcoxi, HET-1^e-(1-6C)alcoxi, HET-3^e-(1-6C)alcoxi, $AR1^c$ -oxi(1-6C)alquilo, HET-1^e-oxi(1-6C)alquilo, HET-3^e-oxi(1-6C)alquilo, $AR1^e$ -(1-6C)alcoxi(1-6C)alquilo, HET-1^e-(1-6C)alcoxi(1-6C)alquilo, y HET-3^e-(1-6C)alcoxi(1-6C)alquilo; en donde cualquier alquilo, R^{8d} , fenilo, $AR1^c$, HET-1^e, HET-3^e o anillo cicloalquilo en cualquiera de los valores anteriores para R^{6a} o R^{6b} puede sustituirse opcionalmente por 1, 2 ó 3 sustituyentes independientemente seleccionados de R^{7c} ,

cada uno de R^{7a} , R^{7b} y R^{7c} se seleccionan independientemente de R^{8e} ,
 $R^{8e}O(CR^{13}R^{14})_{n^-}$, $R^{8e}OCO(CR^{13}R^{14})_{n^-}$, $R^{8e}COO(CR^{13}R^{14})_{n^-}$,
 $HCOO(CR^{13}R^{14})_{n^-}$, $R^{8e}CO(CR^{13}R^{14})_{n^-}$, $HCO(CR^{13}R^{14})_{n^-}$,
 $R^{8e}NHCO(CR^{13}R^{14})_{n^-}$, $H_2NCO(CR^{13}R^{14})_{n^-}$, $(R^{10}R^{11})NCO(CH_2)_{n^-}$,
 $R^{8e}CONH(CR^{13}R^{14})_{n^-}$, $HCONH(CR^{13}R^{14})_{n^-}$, $HCONR^{8e}(CR^{13}R^{14})_{n^-}$,
 $R^{8e}CONR^{8f}(CR^{13}R^{14})_{n^-}$, $R^{8e}OCONH(CR^{13}R^{14})_{n^-}$, $HCONR^{8e}(CR^{13}R^{14})_{n^-}$,
 $R^{8e}OCON(R^{8f})(CR^{13}R^{14})_{n^-}$, $ROCON(H)(CR^{13}R^{14})_{n^-}$,
 $HOCON(R^{8e})(CR^{13}R^{14})_{n^-}$, $R^{8e}(1-6C)alquilo$, halo, ciano, R^{12} , $R^{12}(1-6C)alquilo$, hidroxí, oxo, aminocarbonilo, halo(1-6C)alquilo, halo(1-6C)alcoxi, $R^{8e}S(O)p^{\prime}$ (en donde $p^{\prime}$ es 0, 1 ó 2), amino, amino(1-6C)alquilo, $R^{8e}NH(1-6C)alquilo$, $(R^{8e}R^{8f})N(1-6C)alquilo$, $R^{8e}NHSO_2-$, $(R^{10}R^{11})NSO_2-$, y $R^{8e}SO_2NH-$, donde $n^{\prime}$ es 0 o un entero de desde 1 a 6;

R^{12} es carboxi o una réplica de ácido carboxílico o bioisótero de carboxi;

R^{13} se selecciona de hidrógeno, (1-6C)alquilo, hidroxí, halo, amino, ciano, ((1-3C)alquil)CONH-, carboxi, (1-6C)alcoxi, (1-6C)alcoxycarbonilo, carbamoilo, N-((1-6C)alquil)carbamoilo, halo((1-6C)alquilo) (tal como trifluorometilo), (1-6C)alquilsulfonilo o (1-6C)alquilsulfimilo;

R^{14} es hidrógeno, (1-6C)alquilo o halo(1-6C)alquilo;

y en donde cualquier grupo alquilo o grupo R^{8e} , R^{8f} , R^{10} o R^{11} en un grupo R^{7a} , R^{7b} o R^{7c} se sustituye opcionalmente por 1, 2 ó 3 grupos independientemente seleccionados de R^9 ;

cada R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} y R^{8f} se selecciona independientemente de (1-6C)alquilo, (2-6C)alquenilo, (2-6C)alquinilo y (3-8C)cicloalquilo;

R^{10} y R^{11} se seleccionan independientemente (1-6C)alquilo, (2-6C)alquenilo, (2-6C)alquinilo y (3-8C)cicloalquilo; o R^{10} y R^{11} , junto con el átomo de nitrógeno al cual se unen, forman un anillo heterocíclico de a 4-7 miembros, el cual contiene opcionalmente uno o más heteroátomos adicionales seleccionados de O, N y S, y que se sustituye opcionalmente por desde 1, 2 ó 3 grupos seleccionados de hidroxí, (1-6C)alquilo, (1-6C)haloalquilo, carboxi una réplica de ácido carboxílico o bioisótero del mismo;

R⁹ se selecciona de hidroxilo, halo, amino, ciano, ((1-3C)alquil)CONH-, carboxi, (1-6C)alcoxi, (1-6C)alcoxycarbonilo, carbamoilo, N-((1-6C)alquil)carbamoilo, halo((1-6C)alquil) (tal como trifluorometilo), (1-6C)alquilsulfonilo, (1-6C)alquilsulfinilo,


AR1^a, AR1^b y AR1^c se seleccionan independientemente de fenilo o un sistema de anillo fusionado 6,5-, o 6,6-bicíclico, en donde los dos anillos son carbocíclicos y en donde por lo menos el anillo unido al átomo de nitrógeno enlazante es completamente aromático;

HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e y HET-1^f se seleccionan independientemente de un anillo heteroarilo de a 5 ó 6 miembros que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO₂, y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo;

HET-2^a y HET-2^b se seleccionan independientemente de un sistema de anillo heterocíclico, aromático o parcialmente insaturado, bicíclico 5,5, 5,6, 6,5 o 6,6 fusionado, que contiene 1, 2, 3 ó 4 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en los cuales el anillo directamente unido al átomo de nitrógeno enlazante es aromático, y cualquier anillo del sistema de anillo bicíclico puede ser carbocíclico;

HET-3^b, HET-3^c, HET-3^e se seleccionan independientemente de un anillo heterocíclico de 4, 5, 6 ó 7 miembros, parcial o completamente insaturado, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO₂, y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo; excepto que HET-3^b y HET-3^c no son 3-oxomorfolin-4-ilo.

En una incorporación particular, la presente invención provee un compuesto de fórmula (IB) o su sal farmacéuticamente aceptable,



(IB)

en donde:

R^{1b} se selecciona de $AR1^a$, $HET-1^a$ y $HET-2^a$; en donde R^{1b} se sustituye opcionalmente por 1, 2, 3 ó 4 grupos independientemente seleccionados de R^{3a} ;

R^{3a} se selecciona de halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $R^{82}NHCO-$, $R^{8a}CONH$, $R^{8a}OCONH-$, $R^{8a}OCON(Me)-$, $R^{8a}(1-6C)alquil-$, $-C_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $-OC_6H_{(13-3)}F_a$ (en donde a es 1 a 13), $(1-6C)alquiltio$, $di(1-6C)alquilaminocarbonilo$, $(3-8C)cicloalquiloxi(1-6C)alquilo$, $(3-8C)cicloalquil(1-6C)alcoxi$, $(3-8C)cicloalquil(1-6C)alcoxi(1-6C)alquilo$, $fenilo$, $HET-1^b$, $AR1^b-(1-6C)alcoxi$, $HET-1^b-(1-6C)alcoxi$, $HET-3^b-(1-6C)alcoxi$, $AR1^b-oxi(1-6C)alquilo$, $HET-1^b-oxi(1-6C)alquilo$, $HET-3^b-oxi(1-6C)alquilo$, $AR1^b-(1-6C)alcoxi(1-6C)alquilo$, $HET-1^b-(1-6C)alcoxi(1-6C)alquilo$, y $HET-3^b-(1-6C)alcoxi(1-6C)alquilo$;

en donde cualquier $fenilo$, $AR1^b$, $HET-1^b$, $HET-3$ o anillo $cicloalquilo$ en cualquiera de los valores anteriores para R^{3a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de R^{7d} ;

y en donde cualquier grupo R^{8a} , $(3-8C)cicloalquilo$, o $(1-6C)alquilo$ en cualquiera de las anteriores definiciones para R^{3a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de $(1-6C)alquilo$, $(1-6C)alcoxi$, $(2-6C)alquenilo$, $(2-6C)alquinilo$, $hidroxi$, $ciano$, $carboxi$, $aminocarbonilo$, $-C_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $amino$, $(1-6C)alquilamino$, $di(1-6C)alquilamino$, $(1-6C)alquiltio$, $(1-6C)alquilsulfinilo$, $(1-6C)alquilsulfonilo$, $aminosulfonilo$, $(1-6C)alquilaminosulfonilo$, $di(1-6C)alquilaminosulfonilo$, $(1-6C)alquilsulfonilamino$, $aminocarbonilo$, $(1-6C)alquilaminocarbonilo$, $di(1-$

6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alcoxicarbonilamino, (1-6C)alcoxicarbonil(N-metil)amino y (1-6C)alquilaminocarboniloxi; y/o cualquier grupo R^{8a} , (3-8C)cicloalquilo, o (1-6C)alquilo en cualquiera de las anteriores definiciones para R^{3a} puede sustituirse opcionalmente por 1, 2 ó 3 halo;

R^{2b} es:

- a) un anillo de 6 miembros seleccionado de fenilo, 3-piridilo y 3,5-pirimidinilo; sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o ambas de las posiciones *meta* al enlazamiento con el grupo -Y-NH- (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6c} ; o
- b) un anillo heteroarilo de a 5 miembros, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S); sustituido en un átomo de carbono en una posición que no es adyacente al enlazamiento al grupo -Y-NH-, por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en carbono (donde esté disponible) por 1 ó 2 grupos independientemente seleccionados de R^{6d} ;

R^{4a} se selecciona de:

- a) fenilo;
- b) HET-3^c (en donde cualquier átomo de nitrógeno disponible se sustituye por R^{5a} siempre que, con ello, no se cuaternice);
- c) HET-1^c;
- d) fenoxi, HET-1^c-oxi, R^{8b} O-, fenil(1-6C)alcoxi y HET-1^c-(1-6C)alcoxi;
- e) R^{8b} , R^{8b} O(1-6C)alquilo, amino(1-6C)alquilo, hidroxil(1-6C)alquilo, fenoxil(1-6C)alquilo, HET-1^c-oxil(1-6C)alquilo; y
- f) fenilamino, fenil(1-6C)alquilamino, HET-1^c-amino, HET-1^c-(1-6C)alquilamino, (1-6C)alquilamino, N-fenil-N-(1-6C)alquilamino, N-fenil(1-6C)alquil-N-(1-6C)alquilamino, N-(HET-1^c)-N-(1-6C)alquilamino, N-(HET-1^c-(1-6C)alquil)-N-(1-6C)alquilamino, y di(1-6C)alquilamino;

en donde cualquier alquilo, R^{8b} , fenilo, HET-1^c o anillo HET-3^c en cualquiera de las definiciones a) a f) para R^{4a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de R^{7c} ;

R^{5a} se selecciona de (1-6C)alquilo, (3-8C)cicloalquilo, fenilo (opcionalmente sustituido con 1 ó 2 halo), HET-1^d, $R^{8c}CO-$, fenilcarbonilo (en donde el fenilo se sustituye opcionalmente con 1 ó 2 halo), HET-1^d-carbonilo, (1-6C)alcoxycarbonilo, $R^{8c}NHCO-$, (1-6C)alcoxisulfonilo, (1-6C)alquilaminosulfonilo, (1-6C)alquilsulfinilo y (1-6C)alquilsulfonilo; en donde cualquier grupo (1-6C)alquilo en cualquiera de las definiciones de R^{5a} puede sustituirse opcionalmente por un sustituyente seleccionado de (1-6C)alcoxi, (1-6C)alquilS(O)_{p'}- (en donde p' es 0, 1 ó 2), amino, (1-6C)alquilamino, hidroxí, carboxí, (2-6C)alquenilo, (2-6C)alquinilo, fenilo (opcionalmente sustituido con 1 ó 2 halo) y HET-1^f;

R^{6c} y R^{6d} se seleccionan independientemente de R^{8d} , $R^{8d}O-$, $R^{8d}OCO-$, $R^{8d}COO-$, halo, ciano, halo(1-6C)alquilo, dihalo(1-6C)alquilo, (1-6C)alcoxi(1-6C)alquilo, (1-6C)alcoxi(1-6C)alcoxi, (1-6C)alquilcarbonilo, (1-6C)alquilS(O)_p- (en donde p es 0, 1 ó 2), (1-6C)alquilaminocarbonilo, di(1-6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alquilaminosulfonilo, di(1-6C)alquilaminosulfonilo, (1-6C)alquilsulfonilamino, AR1^c-(1-6C)alcoxi, HET-1^e-(1-6C)alcoxi, HET-3^e-(1-6C)alcoxi, AR1^e-oxi(1-6C)alquilo, HET-1^e-oxi(1-6C)alquilo, HET-3^e-oxi(1-6C)alquilo, AR1^e-(1-6C)alcoxi(1-6C)alquilo, HET-1^e-(1-6C)alcoxi(1-6C)alquilo, y HET-3^e-(1-6C)alcoxi(1-6C)alquilo;

en donde cualquier alquilo, R^{8d} , fenilo, AR1^c, HET-1^c, HET-3^c o anillo cicloalquilo en cualquiera de los valores anteriores para R^{6c} o R^{6d} puede sustituirse opcionalmente por 1, 2 ó 3 sustituyentes independientemente seleccionados de R^{7f} ;

R^{7d} , R^{7e} y R^{7f} se seleccionan independientemente de R^{8c} (opcionalmente sustituido con 1, 2 ó 3 halo o con hidroxí), $R^{8c}O-$ (opcionalmente sustituido con 1, 2 ó 3 halo), $R^{8c}OCO-$, $R^{8c}COO-$, $R^{8c}CO-$, $R^{8c}NHCO-$, $(R^{8c})_2NCO$, $R^{8c}CONH$, $R^{8c}OCONH-$, $R^{8c}OCON(Me)-$, $R^{8c}(1-$

6C)alquil-, halo, ciano, carboxi, carboxi(1-6C)alquilo, hidroxil, aminocarbonilo, $-C_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $-OC_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $R^{8e}S(O)_{p''}$ (en donde p'' es 0, 1 ó 2), amino(1-6C)alquilo, $R^{8e}NH(1-6C)alquilo$, $(R^{8e})_2N(1-6C)alquilo$, $R^{8e}NHSO_2-$, $(R^{8e})_2NSO_2-$, y $R^{8e}SO_2NH-$;

cada R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} y R^{8f} se selecciona de (1-6C)alquilo, (2-6C)alquénilo, (2-6C)alquinilo y (3-8C)cicloalquilo;

$AR1^a$, $AR1^b$ o $AR1^c$ se seleccionan independientemente de fenilo o un sistema de anillo fusionado, bicíclico de 6,5, o 6,6 miembros, en donde los dos anillos son carbocíclicos y en donde por lo menos el anillo enlazado al átomo de nitrógeno enlazante es completamente aromático;


$HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ y $HET-1^f$ se seleccionan independientemente de un anillo heteroarilo de a 5 ó 6 miembros, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO_2 , y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo;

$HET-2^a$ y $HET-2^b$ se seleccionan independientemente de un sistema de anillo heterocíclico aromático o parcialmente insaturado, bicíclico, fusionado de 5,5, 5,6 6,5 o 6,6 miembros, que contiene 1, 2, 3 ó 4 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en los cuales el anillo directamente enlazado al átomo de nitrógeno enlazante es aromático, y cualquiera de los dos anillos del sistema de anillo bicíclico puede ser carbocíclico;


$HET-3^b$, $HET-3^c$, $HET-3^e$ se seleccionan independientemente de un anillo heterocíclico de 4, 5, 6 ó 7 miembros, parcial o completamente insaturado, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO_2 , y en donde cualquier átomo de carbono puede oxidarse

opcionalmente a un grupo carbonilo; excepto que HET-3^b y HET-3^c no son 3-oxomorfolin-4-ilo.

Se entenderá que la definición de R^{2b} en relación con fórmula (IB) como un anillo de 6 miembros, seleccionado de fenilo, 3-piridilo y 3,5-pirimidinilo; sustituido en la posición que es *para* al enlaceamiento con el grupo -Y-NH- con un grupo R^{4a} y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o las dos posiciones que son *meta* al enlaceamiento con el grupo -Y-NH- (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6c}; comprende las siguientes estructuras, para cada definición del sustituyente R^{6c}.



Análogamente, se entenderá que la definición de R^{2b} como un anillo heteroarilo de 5 miembros, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S); sustituido en un átomo de carbono que no es directamente adyacente al enlaceamiento al grupo -Y-NH- por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en carbono (donde esté disponible) por 1 ó 2 grupos independientemente seleccionados de R^{6d}; comprende las siguientes estructuras:



Para evitar dudas, hay que entender que donde en esta especificación se califica un grupo por la expresión 'definido anteriormente' o 'definido de ahora en adelante', el dicho grupo encierra la definición más amplia y que

primero ocurra, así como cada una y todas las definiciones particulares para ese grupo.

Hay que entender que donde los sustituyentes contienen dos sustituyentes en una cadena alquilo, en la cual los dos están enlazados por un heteroátomo (por ejemplo dos sustituyentes alcoxi), entonces estos dos sustituyentes no son sustituyentes en el mismo átomo de carbonote la cadena alquilo.

Ejemplos de $AR1^a$, $AR1^b$ y $AR1^c$ son fenilo, naftilo, indenilo y dihidroindenilo.

Ejemplos de $AR1^a(1-6C)alquilo$, $AR1^b(1-6C)alquilo$ o $AR1^c(1-6C)alquilo$ son bencilo, fenetilo, naftilmetilo y naftiletilo.

Ejemplos de $AR1^a-oxi$, $AR1^b-oxi$ o $AR1^c-oxi$ son fenoxi y naftiloxi. Ejemplos de $AR1^a-oxi(1-6C)alquilo$, $AR1^b-oxi(1-6C)alquil$ o $AR1^c-oxi(1-6C)alquilo$ son fenoximetilo y naftiloxietilo. Ejemplos de $AR1^a-(1-6C)alcoxi$, $AR1^b-(1-6C)alcoxi$ o $AR1^c-(1-6C)alcoxi$ son cualquier valor de $AR1^a$, $AR1^b$ o $AR1^c$ enlazado a cualquier valor de $(1-6C)alcoxi$, por ejemplo benciloxi y naftiloxi. La misma convención se aplica siempre que $AR1^a$, $AR1^b$ o $AR1^c$ se enlaza a otro grupo.

Ejemplos de $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ o $HET-1^f$ son oxazolilo, oxadiazolilo, piridilo, pirimidinilo, imidazolilo, triazolilo, pirazinilo, piridazinilo, pirrolilo, tienilo y furilo.

Otros ejemplos de $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ o $HET-1^f$ incluyen tiazolilo, tetrazolilo, isoxazolilo, pirazolilo, triazinilo, tiadiazolilo, oxadiazolilo y isotiazolilo.

Ejemplos de $HET-1^a-oxi$, $HET-1^b-oxi$, $HET-1^c-oxi$, $HET-1^d-oxi$, $HET-1^e-oxi$ o $HET-1^f-oxi$ son oxazolil-2-oxi, 1,2,4-oxadiazolil-3-oxi, piridil-2-oxi, y pirimidinil-4-oxi. La convención ya ilustrada para $AR1^a$, $AR1^b$ y $AR1^c$ se aplica a $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ o $HET-1^f$ enlazado a cualquier otro grupo.

Ejemplos de $HET-2^a$ o $HET-2^b$ son 1,3-benzodioxol-5-ilo, quinazolinilo, purinilo, indolilo, benzofurilo, benzotienilo, cromanilo, isocromanilo, quinolinilo e isoquinolinilo. Otros ejemplos de $HET-2^a$ o

HET-2^b incluyen quinoxalinilo, benzotiazolilo y benzoxazolilo. Ejemplos de HET-3^b, HET-3^c o HET-3^e son morfolinilo, tiomorfolinilo, piperazinilo, piperidinilo, 2-oxopiperidinilo, pirrolidinilo, 2-oxopirrolidinilo, tetrahidropiranilo, dihidropiranilo, tetrahidrotienilo, azetidinilo, homomorfolinilo, diazepinilo y azepinilo.

Ejemplos particulares de HET-3, HET-3^c o HET-3^e son morfolinilo, tiomorfolinilo, piperazinilo y piperidinilo.

Ejemplos de HET-3^b-oxi, HET-3^c-oxi o HET-3^e-oxi son morfoliniloxi, tiomorfoliniloxi, piperaziniloxi, piperidiniloxi, 2-oxopiperidiniloxi, pirrolidiniloxi, 2-oxopirrolidiniloxi, tetrahidropiraniloxi, dihidropiraniloxi, tetrahidrotieniloxi, azetidiniloxi, homomorfoliniloxi, diazepiniloxi y azepiniloxi. La convención ya ilustrada para AR1^a, AR1^b y AR1^c se aplica a HET-3^b, HET-3^c o HET-3^e enlazado a cualquier otro grupo.

Si no indicado en alguna otra parte, sustituyentes opcionales apropiados para un grupo particular son aquellos especificados para grupos similares en esta solicitud.

Un compuesto de fórmula (I) puede formar sales básicas o ácidas estables y, en tales casos, la administración de un compuesto como una sal puede ser apropiada, y sales farmacéuticamente aceptables pueden elaborarse por métodos convencionales tales como los descritos a continuación. Sales apropiadas farmacéuticamente aceptables incluyen sales de adición de ácido tales como metanosulfonate, tosilato, α -glicerofosfato, fumarato, clorhidrato, citrato, maleato, tartrato y (menos preferiblemente) hidrobromuro. Son igualmente apropiadas las sales formadas con ácido fosfórico y sulfúrico. En otro aspecto sales apropiadas son sales básicas tales como una sal de metal alcalino por ejemplo sodio, una sal de metal alcalinotérreo por ejemplo calcio o magnesio, una sal de amina orgánica por ejemplo trietilamina, morfolina, *N*-metilpiperidina, *N*-etilpiperidina, procaina, dibencilamina, *N,N*-dibenciletilamina, tris-(2-hidroxietil)amina, *N*-metil *D*-glucamina y aminoácidos tales como lisina. Pueden ser más de un catión o anión dependiendo del número de funciones

cargadas y la valencia de los cationes y aniones. Una sal farmacéuticamente aceptable preferida es la sal de sodio.

Sin embargo, para facilitar el aislamiento de la sal durante la preparación, pueden preferirse sales que son menos solubles en el disolvente escogido, sean farmacéuticamente aceptables o no.

En la presente invención hay que entender que un compuesto de la fórmula (I) o su sal puede exhibir el fenómeno de tautomerismo y que los dibujos de las fórmulas en esta especificación pueden representar únicamente una de las posibles formas tautoméricas. Hay que entender que la invención abarca cualquier forma tautomérica que inhiba la actividad DGAT1 y no hay que limitarla simplemente a una forma tautomérica cualquiera utilizada en los dibujos de las fórmulas.

En el arte se conocen diferentes formas de profármacos. Para ejemplos de tales derivados de profármacos, véase:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- d) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- e) N. Kakeya, *et al.*, Chem Pharm Bull, 32, 692 (1984).

Ejemplos de tales profármacos son ésteres escindibles *in vivo* de un compuesto de la invención. Un éster escindible *in vivo* de un compuesto de la invención que contiene un grupo carboxi es, por ejemplo, un éster farmacéuticamente aceptable que se escinde en el cuerpo humano o animal para producir el ácido original. Ésteres apropiados farmacéuticamente aceptables para carboxi incluyen (1-6C)alquil ésteres, por ejemplo metilo o etilo; (1-6C)alcoximetil ésteres, por ejemplo metoximetilo; (1-

6C)alcanoiloximetil ésteres, por ejemplo pivaloiloximetilo; ftalidil ésteres; (3-8C)cicloalcoxicarboniloxi(1-6C)alquil ésteres, por ejemplo 1-ciclohexilcarboniloxietilo; 1,3-dioxolan-2-ilmetil ésteres, por ejemplo 5-metil-1,3-dioxolan-2-ilmetilo; (1-6C)alcoxicarboniloxietil ésteres, por ejemplo 1-metoxycarboniloxietilo; aminocarbonilmetil ésteres y sus versiones mono- o di-N-((1-6C)alquilo), por ejemplo N,N-dimetilaminocarbonilmetil ésteres y N-etilaminocarbonilmetil ésteres; y pueden formarse en cualquier grupo carboxi en los compuestos de esta invención. Un éster escindible *in vivo* de un compuesto de la invención que contiene un grupo hidroxil es, por ejemplo, un éster farmacéuticamente aceptable que se escinde en el cuerpo humano o animal para producir el grupo hidroxil original. Ésteres apropiados farmacéuticamente aceptables para hidroxil incluyen (1-6C)alcanoil ésteres, por ejemplo acetil ésteres; y benzoil ésteres en donde el grupo fenilo puede sustituirse con aminometilo o mono- o di-(1-6C)alquil aminometilo N- sustituido, por ejemplo 4-aminometilbenzoil ésteres y 4-N,N-dimetilaminometilbenzoil ésteres.

Se observará por las personas expertas en el arte que ciertos compuestos de fórmula (I) contienen átomos de carbono y/o azufre asiméricamente sustituidos y, en consecuencia, pueden existir en, y aislarse en, formas óptimamente activas y racémicas. Algunos compuestos pueden exhibir polimorfismo. Hay que entender que la presente invención comprende cualquier forma racémica, ópticamente activa, polimórfica o estereoisomérica, o sus mezclas, forma que posee propiedades útiles en la inhibición de la actividad DGAT1, siendo bien conocido en el arte cómo preparar formas ópticamente activas (por ejemplo, mediante resolución de la forma racémica por técnicas de recristalización, mediante síntesis de materiales de partida óptimamente activos, mediante síntesis quiral, mediante resolución enzimática, mediante biotransformación, o mediante separación cromatográfica usando una fase estacionaria quiral) y cómo determinar la eficacia para la inhibición de la actividad DGAT1 mediante las pruebas estándar descritas más adelante.

Hay que entender asimismo que ciertos compuestos de la fórmula (I) y sus sales pueden existir en formas solvatadas así como no solvatadas tales como, por ejemplo, formas hidratadas. Hay que entender que la invención abarca la totalidad de las formas solvatadas que inhiben la actividad DGAT1.

Como ya se indicó, hemos descubierto una gama de compuestos que poseen excelente actividad inhibidora de DGAT1. Éstos tienen buenas propiedades físicas y/o farmacocinéticas en general. Los siguientes compuestos poseen propiedades farmacéuticas y/o físicas y/o farmacocinéticas preferidas.

Aspectos particulares de la invención comprenden un compuesto de fórmula (I), o su sal farmacéuticamente aceptable, en donde los sustituyentes AR, HET, R¹ a R⁸ y demás sustituyentes ya mencionados tienen los valores definidos anteriormente, o cualquiera de los siguientes valores (los cuales pueden utilizarse donde se considere apropiado con cualquiera de las definiciones e incorporaciones ya divulgadas o que se divulgarán a partir de ahora):

En una incorporación de la invención se provee compuestos de fórmula (I), en una incorporación alternativa se proveen sales farmacéuticamente aceptables de compuestos de fórmula (I). En una incorporación adicional, se proveen profármacos, particularmente ésteres escindibles *in vivo*, de compuestos de fórmula (I). En una incorporación adicional, se proveen sales farmacéuticamente aceptables de profármacos de compuestos de fórmula (I). La referencia incluida en esta solicitud a un compuesto de fórmula (I) debe tomarse en general para aplicarse también a compuestos de las fórmulas (IA), (IB), (IZA), (IZB) y (IZC).

Valores particulares de grupos variables en compuestos de las fórmulas (I), (IA) y (IB) son los siguientes. Tales valores pueden usarse donde se considere apropiado con cualquiera de los demás valores, definiciones, reivindicaciones o incorporaciones definidas anteriormente o más adelante.

- 1) R¹, R^{1a} o R^{1b} es AR^{1a}

- 2) R^1, R^{1a} o R^{1b} es HET-1^a
- 3) R^1, R^{1a} o R^{1b} es HET-2^a
- 4) R^1, R^{1a} o R^{1b} no se sustituye
- 5) R^1 o R^{1a} se sustituye por 1 grupo seleccionado de R^3 o R^{1b} se sustituye por 1 grupo seleccionado de R^{3a}
- 6) R^1 o R^{1a} se sustituye por 2 grupos independientemente seleccionados de R^3 o R^{1b} se sustituye por 2 grupos independientemente seleccionados de R^{3a}
- 7) R^1 o R^{1a} se sustituye por 3 grupos independientemente seleccionados de R^3 o R^{1b} se sustituye por 3 grupos independientemente seleccionados de R^{3a}
- 8) R^1 o R^{1a} se sustituye por 4 grupos independientemente seleccionados de R^3 o R^{1b} se sustituye por 4 grupos independientemente seleccionados de R^{3a}
- 9) R^3 o R^{3a} es halo
- 10) R^3 o R^{3a} se selecciona de $R^{8a}, R^{8a}O-, R^{8a}OCO-, R^{8a}COO-$ y $R^{8a}CO-$
- 11) R^3 o R^{3a} se selecciona de $R^{8a}, R^{8a}O-$ y $R^{8a}OCO-$
- 12) R^3 o R^{3a} se selecciona de $R^{8a}NHCO-, R^{8a}CONH, R^{8a}OCONH-$ y $R^{8a}OCON(Me)-$
- 13) R^3 o R^{3a} se selecciona de $R^{8a}(1-6C)alquil-, haloC_{1-6}alquilo$ o $haloC_{1-6}alcoxi$
- 14) R^3 o R^{3a} se selecciona de $R^{8a}(1-6C)alquil-, -C_6H_{(13-a)}F_a$ (en donde a es 1 a 14), $-OC_6H_{(13-a)}F_a$ (en donde a es 1 a 14)
- 15) R^3 o R^{3a} se selecciona de $(1-6C)alquiltio, alquilcarbonilo$ y $di(1-6C)alquilaminocarbonilo$
- 16) R^3 o R^{3a} se selecciona de $(3-8C)cicloalquiloxi(1-6C)alquilo, (3-8C)cicloalquil(1-6C)alcoxi,$ y $(3-8C)cicloalquil(1-6C)alcoxi(1-6C)alquilo$
- 17) R^3 o R^{3a} se selecciona de halo, $R^{8a}, R^{8a}O-, R^{8a}OCO-, R^{8a}COO-, R^{8a}CO-, R^{8a}NHCO-, R^{8a}CONH, R^{8a}OCONH-, R^{8a}OCON(Me)-, R^{8a}(1-6C)alquil-, halo(1-6C)alquilo$ (tal como $-C_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $halo(1-6C)alcoxi$ (tal como $-OC_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $(1-$

6C)alquiltio, di(1-6C)alquilaminocarbonilo, (3-8C)cicloalquiloxi(1-6C)alquilo, (3-8C)cicloalquil(1-6C)alcoxi, (3-8C)cicloalquil(1-6C)alcoxi(1-6C)alquilo, AR1^b-(1-6C)alcoxi, HET-1^b-(1-6C)alcoxi, HET-3^b-(1-6C)alcoxi, AR1^b-oxi(1-6C)alquilo, HET-1^b-oxi(1-6C)alquilo, HET-3^b-oxi(1-6C)alquilo, AR1^b-(1-6C)alcoxi(1-6C)alquilo, HET-1^b-(1-6C)alcoxi(1-6C)alquilo, y HET-3^b-(1-6C)alcoxi(1-6C)alquilo

18) R³ o R^{3a} se selecciona de fenilo y HET-1^b

19) R³ o R^{3a} es fenilo

20) R³ o R^{3a} se selecciona de AR1^b-(1-6C)alcoxi, y HET-1^b-(1-6C)alcoxi, HET-3^b-(1-6C)alcoxi

21) R³ o R^{3a} se selecciona de AR1^b-oxi(1-6C)alquilo, HET-1^b-oxi(1-6C)alquilo, HET-3^b-oxi(1-6C)alquilo, AR1^b-(1-6C)alcoxi(1-6C)alquilo, HET-1^b-(1-6C)alcoxi(1-6C)alquilo, y HET-3^b-(1-6C)alcoxi(1-6C)alquilo

22) un anillo fenilo, AR1^b, HET-1^b, HET-3^b o cicloalquilo en cualquier valor para R³ se sustituye por 1 ó 2 sustituyentes independientemente seleccionados de R^{7a}, y un anillo fenilo, AR1^b, HET-1^b, HET-3^b o cicloalquilo en cualquier valor para R^{3a} se sustituye por 1 ó 2 sustituyentes independientemente seleccionados de R^{7d}

23) un anillo fenilo, AR1^b, HET-1^b, HET-3^b o cicloalquilo en cualquier valor para R³ o R^{3a} no se sustituye

24) un grupo R^{8a}, (3-8C)cicloalquilo, o (1-6C)alquilo en cualesquiera definiciones para R³ o R^{3a} no se sustituye

25) un grupo R^{8a}, (3-8C)cicloalquilo, o (1-6C)alquilo en cualesquiera definiciones para R³ o R^{3a} se sustituye por 1 ó 2 sustituyentes independientemente seleccionados de (1-6C)alquilo, (1-6C)alcoxi, (2-6C)alquenilo, (2-6C)alquinilo, hidroxilo, ciano, carboxi, aminocarbonilo, -C₆H_(14-a)F_a (en donde a es 1 a 14), amino, (1-6C)alquilamino, di(1-6C)alquilamino, (1-6C)alquiltio, (1-6C)alquilsulfinilo, (1-6C)alquilsulfonilo, aminosulfonilo, (1-6C)alquilaminosulfonilo, di(1-6C)alquilaminosulfonilo, (1-6C)alquilsulfonilamino, aminocarbonilo, (1-6C)alquilaminocarbonilo, di(1-6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alcoxycarbonilamino, (1-6C)alcoxycarbonil

(N-metil)amino y (1-6C)alquilaminocarbonilo; y/o sustituido por 1, 2 ó 3 halo;

26) R^2 o R^{2a} es un anillo de 6 miembros seleccionado de fenilo, 3-piridilo, 5-piridilo y 3,5-pirimidinilo; sustituido en la posición 4 por un grupo seleccionado de R^4 y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o dos posiciones *meta* en relación con el enlazamiento al grupo -Y-NH- (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6a} ; o

27) R^{2b} es un anillo de 6 miembros seleccionado de fenilo, 3-piridilo, 5-piridilo y 3,5-pirimidinilo; sustituido en la posición 4 por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o dos posiciones *meta* en relación con el enlazamiento al grupo -Y-NH- (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6c} ;

28) R^2 , R^{2a} o R^{2b} es fenilo

29) R^2 , R^{2a} o R^{2b} es 3-piridilo

30) R^2 , R^{2a} o R^{2b} es fenilo o 3-piridilo

31) R^2 o R^{2a} es un anillo de 6 miembros sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^4 ; o un grupo R^{2b} es un anillo de 6 miembros sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^{4a}

32) R^2 o R^{2a} es un anillo de 6 miembros sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^4 y sustituido adicionalmente en un átomo de carbono en una posición 3 por R^{6a} ; o R^{2b} es un anillo de 6 miembros sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^{4a} y sustituido adicionalmente en un átomo de carbono en una posición 3 por R^{6c} ;

33) R^2 o R^{2a} es un grupo o sub-fórmula (iv) como ya se definió; o R^{2b} es un anillo heteroarilo de 5 miembros, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya

enlaces O-O, S-S u O-S); sustituido en un átomo de carbono en una posición que no es adyacente al enlazamiento al grupo -Y-NH- por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en carbono (donde esté disponible) por 1 ó 2 grupos independientemente seleccionados de R^{6d}

34) R^2 o R^{2a} es un grupo HET-2^b opcionalmente sustituido con metilo

35) R^4 o R^{4a} es fenilo

36) R^4 o R^{4a} es HET-3^c (en donde cualquier átomo de nitrógeno disponible se sustituye por R^5 siempre que, con ello, no se cuaternice)

37) R^4 o R^{4a} es HET-1^c

38) R^4 o R^{4a} se selecciona de fenoxi, HET-1^c-oxi, R^{8b} O-, fenil(1-6C)alcoxi y HET-1^c-(1-6C)alcoxi

39) R^4 o R^{4a} se selecciona de R^{8b} , R^{8b} O(1-6C)alquilo, amino(1-6C)alquilo, hidroxil(1-6C)alquilo, fenoxil(1-6C)alquilo y HET-1^c-oxil(1-6C)alquilo

40) R^4 o R^{4a} se selecciona de R^{8b} e hidroxil(1-6C)alquilo

41) R^4 o R^{4a} se selecciona de fenilamino, fenil(1-6C)alquilamino, HET-1^c-amino, HET-1^c-(1-6C)alquilamino, (1-6C)alquilamino, N-fenil-N-(1-6C)alquilamino, N-fenil(1-6C)alquil-N-(1-6C)alquilamino, N-(HET-1^c)-N-(1-6C)alquilamino, N-(HET-1^c-(1-6C)alquil)-N-(1-6C)alquilamino, y di(1-6C)alquilamino

42) R^4 o R^{4a} se selecciona de fenilamino, N-fenil-N-(1-6C)alquilamino

43) R^4 o R^{4a} se selecciona de fenilo, HET-3^c, fenoxi, HET-1^c-oxi, R^{8b} O-, fenil(1-6C)alcoxi, HET-1^c-(1-6C)alcoxi, R^{8b} , hidroxil(1-6C)alquilo, fenilamino y N-fenil-N-(1-6C)alquilamino

44) un anillo alquilo, R^{8b} , fenilo, HET-1^c o HET-3^c en cualquiera de las definiciones para R^4 se sustituye por 1 ó 2 sustituyentes independientemente seleccionados de R^{7b} ; o un anillo alquilo, R^{8b} , fenilo, HET-1^c o HET-3^c en cualquiera de las definiciones para R^{4a} se sustituye por 1 ó 2 sustituyentes independientemente seleccionados de R^{7c}

45) un anillo alquilo, R^{8b} , fenilo, HET-1^c o HET-3^c en cualquiera de las definiciones para R^4 o R^{4a} no se sustituye

- 46) R^5 o R^{5a} se selecciona de (1-6C)alquilo y (3-8C)cicloalquilo
- 47) R^5 o R^{5a} es $R^{8c}CO-$
- 48) R^5 o R^{5a} es fenilo (opcionalmente sustituido) o HET-1^d
- 49) R^5 o R^{5a} es fenilcarbonilo (opcionalmente sustituido) o HET-1^d-carbonilo
- 50) R^5 o R^{5a} se selecciona de carboxi, (1-6C)alcoxycarbonilo, $R^{8c}NHCO-$, (1-6C)alcoxisulfonilo, (1-6C)alquilaminosulfonilo, (1-6C)alquilsulfinilo y (1-6C)alquilsulfonilo
- 51) R^5 o R^{5a} se selecciona de (1-6C)alcoxycarbonilo y (1-6C)alquilsulfonilo
- 52) R^5 o R^{5a} se selecciona de (1-6C)alquilo, HET-1, $R^{8c}CO-$, (1-6C)alcoxycarbonilo y (1-6C)alquilsulfonilo
- 53) cualquier grupo (1-6C)alquilo en cualquiera de las definiciones de R^5 o R^{5a} se sustituye por un sustituyente seleccionado de (1-6C)alcoxi, (1-6C)alquilS(O)_p- (en donde *p* es 0, 1 ó 2), amino, (1-6C)alquilamino, hidroxilo, carboxi, (2-6C)alquenilo, (2-6C)alquinilo, fenilo (opcionalmente sustituido) y HET-1^d
- 54) cualquier grupo (1-6C)alquilo en cualquiera de las definiciones de R^5 o R^{5a} no se sustituye
- 55) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de R^{8d} , $R^{8d}O-$, $R^{8d}OCO-$ y $R^{8d}COO-$
- 56) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de halo y ciano
- 57) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de halo(1-6C)alquilo, dihalo(1-6C)alquilo
- 58) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de (1-6C)alcoxi(1-6C)alquilo y (1-6C)alcoxi(1-6C)alcoxi
- 59) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de (1-6C)alquilcarbonilo y (1-6C)alquilS(O)_p- (en donde *p* es 0, 1 ó 2)
- 60) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de (1-6C)alquilaminocarbonilo, di(1-6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alquilaminosulfonilo, di(1-6C)alquilaminosulfonilo y (1-6C)alquilsulfonilamino

- 61) R^{6a} , R^{6b} , R^{6c} o R^{6d} se selecciona de $AR1^c$ -(1-6C)alcoxi, $HET-1^c$ -(1-6C)alcoxi, $HET-3^c$ -(1-6C)alcoxi, $AR1^c$ -oxi(1-6C)alquilo, $HET-1^c$ -oxi(1-6C)alquilo, $HET-3^c$ -oxi(1-6C)alquilo, $AR1^c$ -(1-6C)alcoxi(1-6C)alquilo, $HET-1^c$ -(1-6C)alcoxi(1-6C)alquilo, y $HET-3^c$ -(1-6C)alcoxi(1-6C)alquilo
- 62) cualquier anillo alquilo, R^{8d} , fenilo, $AR1^c$, $HET-1^c$, $HET-3^c$ o cicloalquilo en cualquier valor para R^{6a} , R^{6b} se sustituye por 1, 2 ó 3 sustituyentes independientemente seleccionados de R^{7c} , o en cualquier valor para R^{6c} o R^{6d} se sustituye por 1, 2 ó 3 sustituyentes independientemente seleccionados de R^{7f}
- 63) cualquier anillo alquilo, R^{8d} , fenilo, $AR1^c$, $HET-1^c$, $HET-3^c$ o cicloalquilo en cualquier valor para R^{6a} , R^{6b} , R^{6c} o R^{6d} no se sustituye
- 64) R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} o R^{7f} es R^{8c} (opcionalmente sustituido con 1, 2 ó 3 halo o con hidroxí)
- 65) R^{7a} , R^{7b} , R^{7c} , R^{7d} , R^{7e} o R^{7f} es $R^{8c}O$ - (opcionalmente sustituido con 1, 2 ó 3 halo)
- 66) R^{7a} , R^{7b} o R^{7c} se selecciona de $R^{8c}OCO$ -, $R^{8c}COO$ -, $R^{8c}CO$ -, $R^{8c}NHCO$ -, $R^{10}R^{11}NCO$, $R^{8c}CONH$, $R^{8c}OCONH$ - y $R^{8c}OCON(Me)$ -
- 67) R^{7d} , R^{7e} o R^{7f} se selecciona de $R^{8c}OCO$ -, $R^{8c}COO$ -, $R^{8c}CO$ -, $R^{8c}NHCO$ -, $(R^{8c})_2NCO$, $R^{8c}CONH$, $R^{8c}OCONH$ - y $R^{8c}OCON(Me)$ -
- 68) R^{7a} , R^{7b} o R^{7c} se selecciona de R^{8c} (1-6C)alquil-, halo, ciano, carboxi, carboxi(1-6C)alquilo, hidroxí, aminocarbonilo, halo(1-6C)alquilo, halo(1-6C)alcoxi
- 69) R^{7d} , R^{7e} o R^{7f} se selecciona de R^{8c} (1-6C)alquil-, halo, ciano, carboxi, carboxi(1-6C)alquilo, hidroxí, aminocarbonilo, $-C_6H_{(13-3)}F_a$ (en donde a es 1 a 13), $-OC_6H_{(14-a)}F_a$ (en donde a es 1 a 13)
- 70) R^{7a} , R^{7b} o R^{7c} se selecciona de $R^{8c}S(O)_p$ - (en donde p es 0, 1 ó 2), amino(1-6C)alquilo, $R^{8c}NH(1-6C)alquilo$, $R^{11}R^{10}N(1-6C)alquilo$, $R^{8c}NHSO_2$ -, $R^{11}R^{10}NSO_2$ -, y $R^{8c}SO_2NH$ -
- 71) R^{7d} , R^{7e} o R^{7f} se selecciona de $R^{8c}S(O)_p$ - (en donde p es 0, 1 ó 2), amino(1-6C)alquilo, $R^{8c}NH(1-6C)alquilo$, $(R^{8c})_2N(1-6C)alquilo$, $R^{8c}NHSO_2$ -, $(R^{8c})_2NSO_2$ -, y $R^{8c}SO_2NH$ -

- 72) R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} o R^{8f} se selecciona de (1-6C)alquilo, (2-6C)alquenilo, (2-6C)alquinilo, (3-8C)cicloalquilo
- 73) R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} o R^{8f} se selecciona de (1-6C)alquilo y (3-8C)cicloalquilo
- 74) R^{8a} , R^{8b} , R^{8c} , R^{8d} , R^{8e} o R^{8f} se selecciona de (1-6C)alquilo
- 75) $AR1^a$, $AR1^b$ o $AR1^c$ es fenilo
- 76) $AR1^a$, $AR1^b$ o $AR1^c$ es un sistema de anillo fusionado bicíclico de 6,5, o 6,6 miembros,
- 77) $AR1^a$, $AR1^b$ o $AR1^c$ es naftilo
- 78) $AR1^a$, $AR1^b$ o $AR1^c$ es fenilo, dihidroindenilo o naftilo
- 79) $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ o $HET-1^f$ es un anillo heteroarilo de 5 miembros
- 80) $HET-1^a$, $HET-1^b$, $HET-1^c$, $HET-1^d$, $HET-1^e$ o $HET-1^f$ es un anillo heteroarilo de 6 miembros
- 81) $HET-2^a$ o $HET-2^b$ es un sistema de anillo heterocíclico bicíclico 5,5 o 6,6 fusionado
- 82) $HET-2^a$ o $HET-2^b$ es un sistema de anillo heterocíclico bicíclico es un 5,6-6,5 fusionado
- 83) $HET-3^b$, $HET-3^c$ o $HET-3^e$ es un anillo de 5 miembros
- 84) $HET-3^b$, $HET-3^c$ o $HET-3^e$ es un anillo de 6 miembros
- 85) $HET-3^b$, $HET-3^c$ o $HET-3^e$ es un anillo de 7 miembros
- 86) $HET-3^b$, $HET-3^c$ o $HET-3^e$ es un anillo de 4 miembros
- 87) s es convenientemente 0

En todavía otros compuestos preferidos:

- 88) R^4 es un grupo R^{8b} donde R^{8b} es (3-8C)cicloalquilo tal como ciclohexilo
- 89) R^4 es un grupo R^{8b} donde R^{8b} se sustituye por un sustituyente R^{7b}
- 90) R^{7b} es $H_2NCO(CR^{13}R^{14})_n$ - o $(R^{10}R^{11})NCO(CH_2)_n$
- 91) R^{7b} es $HCONH(CR^{13}R^{14})_n$, $HCONR^{8c}(CR^{13}R^{14})_n$, $R^{8c}CONR^{8f}(CR^{13}R^{14})_n$
- 92) R^{7b} es R^{12} o R^{12} (1-6C)alquilo,
- 93) R^2 o R^{2a} es un grupo ciclohexilo.

En un aspecto adicional de la invención se provee un compuesto de la fórmula (IB') o su sal farmacéuticamente aceptable, el cual un compuesto de la fórmula (IB) como ya se indicó, en donde:

R^{1b} se selecciona de $AR1^a$, HET-1^a y HET-2^a; en donde R^{1b} se sustituye opcionalmente por 1, ó 2 grupos independientemente seleccionados de R^{3a} ;

R^{3a} se selecciona de halo, R^{8a} , $R^{8a}O-$, $R^{8a}OCO-$, $R^{8a}COO-$, $R^{8a}CO-$, $R^{8a}NHCO-$, $R^{8a}CONH-$, $R^{8a}OCONH-$, $R^{8a}OCON(Me)-$, $R^{8a}(1-6C)alquil-$, $-C_6H_{(13-a)}F_a$ (en donde a es 1 a 13), $-OC_6H_{(13-a)}F_a$ (en donde a es 1 a 13), (1-6C)alquiltio, di(1-6C)alquilaminocarbonilo, fenilo, HET-1^b, $AR1^b-(1-6C)alcoxi$, HET-1^b-(1-6C)alcoxi, HET-3^b-(1-6C)alcoxi, $AR1^b-oxi(1-6C)alquilo$, HET-1^b-oxi(1-6C)alquilo, HET-3^b-oxi(1-6C)alquilo, $AR1^b-(1-6C)alcoxi(1-6C)alquilo$, HET-1^b-(1-6C)alcoxi(1-6C)alquilo, y HET-3^b-(1-6C)alcoxi(1-6C)alquilo;

en donde cualquier anillo fenilo, $AR1^b$, HET-1^b, HET-3^b o cicloalquilo en cualquiera de los valores anteriores para R^{3a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de R^{7d} ;

y en donde cualquier grupo R^{8a} , o (1-6C)alquilo en cualquiera de las anteriores definiciones para R^{3a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de (1-6C)alquilo, (1-6C)alcoxi, hidroxilo, carboxi, $-C_6H_{(13-a)}F_a$ (en donde a es 1 a 13), (1-6C)alquilamino, di(1-6C)alquilamino, (1-6C)alquilsulfonilo, (1-6C)alquilaminosulfonilo, di(1-6C)alquilaminosulfonilo, (1-6C)alquilsulfonilamino, aminocarbonilo, (1-6C)alquilaminocarbonilo, di(1-6C)alquilaminocarbonilo, (1-6C)alquilcarbonilamino, (1-6C)alcoxycarbonilamino; y/o cualquier grupo R^{8a} , o (1-6C)alquilo en cualquiera de las anteriores definiciones para R^{3a} puede sustituirse opcionalmente por 1, 2 ó 3 halo;

R^{2b} es:

a) un anillo de 6 miembros seleccionado de fenilo, 3-piridilo y 3,5-pirimidinilo; sustituido en la posición que es *para* al enlazamiento del grupo

-Y-NH- por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o dos posiciones 3 (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6c} ;

R^{4a} se selecciona de:

- a) fenilo;
- b) HET-3^c (en donde cualquier átomo de nitrógeno disponible se sustituye por R^{5a} siempre que, con ello, no se cuaternice);
- c) HET-1^c;
- d) fenoxi, HET-1^c-oxi, $R^{8b}O-$, fenil(1-6C)alcoxi y HET-1^c-(1-6C)alcoxi;
- e) R^{8b} , $R^{8b}O(1-6C)$ alquilo, amino(1-6C)alquilo, hidroxil(1-6C)alquilo, fenoxil(1-6C)alquilo, HET-1^c-oxil(1-6C)alquilo; y
- f) fenilamino, fenil(1-6C)alquilamino, HET-1^c-amino, HET-1^c-(1-6C)alquilamino, (1-6C)alquilamino, N-fenil-N-(1-6C)alquilamino, N-fenil(1-6C)alquil-N-(1-6C)alquilamino, N-(HET-1^c)-N-(1-6C)alquilamino, N-(HET-1^c-(1-6C)alquil)-N-(1-6C)alquilamino, y di(1-6C)alquilamino; en donde cualquier anillo alquilo, R^{8b} , fenilo, HET-1^c o HET-3^c en cualquiera de las definiciones a) a f) para R^{4a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de R^{7c} ;

R^{5a} se selecciona de (1-6C)alquilo, (3-8C)cicloalquilo, fenilo (opcionalmente sustituido con 1 ó 2 halo), HET-1^d, $R^{8c}CO-$, fenilcarbonilo (en donde el fenilo se sustituye opcionalmente con 1 ó 2 halo), HET-1^d-carbonilo, (1-6C)alcoxycarbonilo, $R^{8c}NHCO-$, (1-6C)alcoxisulfonilo, (1-6C)alquilaminosulfonilo, (1-6C)alquilsulfonilo y (1-6C)alquilsulfonilo; en donde cualquier grupo (1-6C)alquilo en cualquiera de las definiciones de R^{5a} puede sustituirse opcionalmente por un sustituyente seleccionado de (1-6C)alcoxi, (1-6C)alquilS(O)_{p'} (en donde p' es 0, 1 ó 2), amino, (1-6C)alquilamino, hidroxil, carboxil, fenilo (opcionalmente sustituido con 1 ó 2 halo) y HET-1^f;

R^{6c} y R^{6d} se seleccionan independientemente de R^{8d} , $R^{8d}O-$, $R^{8d}OCO-$, $R^{8d}COO-$, halo, ciano, halo(1-6C)alquilo, dihalo(1-6C)alquilo, (1-

6C)alcoxi(1-6C)alquilo, (1-6C)alcoxi(1-6C)alcoxi, (1-6C)alquilcarbonilo y (1-6C)alquilS(O)_p- (en donde *p* es 0, 1 ó 2), en donde cualquier alquilo o R^{8d} en cualquiera de los valores anteriores para R^{6c} o R^{6d} puede sustituirse opcionalmente por 1, 2 ó 3 sustituyentes independientemente seleccionados de R^{7f};

R^{7d}, R^{7c} y R^{7f} se seleccionan independientemente de R^{8c} (opcionalmente sustituido con 1, 2 ó 3 halo o con hidroxilo), R^{8c}O- (opcionalmente sustituido con 1, 2 ó 3 halo), halo, ciano, carboxi, carboxi(1-6C)alquilo, hidroxilo, aminocarbonilo, -C₆H_(13-a)F_a (en donde *a* es 1 a 13), -OC₆H_(13-a)F_a (en donde *a* es 1 a 13), R^{8c}S(O)_{p''}- (en donde *p''* es 0, 1 ó 2);

cada R^{8a}, R^{8b}, R^{8c}, R^{8d}, R^{8e} o R^{8f} se selecciona de (1-6C)alquilo;

cada AR1^a, AR1^b o AR1^c es fenilo o un sistema de anillo bicíclico 6,5, ó 6,6 fusionado, en donde los dos anillos son carbocíclicos y en donde por lo menos el anillo enlazado al átomo de nitrógeno enlazante es completamente aromático;

cada HET-1^a, HET-1^b, HET-1^c, HET-1^d, HET-1^e o HET-1^f es un anillo heteroarilo de 5 ó 6 miembros, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO₂, y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo;

cada HET-2^a o HET-2^b es un sistema de anillo heterocíclico aromático o parcialmente saturado, bicíclico, 5,5, 5,6, 6,5 ó 6,6 fusionado, que contiene 1, 2, 3 ó 4 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en los cuales el anillo directamente enlazado al átomo de nitrógeno enlazante es aromático, y cualquiera de los dos anillos del sistema de anillo bicíclico puede ser carboxílico;

HET-3^b, HET-3^c o HET-3^e se selecciona independientemente de un anillo heterocíclico de 4, 5, 6 ó 7 miembros, parcial o completamente saturado, que contiene 1, 2, ó 3 heteroátomos independientemente

seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente en un grupo SO o SO₂, y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo; excepto que HET-3^b, HET-3^c o HET-3^e no es 3-oxomorfolin-4-ilo.

En un aspecto adicional de la invención se provee un compuesto de la fórmula (IB'') o su sal farmacéuticamente aceptable, que es un compuesto de la fórmula (IB) como ya se indicó, en donde:

R^{1b} se selecciona de AR1^a, HET-1^a y HET-2^a; en donde R^{1b} se sustituye opcionalmente por 1, ó 2 grupos independientemente seleccionados de R^{3a};

R^{3a} se selecciona de halo, R^{8a}, R^{8a}O-, R^{8a}OCO-, R^{8a}COO-, R^{8a}CO-, -C₆H_(13-a)F_a (en donde a es 1 a 13), -OC₆H_(13-a)F_a (en donde a es 1 a 13), (1-6C)alquiltio, di(1-6C)alquilaminocarbonilo, fenilo, HET-1^b, AR1^b-(1-6C)alcoxi, HET-1^b-(1-6C)alcoxi, en donde cualquier grupo R^{8a}, o (1-6C)alquilo en cualquiera de las anteriores definiciones para R^{3a} puede sustituirse opcionalmente por 1 ó 2 sustituyentes independientemente seleccionados de (1-6C)alcoxi, hidroxilo, carboxi, -C₆H_(13-a)F_a (en donde a es 1 a 13); y/o por 1, 2 ó 3 halo;

R^{2b} es fenilo o 3-piridilo; sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o las dos posiciones *meta* con respecto al enlazamiento con el grupo -Y-NH- (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6d};

R^{4a} se selecciona de:

- a) fenilo;
- b) HET-3^c (en donde cualquier átomo de nitrógeno disponible se sustituye por R^{5a} siempre que, con ello, no se cuaternice);
- d) fenoxi, HET-1^c-oxi, R^{8b}O-, fenil(1-6C)alcoxi y HET-1^c-(1-6C)alcoxi;
- e) R^{8b}, R^{8b}O(1-6C)alquilo, hidroxilo(1-6C)alquilo; y

f) fenilamino, fenil(1-6C)alquilamino, HET-1^c-amino, HET-1^c-(1-6C)alquilamino, (1-6C)alquilamino, N-fenil-N-(1-6C)alquilamino y di(1-6C)alquilamino;

R^{5a} se selecciona de (1-6C)alquilo, fenilo (opcionalmente sustituido con 1 ó 2 halo), HET-1^d, R^{8c}CO-, fenilcarbonilo (en donde el fenilo se sustituye opcionalmente con 1 ó 2 halo), HET-1^d-carbonilo, (1-6C)alcoxycarbonilo, R^{8c}NHCO-, (1-6C)alcoxisulfonilo, (1-6C)alquilaminosulfonilo, y (1-6C)alquilsulfonilo; en donde cualquier grupo (1-6C)alquilo en cualquiera de las definiciones de R^{5a} puede sustituirse opcionalmente por un sustituyente seleccionado de (1-6C)alcoxi, (1-6C)alquilS(O)p'- (en donde p' es 0, 1 ó 2), amino, (1-6C)alquilamino, hidroxilo, carboxilo, fenilo (opcionalmente sustituido con 1 ó 2 halo) y HET-1^f;

R^{6d} se selecciona de R^{8d}, halo y ciano;

R^{8a}, R^{8b}, R^{8c} o R^{8d} es (1-6C)alquilo;

AR1^a o AR1^b es fenilo o un sistema de anillo fusionado 6,5, ó 6,6 bicíclico, en donde los dos anillos son carbocíclicos y en donde por lo menos el anillo enlazado al átomo de nitrógeno enlazante es completamente aromático;

HET-1^a, HET-1^b, HET-1^c, HET-1^d o HET-1^f es un anillo heteroarilo de 5 ó 6 miembros, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO₂, y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo;

HET-2^a es un sistema de anillo heterocíclico aromático o parcialmente saturado, bicíclico 5,5, 5,6, 6,5 ó 6,6 fusionado, que contiene 1, 2, 3 ó 4 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en los cuales el anillo directamente enlazado al átomo de nitrógeno enlazante es aromático, y cualquiera de los dos anillos del sistema de anillo bicíclico puede ser carboxílico;

HET-3^c es un anillo heterocíclico de 5 ó 6 miembros, parcial o completamente saturado, que contiene 1, 2, ó 3 heteroátomos independientemente seleccionados de O, N y S (siempre que no haya enlaces O-O, S-S u O-S), en donde cualquier átomo de azufre puede oxidarse opcionalmente a un grupo SO o SO₂, y en donde cualquier átomo de carbono puede oxidarse opcionalmente a un grupo carbonilo; excepto que HET-3 no es 3-oxomorfolin-4-ilo.

En un aspecto adicional de la invención se provee un compuesto de la fórmula (IB^m) o su sal farmacéuticamente aceptable, que es un compuesto de la fórmula (IB) como ya se indicó, en donde:

R^{1b} se selecciona de AR1^a y HET-2^a; en donde R^{1b} se sustituye opcionalmente por 1 ó 2 grupos independientemente seleccionados de R^{3a};

R^{3a} se selecciona de halo, R^{8a}, R^{8a}O-, R^{8a}OCO-, -C₆H_(3-a)F_a (en donde a es 1 a 13), y (1-6C)alquiltio;

R^{2b} es fenilo o 3-piridilo; sustituido en la posición que es *para* al enlazamiento con el grupo -Y-NH- por un grupo seleccionado de R^{4a} y, de manera opcional, adicionalmente sustituido en un átomo de carbono en una o las dos posiciones que son *meta* al enlazamiento con el grupo -Y-NH- (donde sea posible) por 1 ó 2 grupos independientemente seleccionados de R^{6c};

R^{4a} se selecciona de:

- a) fenilo;
- b) HET-3^c (en donde cualquier átomo de nitrógeno disponible se sustituye por R^{5a} siempre que, con ello, no se cuaternice);
- d) fenoxi, HET-1^c-oxi, R^{8b}O-, fenil(1-6C)alcoxi y HET-1^c-(1-6C)alcoxi;
- e) R^{8b} e hidroxil(1-6C)alquilo; y
- f) fenilamino y N-fenil-N-(1-6C)alquilamino;

R^{5a} se selecciona de (1-6C)alquilo, bencilo, HET-1^d, R^{8c}CO-, fenilcarbonilo (en donde el fenilo se sustituye opcionalmente con 1 ó 2 halo), HET-1^d-carbonilo, (1-6C)alcoxycarbonilo y (1-6C)alquilsulfonilo;

R^{6c} se selecciona de R^{8d}, halo y ciano;

R^{8a} , R^{8b} , R^{8c} y R^{8d} se seleccionan independientemente de (1-6C)alquilo;

ARI^a se selecciona de fenilo, naftilo, indenilo y dihidroindenilo;

HET-1^c o HET-1^d es piridilo;

HET-2^a es 1,3-benzodioxolilo;

HET-3^c se selecciona de morfolino, morfolinilo, piperazinilo y piperidinilo.

En un aspecto adicional de la invención se provee un compuesto de la fórmula (IB^{'''}), o su sal farmacéuticamente aceptable, que es un compuesto de la fórmula (IB) como ya se indicó, en donde

R^{1b} se selecciona de fenilo, 1-naftilo y 2-naftilo, seleccionado opcionalmente con 1, 2 ó 3 fluoro;

R^{2b} es fenilo o piridilo; sustituido como se describe por un grupo R^{4a} y un grupo R^{6c}

R^{6c} es halo, preferiblemente fluoro o cloro;

R^{4a} es morfolino, acilpiperazinilo o fenilo.

En un aspecto adicional de la invención se provee un compuesto de la fórmula (IB^{'''}), o su sal farmacéuticamente aceptable, que es un compuesto de la fórmula (IB) como ya se indicó, en donde

R^{1b} se selecciona de fenilo, 1-naftilo y 2-naftilo, opcionalmente sustituido con 1, 2 ó 3 fluoro;

R^{2b} es fenilo o piridilo; sustituido como se describe por un grupo R^{4a} y un grupo R^{6c}

R^{6c} es halo, preferiblemente fluoro o cloro;

R^{4a} es morfolino o acilpiperazinilo.

Compuestos preferidos de la invención son uno cualquiera de los siguientes, o sus sales farmacéuticamente aceptables y/o sus profármacos:

N-[4-(4-morfolinil)fenil]-5-[[3-(fenilmetoxi)fenil]amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(4-cloro-3-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(4-clorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2,3-dihidro-1H-inden-5-il)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-morfolinil)fenil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

5-[(3,4-difluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

ácido 3-[[5-[[[4-(4-morfolinil)fenil]amino]carbonil]-1,3,4-oxadiazol-2-il]amino]-benzoico, éster etílico;

5-[(3-etilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-clorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-morfolinil)fenil]-5-(2-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

5-[(3-bromofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3,4-dimetilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-cloro-4-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida,

5-[[3-(metiltio)fenil]amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida,

5-[(4-fluoro-3-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida,

5-[(3,5-dimetilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida,

5-[(2,5-difluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida,

5-[(2,3-difluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida,
5-[(2,4-difluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[[4-(1,1-dimetiletil)fenil]amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(4-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
N-[4-(4-morfolinil)fenil]-5-(fenilamino)-1,3,4-oxadiazol-2-carboxamida;
N-[4-(4-morfolinil)fenil]-5-[[3-(trifluorometil)fenil]amino]-1,3,4-oxadiazol-2-carboxamida;
5-[(4-fluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(2-clorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(2-cloro-4-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(2,3-dimetilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(2-fluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(2-metilfenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(3-fluorofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[[2-(1-metiletil)fenil]amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-[(4-metoxifenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
5-(1,3-benzodioxol-5-ilamino)-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-yodofenil)amino]-N-[4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[3-cloro-4-(4-morfolinil)fenil]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[3-fluoro-4-(4-morfolinil)fenil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[3-cloro-4-(4-morfolinil)fenil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

5-[(3,4-difluorofenil)amino]-N-[3-metil-4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[3-metil-4-(4-morfolinil)fenil]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[3-fluoro-4-(4-morfolinil)fenil]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida,

N-[3-cloro-4-(4-morfolinil)fenil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida,

N-[3-fluoro-4-(4-morfolinil)fenil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[3-metil-4-(4-morfolinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[3-ciano-4-(4-morfolinil)fenil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

5-[(2,5-difluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-morfolinil)-3-piridinil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-fluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-metilfenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

- 5-[[4-(1,1-dimetiletil)fenil]amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(3-clorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-morfolinil)-3-piridinil]-5-(2-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;
- N-[5-cloro-6-(4-morfolinil)-3-piridinil]-5-[(3,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(4-clorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-[5-cloro-6-(4-morfolinil)-3-piridinil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(3,4-difluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(3-bromofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2,3-difluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-[5-cloro-6-(4-morfolinil)-3-piridinil]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(4-metilfenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(4-fluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-morfolinil)-3-piridinil]-5-(fenilamino)-1,3,4-oxadiazol-2-carboxamida;
- 5-(1,3-benzodioxol-5-ilamino)-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(3-metoxifenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2,4-difluorofenil)amino]-N-[6-(4-morfolinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetil-1-piperazinil)fenil]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetil-1-piperazinil)fenil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-metil-1-piperazinil)fenil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetil-1-piperazinil)fenil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetil-1-piperazinil)-3-metilfenil]-5-[(3,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-[4-(1-metiletil)-1-piperazinil]fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-metilfenil)amino]-N-[4-(4-metil-1-piperazinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetil-1-piperazinil)-3-metilfenil]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

ácido 4-[2-fluoro-4-[[[5-(1-naftalenilamino)-1,3,4-oxadiazol-2-il]carbonil]amino]fenil]-1-piperazinacarboxílico, 1,1-dimetiletil éster;

5-[(2-fluorofenil)amino]-N-[4-[4-(fenilmetil)-1-piperazinil]fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetil-1-piperazinil)-3-metilfenil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

ácido 4-[2-fluoro-4-[[[5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il]carbonil]amino]fenil]-1-piperazinacarboxílico, 1,1-dimetiletil éster;

5-[(2-fluorofenil)amino]-N-[4-(4-metil-1-piperazinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

ácido 4-[4-[[[5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il]carbonil]amino]fenil]-1-piperazinacarboxílico, 1,1-dimetiletil éster;

5-[(2-fluorofenil)amino]-N-[4-[4-(4-piridinil)-1-piperazinil]fenil]-1,3,4-oxadiazol-2-carboxamida;

5-(1-naftalenilamino)-N-[4-(1-piperidinil)fenil]-1,3,4-oxadiazol-2-carboxamida;

- 5-[(3-metilfenil)amino]-N-[4-(1-piperidinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-[4-(1-piperidinil)fenil]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-metil-1-piperazinil)-3-piridinil]-5-(1-naftalenilamino)-1,3,4-oxadiazol-2-carboxamida;
- 5-[(3-metilfenil)amino]-N-[6-(4-metil-1-piperazinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-benzoil-1-piperazinil)-3-piridinil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-[4-(3-fluorobenzoil)-1-piperazinil]-3-piridinil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-[6-[4-(metilsulfonyl)-1-piperazinil]-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-[4-(4-fluorobenzoil)-1-piperazinil]-3-piridinil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-[6-(4-metil-1-piperazinil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- ácido 4-[5-[[[5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il]carbonil]amino]-2-piridinil]-1-piperazinacarboxílico, 1,1-dimetiletil éster;
- N-[6-(4-acetil-1-piperazinil)-3-piridinil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-[6-[4-(4-piridinilcarbonil)-1-piperazinil]-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-(6-fenoxi-3-piridinil)-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-[6-(fenilamino)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;
- N-(6-butoxi-3-piridinil)-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(2-fluorofenil)amino]-N-[6-(3-piridiniloxi)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[6-(4-hidroxi)butil]-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(fenilmetoxi)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-(4-butilfenil)-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(2-piridinilmetoxi)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(metilfenilamino)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[1,1'-bifenil]-4-il-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida; y/o

metil {*trans*-4-[4-({5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil [*trans*-4-(4-({5-([3-(benciloxi)fenil]amino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(2,4,5-trifluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil *trans*-4-[4-({5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexanocarboxilato;

metil (*trans*-4-[4-({5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

ácido {*trans*-4-[4-({5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido {*trans*-4-[4-({5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido [*trans*-4-(4-({5-([3-(benciloxi)fenil]amino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

5-[(2-fluorofenil)amino]-N-[6-(4-hidroxibutil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(fenilmetoxi)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-(4-butilfenil)-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(2-piridinilmetoxi)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(metilfenilamino)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[1,1'-bifenil]-4-il-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida; y/o

metil {*trans*-4-[4-({5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil [*trans*-4-(4-({5-([3-(benciloxi)fenil]amino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(2,4,5-trifluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil *trans*-4-[4-({5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexanocarboxilato;

metil (*trans*-4-[4-({5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

ácido {*trans*-4-[4-({5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido {*trans*-4-[4-({5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido [*trans*-4-(4-({5-([3-(benciloxi)fenil]amino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

5-[(2-fluorofenil)amino]-N-[6-(4-hidroxibutil)-3-piridinil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(fenilmetoxi)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-(4-butilfenil)-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(2-piridinilmetoxi)fenil]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-[4-(metilfenilamino)fenil]-1,3,4-oxadiazol-2-carboxamida;

N-[1,1'-bifenil]-4-il-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida; y/o

metil {*trans*-4-[4-({5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil [*trans*-4-(4-({5-([3-(benciloxi)fenil]amino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil (*trans*-4-[4-({5-[(2,4,5-trifluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

metil *trans*-4-[4-({5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexanocarboxilato;

metil (*trans*-4-[4-({5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acetato;

ácido {*trans*-4-[4-({5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido {*trans*-4-[4-({5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido [*trans*-4-(4-({5-([3-(benciloxi)fenil]amino)-1,3,4-oxadiazol-2-il} carbonil) amino}fenil}ciclohexil}acético;

ácido (*trans*-4-{4-[(5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-il) carbonil) amino]fenil}ciclohexil)acético;

(*trans*-4-{4-[(5-[(2,4,5-trifluorofenil)amino]-1,3,4-oxadiazol-2-il) carbonil) amino]fenil}ciclohexil)acetato de sodio;

ácido *trans*-4-{4-[(5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il) carbonil) amino] fenil}ciclohexanocarboxílico;

ácido (*trans*-4-{4-[(5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-il) carbonil) amino]fenil}ciclohexil)acético;

5-[(2-fluorofenil)amino]-N-{4-[4-(2-hidroxietil)ciclohexil]fenil}-1,3,4-oxadiazol-2-carboxamida;

N-{4-[*trans*-4-(2-amino-2-oxoetil)ciclohexil]fenil}-5-[(3-etoxifenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(2-fluorofenil)amino]-N-(3-fenoxipropil)-1,3,4-oxadiazol-2-carboxamida;

5-[(3-metilfenil)amino]-N-(3-fenoxipropil)-1,3,4-oxadiazol-2-carboxamida;

5-[(3-(benciloxi)fenil)amino]-N-(3-fenoxipropil)-1,3,4-oxadiazol-2-carboxamida;

N-(4-ciclohexilfenil)-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-(4-ciclohexilfenil)-5-[(3,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(3-(benciloxi)fenil)amino]-N-(4-ciclohexilfenil)-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(3-isopropoxifenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(4-(metiltio)fenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(4-(benciloxi)fenil)amino]-1,3,4-oxadiazol-2-carboxamida;

5-[(5-bromo-2,4-difluorofenil)amino]-N-(4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;

5-{{2,4-difluoro-5-(trifluorometil)fenil}amino}-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

N-(6-morfolin-4-ilpiridin-3-il)-5-{{3-(trifluorometil)fenil}amino}-1,3,4-oxadiazol-2-carboxamida;

5-{{4-metilpiridin-2-il}amino}-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

5-{{2-fluoro-5-(trifluorometil)fenil}amino}-N-(4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;

5-{{4-fluoro-3-(trifluorometil)fenil}amino}-N-(4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;

5-{{3-isopropoxifenil}amino}-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

N-(6-morfolin-4-ilpiridin-3-il)-5-{{3-fenoxifenil}amino}-1,3,4-oxadiazol-2-carboxamida;

N-(6-morfolin-4-ilpiridin-3-il)-5-{{3-(trifluorometoxi)fenil}amino}-1,3,4-oxadiazol-2-carboxamida;

N-(6-morfolin-4-ilpiridin-3-il)-5-{{3-(piridin-2-ilmetoxi)fenil}amino}-1,3,4-oxadiazol-2-carboxamida;

5-{{3-etoxifenil}amino}-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

N-{{4-(4-acetilpiperazin-1-il)fenil}-5-{{3-etoxifenil}amino}-1,3,4-oxadiazol-2-carboxamida;

5-(bifenil-3-ilamino)-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

5-{{3-(benciloxi)fenil}amino}-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

etil 3-{{5-{{(6-morfolin-4-ilpiridin-3-il)amino}carbonil}-1,3,4-oxadiazol-2-il}amino}benzoato;

terc-butil 3-{{5-{{(6-morfolin-4-ilpiridin-3-il)amino}carbonil}-1,3,4-oxadiazol-2-il}amino}benzoato;

5-{{3-yodofenil}amino}-N-(6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;

N-(6-metoxipiridin-3-il)-5-(1-naftilamino)-1,3,4-oxadiazol-2-carboxamida;
N-(3-fluoro-4-morfolin-4-ilfenil)-5-[(4-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
5-[(2,5-difluorofenil)amino]-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
5-anilino-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
N-(3-fluoro-4-morfolin-4-ilfenil)-5-[(3-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
N-(3-fluoro-4-morfolin-4-ilfenil)-5-[(4-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;
5-[(2,4-difluorofenil)amino]-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
5-[(3-bromofenil)amino]-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
5-[(4-clorofenil)amino]-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
N-(3-fluoro-4-morfolin-4-ilfenil)-5-(2-naftilamino)-1,3,4-oxadiazol-2-carboxamida; 5-[(3,4-difluorofenil)amino]-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
5-[(2,3-difluorofenil)amino]-*N*-(3-fluoro-4-morfolin-4-ilfenil)-1,3,4-oxadiazol-2-carboxamida;
etil 3-[(5-[(3-fluoro-4-morfolin-4-ilfenil)amino]carbonil)-1,3,4-oxadiazol-2-il)amino]benzoato;
N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(4-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(2-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;
N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(3-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(2-metoxifenil)amino]-1,3,4-oxadiazol-2-carboxamida;

- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(4-metoxifenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(2,5-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-anilino-N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;
- 5-[(3-bromofenil)amino]-N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(3-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(4-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-(2,6-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(2-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-(2-naftilamino)-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(3-metoxifenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- 5-[(4-*terc*-butilfenil)amino]-N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-1,3,4-oxadiazol-2-carboxamida;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-[(2,3-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- etil 3-[(5-[(5-cloro-6-morfolin-4-ilpiridin-3-il)amino]carbonil)-1,3,4-oxadiazol-2-il)amino]benzoato;
- N-(5-cloro-6-morfolin-4-ilpiridin-3-il)-5-(1-naftilamino)-1,3,4-oxadiazol-2-carboxamida;
- N-{4-[cis-2,6-dimetilmorfolin-4-il]fenil}-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

- 5-[(3,4-difluorofenil)amino]-N-{4-[cis-2,6-dimetilmorfolin-4-il]fenil}-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-[(2-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-[(3-metil)fenil]amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-[(3,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-[[3-(benciloxi)fenil] amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-(1-naftilamino)-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-[(2,4,5-trifluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[4-(4-acetil-1,4-diazepan-1-il)fenil]-5-[(2,5-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[2-(4-acetilpiperazin-1-il)pirimidin-5-il]-5-[[3-(benciloxi)fenil] amino]-1,3,4-oxadiazol-2-carboxamida;
- N-(2-metil-1,3-benzotiazol-5-il)-5-(1-naftilamino)-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(4-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(3-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(2,5-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-anilino-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(3-bromofenil)amino]-1,3,4-oxadiazol-2-carboxamida;
- N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(3-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(4-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(2-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-(2-naftilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-(piridin-2-ilamino)-1,3,4-oxadiazol-2-carboxamida;

etil 4-{{5-({[6-(4-acetilpiperazin-1-il)piridin-3-il]amino} carbonil)-1,3,4-oxadiazol-2-il}amino} benzoato;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-(1,3-benzodioxol-5-ilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(2,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

etil 3-{{5-({[6-(4-acetilpiperazin-1-il)piridin-3-il]amino} carbonil)-1,3,4-oxadiazol-2-il}amino} benzoato;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(4-*terc*-butilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(3,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(2,3-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(3-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-{{3-(benciloxi)fenil} amino}-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(3,5-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-(1-naftilamino)-1,3,4-oxadiazol-2-carboxamida;



N-[6-(4-acetilpiperazin-1-il)piridin-3-il]-5-[(2,4,5-trifluorofenil) amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(4-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(3-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(2,5-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-anilino-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(3-bromofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(3-fluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(4-metilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(2-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(4-clorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-(2-naftilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-(1,3-benzodioxol-5-ilamino)-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(2,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;

etil-3-{{5-([4-(4-acetilpiperazin-1-il)fenil] amino) carbonil}-1,3,4-oxadiazol-2-il}amino} benzoato;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(4-terc-butilfenil)amino]-1,3,4-oxadiazol-2-carboxamida;

N-[4-(4-acetilpiperazin-1-il)fenil]-5-[(3,4-difluorofenil)amino]-1,3,4-oxadiazol-2-carboxamida;