

CASTELLANOS & CO LTDA

ABOGADOS
Calle 94 A No. 7 A-09
Bogotá, D.C. – COLOMBIA

SUPERINTENDENCIA DE INDUSTRIA Y COMERCIO



No. 14-036579- -00001-0000

Fecha: 2014-02-25 16:02:57 Dep. 2020 DIR.NUEVASCR
Tra. 11 PATENTEIYII Eve: 378 FASENACIONALI
Act. 329 CTOINFORMACION Folios: 383

Señor

Director de la Dirección de Nuevas Creaciones

Superintendencia de Industria y Comercio

Ministerio de Comercio, Industria y Turismo

E.

S.

D.

Re.: Expediente No. 14 036579

Solicitud de Patente PCT entrada en Fase Nacional, para:

“DERIVADOS DE ISOXAZOLINA COMO COMPUESTOS INSECTICIDAS”

Solicitante: **SYNGENTA PARTICIPATIONS AG**

MEMORIAL COMPLEMENTO INFORMACION

Margarita Castellanos Abondano, obrando en mi calidad de apoderada de la sociedad solicitante de la solicitud de Patente citada en la referencia, atentamente me permito presentar la copia de Publicación Internacional No. **PCT/EP2012/066554 de 24/08/2012 (tanto física como en CD)**, esto debido a que al momento de radicar la solicitud en línea no permitió anexar dicho documento, en razón al tamaño del mismo.

En consecuencia, atentamente solicito a Ud. se anexe dicho documento al expediente citado en la referencia, y se continúe con el trámite correspondiente a esta solicitud de Patente.

Del Señor Director,

Margarita Castellanos Abondano
C.C. No. **39.779.654** de Bogotá
T.P.A. No. **70.819** de C.S.J.
Calle 94 A No. 7 A-09 – Bogotá, D.C.
Tel.: 7560770

Bogotá, D.C. **25 FEB. 2014**
MCA/gcb.
P3310
MEMORIAL COMPLEMENTO INFORMACION

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

28 February 2013 (28.02.2013)



(10) International Publication Number

WO 2013/026931 A1

(51) International Patent Classification:

C07D 331/04 (2006.01) A01N 43/36 (2006.01)

C07D 409/12 (2006.01) A01N 43/80 (2006.01)

C07D 413/12 (2006.01)

(21) International Application Number:

PCT/EP2012/066554

(22) International Filing Date:

24 August 2012 (24.08.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11178921.0 25 August 2011 (25.08.2011) EP

12179257.6 3 August 2012 (03.08.2012) EP

12179385.5 6 August 2012 (06.08.2012) EP

PCT/EP2012/065421 7 August 2012 (07.08.2012) EP

(71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CASSAYRE, Jérôme, Yves [FR/CH]; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH). RENOLD, Peter [CH/CH]; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein

(CH). PITTERNA, Thomas [AT/CH]; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH). EL QACEMI, Myriem [FR/CH]; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH).


(74) Agent: THWAITE, Jonathan; Syngenta International AG, WRO 1004-6-22, Schwarzwaldallee 215, CH-4058 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

[Continued on next page]

(54) Title: ISOXAZOLINE DERIVATIVES AS INSECTICIDAL COMPOUNDS



(57) Abstract: The present invention provides compounds of formula (I), wherein G¹ is oxygen; R¹ is hydrogen; R² is group P (P) L is a bond, methylene or ethylene; one of A¹ and A² is S, SO or SO₂ and the other is -C(R⁴)R⁴-R³ is hydrogen or methyl; each R⁴ is independently hydrogen or methyl; Y¹ is C-R⁶, CH or nitrogen; Y² and Y³ are independently CH or nitrogen; wherein no more than two of Y¹, Y² and Y³ are nitrogen and wherein Y² and Y³ are not both nitrogen; R⁵ is hydrogen, halogen, cyano, nitro, NH₂, C₁-C₂alkyl, C₁-C₂haloalkyl, C₃-C₃cycloalkyl, C₃-C₃halocycloalkyl, C₁-C₃alkoxy, C₁-C₂haloalkoxy; R⁶ together with R⁵ forms a -CH=CH-CH=CH- bridge; X² is C-X⁶ or nitrogen; X¹, X³ and X⁶ are independently hydrogen, halogen or trihalomethyl, wherein at least two of X¹, X³ and X⁶ are not hydrogen; X⁴ is trifluoromethyl, difluoromethyl or chlorodifluoromethyl. The invention also provides intermediates useful for the preparation of compounds of formula (I), as well as methods of controlling insects, acarines, nematodes or molluscs using the compounds of formula (I).

WO 2013/026931 A1

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG). **Published:** — with international search report (Art. 21(3))

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))


ISOXAZOLINE DERIVATIVES AS INSECTICIDAL COMPOUNDS

The present invention relates to certain isoxazolines derivatives, to processes and intermediates for preparing these derivatives, to insecticidal, acaricidal, nematocidal and molluscicidal compositions comprising these derivatives and to methods of using these derivatives to control insect, acarine, nematode and mollusc pests. The present invention also relates to use of these compounds in the field of animal health.

Certain isoxazoline derivatives with insecticidal properties are disclosed, for example, in WO2009/08050.

There is an ongoing need to find compounds that exhibit higher levels of activity, broader spectrum of control, higher levels of activity against specific pests, increased safety, reduced impact on the environment. The present invention seeks to address these needs at least in part.

The present invention provides compounds of formula I




wherein

15 G^1 is oxygen;

R^1 is hydrogen;

R^2 is group P



20 L is a bond, methylene or ethylene;

one of A^1 and A^2 is S, SO or SO_2 and the other is $-C(R^4)R^4$;

R^3 is hydrogen or methyl;

each R^4 is independently hydrogen or methyl;

Y^1 , Y^2 and Y^3 are independently CH or nitrogen;

25 wherein no more than two of Y^1 , Y^2 and Y^3 are nitrogen and wherein Y^2 and Y^3 are not both nitrogen;

R^5 is hydrogen, halogen, cyano, nitro, NH_2 , C_1 - C_2 alkyl, C_1 - C_2 haloalkyl, C_3 - C_5 cycloalkyl, C_1 - C_2 halocycloalkyl, C_1 - C_2 alkoxy, C_1 - C_2 haloalkoxy;

X^2 is $C-X^6$ or nitrogen;

X¹, X³ and X⁶ are independently hydrogen, halogen or trihalomethyl, wherein at least two of X¹, X³ and X⁶ are not hydrogen;

X⁴ is trifluoromethyl, difluoromethyl or chlorodifluoromethyl.

5 The compounds of formula (I) may exist in different geometric or optical isomers or tautomeric forms. This invention covers all such isomers and tautomers and mixtures thereof in all proportions as well as isotopic forms such as deuterated compounds. The invention also covers salts and N-oxides of the compounds of the invention.

The compounds of the invention may contain one or more asymmetric carbon atoms, and may
10 exist as enantiomers (or as pairs of diastereoisomers) or as mixtures of such.

Alkyl groups (either alone or as part of a larger group, such as alkoxy-, alkylthio-, alkylsulfinyl-, alkylsulfonyl-, alkylcarbonyl- or alkoxy carbonyl-) can be in the form of a straight or branched chain and are, for example, methyl, ethyl, propyl, prop-2-yl, butyl, but-2-yl, 2-methyl-prop-1-yl or 2-methyl-prop-2-yl. The alkyl groups are preferably C₁-C₆, more preferably C₁-C₄, most preferably C₁-C₃ alkyl groups.

15 Where an alkyl moiety is said to be substituted, the alkyl moiety is preferably substituted by one to four substituents, most preferably by one to three substituents.

Alkylene groups can be in the form of a straight or branched chain and are, for example, -CH₂-, -CH₂-CH₂-, -CH(CH₃)-, -CH₂-CH₂-CH₂-, -CH(CH₃)-CH₂-, or -CH(CH₂CH₃)-. The alkylene groups are preferably C₁-C₃, more preferably C₁-C₂, most preferably C₁ alkylene groups.

20 Alkenyl groups can be in the form of straight or branched chains, and can be, where appropriate, of either the (E)- or (Z)-configuration. Examples are vinyl and allyl. The alkenyl groups are preferably C₂-C₆, more preferably C₂-C₄, most preferably C₂-C₃ alkenyl groups.

Alkynyl groups can be in the form of straight or branched chains. Examples are ethynyl and propargyl. The alkynyl groups are preferably C₂-C₆, more preferably C₂-C₄, most preferably C₂-C₃
25 alkynyl groups.

Halogen is fluorine, chlorine, bromine or iodine.

Haloalkyl groups (either alone or as part of a larger group, such as haloalkoxy-, haloalkylthio-, haloalkylsulfinyl- or haloalkylsulfonyl-) are alkyl groups which are substituted by one or more of the same or different halogen atoms and are, for example, difluoromethyl, trifluoromethyl,
30 chlorodifluoromethyl or 2,2,2-trifluoro-ethyl.

Haloalkenyl groups are alkenyl groups which are substituted by one or more of the same or different halogen atoms and are, for example, 2,2-difluoro-vinyl or 1,2-dichloro-2-fluoro-vinyl.

Haloalkynyl groups are alkynyl groups which are substituted by one or more of the same or different halogen atoms and are, for example, 1-chloro-prop-2-ynyl.

35 Cycloalkyl groups or carbocyclic rings can be in mono- or bi-cyclic form and are, for example, cyclopropyl, cyclobutyl, cyclohexyl and bicyclo[2.2.1]heptan-2-yl. The cycloalkyl groups are preferably C₃-C₈, more preferably C₃-C₆ cycloalkyl groups. Where a cycloalkyl moiety is said to be substituted, the

cycloalkyl moiety is preferably substituted by one to four substituents, most preferably by one to three substituents.

Aryl groups (either alone or as part of a larger group, such as aryl-alkylene-) are aromatic ring systems which can be in mono-, bi- or tricyclic form. Examples of such rings include phenyl, naphthyl, anthracenyl, indenyl or phenanthrenyl. Preferred aryl groups are phenyl and naphthyl, phenyl being most preferred. Where an aryl moiety is said to be substituted, the aryl moiety is preferably substituted by one to four substituents, most preferably by one to three substituents.

Heteroaryl groups (either alone or as part of a larger group, such as heteroaryl-alkylene-) are aromatic ring systems containing at least one heteroatom and consisting either of a single ring or of two or more fused rings. Preferably, single rings will contain up to three heteroatoms and bicyclic systems up to four heteroatoms which will preferably be chosen from nitrogen, oxygen and sulfur. Examples of monocyclic groups include pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl (e.g. 1,2,4 triazolyl), furanyl, thiophenyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl and thiadiazolyl. Examples of bicyclic groups include purinyl, quinolinyl, cinnolinyl, quinoxalinyl, indolyl, indazolyl, benzimidazolyl, benzothiophenyl and benzothiazolyl. Monocyclic heteroaryl groups are preferred, pyridyl being most preferred. Where a heteroaryl moiety is said to be substituted, the heteroaryl moiety is preferably substituted by one to four substituents, most preferably by one to three substituents.

Heterocyclyl groups or heterocyclic rings (either alone or as part of a larger group, such as heterocyclyl-alkylene-) are defined to include heteroaryl groups and in addition their unsaturated or partially unsaturated analogues. Examples of monocyclic groups include isoxazolyl, thietanyl, pyrrolidinyl, tetrahydrofuranyl, [1,3]dioxolanyl, piperidinyl, piperazinyl, [1,4]dioxanyl, and morpholinyl or their oxidised versions such as 1-oxo-thietanyl and 1,1-dioxo-thietanyl. Examples of bicyclic groups include 2,3-dihydro-benzofuranyl, benzo[1,4]dioxolanyl, benzo[1,3]dioxolanyl, chromenyl, and 2,3-dihydro-benzo[1,4]dioxinyl. Where a heterocyclyl moiety is said to be substituted, the heterocyclyl moiety is preferably substituted by one to four substituents, most preferably by one to three substituents.

Preferred values of R^2 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 , X^4 , X^5 and X^6 are, in any combination, as set out below.

Preferably R^2 is thietan-3-yl-, 1-oxo-thietan-3-yl-, 1,1-Dioxo-thietan-3-yl-, 2,2-dimethylthietan-3-yl-, 2,2-dimethyl-1-oxo-thietan-3-yl-, 2,2-dimethyl-1,1-Dioxo-thietan-3-yl-, 3-Methyl-thietan-3-yl-, 3-Methyl-1-oxo-thietan-3-yl-, 3-Methyl-1,1-Dioxo-thietan-3-yl-, thietan-3-ylmethyl-, 1-oxo-thietan-3-ylmethyl-, 1,1-Dioxo-thietan-3-ylmethyl-, thietan-2-ylmethyl-, (1-oxothietan-2-yl)methyl-, (1,1-dioxothietan-2-yl)methyl-, 2-(thietan-3-yl)ethanyl, 2-(1,1-dioxothietan-3-yl)ethanyl, or 2-(1-oxothietan-3-yl)ethanyl more preferably R^2 is 2-(thietan-3-yl)ethanyl, 2-(1,1-dioxothietan-3-yl)ethanyl, 2-(1-oxothietan-3-yl)ethanyl, thietan-3-ylmethyl-, 1-oxo-thietan-3-ylmethyl-, 1,1-Dioxo-thietan-3-ylmethyl-, thietan-3-yl-, 1-oxo-thietan-3-yl-, 1,1-Dioxo-thietan-3-yl-.

Preferably R^5 is hydrogen, chloro, bromo, fluoro, trifluoromethyl, methyl, ethyl, methoxy, nitro, trifluoromethoxy, cyano, cyclopropyl, more preferably R^5 is hydrogen, chloro, bromo, fluoro, trifluoromethyl, methyl, ethyl, nitro, cyano, cyclopropyl, even more preferably R^5 is chloro, bromo, fluoro, methyl, trifluoromethyl.


- 5 Preferably Y^1 is CH, Y^2 is CH, Y^3 is CH, or Y^1 is N, Y^2 is CH, Y^3 is CH, or Y^1 is N, Y^2 is N, Y^3 is CH, or Y^1 is CH, Y^2 is N, Y^3 is CH, or Y^1 is CH, Y^2 is CH, Y^3 is N.

Preferably X^1 is chloro, X^2 is CH, X^3 is chloro, or X^1 is chloro, X^2 is C-F, X^3 is hydrogen, or X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen, or X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen, or X^1 is chloro, X^2 is C-Br, X^3 is chloro, or X^1 is chloro, X^2 is C-F, X^3 is chloro, or X^1 is chloro, X^2 is C-Cl, X^3 is chloro, or X^1 is
 10 chloro, X^2 is C-I, X^3 is chloro, or X^1 is fluoro, X^2 is C-F, X^3 is fluoro, or X^1 is chloro, X^2 is CH, X^3 is bromo, or X^1 is chloro, X^2 is CH, X^3 is fluoro, or X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl, or X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, or X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, or X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, or X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, or X^1 is chloro, X^2 is N, X^3 is chloro, or X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl.

15

The invention also relates to the following intermediates which are useful for the preparation of compounds of formula I:


Compounds of formula X1



20

wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I. Preferred definitions of G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are the same as for the corresponding substituents of compounds of formula I.

Compound of formula X2




wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I.

Preferred definitions of G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are the same as for the

5 corresponding substituents of compounds of formula I.


Compound of formula X3



wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I.

10 Preferred definitions of G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are the same as for the corresponding substituents of compounds of formula I.


Compound of formula X4



15 wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I.

Preferred definitions of G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are the same as for the corresponding substituents of compounds of formula I.

Compound of formula X5



wherein G^1 , R^1 , R^2 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I and X^5 is chloro, bromo, iodo, amino, CHO, CN, OH, C(=O)OH, C(=NOH)H, C(=NOH)Cl, C(=NOH)NH₂, C(=O)CH₃,

- 5 C(=NOH)CH₃, C(=O)CH₂Cl, or C(=O)CH₂Br. Preferred definitions of G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , and Y^3 are the same as for the corresponding substituents of compounds of formula I.

The following embodiments apply to intermediates X1, X2, X3, X4 and X5 as well as compounds of formula I. Embodiments may be combined where possible.

10

In embodiment A1 A^1 is S and A^2 is CH₂.

In embodiment A2 A^1 is SO and A^2 is CH₂.

In embodiment A3 A^1 is SO₂ and A^2 is CH₂.

In embodiment A4 A^1 is CH₂ and A^2 is S.

- 15 In embodiment A5 A^1 is CH₂ and A^2 is SO.

In embodiment A6 A^1 is CH₂ and A^2 is SO₂.

In embodiment B1 A^1 is S and A^2 is CH₂ and L is a bond.

In embodiment B2 A^1 is SO and A^2 is CH₂ and L is a bond.

- 20 In embodiment B3 A^1 is SO₂ and A^2 is CH₂ and L is a bond.

In embodiment B4 A^1 is CH₂ and A^2 is S and L is a bond.

In embodiment B5 A^1 is CH₂ and A^2 is SO and L is a bond.

In embodiment B6 A^1 is CH₂ and A^2 is SO₂ and L is a bond.

In embodiment B7 A^1 is S and A^2 is CH₂ and L is methylene.

- 25 In embodiment B8 A^1 is SO and A^2 is CH₂ and L is methylene.

In embodiment B9 A^1 is SO₂ and A^2 is CH₂ and L is methylene.

In embodiment B10 A^1 is CH₂ and A^2 is S and L is methylene.

In embodiment B11 A^1 is CH₂ and A^2 is SO and L is methylene.

In embodiment B12 A^1 is CH₂ and A^2 is SO₂ and L is methylene.

- 30 In embodiment B13 A^1 is S and A^2 is CH₂ and L is ethylene.

In embodiment B14 A^1 is SO and A^2 is CH₂ and L is ethylene.

In embodiment B15 A^1 is SO₂ and A^2 is CH₂ and L is ethylene.

In embodiment B16 A^1 is CH₂ and A^2 is S and L is ethylene.

In embodiment B17 A^1 is CH₂ and A^2 is SO and L is ethylene.

- 35 In embodiment B18 A^1 is CH₂ and A^2 is SO₂ and L is ethylene.

In embodiment C1 the invention provides compounds of formula I wherein X^2 is $C-X^6$.

- 5 In embodiment C2 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H.

In embodiment C3 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H and each R^4 is hydrogen.

- In embodiment C4 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are
10 C-H, R^3 is hydrogen and each R^4 is hydrogen.

In embodiment D1 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H, R^3 is hydrogen and each R^4 is hydrogen, and A^1 is S and A^2 is CH_2 .

- In embodiment D2 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are
15 C-H, R^3 is hydrogen and each R^4 is hydrogen, and A^1 is SO and A^2 is CH_2 .

In embodiment D3 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H, R^3 is hydrogen and each R^4 is hydrogen, and A^1 is SO_2 and A^2 is CH_2 .

In embodiment D4 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H, R^3 is hydrogen and each R^4 is hydrogen, and A^1 is CH_2 and A^2 is S.

- 20 In embodiment D5 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H, R^3 is hydrogen and each R^4 is hydrogen, and A^1 is CH_2 and A^2 is SO.

In embodiment D6 the invention provides compounds of formula I wherein X^2 is $C-X^6$, Y^1 , Y^2 and Y^3 are C-H, R^3 is hydrogen and each R^4 is hydrogen, and A^1 is CH_2 and A^2 is SO_2 .

- 25 In embodiment E1 the invention provides compounds of formula I wherein R^2 is thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

In embodiment E2 the invention provides compounds of formula I wherein R^2 is 1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

- In embodiment E3 the invention provides compounds of formula I wherein R^2 is 1,1-Dioxo-thietan-3-yl-
30 and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

In embodiment E4 the invention provides compounds of formula I wherein R^2 is 2,2-dimethylthietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

In embodiment E5 the invention provides compounds of formula I wherein R^2 is 2,2-dimethyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

- 35 In embodiment E6 the invention provides compounds of formula I wherein R^2 is 2,2-dimethyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

In embodiment E7 the invention provides compounds of formula I wherein R^2 is 3-Methyl-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.

In embodiment E8 the invention provides compounds of formula I wherein R² is 3-Methyl-1-oxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E9 the invention provides compounds of formula I wherein R² is 3-Methyl-1,1-Dioxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

5 In embodiment E10 the invention provides compounds of formula I wherein R² is thietan-3-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E11 the invention provides compounds of formula I wherein R² is 1-oxo-thietan-3-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E12 the invention provides compounds of formula I wherein R² is 1,1-Dioxo-thietan-3-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

10 In embodiment E13 the invention provides compounds of formula I wherein R² is thietan-2-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E14 the invention provides compounds of formula I wherein R² is (1-oxothietan-2-yl)methyl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

15 In embodiment E15 the invention provides compounds of formula I wherein R² is (1,1-dioxothietan-2-yl)methyl- and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E16 the invention provides compounds of formula I wherein R² is 2-(thietan-3-yl)ethanyl and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E17 the invention provides compounds of formula I wherein R² is 2-(1,1-dioxothietan-3-yl)ethanyl and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E18 the invention provides compounds of formula I wherein R² is 2-(1-oxothietan-3-yl)ethanyl and G¹, R¹, R⁵, Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E19 the invention provides compounds of formula I wherein Y¹ is CH, Y² is CH, Y³ is CH and G¹, R¹, R², R⁵, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

25 In embodiment E20 the invention provides compounds of formula I wherein Y¹ is N, Y² is CH, Y³ is CH and G¹, R¹, R², R⁵, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E21 the invention provides compounds of formula I wherein Y¹ is N, Y² is N, Y³ is CH and G¹, R¹, R², R⁵, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E22 the invention provides compounds of formula I wherein Y¹ is CH, Y² is N, Y³ is CH and G¹, R¹, R², R⁵, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

30 In embodiment E23 the invention provides compounds of formula I wherein Y¹ is CH, Y² is CH, Y³ is N and G¹, R¹, R², R⁵, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E24 the invention provides compounds of formula I wherein R⁵ is hydrogen and G¹, R¹, R², Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

35 In embodiment E25 the invention provides compounds of formula I wherein R⁵ is chloro and G¹, R¹, R², Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

In embodiment E26 the invention provides compounds of formula I wherein R⁵ is bromo and G¹, R¹, R², Y¹, Y², Y³, X¹, X², X³ and X⁴ are as defined for the compound of formula I.

- In embodiment E27 the invention provides compounds of formula I wherein R^5 is fluoro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- In embodiment E28 the invention provides compounds of formula I wherein R^5 is trifluoromethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- 5 In embodiment E29 the invention provides compounds of formula I wherein R^5 is methyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- In embodiment E30 the invention provides compounds of formula I wherein R^5 is ethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- In embodiment E31 the invention provides compounds of formula I wherein R^5 is methoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- 10 In embodiment E32 the invention provides compounds of formula I wherein R^5 is nitro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- In embodiment E33 the invention provides compounds of formula I wherein R^5 is trifluoromethoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- 15 In embodiment E34 the invention provides compounds of formula I wherein R^5 is cyano and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- In embodiment E35 the invention provides compounds of formula I wherein R^5 is cyclopropyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- In embodiment E36 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is chloro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , X^3 and X^4 are as defined for the compound of formula I.
- 20 In embodiment E37 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is hydrogen and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- In embodiment E38 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- 25 In embodiment E39 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- In embodiment E40 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Br, X^3 is chloro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- In embodiment E41 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is chloro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- 30 In embodiment E42 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is chloro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- In embodiment E43 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-I, X^3 is chloro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- 35 In embodiment E44 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-F, X^3 is fluoro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.
- In embodiment E45 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment E46 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is fluoro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment E47 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 5 In embodiment E48 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment E49 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 10 In embodiment E50 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment E51 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 15 In embodiment E52 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is N, X^3 is chloro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment E53 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , and X^3 are as defined for the compound of formula I.

- 20 In embodiment E54 the invention provides compounds of formula I wherein X^4 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , and X^3 are as defined for the compound of formula I.

In embodiment E55 the invention provides compounds of formula I wherein X^4 is difluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , and X^3 are as defined for the compound of formula I.

In embodiment E56 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl

- 25 and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 , and X^3 are as defined for the compound of formula I.

In embodiment F1 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

- 30 In embodiment F2 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F3 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the

- 35 compound of formula I.

In embodiment F4 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 2,2-dimethylthietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F5 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 2,2-dimethyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F6 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 2,2-dimethyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F7 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 3-Methyl-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

10 In embodiment F8 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 3-Methyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F9 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 3-Methyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for
15 the compound of formula I.

In embodiment F10 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F11 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 1-oxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the
20 compound of formula I.

In embodiment F12 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 1,1-Dioxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

25 In embodiment F13 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is thietan-2-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F14 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is (1-oxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the
30 compound of formula I.

In embodiment F15 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is (1,1-dioxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F16 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 2-(thietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the
35 compound of formula I.

In embodiment F17 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 2-(1,1-dioxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F18 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^2 is 2-(1-oxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F19 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

10 In embodiment F20 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F21 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , X^1 , X^2 and X^3 are as defined for the compound of
15 formula I.

In embodiment F22 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F23 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 , X^1 , X^2 and X^3 are as defined for the compound of
20 formula I.

In embodiment F24 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is hydrogen and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

25 In embodiment F25 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is chloro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F26 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is bromo and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F27 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is fluoro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.
30

In embodiment F28 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is trifluoromethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F29 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is methyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.
35

In embodiment F30 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is ethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , X^1 , X^2 and X^3 are as defined for the compound of formula I.

In embodiment F31 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is methoxy and $G^1, R^1, R^2, Y^1, Y^2, Y^3, X^1, X^2$ and X^3 are as defined for the compound of formula I.

In embodiment F32 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is nitro and $G^1, R^1, R^2, Y^1, Y^2, Y^3, X^1, X^2$ and X^3 are as defined for the compound of formula I.

- 5 In embodiment F33 the invention provides compounds of formula I wherein R^5 is trifluoromethoxy and $G^1, R^1, R^2, Y^1, Y^2, Y^3, X^1, X^2$ and X^3 are as defined for the compound of formula I.

In embodiment F34 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and R^5 is cyano and $G^1, R^1, R^2, Y^1, Y^2, Y^3, X^1, X^2$ and X^3 are as defined for the compound of formula I.

- In embodiment F35 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl
10 and R^5 is cyclopropyl and $G^1, R^1, R^2, Y^1, Y^2, Y^3, X^1, X^2$ and X^3 are as defined for the compound of formula I.

In embodiment F36 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is CH, X^3 is chloro and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

- 15 In embodiment F37 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is C-F, X^3 is hydrogen and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

In embodiment F38 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the

- 20 compound of formula I.

In embodiment F39 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

- In embodiment F40 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl
25 and X^1 is chloro, X^2 is C-Br, X^3 is chloro and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

In embodiment F41 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is C-F, X^3 is chloro and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

- 30 In embodiment F42 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is C-Cl, X^3 is chloro and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

- In embodiment F43 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is C-I, X^3 is chloro and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of
35 formula I.

In embodiment F44 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is fluoro, X^2 is C-F, X^3 is fluoro and $G^1, R^1, R^2, R^5, Y^1, Y^2, Y^3$, are as defined for the compound of formula I.

In embodiment F45 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is CH, X^3 is bromo and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , are as defined for the compound of formula I.

In embodiment F46 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is CH, X^3 is fluoro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , are as defined for the compound of formula I.

In embodiment F47 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , are as defined for the compound of formula I.

10 In embodiment F48 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , are as defined for the compound of formula I.

In embodiment F49 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , are as defined for the compound of formula I.

In embodiment F50 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 , are as defined for the compound of formula I.

In embodiment F51 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 are as defined for the compound of formula I.

In embodiment F52 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is chloro, X^2 is N, X^3 is chloro and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 are as defined for the compound of formula I.

25 In embodiment F53 the invention provides compounds of formula I wherein X^4 is chlorodifluoromethyl and X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 are as defined for the compound of formula I.

In embodiment G1 the invention provides compounds of formula I wherein X^2 is C- X^6 , none of X^1 , X^3 and X^6 are hydrogen and R^2 is thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G2 the invention provides compounds of formula I wherein X^2 is C- X^6 , none of X^1 , X^3 and X^6 are hydrogen and R^2 is 1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

35 In embodiment G3 the invention provides compounds of formula I wherein X^2 is C- X^6 , none of X^1 , X^3 and X^6 are hydrogen and R^2 is 1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G4 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 2,2-dimethylthietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G5 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 2,2-dimethyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G6 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 2,2-dimethyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

10 In embodiment G7 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 3-Methyl-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G8 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 3-Methyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as
15 defined for the compound of formula I.

In embodiment G9 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 3-Methyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G10 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for
20 the compound of formula I.

In embodiment G11 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 1-oxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

25 In embodiment G12 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 1,1-Dioxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G13 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is thietan-2-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for
30 the compound of formula I.

In embodiment G14 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is (1-oxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G15 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is (1,1-dioxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as
35 defined for the compound of formula I.

In embodiment G16 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 2-(thietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G17 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 2-(1,1-dioxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G18 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^2 is 2-(1-oxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

10 In embodiment G19 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

In embodiment G20 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the
15 compound of formula I.

In embodiment G21 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

In embodiment G22 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the
20 compound of formula I.

In embodiment G23 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

25 In embodiment G24 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is hydrogen and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G25 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is chloro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound
30 of formula I.

In embodiment G26 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is bromo and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G27 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is fluoro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound
35 of formula I.

In embodiment G28 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is trifluoromethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G29 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is methyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G30 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is ethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

10 In embodiment G31 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is methoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G32 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is nitro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of
15 formula I.

In embodiment G33 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is trifluoromethoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment G34 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is cyano and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound
20 of formula I.

In embodiment G35 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and R^5 is cyclopropyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

25 In embodiment G36 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and X^4 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 are as defined for the compound of formula I.

In embodiment G37 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and X^4 is difluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 are as defined for the
30 compound of formula I.

In embodiment G38 the invention provides compounds of formula I wherein X^2 is $C-X^6$, none of X^1 , X^3 and X^6 are hydrogen and X^4 is chlorodifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , Y^3 are as defined for the compound of formula I.

35 In embodiment H1 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^2 is thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment H2 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is

trifluoromethyl and R² is 1-oxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

In embodiment H3 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 1,1-Dioxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the
5 compound of formula I.

In embodiment H4 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 2,2-dimethylthietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

In embodiment H5 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 2,2-dimethyl-1-oxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as
10 defined for the compound of formula I.

In embodiment H6 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 2,2-dimethyl-1,1-Dioxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

15 In embodiment H7 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 3-Methyl-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

In embodiment H8 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 3-Methyl-1-oxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined
20 for the compound of formula I.

In embodiment H9 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 3-Methyl-1,1-Dioxo-thietan-3-yl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

In embodiment H10 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is thietan-3-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the
25 compound of formula I.

In embodiment H11 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 1-oxo-thietan-3-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

30 In embodiment H12 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is 1,1-Dioxo-thietan-3-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

In embodiment H13 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is thietan-2-ylmethyl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the
35 compound of formula I.

In embodiment H14 the invention provides compounds of formula I wherein X¹ is chloro, X² is CH, X³ is trifluoromethyl and R² is (1-oxothietan-2-yl)methyl- and G¹, R¹, R⁵, Y¹, Y², Y³, and X⁴ are as defined for the compound of formula I.

In embodiment H15 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^2 is (1,1-dioxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment H16 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2-(thietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment H17 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2-(1,1-dioxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

10 In embodiment H18 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2-(1-oxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 , and X^4 are as defined for the compound of formula I.

In embodiment H19 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

In embodiment H20 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

In embodiment H21 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

In embodiment H22 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

25 In embodiment H23 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 , and X^4 are as defined for the compound of formula I.

In embodiment H24 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is hydrogen and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H25 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is chloro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H26 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is bromo and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H27 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is fluoro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H28 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is trifluoromethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H29 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is methyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

10 In embodiment H30 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is ethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H31 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is methoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H32 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is nitro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H33 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is trifluoromethoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

⁴ In embodiment H34 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is cyano and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

25 In embodiment H35 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and R^5 is cyclopropyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment H36 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and X^4 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment H37 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and X^4 is difluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment H38 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl and X^4 is chlorodifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment J1 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J2 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J3 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

10 In embodiment J4 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 2,2-dimethylthietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J5 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 2,2-dimethyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the
15 compound of formula I.

In embodiment J6 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 2,2-dimethyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J7 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 3-Methyl-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound
20 of formula I.

In embodiment J8 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 3-Methyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

25 In embodiment J9 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 3-Methyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J10 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of
30 formula I.

In embodiment J11 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 1-oxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J12 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 1,1-Dioxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the
35 compound of formula I.

In embodiment J13 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is thietan-2-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J14 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is (1-oxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J15 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is (1,1-dioxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

10 In embodiment J16 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 2-(thietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J17 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 2-(1,1-dioxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the
15 compound of formula I.

In embodiment J18 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^2 is 2-(1-oxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J19 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of
20 formula I.

In embodiment J20 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

25 In embodiment J21 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment J22 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of
30 formula I.

In embodiment J23 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment J24 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is hydrogen and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula
35 I.

In embodiment J25 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is chloro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J26 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is bromo and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J27 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is fluoro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 5 In embodiment J28 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is trifluoromethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J29 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is methyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 10 In embodiment J30 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is ethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J31 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is methoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 15 In embodiment J32 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is nitro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment J33 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and R^5 is trifluoromethoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 20 In embodiment J34 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and X^4 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment J35 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and X^4 is difluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

- 25 In embodiment J36 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo and X^4 is chlorodifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

- 30 In embodiment K1 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K2 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

- 35 In embodiment K3 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K4 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2,2-dimethylthietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K5 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2,2-dimethyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K6 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2,2-dimethyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

10 In embodiment K7 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 3-Methyl-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K8 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 3-Methyl-1-oxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K9 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 3-Methyl-1,1-Dioxo-thietan-3-yl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K10 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K11 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 1-oxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

25 In embodiment K12 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 1,1-Dioxo-thietan-3-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K13 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is thietan-2-ylmethyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K14 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is (1-oxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K15 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is (1,1-dioxothietan-2-yl)methyl- and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K16 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2-(thietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K17 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2-(1,1-dioxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K18 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^2 is 2-(1-oxothietan-3-yl)ethanyl and G^1 , R^1 , R^5 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K19 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment K20 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment K21 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment K22 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment K23 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment K24 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is hydrogen and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K25 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is chloro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K26 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is bromo and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K27 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is fluoro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K28 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is trifluoromethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K29 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is methyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K30 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is ethyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K31 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is methoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K32 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is nitro and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K33 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is trifluoromethoxy and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K34 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is cyano and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K35 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and R^5 is cyclopropyl and G^1 , R^1 , R^2 , Y^1 , Y^2 , Y^3 and X^4 are as defined for the compound of formula I.

In embodiment K36 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and X^4 is trifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment K37 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and X^4 is difluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment K38 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl and X^4 is chlorodifluoromethyl and G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 and Y^3 are as defined for the compound of formula I.

In embodiment L1 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L2 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L3 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L4 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L5 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L6 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L7 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L8 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is hydrogen, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L9 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is hydrogen, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L10 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L11 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L12 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L13 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L14 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L15 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L16 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L17 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

10 In embodiment L18 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L19 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L20 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L21 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L22 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

25 In embodiment L23 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L24 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

30 In embodiment L25 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-F, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L26 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L27 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L28 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L29 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L30 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

10 In embodiment L31 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L32 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L33 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L34 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L35 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

25 In embodiment L36 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L37 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L38 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L39 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L40 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L41 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L42 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L43 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L44 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L45 the invention provides compounds of formula I wherein X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L46 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L47 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L48 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L49 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L50 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L51 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L52 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L53 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L54 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L55 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L56 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

10 In embodiment L57 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L58 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L59 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L60 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L61 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

25 In embodiment L62 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L63 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L64 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L65 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L66 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L67 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L68 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

10 In embodiment L69 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L70 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the
15 compound of formula I.

In embodiment L71 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L72 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the
20 compound of formula I.

In embodiment L73 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

25 In embodiment L74 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L75 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the
30 compound of formula I.

In embodiment L76 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L77 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the
35 compound of formula I.

In embodiment L78 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L79 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L80 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

10 In embodiment L81 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L82 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L83 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L84 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

20 In embodiment L85 the invention provides compounds of formula I wherein X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L86 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L87 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

30 In embodiment L88 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

In embodiment L89 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

35 In embodiment L90 the invention provides compounds of formula I wherein X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and G^1 , R^1 , R^2 , R^5 and X^4 are as defined for the compound of formula I.

The compounds in the Tables below illustrate the compounds of the invention.




Table P

	X4	R5	R2
P.001	chlorodifluoromethyl	bromo	(1,1-dioxothietan-2-yl)methyl-
P.002	chlorodifluoromethyl	bromo	(1-oxothietan-2-yl)methyl-
P.003	chlorodifluoromethyl	bromo	1,1-Dioxo-thietan-3-yl-
P.004	chlorodifluoromethyl	bromo	1,1-Dioxo-thietan-3-ylmethyl-
P.005	chlorodifluoromethyl	bromo	cis-1-oxo-thietan-3-yl-
P.006	chlorodifluoromethyl	bromo	cis-1-oxo-thietan-3-ylmethyl-
P.007	chlorodifluoromethyl	bromo	2-(1,1-dioxothietan-3-yl)ethanyl
P.008	chlorodifluoromethyl	bromo	2-(1-oxothietan-3-yl)ethanyl
P.009	chlorodifluoromethyl	bromo	2-(thietan-3-yl)ethanyl
P.010	chlorodifluoromethyl	bromo	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.011	chlorodifluoromethyl	bromo	2,2-dimethyl-1-oxo-thietan-3-yl-
P.012	chlorodifluoromethyl	bromo	2,2-dimethylthietan-3-yl-
P.013	chlorodifluoromethyl	bromo	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.014	chlorodifluoromethyl	bromo	3-Methyl-1-oxo-thietan-3-yl-
P.015	chlorodifluoromethyl	bromo	3-Methyl-thietan-3-yl-
P.016	chlorodifluoromethyl	bromo	thietan-2-ylmethyl-
P.017	chlorodifluoromethyl	bromo	thietan-3-yl-
P.018	chlorodifluoromethyl	bromo	thietan-3-ylmethyl-
P.019	chlorodifluoromethyl	chloro	(1,1-dioxothietan-2-yl)methyl-
P.020	chlorodifluoromethyl	chloro	(1-oxothietan-2-yl)methyl-
P.021	chlorodifluoromethyl	chloro	1,1-Dioxo-thietan-3-yl-
P.022	chlorodifluoromethyl	chloro	1,1-Dioxo-thietan-3-ylmethyl-
P.023	chlorodifluoromethyl	chloro	cis-1-oxo-thietan-3-yl-
P.024	chlorodifluoromethyl	chloro	cis-1-oxo-thietan-3-ylmethyl-
P.025	chlorodifluoromethyl	chloro	2-(1,1-dioxothietan-3-yl)ethanyl
P.026	chlorodifluoromethyl	chloro	2-(1-oxothietan-3-yl)ethanyl
P.027	chlorodifluoromethyl	chloro	2-(thietan-3-yl)ethanyl
P.028	chlorodifluoromethyl	chloro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.029	chlorodifluoromethyl	chloro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.030	chlorodifluoromethyl	chloro	2,2-dimethylthietan-3-yl-
P.031	chlorodifluoromethyl	chloro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.032	chlorodifluoromethyl	chloro	3-Methyl-1-oxo-thietan-3-yl-
P.033	chlorodifluoromethyl	chloro	3-Methyl-thietan-3-yl-

P.034	chlorodifluoromethyl	chloro	thietan-2-ylmethyl-
P.035	chlorodifluoromethyl	chloro	thietan-3-yl-
P.036	chlorodifluoromethyl	chloro	thietan-3-ylmethyl-
P.037	chlorodifluoromethyl	cyano	(1,1-dioxothietan-2-yl)methyl-
P.038	chlorodifluoromethyl	cyano	(1-oxothietan-2-yl)methyl-
P.039	chlorodifluoromethyl	cyano	1,1-Dioxo-thietan-3-yl-
P.040	chlorodifluoromethyl	cyano	1,1-Dioxo-thietan-3-ylmethyl-
P.041	chlorodifluoromethyl	cyano	cis-1-oxo-thietan-3-yl-
P.042	chlorodifluoromethyl	cyano	cis-1-oxo-thietan-3-ylmethyl-
P.043	chlorodifluoromethyl	cyano	2-(1,1-dioxothietan-3-yl)ethanyl
P.044	chlorodifluoromethyl	cyano	2-(1-oxothietan-3-yl)ethanyl
P.045	chlorodifluoromethyl	cyano	2-(thietan-3-yl)ethanyl
P.046	chlorodifluoromethyl	cyano	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.047	chlorodifluoromethyl	cyano	2,2-dimethyl-1-oxo-thietan-3-yl-
P.048	chlorodifluoromethyl	cyano	2,2-dimethylthietan-3-yl-
P.049	chlorodifluoromethyl	cyano	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.050	chlorodifluoromethyl	cyano	3-Methyl-1-oxo-thietan-3-yl-
P.051	chlorodifluoromethyl	cyano	3-Methyl-thietan-3-yl-
P.052	chlorodifluoromethyl	cyano	thietan-2-ylmethyl-
P.053	chlorodifluoromethyl	cyano	thietan-3-yl-
P.054	chlorodifluoromethyl	cyano	thietan-3-ylmethyl-
P.055	chlorodifluoromethyl	cyclopropyl	(1,1-dioxothietan-2-yl)methyl-
P.056	chlorodifluoromethyl	cyclopropyl	(1-oxothietan-2-yl)methyl-
P.057	chlorodifluoromethyl	cyclopropyl	1,1-Dioxo-thietan-3-yl-
P.058	chlorodifluoromethyl	cyclopropyl	1,1-Dioxo-thietan-3-ylmethyl-
P.059	chlorodifluoromethyl	cyclopropyl	cis-1-oxo-thietan-3-yl-
P.060	chlorodifluoromethyl	cyclopropyl	cis-1-oxo-thietan-3-ylmethyl-
P.061	chlorodifluoromethyl	cyclopropyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.062	chlorodifluoromethyl	cyclopropyl	2-(1-oxothietan-3-yl)ethanyl
P.063	chlorodifluoromethyl	cyclopropyl	2-(thietan-3-yl)ethanyl
P.064	chlorodifluoromethyl	cyclopropyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.065	chlorodifluoromethyl	cyclopropyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.066	chlorodifluoromethyl	cyclopropyl	2,2-dimethylthietan-3-yl-
P.067	chlorodifluoromethyl	cyclopropyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.068	chlorodifluoromethyl	cyclopropyl	3-Methyl-1-oxo-thietan-3-yl-
P.069	chlorodifluoromethyl	cyclopropyl	3-Methyl-thietan-3-yl-
P.070	chlorodifluoromethyl	cyclopropyl	thietan-2-ylmethyl-
P.071	chlorodifluoromethyl	cyclopropyl	thietan-3-yl-
P.072	chlorodifluoromethyl	cyclopropyl	thietan-3-ylmethyl-
P.073	chlorodifluoromethyl	ethyl	(1,1-dioxothietan-2-yl)methyl-
P.074	chlorodifluoromethyl	ethyl	(1-oxothietan-2-yl)methyl-
P.075	chlorodifluoromethyl	ethyl	1,1-Dioxo-thietan-3-yl-
P.076	chlorodifluoromethyl	ethyl	1,1-Dioxo-thietan-3-ylmethyl-
P.077	chlorodifluoromethyl	ethyl	cis-1-oxo-thietan-3-yl-
P.078	chlorodifluoromethyl	ethyl	cis-1-oxo-thietan-3-ylmethyl-
P.079	chlorodifluoromethyl	ethyl	2-(1,1-dioxothietan-3-yl)ethanyl

P.080	chlorodifluoromethyl	ethyl	2-(1-oxothietan-3-yl)ethanyl
P.081	chlorodifluoromethyl	ethyl	2-(thietan-3-yl)ethanyl
P.082	chlorodifluoromethyl	ethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.083	chlorodifluoromethyl	ethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.084	chlorodifluoromethyl	ethyl	2,2-dimethylthietan-3-yl-
P.085	chlorodifluoromethyl	ethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.086	chlorodifluoromethyl	ethyl	3-Methyl-1-oxo-thietan-3-yl-
P.087	chlorodifluoromethyl	ethyl	3-Methyl-thietan-3-yl-
P.088	chlorodifluoromethyl	ethyl	thietan-2-ylmethyl-
P.089	chlorodifluoromethyl	ethyl	thietan-3-yl-
P.090	chlorodifluoromethyl	ethyl	thietan-3-ylmethyl-
P.091	chlorodifluoromethyl	fluoro	(1,1-dioxothietan-2-yl)methyl-
P.092	chlorodifluoromethyl	fluoro	(1-oxothietan-2-yl)methyl-
P.093	chlorodifluoromethyl	fluoro	1,1-Dioxo-thietan-3-yl-
P.094	chlorodifluoromethyl	fluoro	1,1-Dioxo-thietan-3-ylmethyl-
P.095	chlorodifluoromethyl	fluoro	cis-1-oxo-thietan-3-yl-
P.096	chlorodifluoromethyl	fluoro	cis-1-oxo-thietan-3-ylmethyl-
P.097	chlorodifluoromethyl	fluoro	2-(1,1-dioxothietan-3-yl)ethanyl
P.098	chlorodifluoromethyl	fluoro	2-(1-oxothietan-3-yl)ethanyl
P.099	chlorodifluoromethyl	fluoro	2-(thietan-3-yl)ethanyl
P.100	chlorodifluoromethyl	fluoro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.101	chlorodifluoromethyl	fluoro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.102	chlorodifluoromethyl	fluoro	2,2-dimethylthietan-3-yl-
P.103	chlorodifluoromethyl	fluoro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.104	chlorodifluoromethyl	fluoro	3-Methyl-1-oxo-thietan-3-yl-
P.105	chlorodifluoromethyl	fluoro	3-Methyl-thietan-3-yl-
P.106	chlorodifluoromethyl	fluoro	thietan-2-ylmethyl-
P.107	chlorodifluoromethyl	fluoro	thietan-3-yl-
P.108	chlorodifluoromethyl	fluoro	thietan-3-ylmethyl-
P.109	chlorodifluoromethyl	hydrogen	(1,1-dioxothietan-2-yl)methyl-
P.110	chlorodifluoromethyl	hydrogen	(1-oxothietan-2-yl)methyl-
P.111	chlorodifluoromethyl	hydrogen	1,1-Dioxo-thietan-3-yl-
P.112	chlorodifluoromethyl	hydrogen	1,1-Dioxo-thietan-3-ylmethyl-
P.113	chlorodifluoromethyl	hydrogen	cis-1-oxo-thietan-3-yl-
P.114	chlorodifluoromethyl	hydrogen	cis-1-oxo-thietan-3-ylmethyl-
P.115	chlorodifluoromethyl	hydrogen	2-(1,1-dioxothietan-3-yl)ethanyl
P.116	chlorodifluoromethyl	hydrogen	2-(1-oxothietan-3-yl)ethanyl
P.117	chlorodifluoromethyl	hydrogen	2-(thietan-3-yl)ethanyl
P.118	chlorodifluoromethyl	hydrogen	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.119	chlorodifluoromethyl	hydrogen	2,2-dimethyl-1-oxo-thietan-3-yl-
P.120	chlorodifluoromethyl	hydrogen	2,2-dimethylthietan-3-yl-
P.121	chlorodifluoromethyl	hydrogen	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.122	chlorodifluoromethyl	hydrogen	3-Methyl-1-oxo-thietan-3-yl-
P.123	chlorodifluoromethyl	hydrogen	3-Methyl-thietan-3-yl-
P.124	chlorodifluoromethyl	hydrogen	thietan-2-ylmethyl-
P.125	chlorodifluoromethyl	hydrogen	thietan-3-yl-

P.126	chlorodifluoromethyl	hydrogen	thietan-3-ylmethyl-
P.127	chlorodifluoromethyl	methoxy	(1,1-dioxothietan-2-yl)methyl-
P.128	chlorodifluoromethyl	methoxy	(1-oxothietan-2-yl)methyl-
P.129	chlorodifluoromethyl	methoxy	1,1-Dioxo-thietan-3-yl-
P.130	chlorodifluoromethyl	methoxy	1,1-Dioxo-thietan-3-ylmethyl-
P.131	chlorodifluoromethyl	methoxy	cis-1-oxo-thietan-3-yl-
P.132	chlorodifluoromethyl	methoxy	cis-1-oxo-thietan-3-ylmethyl-
P.133	chlorodifluoromethyl	methoxy	2-(1,1-dioxothietan-3-yl)ethanyl
P.134	chlorodifluoromethyl	methoxy	2-(1-oxothietan-3-yl)ethanyl
P.135	chlorodifluoromethyl	methoxy	2-(thietan-3-yl)ethanyl
P.136	chlorodifluoromethyl	methoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.137	chlorodifluoromethyl	methoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
P.138	chlorodifluoromethyl	methoxy	2,2-dimethylthietan-3-yl-
P.139	chlorodifluoromethyl	methoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.140	chlorodifluoromethyl	methoxy	3-Methyl-1-oxo-thietan-3-yl-
P.141	chlorodifluoromethyl	methoxy	3-Methyl-thietan-3-yl-
P.142	chlorodifluoromethyl	methoxy	thietan-2-ylmethyl-
P.143	chlorodifluoromethyl	methoxy	thietan-3-yl-
P.144	chlorodifluoromethyl	methoxy	thietan-3-ylmethyl-
P.145	chlorodifluoromethyl	methyl	(1,1-dioxothietan-2-yl)methyl-
P.146	chlorodifluoromethyl	methyl	(1-oxothietan-2-yl)methyl-
P.147	chlorodifluoromethyl	methyl	1,1-Dioxo-thietan-3-yl-
P.148	chlorodifluoromethyl	methyl	1,1-Dioxo-thietan-3-ylmethyl-
P.149	chlorodifluoromethyl	methyl	cis-1-oxo-thietan-3-yl-
P.150	chlorodifluoromethyl	methyl	cis-1-oxo-thietan-3-ylmethyl-
P.151	chlorodifluoromethyl	methyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.152	chlorodifluoromethyl	methyl	2-(1-oxothietan-3-yl)ethanyl
P.153	chlorodifluoromethyl	methyl	2-(thietan-3-yl)ethanyl
P.154	chlorodifluoromethyl	methyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.155	chlorodifluoromethyl	methyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.156	chlorodifluoromethyl	methyl	2,2-dimethylthietan-3-yl-
P.157	chlorodifluoromethyl	methyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.158	chlorodifluoromethyl	methyl	3-Methyl-1-oxo-thietan-3-yl-
P.159	chlorodifluoromethyl	methyl	3-Methyl-thietan-3-yl-
P.160	chlorodifluoromethyl	methyl	thietan-2-ylmethyl-
P.161	chlorodifluoromethyl	methyl	thietan-3-yl-
P.162	chlorodifluoromethyl	methyl	thietan-3-ylmethyl-
P.163	chlorodifluoromethyl	nitro	(1,1-dioxothietan-2-yl)methyl-
P.164	chlorodifluoromethyl	nitro	(1-oxothietan-2-yl)methyl-
P.165	chlorodifluoromethyl	nitro	1,1-Dioxo-thietan-3-yl-
P.166	chlorodifluoromethyl	nitro	1,1-Dioxo-thietan-3-ylmethyl-
P.167	chlorodifluoromethyl	nitro	cis-1-oxo-thietan-3-yl-
P.168	chlorodifluoromethyl	nitro	cis-1-oxo-thietan-3-ylmethyl-
P.169	chlorodifluoromethyl	nitro	2-(1,1-dioxothietan-3-yl)ethanyl
P.170	chlorodifluoromethyl	nitro	2-(1-oxothietan-3-yl)ethanyl
P.171	chlorodifluoromethyl	nitro	2-(thietan-3-yl)ethanyl

P.172	chlorodifluoromethyl	nitro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.173	chlorodifluoromethyl	nitro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.174	chlorodifluoromethyl	nitro	2,2-dimethylthietan-3-yl-
P.175	chlorodifluoromethyl	nitro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.176	chlorodifluoromethyl	nitro	3-Methyl-1-oxo-thietan-3-yl-
P.177	chlorodifluoromethyl	nitro	3-Methyl-thietan-3-yl-
P.178	chlorodifluoromethyl	nitro	thietan-2-ylmethyl-
P.179	chlorodifluoromethyl	nitro	thietan-3-yl-
P.180	chlorodifluoromethyl	nitro	thietan-3-ylmethyl-
P.181	chlorodifluoromethyl	trifluoromethoxy	(1,1-dioxothietan-2-yl)methyl-
P.182	chlorodifluoromethyl	trifluoromethoxy	(1-oxothietan-2-yl)methyl-
P.183	chlorodifluoromethyl	trifluoromethoxy	1,1-Dioxo-thietan-3-yl-
P.184	chlorodifluoromethyl	trifluoromethoxy	1,1-Dioxo-thietan-3-ylmethyl-
P.185	chlorodifluoromethyl	trifluoromethoxy	cis-1-oxo-thietan-3-yl-
P.186	chlorodifluoromethyl	trifluoromethoxy	cis-1-oxo-thietan-3-ylmethyl-
P.187	chlorodifluoromethyl	trifluoromethoxy	2-(1,1-dioxothietan-3-yl)ethanyl
P.188	chlorodifluoromethyl	trifluoromethoxy	2-(1-oxothietan-3-yl)ethanyl
P.189	chlorodifluoromethyl	trifluoromethoxy	2-(thietan-3-yl)ethanyl
P.190	chlorodifluoromethyl	trifluoromethoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.191	chlorodifluoromethyl	trifluoromethoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
P.192	chlorodifluoromethyl	trifluoromethoxy	2,2-dimethylthietan-3-yl-
P.193	chlorodifluoromethyl	trifluoromethoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.194	chlorodifluoromethyl	trifluoromethoxy	3-Methyl-1-oxo-thietan-3-yl-
P.195	chlorodifluoromethyl	trifluoromethoxy	3-Methyl-thietan-3-yl-
P.196	chlorodifluoromethyl	trifluoromethoxy	thietan-2-ylmethyl-
P.197	chlorodifluoromethyl	trifluoromethoxy	thietan-3-yl-
P.198	chlorodifluoromethyl	trifluoromethoxy	thietan-3-ylmethyl-
P.199	chlorodifluoromethyl	trifluoromethyl	(1,1-dioxothietan-2-yl)methyl-
P.200	chlorodifluoromethyl	trifluoromethyl	(1-oxothietan-2-yl)methyl-
P.201	chlorodifluoromethyl	trifluoromethyl	1,1-Dioxo-thietan-3-yl-
P.202	chlorodifluoromethyl	trifluoromethyl	1,1-Dioxo-thietan-3-ylmethyl-
P.203	chlorodifluoromethyl	trifluoromethyl	cis-1-oxo-thietan-3-yl-
P.204	chlorodifluoromethyl	trifluoromethyl	cis-1-oxo-thietan-3-ylmethyl-
P.205	chlorodifluoromethyl	trifluoromethyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.206	chlorodifluoromethyl	trifluoromethyl	2-(1-oxothietan-3-yl)ethanyl
P.207	chlorodifluoromethyl	trifluoromethyl	2-(thietan-3-yl)ethanyl
P.208	chlorodifluoromethyl	trifluoromethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.209	chlorodifluoromethyl	trifluoromethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.210	chlorodifluoromethyl	trifluoromethyl	2,2-dimethylthietan-3-yl-
P.211	chlorodifluoromethyl	trifluoromethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.212	chlorodifluoromethyl	trifluoromethyl	3-Methyl-1-oxo-thietan-3-yl-
P.213	chlorodifluoromethyl	trifluoromethyl	3-Methyl-thietan-3-yl-
P.214	chlorodifluoromethyl	trifluoromethyl	thietan-2-ylmethyl-
P.215	chlorodifluoromethyl	trifluoromethyl	thietan-3-yl-
P.216	chlorodifluoromethyl	trifluoromethyl	thietan-3-ylmethyl-
P.217	difluoromethyl	bromo	(1,1-dioxothietan-2-yl)methyl-

P.218	difluoromethyl	bromo	(1-oxothietan-2-yl)methyl-
P.219	difluoromethyl	bromo	1,1-Dioxo-thietan-3-yl-
P.220	difluoromethyl	bromo	1,1-Dioxo-thietan-3-ylmethyl-
P.221	difluoromethyl	bromo	cis-1-oxo-thietan-3-yl-
P.222	difluoromethyl	bromo	cis-1-oxo-thietan-3-ylmethyl-
P.223	difluoromethyl	bromo	2-(1,1-dioxothietan-3-yl)ethanyl
P.224	difluoromethyl	bromo	2-(1-oxothietan-3-yl)ethanyl
P.225	difluoromethyl	bromo	2-(thietan-3-yl)ethanyl
P.226	difluoromethyl	bromo	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.227	difluoromethyl	bromo	2,2-dimethyl-1-oxo-thietan-3-yl-
P.228	difluoromethyl	bromo	2,2-dimethylthietan-3-yl-
P.229	difluoromethyl	bromo	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.230	difluoromethyl	bromo	3-Methyl-1-oxo-thietan-3-yl-
P.231	difluoromethyl	bromo	3-Methyl-thietan-3-yl-
P.232	difluoromethyl	bromo	thietan-2-ylmethyl-
P.233	difluoromethyl	bromo	thietan-3-yl-
P.234	difluoromethyl	bromo	thietan-3-ylmethyl-
P.235	difluoromethyl	chloro	(1,1-dioxothietan-2-yl)methyl-
P.236	difluoromethyl	chloro	(1-oxothietan-2-yl)methyl-
P.237	difluoromethyl	chloro	1,1-Dioxo-thietan-3-yl-
P.238	difluoromethyl	chloro	1,1-Dioxo-thietan-3-ylmethyl-
P.239	difluoromethyl	chloro	cis-1-oxo-thietan-3-yl-
P.240	difluoromethyl	chloro	cis-1-oxo-thietan-3-ylmethyl-
P.241	difluoromethyl	chloro	2-(1,1-dioxothietan-3-yl)ethanyl
P.242	difluoromethyl	chloro	2-(1-oxothietan-3-yl)ethanyl
P.243	difluoromethyl	chloro	2-(thietan-3-yl)ethanyl
P.244	difluoromethyl	chloro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.245	difluoromethyl	chloro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.246	difluoromethyl	chloro	2,2-dimethylthietan-3-yl-
P.247	difluoromethyl	chloro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.248	difluoromethyl	chloro	3-Methyl-1-oxo-thietan-3-yl-
P.249	difluoromethyl	chloro	3-Methyl-thietan-3-yl-
P.250	difluoromethyl	chloro	thietan-2-ylmethyl-
P.251	difluoromethyl	chloro	thietan-3-yl-
P.252	difluoromethyl	chloro	thietan-3-ylmethyl-
P.253	difluoromethyl	cyano	(1,1-dioxothietan-2-yl)methyl-
P.254	difluoromethyl	cyano	(1-oxothietan-2-yl)methyl-
P.255	difluoromethyl	cyano	1,1-Dioxo-thietan-3-yl-
P.256	difluoromethyl	cyano	1,1-Dioxo-thietan-3-ylmethyl-
P.257	difluoromethyl	cyano	cis-1-oxo-thietan-3-yl-
P.258	difluoromethyl	cyano	cis-1-oxo-thietan-3-ylmethyl-
P.259	difluoromethyl	cyano	2-(1,1-dioxothietan-3-yl)ethanyl
P.260	difluoromethyl	cyano	2-(1-oxothietan-3-yl)ethanyl
P.261	difluoromethyl	cyano	2-(thietan-3-yl)ethanyl
P.262	difluoromethyl	cyano	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.263	difluoromethyl	cyano	2,2-dimethyl-1-oxo-thietan-3-yl-

P.264	difluoromethyl	cyano	2,2-dimethylthietan-3-yl-
P.265	difluoromethyl	cyano	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.266	difluoromethyl	cyano	3-Methyl-1-oxo-thietan-3-yl-
P.267	difluoromethyl	cyano	3-Methyl-thietan-3-yl-
P.268	difluoromethyl	cyano	thietan-2-ylmethyl-
P.269	difluoromethyl	cyano	thietan-3-yl-
P.270	difluoromethyl	cyano	thietan-3-ylmethyl-
P.271	difluoromethyl	cyclopropyl	(1,1-dioxothietan-2-yl)methyl-
P.272	difluoromethyl	cyclopropyl	(1-oxothietan-2-yl)methyl-
P.273	difluoromethyl	cyclopropyl	1,1-Dioxo-thietan-3-yl-
P.274	difluoromethyl	cyclopropyl	1,1-Dioxo-thietan-3-ylmethyl-
P.275	difluoromethyl	cyclopropyl	cis-1-oxo-thietan-3-yl-
P.276	difluoromethyl	cyclopropyl	cis-1-oxo-thietan-3-ylmethyl-
P.277	difluoromethyl	cyclopropyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.278	difluoromethyl	cyclopropyl	2-(1-oxothietan-3-yl)ethanyl
P.279	difluoromethyl	cyclopropyl	2-(thietan-3-yl)ethanyl
P.280	difluoromethyl	cyclopropyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.281	difluoromethyl	cyclopropyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.282	difluoromethyl	cyclopropyl	2,2-dimethylthietan-3-yl-
P.283	difluoromethyl	cyclopropyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.284	difluoromethyl	cyclopropyl	3-Methyl-1-oxo-thietan-3-yl-
P.285	difluoromethyl	cyclopropyl	3-Methyl-thietan-3-yl-
P.286	difluoromethyl	cyclopropyl	thietan-2-ylmethyl-
P.287	difluoromethyl	cyclopropyl	thietan-3-yl-
P.288	difluoromethyl	cyclopropyl	thietan-3-ylmethyl-
P.289	difluoromethyl	ethyl	(1,1-dioxothietan-2-yl)methyl-
P.290	difluoromethyl	ethyl	(1-oxothietan-2-yl)methyl-
P.291	difluoromethyl	ethyl	1,1-Dioxo-thietan-3-yl-
P.292	difluoromethyl	ethyl	1,1-Dioxo-thietan-3-ylmethyl-
P.293	difluoromethyl	ethyl	cis-1-oxo-thietan-3-yl-
P.294	difluoromethyl	ethyl	cis-1-oxo-thietan-3-ylmethyl-
P.295	difluoromethyl	ethyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.296	difluoromethyl	ethyl	2-(1-oxothietan-3-yl)ethanyl
P.297	difluoromethyl	ethyl	2-(thietan-3-yl)ethanyl
P.298	difluoromethyl	ethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.299	difluoromethyl	ethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.300	difluoromethyl	ethyl	2,2-dimethylthietan-3-yl-
P.301	difluoromethyl	ethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.302	difluoromethyl	ethyl	3-Methyl-1-oxo-thietan-3-yl-
P.303	difluoromethyl	ethyl	3-Methyl-thietan-3-yl-
P.304	difluoromethyl	ethyl	thietan-2-ylmethyl-
P.305	difluoromethyl	ethyl	thietan-3-yl-
P.306	difluoromethyl	ethyl	thietan-3-ylmethyl-
P.307	difluoromethyl	fluoro	(1,1-dioxothietan-2-yl)methyl-
P.308	difluoromethyl	fluoro	(1-oxothietan-2-yl)methyl-
P.309	difluoromethyl	fluoro	1,1-Dioxo-thietan-3-yl-

P.310	difluoromethyl	fluoro	1,1-Dioxo-thietan-3-ylmethyl-
P.311	difluoromethyl	fluoro	cis-1-oxo-thietan-3-yl-
P.312	difluoromethyl	fluoro	cis-1-oxo-thietan-3-ylmethyl-
P.313	difluoromethyl	fluoro	2-(1,1-dioxothietan-3-yl)ethanyl
P.314	difluoromethyl	fluoro	2-(1-oxothietan-3-yl)ethanyl
P.315	difluoromethyl	fluoro	2-(thietan-3-yl)ethanyl
P.316	difluoromethyl	fluoro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.317	difluoromethyl	fluoro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.318	difluoromethyl	fluoro	2,2-dimethylthietan-3-yl-
P.319	difluoromethyl	fluoro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.320	difluoromethyl	fluoro	3-Methyl-1-oxo-thietan-3-yl-
P.321	difluoromethyl	fluoro	3-Methyl-thietan-3-yl-
P.322	difluoromethyl	fluoro	thietan-2-ylmethyl-
P.323	difluoromethyl	fluoro	thietan-3-yl-
P.324	difluoromethyl	fluoro	thietan-3-ylmethyl-
P.325	difluoromethyl	hydrogen	(1,1-dioxothietan-2-yl)methyl-
P.326	difluoromethyl	hydrogen	(1-oxothietan-2-yl)methyl-
P.327	difluoromethyl	hydrogen	1,1-Dioxo-thietan-3-yl-
P.328	difluoromethyl	hydrogen	1,1-Dioxo-thietan-3-ylmethyl-
P.329	difluoromethyl	hydrogen	cis-1-oxo-thietan-3-yl-
P.330	difluoromethyl	hydrogen	cis-1-oxo-thietan-3-ylmethyl-
P.331	difluoromethyl	hydrogen	2-(1,1-dioxothietan-3-yl)ethanyl
P.332	difluoromethyl	hydrogen	2-(1-oxothietan-3-yl)ethanyl
P.333	difluoromethyl	hydrogen	2-(thietan-3-yl)ethanyl
P.334	difluoromethyl	hydrogen	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.335	difluoromethyl	hydrogen	2,2-dimethyl-1-oxo-thietan-3-yl-
P.336	difluoromethyl	hydrogen	2,2-dimethylthietan-3-yl-
P.337	difluoromethyl	hydrogen	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.338	difluoromethyl	hydrogen	3-Methyl-1-oxo-thietan-3-yl-
P.339	difluoromethyl	hydrogen	3-Methyl-thietan-3-yl-
P.340	difluoromethyl	hydrogen	thietan-2-ylmethyl-
P.341	difluoromethyl	hydrogen	thietan-3-yl-
P.342	difluoromethyl	hydrogen	thietan-3-ylmethyl-
P.343	difluoromethyl	methoxy	(1,1-dioxothietan-2-yl)methyl-
P.344	difluoromethyl	methoxy	(1-oxothietan-2-yl)methyl-
P.345	difluoromethyl	methoxy	1,1-Dioxo-thietan-3-yl-
P.346	difluoromethyl	methoxy	1,1-Dioxo-thietan-3-ylmethyl-
P.347	difluoromethyl	methoxy	cis-1-oxo-thietan-3-yl-
P.348	difluoromethyl	methoxy	cis-1-oxo-thietan-3-ylmethyl-
P.349	difluoromethyl	methoxy	2-(1,1-dioxothietan-3-yl)ethanyl
P.350	difluoromethyl	methoxy	2-(1-oxothietan-3-yl)ethanyl
P.351	difluoromethyl	methoxy	2-(thietan-3-yl)ethanyl
P.352	difluoromethyl	methoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.353	difluoromethyl	methoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
P.354	difluoromethyl	methoxy	2,2-dimethylthietan-3-yl-
P.355	difluoromethyl	methoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-

P.356	difluoromethyl	methoxy	3-Methyl-1-oxo-thietan-3-yl-
P.357	difluoromethyl	methoxy	3-Methyl-thietan-3-yl-
P.358	difluoromethyl	methoxy	thietan-2-ylmethyl-
P.359	difluoromethyl	methoxy	thietan-3-yl-
P.360	difluoromethyl	methoxy	thietan-3-ylmethyl-
P.361	difluoromethyl	methyl	(1,1-dioxothietan-2-yl)methyl-
P.362	difluoromethyl	methyl	(1-oxothietan-2-yl)methyl-
P.363	difluoromethyl	methyl	1,1-Dioxo-thietan-3-yl-
P.364	difluoromethyl	methyl	1,1-Dioxo-thietan-3-ylmethyl-
P.365	difluoromethyl	methyl	cis-1-oxo-thietan-3-yl-
P.366	difluoromethyl	methyl	cis-1-oxo-thietan-3-ylmethyl-
P.367	difluoromethyl	methyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.368	difluoromethyl	methyl	2-(1-oxothietan-3-yl)ethanyl
P.369	difluoromethyl	methyl	2-(thietan-3-yl)ethanyl
P.370	difluoromethyl	methyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.371	difluoromethyl	methyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.372	difluoromethyl	methyl	2,2-dimethylthietan-3-yl-
P.373	difluoromethyl	methyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.374	difluoromethyl	methyl	3-Methyl-1-oxo-thietan-3-yl-
P.375	difluoromethyl	methyl	3-Methyl-thietan-3-yl-
P.376	difluoromethyl	methyl	thietan-2-ylmethyl-
P.377	difluoromethyl	methyl	thietan-3-yl-
P.378	difluoromethyl	methyl	thietan-3-ylmethyl-
P.379	difluoromethyl	nitro	(1,1-dioxothietan-2-yl)methyl-
P.380	difluoromethyl	nitro	(1-oxothietan-2-yl)methyl-
P.381	difluoromethyl	nitro	1,1-Dioxo-thietan-3-yl-
P.382	difluoromethyl	nitro	1,1-Dioxo-thietan-3-ylmethyl-
P.383	difluoromethyl	nitro	cis-1-oxo-thietan-3-yl-
P.384	difluoromethyl	nitro	cis-1-oxo-thietan-3-ylmethyl-
P.385	difluoromethyl	nitro	2-(1,1-dioxothietan-3-yl)ethanyl
P.386	difluoromethyl	nitro	2-(1-oxothietan-3-yl)ethanyl
P.387	difluoromethyl	nitro	2-(thietan-3-yl)ethanyl
P.388	difluoromethyl	nitro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.389	difluoromethyl	nitro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.390	difluoromethyl	nitro	2,2-dimethylthietan-3-yl-
P.391	difluoromethyl	nitro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.392	difluoromethyl	nitro	3-Methyl-1-oxo-thietan-3-yl-
P.393	difluoromethyl	nitro	3-Methyl-thietan-3-yl-
P.394	difluoromethyl	nitro	thietan-2-ylmethyl-
P.395	difluoromethyl	nitro	thietan-3-yl-
P.396	difluoromethyl	nitro	thietan-3-ylmethyl-
P.397	difluoromethyl	trifluoromethoxy	(1,1-dioxothietan-2-yl)methyl-
P.398	difluoromethyl	trifluoromethoxy	(1-oxothietan-2-yl)methyl-
P.399	difluoromethyl	trifluoromethoxy	1,1-Dioxo-thietan-3-yl-
P.400	difluoromethyl	trifluoromethoxy	1,1-Dioxo-thietan-3-ylmethyl-
P.401	difluoromethyl	trifluoromethoxy	cis-1-oxo-thietan-3-yl-

P.402	difluoromethyl	trifluoromethoxy	cis-1-oxo-thietan-3-ylmethyl-
P.403	difluoromethyl	trifluoromethoxy	2-(1,1-dioxothietan-3-yl)ethanyl
P.404	difluoromethyl	trifluoromethoxy	2-(1-oxothietan-3-yl)ethanyl
P.405	difluoromethyl	trifluoromethoxy	2-(thietan-3-yl)ethanyl
P.406	difluoromethyl	trifluoromethoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.407	difluoromethyl	trifluoromethoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
P.408	difluoromethyl	trifluoromethoxy	2,2-dimethylthietan-3-yl-
P.409	difluoromethyl	trifluoromethoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.410	difluoromethyl	trifluoromethoxy	3-Methyl-1-oxo-thietan-3-yl-
P.411	difluoromethyl	trifluoromethoxy	3-Methyl-thietan-3-yl-
P.412	difluoromethyl	trifluoromethoxy	thietan-2-ylmethyl-
P.413	difluoromethyl	trifluoromethoxy	thietan-3-yl-
P.414	difluoromethyl	trifluoromethoxy	thietan-3-ylmethyl-
P.415	difluoromethyl	trifluoromethyl	(1,1-dioxothietan-2-yl)methyl-
P.416	difluoromethyl	trifluoromethyl	(1-oxothietan-2-yl)methyl-
P.417	difluoromethyl	trifluoromethyl	1,1-Dioxo-thietan-3-yl-
P.418	difluoromethyl	trifluoromethyl	1,1-Dioxo-thietan-3-ylmethyl-
P.419	difluoromethyl	trifluoromethyl	cis-1-oxo-thietan-3-yl-
P.420	difluoromethyl	trifluoromethyl	cis-1-oxo-thietan-3-ylmethyl-
P.421	difluoromethyl	trifluoromethyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.422	difluoromethyl	trifluoromethyl	2-(1-oxothietan-3-yl)ethanyl
P.423	difluoromethyl	trifluoromethyl	2-(thietan-3-yl)ethanyl
P.424	difluoromethyl	trifluoromethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.425	difluoromethyl	trifluoromethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.426	difluoromethyl	trifluoromethyl	2,2-dimethylthietan-3-yl-
P.427	difluoromethyl	trifluoromethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.428	difluoromethyl	trifluoromethyl	3-Methyl-1-oxo-thietan-3-yl-
P.429	difluoromethyl	trifluoromethyl	3-Methyl-thietan-3-yl-
P.430	difluoromethyl	trifluoromethyl	thietan-2-ylmethyl-
P.431	difluoromethyl	trifluoromethyl	thietan-3-yl-
P.432	difluoromethyl	trifluoromethyl	thietan-3-ylmethyl-
P.433	trifluoromethyl	bromo	(1,1-dioxothietan-2-yl)methyl-
P.434	trifluoromethyl	bromo	(1-oxothietan-2-yl)methyl-
P.435	trifluoromethyl	bromo	1,1-Dioxo-thietan-3-yl-
P.436	trifluoromethyl	bromo	1,1-Dioxo-thietan-3-ylmethyl-
P.437	trifluoromethyl	bromo	cis-1-oxo-thietan-3-yl-
P.438	trifluoromethyl	bromo	cis-1-oxo-thietan-3-ylmethyl-
P.439	trifluoromethyl	bromo	2-(1,1-dioxothietan-3-yl)ethanyl
P.440	trifluoromethyl	bromo	2-(1-oxothietan-3-yl)ethanyl
P.441	trifluoromethyl	bromo	2-(thietan-3-yl)ethanyl
P.442	trifluoromethyl	bromo	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.443	trifluoromethyl	bromo	2,2-dimethyl-1-oxo-thietan-3-yl-
P.444	trifluoromethyl	bromo	2,2-dimethylthietan-3-yl-
P.445	trifluoromethyl	bromo	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.446	trifluoromethyl	bromo	3-Methyl-1-oxo-thietan-3-yl-
P.447	trifluoromethyl	bromo	3-Methyl-thietan-3-yl-

P.448	trifluoromethyl	bromo	thietan-2-ylmethyl-
P.449	trifluoromethyl	bromo	thietan-3-yl-
P.450	trifluoromethyl	bromo	thietan-3-ylmethyl-
P.451	trifluoromethyl	chloro	(1,1-dioxothietan-2-yl)methyl-
P.452	trifluoromethyl	chloro	(1-oxothietan-2-yl)methyl-
P.453	trifluoromethyl	chloro	1,1-Dioxo-thietan-3-yl-
P.454	trifluoromethyl	chloro	1,1-Dioxo-thietan-3-ylmethyl-
P.455	trifluoromethyl	chloro	cis-1-oxo-thietan-3-yl-
P.456	trifluoromethyl	chloro	cis-1-oxo-thietan-3-ylmethyl-
P.457	trifluoromethyl	chloro	2-(1,1-dioxothietan-3-yl)ethanyl
P.458	trifluoromethyl	chloro	2-(1-oxothietan-3-yl)ethanyl
P.459	trifluoromethyl	chloro	2-(thietan-3-yl)ethanyl
P.460	trifluoromethyl	chloro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.461	trifluoromethyl	chloro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.462	trifluoromethyl	chloro	2,2-dimethylthietan-3-yl-
P.463	trifluoromethyl	chloro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.464	trifluoromethyl	chloro	3-Methyl-1-oxo-thietan-3-yl-
P.465	trifluoromethyl	chloro	3-Methyl-thietan-3-yl-
P.466	trifluoromethyl	chloro	thietan-2-ylmethyl-
P.467	trifluoromethyl	chloro	thietan-3-yl-
P.468	trifluoromethyl	chloro	thietan-3-ylmethyl-
P.469	trifluoromethyl	cyano	(1,1-dioxothietan-2-yl)methyl-
P.470	trifluoromethyl	cyano	(1-oxothietan-2-yl)methyl-
P.471	trifluoromethyl	cyano	1,1-Dioxo-thietan-3-yl-
P.472	trifluoromethyl	cyano	1,1-Dioxo-thietan-3-ylmethyl-
P.473	trifluoromethyl	cyano	cis-1-oxo-thietan-3-yl-
P.474	trifluoromethyl	cyano	cis-1-oxo-thietan-3-ylmethyl-
P.475	trifluoromethyl	cyano	2-(1,1-dioxothietan-3-yl)ethanyl
P.476	trifluoromethyl	cyano	2-(1-oxothietan-3-yl)ethanyl
P.477	trifluoromethyl	cyano	2-(thietan-3-yl)ethanyl
P.478	trifluoromethyl	cyano	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.479	trifluoromethyl	cyano	2,2-dimethyl-1-oxo-thietan-3-yl-
P.480	trifluoromethyl	cyano	2,2-dimethylthietan-3-yl-
P.481	trifluoromethyl	cyano	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.482	trifluoromethyl	cyano	3-Methyl-1-oxo-thietan-3-yl-
P.483	trifluoromethyl	cyano	3-Methyl-thietan-3-yl-
P.484	trifluoromethyl	cyano	thietan-2-ylmethyl-
P.485	trifluoromethyl	cyano	thietan-3-yl-
P.486	trifluoromethyl	cyano	thietan-3-ylmethyl-
P.487	trifluoromethyl	cyclopropyl	(1,1-dioxothietan-2-yl)methyl-
P.488	trifluoromethyl	cyclopropyl	(1-oxothietan-2-yl)methyl-
P.489	trifluoromethyl	cyclopropyl	1,1-Dioxo-thietan-3-yl-
P.490	trifluoromethyl	cyclopropyl	1,1-Dioxo-thietan-3-ylmethyl-
P.491	trifluoromethyl	cyclopropyl	cis-1-oxo-thietan-3-yl-
P.492	trifluoromethyl	cyclopropyl	cis-1-oxo-thietan-3-ylmethyl-
P.493	trifluoromethyl	cyclopropyl	2-(1,1-dioxothietan-3-yl)ethanyl

P.494	trifluoromethyl	cyclopropyl	2-(1-oxothietan-3-yl)ethanyl
P.495	trifluoromethyl	cyclopropyl	2-(thietan-3-yl)ethanyl
P.496	trifluoromethyl	cyclopropyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.497	trifluoromethyl	cyclopropyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.498	trifluoromethyl	cyclopropyl	2,2-dimethylthietan-3-yl-
P.499	trifluoromethyl	cyclopropyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.500	trifluoromethyl	cyclopropyl	3-Methyl-1-oxo-thietan-3-yl-
P.501	trifluoromethyl	cyclopropyl	3-Methyl-thietan-3-yl-
P.502	trifluoromethyl	cyclopropyl	thietan-2-ylmethyl-
P.503	trifluoromethyl	cyclopropyl	thietan-3-yl-
P.504	trifluoromethyl	cyclopropyl	thietan-3-ylmethyl-
P.505	trifluoromethyl	ethyl	(1,1-dioxothietan-2-yl)methyl-
P.506	trifluoromethyl	ethyl	(1-oxothietan-2-yl)methyl-
P.507	trifluoromethyl	ethyl	1,1-Dioxo-thietan-3-yl-
P.508	trifluoromethyl	ethyl	1,1-Dioxo-thietan-3-ylmethyl-
P.509	trifluoromethyl	ethyl	cis-1-oxo-thietan-3-yl-
P.510	trifluoromethyl	ethyl	cis-1-oxo-thietan-3-ylmethyl-
P.511	trifluoromethyl	ethyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.512	trifluoromethyl	ethyl	2-(1-oxothietan-3-yl)ethanyl
P.513	trifluoromethyl	ethyl	2-(thietan-3-yl)ethanyl
P.514	trifluoromethyl	ethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.515	trifluoromethyl	ethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.516	trifluoromethyl	ethyl	2,2-dimethylthietan-3-yl-
P.517	trifluoromethyl	ethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.518	trifluoromethyl	ethyl	3-Methyl-1-oxo-thietan-3-yl-
P.519	trifluoromethyl	ethyl	3-Methyl-thietan-3-yl-
P.520	trifluoromethyl	ethyl	thietan-2-ylmethyl-
P.521	trifluoromethyl	ethyl	thietan-3-yl-
P.522	trifluoromethyl	ethyl	thietan-3-ylmethyl-
P.523	trifluoromethyl	fluoro	(1,1-dioxothietan-2-yl)methyl-
P.524	trifluoromethyl	fluoro	(1-oxothietan-2-yl)methyl-
P.525	trifluoromethyl	fluoro	1,1-Dioxo-thietan-3-yl-
P.526	trifluoromethyl	fluoro	1,1-Dioxo-thietan-3-ylmethyl-
P.527	trifluoromethyl	fluoro	cis-1-oxo-thietan-3-yl-
P.528	trifluoromethyl	fluoro	cis-1-oxo-thietan-3-ylmethyl-
P.529	trifluoromethyl	fluoro	2-(1,1-dioxothietan-3-yl)ethanyl
P.530	trifluoromethyl	fluoro	2-(1-oxothietan-3-yl)ethanyl
P.531	trifluoromethyl	fluoro	2-(thietan-3-yl)ethanyl
P.532	trifluoromethyl	fluoro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.533	trifluoromethyl	fluoro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.534	trifluoromethyl	fluoro	2,2-dimethylthietan-3-yl-
P.535	trifluoromethyl	fluoro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.536	trifluoromethyl	fluoro	3-Methyl-1-oxo-thietan-3-yl-
P.537	trifluoromethyl	fluoro	3-Methyl-thietan-3-yl-
P.538	trifluoromethyl	fluoro	thietan-2-ylmethyl-
P.539	trifluoromethyl	fluoro	thietan-3-yl-

P.540	trifluoromethyl	fluoro	thietan-3-ylmethyl-
P.541	trifluoromethyl	hydrogen	(1,1-dioxothietan-2-yl)methyl-
P.542	trifluoromethyl	hydrogen	(1-oxothietan-2-yl)methyl-
P.543	trifluoromethyl	hydrogen	1,1-Dioxo-thietan-3-yl-
P.544	trifluoromethyl	hydrogen	1,1-Dioxo-thietan-3-ylmethyl-
P.545	trifluoromethyl	hydrogen	cis-1-oxo-thietan-3-yl-
P.546	trifluoromethyl	hydrogen	cis-1-oxo-thietan-3-ylmethyl-
P.547	trifluoromethyl	hydrogen	2-(1,1-dioxothietan-3-yl)ethanyl
P.548	trifluoromethyl	hydrogen	2-(1-oxothietan-3-yl)ethanyl
P.549	trifluoromethyl	hydrogen	2-(thietan-3-yl)ethanyl
P.550	trifluoromethyl	hydrogen	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.551	trifluoromethyl	hydrogen	2,2-dimethyl-1-oxo-thietan-3-yl-
P.552	trifluoromethyl	hydrogen	2,2-dimethylthietan-3-yl-
P.553	trifluoromethyl	hydrogen	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.554	trifluoromethyl	hydrogen	3-Methyl-1-oxo-thietan-3-yl-
P.555	trifluoromethyl	hydrogen	3-Methyl-thietan-3-yl-
P.556	trifluoromethyl	hydrogen	thietan-2-ylmethyl-
P.557	trifluoromethyl	hydrogen	thietan-3-yl-
P.558	trifluoromethyl	hydrogen	thietan-3-ylmethyl-
P.559	trifluoromethyl	methoxy	(1,1-dioxothietan-2-yl)methyl-
P.560	trifluoromethyl	methoxy	(1-oxothietan-2-yl)methyl-
P.561	trifluoromethyl	methoxy	1,1-Dioxo-thietan-3-yl-
P.562	trifluoromethyl	methoxy	1,1-Dioxo-thietan-3-ylmethyl-
P.563	trifluoromethyl	methoxy	cis-1-oxo-thietan-3-yl-
P.564	trifluoromethyl	methoxy	cis-1-oxo-thietan-3-ylmethyl-
P.565	trifluoromethyl	methoxy	2-(1,1-dioxothietan-3-yl)ethanyl
P.566	trifluoromethyl	methoxy	2-(1-oxothietan-3-yl)ethanyl
P.567	trifluoromethyl	methoxy	2-(thietan-3-yl)ethanyl
P.568	trifluoromethyl	methoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.569	trifluoromethyl	methoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
P.570	trifluoromethyl	methoxy	2,2-dimethylthietan-3-yl-
P.571	trifluoromethyl	methoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.572	trifluoromethyl	methoxy	3-Methyl-1-oxo-thietan-3-yl-
P.573	trifluoromethyl	methoxy	3-Methyl-thietan-3-yl-
P.574	trifluoromethyl	methoxy	thietan-2-ylmethyl-
P.575	trifluoromethyl	methoxy	thietan-3-yl-
P.576	trifluoromethyl	methoxy	thietan-3-ylmethyl-
P.577	trifluoromethyl	methyl	(1,1-dioxothietan-2-yl)methyl-
P.578	trifluoromethyl	methyl	(1-oxothietan-2-yl)methyl-
P.579	trifluoromethyl	methyl	1,1-Dioxo-thietan-3-yl-
P.580	trifluoromethyl	methyl	1,1-Dioxo-thietan-3-ylmethyl-
P.581	trifluoromethyl	methyl	cis-1-oxo-thietan-3-yl-
P.582	trifluoromethyl	methyl	cis-1-oxo-thietan-3-ylmethyl-
P.583	trifluoromethyl	methyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.584	trifluoromethyl	methyl	2-(1-oxothietan-3-yl)ethanyl
P.585	trifluoromethyl	methyl	2-(thietan-3-yl)ethanyl

P.586	trifluoromethyl	methyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.587	trifluoromethyl	methyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.588	trifluoromethyl	methyl	2,2-dimethylthietan-3-yl-
P.589	trifluoromethyl	methyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.590	trifluoromethyl	methyl	3-Methyl-1-oxo-thietan-3-yl-
P.591	trifluoromethyl	methyl	3-Methyl-thietan-3-yl-
P.592	trifluoromethyl	methyl	thietan-2-ylmethyl-
P.593	trifluoromethyl	methyl	thietan-3-yl-
P.594	trifluoromethyl	methyl	thietan-3-ylmethyl-
P.595	trifluoromethyl	nitro	(1,1-dioxothietan-2-yl)methyl-
P.596	trifluoromethyl	nitro	(1-oxothietan-2-yl)methyl-
P.597	trifluoromethyl	nitro	1,1-Dioxo-thietan-3-yl-
P.598	trifluoromethyl	nitro	1,1-Dioxo-thietan-3-ylmethyl-
P.599	trifluoromethyl	nitro	cis-1-oxo-thietan-3-yl-
P.600	trifluoromethyl	nitro	cis-1-oxo-thietan-3-ylmethyl-
P.601	trifluoromethyl	nitro	2-(1,1-dioxothietan-3-yl)ethanyl
P.602	trifluoromethyl	nitro	2-(1-oxothietan-3-yl)ethanyl
P.603	trifluoromethyl	nitro	2-(thietan-3-yl)ethanyl
P.604	trifluoromethyl	nitro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.605	trifluoromethyl	nitro	2,2-dimethyl-1-oxo-thietan-3-yl-
P.606	trifluoromethyl	nitro	2,2-dimethylthietan-3-yl-
P.607	trifluoromethyl	nitro	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.608	trifluoromethyl	nitro	3-Methyl-1-oxo-thietan-3-yl-
P.609	trifluoromethyl	nitro	3-Methyl-thietan-3-yl-
P.610	trifluoromethyl	nitro	thietan-2-ylmethyl-
P.611	trifluoromethyl	nitro	thietan-3-yl-
P.612	trifluoromethyl	nitro	thietan-3-ylmethyl-
P.613	trifluoromethyl	trifluoromethoxy	(1,1-dioxothietan-2-yl)methyl-
P.614	trifluoromethyl	trifluoromethoxy	(1-oxothietan-2-yl)methyl-
P.615	trifluoromethyl	trifluoromethoxy	1,1-Dioxo-thietan-3-yl-
P.616	trifluoromethyl	trifluoromethoxy	1,1-Dioxo-thietan-3-ylmethyl-
P.617	trifluoromethyl	trifluoromethoxy	cis-1-oxo-thietan-3-yl-
P.618	trifluoromethyl	trifluoromethoxy	cis-1-oxo-thietan-3-ylmethyl-
P.619	trifluoromethyl	trifluoromethoxy	2-(1,1-dioxothietan-3-yl)ethanyl
P.620	trifluoromethyl	trifluoromethoxy	2-(1-oxothietan-3-yl)ethanyl
P.621	trifluoromethyl	trifluoromethoxy	2-(thietan-3-yl)ethanyl
P.622	trifluoromethyl	trifluoromethoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.623	trifluoromethyl	trifluoromethoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
P.624	trifluoromethyl	trifluoromethoxy	2,2-dimethylthietan-3-yl-
P.625	trifluoromethyl	trifluoromethoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.626	trifluoromethyl	trifluoromethoxy	3-Methyl-1-oxo-thietan-3-yl-
P.627	trifluoromethyl	trifluoromethoxy	3-Methyl-thietan-3-yl-
P.628	trifluoromethyl	trifluoromethoxy	thietan-2-ylmethyl-
P.629	trifluoromethyl	trifluoromethoxy	thietan-3-yl-
P.630	trifluoromethyl	trifluoromethoxy	thietan-3-ylmethyl-
P.631	trifluoromethyl	trifluoromethyl	(1,1-dioxothietan-2-yl)methyl-

P.632	trifluoromethyl	trifluoromethyl	(1-oxothietan-2-yl)methyl-
P.633	trifluoromethyl	trifluoromethyl	1,1-Dioxo-thietan-3-yl-
P.634	trifluoromethyl	trifluoromethyl	1,1-Dioxo-thietan-3-ylmethyl-
P.635	trifluoromethyl	trifluoromethyl	cis-1-oxo-thietan-3-yl-
P.636	trifluoromethyl	trifluoromethyl	cis-1-oxo-thietan-3-ylmethyl-
P.637	trifluoromethyl	trifluoromethyl	2-(1,1-dioxothietan-3-yl)ethanyl
P.638	trifluoromethyl	trifluoromethyl	2-(1-oxothietan-3-yl)ethanyl
P.639	trifluoromethyl	trifluoromethyl	2-(thietan-3-yl)ethanyl
P.640	trifluoromethyl	trifluoromethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
P.641	trifluoromethyl	trifluoromethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
P.642	trifluoromethyl	trifluoromethyl	2,2-dimethylthietan-3-yl-
P.643	trifluoromethyl	trifluoromethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
P.644	trifluoromethyl	trifluoromethyl	3-Methyl-1-oxo-thietan-3-yl-
P.645	trifluoromethyl	trifluoromethyl	3-Methyl-thietan-3-yl-
P.646	trifluoromethyl	trifluoromethyl	thietan-2-ylmethyl-
P.647	trifluoromethyl	trifluoromethyl	thietan-3-yl-
P.648	trifluoromethyl	trifluoromethyl	thietan-3-ylmethyl-
P.649	chlorodifluoromethyl	bromo	trans-1-oxo-thietan-3-yl-
P.650	chlorodifluoromethyl	chloro	trans-1-oxo-thietan-3-yl-
P.651	chlorodifluoromethyl	cyano	trans-1-oxo-thietan-3-yl-
P.652	chlorodifluoromethyl	cyclopropyl	trans-1-oxo-thietan-3-yl-
P.653	chlorodifluoromethyl	ethyl	trans-1-oxo-thietan-3-yl-
P.654	chlorodifluoromethyl	fluoro	trans-1-oxo-thietan-3-yl-
P.655	chlorodifluoromethyl	hydrogen	trans-1-oxo-thietan-3-yl-
P.656	chlorodifluoromethyl	methoxy	trans-1-oxo-thietan-3-yl-
P.657	chlorodifluoromethyl	methyl	trans-1-oxo-thietan-3-yl-
P.658	chlorodifluoromethyl	nitro	trans-1-oxo-thietan-3-yl-
P.659	chlorodifluoromethyl	trifluoromethoxy	trans-1-oxo-thietan-3-yl-
P.660	chlorodifluoromethyl	trifluoromethyl	trans-1-oxo-thietan-3-yl-
P.661	difluoromethyl	bromo	trans-1-oxo-thietan-3-yl-
P.662	difluoromethyl	chloro	trans-1-oxo-thietan-3-yl-
P.663	difluoromethyl	cyano	trans-1-oxo-thietan-3-yl-
P.664	difluoromethyl	cyclopropyl	trans-1-oxo-thietan-3-yl-
P.665	difluoromethyl	ethyl	trans-1-oxo-thietan-3-yl-
P.666	difluoromethyl	fluoro	trans-1-oxo-thietan-3-yl-
P.667	difluoromethyl	hydrogen	trans-1-oxo-thietan-3-yl-
P.668	difluoromethyl	methoxy	trans-1-oxo-thietan-3-yl-
P.669	difluoromethyl	methyl	trans-1-oxo-thietan-3-yl-
P.670	difluoromethyl	nitro	trans-1-oxo-thietan-3-yl-
P.671	difluoromethyl	trifluoromethoxy	trans-1-oxo-thietan-3-yl-
P.672	difluoromethyl	trifluoromethyl	trans-1-oxo-thietan-3-yl-
P.673	trifluoromethyl	bromo	trans-1-oxo-thietan-3-yl-
P.674	trifluoromethyl	chloro	trans-1-oxo-thietan-3-yl-
P.675	trifluoromethyl	cyano	trans-1-oxo-thietan-3-yl-
P.676	trifluoromethyl	cyclopropyl	trans-1-oxo-thietan-3-yl-
P.677	trifluoromethyl	ethyl	trans-1-oxo-thietan-3-yl-

P.678	trifluoromethyl	fluoro	trans-1-oxo-thietan-3-yl-
P.679	trifluoromethyl	hydrogen	trans-1-oxo-thietan-3-yl-
P.680	trifluoromethyl	methoxy	trans-1-oxo-thietan-3-yl-
P.681	trifluoromethyl	methyl	trans-1-oxo-thietan-3-yl-
P.682	trifluoromethyl	nitro	trans-1-oxo-thietan-3-yl-
P.683	trifluoromethyl	trifluoromethoxy	trans-1-oxo-thietan-3-yl-
P.684	trifluoromethyl	trifluoromethyl	trans-1-oxo-thietan-3-yl-
P.685	chlorodifluoromethyl	bromo	trans-1-oxo-thietan-3-ylmethyl-
P.686	chlorodifluoromethyl	chloro	trans-1-oxo-thietan-3-ylmethyl-
P.687	chlorodifluoromethyl	cyano	trans-1-oxo-thietan-3-ylmethyl-
P.688	chlorodifluoromethyl	cyclopropyl	trans-1-oxo-thietan-3-ylmethyl-
P.689	chlorodifluoromethyl	ethyl	trans-1-oxo-thietan-3-ylmethyl-
P.690	chlorodifluoromethyl	fluoro	trans-1-oxo-thietan-3-ylmethyl-
P.691	chlorodifluoromethyl	hydrogen	trans-1-oxo-thietan-3-ylmethyl-
P.692	chlorodifluoromethyl	methoxy	trans-1-oxo-thietan-3-ylmethyl-
P.693	chlorodifluoromethyl	methyl	trans-1-oxo-thietan-3-ylmethyl-
P.694	chlorodifluoromethyl	nitro	trans-1-oxo-thietan-3-ylmethyl-
P.695	chlorodifluoromethyl	trifluoromethoxy	trans-1-oxo-thietan-3-ylmethyl-
P.696	chlorodifluoromethyl	trifluoromethyl	trans-1-oxo-thietan-3-ylmethyl-
P.697	difluoromethyl	bromo	trans-1-oxo-thietan-3-ylmethyl-
P.698	difluoromethyl	chloro	trans-1-oxo-thietan-3-ylmethyl-
P.699	difluoromethyl	cyano	trans-1-oxo-thietan-3-ylmethyl-
P.700	difluoromethyl	cyclopropyl	trans-1-oxo-thietan-3-ylmethyl-
P.701	difluoromethyl	ethyl	trans-1-oxo-thietan-3-ylmethyl-
P.702	difluoromethyl	fluoro	trans-1-oxo-thietan-3-ylmethyl-
P.703	difluoromethyl	hydrogen	trans-1-oxo-thietan-3-ylmethyl-
P.704	difluoromethyl	methoxy	trans-1-oxo-thietan-3-ylmethyl-
P.705	difluoromethyl	methyl	trans-1-oxo-thietan-3-ylmethyl-
P.706	difluoromethyl	nitro	trans-1-oxo-thietan-3-ylmethyl-
P.707	difluoromethyl	trifluoromethoxy	trans-1-oxo-thietan-3-ylmethyl-
P.708	difluoromethyl	trifluoromethyl	trans-1-oxo-thietan-3-ylmethyl-
P.709	trifluoromethyl	bromo	trans-1-oxo-thietan-3-ylmethyl-
P.710	trifluoromethyl	chloro	trans-1-oxo-thietan-3-ylmethyl-
P.711	trifluoromethyl	cyano	trans-1-oxo-thietan-3-ylmethyl-
P.712	trifluoromethyl	cyclopropyl	trans-1-oxo-thietan-3-ylmethyl-
P.713	trifluoromethyl	ethyl	trans-1-oxo-thietan-3-ylmethyl-
P.714	trifluoromethyl	fluoro	trans-1-oxo-thietan-3-ylmethyl-
P.715	trifluoromethyl	hydrogen	trans-1-oxo-thietan-3-ylmethyl-
P.716	trifluoromethyl	methoxy	trans-1-oxo-thietan-3-ylmethyl-
P.717	trifluoromethyl	methyl	trans-1-oxo-thietan-3-ylmethyl-
P.718	trifluoromethyl	nitro	trans-1-oxo-thietan-3-ylmethyl-
P.719	trifluoromethyl	trifluoromethoxy	trans-1-oxo-thietan-3-ylmethyl-
P.720	trifluoromethyl	trifluoromethyl	trans-1-oxo-thietan-3-ylmethyl-

Table 1P:

Table 1 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table

P.

Table 2P:

Table 2 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 3P:

Table 3 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 4P:

Table 4 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 5P:

Table 5 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 6P:

20 Table 6 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 7P:

25 Table 7 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 8P:

30 Table 8 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 9P:

35 Table 9 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

i Table 10P:

40 Table 10 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 11P:

45 Table 11 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 12P:

50 Table 12 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 13P:

55 Table 13 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 14P:

Table 14 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 15P:

Table 15 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 16P:

Table 16 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 17P:

Table 17 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 18P:

Table 18 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 19P:

Table 19 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 20P:

Table 20 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 21P:

Table 21 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 22P:

Table 22 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 23P:

Table 23 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 24P:

Table 24 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 25P:

Table 25 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 26P:

Table 26 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 27P:

Table 27 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 28P:

Table 28 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 29P:

Table 29 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 30P:

Table 30 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 31P:

Table 31 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 32P:

Table 32 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 33P:

Table 33 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 34P:

45 Table 34 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 35P:

50 Table 35 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 36P:

55 Table 36 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 37P:

Table 37 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 38P:

Table 38 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 39P:

Table 39 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 40P:

Table 40 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 41P:

Table 41 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 42P:

Table 42 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 43P:

Table 43 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 44P:

Table 44 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 45P:

Table 45 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 46P:

Table 46 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 47P:

Table 47 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 48P:

Table 48 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 49P:

Table 49 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 50P:

Table 50 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 51P:

Table 51 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 52P:

Table 52 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 53P:

15 Table 53 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 54P:

20 Table 54 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 55P:

25 Table 55 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 56P:

30 Table 56 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 57P:

35 Table 57 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 58P:

40 Table 58 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 59P:

45 Table 59 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 60P:

50 Table 60 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 61P:

55 Table 61 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 62P:

Table 62 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 63P:

Table 63 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 64P:

Table 64 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 65P:

Table 65 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 66P:

Table 66 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 67P:

Table 67 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 68P:

Table 68 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 69P:

Table 69 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 70P:

Table 70 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 71P:

Table 71 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 72P:

Table 72 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 73P:

Table 73 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 74P:

Table 74 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 75P:

Table 75 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 76P:

Table 76 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 77P:

Table 77 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 78P:

Table 78 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 79P:

Table 79 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 80P:

Table 80 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 81P:

Table 81 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 82P:

Table 82 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 83P:

Table 83 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 84P:

Table 84 P provides 720 compounds of formula (I-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 85P:

Table 85 P provides 720 compounds of formula (I-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 86P:

Table 86 P provides 720 compounds of formula (I-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

10 Table 87P:

Table 87 P provides 720 compounds of formula (I-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 88P:

Table 88 P provides 720 compounds of formula (I-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 89P:

Table 89 P provides 720 compounds of formula (I-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 90P:

Table 90 P provides 720 compounds of formula (I-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.


30 Table 91P:

Table 91 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 92 P:

Table 92 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 93 P:

Table 93 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 94 P:

Table 94 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 95 P:

Table 95 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 96 P:

Table 96 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 97 P:

Table 97 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 98 P:

Table 98 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 99 P:

Table 99 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 100 P:

Table 100 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 101 P:

Table 101 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 102 P:

Table 102 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 103 P:

Table 103 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 104 P:

Table 104 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 105 P:

Table 105 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 106 P:

Table 106 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 107 P:

Table 107 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 108 P:

Table 108 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 109 P:

Table 109 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 110 P:

Table 110 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 111 P:

Table 111 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 112 P:

Table 112 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 113 P:

Table 113 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 114 P:

Table 114 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 115 P:

Table 115 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 116 P:

Table 116 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 117 P:

Table 27 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P

10 Table 118 P:

Table 118 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 119 P:

Table 119 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 120 P:

Table 120 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Cl, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 121 P:

Table 121 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

30 Table 122 P:

Table 122 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 123 P:

Table 123 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 124 P:

Table 124 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

45 Table 125 P:

Table 125 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Br, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

50 Table 126 P:

Table 126 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

55 Table 127 P:

Table 127 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 128 P:

Table 128 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

10 Table 129 P:

Table 129 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 130 P:

Table 130 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-I, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 131 P:

Table 131 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 132 P:

Table 132 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

30 Table 133 P:

Table 133 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 134 P:

Table 134 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 135 P:

Table 135 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

45 Table 136 P:

Table 136 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

50 Table 137 P:

Table 137 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

55 Table 138 P:

Table 138 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 139 P:

Table 139 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 140 P:

Table 140 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 141 P:

Table 141 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 142 P:

Table 142 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 143 P:

Table 143 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 144 P:

Table 144 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 145 P:

Table 145 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 146 P:

Table 146 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 147 P:

Table 147 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 148 P:

Table 148 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 149 P:

Table 149 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 150 P:

Table 150 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 151 P:

Table 151 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 152 P:

Table 152 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 153 P:

Table 153 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 154 P:

Table 154 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 155 P:

Table 155 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 156 P:

Table 156 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 157 P:

Table 157 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 158 P:

Table 158 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 159 P:

Table 159 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 160 P:

Table 160 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 161 P:

Table 161 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

10 Table 162 P:

Table 162 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 163 P:

Table 163 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 164 P:

Table 164 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 165 P:

Table 165 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

30 Table 166 P:

Table 166 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 167 P:

Table 167 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 168 P:

Table 168 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

45 Table 169 P:

Table 169 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

50 Table 170 P:

Table 170 P provides 720 compounds of formula (X1-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

55 Table 171 P:

Table 171 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 172 P:

Table 172 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 173 P:

Table 173 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 174 P:

Table 174 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 175 P:

Table 175 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 176 P:

Table 176 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 177 P:

Table 177 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 178 P:

Table 178 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 179 P:

Table 179 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 180 P:

Table 180 P provides 720 compounds of formula (X1-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.


Table 181 P:

Table 181 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 182 P:

Table 182 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 183 P:

Table 183 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 184 P:

Table 184 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 185 P:

Table 185 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 186 P:

Table 186 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 187 P:

Table 187 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 188 P:

Table 188 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 189 P:

Table 189 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 190 P:

Table 190 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 191 P:

Table 191 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 192 P:

Table 192 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 193 P:

Table 193 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 194 P:

Table 194 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 195 P:

Table 195 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 196 P:

Table 196 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 197 P:

Table 197 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 198 P:

Table 198 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 199 P:

Table 199 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 200 P:

Table 200 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 201 P:

Table 201 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 202 P:

Table 202 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 203 P:

Table 203 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 204 P:

Table 204 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 205 P:

Table 205 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 206 P:

Table 206 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 207 P:

Table 207 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 208 P:

Table 208 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 209 P:

Table 209 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 210 P:

Table 210 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 211 P:

Table 211 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 212 P:

Table 212 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 213 P:

Table 213 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 214 P:

Table 214 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 215 P:

Table 215 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 216 P:

Table 216 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 217 P:

Table 217 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 218 P:

Table 218 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 219 P:

Table 219 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 220 P:

Table 220 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 221 P:

Table 221 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 222 P:

Table 222 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 223 P:

Table 223 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 224 P:

Table 224 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 225 P:

Table 225 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 226 P:

Table 226 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 227 P:

Table 227 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 228 P:

Table 228 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 229 P:

Table 229 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 230 P:

Table 230 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 231 P:

Table 231 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 232 P:

Table 232 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 233 P:

Table 233 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 234 P:

Table 234 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 235 P:

Table 235 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 236 P:

Table 236 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 237 P:

Table 237 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 238 P:

Table 238 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 239 P:

Table 239 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 240 P:

Table 240 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 241 P:

Table 241 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 242 P:

Table 242 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 243 P:

Table 243 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 244 P:

Table 244 P provides 720 compounds of formula (X2-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 245 P:

Table 245 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 246 P:

Table 246 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

10 Table 247 P:

Table 247 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 248 P:

Table 248 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 249 P:

Table 249 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 250 P:

Table 250 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

30 Table 251 P:

Table 251 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 252 P:

Table 252 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 253 P:

Table 253 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

45 Table 254 P:

Table 254 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

50 Table 255 P:

Table 255 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is C-Cl, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

55 Table 256 P:

Table 256 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 257 P:

Table 257 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

10 Table 258 P:

Table 258 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 259 P:

Table 259 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 260 P:

Table 260 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is CH, X^3 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 261 P:

Table 261 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

30 Table 262 P:

Table 262 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 263 P:

Table 263 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 264 P:

Table 264 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

45 Table 265 P:

Table 265 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is N, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

50 Table 266 P:

Table 266 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 267 P:

Table 267 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 268 P:

Table 268 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 269 P:

Table 269 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 270 P:

Table 270 P provides 720 compounds of formula (X2-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is trifluoromethyl, X^2 is N, X^3 is trifluoromethyl, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.


Table 271 P:

Table 271 P provides 720 compounds of formula (X3-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 272 P:

Table 272 P provides 720 compounds of formula (X3-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 273 P:

Table 273 P provides 720 compounds of formula (X3-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 274 P:

Table 274 P provides 720 compounds of formula (X3-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

Table 275 P:

Table 275 P provides 720 compounds of formula (X3-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is chloro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

Table 276 P:

Table 276 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 277 P:

Table 277 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 278 P:

Table 278 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 279 P:

Table 279 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 280 P:

Table 280 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 281 P:

Table 281 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 282 P:

Table 282 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 283 P:

Table 283 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 284 P:

Table 284 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 285 P:

Table 285 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 286 P:

Table 286 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 287 P:

Table 287 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 288 P:

Table 288 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 289 P:

Table 289 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 290 P:

Table 290 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 291 P:

Table 291 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 292 P:

Table 292 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 293 P:

Table 293 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 294 P:

Table 294 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 295 P:

Table 295 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 296 P:

Table 296 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 297 P:

Table 297 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 298 P:

Table 298 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 299 P:

Table 299 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 300 P:

Table 300 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 301 P:

Table 301 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 302 P:

Table 302 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 303 P:

Table 303 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 304 P:

Table 304 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 305 P:

Table 305 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 306 P:

Table 306 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 307 P:

Table 307 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 308 P:

Table 308 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 309 P:

Table 309 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 310 P:

Table 310 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 311 P:

Table 311 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 312 P:

Table 312 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 313 P:

Table 313 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 314 P:

Table 314 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 315 P:

Table 315 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 316 P:

Table 316 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 317 P:

Table 317 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 318 P:

Table 318 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 319 P:

Table 319 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 320 P:

Table 320 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is bromo, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 321 P:

Table 321 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 322 P:

Table 322 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 323 P:

Table 323 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 324 P:

Table 324 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 325 P:

Table 325 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 326 P:

Table 326 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 327 P:

Table 327 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 328 P:

Table 328 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 329 P:

Table 329 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 330 P:

Table 330 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 331 P:

Table 331 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 332 P:

Table 332 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 333 P:

Table 333 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 334 P:

Table 334 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 335 P:

Table 335 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 336 P:

Table 336 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 337 P:

Table 337 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 338 P:

Table 338 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 339 P:

Table 339 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 340 P:

Table 340 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 341 P:

Table 341 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 342 P:

- Table 342 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 5 Table 343 P:
Table 343 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 10 Table 344 P:
Table 344 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 15 Table 345 P:
Table 345 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.
- 20 Table 346 P:
Table 346 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 25 Table 347 P:
Table 347 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 30 Table 348 P:
Table 348 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 35 Table 349 P:
Table 349 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 40 Table 350 P:
Table 350 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.
- 45 Table 351 P:
Table 351 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.
- 50 Table 352 P:
Table 352 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 353 P:

Table 353 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 354 P:

Table 354 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 355 P:

Table 355 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 356 P:

Table 356 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 357 P:

Table 357 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 358 P:

Table 358 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 359 P:

Table 359 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 360 P:

Table 360 P provides 720 compounds of formula (X3-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.




Table 361 P:

Table 361 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 362 P:

Table 362 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 363 P:

Table 363 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 364 P:

Table 364 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 365 P:

Table 365 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 366 P:

Table 366 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 367 P:

Table 367 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 368 P:

Table 368 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 369 P:

Table 369 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 370 P:

Table 370 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

Table 371 P:

Table 371 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 372 P:

Table 372 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 373 P:

Table 373 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 374 P:

Table 374 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 375 P:

Table 375 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 376 P:

Table 376 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 377 P:

Table 377 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 378 P:

Table 378 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 379 P:

Table 19 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 380 P:

Table 380 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 381 P:

Table 381 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 382 P:

Table 382 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 383 P:

Table 383 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 384 P:

Table 384 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 385 P:

Table 385 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-F, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 386 P:

Table 386 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 387 P:

Table 387 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 388 P:

Table 388 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 389 P:

Table 389 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 390 P:

Table 390 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 391 P:

Table 391 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 392 P:

Table 392 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 393 P:

Table 393 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 394 P:

Table 394 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 395 P:

Table 395 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Br, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 396 P:

Table 396 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 397 P:

Table 397 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 398 P:

Table 398 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 399 P:

Table 399 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 400 P:

Table 400 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-I, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 401 P:

Table 401 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 402 P:

Table 402 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 403 P:

Table 403 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 404 P:

Table 404 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is fluoro, X² is C-F, X³ is fluoro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 405 P:

Table 405 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is fluoro, X^2 is C-F, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

5 Table 406 P:

Table 406 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

10 Table 407 P:

Table 407 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

15 Table 408 P:

Table 408 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

20 Table 409 P:

Table 409 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

25 Table 410 P:

Table 410 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is bromo, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

30 Table 411 P:

Table 411 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

35 Table 412 P:

Table 412 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is N, Y^2 is CH, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

40 Table 413 P:

Table 413 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is N, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

45 Table 414 P:

Table 414 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is CH, Y^2 is N, Y^3 is CH and X^4 , R^5 and R^2 have the values listed in the table P.

50 Table 415 P:

Table 415 P provides 720 compounds of formula (X4-A) wherein G^1 is oxygen, R^1 is hydrogen, X^1 is chloro, X^2 is CH, X^3 is fluoro, Y^1 is CH, Y^2 is CH, Y^3 is N and X^4 , R^5 and R^2 have the values listed in the table P.

55 Table 416 P:

Table 416 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 417 P:

Table 417 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 418 P:

Table 418 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 419 P:

Table 419 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 420 P:

Table 420 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 421 P:

Table 421 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 422 P:

Table 422 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 423 P:

Table 423 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 424 P:

Table 424 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 425 P:

Table 425 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 426 P:

Table 426 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55 Table 427 P:

Table 427 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5 Table 428 P:

Table 428 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10 Table 429 P:

Table 429 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

15 Table 430 P:

Table 430 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

20 Table 431 P:

Table 431 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25 Table 432 P:

Table 432 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30 Table 433 P:

Table 433 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35 Table 434 P:

Table 434 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

40 Table 435 P:

Table 435 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

45 Table 436 P:

Table 436 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50 Table 437 P:

Table 437 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

Table 438 P:

Table 438 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 439 P:

Table 439 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

10

Table 440 P:

Table 440 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

15

Table 441 P:

Table 441 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

20

Table 442 P:

Table 442 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

25

Table 443 P:

Table 443 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

30

Table 444 P:

Table 444 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

35

Table 445 P:

Table 445 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is chloro, X² is N, X³ is chloro, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

40

Table 446 P:

Table 446 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

45

Table 447 P:

Table 447 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is N, Y² is CH, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

50

Table 448 P:

Table 448 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is N, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

55

Table 449 P:

Table 449 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is N, Y³ is CH and X⁴, R⁵ and R² have the values listed in the table P.

5

Table 450 P:

Table 450 P provides 720 compounds of formula (X4-A) wherein G¹ is oxygen, R¹ is hydrogen, X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl, Y¹ is CH, Y² is CH, Y³ is N and X⁴, R⁵ and R² have the values listed in the table P.

10




Table X

	R5	R2
X.001	bromo	(1,1-dioxothietan-2-yl)methyl-
X.002	bromo	(1-oxothietan-2-yl)methyl-
X.003	bromo	1,1-Dioxo-thietan-3-yl-
X.004	bromo	1,1-Dioxo-thietan-3-ylmethyl-
X.005	bromo	cis-1-oxo-thietan-3-yl-
X.006	bromo	cis-1-oxo-thietan-3-ylmethyl-
X.007	bromo	2-(1,1-dioxothietan-3-yl)ethanyl
X.008	bromo	2-(1-oxothietan-3-yl)ethanyl
X.009	bromo	2-(thietan-3-yl)ethanyl
X.010	bromo	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.011	bromo	2,2-dimethyl-1-oxo-thietan-3-yl-
X.012	bromo	2,2-dimethylthietan-3-yl-
X.013	bromo	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.014	bromo	3-Methyl-1-oxo-thietan-3-yl-
X.015	bromo	3-Methyl-thietan-3-yl-
X.016	bromo	thietan-2-ylmethyl-
X.017	bromo	thietan-3-yl-
X.018	bromo	thietan-3-ylmethyl-
X.019	chloro	(1,1-dioxothietan-2-yl)methyl-
X.020	chloro	(1-oxothietan-2-yl)methyl-
X.021	chloro	1,1-Dioxo-thietan-3-yl-
X.022	chloro	1,1-Dioxo-thietan-3-ylmethyl-
X.023	chloro	cis-1-oxo-thietan-3-yl-
X.024	chloro	cis-1-oxo-thietan-3-ylmethyl-
X.025	chloro	2-(1,1-dioxothietan-3-yl)ethanyl
X.026	chloro	2-(1-oxothietan-3-yl)ethanyl
X.027	chloro	2-(thietan-3-yl)ethanyl
X.028	chloro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.029	chloro	2,2-dimethyl-1-oxo-thietan-3-yl-

X.030	chloro	2,2-dimethylthietan-3-yl-
X.031	chloro	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.032	chloro	3-Methyl-1-oxo-thietan-3-yl-
X.033	chloro	3-Methyl-thietan-3-yl-
X.034	chloro	thietan-2-ylmethyl-
X.035	chloro	thietan-3-yl-
X.036	chloro	thietan-3-ylmethyl-
X.037	cyano	(1,1-dioxothietan-2-yl)methyl-
X.038	cyano	(1-oxothietan-2-yl)methyl-
X.039	cyano	1,1-Dioxo-thietan-3-yl-
X.040	cyano	1,1-Dioxo-thietan-3-ylmethyl-
X.041	cyano	cis-1-oxo-thietan-3-yl-
X.042	cyano	cis-1-oxo-thietan-3-ylmethyl-
X.043	cyano	2-(1,1-dioxothietan-3-yl)ethanyl
X.044	cyano	2-(1-oxothietan-3-yl)ethanyl
X.045	cyano	2-(thietan-3-yl)ethanyl
X.046	cyano	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.047	cyano	2,2-dimethyl-1-oxo-thietan-3-yl-
X.048	cyano	2,2-dimethylthietan-3-yl-
X.049	cyano	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.050	cyano	3-Methyl-1-oxo-thietan-3-yl-
X.051	cyano	3-Methyl-thietan-3-yl-
X.052	cyano	thietan-2-ylmethyl-
X.053	cyano	thietan-3-yl-
X.054	cyano	thietan-3-ylmethyl-
X.055	cyclopropyl	(1,1-dioxothietan-2-yl)methyl-
X.056	cyclopropyl	(1-oxothietan-2-yl)methyl-
X.057	cyclopropyl	1,1-Dioxo-thietan-3-yl-
X.058	cyclopropyl	1,1-Dioxo-thietan-3-ylmethyl-
X.059	cyclopropyl	cis-1-oxo-thietan-3-yl-
X.060	cyclopropyl	cis-1-oxo-thietan-3-ylmethyl-
X.061	cyclopropyl	2-(1,1-dioxothietan-3-yl)ethanyl
X.062	cyclopropyl	2-(1-oxothietan-3-yl)ethanyl
X.063	cyclopropyl	2-(thietan-3-yl)ethanyl
X.064	cyclopropyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.065	cyclopropyl	2,2-dimethyl-1-oxo-thietan-3-yl-
X.066	cyclopropyl	2,2-dimethylthietan-3-yl-
X.067	cyclopropyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.068	cyclopropyl	3-Methyl-1-oxo-thietan-3-yl-
X.069	cyclopropyl	3-Methyl-thietan-3-yl-
X.070	cyclopropyl	thietan-2-ylmethyl-
X.071	cyclopropyl	thietan-3-yl-
X.072	cyclopropyl	thietan-3-ylmethyl-
X.073	ethyl	(1,1-dioxothietan-2-yl)methyl-
X.074	ethyl	(1-oxothietan-2-yl)methyl-

X.075	ethyl	1,1-Dioxo-thietan-3-yl-
X.076	ethyl	1,1-Dioxo-thietan-3-ylmethyl-
X.077	ethyl	cis-1-oxo-thietan-3-yl-
X.078	ethyl	cis-1-oxo-thietan-3-ylmethyl-
X.079	ethyl	2-(1,1-dioxothietan-3-yl)ethanyl
X.080	ethyl	2-(1-oxothietan-3-yl)ethanyl
X.081	ethyl	2-(thietan-3-yl)ethanyl
X.082	ethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.083	ethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
X.084	ethyl	2,2-dimethylthietan-3-yl-
X.085	ethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.086	ethyl	3-Methyl-1-oxo-thietan-3-yl-
X.087	ethyl	3-Methyl-thietan-3-yl-
X.088	ethyl	thietan-2-ylmethyl-
X.089	ethyl	thietan-3-yl-
X.090	ethyl	thietan-3-ylmethyl-
X.091	fluoro	(1,1-dioxothietan-2-yl)methyl-
X.092	fluoro	(1-oxothietan-2-yl)methyl-
X.093	fluoro	1,1-Dioxo-thietan-3-yl-
X.094	fluoro	1,1-Dioxo-thietan-3-ylmethyl-
X.095	fluoro	cis-1-oxo-thietan-3-yl-
X.096	fluoro	cis-1-oxo-thietan-3-ylmethyl-
X.097	fluoro	2-(1,1-dioxothietan-3-yl)ethanyl
X.098	fluoro	2-(1-oxothietan-3-yl)ethanyl
X.099	fluoro	2-(thietan-3-yl)ethanyl
X.100	fluoro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.101	fluoro	2,2-dimethyl-1-oxo-thietan-3-yl-
X.102	fluoro	2,2-dimethylthietan-3-yl-
X.103	fluoro	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.104	fluoro	3-Methyl-1-oxo-thietan-3-yl-
X.105	fluoro	3-Methyl-thietan-3-yl-
X.106	fluoro	thietan-2-ylmethyl-
X.107	fluoro	thietan-3-yl-
X.108	fluoro	thietan-3-ylmethyl-
X.109	hydrogen	(1,1-dioxothietan-2-yl)methyl-
X.110	hydrogen	(1-oxothietan-2-yl)methyl-
X.111	hydrogen	1,1-Dioxo-thietan-3-yl-
X.112	hydrogen	1,1-Dioxo-thietan-3-ylmethyl-
X.113	hydrogen	cis-1-oxo-thietan-3-yl-
X.114	hydrogen	cis-1-oxo-thietan-3-ylmethyl-
X.115	hydrogen	2-(1,1-dioxothietan-3-yl)ethanyl
X.116	hydrogen	2-(1-oxothietan-3-yl)ethanyl
X.117	hydrogen	2-(thietan-3-yl)ethanyl
X.118	hydrogen	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-

X.119	hydrogen	2,2-dimethyl-1-oxo-thietan-3-yl-
X.120	hydrogen	2,2-dimethylthietan-3-yl-
X.121	hydrogen	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.122	hydrogen	3-Methyl-1-oxo-thietan-3-yl-
X.123	hydrogen	3-Methyl-thietan-3-yl-
X.124	hydrogen	thietan-2-ylmethyl-
X.125	hydrogen	thietan-3-yl-
X.126	hydrogen	thietan-3-ylmethyl-
X.127	methoxy	(1,1-dioxothietan-2-yl)methyl-
X.128	methoxy	(1-oxothietan-2-yl)methyl-
X.129	methoxy	1,1-Dioxo-thietan-3-yl-
X.130	methoxy	1,1-Dioxo-thietan-3-ylmethyl-
X.131	methoxy	cis-1-oxo-thietan-3-yl-
X.132	methoxy	cis-1-oxo-thietan-3-ylmethyl-
X.133	methoxy	2-(1,1-dioxothietan-3-yl)ethanyl
X.134	methoxy	2-(1-oxothietan-3-yl)ethanyl
X.135	methoxy	2-(thietan-3-yl)ethanyl
X.136	methoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.137	methoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
X.138	methoxy	2,2-dimethylthietan-3-yl-
X.139	methoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.140	methoxy	3-Methyl-1-oxo-thietan-3-yl-
X.141	methoxy	3-Methyl-thietan-3-yl-
X.142	methoxy	thietan-2-ylmethyl-
X.143	methoxy	thietan-3-yl-
X.144	methoxy	thietan-3-ylmethyl-
X.145	methyl	(1,1-dioxothietan-2-yl)methyl-
X.146	methyl	(1-oxothietan-2-yl)methyl-
X.147	methyl	1,1-Dioxo-thietan-3-yl-
X.148	methyl	1,1-Dioxo-thietan-3-ylmethyl-
X.149	methyl	cis-1-oxo-thietan-3-yl-
X.150	methyl	cis-1-oxo-thietan-3-ylmethyl-
X.151	methyl	2-(1,1-dioxothietan-3-yl)ethanyl
X.152	methyl	2-(1-oxothietan-3-yl)ethanyl
X.153	methyl	2-(thietan-3-yl)ethanyl
X.154	methyl	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.155	methyl	2,2-dimethyl-1-oxo-thietan-3-yl-
X.156	methyl	2,2-dimethylthietan-3-yl-
X.157	methyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.158	methyl	3-Methyl-1-oxo-thietan-3-yl-
X.159	methyl	3-Methyl-thietan-3-yl-
X.160	methyl	thietan-2-ylmethyl-
X.161	methyl	thietan-3-yl-
X.162	methyl	thietan-3-ylmethyl-
X.163	nitro	(1,1-dioxothietan-2-yl)methyl-

X.164	nitro	(1-oxothietan-2-yl)methyl-
X.165	nitro	1,1-Dioxo-thietan-3-yl-
X.166	nitro	1,1-Dioxo-thietan-3-ylmethyl-
X.167	nitro	cis-1-oxo-thietan-3-yl-
X.168	nitro	cis-1-oxo-thietan-3-ylmethyl-
X.169	nitro	2-(1,1-dioxothietan-3-yl)ethanyl
X.170	nitro	2-(1-oxothietan-3-yl)ethanyl
X.171	nitro	2-(thietan-3-yl)ethanyl
X.172	nitro	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.173	nitro	2,2-dimethyl-1-oxo-thietan-3-yl-
X.174	nitro	2,2-dimethylthietan-3-yl-
X.175	nitro	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.176	nitro	3-Methyl-1-oxo-thietan-3-yl-
X.177	nitro	3-Methyl-thietan-3-yl-
X.178	nitro	thietan-2-ylmethyl-
X.179	nitro	thietan-3-yl-
X.180	nitro	thietan-3-ylmethyl-
X.181	trifluoromethoxy	(1,1-dioxothietan-2-yl)methyl-
X.182	trifluoromethoxy	(1-oxothietan-2-yl)methyl-
X.183	trifluoromethoxy	1,1-Dioxo-thietan-3-yl-
X.184	trifluoromethoxy	1,1-Dioxo-thietan-3-ylmethyl-
X.185	trifluoromethoxy	cis-1-oxo-thietan-3-yl-
X.186	trifluoromethoxy	cis-1-oxo-thietan-3-ylmethyl-
X.187	trifluoromethoxy	2-(1,1-dioxothietan-3-yl)ethanyl
X.188	trifluoromethoxy	2-(1-oxothietan-3-yl)ethanyl
X.189	trifluoromethoxy	2-(thietan-3-yl)ethanyl
X.190	trifluoromethoxy	2,2-dimethyl-1,1-Dioxo-thietan-3-yl-
X.191	trifluoromethoxy	2,2-dimethyl-1-oxo-thietan-3-yl-
X.192	trifluoromethoxy	2,2-dimethylthietan-3-yl-
X.193	trifluoromethoxy	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.194	trifluoromethoxy	3-Methyl-1-oxo-thietan-3-yl-
X.195	trifluoromethoxy	3-Methyl-thietan-3-yl-
X.196	trifluoromethoxy	thietan-2-ylmethyl-
X.197	trifluoromethoxy	thietan-3-yl-
X.198	trifluoromethoxy	thietan-3-ylmethyl-
X.199	trifluoromethyl	(1,1-dioxothietan-2-yl)methyl-
X.200	trifluoromethyl	(1-oxothietan-2-yl)methyl-
X.201	trifluoromethyl	1,1-Dioxo-thietan-3-yl-
X.202	trifluoromethyl	1,1-Dioxo-thietan-3-ylmethyl-
X.203	trifluoromethyl	cis-1-oxo-thietan-3-yl-
X.204	trifluoromethyl	cis-1-oxo-thietan-3-ylmethyl-
X.205	trifluoromethyl	2-(1,1-dioxothietan-3-yl)ethanyl
X.206	trifluoromethyl	2-(1-oxothietan-3-yl)ethanyl
X.207	trifluoromethyl	2-(thietan-3-yl)ethanyl
X.208	trifluoromethyl	2,2-dimethyl-1,1-Dioxo-thietan-3-

		yl-
X.209	trifluoromethyl	2,2-dimethyl-1-oxo-thietan-3-yl-
X.210	trifluoromethyl	2,2-dimethylthietan-3-yl-
X.211	trifluoromethyl	3-Methyl-1,1-Dioxo-thietan-3-yl-
X.212	trifluoromethyl	3-Methyl-1-oxo-thietan-3-yl-
X.213	trifluoromethyl	3-Methyl-thietan-3-yl-
X.214	trifluoromethyl	thietan-2-ylmethyl-
X.215	trifluoromethyl	thietan-3-yl-
X.216	trifluoromethyl	thietan-3-ylmethyl-
X.217	bromo	trans-1-oxo-thietan-3-yl-
X.218	chloro	trans-1-oxo-thietan-3-yl-
X.219	cyano	trans-1-oxo-thietan-3-yl-
X.220	cyclopropyl	trans-1-oxo-thietan-3-yl-
X.221	ethyl	trans-1-oxo-thietan-3-yl-
X.222	fluoro	trans-1-oxo-thietan-3-yl-
X.223	hydrogen	trans-1-oxo-thietan-3-yl-
X.224	methoxy	trans-1-oxo-thietan-3-yl-
X.225	methyl	trans-1-oxo-thietan-3-yl-
X.226	nitro	trans-1-oxo-thietan-3-yl-
X.227	trifluoromethoxy	trans-1-oxo-thietan-3-yl-
X.228	trifluoromethyl	trans-1-oxo-thietan-3-yl-
X.229	bromo	trans-1-oxo-thietan-3-ylmethyl-
X.230	chloro	trans-1-oxo-thietan-3-ylmethyl-
X.231	cyano	trans-1-oxo-thietan-3-ylmethyl-
X.232	cyclopropyl	trans-1-oxo-thietan-3-ylmethyl-
X.233	ethyl	trans-1-oxo-thietan-3-ylmethyl-
X.234	fluoro	trans-1-oxo-thietan-3-ylmethyl-
X.235	hydrogen	trans-1-oxo-thietan-3-ylmethyl-
X.236	methoxy	trans-1-oxo-thietan-3-ylmethyl-
X.237	methyl	trans-1-oxo-thietan-3-ylmethyl-
X.238	nitro	trans-1-oxo-thietan-3-ylmethyl-
X.239	trifluoromethoxy	trans-1-oxo-thietan-3-ylmethyl-
X.240	trifluoromethyl	trans-1-oxo-thietan-3-ylmethyl-

Table 1X:

Table 1X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH, X⁵ is chloro, R⁵ and R² have the values listed in the table X.

5

Table 2 X:

Table 2X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH, X⁵ is bromo, R⁵ and R² have the values listed in the table X.

10 Table 3 X:

Table 3X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH, X⁵ is iodo, R⁵ and R² have the values listed in the table X.

Table 4 X:

15 Table 4X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is CH, Y² is CH, Y³ is CH, X⁵ is amino, R⁵ and R² have the values listed in the table X.

Table 5 X:

Table 5X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is CHO, R^5 and R^2 have the values listed in the table X.

5 Table 6 X:

Table 6X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is CN, R^5 and R^2 have the values listed in the table X.

Table 7 X:

10 Table 7X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is OH, R^5 and R^2 have the values listed in the table X.

Table 8 X:

15 Table 8X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=O)OH, R^5 and R^2 have the values listed in the table X.

Table 9 X:

20 Table 9X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=NOH)H, R^5 and R^2 have the values listed in the table X.

Table 10 X:

Table 10X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=NOH)Cl, R^5 and R^2 have the values listed in the table X.

25 Table 11 X:

Table 11X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=NOH)NH₂, R^5 and R^2 have the values listed in the table X.

Table 12 X:

30 Table 12X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=O)CH₃, R^5 and R^2 have the values listed in the table X.

Table 13 X:

35 Table 13X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=NOH)CH₃, R^5 and R^2 have the values listed in the table X.

Table 14 X:

40 Table 14X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=O)CH₂Cl, R^5 and R^2 have the values listed in the table X.

Table 15 X:

Table 15X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is CH, Y^2 is CH, Y^3 is CH, X^5 is C(=O)CH₂Br, R^5 and R^2 have the values listed in the table X.

45 Table 16 X:

Table 16X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH, X^5 is chloro, R^5 and R^2 have the values listed in the table X.

Table 17 X:

50 Table 17X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH, X^5 is bromo, R^5 and R^2 have the values listed in the table X.

Table 18 X:

55 Table 18X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y^1 is N, Y^2 is CH, Y^3 is CH, X^5 is iodo, R^5 and R^2 have the values listed in the table X.

Table 19 X:

Table 19X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is amino, R⁵ and R² have the values listed in the table X.

5 Table 20 X:

Table 20X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is CHO, R⁵ and R² have the values listed in the table X.

Table 21 X:

10 Table 21X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is CN, R⁵ and R² have the values listed in the table X.

Table 22 X:

15 Table 22X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is OH, R⁵ and R² have the values listed in the table X.

Table 23 X:

20 Table 23X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=O)OH, R⁵ and R² have the values listed in the table X.

Table 24 X:

Table 24X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=NOH)H, R⁵ and R² have the values listed in the table X.

25 Table 25 X:

Table 25X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=NOH)Cl, R⁵ and R² have the values listed in the table X.

Table 26 X:

30 Table 26X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=NOH)NH₂, R⁵ and R² have the values listed in the table X.

Table 27 X:

35 Table 27X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=O)CH₃, R⁵ and R² have the values listed in the table X.

Table 28 X:

40 Table 28X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=NOH)CH₃, R⁵ and R² have the values listed in the table X.

Table 29 X:

Table 29X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=O)CH₂Cl, R⁵ and R² have the values listed in the table X.

45 Table 30 X:

Table 30X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is CH, Y³ is CH, X⁵ is C(=O)CH₂Br, R⁵ and R² have the values listed in the table X.

Table 31 X:

50 Table 31X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is N, Y³ is CH, X⁵ is chloro, R⁵ and R² have the values listed in the table X.

Table 32 X:

55 Table 32X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y¹ is N, Y² is N, Y³ is CH, X⁵ is bromo, R⁵ and R² have the values listed in the table X.

Table 33 X:

Table 33X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is iodo, R⁵ and R² have the values listed in the table X.

5 Table 34 X:

Table 34X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is amino, R⁵ and R² have the values listed in the table X.

Table 35 X:

10 Table 35X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is CHO, R⁵ and R² have the values listed in the table X.

Table 36 X:

15 Table 36X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is CN, R⁵ and R² have the values listed in the table X.

Table 37 X:

20 Table 37X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is OH, R⁵ and R² have the values listed in the table X.

Table 38 X:

Table 38X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=O)OH, R⁵ and R² have the values listed in the table X.

25 Table 39 X:

Table 39X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=NOH)H, R⁵ and R² have the values listed in the table X.

Table 40 X:

30 Table 40X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=NOH)Cl, R⁵ and R² have the values listed in the table X.

Table 41 X:

35 Table 41X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=NOH)NH₂, R⁵ and R² have the values listed in the table X.

Table 42 X:

40 Table 42X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=O)CH₃, R⁵ and R² have the values listed in the table X.

Table 43 X:

Table 43X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=NOH)CH₃, R⁵ and R² have the values listed in the table X.

45 Table 44 X:

Table 44X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=O)CH₂Cl, R⁵ and R² have the values listed in the table X.

Table 45 X:

50 Table 45X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is N, Y2 is N, Y3 is CH, X⁵ is C(=O)CH₂Br, R⁵ and R² have the values listed in the table X.

Table 46 X:

55 Table 46X provides 240 compounds of formula (X5-A) wherein G¹ is oxygen, R¹ is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X⁵ is chloro, R⁵ and R² have the values listed in the table X.

Table 47 X:

Table 47X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is bromo, R^5 and R^2 have the values listed in the table X.

Table 48 X:

- 5 Table 48X provides provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is iodo, R^5 and R^2 have the values listed in the table X.

Table 49 X:

- 10 Table 49X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is amino, R^5 and R^2 have the values listed in the table X.

Table 50 X:

- 15 Table 50X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is CHO, R^5 and R^2 have the values listed in the table X.

Table 51 X:

Table 51X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is CN, R^5 and R^2 have the values listed in the table X.

20 Table 52 X:

Table 52X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is OH, R^5 and R^2 have the values listed in the table X.

Table 53 X:

- 25 Table 53X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=O)OH, R^5 and R^2 have the values listed in the table X.

Table 54 X:

- 30 Table 54X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=NOH)H, R^5 and R^2 have the values listed in the table X.

Table 55 X:

- 35 Table 55X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=NOH)Cl, R^5 and R^2 have the values listed in the table X.

Table 56 X:

Table 56X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=NOH)NH₂, R^5 and R^2 have the values listed in the table X.

40 Table 57 X:

Table 57X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=O)CH₃, R^5 and R^2 have the values listed in the table X.

Table 58 X:

- 45 Table 58X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=NOH)CH₃, R^5 and R^2 have the values listed in the table X.

Table 59 X:

- 50 Table 59X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=O)CH₂Cl, R^5 and R^2 have the values listed in the table X.

Table 60 X:

- 55 Table 60X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is N, Y3 is CH, X^5 is C(=O)CH₂Br, R^5 and R^2 have the values listed in the table X.

Table 61 X:

Table 61X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is chloro, R^5 and R^2 have the values listed in the table X.

Table 62 X:

- 5 Table 62X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is bromo, R^5 and R^2 have the values listed in the table X.

Table 63 X:

- 10 Table 63X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is iodo, R^5 and R^2 have the values listed in the table X.

Table 64 X:

- 15 Table 64X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is amino, R^5 and R^2 have the values listed in the table X.

Table 65 X:

Table 65X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is CHO, R^5 and R^2 have the values listed in the table X.

20 Table 66 X:

Table 66X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is CN, R^5 and R^2 have the values listed in the table X.

Table 67 X:

- 25 Table 67X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is OH, R^5 and R^2 have the values listed in the table X.

Table 68 X:

- 30 Table 68X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=O)OH, R^5 and R^2 have the values listed in the table X.

Table 69 X:

- 35 Table 69X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=NOH)H, R^5 and R^2 have the values listed in the table X.

Table 70 X:

Table 70X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=NOH)Cl, R^5 and R^2 have the values listed in the table X.

40 Table 71 X:

Table 71X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=NOH)NH₂, R^5 and R^2 have the values listed in the table X.

Table 72 X:

- 45 Table 72X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=O)CH₃, R^5 and R^2 have the values listed in the table X.

Table 73 X:

- 50 Table 73X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=NOH)CH₃, R^5 and R^2 have the values listed in the table X.

Table 74 X:

- 55 Table 74X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, Y1 is CH, Y2 is CH, Y3 is N, X^5 is C(=O)CH₂Cl, R^5 and R^2 have the values listed in the table X.

Table 75 X:

Table 75X provides 240 compounds of formula (X5-A) wherein G^1 is oxygen, R^1 is hydrogen, $Y1$ is CH, $Y2$ is CH, $Y3$ is N, X^5 is $C(=O)CH_2Br$, R^5 and R^2 have the values listed in the table X.

The structures of $R2$ indicated in Tables P and X is indicated in the table below:

5

R^2	structure
(1,1-dioxothietan-2-yl)methyl-	
(1-oxothietan-2-yl)methyl-	
1,1-Dioxo-thietan-3-yl-	
1,1-Dioxo-thietan-3-ylmethyl-	
cis-1-oxo-thietan-3-yl-	
cis-1-oxo-thietan-3-ylmethyl-	
2-(1,1-dioxothietan-3-yl)ethanyl	
2-(1-oxothietan-3-yl)ethanyl	
2-(thietan-3-yl)ethanyl	
2,2-dimethyl-1,1-Dioxo-thietan-3-yl-	
2,2-dimethyl-1-oxo-thietan-3-yl-	

2,2-dimethylthietan-3-yl-	
3-Methyl-1,1-Dioxo-thietan-3-yl-	
3-Methyl-1-oxo-thietan-3-yl-	
3-Methyl-thietan-3-yl-	
thietan-2-ylmethyl-	
thietan-3-yl-	
thietan-3-ylmethyl-	
trans-1-oxo-thietan-3-yl-	
trans-1-oxo-thietan-3-ylmethyl-	

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing insects of the family *Curculionidae*, preferably in for use in controlling and/or preventing *Anthonomus grandis*.

- 5 Additional examples of insects from the family of *Curculionidae* are *Anthonomus corvulus*, *Anthonomus elutus*, *Anthonomus elongatus*, *Anthonomus eugenii*, *Anthonomus consors*, *Anthonomus haematopus*, *Anthonomus lecontei*, *Anthonomus molochinus*, *Anthonomus morticinus*, *Anthonomus musculus*, *Anthonomus nigrinus*, *Anthonomus phyllocola*, *Anthonomus pictus*, *Anthonomus pomorum*, *Anthonomus quadrigibbus*, *Anthonomus rectirostris*, *Anthonomus rubi*, *Anthonomus santacruzi*,
 10 *Anthonomus signatus*, *Anthonomus subfasciatus*, and *Anthonomus tenebrosus*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against *Anthonomus grandis* in cotton.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing soil pests.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing corn rootworm, in particular for use against corn root worm from the genus *Diabrotica*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
5 in controlling and/or preventing *Diabrotica virgifera*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Diabrotica barberi*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Diabrotica undecimpunctata howardi*.

10 In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing wireworms, in particular *Agriotes spp.*

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Agriotes spp.* in cereals, potato or corn.

Additional examples of *Agriotes spp.* include *Agriotes lineatus*, *Agriotes obscurus*, *Agriotes*
15 *brevis*, *Agriotes gurgistanus*, *Agriotes sputator*, *Agriotes ustulatus*, *Ctenicera destructor*, and *Limonius californicus*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing grubs, in particular white grubs.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
20 in controlling and/or preventing *Phyllophaga spp.*, particularly on corn, soybean or cotton.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Diloboderus spp.* particularly on corn, soybean or cotton.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Popillia japonica*, particularly on corn, soybean or cotton.

25 Additional examples of white grubs include *Phyllophaga anxia*, *Phyllophaga crinita*, *Phyllophaga subnitida*, *Diloboderus abderus*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing termites, e.g. on sugarcane.

Examples of termites include *Reticulitermes*, *Coptotermes*, *Macrotermes*, *Microtermes*,
30 *Globitermes*. Specific of subterranean termites include *Reticulitermes flavipes*, *Reticulitermes hesperus*, *Reticulitermes virginicus*, *Reticulitermes hageni*, *Reticulitermes speratus*, *Reticulitermes lucifugus*, *Heterotermes aureus*, *Coptotermes formosanus*, *Coptotermes acinaciformis*, *Coptotermes curvignathus*, *Nasutitermes exitiosus*, *Nasutitermes walkeri*, *Mastotermes darwiniensis*, *Schedorhinotermes spp.*, *Macrotermes bellicosus*, *Macrotermes spp.*, *Globitermes sulphureus*, *Odontotermes spp.*. Specific
35 examples of dry wood termites include *Incisitermes minor*, *Marginitermes hubbardi*, *Cryptotermes brevis*, *Kaloterms flavicollis*. Additional examples of termites include *procornitermes spp.* and *procornitermes araujoi*

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing subterranean stinkbugs, e.g. *Scaptocoris spp.*

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Scaptocoris castaneus*, in particular on cereals, soybean or corn.

5 In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing cutworms, e.g. *agrotis spp.*

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Agrotis ipsilon*, particularly on cereals, canola, soybean or corn.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
10 in controlling and/or preventing millipedes, e.g. *Julus spp.*

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Julus spp.*, particularly on cereals, canola, soybean & corn.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing broca gigante, e.g. *Telchin licus*, particularly on sugarcane.

15 In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing whitefly.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing *Bemisia tabaci*, particularly on vegetables, cotton, soybean, or potatoes.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
20 in controlling and/or preventing *Trialeurodes vaporariorum*, particularly on vegetables, cotton, soybean, or potatoes.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use in controlling and/or preventing stinkbugs, in particular *Euschistus spp.*

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
25 in controlling and/or preventing *Euschistus spp.*, particularly in soybean.

Examples of stinkbugs include *Nezara spp.* (e.g. *Nezara viridula*, *Nezara antennata*, *Nezara hilare*), *Piezodorus spp.* (e.g. *Piezodorus guildinii*), *Acrosternum spp.* *Euschistus spp.* (e.g. *Euschistus heros*, *Euschistus servus*), *Halyomorpha halys*, *Plautia crossota*, *Riptortus clavatus*, *Rhopalus msculatus*, *Antestiopsis orbitalis*, *Dichelops spp.* (e.g. *Dichelops furcatus*, *Dichelops melacanthus*), *Eurygaster spp.*
30 (e.g. *Eurygaster intergriceps*, *Eurygaster maura*), *Oebalus spp.* (e.g. *Oebalus mexicana*, *Oebalus poecilus*, *Oebalus pugnase*, *Scotinophara spp.* (e.g. *Scotinophara lurida*, *Scotinophara coarctata*). Preferred targets include *Antestiopsis orbitalis*, *Dichelops furcatus*, *Dichelops melacanthus*, *Euschistus heros*, *Euschistus servus*, *Nezara viridula*, *Nezara hilare*, *Piezodorus guildinii*, *Halyomorpha halys*. In one embodiment the stinkbug target is *Nezara viridula*, *Piezodorus spp.*, *Acrosternum spp.*, *Euschistus*
35 *heros*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against rice pests.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against stemborer, particularly in rice.

Examples of stemborers include *Chilo sp*, *Chilo suppressalis*, *Chilo polychrysus*, *Chilo auricilius*, *Scirpophaga spp.*, *Scirpophaga incertulas*, *Scirpophaga innotata*, *Scirpophaga nivella*

5 *Sesamia sp*, *Sesamia inferens*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against leaffolder, particularly in rice.

Examples of leaffolders include *Cnaphalocrocis spp.*, *Cnaphalocrocis medinalis*, *Marasmia spp.*, *Marasmia patnalis*, *Marasmia exigua*.

10 In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against hoppers, particularly in rice.

Examples of Hoppers include *Nephotettix spp.*, *Nephotettix virescens*, *Nephotettix nigropictus*, *Nephotettix malayanus*, *Nephotettix cincticeps*, *Nilaparvata lugens*, *Sogatella furcifera*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
15 against gallmidge, particularly in rice.

Examples of Gall midge include *Orseolia sp*, *Orseolia oryzae*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against whorl maggot, particularly in rice.

Examples of whorl maggots include *Hydrellia sp*, *Hydrellia philippina*.

20 In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against Rice bugs, particularly in rice.

Examples of rice bugs include *Leptocorisa sp*, *Leptocorisa oratorius*, *Leptocorisa chinensis*, *Leptocorisa acuta*.


In one embodiment the invention provides a compound selected from Tables 1P to 90P for use
25 against Black bugs, particularly in rice.

Examples of Black bugs include *Scotinophara sp*, *Scotinophara coarctata*, *Scotinophara lurida*, *Scotinophara latiuscula*.

In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against *plutella spp.*.

30 In one embodiment the invention provides a compound selected from Tables 1P to 90P for use against *Plutella xylostella*, particularly in brassica crops.

Compounds of formula I include at least one chiral centre and may exist as compounds of formula I* or compounds of formula I**. Compounds I* and I** are enantiomers if there is no other
35 chiral center or epimers otherwise.




mixtures of compounds I* and I** in any ratio e.g. in a molar ratio of 1:99 to 99:1, e.g. 10:1 to 1:10, e.g. a substantially 50:50 molar ratio. In an enantiomerically (or epimerically) enriched mixture of formula I**, the molar proportion of compound I** compared to the total amount of both enantiomers is for example greater than 50%, e.g. at least 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or at least 99%.

Likewise, in enantiomerically (or epimerically) enriched mixture of formula I*, the molar proportion of the compound of formula I* compared to the total amount of both enantiomers (or epimerically) is for example greater than 50%, e.g. at least 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or at least 99%.

Enantiomerically (or epimerically) enriched mixtures of formula I** are preferred. Where compounds of formula I contain an SO group, cis SO groups are preferred. When compounds of formula I** contain an SO group, cis SO groups are preferred. Each of the compounds disclosed in Tables 1P to 270P represents a specific disclosure of a compound with the R configuration (i.e. at the corresponding position indicated for compounds I* and I**) and a specific disclosure of a compound having the S configuration (i.e. at the corresponding position indicated for compounds I* and I**).

Compounds of formula I can be prepared according to the schemes illustrated in WO2009/08050, which is incorporated herein by reference. Further details regarding possible routes to the compounds of the invention are given below.

Scheme 1



1) Compounds of formula (I) can be prepared by reacting a compound of formula (V) wherein R is OH, C₁-C₆alkoxy or Cl, F or Br, with an amine of formula (XX), wherein L is methylene or ethylene,

as shown in **Scheme 1**. When R is OH such reactions are usually carried out in the presence of a coupling reagent, such as N,N'-dicyclohexylcarbodiimide ("DCC"), 1-ethyl-3-(3-dimethylamino-propyl)-carbodiimide hydrochloride ("EDC") or bis(2-oxo-3-oxazolidinyl)phosphonic chloride ("BOP-Cl"), in the presence of a base, and optionally in the presence of a nucleophilic catalyst, such as hydroxybenzotriazole 5 ("HOBT"). When R is Cl, such reactions are usually carried out in the presence of a base, and optionally in the presence of a nucleophilic catalyst. It is possible to conduct the reaction in a biphasic system comprising an organic solvent, preferably ethyl acetate, and an aqueous solvent, preferably a solution of sodium hydrogen carbonate. When R is C₁-C₆alkoxy it is sometimes possible to convert the ester directly to the amide by heating the ester and amine together in a thermal process. Suitable bases include pyridine, 10 triethylamine, 4-(dimethylamino)-pyridine ("DMAP") or diisopropylethylamine (Hunig's base). Preferred solvents are N,N-dimethylacetamide, tetrahydrofuran, dioxane, 1,2-dimethoxyethane, ethyl acetate and toluene. The reaction is carried out at a temperature of from 0°C to 100°C, preferably from 15°C to 30°C, in particular at ambient temperature. Amines of formula (XX) are either known in the literature or can be prepared using methods known to a person skilled in the art. Some of these methods are described in the 15 preparation examples.


2) Acid halides of formula (V), wherein R is Cl, F or Br, may be made from carboxylic acids of formula (V), wherein R is OH, under standard conditions, as described for example in WO2008/128711.

3) Carboxylic acids of formula (V), wherein R is OH, may be formed from esters of formula (V), wherein R is C₁-C₆alkoxy as described for example in WO2009/072621.

20 4) Compounds of formula (I) can be prepared by reacting a compound of formula (VI) wherein X^B is a leaving group, for example a halogen, such as bromo, with carbon monoxide and an amine of formula (X), in the presence of a catalyst, such as palladium(II) acetate or bis-(triphenylphosphine)palladium(II) dichloride, optionally in the presence of a ligand, such as triphenylphosphine, and a base, such as sodium carbonate, pyridine, triethylamine, 4-(dimethylamino)- 25 pyridine ("DMAP") or diisopropylethylamine (Hunig's base), in a solvent, such as water, N,N-dimethylformamide or tetrahydrofuran. The reaction is carried out at a temperature from 50°C to 200°C, preferably from 100°C to 150°C. The reaction is carried out at a pressure of from 50 to 200 bar, preferably from 100 to 150 bar.

5) Compounds of formula (VI) wherein X^B is a leaving group, e.g. halogen, C₁-C₈alkoxy, C₁-C₈alkylsulfonyloxy, C₁-C₈haloalkylsulfonyloxy, C₁-C₈arylsulfonyloxy, optionally substituted C₁-C₈arylsulfonyloxy (aryl is preferably phenyl), diazonium salts (e.g. X^B is -N₂⁺ Cl⁻, -N₂⁺ BF₄⁻, -N₂⁺ Br⁻, -N₂⁺ PF₆⁻), phosphonate esters (e.g. -OP(O)(OR^x)₂, wherein R^x is methyl or ethyl), preferably bromo, iodo, chloro, trifluoromethylsulfoxy, p-toluenesulfoxy, diazonium chloride, preferably halogen, more preferably bromo, can be made by a various of methods, for example as described in WO2009/080250.

35 **Scheme 2**



6) Thietan-3-nitrile (III) can be obtained by reaction of epithiochlorhydrin (II) with a cyanide M-CN, such as sodium cyanide or potassium cyanide in the presence of water, preferably as a co-solvent with an organic solvent such as benzene or tetrahydrofuran, at a temperature of 20°C to 100°C, preferably 40-60°C preferably around 50°C.

Water is preferably used as a solvent, more preferably as a co-solvent with an organic solvent, preferably a water-immiscible organic solvent, e.g. such that the reaction then takes place in a biphasic system. The organic co-solvent is preferred to be aprotic, and is more preferably chosen from pentane, hexane, heptane, cyclohexane, benzene, toluene, xylene, tetrahydrofuran, ethyl acetate, diethyl ether, or methyl-tert-butyl ether. The reaction can be performed with an excess of epithiochlorhydrin or with an excess of M-CN, preferably in stoichiometric ratio or slight excess of one or the other reagent.

7) Compounds of formula (IV) can be prepared by reaction of compounds of formula (III) with a suitable reducing reagent. The most suitable, but not exclusive, method is the use of metal hydride reagents, such as lithium aluminum hydride or borane, in the presence or not of cocatalysts. Other methods that can be carried out involve the hydrogenation in the presence of Raney Ni, or palladium for instance. The most common solvents for this reaction are alcohols such as methanol or ethanol, tetrahydrofuran, toluene, ethers, such as diethyl ether or methyl-tert-butyl ether. In most cases it is advantageous to conduct the reaction at dilution between 0.1 M to 1 M, preferably 0.3 M to 0.5 M, at a temperature of 0°C to 100°C, preferably 0-80°C preferably around 20°C, and the reaction time in most cases is between 30 minutes and 12 hours.

Scheme 3



8) Compounds of formula (VIII) can be obtained by various methods from the thietanone (VII). For example, they can be prepared by performing a Wittig reaction or a related reaction, using (triphenylphosphoranylidene)acetonitrile or (Cyanomethyl)diethoxyphosphine oxide as a reagent or following a procedure in analogy to the one described in Organic Letters 2010, 12(9), 1944-1947 and Journal of the American Chemical Society 2009, 131(8), 2786-2787. Such reactions are usually performed in a solvent, such as toluene or dichloromethane, at a temperature of from 0°C to 150°C, preferably from 0°C to 50°C. Thietan-3-one (VII) can be prepared according to known methods, for example described in Synlett, (11), 783-4; 1991 or in WO 2007046548.


9) Compounds of formula (IX) can be prepared by reaction of compounds of formula (VIII) with a suitable reducing reagent. The most suitable, but not exclusive, method is the use of metal hydride reagents, such as sodium borohydride. Other methods that can be carried out involve the hydrogenation under a hydrogen atmosphere in the presence of palladium for instance. The most common solvents for this reaction are alcohols such as methanol or ethanol. In most cases it is advantageous to conduct the reaction at dilution between 0.1 M to 1 M, preferably 0.3 M to 0.5 M, at a temperature of 0°C to 100°C, preferably 0-80°C preferably around 20°C, and the reaction time in most cases is between 30 minutes and 12 hours.

10) Compounds of formula (X) can be prepared by reaction of compounds of formula (IX) with a suitable reducing reagent. The most suitable, but not exclusive, method is the use of metal hydride reagents, such as lithium aluminum hydride or borane, in the presence or not of cocatalysts. Other methods that can be carried out involve the hydrogenation in the presence of Raney Ni, or palladium for instance. The most common solvents for this reaction are alcohols such as methanol or ethanol, tetrahydrofuran, toluene, ethers, such as diethyl ether or methyl tert-butyl ether. In most cases it is advantageous to conduct the reaction at dilution between 0.1 M to 1 M, preferably 0.3 M to 0.5 M, at a temperature of 0°C to 100°C, preferably 0-80°C preferably around 20°C, and the reaction time in most cases is between 30 minutes and 12 hours.

11) Alternatively, compounds of formula (X) may be obtained directly by reduction of compounds of formula (VIII), for example under hydrogenation conditions, e.g. using hydrogen and a metal hydrogenation catalyst, e.g. Palladium, Rhodium, or Platinum.

For the preparation of the amines, a few other routes can be followed, some of which are represented in schemes 4,5, 6 and 7

Scheme 4



12) Compound (IV) may be prepared by reduction of a nitromethylene compound of formula (XI). Reducing agents suitable for this reduction include for example metal hydride reagents such as sodium borohydride or lithium aluminium hydride optionally in the presence of a catalyst, for example nickel chloride. Hydrogenation in the presence of metal catalyst such as raney nickel or palladium on


charcoal is an alternative for such reductions. Examples of such methods can be found in Journal of the American Chemical Society (2003), 125(40), 12125-12136, US patent n° 20050261327, or in Journal of Medicinal Chemistry (2010), 53(7), 2942-2951.

13) Compounds of formula (XI) can be prepared by reaction of the ketone of formula (VII) with 5 nitromethane under standard conditions (Henry reaction), as described for example in Angewandte Chemie, International Edition, 45(46), 7736-7739; 2006 or in Journal of Medicinal Chemistry, 53(8), 3227-3246; 2010.

14) Alternatively, the amine of formula (IV) may be obtained from the carboxylic acid of formula (XIII) using Schmidt or Curtius rearrangement. Schmidt rearrangement involves treatment with hydrazoic acid under a range of possible conditions known to the person skilled of the art, for example as described in Journal of Organic Chemistry, 58(6), 1372-6; 1993 for the conversion of bicyclo[1.1.1]pentane-2-carboxylic acid to bicyclo[1.1.1]pentane-2-amine. Curtius rearrangement can be carried out under different possible conditions known to the person skilled in the art, for example treatment of (XIII) with diphenylphosphoryl azide followed by heating and reaction with an alcohol such as benzyl alcohol or tert-butanol; the corresponding ester is obtained and subsequently deprotected to the amine by hydrogenolysis (benzyl ester) or treatment with trifluoroacetic acid (tert-butyl ester). Such conditions are for example described in Journal of Organic Chemistry, 75(17), 5941-5952, 2010 or Tetrahedron: Asymmetry, 14(23), 3773-3778; 2003.


15) The acid of formula (XII) may derive from the ketone of formula (VII) after homologation of the ketone (VII) to an ester of formula (XII) followed by hydride reduction or hydrogenation of the double bond. Similar synthetic sequences from related substrates are described for example in Chemical & Pharmaceutical Bulletin, 52(6), 675-687; 2004, Synlett (2005), (10), 1559-1562, WO 2005019221, or WO 2010031735.

Scheme 5




16) Amines of formula (IV) and (X) may be converted to further amine intermediates of formula (XIV) and (XV) wherein n is 1 or 2. The amines may be directly oxidized or first protected, oxidized and then deprotected.

Scheme 6



17) The amine of formula (XVII) may be obtained from a compound of formula (XVI) wherein each X^B is independently hydroxy, halogen or a leaving group such as mesylate, tosylate or triflate. The sequence is then similar to that described for example in WO 2007080131 for the conversion of serinol to thietan-3-ylamine. Compounds of formula (XVI) are either known compounds or can be prepared by known methods to the person skilled in the art. Compounds of formula XVII may be oxidised as described in 16).

Scheme 7



10

18) Similarly, the amine of formula (IV) and (X) may be obtained from compounds of formula (XVIII) and (XIX) wherein each X^B is independently hydroxy, halogen or a leaving group such as mesylate, tosylate or triflate. Compounds of formula (XVIII) and (XIX) are either known compounds or can be prepared by known methods to the person skilled in the art.

15

Protecting a crop of useful plants from insects, acarines, nematodes or molluscs, means e.g. controlling the population of insects, acarines, nematodes or molluscs in said crop of useful plants, e.g. such that the population of said insects, acarines, nematodes or molluscs is less than would be present in the absence of said compound, preferably significantly less, e.g. at least 5, 10, 15, 20, 25, 30, 35, 40, 45 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or even at least 99% less than in the absence of said compound.

The compounds of formula (I) can be used to combat and control infestations of insect pests such as Lepidoptera, Diptera, Hemiptera, Thysanoptera, Orthoptera, Dictyoptera, Coleoptera, Siphonaptera, Hymenoptera and Isoptera and also other invertebrate pests, for example, acarine, nematode and mollusc pests. Insects, acarines, nematodes and molluscs are hereinafter collectively referred to as pests. The pests which may be combated and controlled by the use of the compound of the invention include those pests associated with agriculture (which term includes the growing of crops for food and fiber products),

horticulture and animal husbandry, companion animals, forestry and the storage of products of vegetable origin (such as fruit, grain and timber); those pests associated with the damage of man-made structures and the transmission of diseases of man and animals; and also nuisance pests (such as flies). The compounds of the invention may be used for example on turf, ornamentals, such as flowers, shrubs,

- 5 broad-leaved trees or evergreens, for example conifers, as well as for tree injection, pest management and the like. Compositions comprising the compound of formula I may be used on ornamental garden plants (e.g. flowers, shrubs, broad-leaved trees or evergreens), e.g. to control aphids, whitefly, scales, mealybug, beetles and caterpillars. Compositions comprising the compound of formula I may be used on garden plants (e.g. flowers, shrubs, broad-leaved trees or evergreens), on indoor plants (e.g. flowers and shrubs)
- 10 and on indoor pest e.g. to control aphids, whitefly, scales, mealybug, beetles and caterpillars.

Furthermore, the compounds of the invention may be effective against harmful insects, without substantially imposing any harmful side effects to cultivated plants. Application of the compounds of the invention may increase the harvest yields, and may improve the quality of the harvested material. The compounds of the invention may have favourable properties with respect to amount applied, residue

15 formulation, selectivity, toxicity, production methodology, high activity, wide spectrum of control, safety, control of resistant organisms, e.g. pests that are resistant to organic phosphorus agents and/or carbamate agents.

Examples of pest species which may be controlled by the compounds of formula (I) include: coleopterans, for example, *Callosobruchus chinensis*, *Sitophilus zeamais*, *Tribolium castaneum*,

20 *Epilachna vigintioctomaculata*, *Agriotes fuscicollis*, *Anomala rufocuprea*, *Leptinotarsa decemlineata*, *Diabrotica* spp., *Monochamus alternatus*, *Lissorhoptrus oryzophilus*, *Lyctus bruneus*, *Aulacophora femoralis*; lepidopterans, for example, *Lymantria dispar*, *Malacosoma neustria*, *Pieris rapae*, *Spodoptera litura*, *Mamestra brassicae*, *Chilo suppressalis*, *Pyrausta nubilalis*, *Ephestia cautella*, *Adoxophyes orana*, *Carpocapsa pomonella*, *Agrotis fucosa*, *Galleria mellonella*, *Plutella maculipennis*,

25 *Heliothis virescens*, *Phyllocnistis citrella*; hemipterans, for example, *Nephotettix cincticeps*, *Nilaparvata lugens*, *Pseudococcus comstocki*, *Unaspis yanonensis*, *Myzus persicas*, *Aphis pomi*, *Aphis gossypii*, *Rhopalosiphum pseudobrassicae*, *Stephanitis nashi*, *Nezara* spp., *Trialeurodes vaporariorum*, *Psylla* spp.; thysanopterans, for example, *Thrips palmi*, *Franklinella occidentalis*; orthopterans, for example, *Blattella germanica*, *Periplaneta americana*, *Gryllotalpa africana*, *Locusta migratoria migratorioides*; isopterans,

30 for example, *Reticulitermes speratus*, *Coptotermes formosanus*; dipterans, for example, *Musca domestica*, *Aedes aegypti*, *Hylemia platura*, *Culex pipiens*, *Anopheles sinensis*, *Culex tritaeniorhynchus*, *Liriomyza trifolii*; acari, for example, *Tetranychus cinnabarinus*, *Tetranychus urticae*, *Panonychus citri*, *Aculops pelekassi*, *Tarsonemus* spp.; nematodes, for example, *Meloidogyne incognita*, *Bursaphelenchus ligniculus* Mamiya et Kiyohara, *Aphelenchoides besseyi*, *Heterodera glycines*, *Pratylenchus* spp..

35 Examples of further pest species which may be controlled by the compounds of formula (I) include: from the order of the Anoplura (*Phthiraptera*), for example, *Damalinia* spp., *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Trichodectes* spp.; from the class of the Arachnida, for example, *Acarus siro*, *Aceria sheldoni*, *Aculops* spp., *Aculus* spp., *Amblyomma* spp., *Argas* spp., *Boophilus* spp.,

- Brevipalpus* spp., *Bryobia praetiosa*, *Chorioptes* spp., *Dermanyssus gallinae*, *Eotetranychus* spp.,
Epitrimerus pyri, *Eutetranychus* spp., *Eriophyes* spp., *Hemitarsonemus* spp., *Hyalomma* spp., *Ixodes* spp.,
Latrodectus mactans, *Metatetranychus* spp., *Oligonychus* spp., *Ornithodoros* spp., *Panonychus* spp.,
Phyllocoptruta oleivora, *Polyphagotarsonemus latus*, *Psoroptes* spp., *Rhipicephalus* spp., *Rhizoglyphus*
5 spp., *Sarcoptes* spp., *Scorpio maurus*, *Stenotarsonemus* spp., *Tarsonemus* spp., *Tetranychus* spp., *Vasates*
lycopersici; from the class of the *Bivalva*, for example, *Dreissena* spp.; from the order of the *Chilopoda*,
for example, *Geophilus* spp., *Scutigera* spp.; from the order of the *Coleoptera*, for example,
Acanthoscehdes obtectus, *Adoretus* spp., *Agelastica alni*, *Agriotes* spp., *Amphimallon solstitialis*,
Anobium punctatum, *Anoplophora* spp., *Anthonomus* spp., *Anthrenus* spp., *Apogonia* spp., *Atomaria* spp.,
10 *Attagenus* spp., *Bruchidius obtectus*, *Bruchus* spp., *Ceuthorhynchus* spp., *Cleonus mendicus*, *Conoderus*
spp., *Cosmopolites* spp., *Costelytra zealandica*, *Curculio* spp., *Cryptorhynchus lapathi*, *Dermestes* spp.,
Diabrotica spp., *Epilachna* spp., *Faustinus cubae*, *Gibbium psylloides*, *Heteronychus arator*,
Hylamorpha elegans, *Hylotrupes bajulus*, *Hypera postica*, *Hypothenemus* spp., *Lachnosterna*
consanguinea, *Leptinotarsa decemlineata*, *Lissorhoptrus oryzophilus*, *Lixus* spp., *Lyctus* spp., *Meligethes*
15 *aeneus*, *Melolontha melolontha*, *Migdolus* spp., *Monochamus* spp., *Naupactus xanthographus*, *Niptus*
hololeucus, *Oryctes rhinoceros*, *Oryzaephilus surinamensis*, *Otiorrhynchus sulcatus*, *Oxycetonia*
jucunda, *Phaedon cochleariae*, *Phyllophaga* spp., *Popillia japonica*, *Premnotrypes* spp., *Psylliodes*
chrysocephala, *Ptinus* spp., *Rhizobius ventralis*, *Rhizopertha dominica*, *Sitophilus* spp., *Sphenophorus*
spp., *Sternechus* spp., *Symphyletes* spp., *Tenebrio molitor*, *Tribolium* spp., *Trogoderma* spp., *Tychius*
20 spp., *Xylotrechus* spp., *Zabrus* spp.; from the order of the *Collembola*, for example, *Onychiurus armatus*;
from the order of the *Dermaptera*, for example, *Forficula auricularia*; from the order of the *Diplopoda*,
for example, *Blaniulus guttulatus*; from the order of the *Diptera*, for example, *Aedes* spp., *Anopheles* spp.,
Bibio hortulanus, *Calliphora erythrocephala*, *Ceratitis capitata*, *Chrysomyia* spp., *Cochliomyia* spp.,
Cordylobia anthropophaga, *Culex* spp., *Cuterebra* spp., *Dacus oleae*, *Dermatobia hominis*, *Drosophila*
25 spp., *Fannia* spp., *Gastrophilus* spp., *Hylemyia* spp., *Hyppobosca* spp., *Hypoderma* spp., *Liriomyza* spp.,
Lucilia spp., *Musca* spp., *Nezara* spp., *Oestrus* spp., *Oscinella frit*, *Pegomyia hyoscyami*, *Phorbia* spp.,
Stomoxys spp., *Tabanus* spp., *Tannia* spp., *Tipula paludosa*, *Wohlfahrtia* spp.; from the class of the
Gastropoda, for example, *Arion* spp., *Biomphalaria* spp., *Bulinus* spp., *Deroceras* spp., *Galba* spp.,
Lymnaea spp., *Oncomelania* spp., *Succinea* spp.; from the class of the *helminths*, for example,
30 *Ancylostoma duodenale*, *Ancylostoma ceylanicum*, *Ancylostoma braziliensis*, *Ancylostoma* spp., *Ascaris*
lubricoides, *Ascaris* spp., *Brugia malayi*, *Brugia timori*, *Bunostomum* spp., *Chabertia* spp., *Clonorchis*
spp., *Cooperia* spp., *Dicrocoelium* spp., *Dictyocaulus filaria*, *Diphyllbothrium latum*, *Dracunculus*
medinensis, *Echinococcus granulosus*, *Echinococcus multilocularis*, *Enterobius vermicularis*, *Faciola*
spp., *Haemonchus* spp., *Heterakis* spp., *Hymenolepis nana*, *Hyostrogylus* spp., *Loa Loa*, *Nematodirus*
35 spp., *Oesophagostomum* spp., *Opisthorchis* spp., *Onchocerca volvulus*, *Ostertagia* spp., *Paragonimus*
spp., *Schistosomen* spp., *Strongyloides fuelleborni*, *Strongyloides stercoralis*, *Strongyloides* spp., *Taenia*
saginata, *Taenia solium*, *Trichinella spiralis*, *Trichinella nativa*, *Trichinella britovi*, *Trichinella nelsoni*,
Trichinella pseudospiralis, *Trichostrongylus* spp., *Trichuris trichuria*, *Wuchereria bancrofti*; it may be

- furthermore possible to control protozoa, such as *Eimeria*; from the order of the *Heteroptera*, for example, *Anasa tristis*, *Antestiopsis* spp., *Blissus* spp., *Calocoris* spp., *Campylomma livida*, *Cavelerius* spp., *Cimex* spp., *Creontiades dilutus*, *Dasyneus piperis*, *Dichelops furcatus*, *Diconocoris hewetti*, *Dysdercus* spp., *Euschistus* spp., *Eurygaster* spp., *Heliopeltis* spp., *Horcias nobilellus*, *Leptocoris* spp.,
- 5 *Leptoglossus phyllopus*, *Lygus* spp., *Macropes excavatus*, *Miridae*, *Nezara* spp., *Oebalus* spp., *Pentomidae*, *Piesma quadrata*, *Piezodorus* spp., *Psallus seriatus*, *Pseudacysta perseae*, *Rhodnius* spp., *Sahlbergella singularis*, *Scotinophora* spp., *Stephanitis nashi*, *Tibraca* spp., *Triatoma* spp.; from the order of the *Homoptera*, for example, *Acyrtosiphon* spp., *Aeneolamia* spp., *Agonosцена* spp., *Aleurodes* spp., *Aleurolobus barodensis*, *Aleurothrixus* spp., *Amrasca* spp., *Anuraphis cardui*, *Aonidiella* spp.,
- 10 *Aphanostigma piri*, *Aphis* spp., *Arboridia apicalis*, *Aspidiella* spp., *Aspidiotus* spp., *Atanus* spp., *Aulacorthum solani*, *Bemisia* spp., *Brachycaudus helichrysi*, *Brachycolus* spp., *Brevicoryne brassicae*, *Calligypona marginata*, *Carneocephala fulgida*, *Ceratovacuna lanigera*, *Cercopidae*, *Ceroplastes* spp., *Chaetosiphon fragaefolii*, *Chionaspis tegalensis*, *Chlorita onukii*, *Chromaphis juglandicola*, *Chrysomphalus ficus*, *Cicadulina mbila*, *Coccoxymytilus halli*, *Coccus* spp., *Cryptomyzus ribis*, *Dalbulus*
- 15 spp., *Dialeurodes* spp., *Diaphorina* spp., *Diaspis* spp., *Doralis* spp., *Drosicha* spp., *Dysaphis* spp., *Dysmicoccus* spp., *Empoasca* spp., *Eriosoma* spp., *Erythroneura* spp., *Euscelis bilobatus*, *Geococcus coffeae*, *Homalodisca coagulata*, *Hyalopterus arundinis*, *Icerya* spp., *Idiocerus* spp., *Idioscopus* spp., *Laodelphax striatellus*, *Lecanium* spp., *Lepidosaphes* spp., *Lipaphis erysimi*, *Macrosiphum* spp., *Mahanarva fimbriolata*, *Melanaphis sacchari*, *Metcalfiella* spp., *Metopolophium dirhodum*, *Monellia*
- 20 *costalis*, *Monelliopsis pecanis*, *Myzus* spp., *Nasonovia ribisnigri*, *Nephotettix* spp., *Nilaparvata lugens*, *Oncometopia* spp., *Orthezia praelonga*, *Parabemisia myricae*, *Paratrioza* spp., *Parlatoria* spp., *Pemphigus* spp., *Peregrinus maidis*, *Phenacoccus* spp., *Phloeomyzus passerinii*, *Phorodon humuli*, *Phylloxera* spp., *Pinnaspis aspidistrae*, *Planococcus* spp., *Protopulvinaria pyriformis*, *Pseudaulacaspis pentagona*, *Pseudococcus* spp., *Psylla* spp., *Pteromalus* spp., *Pyrilla* spp., *Quadrastpidiotus* spp., *Quesada*
- 25 *gigas*, *Rastrococcus* spp., *Rhopalosiphum* spp., *Saissetia* spp., *Scaphoides titanus*, *Schizaphis graminum*, *Selenaspis articulatus*, *Sogata* spp., *Sogatella furcifera*, *Sogatodes* spp., *Stictocephala festina*, *Tenalaphara malayensis*, *Tinocallis caryaefoliae*, *Tomaspis* spp., *Toxoptera* spp., *Trialeurodes vaporariorum*, *Trioza* spp., *Typhlocyba* spp., *Unaspis* spp., *Viteus vitifolii*; from the order of the *Hymenoptera*, for example, *Diprion* spp., *Hoplocampa* spp., *Lasius* spp., *Mono-* *morium pharaonis*,
- 30 *Vespa* spp.; from the order of the *Isopoda*, for example, *Armadillidium vulgare*, *Oniscus asellus*, *Porcellio scaber*; from the order of the *Isoptera*, for example, *Reticulitermes* spp., *Odontotermes* spp.; from the order of the *Lepidoptera*, for example, *Acronicta major*, *Aedia leucomelas*, *Agrotis* spp., *Alabama argillacea*, *Anticarsia* spp., *Barathra brassicae*, *Bucculatrix thurberiella*, *Bupalus piniarius*, *Cacoecia podana*, *Capua reticulana*, *Carpocapsa pomonella*, *Cheimatobia brumata*, *Chilo* spp.,
- 35 *Choristoneura fumiferana*, *Clysia ambiguella*, *Cnaphalocerus* spp., *Earias insulana*, *Ephestia kuehniella*, *Euproctis chrysorrhoea*, *Euxoa* spp., *Feltia* spp., *Galleria mellonella*, *Helicoverpa* spp., *Heliiothis* spp., *Hofmannophila pseudospretella*, *Homona magnanima*, *Hyponomeuta padella*, *Laphygma* spp., *Lithocolletis blancardella*, *Lithophane antennata*, *Loxagrotis albicosta*, *Lymantria* spp., *Malacosoma*

- neustria*, *Mamestra brassicae*, *Mocis repanda*, *Mythimna separata*, *Oria* spp., *Oulema oryzae*, *Panolis flammea*, *Pectinophora gossypiella*, *Phyllocnistis citrella*, *Pieris* spp., *Plutella xylostella*, *Prodenia* spp., *Pseudaletia* spp., *Pseudoplusia includens*, *Pyrausta nubilalis*, *Spodoptera* spp., *Thermesia gemmatilis*, *Tinea pellionella*, *Tineola bisselliella*, *Tortrix viridana*, *Trichoplusia* spp.; from the order of the
- 5 *Orthoptera*, for example, *Acheta domesticus*, *Blatta orientalis*, *Blattella germanica*, *Gryllotalpa* spp., *Leucophaea maderae*, *Locusta* spp., *Melanoplus* spp., *Periplaneta americana*, *Schistocerca gregaria*; from the order of the *Siphonaptera*, for example, *Ceratophyllus* spp., *Xenopsylla cheopis*. From the order of the *Symphyla*, for example, *Scutigera* spp.; from the order of the *Thysanoptera*, for example, *Baliothrips bififormis*, *Enneothrips flavens*, *Frankliniella* spp., *Heliothrips* spp., *Hercinothrips femoralis*,
- 10 *Kakothrips* spp., *Rhipiphorothrips cruentatus*, *Scirtothrips* spp., *Taeniothrips cardamoni*, *Thrips* spp.; from the order of the *Thysanura*, for example, *Lepisma saccharina*. The phytoparasitic nematodes include, for example, *Anguina* spp., *Aphelenchoides* spp., *Belonoaimus* spp., *Bursaphelenchus* spp., *Ditylenchus dipsaci*, *Globodera* spp., *Helicotylenchus* spp., *Heterodera* spp., *Longidorus* spp., *Meloidogyne* spp., *Pratylenchus* spp., *Radopholus similis*, *Rotylenchus* spp., *Trichodorus* spp.,
- 15 *Tylenchorhynchus* spp., *Tylenchulus* spp., *Tylenchulus semipenetrans*, *Xiphinema* spp.
- In particular, the compounds of the invention may be used to control the following pest species:
- Myzus persicae* (aphid), *Aphis gossypii* (aphid), *Aphis fabae* (aphid), *Lygus* spp. (capsids), *Dysdercus* spp. (capsids), *Nilaparvata lugens* (planthopper), *Nephotettix inciticeps* (leafhopper), *Nezara* spp. (stinkbugs), *Euschistus* spp. (stinkbugs), *Leptocoris* spp. (stinkbugs), *Frankliniella occidentalis*
- 20 (thrip), *Thrips* spp. (thrips), *Leptinotarsa decemlineata* (Colorado potato beetle), *Anthonomus grandis* (boll weevil), *Aonidiella* spp. (scale insects), *Trialeurodes* spp. (white flies), *Bemisia tabaci* (white fly), *Ostrinia nubilalis* (European corn borer), *Spodoptera littoralis* (cotton leafworm), *Heliothis virescens* (tobacco budworm), *Helicoverpa armigera* (cotton bollworm), *Helicoverpa zea* (cotton bollworm), *Sylepta derogata* (cotton leaf roller), *Pieris brassicae* (white butterfly), *Plutella xylostella* (diamond back
- 25 moth), *Agrotis* spp. (cutworms), *Chilo suppressalis* (rice stem borer), *Locusta migratoria* (locust), *Chortiocetes terminifera* (locust), *Diabrotica* spp. (rootworms), *Panonychus ulmi* (European red mite), *Panonychus citri* (citrus red mite), *Tetranychus urticae* (two-spotted spider mite), *Tetranychus cinnabarinus* (carmine spider mite), *Phyllocoptruta oleivora* (citrus rust mite), *Polyphagotarsonemus latus* (broad mite), *Brevipalpus* spp. (flat mites), *Boophilus microplus* (cattle tick), *Dermacentor*
- 30 *variabilis* (American dog tick), *Ctenocephalides felis* (cat flea), *Liriomyza* spp. (leafminer), *Musca domestica* (housefly), *Aedes aegypti* (mosquito), *Anopheles* spp. (mosquitoes), *Culex* spp. (mosquitoes), *Lucillia* spp. (blowflies), *Blattella germanica* (cockroach), *Periplaneta americana* (cockroach), *Blatta orientalis* (cockroach), termites of the *Mastotermitidae* (for example *Mastotermes* spp.), the *Kalotermitidae* (for example *Neotermes* spp.), the *Rhinotermitidae* (for example *Coptotermes formosanus*,
- 35 *Reticulitermes flavipes*, *R. speratu*, *R. virginicus*, *R. hesperus*, and *R. santonensis*) and the *Termitidae* (for example *Globitermes sulfureus*), *Solenopsis geminata* (fire ant), *Monomorium pharaonis* (pharaoh's ant), *Damalinea* spp. and *Linognathus* spp. (biting and sucking lice), *Meloidogyne* spp. (root knot nematodes), *Globodera* spp. and *Heterodera* spp. (cyst nematodes), *Pratylenchus* spp. (lesion nematodes),

Rhodopholus spp. (banana burrowing nematodes), *Tylenchulus* spp. (citrus nematodes), *Haemonchus contortus* (barber pole worm), *Caenorhabditis elegans* (vinegar eelworm), *Trichostrongylus* spp. (gastro intestinal nematodes) and *Deroceras reticulatum* (slug).

The compound of formula I may be used for pest control on various plants, including soybean
 5 (e.g. in some cases 10-70g/ha), corn (e.g. in some cases 10-70g/ha), sugarcane (e.g. in some cases 20-200g/ha), alfalfa (e.g. in some cases 10-70g/ha), brassicas (e.g. in some cases 10-50g/ha), oilseed rape (e.g. canola) (e.g. in some cases 20-70g/ha), potatoes (including sweet potatoes) (e.g. in some cases 10-70g/ha), cotton (e.g. in some cases 10-70g/ha), rice (e.g. in some cases 10-70g/ha), coffee (e.g. in some cases 30-150g/ha), citrus (e.g. in some cases 60-200g/ha), almonds (e.g. in some cases 40-180g/ha),
 10 fruiting vegetables, cucurbits and pulses (e.g. tomatoes, pepper, chili, eggplant, cucumber, squash etc.) (e.g. in some cases 10-80g/ha), tea (e.g. in some cases 20-150g/ha), bulb vegetables (e.g. onion, leek etc.) (e.g. in some cases 30-90g/ha), grapes (e.g. in some cases 30-180g/ha), pome fruit (e.g. apples, pears etc.) (e.g. in some cases 30-180g/ha), and stone fruit (e.g. pears, plums etc.) (e.g. in some cases 30-180g/ha).

The compounds of the invention may be used for pest control on various plants, including
 15 soybean, corn, sugarcane, alfalfa, brassicas, oilseed rape (e.g. canola), potatoes (including sweet potatoes), cotton, rice, coffee, citrus, almonds, fruiting vegetables, cucurbits and pulses (e.g. tomatoes, pepper, chili, eggplant, cucumber, squash etc.), tea, bulb vegetables (e.g. onion, leek etc.), grapes, pome fruit (e.g. apples, pears etc.), stone fruit (e.g. pears, plums etc.), and cereals.

The compounds of the invention may be used on soybean to control, for example, *Elasmopalpus lignosellus*, *Diloboderus abderus*, *Diabrotica speciosa*, *Trialeurodes* spp., *Bemisia* spp., aphids, *Sternechus subsignatus*, *Formicidae*, *Agrotis ypsilon*, *Julus* spp., *Murgantia* spp., *Halyomorpha* spp., *Thyanta* spp., *Megascelis* spp., *Procornitermes* spp., *Gryllotalpidae*, *Nezara viridula*, *Piezodorus* spp., *Acrosternum* spp., *Neomegalotomus* spp., *Cerotoma trifurcata*, *Popillia japonica*, *Edessa* spp., *Liogenys fuscus*, stalk borer, *Scaptocoris castanea*, *phyllophaga* spp., *Migdolus* spp., *Pseudoplusia includens*,
 25 *Anticarsia gemmatilis*, *Epinotia* spp., *Rachiplusia* spp., *Spodoptera* spp. (e.g. *Spodoptera frugiperda*), *Bemisia tabaci*, *Tetranychus* spp., *Agriotes* spp., *Euschistus* spp. (e.g. *Euschistus heros*). The compounds of the invention are preferably used on soybean to control *Diloboderus abderus*, *Diabrotica speciosa*, *Trialeurodes* spp., *Bemisia* spp., *Nezara viridula*, *Piezodorus* spp., *Acrosternum* spp., *Cerotoma trifurcata*, *Popillia japonica*, *Euschistus heros*, *Scaptocoris castanea*, *phyllophaga* spp., *Migdolus* spp.,
 30 *Agriotes* spp., *Euschistus* spp..

The compounds of the invention may be used on corn to control, for example, *Euschistus* spp. (e.g. *Euschistus heros*), *Dichelops furcatus*, *Diloboderus abderus*, *Thyanta* spp., *Elasmopalpus lignosellus*, *Halyomorpha* spp., *Spodoptera frugiperda*, *Nezara viridula*, *Cerotoma trifurcata*, *Popillia japonica*, *Agrotis ypsilon*, *Diabrotica speciosa*, aphids, *Heteroptera*, *Procornitermes* spp., *Scaptocoris castanea*, *Formicidae*, *Julus* spp., *Dalbulus maidis*, *Diabrotica* spp. (e.g. *Diabrotica virgifera*), *Mocis latipes*, *Bemisia tabaci*, *heliethis* spp., *Tetranychus* spp., *thrips* spp., *phyllophaga* spp., *Migdolus* spp., *scaptocoris* spp., *Liogenys fuscus*, *Spodoptera* spp., *Ostrinia* spp., *Sesamia* spp., wireworms, *Agriotes* spp., *Halotydeus destructor*. The compounds of the invention are preferably used on corn to control

Euschistus spp., (c.g. *Euschistus heros*), *Dichelops furcatus*, *Diloboderus abderus*, *Nezara viridula*, *Cerotoma trifurcata*, *Popillia japonica*, *Diabrotica* spp. (c.g. *Diabrotica speciosa*, *Diabrotica virgifera*), *Tetranychus* spp., *Thrips* spp., *Phyllophaga* spp., *Migdolus* spp., *Scaptocoris* spp., *Agriotes* spp..

The compounds of the invention may be used on sugar cane to control, for example,

- 5 *Sphenophorus* spp., termites, *Migdolus* spp., *Diloboderus* spp., *Telchin licus*, *Diatrea saccharalis*, *Mahanarva* spp., Mealybugs, *Chilo* spp..

The compounds of the invention may be used on alfalfa to control, for example, *Hypera brunneipennis*, *Hypera postica*, *Colias eurytheme*, *Collops* spp., *Empoasca solana*, *Epitrix* spp., *Geocoris* spp., *Lygus hesperus*, *Lygus lineolaris*, *Spissistilus* spp., *Spodoptera* spp., Aphids, *Trichoplusia ni*. The

- 10 compounds of the invention are preferably used on alfalfa to control *Hypera brunneipennis*, *Hypera postica*, *Empoasca solana*, *Epitrix* spp., *Lygus hesperus*, *Lygus lineolaris*, *Trichoplusia ni*.

The compounds of the invention may be used on brassicas to control, for example, *Chrysodeixis* spp., *Plutella xylostella*, *Pieris* spp. (c.g. *Pieris brassicae*, *Pieris rapae*, *Pieris napi*), *Mamestra* spp. (c.g. *Mamestra brassicae*), *Plusia* spp., *Trichoplusia* spp. (c.g. *Trichoplusia ni*), *Phyllotreta* spp. (c.g.

- 15 *Phyllotreta cruciferae*, *Phyllotreta striolata*), *Spodoptera* spp., *Empoasca* spp., *thrips* spp., *Delia* spp., *Murgantia* spp., *Trialeurodes* spp., *Bemisia* spp., *Microtheca* spp., Aphids, *Chaetocnema* spp., *Psylliodes* spp. (c.g. *Psylliodes chrysocephala*). The compounds of the invention are preferably used on brassicas to control *Plutella xylostella*, *Pieris* spp., *Plusia* spp., *Trichoplusia ni*, *Phyllotreta* spp., *Thrips* spp., *Chaetocnema* spp..

- 20 The compounds of the invention may be used on oil seed rape, e.g. canola, to control, for example, *Meligethes* spp. (c.g. *Meligethes aeneus*), *Ceutorhynchus* spp., (c.g. *Ceutorhynchus assimilis*, *Ceutorhynchus napi*), *Halotydeus destructor*, *Psylliodes* spp. (c.g. *Psylliodes chrysocephala*), *Phyllotreta* spp. (c.g. *Phyllotreta cruciferae*, *Phyllotreta striolata*), *Chaetocnema* spp..

- The compounds of the invention may be used on potatoes, including sweet potatoes, to control, 25 for example, *Empoasca* spp., *Leptinotarsa* spp., *Diabrotica speciosa*, *Phthorimaea* spp., *Paratrioza* spp., *Maladera matrida*, *Agriotes* spp., Aphids, wireworms. The compounds of the invention are preferably used on potatoes, including sweet potatoes, to control *Empoasca* spp., *Leptinotarsa* spp., *Diabrotica speciosa*, *Phthorimaea* spp., *Paratrioza* spp., *Agriotes* spp..

- The compounds of the invention may be used on cotton to control, for example, *Anthonomus* 30 *grandis*, *Pectinophora* spp., *heliopsis* spp., *Spodoptera* spp., *Tetranychus* spp. (c.g. *Tetranychus urticae*), *Empoasca* spp., *Thrips* spp. (c.g. *Thrips tabaci*, *Thrips palmi*), *Bemisia tabaci*, *Trialeurodes* spp., Aphids, *Lygus* spp. (c.g. *Lygus lineolaris*, *Lygus Hesperus*), *phyllophaga* spp., *Scaptocoris* spp., *Austroasca viridigrisea*, *Creontiades* spp., *Nezara* spp., *Piezodorus* spp., *Halotydeus destructor*, *Oxycaraenus hyalinipennis*, *Dysdercus cingulatus*, *Amrasca* spp. (c.g. *Amrasca biguttula biguttula*), *Frankliniella* spp. 35 (c.g. *Frankliniella schultzei*), *Scirtothrips* spp. (c.g. *Scirtothrips dorsalis*), *Anaphothrips* spp., *Polyphagotarsonemus latus*. The compounds of the invention are preferably used on cotton to control *Anthonomus grandis*, *Tetranychus* spp., *Empoasca* spp., *thrips* spp., *Lygus* spp., *phyllophaga* spp., *Scaptocoris* spp..

The compounds of the invention may be used on rice to control, for example, *Leptocorisa* spp. (e.g. *Leptocorisa oratorius*, *Leptocorisa chinensis*, *Leptocorisa acuta*), *Cnaphalocrosis* spp., *Chilo* spp. (e.g. *Chilo suppressalis*, *Chilo polychrysus*, *Chilo auricilius*), *Scirpophaga* spp. (e.g. *Scirpophaga incertulas*, *Scirpophaga innotata*, *Scirpophaga nivella*), *Lissorhoptrus* spp., *Oebalus pugnax*,

- 5 *Scotinophara* spp. (e.g. *Scotinophara coarctata*, *Scotinophara lurida*, *Scotinophara latiuscula*), *Nephotettix* spp. (e.g. *Nephotettix malayanus*, *Nephotettix nigropictus*, *Nephotettix parvus*, *Nephotettix virescens*, *Nephotettix cincticeps*), Mealybugs, *Sogatella furcifera*, *Nilaparvata lugens*, *Orseolia* spp. (e.g. *Orseolia oryzae*), *Cnaphalocrocis medinalis*, *Marasmia* spp. (e.g. *Marasmia patnalis*, *Marasmia exigua*), *Stenchaetothrips biformis*, *Thrips* spp., *Hydrellia* spp. (e.g. *Hydrellia philippina*), Grasshoppers, *Pomacea*
- 10 *canaliculata*, *Scirpophaga innotata*, *Sesamia inferens*, *Laodelphax striatellus*, *Nymphula depunctalis*, *Oulema oryzae*, Stinkbugs. The compounds of the invention are preferably used on rice to control *Leptocorisa* spp., *Lissorhoptrus* spp., *Oebalus pugnax*, *Nephotettix* spp. (e.g. *Nephotettix malayanus*, *Nephotettix nigropictus*, *Nephotettix parvus*, *Nephotettix virescens*, *Nephotettix cincticeps*), *Sogatella furcifera*, *Stenchaetothrips biformis*, *Thrips* spp., *Hydrellia* spp. (e.g. *Hydrellia philippina*),
- 15 Grasshoppers, *Pomacea canaliculata*, *Scirpophaga innotata*, *Chilo* spp., *Oulema oryzae*.

The compounds of the invention may be used on coffee to control, for example, *Hypothenemus* spp. (e.g. *Hypothenemus Hampei*), *Perileucoptera Coffeella*, *Tetranychus* spp., *Brevipalpus* spp., Mealybugs. The compounds of the invention are preferably used on coffee to control *Hypothenemus Hampei*, *Perileucoptera Coffeella*.

- 20 The compounds of the invention may be used on citrus to control, for example, *Panonychus citri*, *Phyllocoptruta oleivora*, *Brevipalpus* spp. (e.g. *Brevipalpus californicus*, *Brevipalpus phoenicis*), *Diaphorina citri*, *Scirtothrips* spp. (e.g. *Scirtothrips dorsalis*), *Thrips* spp., *Unaspis* spp., *Ceratitidis capitata*, *Phyllocnistis* spp. (e.g. *Phyllocnistis citrella*), Aphids, Hardscales, Softscales, Mealybugs. The compounds of the invention are preferably used on citrus to control *Panonychus citri*,
- 25 *Phyllocoptruta oleivora*, *Brevipalpus* spp., *Diaphorina citri*, *Scirtothrips* spp., *thrips* spp., *Phyllocnistis* spp..

The compounds of the invention may be used on almonds to control, for example, *Amyelois transitella*, *Tetranychus* spp.

- The compounds of the invention may be used on fruiting vegetables, cucurbits and pulses,
- 30 including tomatoes, pepper, chili, eggplant, cucumber, squash etc., to control, for example, *Thrips* spp., *Tetranychus* spp. (e.g. *Tetranychus urticae*), *Polyphagotarsonemus* spp. (e.g. *Polyphagotarsonemus latus*), *Aculops* spp. (e.g. *Aculops lycopersici*), *Empoasca* spp. (e.g. *Empoasca fabae*), *Spodoptera* spp., *heliolithis* spp., *Tuta absoluta*, *Liriomyza* spp. (e.g. *Liriomyza brassicae*, *Liriomyza bryoniae*, *Liriomyza huidobrensis*, *Liriomyza sativae*, *Liriomyza trifolii*), *Bemisia tabaci*, *Trialeurodes* spp., Aphids,
- 35 *Paratrioza* spp., *Frankliniella* spp. (e.g. *Frankliniella occidentalis*, *Frankliniella intonsa*, *Frankliniella hispinosa*), *Spodoptera* spp. (e.g. *Spodoptera exigua*, *Spodoptera littoralis*, *Spodoptera litura*, *Spodoptera frugiperda*, *Spodoptera eridania*), *Anthonomus* spp., *Phyllotreta* spp., *Amrasca* spp. (e.g. *Amrasca biguttula biguttula*), *Epilachna* spp., *Halyomorpha* spp., *Scirtothrips* spp., *Leucinodes* spp. (e.g.

Leucinodes orbonalis), *Neoleucinodes* spp. (e.g. *Neoleucinodes elegantalis*), *Maruca* spp., Fruit flies, Stinkbugs, *Lepidopteras*, *Coleopteras*, *Helicoverpa* spp. (e.g. *Helicoverpa armigera*), *Heliothis* spp. (e.g. *Heliothis virescens*), *Paratrioza* spp. (e.g. *Paratrioza cockerelli*), The compounds of the invention are preferably used on fruiting vegetables, cucurbits and pulses, including tomatoes, pepper, chili, eggplant, 5 cucumber, squash etc., to control *Thrips* spp., *Tetranychus* spp., *Polyphagotarsonemus* spp., *Aculops* spp., *Empoasca* spp., *Spodoptera* spp., *heliothis* spp., *Tuta absoluta*, *Liriomyza* spp., *Paratrioza* spp., *Frankliniella occidentalis*, *Frankliniella* spp., *Amrasca* spp., *Scirtothrips* spp., *Leucinodes* spp., *Neoleucinodes* spp..

The compounds of the invention may be used on tea to control, for example, *Pseudaulecaspis* 10 spp., *Empoasca* spp., *Scirtothrips* spp., *Caloptilia theivora*, *Tetranychus* spp.. The compounds of the invention are preferably used on tea to control *Empoasca* spp., *Scirtothrips* spp..

The compounds of the invention may be used on bulb vegetables, including onion, leek etc. to control, for example, *Thrips* spp., *Spodoptera* spp., *heliothis* spp.. The compounds of the invention are preferably used on bulb vegetables, including onion, leek etc. to control *Thrips* spp..

15 The compounds of the invention may be used on grapes to control, for example, *Empoasca* spp., *Lobesia* spp., *Eupoecilia ambiguella*, *Frankliniella* spp., *Thrips* spp., *Tetranychus* spp., *Rhipiphorothrips Cruentatus*, *Eotetranychus Willamettei*, *Erythroneura Elegantula*, *Scaphoides* spp., *Scelodonta strigicollis*, Mealybugs. The compounds of the invention are preferably used on grapes to control *Frankliniella* spp., *Thrips* spp., *Tetranychus* spp., *Rhipiphorothrips Cruentatus*, *Scaphoides* spp..

20 The compounds of the invention may be used on pome fruit, including apples, pears etc., to control, for example, *Cacopsylla* spp., *Psylla* spp., *Panonychus ulmi*, *Cydia pomonella*, *Lepidopteras*, Aphids, Hardscales, Softscales. The compounds of the invention are preferably used on pome fruit, including apples, pears etc., to control *Cacopsylla* spp., *Psylla* spp., *Panonychus ulmi*.

The compounds of the invention may be used on stone fruit to control, for example, *Grapholita* 25 *molesta*, *Scirtothrips* spp., *Thrips* spp., *Frankliniella* spp., *Tetranychus* spp., Aphids, Hardscales, Softscales, Mealybugs. The compounds of the invention are preferably used on stone fruit to control *Scirtothrips* spp., *Thrips* spp., *Frankliniella* spp., *Tetranychus* spp..

The compounds of the invention may be used on cereals to control, for example, Aphids, Stinkbugs, earthmites, *Eurygaster integriceps*, *Zabrus tenebrioides*, *Anisoplia austriaca*, *Chaetocnema* 30 *aridula*, *Phyllotreta* spp., *Oulema melanopus*, *Oscinella* spp., *Delia* spp., *Mayetiola* spp., *Contarinia* spp., *Cephus* spp., *Steneotarsonemus* spp., *Apamea* spp..

In another embodiment compounds of formula I may be used on rice to control *Baliothrips* *biformis* (Thrips), *Chilo* spp. (e.g. *Chilo polychrysus* (Dark headed striped borer), *Chilo suppressalis* (Rice stemborer), *Chilo indicus* (Paddy stem borer), *Chilo polychrysus* (Dark-headed rice borer), *Chilo* 35 *suppressalis* (Stripe stem borer)), *Cnaphalocrocis medinalis* (Rice leaf folder), *Dicladispa armigera* (Hispa), *Hydrellia philipina* (Rice whorl-maggot), *Laodelphax* spp. (Smaller brown planthopper) (e.g. *Laodelphax striatellus*), *Lema oryzae* (Rice leafbeetle), *Leptocorsia acuta* (Rice bug), *Leptocorsia oratorius* (rice bug), *Lissorhoptrus oryzophilus* (rice water weevil), *Mythemina separata* (armyworm),

- Nephotettix* spp. (Green leafhopper) (e.g. *Nephotettix cincticeps*, *Nephotettix malayanus*, *Nephotettix nigropictus*, *Nephotettix parvus*, *Nephotettix virescens*), *Nilaparvata lugens* (Brown Planthopper), *Nymphula depunctalis* (Rice caseworm), *Orseolia oryzae* (Rice Gall midge), *Oulema oryzae* (Rice leafbeetle), *Scirpophaga incertulas* (Yellow Stemborer), *Scirpophaga innotata* (White Stemborer),
- 5 *Scotinophara coarctata* (Rice black bug), *Sogaella frucifera* (White-backed planthopper), *Steneotarsonemus spinki*.

The compounds of the invention may be used to control animal housing pests including: Ants, Bedbugs (adult), Bees, Beetles, Boxelder Bugs, Carpenter Bees, Carpet Beetles, Centipedes, Cigarette, Beetles, Clover Mites, Cockroaches, Confused Flour Beetle, Crickets, Earwigs, Firebrats, Fleas, Flies,

10 Lesser Grain Borers, Millipedes, Mosquitoes, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Silverfish, Sowbugs, Spiders, Termites, Ticks, Wasps, Cockroaches, Crickets, Flies, Litter Beetles (such as Darkling, Hide, and Carrion), Mosquitoes, Pillbugs, Scorpions, Spiders, Spider Mites (Twospotted, Spruce), Ticks.

The compounds of the invention may be used to control ornamental pests including: Ants

15 (Including Imported fire ants), Armyworms, Azalea caterpillars, Aphids, Bagworms, Black vine weevils (adult), Boxelder bugs, Budworms, California oakworms, Cankerworms, Cockroaches, Crickets, Cutworms, Eastern tent caterpillars, Elm leaf beetles, European sawflies, Fall webworms, Flea beetles, Forest tent caterpillars, Gypsy moth larvae, Japanese beetles (adults), June beetles (adults), Lace bugs, Leaf-feeding caterpillars, Leafhoppers, Leafminers (adults), Leaf rollers, Leaf skeletonizers, Midges,

20 Mosquitoes, Oleander moth larvae, Pillbugs, Pine sawflies, Pine shoot beetles, Pinetip moths, Plant bugs, Root weevils, Sawflies, Scale insects (crawlers), Spiders, Spittlebugs, Striped beetles, Striped oakworms, Thrips, Tip moths, Tussock moth larvae, Wasps, Broadmites, Brown softscales, California redscales (crawlers), Clover mites, Mealybugs, Pineneedlescales (crawlers), Spider mites, Whiteflies

The compounds of the invention may be used to control turf pests including: Ants (Including

25 Imported fire ants, Armyworms, Centipedes, Crickets, Cutworms, Earwigs, Fleas (adult), Grasshoppers, Japanese beetles (adult), Millipedes, Mites, Mosquitoes (adult), Pillbugs, Sod webworms, Sow bugs, Ticks (including species which transmit Lyme disease), Bluegrass billbugs (adult), Black turfgrass ataenius (adult), Chiggers, Fleas (adult), Grubs (suppression), Hyperodes weevils (adult), Mole crickets (nymphs and young adults), Mole Crickets (mature adults), Chinch Bugs.

30 The compounds of formula (I), in particular those in the tables above, may be used for soil applications, including as a seed application, to target at least the following: sucking pests such as aphids, thrips, brown plant hopper (e.g. on rice), sting bugs, white flies (e.g. on cotton and vegetables), mites; on soil pests such as corn root worm, wireworms, white grubs, zabrus, termites (e.g. on sugar cane, soy, pasture), maggots, cabbage root fly, red legged earth mite; on lepidoptera, such as spodoptera, cutworms,

35 *elasmoplus*, *plutella* (e.g. brassica), stem borers, leaf miners, flea beetle, *Sternechus*; on nematocides, such as *Heterodera glycines* (e.g. on soybean), *Pratylenchus brachyurus* (e.g. on corn), *P. zaeae* (e.g. on corn), *P. penetrans* (e.g. on corn), *Meloidogyne incognita* (e.g. on vegetables), *Heterodera schachtii*

(e.g. on sugar beet), *Rotylenchus reniformis* (e.g. on cotton), *Heterodera avenae* (e.g. on cereals), *Pratylenchus neglectus* (e.g. on cereals), *thornei* (e.g. on cereals).

The compounds of formula (I), in particular those in the tables above may be used for seed applications at least on the following: soil grubs for corn, soybeans, sugarcane: *Migdolus spp*;

5 *Phyllophaga spp.*; *Diloboderus spp*; *Cyclocephala spp*; *Lyogenys fuscus*; sugarcane weevils:

Sphenophorus levis & *Metamasius hemipterus*; termites for soybeans, sugarcane, pasture, others:

Heterotermes tenuis; *Heterotermes longiceps*; *Cornitermes cumulans*; *Procornitermes triacifer* ;

, *Neocapritermes opacus*; *Neocapritermes parvus*; corn root worms for corn and potatoes: *Diabrotica spp.*,

seed Maggot: *Delia platura*; soil stinkbugs: *Scaptocoris castanea*; wireworms: *Agriotes spp*; *Athous spp*

10 *Hipnoides bicolor*; *Ctenicera destructor*; *Limonijs canu*; *Limonijs californicus*; rice water weevil:

Lissorhoptrus oryzophilus; Red Legged earth mites: *Halotydeus destructor*.

The invention therefore provides a method of combating and/or controlling an animal pest, e.g. an invertebrate animal pest, which comprises applying to the pest, to a locus of the pest, or to a plant susceptible to attack by the pest a pesticidally effective amount of a compound of formula (I). In

15 particular, the invention provides a method of combating and/or controlling insects, acarines, nematodes or molluscs which comprises applying an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I), or a composition containing a compound of formula (I), to a pest, a locus of pest, preferably a plant, or to a plant susceptible to attack by a pest, The compounds of formula (I) are preferably used against insects, acarines or nematodes.

20 The term "plant" as used herein includes seedlings, bushes and trees. Crops are to be understood as also including those crops which have been rendered tolerant to herbicides or classes of herbicides (e.g. ALS-, GS-, EPSPS-, PPO- and HPPD-inhibitors) by conventional methods of breeding or by genetic engineering. An example of a crop that has been rendered tolerant to imidazolinones, e.g. imazamox, by conventional methods of breeding is Clearfield® summer rape (canola). Examples of crops that have been
25 rendered tolerant to herbicides by genetic engineering methods include e.g. glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady® and LibertyLink®.

The compounds of the invention may be applied to plant parts. Plant parts are to be understood as meaning all parts and organs of plants above and below the ground, such as shoot, leaf, flower and root,
30 examples which may be mentioned being leaves, needles, stalks, stems, flowers, fruit bodies, fruits, seeds, roots, tubers and rhizomes. The plant parts also include harvested material, and vegetative and generative propagation material, for example-cuttings, tubers, rhizomes, offshoots and seeds. Treatment according to the invention of the plants and plant parts with the active compounds is carried out directly or by allowing the compounds to act on their surroundings, habitat or storage space by the customary treatment methods,
35 for example by immersion, spraying, evaporation, fogging, scattering, painting on, injecting and, in the case of propagation material, in particular in the case of seed, also by applying one or more coats.

The term "plant" as used herein includes seedlings, bushes and trees. Crops are to be understood as also including those crops which have been rendered tolerant to herbicides or classes of herbicides (e.g.

ALS-, GS-, EPSPS-, PPO- and HPPD-inhibitors) by conventional methods of breeding or by genetic engineering. An example of a crop that has been rendered tolerant to imidazolinones, e.g. imazamox, by conventional methods of breeding is Clearfield® summer rape (canola). Examples of crops that have been rendered tolerant to herbicides by genetic engineering methods include e.g. glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady® and LibertyLink®.

Compounds of formula I may be used on transgenic plants (including cultivars) obtained by genetic engineering methods and/or by conventional methods. These are understood as meaning plants having novel properties ("traits") which have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive "synergistic" effects.

Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, higher quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

The preferred transgenic plants or plant cultivars which are to be treated according to the invention include all plants which, by virtue of the genetic modification, received genetic material which imparts particularly advantageous, useful traits to these plants. Examples of such traits are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, higher quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products.

Further and particularly emphasized examples of such traits are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds.

Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soybean, potatoes, sugar beet, tomatoes, peas and other vegetable varieties, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes).

Compounds of formula I may be used on transgenic plants that are capable of producing one or more pesticidal proteins which confer upon the transgenic plant tolerance or resistance to harmful pests, e.g. insect pests, nematode pests and the like. Such pesticidal proteins include, without limitation, Cry proteins from *Bacillus thuringiensis* Cry1Ab, Cry1Ac, Cry1F, Cry2Ab, Cry2Ae, Cry3A, Cry3Bb, or Cry9C; engineered proteins such as modified Cry3A (US Patent 7,030,295) or Cry1A.105; or vegetative insecticidal proteins such as Vip1, Vip2 or Vip3. A full list of Bt Cry proteins and VIPs useful in the

invention can be found on the worldwide web at *Bacillus thuringiensis* Toxin Nomenclature Database maintained by the University of Sussex (*see also*, Crickmore *et al.* (1998) *Microbiol. Mol. Biol. Rev.* 62:807-813). Other pesticidal proteins useful in the invention include proteins of bacteria colonizing nematodes, e.g. *Photorhabdus spp.* or *Xenorhabdus spp.*; toxins produced by animals, such as scorpion toxins, arachnid toxins, wasp toxins, or other insect-specific neurotoxins; toxins produced by fungi, such as *Streptomyces* toxins, plant lectins, such as pea or barley lectins; agglutinins; proteinase inhibitors, such as trypsin inhibitors, serine protease inhibitors, patatin, cystatin or papain inhibitors; ribosome-inactivating proteins (RIP), such as ricin, maize-RIP, abrin, luffin, saporin or bryodin; steroid metabolism enzymes, such as 3-hydroxysteroid oxidase, ecdysteroid-IDP-glycosyl-transferase, cholesterol oxidases, ecdysone inhibitors or HMG-CoA-reductase; ion channel blockers, such as blockers of sodium or calcium channels; juvenile hormone esterase; diuretic hormone receptors (helicokinin receptors); stilben synthase, bibenzyl synthase, chitinases or glucanases. Further examples of such pesticidal proteins or transgenic plants capable of synthesizing such proteins are disclosed, e.g., in EP-A 374753, WO 93/007278, WO 95/34656, EP-A 427529, EP-A 451878, WO 03/18810 and WO 03/52073. The methods for producing such transgenic plants are generally known to the person skilled in the art and some of which are commercially available such as Agrisure®CB (corn producing Cry1Ab), Agrisure®RW (corn producing mCry3A), Agrisure® Viptera (corn hybrids producing Vip3Aa); Agrisure300GT (corn hybrids producing Cry1Ab and mCry3A); YieldGard® (corn hybrids producing the Cry1Ab protein), YieldGard® Plus (corn hybrids producing Cry1Ab and Cry3Bb1), Genuity® SmartStax® (corn hybrids with Cry1A.105, Cry2Ab2, Cry1F, Cry34/35, Cry3Bb) ; Herculex® I (corn hybrids producing Cry1Fa) and Herculex®RW (corn hybrids producing Cry34Ab1, Cry35Ab1 and the enzyme Phosphinothricin-N-Acetyltransferase [PAT]) ; NuCOTN®33B (cotton cultivars producing Cry1Ac), Bollgard®I (cotton cultivars producing Cry1Ac), Bollgard®II (cotton cultivars producing Cry1Ac and Cry2Ab2) and VIPCOT® (cotton cultivars producing a Vip3Aa). Soybean Cyst Nematode resistance soybean (SCN® - Syngenta) and soybean with Aphid resistant trait (AMT®) are also of interest.

Further examples of such transgenic crops are:

1. **Bt11 Maize** from Syngenta Seeds SAS, Chemin de l'Hobit 27, F-31 790 St. Sauveur, France, registration number C/FR/96/05/10. Genetically modified *Zea mays* which has been rendered resistant to attack by the European corn borer (*Ostrinia nubilalis* and *Sesamia nonagrioides*) by transgenic expression of a truncated CryIA(b) toxin. Bt11 maize also transgenically expresses the enzyme PAT to achieve tolerance to the herbicide glufosinate ammonium.
2. **Bt176 Maize** from Syngenta Seeds SAS, Chemin de l'Hobit 27, F-31 790 St. Sauveur, France, registration number C/FR/96/05/10. Genetically modified *Zea mays* which has been rendered resistant to attack by the European corn borer (*Ostrinia nubilalis* and *Sesamia nonagrioides*) by transgenic expression of a CryIA(b) toxin. Bt176 maize also transgenically expresses the enzyme PAT to achieve tolerance to the herbicide glufosinate ammonium.
3. **MIR604 Maize** from Syngenta Seeds SAS, Chemin de l'Hobit 27, F-31 790 St. Sauveur, France, registration number C/FR/96/05/10. Maize which has been rendered insect-resistant by transgenic

expression of a modified CryIIIA toxin. This toxin is Cry3A055 modified by insertion of a cathepsin-D-protease recognition sequence. The preparation of such transgenic maize plants is described in WO 03/018810.

4. **MON 863 Maize** from Monsanto Europe S.A. 270-272 Avenue de Tervuren, B-1150 Brussels, Belgium, registration number C/DE/02/9. MON 863 expresses a CryIIIB(b1) toxin and has resistance to certain Coleoptera insects.

5. **IPC 531 Cotton** from Monsanto Europe S.A. 270-272 Avenue de Tervuren, B-1150 Brussels, Belgium, registration number C/ES/96/02.

6. **1507 Maize** from Pioneer Overseas Corporation, Avenue Tedesco, 7 B-1160 Brussels, Belgium, registration number C/NL/00/10. Genetically modified maize for the expression of the protein Cry1F for achieving resistance to certain Lepidoptera insects and of the PAT protein for achieving tolerance to the herbicide glufosinate ammonium.

7. **NK603 × MON 810 Maize** from Monsanto Europe S.A. 270-272 Avenue de Tervuren, B-1150 Brussels, Belgium, registration number C/GB/02/M3/03. Consists of conventionally bred hybrid maize varieties by crossing the genetically modified varieties NK603 and MON 810. NK603 × MON 810 Maize transgenically expresses the protein CP4 EPSPS, obtained from *Agrobacterium sp.* strain CP4, which imparts tolerance to the herbicide Roundup® (contains glyphosate), and also a CryIA(b) toxin obtained from *Bacillus thuringiensis subsp. kurstaki* which brings about tolerance to certain Lepidoptera, include the European corn borer.

Further examples of transgenic plants, and of very high interest, are those carrying traits conferring resistance to 2,4D (e.g. Enlist®) (e.g. WO 2011/066384), glyphosate (e.g. Roundup Ready®, Roundup Ready 2 Yield®), sulfonylurea (e.g. STS®), glufosinate (e.g. Liberty Link®, Ignite®), Dicamba (Monsanto), HPPD tolerance (e.g. isoxaflutole herbicide) (Bayer CropScience, Syngenta). Double or triple stacks of any of the traits described here are also of interest, including glyphosate and sulfonyl-urea tolerance (e.g. Optimum GAT®), plants stacked with STS® and Roundup Ready® or plants stacked with STS® and Roundup Ready 2 Yield®, dicamba and glyphosate tolerance (Monsanto). Of particular interest are soybean plants carrying traits conferring resistance to 2,4D (e.g. Enlist®), glyphosate (e.g. Roundup Ready®, Roundup Ready 2 Yield®), sulfonylurea (e.g. STS®), glufosinate (e.g. Liberty Link®, Ignite®), Dicamba (Monsanto) HPPD tolerance (e.g. isoxaflutole herbicide) (Bayer CropScience, Syngenta). Double or triple stack in soybean plants of any of the traits described here are also of interest, including glyphosate and sulfonyl-urea tolerance (e.g. Optimum GAT®, plants stacked with STS® and Roundup Ready® or Roundup Ready 2 Yield®), dicamba and glyphosate tolerance (Monsanto).

Transgenic crops of insect-resistant plants are also described in BATS (Zentrum für Biosicherheit und Nachhaltigkeit, Zentrum BATS, Clarastrasse 13, 4058 Basel, Switzerland) Report 2003, (<http://bats.ch>).

Examples of cotton transgenic events include MON 531 / 757 / 1076 (Bollgard I® - Monsanto), MON1445 (Roundup ready cotton® - Monsanto), MON531 × MON1445 (Bollgard I + RR® - Monsanto), MON15985 (Genuity Bollgard II cotton® - Monsanto), MON88913 (Genuity RR FLEX

cotton® – Monsanto), MON15985 x MON1445 (Genuity Bollgard II + RR FELX cotton® – Monsanto), MON15983 x MON88913 (Genuity Bollgard II + RR FLEX cotton® – Monsanto), MON15985 (FibreMax Bollgard II Cotton® – Monsanto), LL25 (FibreMax LL cotton® – BCS Stoneville), GHB614 (FibreMax GlyTol cotton® – BCS Stoneville), LL25 x MON15985 (FibreMax LL Bollgard II cotton® – BCS Stoneville / Monsanto), GHB614 x LL25 (FibreMax LL GlyTol cotton® – BCS Stoneville), GHB614 x LL25 x MON15985 (FibreMax RR GlyTol Bollgard II cotton® – BCS Stoneville), MON88913 x MON15985 (FibreMax LL GlyTol Bollgard II cotton® – Monsanto), MON88913 (FibreMax RR Flex cotton® – Monsanto), GHB119 + T304-40 (Twinlink® – BCS Stoneville), GHB119 + T304-40 x LL25 x GHB614 (Twinlink LL GT® – BCS Stoneville), 3006-210-23 x 281-24-236

10 (PhytoGen Widestrike Insect Protection® – Dow), 3006-210-23 x 281-24-236 x MON88913 (PhytoGen Widestrike Insect Protection + RR FLEX –® Dow / Monsanto), 3006-210-23 x 281-24-236 x MON1445 ((PhytoGen Widestrike Insect Protection + RR® – Dow / Monsanto), MON1445 (PhytoGen Roundup Ready® – Monsanto), MON88913 (PhytoGen Roundup Ready FLEX® – Monsanto), COT102 x COT67B (Vipcot® – Syngenta), COT102 x COT67B x MON88913 (Vipcot RR FLEX® – Syngenta / Monsanto), 281-24-236 (Dow), 3006-210-23 (Dow), COT102 (Syngenta), COT67B (Syngenta), T304-40 (BCS Stoneville).

Examples of Soy transgenic events include MON87701 x MON89788 (Genuity Roundup ready 2 Yield soybeans® – Monsanto), MON89788 (Roundup Ready2Yield®, RR2Y® – Monsanto), MON87708 (Monsanto), 40-3-2 (Roundup Ready®, RR1® – Monsanto), MON87701 (Monsanto), DAS-68416 (Enlist Weed Control System® – Dow), DP356043 (Optimum GAT® – Pioneer), A5547-127 (LibertyLink soybean® – Bayercropscience), A2704-12 (Bayercropscience), GU262 (Bayercropscience), W62 W98 (Bayercropscience), CRV127 (Cultivance® – BASF / EMBRAPA) SYHT0H2 (WO2012/082548).

Examples of Maize transgenic events include T25 (LibertyLink®, LL® – Bayercropscience), DHT-1 (Dow), TC1507 (Herculex I® – Dow), DAS59122-7 (Herculex RW® – Dow), TC1507 + DAS59122-7 – Herculex Xtra® – Dow), TC1507 x DAS-59122-7 x NK603 (Herculex Xtra + RR® – Dow), TC1507 x DAS-59122- x MON88017 x MON89034 (Genuity Smartstax corn®, Genuity Smartstax RIB complete® – Monsanto / Dow), MON89034 x NK603 (Genuity VT double PRO® – Monsanto), MON89034 + MON88017 (Genuity VT Triple PRO® – Monsanto), NK603 (Roundup Ready 2®, RR2® – Monsanto), MON810 (YieldGard BT®, Yieldgard cornborer® – Monsanto), MON810 x NK603 (YieldGard cornborer RR Corn 2® – Monsanto), MON810 x MON863 (YieldGard Plus® – Monsanto), MON863 x MON810 x NK603 (YieldGard Plus + RR Corn2® / YieldGard RR Maize® – Monsanto), MON863 x NK603 (YieldGard Rotworm + RR Corn 2® – Monsanto), MON863 (YieldBard RW® – Monsanto), MON89034 (YieldGard RW® – Monsanto), MON88017 (YieldGard VT RW® – Monsanto), MON810 + MON88017 (YieldGard VT Triple® – Monsanto), MON88017 + MON89034 (YieldGard VT Triple Pro® – Monsanto), Bt11 + MIR604 + GA21 (Agrisure 3000® – Syngenta), Bt11 + TC1507 + MIR604 + 5307 + GA21 (Syngenta), Bt11 + TC1507 + MIR604 + DAS59122 + GA21 (Agrisure 3122® – Syngenta), BT11 (Agrisure CB® – Syngenta), GA21 – (Agrisure GT® – Syngenta), MIR604 (Agrisure RW® – Syngenta), Bt11 + MIR162 (Agrisure TL VIP® – Syngenta), BT11 + MIR162

+ GA21 (Agrisure Viptra 3110® – Syngenta), BT11 + MIR162 + MIR604 (Agrisure TM 3100® – Syngenta), Event3272 + BT11 + MIR604 + GA21 (Syngenta), BT11 + MIR1692 + MIR604 + GA21 (Agrisure Viptera 3111® – Syngenta), BT11 + MIR 162 + TC1507 + GA21 (Agrisure Viptera 3220® – Syngenta), BT11 + MIR162 + TC1507 + MIR604 + 5307 + GA21 (Agrisure Viptera 3222® – Syngenta),
5 MIR162 (Syngenta), BT11 + GA21 + MIR162 + MIR604 + 5307 (Syngenta), 5307 (Syngenta).

In order to apply a compound of formula (I) as an insecticide, acaricide, nematocide or molluscicide to a pest, a locus of pest, or to a plant susceptible to attack by a pest, a compound of formula (I) is usually formulated into a composition which includes, in addition to the compound of formula (I), a
10 suitable inert diluent or carrier and, optionally, a surface active agent (SFA). SFAs are chemicals which are able to modify the properties of an interface (for example, liquid/solid, liquid/air or liquid/liquid interfaces) by lowering the interfacial tension and thereby leading to changes in other properties (for example dispersion, emulsification and wetting). It is preferred that all compositions (both solid and liquid formulations) comprise, by weight, 0.0001 to 95%, more preferably 1 to 85%, for example 5 to 60%, of a
15 compound of formula (I). The composition is generally used for the control of pests such that a compound of formula (I) is applied at a rate of from 0.1g to 10kg per hectare, preferably from 1g to 6kg per hectare, more preferably from 1g to 1kg per hectare.

When used in a seed dressing, a compound of formula (I) is generally used at a rate of 0.0001g to
10g (for example 0.001g or 0.05g), preferably 0.005g to 10g, more preferably 0.005g to 4g, per kilogram
20 of seed.

In another aspect the present invention provides a composition comprising a pesticidally effective amount of a compound of formula (I), in particular an insecticidal, acaricidal, nematocidal or molluscicidal composition comprising an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I) and a suitable carrier or diluent therefor. The composition
25 is preferably an insecticidal, acaricidal, nematocidal or molluscicidal composition.

The compositions can be chosen from a number of formulation types, including dustable powders (DP), soluble powders (SP), water soluble granules (SG), water dispersible granules (WG), wettable powders (WP), granules (GR) (slow or fast release), soluble concentrates (SL), oil miscible liquids (OL), ultra low volume liquids (UL), emulsifiable concentrates (EC), dispersible concentrates (DC), emulsions
30 (both oil in water (EW) and water in oil (EO)), micro-emulsions (ME), suspension concentrates (SC), aerosols, fogging/smoke formulations, capsule suspensions (CS) and seed treatment formulations. The formulation type chosen in any instance will depend upon the particular purpose envisaged and the physical, chemical and biological properties of the compound of formula (I).

Dustable powders (DP) may be prepared by mixing a compound of formula (I) with one or more
35 solid diluents (for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulfur, lime, flours, talc and other organic and inorganic solid carriers) and mechanically grinding the mixture to a fine powder.

Soluble powders (SP) may be prepared by mixing a compound of formula (I) with one or more water-soluble inorganic salts (such as sodium bicarbonate, sodium carbonate or magnesium sulfate) or one or more water-soluble organic solids (such as a polysaccharide) and, optionally, one or more wetting agents, one or more dispersing agents or a mixture of said agents to improve water

5 dispersibility/solubility. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water soluble granules (SG).

Wettable powders (WP) may be prepared by mixing a compound of formula (I) with one or more solid diluents or carriers, one or more wetting agents and, preferably, one or more dispersing agents and, optionally, one or more suspending agents to facilitate the dispersion in liquids. The mixture is then
10 ground to a fine powder. Similar compositions may also be granulated to form water dispersible granules (WG).

Granules (GR) may be formed either by granulating a mixture of a compound of formula (I) and one or more powdered solid diluents or carriers, or from pre-formed blank granules by absorbing a compound of formula (I) (or a solution thereof, in a suitable agent) in a porous granular material (such as
15 pumice, attapulgite clays, fuller's earth, kieselguhr, diatomaceous earths or ground corn cobs) or by adsorbing a compound of formula (I) (or a solution thereof, in a suitable agent) on to a hard core material (such as sands, silicates, mineral carbonates, sulfates or phosphates) and drying if necessary. Agents which are commonly used to aid absorption or adsorption include solvents (such as aliphatic and aromatic petroleum solvents, alcohols, ethers, ketones and esters) and sticking agents (such as polyvinyl acetates,
20 polyvinyl alcohols, dextrans, sugars and vegetable oils). One or more other additives may also be included in granules (for example an emulsifying agent, wetting agent or dispersing agent).

Dispersible Concentrates (DC) may be prepared by dissolving a compound of formula (I) in water or an organic solvent, such as a ketone, alcohol or glycol ether. These solutions may contain a surface active agent (for example to improve water dilution or prevent crystallization in a spray tank).

25 Emulsifiable concentrates (EC) or oil-in-water emulsions (EW) may be prepared by dissolving a compound of formula (I) in an organic solvent (optionally containing one or more wetting agents, one or more emulsifying agents or a mixture of said agents). Suitable organic solvents for use in ECs include aromatic hydrocarbons (such as alkylbenzenes or alkylnaphthalenes, exemplified by SOLVESSO 100, SOLVESSO 150 and SOLVESSO 200; SOLVESSO is a Registered Trade Mark), ketones (such as
30 cyclohexanone or methylcyclohexanone) and alcohols (such as benzyl alcohol, furfuryl alcohol or butanol), *N*-alkylpyrrolidones (such as *N*-methylpyrrolidone or *N*-octylpyrrolidone), dimethyl amides of fatty acids (such as C₈-C₁₀ fatty acid dimethylamide) and chlorinated hydrocarbons. An EC product may spontaneously emulsify on addition to water, to produce an emulsion with sufficient stability to allow spray application through appropriate equipment. Preparation of an EW involves obtaining a compound
35 of formula (I) either as a liquid (if it is not a liquid at room temperature, it may be melted at a reasonable temperature, typically below 70°C) or in solution (by dissolving it in an appropriate solvent) and then emulsifying the resultant liquid or solution into water containing one or more SFAs, under high shear, to produce an emulsion. Suitable solvents for use in EWs include vegetable oils, chlorinated hydrocarbons

(such as chlorobenzenes), aromatic solvents (such as alkylbenzenes or alkylnaphthalenes) and other appropriate organic solvents which have a low solubility in water.

Microemulsions (ME) may be prepared by mixing water with a blend of one or more solvents with one or more SFAs, to produce spontaneously a thermodynamically stable isotropic liquid formulation. A compound of formula (I) is present initially in either the water or the solvent/SFA blend. Suitable solvents for use in MEs include those hereinbefore described for use in ECs or in EWs. An ME may be either an oil-in-water or a water-in-oil system (which system is present may be determined by conductivity measurements) and may be suitable for mixing water-soluble and oil-soluble pesticides in the same formulation. An ME is suitable for dilution into water, either remaining as a microemulsion or forming a conventional oil-in-water emulsion.

Suspension concentrates (SC) may comprise aqueous or non-aqueous suspensions of finely divided insoluble solid particles of a compound of formula (I). SCs may be prepared by ball or bead milling the solid compound of formula (I) in a suitable medium, optionally with one or more dispersing agents, to produce a fine particle suspension of the compound. One or more wetting agents may be included in the composition and a suspending agent may be included to reduce the rate at which the particles settle. Alternatively, a compound of formula (I) may be dry milled and added to water, containing agents hereinbefore described, to produce the desired end product.

Aerosol formulations comprise a compound of formula (I) and a suitable propellant (for example *n*-butane). A compound of formula (I) may also be dissolved or dispersed in a suitable medium (for example water or a water miscible liquid, such as *n*-propanol) to provide compositions for use in non-pressurized, hand-actuated spray pumps.

A compound of formula (I) may be mixed in the dry state with a pyrotechnic mixture to form a composition suitable for generating, in an enclosed space, a smoke containing the compound.

Capsule suspensions (CS) may be prepared in a manner similar to the preparation of EW formulations but with an additional polymerization stage such that an aqueous dispersion of oil droplets is obtained, in which each oil droplet is encapsulated by a polymeric shell and contains a compound of formula (I) and, optionally, a carrier or diluent therefor. The polymeric shell may be produced by either an interfacial polycondensation reaction or by a coacervation procedure. The compositions may provide for controlled release of the compound of formula (I) and they may be used for seed treatment. A compound of formula (I) may also be formulated in a biodegradable polymeric matrix to provide a slow, controlled release of the compound.

A composition may include one or more additives to improve the biological performance of the composition (for example by improving wetting, retention or distribution on surfaces; resistance to rain on treated surfaces; or uptake or mobility of a compound of formula (I)). Such additives include surface active agents, spray additives based on oils, for example certain mineral oils or natural plant oils (such as soy bean and rape seed oil), and blends of these with other bio-enhancing adjuvants (ingredients which may aid or modify the action of a compound of formula (I)).

A compound of formula (I) may also be formulated for use as a seed treatment, for example as a powder composition, including a powder for dry seed treatment (DS), a water soluble powder (SS) or a water dispersible powder for slurry treatment (WS), or as a liquid composition, including a flowable concentrate (FS), a solution (LS) or a capsule suspension (CS). The preparations of DS, SS, WS, FS and 5 LS compositions are very similar to those of, respectively, DP, SP, WP, SC and DC compositions described above. Compositions for treating seed may include an agent for assisting the adhesion of the composition to the seed (for example a mineral oil or a film-forming barrier).

Wetting agents, dispersing agents and emulsifying agents may be surface SFAs of the cationic, anionic, amphoteric or non-ionic type.

10 Suitable SFAs of the cationic type include quaternary ammonium compounds (for example cetyltrimethyl ammonium bromide), imidazolines and amine salts.

Suitable anionic SFAs include alkali metals salts of fatty acids, salts of aliphatic monoesters of sulfuric acid (for example sodium lauryl sulfate), salts of sulfonated aromatic compounds (for example sodium dodecylbenzenesulfonate, calcium dodecylbenzenesulfonate, butylnaphthalene sulfonate and 15 mixtures of sodium di-*isopropyl*- and tri-*isopropyl*-naphthalene sulfonates), ether sulfates, alcohol ether sulfates (for example sodium laureth-3-sulfate), ether carboxylates (for example sodium laureth-3-carboxylate), phosphate esters (products from the reaction between one or more fatty alcohols and phosphoric acid (predominately mono-esters) or phosphorus pentoxide (predominately di-esters), for example the reaction between lauryl alcohol and tetraphosphoric acid; additionally these products may be 20 ethoxylated), sulfosuccinamates, paraffin or olefine sulfonates, taurates and lignosulfonates.

Suitable SFAs of the amphoteric type include betaines, propionates and glycines.

Suitable SFAs of the non-ionic type include condensation products of alkylene oxides, such as ethylene oxide, propylene oxide, butylene oxide or mixtures thereof, with fatty alcohols (such as oleyl alcohol or cetyl alcohol) or with alkylphenols (such as octylphenol, nonylphenol or octylcresol); partial 25 esters derived from long chain fatty acids or hexitol anhydrides; condensation products of said partial esters with ethylene oxide; block polymers (comprising ethylene oxide and propylene oxide); alkanolamides; simple esters (for example fatty acid polyethylene glycol esters); amine oxides (for example lauryl dimethyl amine oxide); and lecithins.

Suitable suspending agents include hydrophilic colloids (such as polysaccharides, 30 polyvinylpyrrolidone or sodium carboxymethylcellulose) and swelling clays (such as bentonite or attapulgate).

A compound of formula (I) may be applied by any of the known means of applying pesticidal compounds. For example, it may be applied, formulated or unformulated, to the pests or to a locus of the pests (such as a habitat of the pests, or a growing plant liable to infestation by the pests) or to any part of 35 the plant, including the foliage, stems, branches or roots, to the seed before it is planted or to other media in which plants are growing or are to be planted (such as soil surrounding the roots, the soil generally, paddy water or hydroponic culture systems), directly or it may be sprayed on, dusted on, applied by dipping, applied as a cream or paste formulation, applied as a vapor or applied through distribution or

incorporation of a composition (such as a granular composition or a composition packed in a water-soluble bag) in soil or an aqueous environment.

A compound of formula (I) may also be injected into plants or sprayed onto vegetation using electrodynamic spraying techniques or other low volume methods, or applied by land or aerial irrigation systems.

Compositions for use as aqueous preparations (aqueous solutions or dispersions) are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, the concentrate being added to water before use. These concentrates, which may include DCs, SCs, ECs, EWs, MEs, SGs, SPs, WPs, WGs and CSs, are often required to withstand storage for prolonged periods and, after such storage, to be capable of addition to water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. Such aqueous preparations may contain varying amounts of a compound of formula (I) (for example 0.0001 to 10%, by weight) depending upon the purpose for which they are to be used.

A compound of formula (I) may be used in mixtures with fertilizers (for example nitrogen-, potassium- or phosphorus-containing fertilizers). Suitable formulation types include granules of fertilizer. The mixtures preferably contain up to 25% by weight of the compound of formula (I).

The invention therefore also provides a fertilizer composition comprising a fertilizer and a compound of formula (I).

The compositions of this invention may contain other compounds having biological activity, for example micronutrients or compounds having fungicidal activity or which possess plant growth regulating, herbicidal, insecticidal, nematocidal or acaricidal activity.

The compound of formula (I) may be the sole active ingredient of the composition or it may be admixed with one or more additional active ingredients such as a pesticide, e.g. a insecticide, fungicide or herbicide, or a synergist or plant growth regulator where appropriate. An additional active ingredient may provide a composition having a broader spectrum of activity or increased persistence at a locus; synergize the activity or complement the activity (for example by increasing the speed of effect or overcoming repellency) of the compound of formula (I); or help to overcome or prevent the development of resistance to individual components. The particular additional active ingredient will depend upon the intended utility of the composition.

Examples of suitable pesticides include the following (where "Tx" means a compound of formula (I), and in particular a compound selected from Tables 1P to 90P and 1Q to 36Q, which may result in a synergistic combination with the given active ingredient):

a) Pyrethroids, such as permethrin +Tx, cypermethrin+Tx, fenvalerate+Tx, esfenvalerate+Tx, deltamethrin+Tx, cyhalothrin+Tx (in particular lambda-cyhalothrin+Tx and gamma cyhalothrin+Tx), bifenthrin+Tx, fenpropathrin+Tx, cyfluthrin+Tx, tefluthrin+Tx, fish safe pyrethroids+Tx (for example ethofenprox+Tx), natural pyrethrin+Tx, tetramethrin+Tx, S-bioallethrin+Tx, fenfluthrin+Tx, prallethrin+Tx, acrinathirin+Tx, ctofenprox +Tx or 5-benzyl-3-furylmethyl-(E)-(1R,3S)-2,2-dimethyl-3-(2-oxothiolan-3-ylidenemethyl)cyclopropane carboxylate+Tx;

- b) Organophosphates, such as profenofos+Tx, sulprofos+Tx, acephate+Tx, methyl parathion+Tx, azinphos-methyl+Tx, demeton-s-methy+Tx I, heptenophos+Tx, thiometon+Tx, fenamiphos+Tx, monocrotophos+Tx, profenofos+Tx, triazophos+Tx, methamidophos+Tx, dimethoate+Tx, phosphamidon+Tx, malathion+Tx, chlorpyrifos+Tx, phosalone+Tx, terbufos+Tx, fensulfothion+Tx, 5 fonofos+Tx, phorate+Tx, phoxim+Tx, pirimiphos-methyl+Tx, pirimiphos-ethyl+Tx, fenitrothion+Tx, fosthiazate+Tx or diazinon+Tx;
- c) Carbamates (including aryl carbamates), such as pirimicarb+Tx, triazamate+Tx, cloethocarb+Tx, carbofuran+Tx, furathiocarb+Tx, ethiofencarb+Tx, aldicarb+Tx, thiofurox+Tx, carbosulfan+Tx, bendiocarb+Tx, fenobucarb+Tx, propoxur+Tx, methomyl+Tx or oxamyl+Tx;
- 10 d) Benzoyl ureas, such as diflubenzuron+Tx, triflumuron+Tx, hexaflumuron+Tx, flufenoxuron+Tx, diafenthiuron+Tx, lufeneron+Tx, novaluron+Tx, noviflumuron+Tx or chlorfluazuron+Tx;
- c) Organic tin compounds, such as cyhexatin+Tx, fenbutatin oxide+Tx or azocyclotin+Tx;
- f) Pyrazoles, such as tebufenpyrad+Tx, tolfenpyrad+Tx, ethiprole+Tx, pyriprole+Tx, fipronil+Tx, and fenpyroximate+Tx;
- 15 g) Macrolides, such as avermectins or milbemycins, for example abamectin+Tx, emamectin benzoate+Tx, ivermectin+Tx, milbemycin+Tx, spinosad+Tx, azadirachtin+Tx, milbemectin+Tx, lepimectin+Tx or spinetoram+Tx;
- h) Hormones+Tx or pheromones+Tx;
- i) Organochlorine compounds, such as endosulfan+Tx (in particular alpha-endosulfan+Tx), benzene 20 hexachloride+Tx, DDT+Tx, chlordane+Tx or dieldrin+Tx;
- j) Amidines, such as chlordimeform+Tx or amitraz+Tx;
- k) Fumigant agents, such as chloropicrin+Tx, dichloropropane+Tx, methyl bromide+Tx or metam+Tx;
- l) Neonicotinoid compounds, such as imidacloprid+Tx, thiacloprid+Tx, acetamiprid+Tx, nitenpyram+Tx, dinotefuran+Tx, thiamethoxam+Tx, clothianidin+Tx, or nithiazine+Tx;
- 25 m) Diacylhydrazines+Tx, such as tebufenozide+Tx, chromafenozide+Tx or methoxyfenozide+Tx;
- n) Diphenyl ethers, such as diofenolan+Tx or pyriproxifen+Tx;
- o) Pyrazolines such as Indoxacarb+Tx or metaflumizone+Tx;
- p) Ketoenols, such as Spirotetramat+Tx, spiroticlofen+Tx or spiromesifen+Tx;
- q) Diamides, such as flubendiamide+Tx, chlorantraniliprole +Tx (Rynaxypyr®) or cyantraniliprole+Tx;
- 30 r) Essential oils such as Bugoil® - (PlantImpact); or
- s) a comopund selected from buprofezine+Tx, flonicamid+Tx, acequinocy+Tx I, bifenazate+Tx, cyenopyrafen+Tx, cyflumetofen+Tx, etoxazole+Tx, flometoquin+Tx, fluacrypyrim+Tx, fluensulfone+Tx, flufenerim+Tx, flupyradifuone+Tx, harpin+Tx, iodomethane+Tx, dodecadienol+Tx, pyridaben+Tx, pyridalyl+Tx, pyrimidifen+Tx, flupyradifurone+Tx, 4-[(6-Chloro-pyridin-3-ylmethyl)-(2,2-difluoro- 35 ethyl)-amino]-5H-furan-2-one (DE 102006015467), CAS: 915972-17-7 +Tx (WO 2006129714; WO2011/147953; WO2011/147952), CAS: 26914-55-8 (WO 2007020986), chlorfenapyr+Tx, pymetrozine+Tx, sulfoxaflor+Tx and pyrifluquinazon+Tx.

In addition to the major chemical classes of pesticide listed above, other pesticides having particular targets may be employed in the composition, if appropriate for the intended utility of the composition. For instance, selective insecticides for particular crops, for example stemborer specific insecticide (combinations such as cartap +Tx) or hopper specific insecticides (combinations such as buprofezin +Tx) for use in rice may be employed. Alternatively insecticides or acaricides specific for particular insect species/stages may also be included in the compositions (for example acaricidal ovo-larvicides, to give combinations such as clofentezine +Tx, flubenzimine +Tx, hexythiazox +Tx or tetradifon +Tx; acaricidal motilicides, to give combinations such as dicofol +Tx or propargite +Tx; acaricides, to give combinations such as bromopropylate +Tx or chlorobenzilate +Tx; or growth regulators, such as hydramethylnon +Tx, cyromazine +Tx, methoprene +Tx, chlorfluazuron +Tx or diflubenzuron +Tx).

Examples of fungicidal compounds and combinations which may be included in the composition of the invention are (E)-N-methyl-2-[2-(2,5-dimethylphenoxy)methyl]phenyl]-2-methoxy-iminoacetamide (SSF-129) +Tx, 4-bromo-2-cyano-N,N-dimethyl-6-trifluoromethylbenzimidazole-1-sulfonamide +Tx, α -[N-(3-chloro-2,6-xylyl)-2-methoxyacetamido]- γ -butyrolactone +Tx, 4-chloro-2-cyano-N,N-dimethyl-5-p-tolylimidazole-1-sulfonamide (IKF-916, cyamidazosulfamid) +Tx, 3-5-dichloro-N-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide (RH-7281, zoxamide) +Tx, N-allyl-4,5,-dimethyl-2-trimethylsilylthiophene-3-carboxamide (MON65500) +Tx, N-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy)propionamide (AC382042) +Tx, N-(2-methoxy-5-pyridyl)-cyclopropane carboxamide +Tx, acibenzolar (CGA245704) (e.g. acibenzolar-S-methyl) +Tx, alanycarb +Tx, aldimorph +Tx, anilazine +Tx, azaconazole +Tx, azoxystrobin +Tx, benalaxyl +Tx, benomyl +Tx, benthiavalicarb +Tx, biloxazol +Tx, bitertanol +Tx, bixafen +Tx, blastidicin S +Tx, boscalid +Tx, bromuconazole +Tx, bupirimate +Tx, captafol +Tx, captan +Tx, carbendazim +Tx, carbendazim +Tx, chlorhydrate +Tx, carboxin +Tx, carpropamid +Tx, carvone +Tx, CGA41396 +Tx, CGA41397 +Tx, chinomethionate +Tx, chlorothalonil +Tx, chlorozolate +Tx, clozylacon +Tx, copper containing compounds to give combinations such as copper oxychloride +Tx, copper oxyquinolate +Tx, copper sulfate +Tx, copper tallate +Tx and Bordeaux mixture +Tx, cyclufenamid +Tx, cymoxanil +Tx, cyproconazole +Tx, cyprodinil +Tx, debacarb +Tx, di-2-pyridyl disulfide 1,1'-dioxide +Tx, dichlofluanid +Tx, diclomezine +Tx, dicloran +Tx, diethofencarb +Tx, difenoconazole +Tx, difenzoquat +Tx, diflumetorim +Tx, O,O-di-iso-propyl-S-benzyl thiophosphate +Tx, dimefluazole +Tx, dimetconazole +Tx, dimethomorph +Tx, dimethirimol +Tx, diniconazole +Tx, dinocap +Tx, dithianon +Tx, dodecyl dimethyl ammonium chloride +Tx, dodemorph +Tx, dodine +Tx, doguadine +Tx, edifenphos +Tx, epoxiconazole +Tx, ethirimo +Tx, ethyl-(Z)-N-benzyl-N-([methyl(methyl-thioethylideneaminooxycarbonyl)amino]thio)- β -alaninate +Tx, etridiazole +Tx, famoxadone +Tx, fenamidone (RPA407213) +Tx, fenarimol +Tx, fenbuconazole +Tx, fenfuram +Tx, fenhexamid (KBR2738) +Tx, fenciclonil +Tx, fenpropidin +Tx, fenpropimorph +Tx, fentin acetate +Tx, fentin hydroxide +Tx, ferbam +Tx, ferimzone +Tx, fluazinam +Tx, fludioxonil +Tx, flumetover +Tx, fluopyram +Tx, fluoxastrobin +Tx, fluoroimide +Tx, fluquinconazole +Tx, flusilazole +Tx, flutolanil +Tx, flutriafol +Tx, fluxapyroxad +Tx, folpet +Tx,

- fuberidazole +Tx, furalaxyl +Tx, furametpyr +Tx, guazatine, +Tx hexaconazole +Tx, hydroxyisoxazole +Tx, hymexazole +Tx, imazalil +Tx, imibenconazole +Tx, iminoctadine +Tx, iminoctadine triacetate +Tx, ipconazole +Tx, iprobenfos +Tx, iprodione +Tx, iprovalicarb (SZX0722) +Tx, isopropanyl butyl carbamate +Tx, isoprothiolane +Tx, isopyrazam +Tx, kasugamycin +Tx, kresoxim-methyl +Tx,
- 5 LY186054 +Tx, LY211795 +Tx, LY248908 +Tx, mancozeb +Tx, mandipropamid +Tx, maneb +Tx, mefenoxam +Tx, metalaxyl +Tx, mepanipyrim +Tx, mepronil +Tx, metalaxyl +Tx, metconazole +Tx, metiram +Tx, metiram-zinc +Tx, metominostrobin +Tx, myclobutanil +Tx, neoasozin +Tx, nickel dimethyldithiocarbamate +Tx, nitrothal-*isopropyl* +Tx, nuarimol +Tx, ofurace +Tx, organomercury compounds, +Tx oxadixyl +Tx, oxasulfuron +Tx, oxolinic acid +Tx, oxpoconazole +Tx, oxycarboxin
- 10 +Tx, pefurazoate +Tx, penconazole +Tx, pencycuron +Tx, penflufen +Tx, penthiopyrad +Tx, phenazin oxide +Tx, phosetyl-Al +Tx, phosphorus acids +Tx, phthalide +Tx, picoxystrobin (ZA1963) +Tx, polyoxinD +Tx, polyram +Tx, probenazole +Tx, prochloraz +Tx, procymidone +Tx, propamocarb +Tx, propiconazole +Tx, propineb +Tx, propionic acid +Tx, prothioconazole +Tx, pyrazophos +Tx, pyrifenoxy +Tx, pyrimethanil +Tx, pyraclostrobin +Tx, pyroquilon +Tx, pyroxyfur +Tx, pyrrolnitrin +Tx, quaternary
- 15 ammonium compounds +Tx, quinomethionate +Tx, quinoxifen +Tx, quintozone +Tx, sedaxane +Tx, sipconazole (F-155) +Tx, sodium pentachlorophenate +Tx, spiroxamine +Tx, streptomycin +Tx, sulfur +Tx, tebuconazole +Tx, tecloftalam +Tx, tecnazene +Tx, tetraconazole +Tx, thiabendazole +Tx, thifluzamid +Tx, 2-(thiocyanomethylthio)benzothiazole +Tx, thiophanate-methyl +Tx, thiram +Tx, timibenconazole +Tx, tolclofos-methyl +Tx, tolylfluanid +Tx, triadimefon +Tx, triadimenol +Tx,
- 20 triazbutil +Tx, triazoxide +Tx, tricyclazole +Tx, tridemorph +Tx, trifloxystrobin (CGA279202) +Tx, triforine +Tx, triflumizole +Tx, triticonazole +Tx, validamycin A +Tx, vapam +Tx, vinclozolin +Tx, zineb +Tx and ziram +Tx, N-[9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide [1072957-71-1] +Tx, 1-methyl-3-difluoromethyl-1H-pyrazole-4-carboxylic acid (2-dichloromethylene-3-ethyl-1-methyl-indan-4-yl)-amide +Tx, and 1-
- 25 methyl-3-difluoromethyl-4H-pyrazole-4-carboxylic acid [2-(2,4-dichloro-phenyl)-2-methoxy-1-methyl-ethyl]-amide +Tx.

The active ingredients combinations described above comprising a compound selected of the invention, in particular from Tables 1P to 90P and 1Q to 36Q and an active ingredient as described above are preferably combined in a mixing ratio of from 100:1 to 1:6000, especially from 50:1 to 1:50,

30 more especially in a ratio of from 20:1 to 1:20, even more especially from 10:1 to 1:10, very especially from 5:1 and 1:5, special preference being given to a ratio of from 2:1 to 1:2, and a ratio of from 4:1 to 2:1 being likewise preferred, above all in a ratio of 1:1, or 5:1, or 5:2, or 5:3, or 5:4, or 4:1, or 4:2, or 4:3, or 3:1, or 3:2, or 2:1, or 1:5, or 2:5, or 3:5, or 4:5, or 1:4, or 2:4, or 3:4, or 1:3, or 2:3, or 1:2, or 1:600, or 1:300, or 1:150, or 1:35, or 2:35, or 4:35, or 1:75, or 2:75, or 4:75, or 1:6000, or 1:3000, or 1:1500, or

35 1:350, or 2:350, or 4:350, or 1:750, or 2:750, or 4:750. Those mixing ratios are understood to include, on the one hand, ratios by weight and also, on other hand, molar ratios.

In addition, biological agents may be included in the composition of the invention e.g. *Bacillus* species such as *Bacillus firmus*+Tx, *Bacillus cereus*+Tx, *Bacillus subtilis*+Tx, and *Pasteuria* species such

as *Pasteuria penetrans* +Tx and *Pasteuria nishizawae*+Tx. A suitable *Bacillus firmus* strain is strain CNCM I-1582 which is commercially available as BioNem™. A suitable *Bacillus cereus* strain is strain CNCM I-1562. Of both *Bacillus* strains more details can be found in US 6,406,690. Other biological organisms that may be included in the compositions of the invention are bacteria such as *Streptomyces* 5 *spp.* such as *S. avermitilis*, and fungi such as *Pochonia spp.* such as *P. chlamydosporia*+Tx. Also of interest are *Metarhizium spp.* such as *M. anisopliae*+Tx; *Pochonia spp.* such as *P. chlamydosporia*+Tx.

The compounds of formula (I) may be mixed with soil, peat or other rooting media for the protection of plants against seed-borne, soil-borne or foliar fungal diseases.

Examples of suitable synergists for use in the compositions include piperonyl butoxide, sesamex, 10 safroxan and dodecyl imidazole.

Suitable herbicides and plant-growth regulators for inclusion in the compositions will depend upon the intended target and the effect required.

An example of a rice selective herbicide which may be included is propanil. An example of a plant growth regulator for use in cotton is PLX™.

15 Some mixtures may comprise active ingredients which have significantly different physical, chemical or biological properties such that they do not easily lend themselves to the same conventional formulation type. In these circumstances other formulation types may be prepared. For example, where one active ingredient is a water insoluble solid and the other a water insoluble liquid, it may nevertheless be possible to disperse each active ingredient in the same continuous aqueous phase by dispersing the 20 solid active ingredient as a suspension (using a preparation analogous to that of an SC) but dispersing the liquid active ingredient as an emulsion (using a preparation analogous to that of an EW). The resultant composition is a suspoemulsion (SE) formulation.

The compounds of the invention are also useful in the field of animal health, e.g. they may be used against parasitic invertebrate pests, more preferably against parasitic invertebrate pests in or on an 25 animal. Examples of pests include nematodes, trematodes, cestodes, flies, mites, ticks, lice, fleas, true bugs and maggots. The animal may be a non-human animal, e.g. an animal associated with agriculture, e.g. a cow, a pig, a sheep, a goat, a horse, or a donkey, or a companion animal, e.g. a dog or a cat.

In a further aspect the invention provides a compound of the invention for use in a method of therapeutic treatment.

30 In a further aspect the invention relates to a method of controlling parasitic invertebrate pests in or on an animal comprising administering a pesticidally effective amount of a compound of the invention. The administration may be for example oral administration, parenteral administration or external administration, e.g. to the surface of the animal body. In a further aspect the invention relates to a compound of the invention for controlling parasitic invertebrate pests in or on an animal. In a further 35 aspect the invention relates to use of a compound of the invention in the manufacture of a medicament for controlling parasitic invertebrate pests in or on an animal

In a further aspect, the invention relates to a method of controlling parasitic invertebrate pests comprising administering a pesticidally effective amount of a compound of the invention to the environment in which an animal resides.

In a further aspect the invention relates to a method of protecting an animal from a parasitic
5 invertebrate pest comprising administering to the animal a pesticidally effective amount of a compound of the invention. In a further aspect the invention relates to a compound of the invention for use in protecting an animal from a parasitic invertebrate pest. In a further aspect the invention relates to use of a compound of the invention in the manufacture of a medicament for protecting an animal from a parasitic invertebrate pest.

10 In a further aspect the invention provides a method of treating an animal suffering from a parasitic invertebrate pest comprising administering to the animal a pesticidally effective amount of a compound of the invention. In a further aspect the invention relates to a compound of the invention for use in treating an animal suffering from a parasitic invertebrate pest. In a further aspect the invention relates to use of a compound of the invention in the manufacture of a medicament for treating an animal
15 suffering from a parasitic invertebrate pest.

In a further aspect, the invention provides a pharmaceutical composition comprising a compound of the invention and a pharmaceutically suitable excipient.

The compounds of the invention may be used alone or in combination with one or more other biologically active ingredients.

20 In one aspect the invention provides a combination product comprising a pesticidally effective amount of a component A and a pesticidally effective amount of component B wherein component A is a compound of the invention and component B is a compound as described below.

The compounds of the invention may be used in combination with anthelmintic agents. Such anthelmintic agents include, compounds selected from the macrocyclic lactone class of compounds such
25 as ivermectin, avermectin, abamectin, emamectin, eprinomectin, doramectin, selamectin, moxidectin, nemadectin and milbemycin derivatives as described in EP- 357460, EP-444964 and EP-594291. Additional anthelmintic agents include semisynthetic and biosynthetic avermectin/milbemycin derivatives such as those described in US-5015630, WO-9415944 and WO-9522552. Additional anthelmintic agents include the benzimidazoles such as albendazole, cambendazole, fenbendazole, flubendazole,
30 mebendazole, oxfendazole, oxbendazole, parbendazole, and other members of the class. Additional anthelmintic agents include imidazothiazoles and tetrahydropyrimidines such as tetramisole, levamisole, pyrantel pamoate, oxfendazole, morantel. Additional anthelmintic agents include flukicides, such as triclabendazole and clorsulon and the cestocides, such as praziquantel and epsiprantel.

The compounds of the invention may be used in combination with derivatives and analogues of
35 the paraherquamide/marcfortine class of anthelmintic agents, as well as the antiparasitic oxazolines such as those disclosed in US-5478855, US- 4639771 and DE-19520936.

The compounds of the invention may be used in combination with derivatives and analogues of the general class of dioxomorpholine antiparasitic agents as described in WO-9615121 and also with

anthelmintic active cyclic depsipeptides such as those described in WO-9611945, WO-9319053, WO-9325543, EP-626375, EP-382173, WO-9419334, EP-382173, and EP-503538.

The compounds of the invention may be used in combination with other ectoparasiticides; for example, fipronil; pyrethroids; organophosphates; insect growth regulators such as lufenuron; ecdysone agonists such as tebufenozide and the like; neonicotinoids such as imidacloprid and the like.

The compounds of the invention may be used in combination with terpene alkaloids, for example those described in International Patent Application Publication Numbers WO95/19363 or WO04/72086, particularly the compounds disclosed therein.

Other examples of such biologically active compounds that the compounds of the invention may be used in combination with include but are not restricted to the following:

Organophosphates: acephate, azamethiphos, azinphos-ethyl, azinphos-methyl, bromophos, bromophos-ethyl, cadusafos, chlorethoxyphos, chlorpyrifos, chlorfenvinphos, chlormephos, demeton, demeton-S-methyl, demeton-S-methyl sulphone, dialifos, diazinon, dichlorvos, dicrotophos, dimethoate, disulfoton, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fensulfothion, fenthion, flupyrzofos, fonofos, formothion, fosthiazate, heptenophos, isazophos, isothioate, isoxathion, malathion, methacriphos, methamidophos, methidathion, methyl-parathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, paraoxon, parathion, parathion-methyl, phenthoate, phosalone, phosfolan, phosphocarb, phosmet, phosphamidon, phorate, phoxim, pirimiphos, pirimiphos-methyl, profenofos, propaphos, proetamphos, prothiofos, pyraclofos, pyridapenthion, quinalphos, sulprophos, temephos, terbufos, tebupirimfos, tetrachlorvinphos, thimeton, triazophos, trichlorfon, vamidothion.

Carbamates: alanycarb, aldicarb, 2-sec-butylphenyl methylcarbamate, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenoxycarb, fenthioncarb, furathiocarb, HCN-801, isoprocarb, indoxacarb, methiocarb, methomyl, 5-methyl-m-cumenylbutyryl(methyl)carbamate, oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, triazamate, UC-51717.

Pyrethroids: acrinathin, allethrin, alphametrin, 5-benzyl-3-furylmethyl (E) - (1 R)-cis-2,2-dimethyl-3-(2-oxothiolan-3-ylidenemethyl)cyclopropanecarboxylate, bifenthrin, beta-cyfluthrin, cyfluthrin, a-cypermethrin, beta-cypermethrin, bioallethrin, bioallethrin((S)-cyclopentylisomer), bioresmethrin, bifenthrin, NCI-85193, cycloprothrin, cyhalothrin, cythithrin, cyphenothrin, deltamethrin, empenthrin, esfenvalerate, ethofenprox, fenfluthrin, fenpropathrin, fenvalerate, flucythrinate, flumethrin, fluvalinate (D isomer), imiprothrin, cyhalothrin, lambda-cyhalothrin, permethrin, phenothrin, prallethrin, pyrethrins (natural products), resmethrin, tetramethrin, transfluthrin, theta-cypermethrin, silafluofen, t-fluvalinate, tefluthrin, tralomethrin, Zeta-cypermethrin.

Arthropod growth regulators: a) chitin synthesis inhibitors: benzoylureas: chlorfluazuron, diflubenzuron, fluazuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron, buprofezin, diofenolan, hexythiazox, etoxazole, chlorfentazine; b) ecdysone antagonists: halofenozide, methoxyfenozide, tebufenozide; c) juvenoids: pyriproxyfen, methoprene (including S-methoprene), fenoxycarb; d) lipid biosynthesis inhibitors: spiroticlofen.

Other antiparasitics: acequinocyl, amitraz, AKD-1022, ANS-118, azadirachtin, *Bacillus thuringiensis*, bensultap, bifenazate, binapacryl, bromopropylate, BTG-504, BTG-505, camphechlor, cartap, chlorobenzilate, chlordimeform, chlorfenapyr, chromafenozide, clothianidine, cyromazine, diaclofen, diafenthiuron, DBI-3204, dinactin, dihydroxymethyldihydroxypyrrolidine, dinobuton, dinocap, 5 endosulfan, ethiprole, ethofenprox, fenazaquin, flumite, MTI- 800, fenpyroximate, fluacrypyrim, flubenzimine, flubrocyclopyrrolate, flufenazine, flufenprox, fluproxyfen, halofenprox, hydramethylnon, IKI-220, kanemite, NC-196, neem guard, nidinorterfuran, nitenpyram, SD-35651, WL-108477, piridaryl, propargite, prothifluorbutyl, pymethrozine, pyridaben, Buprofezin, pyrimidifen, NC-1111, R-195, RH-0345, RH-2485, RYI-210, S-1283, S-1833, SI-8601, silafluofen, silomazine, spinosad, tebufenpyrad, tetradifon, 10 tetranactin, thiacloprid, thiocyclam, thiamethoxam, tolfenpyrad, triazamate, triethoxyspinosyn, trinactin, verbutin, vertalec, YI-5301.

Fungicides: acibenzolar, aldimorph, ampropylfos, andoprim, azaconazole, azoxystrobin, benalaxyl, benomyl, bialaphos, blasticidin-S, Bordeaux mixture, bromuconazole, bupirimate, carpropamid, captan, captan, carbendazim, chlorfenazole, chloroneb, chloropicrin, chlorothalonil, 15 chlozolinate, copper oxychloride, copper salts, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, cyprofuram, RH-7281, diclocymet, diclobutrazole, diclomezine, dicloran, difenoconazole, RP-407213, dimethomorph, domoxystrobin, diniconazole, diniconazole-M, dodine, edifenphos, epoxiconazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fencaramid, fenpiclonil, fenpropidin, fenpropimorph, fentin acetate, fluazinam, fludioxonil, flumetover, flumorf/flumorlin, fentin hydroxide, 20 fluoxastrobin, fluquinconazole, flusilazole, flutolanil, flutriafol, folpet, fosetyl- aluminium, furalaxyl, furametapir, hexaconazole, ipconazole, iprobenfos, iprodione, isoprothiolane, kasugamycin, kresoxim-methyl, mancozeb, maneb, mefenoxam, mepronil, metalaxyl, metconazole, metominostrobin/fenominostrobin, metrafenone, myclobutanil, neo-asozin, nicobifen, orysastrobin, oxadixyl, penconazole, pencycuron, probenazole, prochloraz, propamocarb, propiconazole, proquinazid, 25 prothioconazole, pyrifenoxy, pyraclostrobin, pyrimethanil, pyroquilon, quinoxifen, spiroxamine, sulfur, tebuconazole, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tiadinil, triadimefon, triadimenol, tricyclazole, trifloxystrobin, triticonazole, validamycin, vinclozin.

Biological agents: *Bacillus thuringiensis* ssp *aizawai*, kurstaki, *Bacillus thuringiensis* delta endotoxin, baculovirus, entomopathogenic bacteria, virus and fungi.

30 Bactericides: chlortetracycline, oxytetracycline, streptomycin.

Other biological agents: enrofloxacin, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, omeprazole, tiamulin, benazepril, pyriprole, cefquinome, florfenicol, buserelin, cefovecin, tulathromycin, ceftiofur, carprofen, metaflumizone, praziquantel, triclofenazole.

When used in combination with other active ingredients, the compounds of the invention are 35 preferably used in combination with the following (where "Tx" means a compound of formula (I), and in particular a compound selected from Tables 1P to 90P and 1Q to 36Q, which may result in a synergistic combination with the given active ingredient): imidacloprid + Tx, enrofloxacin + Tx, praziquantel + Tx, pyrantel embonate + Tx, febantel + Tx, penethamate + Tx, moloxicam + Tx, cefalexin + Tx, kanamycin +

Tx, pimobendan + Tx, clenbuterol + Tx, fipronil + Tx, ivermectin + Tx, omeprazole + Tx, tiamulin + Tx, benazepril + Tx, milbemycin + Tx, cyromazine + Tx, thiamethoxam + Tx, pyriprole + Tx, deltamethrin + Tx, cefquinome + Tx, florfenicol + Tx, buserelin + Tx, cefovecin + Tx, tulathromycin + Tx, ceftiour + Tx, selamectin + Tx, carprofen + Tx, metaflumizone + Tx, moxidectin + Tx, methoprene (including S-
 5 methoprene) + Tx, clorsulon + Tx, pyrantel + Tx, amitraz + Tx, triclabendazole + Tx, avermectin + Tx, abamectin + Tx, emamectin + Tx, eprinomectin + Tx, doramectin + Tx, selamectin + Tx, nemadectin + Tx, albendazole + Tx, cambendazole + Tx, fenbendazole + Tx, flubendazole + Tx, mebendazole + Tx, oxfendazole + Tx, oxibendazole + Tx, parbendazole + Tx, tetramisole + Tx, levamisole + Tx, pyrantel pamoate + Tx, oxantel + Tx, morantel + Tx, triclabendazole + Tx, epsiprantel + Tx, fipronil + Tx,
 10 lufenuron + Tx, ecdysone + Tx or tebufenozide + Tx; more preferably, enrofloxacin + Tx, praziquantel + Tx, pyrantel embonate + Tx, febantel + Tx, penethamate + Tx, moloxicam + Tx, cefalexin + Tx, kanamycin + Tx, pimobendan + Tx, clenbuterol + Tx, omeprazole + Tx, tiamulin + Tx, benazepril + Tx, pyriprole + Tx, cefquinome + Tx, florfenicol + Tx, buserelin + Tx, cefovecin + Tx, tulathromycin + Tx, ceftiour + Tx, selamectin + Tx, carprofen + Tx, moxidectin + Tx, clorsulon + Tx, pyrantel + Tx,
 15 eprinomectin + Tx, doramectin + Tx, selamectin + Tx, nemadectin + Tx, albendazole + Tx, cambendazole + Tx, fenbendazole + Tx, flubendazole + Tx, mebendazole + Tx, oxfendazole + Tx, oxibendazole + Tx, parbendazole + Tx, tetramisole + Tx, levamisole + Tx, pyrantel pamoate + Tx, oxantel + Tx, morantel + Tx, triclabendazole + Tx, epsiprantel + Tx, lufenuron + Tx or ecdysone + Tx; even more preferably enrofloxacin + Tx, praziquantel + Tx, pyrantel embonate + Tx, febantel + Tx,
 20 penethamate + Tx, moloxicam + Tx, cefalexin + Tx, kanamycin + Tx, pimobendan + Tx, clenbuterol + Tx, omeprazole + Tx, tiamulin + Tx, benazepril + Tx, pyriprole + Tx, cefquinome + Tx, florfenicol + Tx, buserelin + Tx, cefovecin + Tx, tulathromycin + Tx, ceftiour + Tx, selamectin + Tx, carprofen + Tx, moxidectin + Tx, clorsulon + Tx or pyrantel + Tx.

Examples of ratios of compounds of formula I to mixing partner include 100:1 to 1:6000, 50:1 to
 25 1:50, 20:1 to 1:20, even more especially from 10:1 to 1:10, 5:1 to 1:5, 2:1 to 1:2, 4:1 to 2:1, 1:1, or 5:1, or 5:2, or 5:3, or 5:4, or 4:1, or 4:2, or 4:3, or 3:1, or 3:2, or 2:1, or 1:5, or 2:5, or 3:5, or 4:5, or 1:4, or 2:4, or 3:4, or 1:3, or 2:3, or 1:2, or 1:600, or 1:300, or 1:150, or 1:35, or 2:35, or 4:35, or 1:75, or 2:75, or 4:75, or 1:6000, or 1:3000, or 1:1500, or 1:350, or 2:350, or 4:350, or 1:750, or 2:750, or 4:750. Those
 mixing ratios are understood to include, on the one hand, ratios by weight and also, on other hand, molar
 30 ratios.

Of particular note is a combination where the additional active ingredient has a different site of action from the compound of formula I. In certain instances, a combination with at least one other parasitic invertebrate pest control active ingredient having a similar spectrum of control but a different site of action will be particularly advantageous for resistance management. Thus, a combination product
 35 of the invention may comprise a pesticidally effective amount of a compound of formula I and pesticidally effective amount of at least one additional parasitic invertebrate pest control active ingredient having a similar spectrum of control but a different site of action.

One skilled in the art recognizes that because in the environment and under physiological conditions salts of chemical compounds are in equilibrium with their corresponding non salt forms, salts share the biological utility of the non salt forms.

Thus a wide variety of salts of compounds of the invention (and active ingredients used in combination with the active ingredients of the invention) may be useful for control of invertebrate pests and animal parasites. Salts include acid-addition salts with inorganic or organic acids such as hydrobromic, hydrochloric, nitric, phosphoric, sulfuric, acetic, butyric, fumaric, lactic, maleic, malonic, oxalic, propionic, salicylic, tartaric, 4-toluenesulfonic or valeric acids.

The compounds of the invention also include N-oxides. Accordingly, the invention comprises combinations of compounds of the invention including N-oxides and salts thereof and an additional active ingredient including N-oxides and salts thereof.

The compositions for use in animal health may also contain formulation auxiliaries and additives, known to those skilled in the art as formulation aids (some of which may be considered to also function as solid diluents, liquid diluents or surfactants). Such formulation auxiliaries and additives may control: pH (buffers), foaming during processing (antifoams such polyorganosiloxanes), sedimentation of active ingredients (suspending agents), viscosity (thixotropic thickeners), in-container microbial growth (antimicrobials), product freezing (antifreezes), color (dyes/pigment dispersions), wash-off (film formers or stickers), evaporation (evaporation retardants), and other formulation attributes. Film formers include, for example, polyvinyl acetates, polyvinyl acetate copolymers, polyvinylpyrrolidone-vinyl acetate copolymer, polyvinyl alcohols, polyvinyl alcohol copolymers and waxes. Examples of formulation auxiliaries and additives include those listed in McCutcheon's Volume 2: Functional Materials, annual International and North American editions published by McCutcheon's Division, The Manufacturing Confectioner Publishing Co.; and PCT Publication WO 03/024222.

The compounds of the invention can be applied without other adjuvants, but most often application will be of a formulation comprising one or more active ingredients with suitable carriers, diluents, and surfactants and possibly in combination with a food depending on the contemplated end use. One method of application involves spraying a water dispersion or refined oil solution of the combination products. Compositions with spray oils, spray oil concentrations, spreader stickers, adjuvants, other solvents, and synergists such as piperonyl butoxide often enhance compound efficacy. Such sprays can be applied from spray containers such as a can, a bottle or other container, either by means of a pump or by releasing it from a pressurized container, e.g., a pressurized aerosol spray can. Such spray compositions can take various forms, for example, sprays, mists, foams, fumes or fog. Such spray compositions thus can further comprise propellants, foaming agents, etc. as the case may be. Of note is a spray composition comprising a pesticidally effective amount of a compound of the invention and a carrier. One embodiment of such a spray composition comprises a pesticidally effective amount of a compound of the invention and a propellant. Representative propellants include, but are not limited to, methane, ethane, propane, butane, isobutane, butene, pentane, isopentane, neopentane, pentene, hydrofluorocarbons, chlorofluorocarbons, dimethyl ether, and mixtures of the foregoing. Of note is a spray composition (and a

method utilizing such a spray composition dispensed from a spray container) used to control at least one parasitic invertebrate pest selected from the group consisting of mosquitoes, black flies, stable flies, deer flies, horse flies, wasps, yellow jackets, hornets, ticks, spiders, ants, gnats, and the like, including individually or in combinations.

5 The controlling of animal parasites includes controlling external parasites that are parasitic to the surface of the body of the host animal (e.g., shoulders, armpits, abdomen, inner part of the thighs) and internal parasites that are parasitic to the inside of the body of the host animal (e.g., stomach, intestine, lung, veins, under the skin, lymphatic tissue). External parasitic or disease transmitting pests include, for example, chiggers, ticks, lice, mosquitoes, flies, mites and fleas. Internal parasites include heartworms,
10 hookworms and helminths. The compounds of the invention may be particularly suitable for combating external parasitic pests. The compounds of the invention may be suitable for systemic and/or non-systemic control of infestation or infection by parasites on animals.

 The compounds of the invention may be suitable for combating parasitic invertebrate pests that infest animal subjects including those in the wild, livestock and agricultural working animals. Livestock
15 is the term used to refer (singularly or plurally) to a domesticated animal intentionally reared in an agricultural setting to make produce such as food or fiber, or for its labor; examples of livestock include cattle, sheep, goats, horses, pigs, donkeys, camels, buffalo, rabbits, hens, turkeys, ducks and geese (e.g., raised for meat, milk, butter, eggs, fur, leather, feathers and/or wool), cultured fish, honeybees. By combating parasites, fatalities and performance reduction (in terms of meat, milk, wool, skins, eggs, etc.)
20 are reduced, so that applying the compounds of the invention allows more economic and simple husbandry of animals.

 By controlling these pests it is intended to reduce deaths and improve performance (in the case of meat, milk, wool, hides, eggs, honey and the like) and health of the host animal. Also, controlling parasites may help to prevent the transmittance of infectious agents, the term "controlling" referring to the
25 veterinary field, meaning that the active compounds are effective in reducing the incidence of the respective parasite in an animal infected with such parasites to innocuous levels, e.g. the active compound is effective in killing the respective parasite, inhibiting its growth, or inhibiting its proliferation.

 The compounds of the invention may be suitable for combating parasitic invertebrate pests that infest companion animals and pets (e.g., dogs, cats, pet birds and aquarium fish), research and
30 experimental animals (e.g., hamsters, guinea pigs, rats and mice), as well as animals raised for/in zoos, wild habitats and/or circuses.

 In an embodiment of this invention, the animal is preferably a vertebrate, and more preferably a mammal, avian or fish. In a particular embodiment, the animal subject is a mammal (including great apes, such as humans). Other mammalian subjects include primates (e.g., monkeys), bovine (e.g., cattle or dairy
35 cows), porcine (e.g., hogs or pigs), ovine (e.g., goats or sheep), equine (e.g., horses), canine (e.g., dogs), feline (e.g., house cats), camels, deer, donkeys, buffalos, antelopes, rabbits, and rodents (e.g., guinea pigs, squirrels, rats, mice, gerbils, and hamsters). Avians include Anatidae (swans, ducks and geese), Columbidae (e.g., doves and pigeons), Phasianidae (e.g., partridges, grouse and turkeys), Thesienidae

(e.g., domestic chickens), Psittacines (e.g., parakeets, macaws, and parrots), game birds, and ratites (e.g., ostriches).

Birds treated or protected by the compounds of the invention can be associated with either commercial or noncommercial aviculture. These include Anatidae, such as swans, geese, and ducks, 5 Columbidae, such as doves and domestic pigeons, Phasianidae, such as partridge, grouse and turkeys, Thesienidae, such as domestic chickens, and Psittacines, such as parakeets, macaws and parrots raised for the pet or collector market, among others.

For purposes of the present invention, the term "fish" is understood to include without limitation, the Teleosti grouping of fish, i.e., teleosts. Both the Salmoniformes order (which includes the Salmonidae 10 family) and the Perciformes order (which includes the Centrarchidae family) are contained within the Teleosti grouping. Examples of potential fish recipients include the Salmonidae, Serranidae, Sparidae, Cichlidae, and Centrarchidae, among others.

Other animals are also contemplated to benefit from the inventive methods, including marsupials (such as kangaroos), reptiles (such as farmed turtles), and other economically important domestic animals 15 for which the inventive methods are safe and effective in treating or preventing parasite infection or infestation.

Examples of parasitic invertebrate pests controlled by administering a pesticidally effective amount of the compounds of the invention to an animal to be protected include ectoparasites (arthropods, acarines, etc.) and endoparasites (helminths, e.g., nematodes, trematodes, cestodes, acanthocephalans, etc. 20 and protozoae, such as coccidia).

The disease or group of diseases described generally as helminthiasis is due to infection of an animal host with parasitic worms known as helminths. The term 'helminths' is meant to include nematodes, trematodes, cestodes and acanthocephalans. Helminthiasis is a prevalent and serious economic problem with domesticated animals such as swine, sheep, horses, cattle, goats, dogs, cats and 25 poultry.

Among the helminths, the group of worms described as nematodes causes widespread and at times serious infection in various species of animals.

Nematodes that are contemplated to be treated by the compounds of the invention include, without limitation, the following genera: *Acanthocheilonema*, *Aelurostrongylus*, *Ancylostoma*, 30 *Angiostrongylus*, *Ascaridia*, *Ascaris*, *Brugia*, *Bunostomum*, *Capillaria*, *Chabertia*, *Cooperia*, *Crenosoma*, *Dictyocaulus*, *Diectophyme*, *Dipetalonema*, *Diphyllbothrium*, *Dirofilaria*, *Dracunculus*, *Enterobius*, *Filaroides*, *Haemonchus*, *Heterakis*, *Lagochilascaris*, *Loa*, *Mansonella*, *Muellerius*, *Necator*, *Nematodirus*, *Oesophagostomum*, *Ostertagia*, *Oxyuris*, *Parafilaria*, *Parascaris*, *Physaloptera*, *Protostrongylus*, *Setaria*, *Spirocerca*, *Stephanofilaria*, *Strongyloides*, *Strongylus*, *Thelazia*, *Toxascaris*, 35 *Toxocara*, *Trichinella*, *Trichonema*, *Trichostrongylus*, *Trichuris*, *Uncinaria* and *Wuchereria*.

Of the above, the most common genera of nematodes infecting the animals referred to above are *Haemonchus*, *Trichostrongylus*, *Ostertagia*, *Nematodirus*, *Cooperia*, *Ascaris*, *Bunostomum*, *Oesophagostomum*, *Chabertia*, *Trichuris*, *Strongylus*, *Trichonema*, *Dictyocaulus*, *Capillaria*, *Heterakis*,

Toxocara, *Ascaridia*, *Oxyuris*, *Ancylostoma*, *Uncinaria*, *Toxascaris* and *Parascaris*. Certain of these, such as *Nematodirus*, *Cooperia* and *Oesophagostomum* attack primarily the intestinal tract while others, such as *Haemonchus* and *Ostertagia*, are more prevalent in the stomach while others such as *Dictyocaulus* are found in the lungs. Still other parasites may be located in other tissues such as the heart
5 and blood vessels, subcutaneous and lymphatic tissue and the like.

Trematodes that are contemplated to be treated by the invention and by the inventive methods include, without limitation, the following genera: *Alaria*, *Fasciola*, *Nanophyetus*, *Opisthorchis*, *Paragonimus* and *Schistosoma*.

Cestodes that are contemplated to be treated by the invention and by the inventive methods
10 include, without limitation, the following genera: *Diphyllbothrium*, *Diplydium*, *Spirometra* and *Taenia*.

The most common genera of parasites of the gastrointestinal tract of humans are *Ancylostoma*, *Necator*, *Ascaris*, *Strongyloides*, *Trichinella*, *Capillaria*, *Trichuris* and *Enterobius*. Other medically important genera of parasites which are found in the blood or other tissues and organs outside the gastrointestinal tract are the filarial worms such as *Wuchereria*, *Brugia*, *Onchocerca* and *Loa*, as well
15 as *Dracunculus* and extra intestinal stages of the intestinal worms *Strongyloides* and *Trichinella*.

Numerous other helminth genera and species are known to the art, and are also contemplated to be treated by the compounds of the invention. These are enumerated in great detail in Textbook of Veterinary Clinical Parasitology, Volume 1, Helminths, E. J. L. Soulsby, F. A. Davis Co., Philadelphia, Pa.; Helminths, Arthropods and Protozoa, (6th Edition of Monnig's Veterinary Helminthology and
20 Entomology), E. J. L. Soulsby, Williams and Wilkins Co., Baltimore, Md.

The compounds of the invention may be effective against a number of animal ectoparasites (e.g., arthropod ectoparasites of mammals and birds in particular insects such as flies (stinging and licking), parasitic fly larvae, lice, hair lice, bird lice, fleas and the like; or acarids, such as ticks, for examples hard ticks or soft ticks, or mites, such as scab mites, harvest mites, bird mites and the like).

25 Insect and acarine pests include, e.g., biting insects such as flies and mosquitoes, mites, ticks, lice, fleas, true bugs, parasitic maggots, and the like.

Adult flies include, e.g., the horn fly or *Haematobia irritans*, the horse fly or *Tabanus* spp., the stable fly or *Stomoxys calcitrans*, the black fly or *Simulium* spp., the deer fly or *Chrysops* spp., the louse fly or *Melophagus ovinus*, and the tsetse fly or *Glossina* spp. Parasitic fly maggots include, e.g., the bot
30 fly (*Oestrus ovis* and *Cuterebra* spp.), the blow fly or *Phaenicia* spp., the screwworm or *Cochliomyia hominivorax*, the cattle grub or *Hypoderma* spp., the fleece worm and the *Gastrophilus* of horses. Mosquitoes include, for example, *Culex* spp., *Anopheles* spp. and *Aedes* spp.

Mites include *Mesostigmaphatalpha* spp. e.g., *mesostigmatids* such as the chicken mite, *Dermaphanyssus galphallinalphae*; itch or scab mites such as *Sarcoptidae* spp. for example,
35 *Salpharoptes scalphabiei*; mange mites such as *Psoroptidae* spp. including *Chorioptes bovis* and *Psoroptes ovis*; chiggers e.g., *Trombiculidae* spp. for example the North American chigger, *Trombiculalpha alphalfreddugesii*.

Ticks include, e.g., soft-bodied ticks including Argasidae spp. for example *Argasidaphas* spp. and *Ornithodoros* spp.; hard-bodied ticks including Ixodidae spp., for example *Rhipicephalaphalus sanguineus*, *Dermacentor variabilis*, *Dermacentor andersoni*, *Amblyomma americanum*, *Ixodes scapularis* and other *Rhipicephalus* spp. (including the former *Boophilus* genera).

5 Lice include, e.g., sucking lice, e.g., *Menopon* spp.

and *Bovicola* spp.; biting lice, e.g., *Haematopinus* spp., *Linognathus* spp. and *Solenopotes* spp.

Fleas include, e.g., *Ctenocephalides* spp., such as dog flea (*Ctenocephalides canis*) and cat flea (*Ctenocephalides felis*); *Xenopsylla* spp. such as oriental rat flea (*Xenopsylla cheopis*); and *Pulex* spp. such as human flea (*Pulex irritans*).

10 True bugs include, e.g., *Cimicidae* or e.g., the common bed bug (*Cimex lectularius*); *Triatominae* spp. including triatomid bugs also known as kissing bugs; for example *Rhodnius prolixus* and *Triatoma* spp.

Generally, flies, fleas, lice, mosquitoes, gnats, mites, ticks and helminths cause tremendous losses to the livestock and companion animal sectors. Arthropod parasites also are a nuisance to humans and can
15 vector disease-causing organisms in humans and animals.

Numerous other parasitic invertebrate pests are known to the art, and are also contemplated to be treated by the compounds of the invention. These are enumerated in great detail in Medical and Veterinary Entomology, D. S. Kettle, John Wiley AND Sons, New York and Toronto; Control of Arthropod Pests of Livestock: A Review of Technology, R. O. Drummand, J. E. George, and S. E. Kunz,
20 CRC Press, Boca Raton, Fla.

The compounds of the invention may also be effective against ectoparasites, e.g. insects such as flies (stinging and licking), parasitic fly larvae, lice, hair lice, bird lice, fleas and the like; or acarids, such as ticks, for examples hard ticks or soft ticks, or mites, such as scab mites, harvest mites, bird mites and the like. These include e.g. flies such as *Haematobia (Lyperosia) irritans* (horn fly), *Simulium* spp.
25 (blackfly), *Glossina* spp. (tsetse flies), *Hydrotaea irritans* (head fly), *Musca autumnalis* (face fly), *Musca domestica* (house fly), *Morellia simplex* (sweat fly), *Tabanus* spp. (horse fly), *Hypoderma bovis*, *Hypoderma lineatum*, *Lucilia sericata*, *Lucilia cuprina* (green blowfly), *Calliphora* spp. (blowfly), *Protophormia* spp., *Oestrus ovis* (nasal botfly), *Culicoides* spp. (midges), *Hippobosca equine*, *Gastrophilus intestinalis*, *Gastrophilus haemorrhoidalis* and *Gastrophilus nasalis*; lice such as *Bovicola*
30 (*Damalinia*) *bovis*, *Bovicola equi*, *Haematopinus asini*, *Felicola subrostratus*, *Heterodoxus spiniger*, *Lignonathus setosus* and *Trichodectes canis*; keds such as *Melophagus ovinus*; and mites such as *Psoroptes* spp., *Sarcoptes scabiei*, *Chorioptes bovis*, *Demodex equi*, *Cheyletiella* spp., *Notoedres cati*, *Trombicula* spp. and *Otodectes cyanotis* (ear mites).

Examples of species of animal health pesets include those from the order of the *Anoplurida*, for
35 example *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Phthirus* spp., *Solenopotes* spp.; particular examples are: *Linognathus setosus*, *Linognathus vituli*, *Linognathus ovillus*, *Linognathus oviformis*, *Linognathus pedalis*, *Linognathus stenopsis*, *Haematopinus asini macrocephalus*, *Haematopinus eurysternus*, *Haematopinus suis*, *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Phylloera*

- vastatrix*, *Phthirus pubis*, *Solenopotes capillatus*; from the order of the *Mallophagida* and the suborders *Amblycerina* and *Ischnocerina*, for example *Trimenopon* spp., *Menopon* spp., *Trinoton* spp., *Bovicola* spp., *Werneckiella* spp., *Lepikentron* spp., *Damalina* spp., *Trichodectes* spp., *Felicola* spp.; particular examples are: *Bovicola bovis*, *Bovicola ovis*, *Bovicola limbata*, *Damalina bovis*, *Trichodectes canis*.
- 5 *Felicola subrostratus*, *Bovicola caprae*, *Lepikentron ovis*, *Werneckiella equi*; from the order of the *Diptera* and the suborders *Nematocerina* and *Brachycerina*, for example *Aedes* spp., *Anopheles* spp., *Culex* spp., *Simulium* spp., *Eusimulium* spp., *Phlebotomus* spp., *Lutzomyia* spp., *Culicoides* spp., *Chrysops* spp., *Odagnia* spp., *Wilhelmia* spp., *Hybomitra* spp., *Atylotus* spp., *Tabanus* spp., *Haematopota* spp., *Philipomyia* spp., *Braula* spp., *Musca* spp., *Hydrotaea* spp., *Stomoxys* spp., *Haematobia* spp.,
- 10 *Morellia* spp., *Fannia* spp., *Glossina* spp., *Calliphora* spp., *Lucilia* spp., *Chrysomyia* spp., *Wohlfahrtia* spp., *Sarcophaga* spp., *Oestrus* spp., *Hypoderma* spp., *Gasterophilus* spp., *Hippobosca* spp., *Lipoptena* spp., *Melophagus* spp., *Rhinoestrus* spp., *Tipula* spp.; particular examples are: *Aedes aegypti*, *Aedes albopictus*, *Aedes taeniorhynchus*, *Anopheles gambiae*, *Anopheles maculipennis*, *Calliphora erythrocephala*, *Chrysosoma pluvialis*, *Culex quinquefasciatus*, *Culex pipiens*, *Culex tarsalis*, *Fannia*
- 15 *canicularis*, *Sarcophaga carnaria*, *Stomoxys calcitrans*, *Tipula paludosa*, *Lucilia cuprina*, *Lucilia sericata*, *Simulium reptans*, *Phlebotomus papatasi*, *Phlebotomus longipalpis*, *Odagnia ornata*, *Wilhelmia equina*, *Boophthora erythrocephala*, *Tabanus bromius*, *Tabanus spodopterus*, *Tabanus atratus*, *Tabanus sudeticus*, *Hybomitra ciurea*, *Chrysops caecutiens*, *Chrysops relictus*, *Haematopota pluvialis*, *Haematopota italica*, *Musca autumnalis*, *Musca domestica*, *Haematobia irritans irritans*, *Haematobia*
- 20 *irritans exigua*, *Haematobia stimulans*, *Hydrotaea irritans*, *Hydrotaea albipuncta*, *Chrysomya chloropyga*, *Chrysomya bezziana*, *Oestrus ovis*, *Hypoderma bovis*, *Hypoderma lineatum*, *Przhevalskiana silenus*, *Dermatobia hominis*, *Melophagus ovinus*, *Lipoptena capreoli*, *Lipoptena cervi*, *Hippobosca variegata*, *Hippobosca equina*, *Gasterophilus intestinalis*, *Gasterophilus haemorroidalis*, *Gasterophilus inermis*, *Gasterophilus nasalis*, *Gasterophilus nigricornis*, *Gasterophilus pecorum*, *Braula coeca*; from
- 25 the order of the *Siphonaptera*, for example *Pulex* spp., *Ctenocephalides* spp., *Tunga* spp., *Xenopsylla* spp., *Ceratophyllus* spp.; particular examples are: *Ctenocephalides canis*, *Ctenocephalides felis*, *Pulex irritans*, *Tunga penetrans*, *Xenopsylla cheopis*; from the order of the *Heteropterida*, for example *Cimex* spp., *Triatoma* spp., *Rhodnius* spp., *Panstrongylus* spp.; from the order of the *Blattarida*, for example *Blatta orientalis*, *Periplaneta americana*, *Blattella germanica*, *Supella* spp. (e.g. *Supella longipalpa*);
- 30 from the subclass of the *Acari* (*Acarina*) and the orders of the *Meta-* and *Mesostigmata*, for example *Argas* spp., *Ornithodoros* spp., *Otobius* spp., *Ixodes* spp., *Amblyomma* spp., *Rhipicephalus* (*Boophilus*) spp. *Dermacentor* spp., *Haemophysalis* spp., *Hyalomma* spp., *Dermanyssus* spp., *Rhipicephalus* spp. (the original genus of multi host ticks) *Ornithonyssus* spp., *Pneumonyssus* spp., *Raillietia* spp., *Pneumonyssus* spp., *Sternostoma* spp., *Varroa* spp., *Acarapis* spp.; particular examples are: *Argas persicus*, *Argas*
- 35 *reflexus*, *Ornithodoros moubata*, *Otobius megnini*, *Rhipicephalus* (*Boophilus*) *microplus*, *Rhipicephalus* (*Boophilus*) *decoloratus*, *Rhipicephalus* (*Boophilus*) *annulatus*, *Rhipicephalus* (*Boophilus*) *calceratus*, *Hyalomma anatolicum*, *Hyalomma aegypticum*, *Hyalomma marginatum*, *Hyalomma transiens*, *Rhipicephalus evertsi*, *Ixodes ricinus*, *Ixodes hexagonus*, *Ixodes canisuga*, *Ixodes pilosus*, *Ixodes*

- rubicundus*, *Ixodes scapularis*, *Ixodes holocyclus*, *Haemaphysalis concinna*, *Haemaphysalis punctata*, *Haemaphysalis cinnabarina*, *Haemaphysalis otophila*, *Haemaphysalis leachi*, *Haemaphysalis longicorni*, *Dermacentor marginatus*, *Dermacentor reticulatus*, *Dermacentor pictus*, *Dermacentor albipictus*, *Dermacentor andersoni*, *Dermacentor variabilis*, *Hyalomma mauritanicum*, *Rhipicephalus sanguineus*,
5 *Rhipicephalus bursa*, *Rhipicephalus appendiculatus*, *Rhipicephalus capensis*, *Rhipicephalus turanicus*, *Rhipicephalus zambeziensis*, *Amblyomma americanum*, *Amblyomma variegatum*, *Amblyomma maculatum*, *Amblyomma hebraeum*, *Amblyomma cajennense*, *Dermanyssus gallinae*, *Ornithonyssus bursa*, *Ornithonyssus sylviarum*, *Varroa jacobsoni*; from the order of the Actinedida (Prostigmata) and Acaridida (Astigmata), for example *Acarapis* spp., *Cheyletiella* spp., *Ornithocheyletia* spp., *Myobia* spp.,
10 *Psorergates* spp., *Demodex* spp., *Trombicula* spp., *Listrophorus* spp., *Acarus* spp., *Tyrophagus* spp., *Caloglyphus* spp., *Hypodectes* spp., *Pterolichus* spp., *Psoroptes* spp., *Chorioptes* spp., *Otodectes* spp., *Sarcoptes* spp., *Notoedres* spp., *Knemidocoptes* spp., *Cytodites* spp., *Laminosioptes* spp.; particular examples are: *Cheyletiella yasguri*, *Cheyletiella blakei*, *Demodex canis*, *Demodex bovis*, *Demodex ovis*, *Demodex caprae*, *Demodex equi*, *Demodex caballi*, *Demodex suis*, *Neotrombicula autumnalis*,
15 *Neotrombicula desaleri*, *Neoschongastia xerothermobia*, *Trombicula akamushi*, *Otodectes cynotis*, *Notoedres cati*, *Sarcoptes canis*, *Sarcoptes bovis*, *Sarcoptes ovis*, *Sarcoptes rupicaprae* (*S. caprae*), *Sarcoptes equi*, *Sarcoptes suis*, *Psoroptes ovis*, *Psoroptes cuniculi*, *Psoroptes equi*, *Chorioptes bovis*, *Psoergates ovis*, *Pneumonyssoides mange*, *Pneumonyssoides caninum*, *Acarapis woodi*, *Gasterophilus* spp., *Stomoxys* spp., *Trichodectes* spp., *Rhodnius* spp., *Ctenocephalides canis*, *Cimex lecturarius*,
20 *Ctenocephalides felis*, *Lucilia cuprina*; examples of acari include *Ornithodoros* spp., *Ixodes* spp., *Boophilus* spp..

Treatments of the invention are by conventional means such as by enteral administration in the form of, for example, tablets, capsules, drinks, drenching preparations, granulates, pastes, boli, feed-through procedures, or suppositories; or by parenteral administration, such as, for example, by injection
25 (including intramuscular, subcutaneous, intravenous, intraperitoneal) or implants; or by nasal administration; or by dermal application in the form of, for example, bathing or dipping, spraying, pouring-on and spotting-on, washing, dusting, and with the aid of active-compound-comprising shaped articles such as collars, ear tags, tail tags, limb bands, halters, marking devices and the like.

When compounds of the invention are applied in combination with an additional biologically
30 active ingredient, they may be administered separately e.g. as separate compositions. In this case, the biologically active ingredients may be administered simultaneously or sequentially. Alternatively, the biologically active ingredients may be components of one composition.

The compounds of the invention may be administered in a controlled release form, for example in subcutaneous or orally administered slow release formulations.

35 Typically a parasitocidal composition according to the present invention comprises a compound of the invention, optionally in combination with an additional biologically active ingredient, or N-oxides or salts thereof, with one or more pharmaceutically or veterinarily acceptable carriers comprising excipients and auxiliaries selected with regard to the intended route of administration (e.g., oral or

parenteral administration such as injection) and in accordance with standard practice. In addition, a suitable carrier is selected on the basis of compatibility with the one or more active ingredients in the composition, including such considerations as stability relative to pH and moisture content. Therefore of note are compounds of the invention for protecting an animal from an invertebrate parasitic pest
5 comprising a parasitically effective amount of a compound of the invention, optionally in combination with an additional biologically active ingredient and at least one carrier.

For parenteral administration including intravenous, intramuscular and subcutaneous injection, the compounds of the invention can be formulated in suspension, solution or emulsion in oily or aqueous vehicles, and may contain adjuncts such as suspending, stabilizing and/or dispersing agents.

10 The compounds of the invention may also be formulated for bolus injection or continuous infusion. Pharmaceutical compositions for injection include aqueous solutions of water-soluble forms of active ingredients (e.g., a salt of an active compound), preferably in physiologically compatible buffers containing other excipients or auxiliaries as are known in the art of pharmaceutical formulation. Additionally, suspensions of the active compounds may be prepared in a lipophilic vehicle. Suitable
15 lipophilic vehicles include fatty oils such as sesame oil, synthetic fatty acid esters such as ethyl oleate and triglycerides, or materials such as liposomes.

Aqueous injection suspensions may contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers. Alternatively, the active
20 ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water, before use.

In addition to the formulations described supra, the compounds of the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular or subcutaneous injection.

25 The compounds of the invention may be formulated for this route of administration with suitable polymeric or hydrophobic materials (for instance, in an emulsion with a pharmacologically acceptable oil), with ion exchange resins, or as a sparingly soluble derivative such as, without limitation, a sparingly soluble salt.

For administration by inhalation, the compounds of the invention can be delivered in the form of
30 an aerosol spray using a pressurized pack or a nebulizer and a suitable propellant, e.g., without limitation, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane or carbon dioxide. In the case of a pressurized aerosol, the dosage unit may be controlled by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or
35 starch.

The compounds of the invention may have favourable pharmacokinetic and pharmacodynamic properties providing systemic availability from oral administration and ingestion. Therefore after ingestion by the animal to be protected, parasitically effective concentrations of a compound of the

invention in the bloodstream may protect the treated animal from blood-sucking pests such as fleas, ticks and lice. Therefore of note is a composition for protecting an animal from an invertebrate parasite pest in a form for oral administration (i.e. comprising, in addition to a parasitically effective amount of a compound of the invention, one or more carriers selected from binders and fillers suitable for oral
5 administration and feed concentrate carriers).

For oral administration in the form of solutions (the most readily available form for absorption), emulsions, suspensions, pastes, gels, capsules, tablets, boluses, powders, granules, rumen-retention and feed/water/lick blocks, the compounds of the invention can be formulated with binders/fillers known in the art to be suitable for oral administration compositions, such as sugars and sugar derivatives (e.g.,
10 lactose, sucrose, mannitol, sorbitol), starch (e.g., maize starch, wheat starch, rice starch, potato starch), cellulose and derivatives (e.g., methylcellulose, carboxymethylcellulose, ethylhydroxycellulose), protein derivatives (e.g., zein, gelatin), and synthetic polymers (e.g., polyvinyl alcohol, polyvinylpyrrolidone). If desired, lubricants (e.g., magnesium stearate), disintegrating agents (e.g., cross-linked polyvinylpyrrolidinone, agar, alginic acid) and dyes or pigments can be added. Pastes and gels often also
15 contain adhesives (e.g., acacia, alginic acid, bentonite, cellulose, xanthan gum, colloidal magnesium aluminum silicate) to aid in keeping the composition in contact with the oral cavity and not being easily ejected.

In one embodiment a composition of the present invention is formulated into a chewable and/or edible product (e.g., a chewable treat or edible tablet). Such a product would ideally have a taste, texture
20 and/or aroma favored by the animal to be protected so as to facilitate oral administration of the compounds of the invention.

If the parasiticide compositions are in the form of feed concentrates, the carrier is typically selected from high-performance feed, feed cereals or protein concentrates.

Such feed concentrate-containing compositions can, in addition to the parasiticide active ingredients,
25 comprise additives promoting animal health or growth, improving quality of meat from animals for slaughter or otherwise useful to animal husbandry.

These additives can include, for example, vitamins, antibiotics, chemotherapeutics, bacteriostats, fungistats, coccidiostats and hormones.

The compound of the invention may also be formulated in rectal compositions such as
30 suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

The formulations for the method of this invention may include an antioxidant, such as BHT (butylated hydroxytoluene). The antioxidant is generally present in amounts of at 0.1- 5 percent (wt/vol). Some of the formulations require a solubilizer, such as oleic acid, to dissolve the active agent, particularly
35 if spinosad is included. Common spreading agents used in these pour-on formulations include isopropyl myristate, isopropyl palmitate, caprylic/capric acid esters of saturated C₁₂-C₁₈ fatty alcohols, oleic acid, oleic ester, ethyl oleate, triglycerides, silicone oils and dipropylene glycol methyl ether. The pour-on formulations for the method of this invention are prepared according to known techniques. Where the

pour-on is a solution, the parasiticide/insecticide is mixed with the carrier or vehicle, using heat and stirring if required. Auxiliary or additional ingredients can be added to the mixture of active agent and carrier, or they can be mixed with the active agent prior to the addition of the carrier. Pour-on formulations in the form of emulsions or suspensions are similarly prepared using known techniques.

5 Other delivery systems for relatively hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well-known examples of delivery vehicles or carriers for hydrophobic drugs. In addition, organic solvents such as dimethylsulfoxide may be used, if needed.

The rate of application required for effective parasitic invertebrate pest control (e.g. "pesticidally effective amount") will depend on such factors as the species of parasitic invertebrate pest to be
10 controlled, the pest's life cycle, life stage, its size, location, time of year, host crop or animal, feeding behavior, mating behavior, ambient moisture, temperature, and the like. One skilled in the art can easily determine the pesticidally effective amount necessary for the desired level of parasitic invertebrate pest control.

In general for veterinary use, the compounds of the invention are administered in a pesticidally
15 effective amount to an animal, particularly a homeothermic animal, to be protected from parasitic invertebrate pests.

A pesticidally effective amount is the amount of active ingredient needed to achieve an observable effect diminishing the occurrence or activity of the target parasitic invertebrate pest. One skilled in the art will appreciate that the pesticidally effective dose can vary for the various compounds
20 and compositions useful for the method of the present invention, the desired pesticidal effect and duration, the target parasitic invertebrate pest species, the animal to be protected, the mode of application and the like, and the amount needed to achieve a particular result can be determined through simple experimentation.

For oral or parenteral administration to animals, a dose of the compositions of the present
25 invention administered at suitable intervals typically ranges from about 0.01 mg/kg to about 100 mg/kg, and preferably from about 0.01 mg/kg to about 30 mg/kg of animal body weight.

Suitable intervals for the administration of the compositions of the present invention to animals range from about daily to about yearly. Of note are administration intervals ranging from about weekly to about once every 6 months. Of particular note are monthly administration intervals (i.e. administering the
30 compounds to the animal once every month).


It has now been found, surprisingly, that the animal health active ingredient mixtures according to the invention not only delivers about the additive enhancement of the spectrum of action with respect to the pest to be controlled that was in principle to be expected but achieves a synergistic effect which can extend the range of action of the component A and of the component B in two ways. Firstly, the rates of
35 application of the component A and of the component B are lowered whilst the action remains equally good. Secondly, the active ingredient mixture still achieves a high degree of pest control, sometimes even where the two individual components have become totally ineffective in such a low application rate range. This allows increased safety in use.

This synergistic effect applies in particular to mixtures where component A is a compound of formula I, in particular the compounds listed in Tables 1P to 90P, and component B is imidacloprid, enrofloxacin, praziquantel, pyrantel embonate, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, fipronil, ivermectin, omeprazole, tiamulin, benazepril, milbemycin, cyromazine, thiamethoxam, pyriprole, deltamethrin, cefquinome, florfenicol, buserelin, cefovecin, tulathromycin, ceftiofur, selamectin, carprofen, metaflumizone, moxidectin, methoprene (including S-methoprene), clorsulon, pyrantel, amitraz, triclabendazole, avermectin, abamectin, emamectin, eprinomectin, doramectin, selamectin, nemadectin, albendazole, cambendazole, fenbendazole, flubendazole, mebendazole, oxfendazole, oxibendazole, parbendazole, tetramisole, levamisole, pyrantel pamoate, oxantel, morantel, triclabendazole, epsiprantel, fipronil, lufenuron, ecdysone or tebufenozide. More preferably, component B is enrofloxacin, praziquantel, pyrantel embonate, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, omeprazole, tiamulin, benazepril, pyriprole, cefquinome, florfenicol, buserelin, cefovecin, tulathromycin, ceftiofur, selamectin, carprofen, moxidectin, clorsulon, pyrantel, eprinomectin, doramectin, selamectin, nemadectin, albendazole, cambendazole, fenbendazole, flubendazole, mebendazole, oxfendazole, oxibendazole, parbendazole, tetramisole, levamisole, pyrantel pamoate, oxantel, morantel, triclabendazole, epsiprantel, lufenuron or ecdysone. Even more preferably, component B is enrofloxacin, praziquantel, pyrantel embonate, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, omeprazole, tiamulin, benazepril, pyriprole, cefquinome, florfenicol, buserelin, cefovecin, tulathromycin, ceftiofur, selamectin, carprofen, moxidectin, clorsulon or pyrantel.

The following Examples illustrate, but do not limit, the invention. Documents referred to herein are incorporated by reference.


Examples

25 Preparation of *t*-butyl 4-hydroxyiminomethyl-2-bromobenzoate




To a solution of *t*-butyl 4-formyl-2-bromobenzoate (23 g, 80 mmol) in isopropanol:water (8:2, 160 ml) at 10°C was slowly added a solution of sodium acetate (8.2 g, 100 mmol) in water (20 ml) and hydroxylamine hydrochloride (7.23 g, 100 mmol) in water (20 ml). The reaction mixture was stirred for 3 hours at room temperature. After completion of reaction (monitored by TLC) the solvent was evaporated under reduced pressure. The reaction mixture was diluted with water (40 ml) and the aqueous layer was extracted with ethyl acetate (3 x 150 ml). The combined organic layers were dried over sodium sulphate and concentrated to give 4-hydroxyiminomethyl-2-bromobenzoate which was purified by column chromatography (24 g, Yield 99%). ^1H NMR (400 MHz, CDCl_3) 1.63 (s, 9H), 7.52 (dd, 1H), 7.69 (d, 1H), 7.84 (d, 1H), 8.06 (s, 1H) ppm.

Preparation of *t*-butyl 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-bromobenzoate



- 5 To a solution of *t*-butyl 4-hydroxyiminomethyl-2-bromobenzoate (24 g, 8 mmol) in N,N-dimethylformamide (180 ml) was added N-chlorosuccinimide (11.7 g, 84 mmol) and the reaction mixture was stirred at 50°C for 3 hours under a nitrogen atmosphere. The reaction mixture was then cooled to room temperature, treated with 1,3-dichloro-5-(3,3,3-trifluoroprop-1-en-2-yl)benzene (23.1 g, 96 mmol, in 50 ml of N,N-dimethylformamide) (prepared according to WO 2005/085216) and triethylamine
- 10 (22 ml, 160 mmol, dissolved in 22 ml of N,N-dimethylformamide) and stirred for 16 hours. The reaction mixture was then diluted with water (40 ml) and extracted with ethyl acetate (3 x 150 ml). The combined organic layers were dried over sodium sulphate, concentrated under reduced pressure and purified by column chromatography to give *t*-butyl 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-bromobenzoate (30 g, Yield 69.7 %). ¹H NMR (400 MHz, CDCl₃): 1.59 (s, 9H), 3.68 (d,
- 15 1H), 4.06 (d, 1H), 7.43 (d, 2H), 7.50 (d, 1H), 7.68 (dd, 1H), 7.72 (dd, 1H), 7.87 (d, 1H) ppm.


Preparation of *t*-butyl 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-ethylbenzoate



- 20 To the solution of *t*-butyl 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-bromobenzoate (1.5 g, 2.7 mmol) in tetrahydrofuran (30 ml), cesium carbonate (1.8 g, 5.5 mmol) and 1,1'-Bis(diphenylphosphino) ferrocene]dichloropalladium(II) (0.12 g, 0.13 mmol) were added. A triethylborane solution (1M in THF) (5.5 ml, 5.5 mmol) was then slowly added and the reaction mixture was stirred at 75 °C for 4 hours. After completion of the reaction, the solvent was evaporated under
- 25 reduced pressure. The reaction mixture was diluted with water (40 ml) and extracted with ethyl acetate (2 x 50 ml). The combined organic layers were dried over sodium sulphate, concentrated under reduced pressure and purified by column chromatography to give *t*-butyl 4-[5-(3,5-dichlorophenyl)-5-


trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-ethylbenzoate (1 g, Yield 74%). ^1H NMR (400 MHz, CDCl_3); 1.28 (t, 3H), 1.58 (s, 9H), 2.90 (q, 2H), 3.70 (d, 1H), 4.10 (d, 1H), 7.42-7.53 (m, 5H), 7.78 (d, 1H) ppm.

5 Preparation of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-ethylbenzoic acid



To a solution of *t*-butyl 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-ethylbenzoate (1g, 2 mmol) in dichloromethane (15ml) at 0°C was added trifluoroacetic acid (1.28 ml, 16mmol) under a nitrogen atmosphere. The reaction mixture was stirred for 24 hours at room temperature. After completion of the reaction, the solvent was evaporated under reduced pressure. The reaction mixture was diluted with water (40ml) and extracted with dichloromethane (2 x 50 ml). The combined organic layer were dried over sodium sulphate, concentrated under reduced pressure and purified by column chromatography to give 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-ethylbenzoic acid (0.88 g, Yield 100%). ^1H NMR (400 MHz, CDCl_3); 1.28 (t, 3H), 3.02 (q, 2H), 3.49 (d, 1H), 3.72 (d, 1H), 7.43-7.60 (m, 5H), 8.05 (d, 1H) ppm.

Preparation of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-ethyl-N-(thietan-3-yl)benzamide B69




20


To a solution of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-ethylbenzoic acid (0.3 g, 0.69 mmol) in dichloromethane (30 ml) containing 1 drop of N,N-dimethylformamide cooled at 0 °C, was added oxalyl chloride (0.07 ml, 0.83 mmol). The reaction mixture was stirred for 2 hours at room temperature under nitrogen and then was concentrated to dryness. The residue was dissolved in dichloromethane (30 ml) and treated with thietan-3-amine (0.074 g, 0.83 mmol) and triethylamine (0.23 ml, 1.7 mmol) in dichloromethane (20 ml) at 0 °C. The reaction mixture was stirred at room temperature overnight. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (40 ml). The aqueous layer was extracted with dichloromethane (2 x 50 ml). The combined organic layer was dried over sodium sulphate and concentrated under reduced pressure. The compound was

purified by column chromatography to give the title compound B69 as solid (0.24 g, Yield 68.7 %). ^1H NMR (400 MHz, CDCl_3): 1.25 (t, 3H), 2.80 (2H, q), 3.35 (t, 2H), 3.50 (t, 2H), 3.70 (d, 1H), 4.10 (d, 1H), 5.35-5.46 (1H, m), 6.25 (d, 1H), 7.35 (d, 1H), 7.39-7.43 (1H, m), 7.46-7.55 (3H, m), 7.62 (1H, s) ppm.

5 Nitration of 4-formylbenzoic acid:



Major



Minor


A solution of 4-formylbenzoic acid (0.2 g, 1.3mmol) in sulfuric acid (0.5 ml) was cooled to 0°C , treated with nitric acid (0.1 ml) over 5 min, and stirred for 20 hours at room temperature. The mixture was poured into an ice-cooled water and filtered to give a white solid. Purification by chromatography

10 (hexane/ethyl acetate 1:4 as eluent) provided two regio-isomers; 4-formyl-2-nitrobenzoic acid (major) and 4-formyl-3-nitrobenzoic acid (minor).

Data for the major isomer: ^1H -NMR (400 MHz, DMSO): 10.28 (1H, s), 8.48 (1H, d), 8.33-8.35 (1H, d), 7.93 (1H, d) ppm.


Data for the minor isomer: ^1H -NMR (400 MHz, DMSO): 10.03 (1H, s), 8.07 (1H, s), 8.05-8.06 (1H, m), 15 7.72 (1H, d) ppm.

Preparation of methyl 4-dimethoxymethyl-2-nitrobenzoate




A solution of 4-formyl-2-nitrobenzoic acid (0.1 g, 0.5 mmol) in thionyl chloride (2 ml) was refluxed for 20 hours. The reaction mixture was then cooled, the solvent was evaporated then methanol (1 ml) was added. The reaction mixture was stirred for 10 min at room temperature. Evaporation of excess of methanol gave methyl 4-dimethoxymethyl-2-nitrobenzoate (0.1 g). ^1H -NMR (400 MHz, CDCl_3): 8.41 (1H, d), 8.22 (1H, m), 7.86 (1H, d), 5.90 (1H, s), 3.94 (3H, s), 3.38 (6H, s) ppm.

25 Preparation of methyl 4-formyl-2-nitrobenzoate



A solution of methyl 4-dimethoxymethyl-2-nitrobenzoate (2 g, 7.8 mmol) in acetone (25 ml) was treated dropwise with hydrochloric acid (6M, 20 ml) and stirred for 2 hours. The solvent was removed under vacuo, the residue was then treated with water (25 ml) and extracted with ethyl acetate (2 X 50 ml). The combined organic layers were dried over sodium sulfate and concentrated to give methyl 4-formyl-2-nitrobenzoate (1.5 g). ¹H-NMR (400 MHz, CDCl₃): 10.10 (1H, s), 8.39-8.41 (1H, m), 8.16 (1H, d), 7.88 (1H, d), 3.94 (3H, s) ppm.


Preparation of methyl 4-hydroxyiminomethyl-2-nitrobenzoate



10

A solution of methyl 4-formyl-2-nitrobenzoate (2.0 g, 9.4 mmol) in isopropanol (50 ml) and water (25 ml) was treated with hydroxylamine hydrochloride (0.7 g) and sodium acetate (0.72 g) and stirred for 2 hours at room temperature. The solvent was then evaporated and the residue was treated with water (25 ml) and extracted with ethyl acetate (2 X 50 ml). The combined organic layers were dried over sodium sulfate and concentrated to give methyl 4-hydroxyimino-2-nitrobenzoate (1.8 g). ¹H-NMR (400 MHz, CDCl₃): 11.90 (1H, s), 8.20 (1H, d), 8.00 (1H, d), 7.89 (1H, d), 3.85 (3H, s) ppm.

Preparation of methyl 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoate




20

A solution of methyl 4-(hydroxyiminomethyl-2-nitrobenzoate (1.0 g, 4.4 mmol) in N,N-dimethylformamide (10 ml) was treated with N-chlorosuccinimide (0.85 g, 4.9 mmol) and stirred at 50 °C for 3 hours under a nitrogen atmosphere. The reaction mixture was cooled to room temperature, treated


with 1,3-dichloro-5-(3,3,3-trifluoroprop-1-en-2-yl)benzene (1.18g, 4.9 mmol) (prepared according to WO 2005/085216) and triethylamine (0.5ml), stirred at room temperature for 16 hours, treated with water (30 ml), and extracted with ethyl acetate (2 X 50 ml). The combined organic layers were dried over sodium sulfate, concentrated and purified by column chromatography (hexane/ethyl acetate 1:4 as eluent) to give methyl 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoate (1.2 g). ¹H-NMR (400 MHz, CDCl₃): 8.13 (1H, d), 7.95-7.98 (1H, m), 7.68 (1H, d), 7.50 (2H, m), 7.44-7.46 (1H, m), 4.10 (1H, d), 3.95 (3H, s), 3.73 (1H, d) ppm.

10 Preparation of 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoic acid



A solution of methyl 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoate (1 g, 0.43 mmol) and lithium hydroxide (73 mg) in tetrahydrofuran (20 ml) and water (10 ml) was stirred for 1 hour at room temperature. Tetrahydrofuran was then removed and the residue was extracted with ethyl acetate (2 x 25 ml). The combined organic layers were dried over sodium sulfate and concentrated to give 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoic acid (0.75 g), ¹H-NMR (400 MHz, DMSO): 7.85-7.87 (2H, m), 7.82 (1H, m), 7.66-7.69 (1H, m), 7.61-7.62 (2H, m), 4.22-4.46 (2H, m) ppm.

20 Preparation of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-nitro-N-(thietan-3-yl)benzamide




Oxalyl chloride (2 ml) was added dropwise to a solution of 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoic acid (0.1 g, 2.00 mmol) in dichloromethane (5 ml) containing 1 drop of N,N-dimethylformamide, stirred at room temperature under a nitrogen atmosphere for 4-6 hours, and concentrated. The residue was dissolved in tetrahydrofuran (10 ml), treated with thietan-3-amine (1 equivalent), triethylamine (2 equivalents) and stirred for 16 hours under a nitrogen atmosphere. The

reaction mixture was concentrated and purified by column chromatography to get 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-nitro-N-(thietan-3-yl)benzamide (0.47 g). ¹H-NMR (400 MHz, CDCl₃): 8.2 (1H, d), 8.01 (1H, m), 7.57 (1H, d), 7.45-7.50 (3H, m), 6.58 (1H, d), 5.33-5.40 (1H, m), 4.12 (1H, d), 3.75 (1H, d), 3.38-3.50 (4H, m) ppm.


5

Preparation of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-amino-N-(thietan-3-yl)benzamide




A suspension of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-nitro-N-(thietan-3-yl)benzamide (40 mg, 0.07 mmol) and iron powder (26 mg) in isopropanol (10 ml) and acetic acid (1 ml) was heated at 70°C for 3 hours. Isopropanol was evaporated and the residue was treated with water (10 ml) and extracted with dichloromethane (25 ml). The combined organic layers were dried over sodium sulfate, concentrated, and purified by column chromatography (hexane/ethyl acetate as eluent) to give 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-amino-N-(thietan-3-yl)benzamide (15 mg). ¹H-NMR (400 MHz, CDCl₃): 7.49 (2H, d), 7.42 (1H, m), 7.34 (1H, d), 6.92-6.94 (2H, m), 6.48 (1H, d) 5.18-5.20 (1H, m), 4.03 (1H, d), 3.64 (1H, d), 3.31-3.41 (4H, m), 0.82 (3H, m) ppm.

Preparation of methyl 4-formyl-3-nitrobenzoate



Following the same procedure as described for the preparation of methyl 4-formyl-2-nitrobenzoate, methyl 4-formyl-3-nitrobenzoate could be prepared. ¹H-NMR (400 MHz, CDCl₃): 10.44 (1H, s), 8.75 (1H, d), 8.16 (1H, d), 8.39-8.42 (1H, m), 7.99 (1H, d), 4.00 (3H, s) ppm.


Preparation of methyl 4-hydroxyiminomethyl-3-nitrobenzoate



Following the same procedure as described for the preparation of methyl 4-hydroxyiminomethyl-2-nitrobenzoate, methyl 4-hydroxyiminomethyl-3-nitrobenzoate was obtained. $^1\text{H-NMR}$ (400 MHz, CDCl_3): 12.08 (1H s), 8.45 (2H, m), 8.23-8.25 (1H, m), 8.03-8.05 (1H, m), 3.91 (3H, s) ppm.

5

Preparation of methyl 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-3-nitrobenzoate



Following the same procedure as described for the preparation of methyl 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitrobenzoate, the title compound could be obtained. $^1\text{H-NMR}$ (400 MHz, CDCl_3): 8.73 (1H, d), 8.33-8.36 (1H, m), 7.68 (1H, d), 7.44-7.46 (3H, m), 3.96-4.01 (4H, m), 3.69 (1H, d) ppm.

1

Preparation of 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-3-nitro-benzoic acid


15 acid



Following the same procedure as described for the preparation of 4-[3-(3,5-dichlorophenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-5-yl]-2-nitro-benzoic acid, the title compound could be obtained. $^1\text{H-NMR}$ (400 MHz, DMSO): 8.40 (1H, d), 8.19-8.21 (1H, m), 7.81-7.82 (1H, m), 7.70 (1H, d), 7.61 (2H, d), 4.17-4.32 (2H, m) ppm.


20

Preparation of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-3-nitro-N-(thietan-3-yl)benzamide



Following the same procedure as described for the preparation of 4-[5-(3,5-dichlorophenyl)-5-trifluoromethyl-4H-isoxazol-3-yl]-2-nitro-N-(thietan-3-yl)benzamide, the title compound could be obtained. $^1\text{H-NMR}$ (400 MHz, CDCl_3): 8.47 (1H, d), 8.11-8.13 (1H, m), 7.66 (1H, d), 7.45 (3H, d), 6.98 (1H, d), 5.40-5.46 (1H, m), 4.11 (1H, d), 3.98 (1H, d), 3.39-3.48 (4H, m) ppm.


Preparation of 4 thietane-3-carbonitrile



To a solution of epithiochlorohydrine (2.7 g) in benzene (10 ml) was added at room temperature a solution of potassium cyanide (4 g) in water (10 ml). The resulting mixture was heated to 50°C for 12 hours. The mixture was extracted with benzene then the organic phase was washed with aqueous saturated sodium hydrogenocarbonate solution then water and brine. The organic phase was dried with sodium sulphate then the solvent was evaporated to dryness to give the crude residue as a yellow oil (1.76g). 1 g of the crude was purified by a flash chromatography to obtain the title product as a brown solid (0.5 g). $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): 3.25-3.35 (m, 2H), 3.65-3.75 (m, 2H), 4.10-4.25 (m, 1H). $^1\text{H-NMR}$ (CDCl_3 , 100 MHz) 27.3 (1C), 28.7 (2C), 119.7 (1C). GCMS (Method C): $\text{rt} = 5.07$ min (87%) m/z : $[\text{M-CN}]^+ = 73$; $[\text{M}+1]^+ = 100$.

20


Preparation of 4 thietane-3-carbonitrile



To a solution of epithiochlorohydrine (5.4 g) in tetrahydrofuran (20 ml) was added at room temperature a solution of potassium cyanide (4.9 g) in water (20 ml). The resulting mixture was heated to 50°C for 12 hr. The mixture was extracted with tetrahydrofuran then the organic phase was washed with aqueous saturated sodium hydrogenocarbonate solution then water and brine. The organic phase was dried with sodium sulphate then the solvent was evaporated to dryness to give the crude title product as a violet oil (3.75g), which was analysed by NMR and GCMS and contained ca 10% of 2-aminothiophene.

Preparation of thietan-3-ylmethanamine

To a solution of lithium aluminium hydride (540 mg) in diethyl ether (30ml) was added in a suspension of
5 thietane-3-carbonitrile (1.4 g) in diethyl ether (30ml) at 0°C then the mixture was stirred at room
temperature for 30 min. The mixture was then diluted with ether (50ml), cooled to 0°C then water (0.55
ml), 15% aqueous sodium hydroxide (0.55ml) and water (1.60 ml) were added. The mixture was allowed
to warm to room temperature and stirred for 15 min. Anhydrous magnesium sulphate was added and
stirring was continued for 15 min, then the suspension was filtered on a celite pad. The solvent was
10 evaporated to dryness to give the crude title product as an oil (980 mg). ¹H-NMR (CDCl₃, 400 MHz):
3.21 (m, 3H); 3.00 (m, 2H); 2.86 (m, 2H).

Preparation of 2-(thietan-3-ylidene)acetonitrile

15 To a solution of thietan-3-one (900 mg) in 10 ml of dry methylene chloride was added a solution of
(triphenylphosphoranylidene)acetonitrile (3.31 g) in 20 ml of dry methylene chloride at 0°C. The solution
was allowed to warm to room temperature and after stirring for 15 min the solvent was removed under
reduced pressure. The residue was purified by a flash chromatography (silica gel: Dichloromethane as an
eluant) to give the title compound (785 mg) as a yellow-brown oil. GC/MS (Method C): rt = 2.89 min
20 (54.7 %) m/z: [M+1]⁺ = 112. ¹H-NMR (CDCl₃, 400 MHz): 5.11 (m, 1H), 4.15 (m, 2H), 4.03 (m, 2H).

Preparation of 2-(thietan-3-yl)acetonitrile

To a stirred solution of 2-(thietan-3-ylidene)acetonitrile (2.775 g) in dry methanol (250 mL), cooled to
25 0°C, was added sodium borohydride (9.250 g) in small portions over 30 min. The reaction mixture was
allowed to warm to room temperature and stirred for a further hour.

The solvent was removed under reduced pressure. The purple residue was dissolved in ethyl acetate (50
ml) and extracted with saturated sodium hydrogenocarbonate (2x50 ml). The organic layer, dried over
sodium sulphate, filtered and the solvent was removed under reduced pressure to yield a oil yellow. The
30 residue was purified by a flash chromatography (silica gel: cyclohexane/ethyl acetate as an eluant) to give
the title compound (2.2 g) as a yellow oil. GC/MS (Method C): rt = 5.99 min (100 %) m/z: [M+1]⁺ = 114.
¹H-NMR (CDCl₃, 400 MHz): 3.56 (m, 1H), 3.35 (dd, 2H), 3.09 (dd, 2H), 2.55 (d, 2H).

Preparation of 2-(thietan-3-yl)ethanamine

To a solution of lithium aluminium hydride (76 mg) in diethyl ether (4 ml) was added in a suspension of
 5 2-(thietan-3-yl)acetonitrile (226 mg) in diethyl ether (30ml) at 0°C then the mixture was stirred at room
 temperature until the end of the gas evolution. The mixture was then diluted with ether (6 ml), cooled to
 0°C then water (0.08 ml), 15% aqueous sodium hydroxide (0.08ml) and water (0.25 ml) were added. The
 mixture was allowed to warm to room temperature and stirred for 15 min. Anhydrous magnesium
 sulphate was added and stirring was continued for 15 min, then the suspension was filtered on a celite
 10 pad. The solvent was evaporated to dryness to give the crude title product as a brown oil (196 mg). ¹H-
 NMR (CDCl₃, 400 MHz): 3.4-3.35 (m, 1H); 3.20-3.10 (2H); 3.1-3.0 (m, 2H); 2.6 (m, 2H). GC/MS
 (Method C): rt = 5.80 min (90 %) m/z: [M+1]⁺ = 118.

Preparation of 4-tert-Butoxycarbonylamino-3-hydroxy-butyric acid

15

To a solution of 4-amino-3-hydroxybutyric acid (1.9 g) in aqueous 1M sodium hydroxide (35 ml) di-tert-
 butyl dicarbonate dissolved in dioxane (10ml) was added drop wise. The reaction mixture was stirred at
 ambient temperature for 72 hours. Dioxane was evaporated and the pH was adjusted to 2-3 by adding 1M
 hydrochloric acid. The mixture was extracted three times with ethyl acetate. The organic phases were
 20 combined, dried over anhydrous sodium sulfate and concentrated. The residue was purified over silica gel
 (ethyl acetate / methanol 1:1) to give 4-tert-butoxycarbonylamino-3-hydroxy-butyric acid (3.42 g).
¹H-NMR (CDCl₃, 400 MHz): 5.05 (s, 1H), 4.07 (s, 1H), 3.28 (m, 1H), 3.10 (s, 1H), 2.48 (m, 2H), 1.40 (s,
 9H).

25 Preparation of (2,4-Dihydroxy-butyl)-carbamic acid tert-butyl ester

To a borane tetrahydrofuran complex solution 1M in tetrahydrofuran (15ml) was added at 0°C under an
 atmosphere of argon a solution of 4-tert-butoxycarbonylamino-3-hydroxy-butyric acid (1.5 g) in
 tetrahydrofuran (35ml). The reaction mixture was stirred at 0°C for 3 hours. The reaction was quenched
 30 by drop wise addition of 10% acetic acid in methanol (30ml). Ethyl acetate was added (250ml) and the
 mixture was washed with 1M hydrochloric acid (60ml) and 1M sodium hydrogenocarbonate (80ml). The
 organic phase was dried over anhydrous sodium sulfate and concentrated. The residue was purified over

silica gel (ethyl acetate / methanol 9:1) to give (2,4-dihydroxy-butyl)-carbamic acid tert-butyl ester (0.83 g).

¹H-NMR (CDCl₃, 400 MHz): 5.02 (s, 1H), 3.95 (s, 1H), 3.88 (m, 2H), 3.25 (m, 1H), 3.15 (m, 1H), 1.70 (m, 2H), 1.45 (s, 9H).

5

Preparation of Methanesulfonic acid 1-(tert-butoxycarbonylamino-methyl)-3-methanesulfonyloxy-propyl ester




To a solution of (2,4-Dihydroxy-butyl)-carbamic acid tert-butyl ester in dichloromethane (15 ml) was added triethylamine (1.96 ml). The reaction mixture was cooled to 0°C and methanesulfonyl chloride (0.82 ml) was added drop wise. The mixture was stirred at 0°C for 2 hours. The mixture was filtered and dichloromethane was added (200ml). The mixture was washed with 0.2M hydrochloric acid (50 ml), water (30 ml) and 0.2M sodium carbonate (50ml). The organic phase was dried over anhydrous sodium sulfate and concentrated to give crude methanesulfonic acid 1-(tert-butoxycarbonylamino-methyl)-3-

15 methanesulfonyloxy-propyl ester (1.24 g) which was used in the next step without further purification.

¹H-NMR (CDCl₃, 400 MHz): 4.95 (m, 1H), 3.85 (m, 2H), 3.95 (m, 2H), 3.10 (m, 6H), 2.15 (m, 2H), 1.45 (s, 9H).

Preparation of Thietan-2-ylmethyl-carbamic acid tert-butyl ester



20

To a solution of crude methanesulfonic acid 1-(tert-butoxycarbonylamino-methyl)-3-methanesulfonyloxy-propyl ester (1.24 g) in methanol (25 ml) was added sodium sulfide monohydrate (0.84 g). The reaction mixture was stirred at 40°C for 3 hours. The methanol was partially removed by distillation. Water was added and the mixture was extracted three times with ethyl acetate. The organic phases were combined, dried over anhydrous sodium sulfate and concentrated. The residue was purified over silica gel (ethyl acetate / heptanes, from 1:4 to 1:1) to give thietan-2-ylmethyl-carbamic acid tert-butyl ester (0.15 g).

¹H-NMR (CDCl₃, 400 MHz): 4.95 (s, 1H), 3.70 (m, 1H), 3.40 (m, 2H), 3.15 (m, 1H), 3.05 (m, 1H), 2.95 (m, 1H), 2.60 (m, 1H), 1.45 (s, 9H).

30

Preparation of Thietan-2-yl-methylamine hydrochloride




To a solution of thietan-2-ylmethyl-carbamic acid tert-butyl ester (2 g) in diethyl ether (20 ml) was added 2M hydrochloric acid in diethyl ether (24.5 ml) at 0°C. The reaction mixture was allowed to warm to ambient and stirred for 72 hours. The diethyl ether was removed by distillation. The residue was washed 5 with hexane and filtered to give thietan-2-yl-methanamine hydrochloride (1.11 g).

¹H-NMR (D₂O, 400 MHz): 4.95 (m, 1H), 3.35 (m, 2H), 3.25-3.05 (m, 3H), 2.65 (m, 1H).


Method for preparing compounds of the invention from a carboxylic acid

4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-ylmethyl)benzamide




To a suspension of 4-[5-(3,5-dichloro-phenyl)-5-trifluoromethyl-4,5-dihydro-isoxazol-3-yl]-benzoic acid (836 mg, prepared as described for example in WO200/9080250) in dichloromethane (30 ml) was added oxalyl chloride (280 mg) and then one drop of N,N-dimethylformamide. The reaction mixture stirred at room temperature until the solids were dissolved, and the solvent was evaporated in vacuo to give a solid, which was dissolved in dichloromethane (10 ml) and the resulting solution added dropwise to a solution of triethylamine (0.5 ml) and thietan-3-ylmethanamine (example B, 250 mg) in dichloromethane (20ml) at room temperature, under argon. The reaction was stirred overnight at room 20 temperature, diluted with water, and the organic layer washed with 1N hydrochloric acid solution, then with 1N sodium hydroxide solution, then with brine. The organic phase dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Purification by column chromatography (eluent heptane / ethyl acetate) afforded the title compound as a solid (570 mg). ¹H-NMR (CDCl₃, 400 MHz): 7.60-7.40 (m, 6H), 6.02 (m, 1H), 4.10 (d, 1H), 3.70 (d, 1H), 3.65 (t, 2H), 3.55 (m, 1H), 3.33 (t, 2H), 3.05 (dd, 2H), 2.50 (s, 25 3H).

Preparation of 4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(1,1-dioxo-thietan-3-ylmethyl)benzamide



To a solution of 4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-ylmethyl)benzamide (505 mg) in dichloromethane (50 ml) at 0°C was added a solution of sodium hydrogenocarbonate (504 mg) in water (10 ml) then 3-chloroperbenzoic acid (70% w/w, 615 mg). The resulting mixture was allowed to stir at room temperature for 1 hour. The reaction mixture was then extracted with dichloromethane then the organic phase was washed twice with water, dried over anhydrous sodium sulfate, filtered and evaporated to give the crude product as a solid. Purification by chromatography using the ISCO Rf machine (dichloromethane then 5% methanol) afforded the title product (300 mg) as a white solid. ¹H-NMR (CDCl₃, 400 MHz): 7.60-7.40 (m, 6H), 6.25 (m, 1H), 4.30 (dd, 2H), 4.10 (d, 1H), 4.00 (dd, 2H), 3.75 (t, 2H), 3.72 (d, 1H), 3.00 (m, 1H), 2.50 (s, 3H).

General method BOP T°C for preparing the compounds of the invention in parallel




15 This general method was used to prepare a number of compounds in parallel.

To a solution of the appropriate carboxylic acid (30 μmol), for example 4-[5-[3,5-bis(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid (made as described in WO2009080250) for Compound A1 of Table A, in *N,N*-dimethylacetamide ("DMA") (0.4 ml) was added a solution of the appropriate amine (36 μmol), for example 3-methyl-thietan-3-ylamine (made as described in WO 2007/080131) for Compound A1 of Table A, in *N,N*-

dimethylacetamide (0.145 ml) followed by diisopropylethylamine (Hunig's base) (0.04 ml, 60 μ mol) and a solution of bis(2-oxo-3-oxazolidinyl)phosphonic chloride ("BOP-Cl") (15.3 mg) in *N,N*-dimethylacetamide (0.2 ml). The reaction mixture was stirred for 16 hours at T°C. Then the mixture was diluted with acetonitrile (0.6 ml) and a sample was used for the LC-MS analysis. The remaining mixture was further diluted with acetonitrile / *N,N*-dimethylformamide (4:1) (0.8 ml) and purified by HPLC to give the desired compound.


General Suzuki method for preparing the compounds of the invention in parallel




10 This general method was used to prepare a number of compounds in parallel.

In a microwave vial, the appropriate boronic acid (2eq.), for example (3-ethoxyphenyl)boronic acid, is introduced. Then, a suspension of the appropriate halogeno compound (0.5 mL from a solution of 0.019 mmol in 9.5 mL of acetonitrile), for example 4-[5-(3-bromo-5-chloro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(1-oxothietan-3-yl)benzamide, for Compound A103 of Table A, is added followed by 0.3 mL of acetonitrile, potassium carbonate (14eq.) and a suspension of $\text{PdCl}_2(\text{PPh}_3)_2$ (0.2 mL from a solution of 56.7 mg in 5 mL of acetonitrile). The vials are flushed with argon and sealed, the suspension was heated for 700s at 120°C in a microwave. After removing the solvent of the reaction, the crude residue was extracted with 2 mL of ethyl acetate and 2 mL of water. The organic layer is then filtered and then concentrated under vacuo. The crude residue was dissolved in 0.8 mL of DMF and purified on preparative chromatography to give the desired compound.

Preparation of 2-methyl-N-(1-oxothietan-3-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide (C17)




Step A: tert-butyl 2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoate



A 1500 mL reaction-vessel was charged at RT with 1-Anthracen-9-ylmethyl quininium chloride
 5 13.77 mmol, 7.59 g) and tert-butyl 2-methyl-4-[(Z)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-
 2-enoyl]benzoate (68.87 mmol, 34 g) and dissolved in chloroform (700 mL). The mixture
 cooled down to -30°C. Then, a solution of hydroxylamine (50 mass% in H₂O, 9.1 g) in water
 (50.9 mL) and cesium hydroxide hydrate (185.9 mmol, 32.88 g) in water (60 mL) were added
 simultaneously dropwise to the reaction mixture over 30 min. After the addition the reaction was
 10 kept 16h at -30°C after which time the conversion was complete.

The reaction was brought to room temperature diluted with dichloromethane and washed three
 times with water. The organic phase was dried over MgSO₄ and evaporated to yield crude title
 product. The product was purified over a silica gel column (eluent: EtOAc /Heptane) to obtain
 39.0 g of the title product. LCMS (Method GR): RT 1.44 min, no ionisation: ¹H NMR (400
 15 MHz, CDCl₃) 16.0 (s, 9H), 2.60 (s, 3H), 3.70 (d, 1H), 4.10-4.20 (m, 2H), 7.25-7.90 (m, 5H), ¹⁹F-
 NMR (CDCl₃, 376 MHz): -79.40.


Step B: 2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoic acid



20 To a solution of tert-butyl 2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-
 isoxazol-3-yl]benzoate (64.87 mmol, 33 g) in dichloromethane (300 mL) was added
 trifluoroacetic acid (389.2 mmol, 44.6 g) at 0°C. After addition the reaction was brought to
 room temperature and the mixture stirred over night after which time LC-MS analysis showed a
 complete conversion. The crude mixture was concentrated and the residue dissolved in 300 mL
 25 of ethylacetate, washed with brine, dried over MgSO₄ and evaporated to yield crude title product
 which was purified over a silica gel column (eluent: CH₂Cl₂ / EtOAc) to obtain the desired
 product as pale yellow foam. LCMS (Method GR) RT 1.23 min, [M+H]⁺ 450/452; ¹H-NMR
 (CDCl₃, 400 MHz): 2.7 (s, 3H), 3.70 (d, 2H), 4.10 (d, 2H), 7.60 (s, 3H), 7.01 (d, 1H), 8.15 (d,


2H), 7.01 (d, 1H), 8.15 (d, 2H), 11.40 (s, 1H); ^{19}F -NMR (CDCl_3 , 376 MHz): -79.38.

Step C: 2-methyl-N-(thietan-3-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide



To a solution of 2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoic acid (28.72 mmol, 13 g) in dichloromethane (100mL) and N,N-dimethylformamide (0.2867 mmol, 0.0210 g) was added oxalyl chloride (4.836 g) dropwise at room temperature and under argon (the liberated hydrochloric acid gas was scrubbed with a sodium hydroxide 0.5N solution). The mixture was stirred at room temperature for 1 h and at 30°C for 20 min until the gas evolution ceased. The reaction mixture was concentrated to dryness and redissolved in dichloromethane (100mL), triethylamine (71.81 mmol, 7.340 g) was added at 0-5 °C under argon followed by dropwise addition of thietan-3-amine (28.72 mmol, 2.561 g). The mixture was allowed to warm to room temperature and stirred for 16 h. The mixture was extracted with aqueous hydrochloric acid (0.5N), water, brine, dried over MgSO_4 , filtered and evaporated affording the crude product. Purification over silica gel (eluent: CH_2Cl_2 / EtOAc) yielded the title compound. LCMS (Method GR) RT 1.25 min, $[\text{M}+\text{H}]^+$ 523/525; ^1H -NMR (CDCl_3 , 400 MHz): 2.45 (s, 3H), 3.35 (m, 2H), 3.50 (m, 2H), 3.60 (d, 1H), 7.41 (d, 1H), 5.40 (q, 1H), 6.20 (d, NH), 7.40 (s, 1H), 7.51 (d, 1H), 7.40 (s, 1H), 7.50 (s, 2H), 7.65 (s, 2H). ^{19}F -NMR (CDCl_3 , 376 MHz): -79.43.

Step D: 2-methyl-N-(1-oxidothietan-1-ium-3-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide (C17)



To a suspension of 2-methyl-N-(thietan-3-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide in dichloromethane (92 mL) was added m-CPBA (24.3 mmol, 5.58 g) portion wise at 0-5 °C, and the reaction mixture was stirred for 10 min. A saturated $\text{Na}_2\text{S}_2\text{O}_3$ solution was added, and the mixture was stirred for 10 min. The organic layer


was washed with sat. NaHCO_3 solution, water and brine, dried over MgSO_4 , filtered and evaporated. Purification over silica gel (cluent: CH_2Cl_2 / EtOAc) yielded the title compound. LCMS (Method GR) 1.13 min, M^+ 539/541; ^1H NMR (400 MHz, CDCl_3) 2.50 (s, 3H), 3.28 (m, 1H), 3.70 (d, 1H), 4.10-4.20 (m, 4H), 4.70 (m, 1H), 6.50 (d, 1H), 7.40-7.70 (1d + 4s, 5H), ^{19}F -NMR (CDCl_3 , 376 MHz): -79.43.

Chiral HPLC method

Column: Daicel CHIRALPAK® IB, $3\mu\text{m}$, 0.46cm x 10cm


Mobile phase: Heptan/EtOH 80/20

10 Retention times of eluting isomers: 4.85 min (8.2%) 5.66 min (91.8%).




Similarly, N-(1,1-dioxothietan-3-yl)-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide (C19) could be obtained: LCMS (Method GR) 1.17 min, M^+ 553/555; ^{19}F -NMR (CDCl_3 , 376 MHz): -79.44.

Preparation of 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(1,1-dioxothietan-3-yl)-2-methyl-benzamide (C18)



20

Step A: *tert*-butyl 4-[(*E*)-3-(3,5-dichloro-4-fluoro-phenyl)-4,4,4-trifluoro-but-2-enoyl]-2-methyl-benzoate




A 250 mL reaction-vessel equipped with a Dean-Stark apparatus was charged with *tert*-butyl 4-[(*E*)-3-(3,5-dichloro-4-fluoro-phenyl)-4,4,4-trifluoro-but-2-enoyl]-2-methyl-benzoate (19.7 g, 45.3 mmol) and

dissolved in toluene (120 mL) followed by addition of tert-butyl 4-acetyl-2-methyl-benzoate (11.7 g, 49.8 mmol) and triethylamine (0.638 mL). Then potassium carbonate (4 g) was added reaction stirred at reflux for 4 h at which time another portion of potassium carbonate (4 g) was added and the reaction mixture was stirred over night at reflux. LC-MS showed an almost complete conversion to desired product. The reaction mixture was concentrated to 50 mL and diluted with 500 mL dichloromethane and washed with water two times. The organic layer was dried over MgSO_4 , evaporated to yield 24.2 g of the crude title product. The product was purified over a silica gel column (eluent: EtOAc /Heptane) to obtain 12.8 g of the title product.

LCMS (Method GR) RT 1.38 min; ^1H -NMR (CDCl_3 , 400 MHz): 1.6 (s, 9H), 2.6 (s, 3H), 7.2 (d, 2H), 7.40 (s, 1H), 7.6 (s, 2H), 7.8 (d, 1H). ^{19}F NMR (CDCl_3 , 376 MHz): -66.68, -113.

Step B: tert-butyl 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoate




A 1500 mL reaction-vessel was charged at room temperature with 1-Anthracen-9-ylmethyl quininium chloride (2.93 g, 5.32 mmol) and tert-butyl 4-[(E)-3-(3,5-dichloro-4-fluoro-phenyl)-4,4,4-trifluoro-but-2-enoyl]-2-methyl-benzoate (12.7 g, 26.6 mmol) and dissolved in chloroform (700 mL). The mixture cooled down to -20°C . Then, a solution hydroxylamine (50 mass% in water, 3.3 mL) in water (16.7 mL) and cesium hydroxide (10.8 g, 71.8 mmol) in water (20 mL) were added simultaneously dropwise to the reaction mixture over 12 min. After the addition the reaction was kept 1.5 h at -20°C after which time the conversion was complete.

The reaction was brought to room temperature diluted with dichloromethane and washed three times with water. The organic phase was dried over MgSO_4 and evaporated to yield 24.2 g of the crude title product. The product was purified over a silica gel column (eluent: EtOAc /Heptane) to obtain 12.6 gr of the title product.

LCMS (Method GR) Rt 1.4 min.; ^1H -NMR (CDCl_3 , 400 MHz): 1.6 (s, 9H), 2.6 (s, 3H), 3.7 (d, 1H), 4.2 (d, 1H), 7.40 (d, 2H), 7.6 (d, 2H), 7.8 (d, 1H). ^{19}F -NMR (CDCl_3 , 376 MHz): -79.6, -113.

Step C: 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid




A 100mL vessel was charged with tert-butyl 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoate (3.1 g, 6.3 mmol) and dissolved in CH_2Cl_2 (30 mL) and then cooled down to 0°C . To this mixture 2,2,2-trifluoroacetic acid (4.3 g, 38 mmol) was added over 10min. After addition the reaction was brought to room temperature and the mixture stirred over night after which time LC-MS analysis showed a complete conversion.

The crude mixture was concentrated and the residue dissolved in 300mL of ethylacetate, washed with brine, dried over MgSO_4 and evaporated to yield 2.7g of the crude title product which was directly used for the step D.

10 LCMS (Method GR) RT 1.19 min, $[\text{M}-\text{H}]^+ 434$; $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): 2.6 (s, 3H), 3.68 (d, 1H), 4.10 (d, 1H), 7.6 (t, 4H), 8.2 (d, 1H), 11.5(s,1H) $^{19}\text{F-NMR}$ (CDCl_3 , 376 MHz): -79.6, -114.

Step D: 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(1,1-dioxothietan-3-yl)-2-methyl-benzamide (C18)



15

A 100mL vessel was charged with 1,1-dioxothietan-3-amine hydrochloride (1.2g) in 25mL of N,N-dimethylformamide followed by successive addition of triethylamine (2.1 mL), hydroxyazabenzotriazole (1g), 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (1.4g) and 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid (2.6g) dissolved in N,N-dimethylformamide (5mL). The mixture was then stirred overnight at room temperature. The reaction mixture was diluted in 600mL of ethyl acetate, washed 3-times with 500mL of brine, then with 0.5N HCl cold, dried over MgSO_4 and evaporated to yield 4.6g of the crude title compound. The product was purified over a silica gel column (eluent: CH_2Cl_2 / EtOAc) to yield 2g of the title compound.

20 LCMS (Method GR) RT 1.13 min, $[\text{M}-\text{H}]^+ 536, 537, 539, 541$
 $^1\text{H-NMR}$ (CDCl_3 , 400 MHz): 2.50 (s, 3H), 3.7 (d, 1H), 4.02 (m, 3H), 4.7 (m, 2H), 4.9 (m, 1H), 6.5 (d, 1H), 7.4 (d, 1H), 7.5 (d, 2H), 7.6 (d, 2H). $^{19}\text{F-NMR}$ (CDCl_3 , 376 MHz): -79.4, -115. Mp: 134°C - 136°C .

Chiral HPLC method


30 Column: Daicel CHIRALPAK® IB, $3\mu\text{m}$, $0.46\text{cm} \times 10\text{cm}$

Mobile phase: Hept/EtOH/DEA: 80/20 /0.1%

Flow rate: 1.0 ml/min

Retention times of eluting isomers: 7.15min (11.3%) 9.31min (87.1%).

5 Preparation of 2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(1-oxidothietan-1-ium-3-yl)benzamide (C15)



^1H NMR (DMSO- d_6 , 400MHz): δ = 9.17 (d, J =7.3 Hz, 1 H), 7.82 - 7.87 (m, 2 H), 7.79 (dd, J =7.9, 1.7 Hz, 1 H), 7.56 - 7.67 (m, 3 H), 4.44 (s, 1 H), 4.38 (s, 2 H), 4.05 - 4.19 (m, 2 H), 3.22 ppm (dd, J =12.3, 10.1 Hz, 2 H).

Chiral HPLC analysis:

HPLC: Waters UPLC - HClass

DAD Detector Waters UPLC

Column: Daicel CHIRALPAK[®] IB, 3 μ m, 0.46cm x 10cm

15 Mobile phase: Heptan/EtOH 90/10

Flow rate: 1 ml/min


Detection: 266nm

Sample concentration: 1mg/mL in Heptan/2-Propanol 50/50

Injection: 2 μ L

20 1st eluting isomer: 13.99 min (25.9 %), 2nd eluting isomer: 15.51 min (74.1%)

Preparation of 5-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-3-methyl-N-(1-oxidothietan-1-ium-3-yl)pyridine-2-carboxamide (C16)



25

^1H NMR (CDCl₃, 400MHz): δ = 8.69 (d, J =1.5 Hz, 1 H), 8.52 (d, J =8.1 Hz, 1 H), 7.83 - 7.94 (m, 1 H), 7.48 - 7.57 (m, 2 H), 7.40 - 7.48 (m, 1 H), 4.58 (d, J =7.7 Hz, 1 H), 4.05 - 4.24 (m, 3 H), 3.74 (d, J =17.2 Hz, 1 H), 3.25 - 3.43 (m, 2 H), 2.77 ppm (s, 3 H). m.p.=221-240°C.

30 **Chiral HPLC analysis:**

HPLC: Waters UPLC - HClass

DAD Detector Waters UPLC

Column: Daicel CHIRALPAK® IB, 3µm, 0.46cm x 10cm

Mobile phase: EtOH/MeOH 50/50

5 Flow rate: 1 ml/min

Detection: 266nm

Sample concentration: 1mg/mL in Heptan/2-Propanol 50/50

Injection: 2µL

1st eluting isomer: 2.82 min (83.6 %), 2nd eluting isomer: 4.10 min (16.4%)

10

GC/MS methods:

Method C

Volatile CI/EI	GCMS was conducted on a Thermo, MS: DSQ and GC: TRACE GC ULTRA with a column from Zebron phenomenex: Phase ZB-5ms 15 m, diam: 0.25 mm, 0.25 µm, H ₂ flow 1.7 ml/min, temp injector: 250°C, temp detector: 220°C, method: hold 2 min at 40 °C, 25°C/min until 320°C, hold 1min12s at 320°C, total time 15min. CI reagent gas: Methane, flow 1ml/min
-----------------------	--

LC/MS methods:

15 Method A:

ACQUITY SQD Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Ionisation method: Electrospray

Polarity: positive ions

Capillary (kV) 3.00, Cone (V) 20.00, Extractor (V) 3.00, Source Temperature (°C) 150, Desolvation

20 Temperature (°C) 400, Cone Gas Flow (L/Hr) 60, Desolvation Gas Flow (L/Hr) 700

Mass range: 100 to 800 Da

DAD Wavelength range (nm): 210 to 400

Method Waters ACQUITY UPLC with the following HPLC gradient conditions

(Solvent A: Water/Methanol 9:1, 0.1% formic acid and Solvent B: Acetonitrile, 0.1% formic acid)

25 Type of column: Waters ACQUITY UPLC HSS T3; Column length: 30 mm; Internal diameter of column: 2.1 mm; Particle Size: 1.8 micron; Temperature: 60°C.

Time (minutes)	A (%)	B (%)	Flow rate (ml/min)
0	100	0	0.75
2.5	0	100	0.75
2.8	0	100	0.75
3	100	0	0.75

Method B:

ZQ Mass Spectrometer from Waters (single quadrupole mass spectrometer), ionization method: electrospray, polarity: positive ionization, capillary (kV) 3.00, cone (V) 30.00, extractor (V) 3.00, source temperature (°C) 100, desolvation temperature (°C) 200, cone gas flow (L/Hr) 200, desolvation gas flow 5 (L/Hr) 250, mass range: 150 to 800 Da.

1100er Series HPLC from Agilent: quaternary pump, heated column compartment and diode-array detector.

Column: Waters Atlantis dc18; length: 20 mm; internal diameter: 3 mm; particle size: 3 µm, temperature (°C) 40, DAD wavelength range (nm): 200 to 500, solvent gradient: A = 0.1% of formic acid in water 10 and B: 0.1% of formic acid in acetonitrile.

Time (minutes)	A (%)	B (%)	Flow rate (ml/min)
0.0	90	10	1.7
5.5	0.0	100	1.7
5.8	0.0	100	1.7
5.9	90	10	1.7

Method C:

ZMD Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Ionisation method: Electrospray

15 Polarity: positive ions

Capillary (kV) 3.00, Cone (V) 30.00, Extractor (V) 3.00, Source Temperature (°C) 150, Desolvation Temperature (°C) 320, Cone Gas Flow (L/Hr) 50, Desolvation Gas Flow (L/Hr) 400

Mass range: 150 to 800 Da

Column: Waters Atlantis dc18; length: 20 mm; internal diameter: 3 mm; particle size: 3 µm, temperature 20 (°C) 40, DAD wavelength range (nm): 200 to 500,

Waters Alliance 2795 LC with the following HPLC gradient conditions (Solvent A: 0.1% of formic acid in water and Solvent B: 0.1% of formic acid in acetonitrile)

Time (minutes)	A (%)	B (%)	Flow rate (ml/min)
0	80	20	1.7
5	0	100	1.7
5.6	0	100	1.7
6	80	20	1.7

Method D:

25 ZQ Mass Spectrometer from Waters (single quadrupole mass spectrometer), ionization method: electrospray, polarity: positive ionization, capillary (kV) 3.00, cone (V) 30.00, extractor (V) 3.00, source temperature (°C) 100, desolvation temperature (°C) 200, cone gas flow (L/Hr) 200, desolvation gas flow (L/Hr) 250, mass range: 150 to 800 Da.

1100er Series HPLC from Agilent: quaternary pump, heated column compartment and diode-array detector.

Column: Waters Atlantis dc18; length: 20 mm; internal diameter: 3 mm; particle size: 3 μ m, temperature ($^{\circ}$ C) 40, DAD wavelength range (nm): 200 to 500, solvent gradient: A = 0.1% of formic acid in water

5 and B: 0.1% of formic acid in acetonitrile.

Time (minutes)	A (%)	B (%)	Flow rate (ml/min)
0	80	20	1.7
2.5	0	100	1.7
2.8	0	100	1.7
2.9	80	20	1.7

Method E:

ZMD Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Ionisation method: Electrospray

10 Polarity: positive ions

Capillary (kV) 3.00, Cone (V) 30.00, Extractor (V) 3.00, Source Temperature ($^{\circ}$ C) 150, Desolvation Temperature ($^{\circ}$ C) 320, Cone Gas Flow (L/Hr) 50, Desolvation Gas Flow (L/Hr) 400

Mass range: 150 to 800 Da

Column: Waters Atlantis dc18; length: 20 mm; internal diameter: 3 mm; particle size: 3 μ m, temperature

15 ($^{\circ}$ C) 40, DAD wavelength range (nm): 200 to 500,

Waters Alliance 2795 LC with the following HPLC gradient conditions (Solvent A: 0.1% of formic acid in water and Solvent B: 0.1% of formic acid in acetonitrile)

Time (minutes)	A (%)	B (%)	Flow rate (ml/min)
0	80	20	1.7
2.5	0	100	1.7
2.8	0	100	1.7
2.9	80	20	1.7

Method e:

20 ZQ Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Instrument Parameter:

Ionisation method: Electrospray

Polarity: positive ions

Capillary (kV) 3.00, Cone (V) 30.00, Extractor (V) 2.00, Source Temperature ($^{\circ}$ C) 100, Desolvation

25 Temperature ($^{\circ}$ C) 250, Cone Gas Flow (L/Hr) 50, Desolvation Gas Flow (L/Hr) 400

Mass range: 150 to 1000 Da

HP 1100 HPLC from Agilent: solvent degasser, quaternary pump (ZCQ) / binary pump (ZDQ), heated column compartment and diode-array detector.

Column: Phenomenex Gemini C18, 3 μ m, 30 x 3 mm,

Temp: 60 °C

DAD Wavelength range (nm): 200 to 500

Solvent Gradient:

5 A = water + 0.05 % HCOOH

B= Acetonitril/Methanol (4:1, v:v) + 0.04 % HCOOH

	Time	A%	B%	Flow (ml/min)
	0.00	95.0	5.0	1.700
	2.00	0.0	100.0	1.700
10	2.80	0.0	100.0	1.700
	2.90	95.0	5.0	1.700
	3.00	95.0	5.0	1.700

Method h:

15 ZMD Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Instrument Parameter:

Ionisation method: Electrospray

Polarity: positive (negative) ions

Capillary (kV) 3.80, Cone (V) 30.00, Extractor (V) 3.00, Source Temperature (°C) 150, Desolvation

20 Temperature (°C) 350, Cone Gas Flow (L/Hr) OFF, Desolvation Gas Flow (L/Hr) 600

Mass range: 100 to 900 Da

HP 1100 HPLC from Agilent: solvent degasser, binary pump, heated column compartment and diode-array detector.

Column: Phenomenex Gemini C18, 3 μ m, 30 x 3 mm,

25 Temp: 60 °C

DAD Wavelength range (nm): 200 to 500

Solvent Gradient:

A = water + 0.05 % HCOOH

B= Acetonitril/Methanol (4:1, v:v) + 0.04 % HCOOH

	Time	A%	B%	Flow (ml/min)
	0.00	95.0	5.0	1.700
	2.00	0.0	100.0	1.700
	2.80	0.0	100.0	1.700
	2.90	95.0	5.0	1.700
35	3.00	95.0	5.0	1.700

Method i:

ZQ Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Instrument Parameter:

Ionisation method: Electrospray

Polarity: positive or negative ions

Capillary (kV) 3.00, Cone (V) 30.00 V, Extractor (V) 2.00, Source Temperature (°C) 100, Desolvation

5 Temperature (°C) 250, Cone Gas Flow (L/Hr) 50, Desolvation Gas Flow (L/Hr) 400

Mass range: 100 to 900 Da

HP 1100 HPLC from Agilent: solvent degasser, quaternary pump (ZCQ) / binary pump (ZDQ), heated column compartment and diode-array detector.

Column: Phenomenex Gemini C18, 3 μ m, 30 x 3 mm,

10 Temp: 60 °C

DAD Wavelength range (nm): 200 to 500

Solvent Gradient:

A = water + 0.05 % HCOOH

B= Acetonitril/Methanol (4:1, v:v) + 0.04 % HCOOH

15	Time	A%	B%	Flow (ml/min)
	0.00	95.0	5.0	1.700
	2.00	0.0	100.0	1.700
	2.80	0.0	100.0	1.700
	2.90	95.0	5.0	1.700
20	3.00	95.0	5.0	1.700

Method k:

SQD Mass Spectrometer from Waters (Single quadrupole mass spectrometer)

Instrument Parameter:

25 Ionization method: Electrospray

Polarity: positive and negative ions

Capillary: 3.00 kV

Cone: 45.00 V

Extractor: 2.00 V

30 Source Temperature: 150°C,

Desolvation Temperature: 250°C

Cone Gas Flow: 0 L/Hr

Desolvation Gas Flow: 650 L/Hr

Mass range: 100 to 900 Da

35 Acquity UPLC from Waters:

Binary pump, heated column compartment and diode-array detector.

Solvent degasser, binary pump, heated column compartment and diode-array detector.

Column: Phenomenex Gemini C18, 3 μ m, 30 x 2 mm,

Temp: 60 °C

DAD Wavelength range (nm): 210 to 500

Solvent Gradient:

A = H2O + 5% MeOH + 0.05 % HCOOH

5 B= Acetonitril + 0.05 % HCOOH

Time	A%	B%	Flow (ml/min)
0.00	100	0	0.850
1.2	0	100.0	0.850
1.50	0	100.0	0.850

Method GR

MS	ACQUITY SQD Mass Spectrometer from Waters (Single quadrupole mass spectrometer) Ionisation method: Electrospray Polarity: positive ions Capillary (kV) 3.00, Conc (V) 20.00, Extractor (V) 3.00, Source Temperature (°C) 150, Desolvation Temperature (°C) 400, Cone Gas Flow (L/Hr) 60, Desolvation Gas Flow (L/Hr) 700 Mass range: 100 to 800 Da DAD Wavelength range (nm): 210 to 400																				
LC	Method Waters ACQUITY UPLC with the following HPLC gradient conditions (Solvent A: Water/Methanol 9:1,0.1% formic acid and Solvent B: Acetonitrile,0.1% formic acid) <table><tr><td>Time (min)</td><td>A%</td><td>B%</td><td>Flow (ml/min)</td></tr><tr><td>0</td><td>100</td><td>0</td><td>0.75</td></tr><tr><td>2.5</td><td>0</td><td>100</td><td>0.75</td></tr><tr><td>2.8</td><td>0</td><td>100</td><td>0.75</td></tr><tr><td>3.0</td><td>100</td><td>0</td><td>0.75</td></tr></table>	Time (min)	A%	B%	Flow (ml/min)	0	100	0	0.75	2.5	0	100	0.75	2.8	0	100	0.75	3.0	100	0	0.75
Time (min)	A%	B%	Flow (ml/min)																		
0	100	0	0.75																		
2.5	0	100	0.75																		
2.8	0	100	0.75																		
3.0	100	0	0.75																		

10 Tables A, B and C disclose compounds of the invention. Some reference compounds are also disclosed.

Table A:




Table A discloses compounds of formula (I-a) where R⁵ is methyl, Y¹, Y², Y³ are CH, G¹ is oxygen and R¹ is hydrogen.

	X1	X2	X3	X4	R2	LC/MS Method	retention time (min)	observed mass	Preparation method
A001	C-CF3	CH	C-CF3	CF3	3-Methyl- thietan-3- yl-	E	2.16	570.78	BOP 100°C
A002	C-CF3	CH	C-CF3	CF3	thietan-3- yl-	E	2.06	556.77	BOP 100°C
A003	C-CF3	CH	CH	CF3	3-Methyl- thietan-3- yl-	E	1.98	502.84	BOP 100°C
A004	C-CF3	CH	CH	CF3	thietan-3- yl-	E	1.89	488.84	BOP 100°C
A005	CH	C-CN	CH	CF3	1,1-Dioxo- thietan-3- yl-	B	2.67	ES- 475.89	BOP 100°C
A006	CH	C-CN	CH	CF3	3-Methyl- thietan-3- yl-	B	3.27	ES- 457.83	BOP 100°C
A007	CH	C-CN	CH	CF3	1-oxo- thietan-3- yl-	B	2.43	461.9	BOP 100°C
A008	CH	C-CN	CH	CF3	thietan-3- yl-	B	3.04	445.86	BOP 100°C
A009	CH	C- SCH3	CH	CF3	1,1-Dioxo- thietan-3- yl-	B	2.98	498.87	BOP 100°C
A010	CH	C- SCH3	CH	CF3	3-Methyl- thietan-3- yl-	B	3.53	480.93	BOP 100°C

A011	CH	C-SCH ₃	CH	CF ₃	1-oxo-thietan-3-yl-	B	2.76	482.88	BOP 100°C
A012	CH	C-SCH ₃	CH	CF ₃	thietan-3-yl-	B	3.35	466.9	BOP 100°C
A013	C-Cl	CH	C-Cl	cyclopropyl	1,1-Dioxo-thietan-3-yl-	B	3.29	492.83	BOP 100°C
A014	C-Cl	CH	C-Cl	cyclopropyl	1-oxo-thietan-3-yl-	B	3.01	476.84	BOP 100°C
A015	C-Cl	CH	C-Cl	cyclopropyl	thietan-3-yl-	B	3.65	460.86	BOP 100°C
A016	C-SCH ₃	CH	CH	CF ₃	1,1-Dioxo-thietan-3-yl-	B	2.96	498.9	BOP 100°C
A017	C-SCH ₃	CH	CH	CF ₃	3-Methyl-thietan-3-yl-	B	3.51	480.93	BOP 100°C
A018	C-SCH ₃	CH	CH	CF ₃	1-oxo-thietan-3-yl-	B	2.73	482.89	BOP 100°C
A019	C-SCH ₃	CH	CH	CF ₃	thietan-3-yl-	B	3.32	466.9	BOP 100°C
A020	C-F	C-Cl	CH	CF ₃	1,1-Dioxo-thietan-3-yl-	B	3.11	504.86	BOP 100°C
A021	C-F	C-Cl	CH	CF ₃	3-Methyl-thietan-3-yl-	B	3.64	486.9	BOP 100°C
A022	C-F	C-Cl	CH	CF ₃	1-oxo-thietan-3-yl-	B	2.87	488.85	BOP 100°C
A023	C-F	C-Cl	CH	CF ₃	thietan-3-yl-	B	3.47	472.87	BOP 100°C
A024	C-Cl	CH	C-F	CF ₃	1,1-Dioxo-thietan-3-	B	3.16	504.85	BOP 100°C

					yl-				
A025	C-Cl	CH	C-F	CF3	3-Methyl-thietan-3-yl-	B	3.67	486.9	BOP 100°C
A026	C-Cl	CH	C-F	CF3	1-oxo-thietan-3-yl-	B	2.9	488.86	BOP 100°C
A027	C-Cl	CH	C-F	CF3	thietan-3-yl-	B	3.49	472.87	BOP 100°C
A028	C-Cl	C-F	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	3.09	504.83	BOP 100°C
A029	C-Cl	C-F	CH	CF3	3-Methyl-thietan-3-yl-	B	3.61	486.8	BOP 100°C
A030	C-Cl	C-F	CH	CF3	1-oxo-thietan-3-yl-	B	2.85	488.79	BOP 100°C
A031	C-Cl	C-F	CH	CF3	thietan-3-yl-	B	3.43	472.8	BOP 100°C
A032	CH	C- CF3	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	3.15	520.88	BOP 100°C
A033	CH	C- CF3	CH	CF3	3-Methyl-thietan-3-yl-	B	3.64	502.9	BOP 100°C
A034	CH	C- CF3	CH	CF3	1-oxo-thietan-3-yl-	B	2.93	504.9	BOP 100°C
A035	CH	C- CF3	CH	CF3	thietan-3-yl-	B	3.45	488.84	BOP 100°C
A036	C-Cl	C-Cl	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	3.27	520.84	BOP 100°C
A037	C-Cl	C-Cl	CH	CF3	3-Methyl-thietan-3-yl-	B	3.78	502.82	BOP 100°C

A038	C-Cl	C-Cl	CH	CF3	1-oxo-thietan-3-yl-	B	3.04	504.77	BOP 100°C
A039	C-Cl	C-Cl	CH	CF3	thietan-3-yl-	B	3.59	488.77	BOP 100°C
A040	C-Cl	CH	C-OCH3	CF3	thietan-3-yl-	*			
A041	C-SO2CH3	CH	CH	CF3	thietan-3-yl-	*			
A042	C-F	C-F	C-F	CF3	thietan-3-yl-	*			
A043	C-SOCH3	CH	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	2.08	514.85	BOP 100°C
A044	C-SOCH3	CH	CH	CF3	3-Methyl-thietan-3-yl-	B	2.63	496.91	BOP 100°C
A045	C-SOCH3	CH	CH	CF3	1-oxo-thietan-3-yl-(cis)	B	1.91	498.86	BOP 100°C
A046	C-SOCH3	CH	CH	CF3	thietan-3-yl-	B	2.44	482.88	BOP 100°C
A047	CH	C-Cl	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	3.03	486.85	BOP 100°C
A048	CH	C-Cl	CH	CF3	3-Methyl-thietan-3-yl-	B	3.56	468.9	BOP 100°C
A049	CH	C-Cl	CH	CF3	1-oxo-thietan-3-yl-(cis)	B	2.82	470.87	BOP 100°C
A050	CH	C-Cl	CH	CF3	thietan-3-yl-	B	3.39	454.88	BOP 100°C
A051	C-OCF3	CH	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	3.2	536.88	BOP 100°C

A052	C-OCF ₃	CH	CH	CF ₃	3-Methyl-thietan-3-yl-	B	3.7	518.92	BOP 100°C
A053	C-OCF ₃	CH	CH	CF ₃	1-oxo-thietan-3-yl-(cis)	B	2.98	520.88	BOP 100°C
A054	C-OCF ₃	CH	CH	CF ₃	thietan-3-yl-	B	3.54	504.9	BOP 100°C
A055	CH	C-Br	CH	CF ₃	1,1-Dioxo-thietan-3-yl-	B	3.09	530.79	BOP 100°C
A056	CH	C-Br	CH	CF ₃	3-Methyl-thietan-3-yl-	B	3.6	512.84	BOP 100°C
A057	CH	C-Br	CH	CF ₃	thietan-3-yl-	B	3.44	499.81	BOP 100°C
A058	CH	C-OCF ₃	CH	CF ₃	thietan-3-yl-	B	3.53	504.8	BOP 100°C
A059	CH	C-OCF ₃	CH	CF ₃	1,1-Dioxo-thietan-3-yl-	B	3.2	536.7	BOP 100°C
A060	CH	C-OCF ₃	CH	CF ₃	3-Methyl-thietan-3-yl-	B	3.67	518.5	BOP 100°C
A061	CH	C-OCF ₃	CH	CF ₃	1-oxo-thietan-3-yl-(cis)	B	2.98	520.4	BOP 100°C
A062	C-Cl	CH	C-CF ₃	CF ₃	1,1-Dioxo-thietan-3-yl-	B	3.38	554.9	BOP 100°C
A063	C-Cl	CH	C-CF ₃	CF ₃	3-Methyl-thietan-3-yl-	B	3.87	536.9	BOP 100°C
A064	C-Cl	CH	C-CF ₃	CF ₃	1-oxo-thietan-3-yl-(cis)	B	3.16	538.8	BOP 100°C
A065	C-Cl	CH	C-CF ₃	CF ₃	thietan-3-yl-	B	3.69	522.8	BOP 100°C

A066	C-F	N	CH	CF3	1,1-Dioxo-thietan-3-yl-	B	2.39	472	BOP 100°C
A067	C-F	N	CH	CF3	3-Methyl-thietan-3-yl-	B	2.98	453.9	BOP 100°C
A068	C-F	N	CH	CF3	1-oxo-thietan-3-yl-(cis)	B	2.21	454.9	BOP 100°C
A069	C-F	N	CH	CF3	thietan-3-yl-	B	2.78	439.9	BOP 100°C
A070	N	CH	N	CF3	1,1-Dioxo-thietan-3-yl-	C	1.16	454.97	BOP 100°C
A071	N	CH	N	CF3	3-Methyl-thietan-3-yl-	C	1.83	437.02	BOP 100°C
A072	N	CH	N	CF3	1-oxo-thietan-3-yl-(cis)	C	0.96	439	BOP 100°C
A073	N	CH	N	CF3	thietan-3-yl-	C	1.58	423	BOP 100°C
A074	C-Cl	N	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	C	2.36	521.91	BOP 100°C
A075	C-Cl	N	C-Cl	CF3	3-Methyl-thietan-3-yl-	C	2.88	503.92	BOP 100°C
A076	C-Cl	N	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	C	2.08	505.91	BOP 100°C
A077	C-Cl	N	C-Cl	CF3	thietan-3-yl-	C	2.72	489.92	BOP 100°C
A078	C-Cl	CH	C-Br	CF3	1,1-Dioxo-thietan-3-yl-	C	3.4	566.74	BOP 100°C
A079	C-Cl	CH	C-Br	CF3	3-Methyl-thietan-3-	C	3.9	548.81	BOP 100°C

					yl-				
A080	C-Cl	CH	C-Br	CF3	1-oxo-thietan-3-yl-(cis)	C	3.1	550.77	BOP 100°C
A081	C-Cl	CH	C-Br	CF3	thietan-3-yl-	C	3.7	534.78	BOP 100°C
A082	C-CN	CH	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	C	2.05	511.96	BOP 100°C
A083	C-CN	CH	C-Cl	CF3	3-Methyl-thietan-3-yl-	C	2.49	494.01	BOP 100°C
A084	C-CN	CH	C-Cl	CF3	thietan-3-yl-	C	2.34	479.99	BOP 100°C
A085	C-CN	CH	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	C	1.87	495.99	BOP 100°C
A086	C-3,5-dibromophenyl	CH	C-Br	CF3	thietan-3-yl-	*			
A087	C-3,5-dibromophenyl	CH	C-Br	CF3	1-oxo-thietan-3-yl-	*			
A088	C-Cl	C-Br	C-Cl	CF3	thietan-3-yl-	*			
A089	C-Cl	C-Br	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	*			
A090	C-Cl	C-Br	C-Cl	CF3	1-oxo-thietan-3-yl-	*			
A091	C-3,5-dibromophenyl	CH	C-Br	CF3	1,1-Dioxo-thietan-3-yl-	*			

A092	C-Cl	CH	C-CH ₃	CF ₃	1,1-Dioxo-thietan-3-yl-	B	3.36	501	BOP 100°C
A093	C-Cl	CH	C-CH ₃	CF ₃	3-Methyl-thietan-3-yl-	B	3.94	483	BOP 100°C
A094	C-Cl	CH	C-CH ₃	CF ₃	1-oxo-thietan-3-yl-(cis)	B	3.11	485	BOP 100°C
A095	C-Cl	CH	C-CH ₃	CF ₃	thietan-3-yl-	B	3.72	469	BOP 100°C
A096	C-Cl	CH	C-Cl	CF ₃	thietan-2-ylmethyl-	*			
A097	C-Cl	CH	C-Cl	CF ₃	(1-oxothietan-2-yl)methyl-	e	1.96	519	
A098	C-Cl	CH	C-Cl	CF ₃	(1,1-dioxothieta n-2-yl)methyl	e	1.97	535	
A099	C-CN	CH	C-CN	CF ₃	1,1-Dioxo-thietan-3-yl-	B	2.73	503.1	BOP 100°C
A100	C-CN	CH	C-CN	CF ₃	3-Methyl-thietan-3-yl-	B	3.13	485.1	BOP 100°C
A101	C-CN	CH	C-CN	CF ₃	thietan-3-yl-	B	3.07	471.1	BOP 100°C
A102	C-CN	CH	C-CN	CF ₃	1-oxo-thietan-3-yl-(cis)	B	2.5	487.1	BOP 100°C
A103	C-Cl	CH	C-3-ethoxyp henyl	CF ₃	1-oxo-thietan-3-yl-	D	2.14	591	Suzuki
A104	C-Cl	CH	C-CHCH ₂	CF ₃	1-oxo-thietan-3-yl-	D	1.88	496.97	Suzuki

A105	C-Cl	CH	1H-indol-5-yl	CF3	1-oxo-thietan-3-yl-	D	1.9	585.98	Suzuki
A106	C-Cl	CH	C-4-fluorophenyl	CF3	1-oxo-thietan-3-yl-	D	2.12	564.85	Suzuki
A107	C-Cl	CH	C-4-chlorophenyl	CF3	1-oxo-thietan-3-yl-	D	2.12	580.93	Suzuki
A108	C-Cl	CH	C-3-trifluoromethylphenyl	CF3	1-oxo-thietan-3-yl-	D	2.17	614.96	Suzuki
A109	C-Cl	CH	C-2-chlorophenyl	CF3	1-oxo-thietan-3-yl-	D	2.11	580.92	Suzuki
A110	C-Cl	CH	C-3,5-bis(trifluoromethyl)phenyl	CF3	1-oxo-thietan-3-yl-	D	2.31	682.84	Suzuki
A111	C-Cl	CH	C-CH ₂ CH(CH ₃) ₂	CF3	1-oxo-thietan-3-yl-	D	2.12	526.98	Suzuki
A112	C-Cl	CH	C-3,5-dichlorophenyl	CF3	1-oxo-thietan-3-yl-	D	2.32	614.82	Suzuki
A113	C-Cl	CH	C-4-cyanophenyl	CF3	1-oxo-thietan-3-yl-	D	1.95	571.96	Suzuki
A114	C-Cl	CH	C-3,5-difluorophenyl	CF3	1-oxo-thietan-3-yl-	D	2.12	582.94	Suzuki
A115	C-Cl	CH	2-chloro-4-pyridyl	CF3	1-oxo-thietan-3-yl-	D	1.9	581.9	Suzuki

A116	C-Cl	CH	C-3-chloro-4-methylphenyl	CF3	1-oxo-thietan-3-yl-	D	2.29	594.95	Suzuki
A117	C-Cl	CH	C-3-chloro-2-fluorophenyl	CF3	1-oxo-thietan-3-yl-	D	2.12	598.84	Suzuki
A118	C-Cl	CH	methylimidazol-4-yl	CF3	1-oxo-thietan-3-yl-	D	1.11	550.96	Suzuki
A119	C-Cl	CH	oxazol-5-yl	CF3	1-oxo-thietan-3-yl-	D	1.68	537.93	Suzuki
A120	C-Cl	CH	1,3,5-trimethylpyrazol-4-yl	CF3	1-oxo-thietan-3-yl-	D	1.72	578.96	Suzuki
A121	C-Cl	CH	3-chloro-2-thienyl	CF3	1-oxo-thietan-3-yl-	D	2.11	586.85	Suzuki
A122	C-Cl	C-Cl	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	B	3.8	554.98	BOP 90°C
A123	C-Cl	C-Cl	C-Cl	CF3	3-Methyl-thietan-3-yl-	B	4.34	537	BOP 90°C
A124	C-Cl	C-Cl	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	B	3.61	538.92	BOP 90°C
A125	C-Cl	C-Cl	C-Cl	CF3	thietan-3-yl-	B	4.04	522.98	BOP 90°C
A126	C-Cl	C-CN	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	B	3.5	544.3	BOP 80°C

A127	C-Cl	C-CN	C-Cl	CF3	3-Methyl-thietan-3-yl-	B	4.01	526.3	BOP 80°C
A128	C-Cl	C-CN	C-Cl	CF3	thietan-3-yl-	B	3.83	512.3	BOP 80°C
A129	C-Cl	C-CN	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	B	3.28	528.3	BOP 80°C
A130	C-Cl	CH	C-Cl	CF3	thietan-3-ylmethyl-	*			
A131	C-Cl	CH	C-Cl	CF3	2-(thietan-3-yl)ethanyl	*			
A132	C-Cl	CH	C-Cl	CF3	1,1-Dioxo-thietan-3-ylmethyl	*			
A133	C-Cl	CH	C-Cl	CF3	2-(1,1-dioxothieta n-3-yl)ethanyl	*			
A134	C-Cl	C-OCH 3	C-Cl	CF3	thietan-3-yl-	i	2.09	ES+: 519/521	
A135	C-Cl	C-OCH 3	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	*			
A136	C-Cl	C-I	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	A	1.95	646.45	BOP 80°C
A137	C-Cl	C-I	C-Cl	CF3	3-Methyl-thietan-3-yl-	A	2.19	629.44	BOP 80°C
A138	C-Cl	C-I	C-Cl	CF3	thietan-3-yl-	A	2.11	615.14	BOP 80°C
A139	C-Cl	C-F	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	A	1.85	539.37	BOP 80°C

A140	C-Cl	C-F	C-Cl	CF3	3-Methyl-thietan-3-yl-	A	2.12	520.65	BOP 80°C
A141	C-Cl	C-F	C-Cl	CF3	thietan-3-yl-				
A142	C-Cl	C-SCH3	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	A	1.9	567.29	BOP 80°C
A143	C-Cl	C-SCH3	C-Cl	CF3	3-Methyl-thietan-3-yl-	A	2.17	549.13	BOP 80°C
A144	C-Cl	C-SCH3	C-Cl	CF3	thietan-3-yl-				
A145	C-Cl	C-I	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	A	1.84	630.99	BOP 80°C
A146	C-Cl	C-F	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	A	1.75	523.12	BOP 80°C
A147	C-Cl	C-SCH3	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	A	1.8	551.35	BOP 80°C
A148	C-Cl	C-OH	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	*			
A149	C-Cl	C-SOCH3	C-Cl	CF3	1,1-Dioxo-thietan-3-yl-	A	1.37	582.91	BOP 80°C
A150	C-Cl	C-SOCH3	C-Cl	CF3	3-Methyl-thietan-3-yl-	A	1.67	565.12	BOP 80°C
A151	C-Cl	C-SOCH3	C-Cl	CF3	1-oxo-thietan-3-yl-(cis)	A	1.28	567.09	BOP 80°C
A152	C-Cl	C-SOCH3	C-Cl	CF3	1-oxo-thietan-3-yl-	A	1.57	551.11	BOP 80°C

A153	C-Cl	C-SO ₂ CH ₃	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	A	1.46	598.93	BOP 80°C
A154	C-Cl	C-SO ₂ CH ₃	C-Cl	CF ₃	3-Methyl-thietan-3-yl-	A	1.75	581.52	BOP 80°C
A155	C-Cl	C-SO ₂ CH ₃	C-Cl	CF ₃	1-oxo-thietan-3-yl-(cis)	A	1.37	583.08	BOP 80°C
A156	C-Cl	C-SO ₂ CH ₃	C-Cl	CF ₃	thietan-3-yl-	A	1.65	567.08	BOP 80°C
A157	C-Cl	C-Cl	C-CF ₃	CF ₃	1,1-Dioxo-thietan-3-yl-	A	1.95	589.49	BOP 80°C
A158	C-Cl	C-Cl	C-CF ₃	CF ₃	3-Methyl-thietan-3-yl-	A	2.19	570.65	BOP 80°C
A159	C-Cl	C-Cl	C-CF ₃	CF ₃	1-oxo-thietan-3-yl-(cis)	A	1.86	573.14	BOP 80°C
A160	C-Cl	C-Cl	C-CF ₃	CF ₃	thietan-3-yl-	A	2.11	557.39	BOP 80°C
A161	C-Cl	C-OCOCH ₃	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	*			
A162	C-Cl	C-OSO ₂ CH ₃	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	*			
A163	C-Cl	C-OSO ₂ -4-methylphenyl	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	*			
A164	C-Cl	C-OSO ₂ -2-	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	*			

		nitro-phenyl							
A165	C-Cl	C-OSO ₂ CF ₃	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	h	2.13	ES+: 669/671	
A166	C-CF ₃	C-Cl	C-CF ₃	CF ₃	1,1-Dioxo-thietan-3-yl-	A	1.96	623.35	BOP 80°C
A167	C-CF ₃	C-Cl	C-CF ₃	CF ₃	3-Methyl-thietan-3-yl-	A	2.18	605.18	BOP 80°C
A168	C-CF ₃	C-Cl	C-CF ₃	CF ₃	1-oxo-thietan-3-yl-(cis)	A	1.88	607.24	BOP 80°C
A169	C-CF ₃	C-Cl	C-CF ₃	CF ₃	thietan-3-yl-	A	2.11	591.22	BOP 80°C
A170	C-Cl	C-SCH ₂ CF ₃	C-Cl	CF ₃	1,1-Dioxo-thietan-3-yl-	A	1.9	635.1	BOP 80°C
A171	C-Cl	C-SCH ₂ CF ₃	C-Cl	CF ₃	3-Methyl-thietan-3-yl-	A	2.14	616.99	BOP 80°C
A172	C-Cl	C-SCH ₂ CF ₃	C-Cl	CF ₃	1-oxo-thietan-3-yl-(cis)	A	1.82	619.2	BOP 80°C
A173	C-Cl	C-SCH ₂ CF ₃	C-Cl	CF ₃	thietan-3-yl-	A	2.06	603.26	BOP 80°C
A174	C-Cl	C-SCH ₂ CF ₃	C-SCH ₂ CF ₃	CF ₃	1-oxo-thietan-3-yl-	*			
A175	C-Cl	C-SOCH ₂ CF ₃	C-Cl	CF ₃	1-oxo-thietan-3-yl-	*			
A176	C-Cl	C-SCF ₃	C-Cl	CF ₃	1-oxo-thietan-3-	*			

					yl-				
A177	C-Cl	C-SCH2 Cl	C-Cl	CF3	1-oxo- thietan-3- yl-	*			
A178	C-Cl	CH	C-Cl	CF3	2-(1- oxothietan -3- yl)ethanyl	i	2.04	ES+: 533/534	
A179	C-Cl	C-SCH2 Cl	C-Cl	CF3	1,1-Dioxo- thietan-3- yl-	*			
A180	C-Cl	C-SCF3	C-Cl	CF3	1,1-Dioxo- thietan-3- yl-	*			
A181	C-Cl	C-SOC H2Cl	C-Cl	CF3	1,1-Dioxo- thietan-3- yl-	*			
A182	C-Cl	C-SO2C H2Cl	C-Cl	CF3	1,1-Dioxo- thietan-3- yl-	*			
A183	C-Cl	C-SOCF 3	C-Cl	CF3	1,1-Dioxo- thietan-3- yl-	*			
A184	C-Cl	CH	C-Cl	CF3	2-(3- methoxyth ietan-3- yl)ethyl	i	2.22	ES+: 547/549	
A185	C-Cl	oxazo l-5-yl-	C-Cl	CF3	1-oxo- thietan-3- yl-	*			
A186	C-Cl	C-4 chloro pheny l	C-Cl	CF3	1-oxo- thietan-3- yl-	*			
A187	C-Cl	C-CHC H2	C-Cl	CF3	1-oxo- thietan-3- yl-	*			

A188	C-Cl	CH	C-Cl	CF ₂ C 1	1,1-Dioxo- thietan-3- yl-	A	1.85	536.88	BOP 60°C
A189	C-Cl	CH	C-Cl	CF ₂ C 1	3-Methyl- thietan-3- yl-	A	2.11	519.2	BOP 60°C
A190	C-Cl	CH	C-Cl	CF ₂ C 1	1-oxo- thietan-3- yl-(cis)	A	1.75	521.19	BOP 60°C
A191	C-Cl	CH	C-Cl	CF ₂ C 1	thietan-3- yl-	A	2.02	505.14	BOP 60°C
A192	C-Cl	CH	C-Cl	CF ₃	1-oxo- thietan-3- ylmethyl-	k	1.02	ES-: 519/521	
A193	C-Cl	CH	C-Cl	CF ₂ H	1,1-Dioxo- thietan-3- yl-	k	1.01	ES-: 501/503	
A194	C-Cl	CH	C-Cl	CH ₃	1- Cyanoimin o-1- thietan-3- yl	e	2.16	529	
A195	C-Cl	CH	C-Cl	CH ₃	3-tert- butyl azetidine- 1- carboxylat e	B, *	2.23 min	ES-: 570/572	
A196	C-Cl	CH	C-Cl	CH ₃	azetidin-3-	B, *	1.43 min	ES-: 470/472	
A197	C-Cl	CH	C-Cl	CH ₃	1- methylazet idin-3-	B, *	1.43 min	ES-: 484/486	
A198	C-Cl	CH	C-Cl	CH ₃	1-methyl- 1-oxido- azetidin-1- ium-3-	B	1.53 min	ES-: 500/502	

A199	C-Cl	CH	C-Cl	CH3	1-Cyanoimin o-1-oxo-1- thietan-3- yl	e	2.02	545	
A200	C-Cl	CH	C-Cl	CH3	(1,1-Dioxo-1- thietan-3- ylcarbamo yl)-methyl	E	2.46	577.85	BOP-
A201	C-Cl	CH	C-Cl	CH3	(3-Methyl- thietan-3- ylcarbamo yl)-methyl	E	2.9	559.97	BOP-
A202	C-Cl	CH	C-Cl	CH3	(1-oxo-1- thietan-3- ylcarbamo yl)-methyl	E	2.23	561.85	BOP-
A203	C-Cl	CH	C-Cl	CH3	(1-thietan- 3- ylcarbamo yl)-methyl	E	2.7	545.86	BOP-
A204	C-Cl	CH	C-Cl	CH3	1-Oxo-1- (pyridin-2- ylimino)- 1-thietan- 3-yl	e	1.92	597	BOP
A205	C-Cl	CH	C-Cl	CH3	3-(azetidin- 1-yl- phenyl- methanone)	B, *	2.09 min	ES: 574/576	
A206	C-Cl	CH	C-Cl	CH3	3-(methyl azetidine- 1- carboxylat e)	B, *	2.04 min	ES: 528/530	

A207	C-Cl	CH	C-Cl	CH3	3-Hydroxyimino-cyclobutyl	i	1.97	500	
A208	C-Cl	CH	C-Cl	CH3	3-Oxo-cyclobutyl	i	2.02	485	
A209	C-Cl	CH	C-Cl	CH3	1-(1,1-Dioxo-1-thietan-3-ylcarbamo-yl)-ethyl-	B	3.51	591.9	BOP-
A210	C-Cl	CH	C-Cl	CH3	1-(1-Oxo-1-thietan-3-ylcarbamo-yl)-ethyl-	B	3.3	575.9	BOP-
A211	C-Cl	CH	C-Cl	CH3	1-(1-Thietan-3-ylcarbamo-yl)-ethyl-	B	3.83	559.9	BOP-
A212	C-Cl	CH	C-Cl	CH3	1-(1,1-Dioxo-1-thietan-3-ylcarbamo-yl)-2-phenyl-ethyl	B	3.85	667.8	BOP-
A213	C-Cl	CH	C-Cl	CH3	1-(3-Methyl-thietan-3-ylcarbamo-yl)-2-phenyl-ethyl	B	4.35	649.9	BOP-
A214	C-Cl	CH	C-Cl	CH3	1-(1-Oxo-1-thietan-3-ylcarbamo-yl)-ethyl-	B	3.71	651.9	BOP-

					yl)-2-phenyl-ethyl				
A215	C-Cl	CH	C-Cl	CH3	1-(1-Thietan-3-ylcarbamo-yl)-2-phenyl-ethyl	B	4.21	635.9	BOP-
A216	C-Cl	CH	C-Cl	CH3	3-Methoxyimino-cyclobutyl	i	2.08	514	
A217	C-Cl	CH	C-Cl	CH3	3-Benzoyloxyimino-cyclobutyl	i	2.33	590	
A218	C-Cl	CH	C-Cl	CH3	3-(2,2,2-Trifluoroethoxyimino)-cyclobutyl	i	2.21	582	
A219	C-Cl	CH	C-Cl	CH3	1-Benzoyloxy-2-oxo-azetidin-3-yl	i	2.21	592	
A220	C-Cl	CH	C-Cl	CH3	3-Ethoxyimino-cyclobutyl	k	1.16	528	

*: NMR data as follows: ¹H-NMR (400 MHz, CDCl₃, in ppm)

A040: 2.47 (s, 3H), 3.40 (t, 2H), 3.50 (t, 2H), 3.72 (d, 1H), 3.85 (s, 3H), 4.06 (d, 1H), 5.43 (m, 1H), 6.19 (d, 1H), 6.95 (m, 1H), 7.08 (s, 1H), 7.17 (s, 1H), 7.41 (d, 1H), 7.53 (m, 2H).

A041: 2.48 (s, 3H), 3.11 (s, 3H), 3.40 (t, 2H), 3.51 (t, 2H), 3.78 (d, 1H), 4.15 (d, 1H), 5.43 (m, 1H), 6.19 (d, 1H), 7.43 (d, 1H), 7.54 (m, 2H), 7.71 (t, 1H), 7.98 (d, 1H), 8.04 (d, 1H), 8.18 (s, 1H).

A042: 2.48 (s, 3H), 3.40 (t, 2H), 3.50 (t, 2H), 3.66 (d, 1H), 4.08 (d, 1H), 5.43 (m, 1H), 6.19 (d, 1H), 7.32 (d, 1H), 7.42 (d, 1H), 7.43 (s, 1H), 7.52 (m, 2H).

- A086: 2.48 (s, 3H), 3.40 (t, 2H), 3.48 (t, 2H), 3.77 (d, 1H), 4.14 (d, 1H), 5.43 (m, 1H), 6.18 (d, 1H), 7.42(d, 1H), 7.55 (m, 2H), 7.64 (s, 2H), 7.70 (s, 1H), 7.72 (d, 2H), 7.78 (s, 1H).
- A087: 2.46 (s, 3H), 3.31 (m, 2H), 3.78 (d, 1H), 4.14 (d, 1H), 4.19 (m, 2H), 4.69 (m, 1H), 6.63 (d, 1H), 7.43 (d, 1H), 7.54 (s, 2H), 7.63 (s, 2H), 7.69 (s, 1H), 7.73 (d, 2H), 7.77 (s, 1H).
- 5 A088: 2.44 (s, 3H), 3.39 (t, 2H), 3.46 (t, 2H), 3.70 (d, 1H), 4.09 (d, 1H), 5.39 (m, 1H), 6.41 (d, 1H), 7.38 (d, 1H), 7.49 (m, 2H), 7.63 (s, 2H).
- A089: 2.49 (s, 3H), 3.70 (d, 1H), 4.05 (m, 2H), 4.09 (d, 1H), 4.63 (m, 2H), 4.90 (m, 1H), 6.53 (d, 1H), 7.45 (d, 1H), 7.54 (d, 1H), 7.55 (s, 1H), 7.64 (s, 2H).
- A090: 2.48 (s, 3H), 3.23 (m, 2H), 3.70 (d, 1H), 4.09 (d, 1H), 4.22 (m, 2H), 4.69 (m, 1H), 6.25 (d, 1H), 10 7.44 (d, 1H), 7.54 (m, 2H), 7.63 (s, 2H).
- A091: 2.50 (s, 3H), 3.78 (d, 1H), 4.04 (m, 2H), 4.15 (d, 1H), 4.64 (m, 2H), 4.90 (m, 1H), 6.51 (d, 1H), 7.46 (d, 1H), 7.56 (d, 1H), 7.58 (s, 1H), 7.64 (s, 2H), 7.70 (s, 1H), 7.74 (d, 2H), 7.78 (s, 1H).
- A096: 2.50 (s, 3H), 2.74 (m, 1H), 3.05 (m, 2H), 3.22 (m, 1H), 3.73 (m, 3H), 4.00 (m, 1H), 4.09 (d, 1H), 6.27 (m, 1H), 7.44 (m, 2H), 7.53 (m, 4H).
- 15 A130: 7.55 (m, 4H), 7.50-7.40(m, 2H), 6.0(t br, 1H), 4.15-4.10(d, 1H), 3.75-3.70(d, 1H), 3.65-3.6(m, 2H CH2), 3.55-3.5(m, 1H), 3.40-3.30(t, 2H), 3.10-3.05(q, 2H), 2.50 (s, 3H)
- A131: 7.55(m, 4H), 7.50-7.40(m, 2H), 5.80(t br, 1H), 4.10(d, 2H), 3.40(m, 3H), 3.25(t, 2H), 3.10(t, 2H), 2.50 (s, 3H), 1.90(q, 2H),
- A132: 7.55-7.50(m, 4H), 7.45-7.40(m, 2H), 6.20 (t br, 1H), 4.30-4.25 (m, 2H), 4.10-4.05(d, 1H), 4.0-3.9 20 (m, 2H), 3.75-3.65 (m, 3H), 3.0 (m, 1H), 2.50 (s, 3H)
- A133: 7.55-7.50(m, 4H), 7.45-7.40(m, 1H), 7.35(m, 1H), 5.9(t br, 1H), 4.30-4.25(m, 2H), 4.10-4.05(d, 1H), 3.9-3.8(m, 2H), 3.75-3.65(d, 1H), 3.5-3.4(q, 2H), 2.6(m, 1H), 2.45 (s, 3H), 2.0 (q, 2H).
- A135: 7.56 (m, 4H), 7.45 (d, 1H), 6.49 (d, 1H), 4.9 (m, 1H), 4.64 (m, 2H), 4.06 (m, 3H), 3.94 (s, 3H), 3.70 (d, 1H), 2.49 (s, 3H).
- 25 A148: 7.56 (m, 4H), 7.46 (d, 1H), 6.44 (d, 1H), 5.99 (m, 1H), 4.92 (m, 1H), 4.64 (m, 2H), 4.06 (m, 3H), 3.70 (d, 1H), 2.50 (s, 3H).
- A161: 7.63 (s, 2H), 7.48 (m, 2H), 7.39 (d, 1H), 6.93 (d, 1H), 4.82 (m, 1H), 4.56 (m, 2H), 4.08 (m, 3H), 3.73 (d, 1H), 2.42 (s, 3H), 2.41 (s, 3H)
- A162: 7.68 (s, 2H), 7.55 (m, 2H), 7.47 (d, 1H), 6.58 (d, 1H), 4.91 (m, 1H), 4.63 (m, 2H), 4.05 (m, 3H), 30 3.72 (d, 1H), 3.50 (s, 3H), 2.49 (s, 3H)
- A163: 7.91 (d, 2H), 7.61 (s, 2H), 7.55 (m, 2H), 7.38 (m, 3H), 6.84 (d, 1H), 4.84 (m, 1H), 4.59 (m, 2H), 4.09 (m, 3H), 3.73 (d, 1H), 2.49 (s, 3H), 2.45 (s, 3H)
- A164: 8.16 (d, 1H), 7.94 (m, 2H), 7.90 (m, 1H), 7.63 (s, 2H), 7.50 (m, 2H), 7.42 (d, 1H), 6.78 (d, 1H), 4.85 (m, 1H), 4.59 (m, 2H), 4.11 (m, 3H), 3.75 (d, 1H), 2.45 (s, 3H).
- 35 A174: 2.48 (s, 3H), 3.25 (m, 2H), 3.49 (q, 2H), 3.62 (q, 2H), 3.69 (d, 1H), 4.11 (d, 1H), 4.22 (m, 2H), 4.69 (m, 1H), 6.30 (d, 1H), 7.44 (d, 1H), 7.52 (m, 2H), 7.55 (s, 1H), 7.58 (s, 1H).
- A175: 2.48 (s, 3H), 3.24 (m, 2H), 3.69 (dd, 1H), 3.94 (m, 1H), 4.13 (d, 1H), 4.21 (m, 1H), 4.21 (m, 2H), 4.72 (m, 1H), 6.46 (d, 1H), 7.45 (d, 1H), 7.54 (m, 2H), 7.68 (s, 2H).

- A176: 2.48 (s, 3H), 3.26 (m, 2H), 3.73 (d, 1H), 4.12 (d, 1H), 4.22 (m, 2H), 4.70 (m, 1H), 6.35 (d, 1H), 7.44 (d, 1H), 7.54 (m, 2H), 7.76 (s, 2H).
- A177: 2.48 (s, 3H), 3.25 (m, 2H), 3.73 (d, 1H), 4.11 (d, 1H), 4.21 (m, 2H), 4.61 (m, 1H), 4.98 (s, 2H), 6.33 (d, 1H), 7.44 (d, 1H), 7.55 (m, 2H), 7.70 (s, 2H).
- 5 A179: 2.50 (s, 3H), 3.73 (d, 1H), 4.03 (m, 2H), 4.11 (d, 1H), 4.64 (m, 2H), 4.91 (m, 1H), 4.98 (s, 2H), 6.45 (d, 1H), 7.47 (d, 1H), 7.56 (d, 1H), 7.57 (s, 1H), 7.71 (s, 2H).
- A180: 2.49 (s, 3H), 3.73 (d, 1H), 4.03 (m, 2H), 4.13 (d, 1H), 4.64 (m, 2H), 4.91 (m, 1H), 6.50 (d, 1H), 7.46 (d, 1H), 7.55 (d, 1H), 7.56 (s, 1H), 7.76 (s, 2H).
- A181: 2.49 (s, 3H), 3.73 (dd, 1H), 4.04 (m, 2H), 4.14 (d, 1H), 4.63 (m, 2H), 4.87 (d, 1H), 4.90 (m, 1H), 10 5.05 (d, 1H), 6.59 (d, 1H), 7.46 (d, 1H), 7.54 (d, 1H), 7.55 (s, 1H), 7.67 (s, 2H).
- A182: 2.50 (s, 3H), 3.73 (d, 1H), 4.03 (m, 2H), 4.15 (d, 1H), 4.65 (m, 2H), 4.90 (m, 1H), 4.92 (s, 2H), 6.48 (d, 1H), 7.47 (d, 1H), 7.55 (d, 1H), 7.56 (s, 1H), 7.78 (s, 2H).
- A183: 2.50 (s, 3H), 3.73 (d, 1H), 4.04 (m, 2H), 4.16 (d, 1H), 4.64 (m, 2H), 4.90 (m, 1H), 6.47 (d, 1H), 7.47 (d, 1H), 7.56 (d, 2H), 7.83 (s, 2H).
- 15 A185: 2.49 (s, 3H), 3.24 (m, 2H), 3.74 (d, 1H), 4.13 (d, 1H), 4.22 (m, 2H), 4.71 (m, 1H), 6.39 (d, 1H), 7.35 (s, 1H), 7.45 (d, 1H), 7.56 (m, 2H), 7.70 (s, 2H), 8.07 (s, 1H).
- A186: 2.49 (s, 3H), 3.23 (m, 2H), 3.76 (d, 1H), 4.13 (d, 1H), 4.22 (m, 2H), 4.70 (m, 1H), 6.28 (d, 1H), 7.20 (d, 2H), 7.46 (m, 3H), 7.57 (m, 2H), 7.68 (s, 2H).
- A187: 2.47 (s, 3H), 3.28 (m, 2H), 3.71 (d, 1H), 4.08 (d, 1H), 4.20 (m, 2H), 4.70 (m, 1H), 5.81 (dd, 2H), 20 6.51 (d, 1H), 6.70 (dd, 1H), 7.41 (d, 1H), 7.54 (m, 2H), 7.58 (s, 2H).
- A195: 1.19 (s, 9H), 2.16 (s, 3H), 3.50 (d, 1H), 3.58 (m, 2H), 3.85 (m, 1H), 4.02 (dt, 2H), 4.50 (m, 1H), 6.75 (d, 1H), 7.10-7.30 (m, 6H).
- A196: 2.30 (s, 3H), 3.46 (br t, 2H), 3.52 (d, 1H), 3.91 (t, 2H), 3.98 (d, 1H), 4.80 (m, 1H), 7.00 (d, 1H), 7.30-7.45 (m, 6H).
- 25 A197: 2.30 (s, 3H), 2.40 (s, 3H), 3.05 (m, 2H), 3.60 (t, 2H), 3.65 (d, 1H), 4.01 (d, 1H), 4.51 (m, 1H), 6.38 (m, 1H), 7.30-7.50 (m, 6H).
- A205: 2.45 (s, 3H), 3.69 (d, 1H), 4.05 (d, 1H), 4.15 (m, 2H), 4.55 (m, 2H), 4.90 (m, 1H), 6.51 (d, 1H), 7.30-7.60 (m, 11H).
- A206: 2.45 (s, 3H), 3.69 (s, 3H), 3.70 (d, 1H), 3.90 (m, 2H), 4.07 (d, 1H), 4.40 (t, 2H), 4.72 (m, 1H), 30 6.20 (d, 1H), 7.40-7.60 (m, 6H).

Table B:

Table B discloses compounds of formula (I-a) where X^1 and X^3 are chloro, X^2 is CH, X^4 is trifluoromethyl, G^1 is oxygen and R^1 is hydrogen.

35

	R2	Y1	Y2	Y3	R5	LC/MS Metho d	retentio n time (min)	observed mass	Preparatio n method
--	----	----	----	----	----	---------------------	-----------------------------	------------------	------------------------

B01	3-Methyl-thietan-3-yl-	N	N	H	CF3	E	2.08	ES-556.72	BOP 100°C
B02	thietan-3-yl-	N	N	H	CF3	E	1.99	ES-542.75	BOP 100°C
B03	3-Methyl-thietan-3-yl-	CH	CH	CH	CF3	E	2.16	ES-554.8	BOP 100°C
B04	thietan-3-yl-	CH	CH	CH	CF3	E	2.08	ES-540.8	BOP 100°C
B05	3-Methyl-thietan-3-yl-	CH	CH	CH	Br	E	2.09	566.81	BOP 100°C
B06	thietan-3-yl-	CH	CH	CH	Br	*			
B07	3-Methyl-thietan-3-yl-	CH	CH	CH	CN	E	2.13	514.24	BOP 100°C
B08	thietan-3-yl-	CH	CH	CH	CN	E	2.2	500.72	BOP 100°C
B09	thietan-3-yl-	CH	CH	C-Br	CH3	*			
B10	thietan-3-yl-	CH	CH	C-I	CH3	*			
B11	thietan-3-yl-	CH	C-Br	CH	CH3	*			
B12	thietan-3-yl-	CH	C-Cl	CH	CH3	*			
B13	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	F	B	3.11	524.84	BOP 80°C
B14	3-Methyl-thietan-3-yl-	CH	CH	CH	F	B	3.61	506.88	BOP 80°C
B15	1-oxo-thietan-3-yl-	CH	CH	CH	F	B	2.92	508.86	BOP 80°C
B16	thietan-3-yl-	CH	CH	CH	F	*			
B17	1,1-Dioxo-thietan-3-yl-	CH	CH	CH3	CH3	B	3.38	534.84	BOP 100°C
B18	3-Methyl-thietan-3-yl-	CH	CH	CH3	CH3	B	3.91	516.86	BOP 100°C
B19	1-oxo-thietan-3-yl-(cis)	CH	CH	CH3	CH3	B	3.13	518.86	BOP 100°C

B20	thietan-3-yl-	CH	CH	CH3	CH3	B	3.74	502.88	BOP 100°C
B21	1,1-Dioxo-thietan-3-yl-	C-F	CH	CH	H	B	3.35	ES-522.82	BOP 100°C
B22	3-Methyl-thietan-3-yl-	C-F	CH	CH	H	B	3.94	506.78	BOP 100°C
B23	1-oxo-thietan-3-yl-	C-F	CH	CH	H	B	3.14	508.74	BOP 100°C
B24	thietan-3-yl-	C-F	CH	CH	H	B	3.74	492.76	BOP 100°C
B25	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	OCH ₃	B	3.43	536.76	BOP 100°C
B26	3-Methyl-thietan-3-yl-	CH	CH	CH	OCH ₃	B	4.04	518.8	BOP 100°C
B27	1-oxo-thietan-3-yl-	CH	CH	CH	OCH ₃	B	3.17	520.76	BOP 100°C
B28	thietan-3-yl-	CH	CH	CH	OCH ₃	B	3.84	504.77	BOP 100°C
B29	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	Cl	B	3.37	540.82	BOP 100°C
B30	3-Methyl-thietan-3-yl-	CH	CH	CH	Cl	B	3.89	522.79	BOP 100°C
B31	1-oxo-thietan-3-yl-	CH	CH	CH	Cl	B	3.13	524.74	BOP 100°C
B32	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	cyclopropyl	B	3.8	546.82	BOP 100°C
B33	3-Methyl-thietan-3-yl-	CH	CH	CH	cyclopropyl	B	4.2	528.87	BOP 100°C
B34	1-oxo-thietan-3-yl-(cis)	CH	CH	CH	cyclopropyl	B	3.39	530.84	BOP 100°C
B35	thietan-3-yl-	CH	CH	CH	cyclopropyl	B	3.99	514.86	BOP 100°C
B36	1,1-Dioxo-thietan-3-yl-	CH	CH	N	CH3	B	3.55	521.79	BOP 100°C

B37	3-Methyl-thietan-3-yl-	CH	CH	N	CH3	B	4.32	503.85	BOP 100°C
B38	thietan-3-yl-	CH	CH	N	CH3	B	4.05	489.83	BOP 100°C
B39	1-oxo-thietan-3-yl-(cis)	CH	CH	N	CH3	B	3.23	505.81	BOP 100°C
B40	1,1-Dioxo-thietan-3-yl-	CH	CH	N	H	B	3.45	507.8	BOP 100°C
B41	3-Methyl-thietan-3-yl-	CH	CH	N	H	B	4.02	489.88	BOP 100°C
B42	thietan-3-yl-	CH	CH	N	H	B	3.84	475.82	BOP 100°C
B43	1-oxo-thietan-3-yl-(trans)	CH	CH	N	H	B	3.1	491.79	BOP 100°C
B44	1,1-Dioxo-thietan-3-yl-	CH	N	CH	CH3	B	3.33	521.92	BOP 100°C
B45	3-Methyl-thietan-3-yl-	CH	N	CH	CH3	B	3.9	503.92	BOP 100°C
B46	thietan-3-yl-	CH	N	CH	CH3	B	3.68	489.93	BOP 100°C
B47	1,1-Dioxo-thietan-3-yl-	N	CH	CH	CH3	B	3.36	521.92	BOP 100°C
B48	3-Methyl-thietan-3-yl-	N	CH	CH	CH3	B	3.96	503.98	BOP 100°C
B49	thietan-3-yl-	N	CH	CH	CH3	B	3.77	489.96	BOP 100°C
B50	1-oxo-thietan-3-yl-(cis)	CH	N	CH	CH3	B	3.09	505.91	BOP 100°C
B51	1-oxo-thietan-3-yl-(cis)	N	CH	CH	CH3	B	3.13	505.94	BOP 100°C
B52	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	COC H3	B	3.39	549	BOP 100°C
B53	1-oxo-thietan-3-yl-(cis)	CH	CH	CH	COC H3	B	3.16	533	BOP 100°C
B54	thietan-3-yl-	CH	CH	CH	COC H3	B	3.77	517	BOP 100°C
B55	thietan-3-yl-	CH	NO2	CH	H	*			

B56	thietan-3-yl-	CH	CH	CH	NO2	*			
B57	thietan-3-yl-	CH	CH	CH	NH2	*			
B58	1-oxo-thietan-3-yl-(cis)	CH	CH	CH	CF3	e	2	557	
B59	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	H	A	1.79	507.46	BOP 80°C
B60	3-Methyl-thietan-3-yl-	CH	CH	CH	H	A	2.05	489.34	BOP 80°C
B61	thietan-3-yl-	CH	CH	CH	H				
B62	1-oxo-thietan-3-yl-(cis)	CH	CH	CH	H	A	1.69	491.27	BOP 80°C
B63	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	CH2S CH3	*			
B64	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	CH2 OCO C6H5	*			
B65	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	CH2S O2C H3	*			
B66	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	CH2S OCH 3	*			
B67	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	CH2 OCO CH3	*			
B68	1,1-Dioxo-thietan-3-yl-	CH	CH	CH	CH2 OH	*			
B69	thietan-3-yl-	CH	CH	CH	CH2C H3	*			

*: NMR data as follows: ¹H-NMR (400 MHz, CDCl₃, in ppm)

B06: 7.9 (s, 1H), 7.65 (m, 1H), 7.60 (m, 1H), 7.5 (m, 2H), 7.45 (m, 1H), 6.5 (dd br, 1H), 5.4 (m, 1H), 4.1-4.05 (d, 1H), 3.75-3.65 (d, 1H), 3.5-3.4 (m, 4H).

B09: 7.6 (s, 1H), 7.45-7.35 (m, 4H), 6.15 (dd br, 1H), 5.4-5.3 (m, 1H), 3.9 (d, 1H), 3.6 (d, 1H), 3.4-3.3 (m, 4H), 2.30 (s, 3H)

B10: 7.8 (s, 1H), 7.42 (m, 3H), 7.35 (m, 1H), 6.1 (dd br, 1H), 5.4-5.3 (m, 1H), 3.9 (d, 1H), 3.6 (d, 1H), 3.4 (t, 2H), 3.3 (d, 2H), 2.30 (s, 3H).

5 **B11:** 7.6 (s, 1H), 7.5 (m, 2H), 7.45 (m, 1H), 7.4 (m, 1H); 6.3 (dd br, 1H), 5.4 (m, 1H), 4.3-4.3 (d, 1H), 3.9-3.8 (d, 1H), 3.5 (t, 2H), 3.4 (d, 2H), 2.40 (s, 3H).

B12: 7.5 (s, 1H), 7.4 (m, 2H), 7.35 (m, 2H), 6.2-6.1 (dd br, 1H), 5.3 (m, 1H), 4.2-4.15 (d, 1H), 3.7-3.8 (d, 1H), 3.9 (t, 2H), 3.3 (d, 2H), 2.30 (s, 3H).

B16: 8.1 (m, 1H), 7.5-7.35 (m, 4H), 7.0 (m, 1H), 5.45-5.35 (m, 1H), 4.0 (d, 1H), 3.6 (d, 1H), 3.45-3.25 (m, 4H).

B55: 8.47 (1H, d), 8.11-8.13(1H, m), 7.66 (1H, d), 7.45 (3H, d), 6.98 (1H, d), 5.40-5.46 (1H, m), 4.11 (1H, d), 3.98 (1H, d), 3.39-3.48 (4H, m).

B56: 8.2 (1H, d), 8.01 (1H, m), 7.57 (1H, d), 7.45-7.50 (3H, m), 6.58 (1H, d), 5.33-5.40 (1H, m), 4.12 (1H, d), 3.75 (1H, d), 3.38-3.50 (4H, m)

15 **B57:** 7.49 (2H, d), 7.42 (1H, m), 7.34 (1H, d), 6.92-6.94 (2H, m), 6.48 (1H, d) 5.18-5.20 (1H, m), 4.03 (1H, d), 3.64 (1H, d), 3.31-3.41 (4H, m), 0.82 (3H, m).

B63: 7.53 - 7.66 (m, 4 H), 7.51 (m, 2 H), 7.38 - 7.47 (m, 1 H), 4.79 (m, 1H), 4.60 (m, 2 H), 4.04 - 4.20 (m, 3 H), 3.91 (s, 2 H), 3.72 (d, 1 H), 2.10 (s, 3 H).

B64: (DMSO-d₆, 400MHz) 7.91 - 8.04 (m, 3 H), 7.75 - 7.86 (m, 2 H), 7.61 - 7.74 (m, 4 H), 7.46 - 7.59 (m, 2 H), 5.53 (s, 2 H), 4.38 - 4.59 (m, 4 H), 4.28 - 4.38 (m, 1 H), 4.10 - 4.24 (m, 2 H).

B65: 3.22 (s, 3H), 3.72 (d, 1H), 4.08 (m, 3H), 4.59 (m, 3H), 4.68 (s, 2H), 7.45 (d, 1H), 7.51 (d, 2H), 7.73 (d, 1H), 7.77 (d, 1H), 7.85 (d, 1H), 8.11 (m, 1H).

B66: 2.98 (d, 3H), 3.73 (d, 1H), 3.96 (d, 1H), 4.11 (m, 1H), 4.29 (m, 2H), 4.50 (m, 3H), 4.78 (d, 1H), 7.45 (m, 1H), 7.52 (m, 2H), 7.65 (d, 1H), 7.81 (m, 2H), 9.67 (m, 1H).

25 **B67:** 2.14 (s, 3H), 3.72 (d, 1H), 4.13 (m, 3H), 4.60 (m, 2H), 4.82 (m, 1H), 5.28 (s, 2H), 7.45 (d, 1H), 7.52 (d, 2H), 7.63 (d, 1H), 7.68 (d, 1H), 7.81 (m, 2H).

B68: 3.73 (d, 1H), 4.10 (m, 3H), 4.63 (m, 2H), 4.73 (s, 2H), 4.83 (m, 1H), 7.45 (d, 1H), 7.52 (d, 2H), 7.71 (s, 2H), 7.78 (d, 1H), 8.22 (m, 1H).

B69: 1.25 (t, 3H), 2.80 (2H, q), 3.35 (t, 2H), 3.50 (t, 2H), 3.70 (d, 1H), 4.10 (d, 1H), 5.35-5.46 (1H, m), 6.25 (d, 1H), 7.35 (d, 1H), 7.39-7.43 (1H, m), 7.46-7.55 (3H, m), 7.62 (1H, s).

	Chemical name	LC/MS Method	retention time (min)	observed mass	Preparation method
C001	tert-butyl N-[1-[4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzoyl]imino-1-oxo-thietan-3-yl]carbamate	e	2.21	620	

C002	2-methyl-4-[5-(1H-pyrrol-2-yl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(thietan-3-yl)benzamide	*			
C003	4-[5-(4-tert-butyl-1H-pyrrol-2-yl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-yl)benzamide	*			
C004	4-[5-[1-(2,2-difluoroethyl)pyrrol-3-yl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-yl)benzamide	*			
C005	4-[5-[4-tert-butyl-1-(2,2-difluoroethyl)pyrrol-3-yl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-yl)benzamide	*			
C006	2-methyl-4-[5-[1-methyl-5-(trifluoromethyl)pyrrol-3-yl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(thietan-3-yl)benzamide	*			
C007	4-[5-[1-(2,2-difluoroethyl)-5-(trifluoromethyl)pyrrol-3-yl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-yl)benzamide	*			
C008	N-(1,1-dioxothietan-3-yl)-2-methyl-4-[5-(trifluoromethyl)-5-[5-(trifluoromethyl)-1H-pyrrol-3-yl]-4H-isoxazol-3-yl]benzamide	A	1.44	510.42	BOP 80°C
C009	2-methyl-N-(1-oxothietan-3-yl)-4-[5-(trifluoromethyl)-5-[5-(trifluoromethyl)-1H-pyrrol-3-yl]-4H-isoxazol-3-yl]benzamide	A	1.35	494.28	BOP 80°C
C010	2-methyl-N-(thietan-3-yl)-4-[5-(trifluoromethyl)-5-[5-(trifluoromethyl)-1H-pyrrol-3-yl]-4H-isoxazol-3-yl]benzamide	A	1.61	477.72	BOP 80°C
C011	4-[5-(3,5-dichlorophenyl)-2-oxido-5-(trifluoromethyl)-4H-isoxazol-2-ium-3-yl]-2-methyl-N-(thietan-3-yl)benzamide	E, *	2.03 min	ES ⁺ : 505/507 (MH ⁺)	
C012	4-[5-(3,5-dichlorophenyl)-2-oxido-5-(trifluoromethyl)-4H-isoxazol-2-ium-3-yl]-N-(1,1-dioxothietan-3-yl)-2-methyl-benzamide	E, *	1.92 min	ES ⁺ : 537/539 (MH ⁺)	
C013	[4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-phenyl]-(6-thia-2-azaspiro[3.3]heptan-2-yl)methanone	B, *	2.18 min	ES ⁺ : 515/517	

C014	4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[3-(trifluoromethyl)thietan-3-yl]benzamide	*			
C015	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(1-oxidothietan-1-ium-3-yl)benzamide	*See above			
C016	5-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-3-methyl-N-(1-oxothietan-3-yl)pyridine-2-carboxamide	*See above			
C017	2-methyl-N-(1-oxothietan-3-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	*See above			
C018	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(1,1-dioxothietan-3-yl)-2-methyl-benzamide	*See above			
C019	N-(1,1-dioxothietan-3-yl)-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	*See above			
C020	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(1,1-dioxothietan-3-yl)methyl]-2-methyl-benzamide	*			
C021	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(thietan-3-ylmethyl)benzamide	*			

C002: 2.46 (s, 3H); 3.41 (t, 2H); 3.49 (t, 2H); 3.67 (d, 1H); 3.94 (d, 1H); 5.43 (m, 1H); 6.26 (d, 1H); 6.34 (s, 1H); 6.83 (m, 1H); 7.02 (m, 1H); 7.39 (d, 1H); 7.52 (d, 1H); 7.53 (s, 1H); 8.38 (s, 1H).

C003: 1.30 (s, 9H); 2.47 (s, 3H); 3.40 (t, 2H); 3.50 (t, 2H); 3.68 (d, 1H); 3.89 (d, 1H); 5.43 (m, 1H); 6.01 (s, 1H); 6.21 (d, 1H); 6.86 (s, 1H); 7.40 (d, 1H); 7.53 (d, 1H); 7.54 (s, 1H); 8.04 (s, 1H).

C004: 2.46 (s, 3H); 3.41 (t, 2H); 3.49 (t, 2H); 3.65 (d, 1H); 3.92 (d, 1H); 4.20 (td, 2H); 5.42 (m, 1H); 5.93 (tt, 1H); 6.30 (s, 1H); 6.31 (d, 1H); 6.71 (s, 1H); 6.91(s, 1H); 7.39 (d, 1H); 7.50 (d,1H); 7.51 (s, 1H).

C005: 1.36 (s, 9H); 2.47 (s, 3H); 3.40 (t, 2H); 3.50 (t, 2H); 3.65 (d, 1H); 3.88 (d, 1H); 4.37 (td, 2H); 5.43 (m, 1H); 5.93 (tt, 1H); 6.03 (s, 1H); 6.20 (d, 1H); 6.83 (s, 1H); 7.41 (d, 1H); 7.54 (m, 2H).

10 C006: 2.48 (s, 3H); 3.41 (t, 2H); 3.50 (t, 2H); 3.62 (d, 1H); 3.72 (s, 3H); 3.95 (d, 1H); 5.45 (m, 1H); 6.22 (m, 1H); 6.67 (s, 1H); 6.95 (s, 1H); 7.42 (d, 1H); 7.52 (m, 2H).

C007: 2.48 (s, 3H); 3.41 (t, 2H); 3.50 (t, 2H); 3.62 (d, 1H); 3.95 (d, 1H); 4.35 (m, 2H); 5.45 (m, 1H); 5.95 (tt, 1H); 6.22 (m, 1H); 6.78 (s, 1H); 7.11 (s, 1H); 7.42 (d, 1H); 7.52 (m, 2H).

C011: 2.47 (s, 3H), 3.35-3.55 (m, 4H), 3.76 (d, 1H), 4.21 (d, 1H), 5.40 (m, 1H), 6.20 (br d, 1H), 7.40-7.80 (m, 6H).

C012: 2.50 (s, 3H), 3.78 (d, 1H), 4.04 (m, 2H), 4.21 (d, 1H), 4.53 (m, 2H), 4.89 (m, 1H), 6.45 (br d, 1H), 7.45-7.80 (m, 6H).

5 C013: 2.39 (s, 3H), 3.30 (d, 2H), 3.41 (d, 2H), 3.69 (s, 3H), 3.92 (s, 2H), 4.09 (d, 1H), 4.22 (s, 1H), 4.72 (m, 1H), 7.20-7.60 (m, 6H).

C014: 7.40 - 7.60 (m, 6 H), 5.91 (s, 1 H), 4.09 (d, J=17.2 Hz, 1 H), 3.79 - 3.70 (m, 5 H), 2.50 ppm (s, 3 H)

C020: 2.38 (s, 3H), 2.90 (m, 1H), 3.60-3.70 (m, 3H), 3.85-3.92 (m, 2H), 4.00 (d, 1H), 4.19 (m, 2H), 6.20 (br t, 1H), 7.30-7.50 (m, 6H)

C021: 2.48 (s, 3H), 3.07 (m, 2H), 3.32 (m, 2H), 3.55 (m, 1H), 3.64 (t, 2H), 3.71 (d, 1H), 4.11 (d, 1H), 6.10 (br t, 1H), 7.40-7.60 (m, 6H)

Biological examples

15 This Example illustrates the insecticidal and acaricidal properties of compounds of formula (I). Tests were performed as follows:

Spodoptera littoralis (Egyptian cotton leafworm):

Cotton leaf discs were placed on agar in a 24-well microtiter plate and sprayed with test solutions at an application rate of 200 ppm. After drying, the leaf discs were infested with 5 L1 larvae. The samples were checked for mortality, feeding behavior, and growth regulation 3 days after treatment (DAT).

The following compounds gave at least 80% control of *Spodoptera littoralis*: A001, A002, A003, A004, A013, A014, A015, A020, A021, A022, A023, A024, A025, A026, A027, A028, A029, A030, A031, A032, A033, A034, A035, A036, A037, A038, A039, A040, A042, A047, A048, A049, A050, A051, A052, A053, A054, A055, A056, A057, A058, A059, A061, A062, A063, A064, A065, A074, A075, A076, A077, A078, A079, A080, A081, A082, A083, A084, A085, A086, A087, A088, A089, A090, A091, A092, A093, A094, A095, A096, A097, A098, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A145, A146, A148, A157, A158, A159, A160, A162, A163, A165, A166, A167, A168, A169, A170, A172, A174, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A188, A189, A190, A191, A192, A193, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, B03, B04, B05, B06, B08, B09, B10, B11, B12, B13, B14, B15, B16, B17, B18, B19, B20, B21, B22, B23, B24, B25, B26, B27, B28, B29, B30, B31, B32, B33, B34, B35, B36, B37, B38, B39, B40, B41, B42, B43, B44, B45, B46, B47, B49, B50, B51, B53, B55, B56, B57, B58, B59, B61, B62, B63, B64, B67, B68, B69, C005, C006, C007, C011, C012, C14, C15, C16, C17, C18, C20

Heliothis virescens (Tobacco budworm):

Eggs (0-24 h old) were placed in 24-well microtiter plate on artificial diet and treated with test solutions at an application rate of 200 ppm (concentration in well 18 ppm) by pipetting. After an incubation period of 4 days, samples were checked for egg mortality, larval mortality, and growth regulation.

5 The following compounds gave at least 80% control of *Heliothis virescens*:

A001, A002, A003, A004, A013, A014, A015, A016, A017, A018, A019, A020, A021, A022, A023, A024, A025, A026, A027, A028, A029, A030, A031, A032, A033, A034, A035, A036, A037, A038, A039, A040, A042, A047, A048, A049, A050, A051, A052, A053, A054, A055, A056, A057, A058, A059, A060, A061, A062, A063, A064, A065, A068, A074, A075, A076, A077, A078, A079, A080, A081, A082, A083, A084, A085, A086, A087, A088, A089, A090, A091, A092, A093, A094, A095, A096, A097, A098, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A153, A157, A158, A159, A160, A162, A163, A165, A166, A167, A168, A169, A170, A171, A172, A174, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A191, A192, A193, A194, A196, A198, A200, A201, A202, A203, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, B03, B04, B05, B06, B08, B09, B10, B11, B12, B13, B14, B15, B16, B17, B18, B19, B20, B21, B22, B23, B24, B25, B26, B27, B28, B29, B30, B31, B32, B33, B34, B35, B36, B37, B38, B39, B40, B41, B42, B43, B44, B45, B46, B47, B49, B50, B51, B52, B53, B55, B56, B57, B58, B59, B61, B62, B63, B64, B67, B68, B69, C003, C005, C007, C008, C011, C012, C14, C15, C16, C17, C18, C20

Plutella xylostella (Diamond back moth):

25 24-well microtiter plate (MTP) with artificial diet was treated with test solutions at an application rate of 200 ppm (concentration in well 18 ppm) by pipetting. After drying, the MTP's were infested with L2 larvae (7-12 per well). After an incubation period of 6 days, samples were checked for larval mortality and growth regulation.

The following compounds gave at least 80% control of *Plutella xylostella*:

30 A001, A002, A003, A004, A005, A006, A007, A008, A009, A010, A011, A012, A013, A014, A015, A016, A017, A018, A019, A020, A021, A022, A023, A024, A025, A026, A027, A028, A029, A030, A031, A032, A033, A034, A035, A036, A037, A038, A039, A040, A041, A042, A043, A044, A045, A046, A047, A048, A049, A050, A051, A052, A053, A054, A055, A056, A057, A058, A059, A060, A061, A062, A063, A064, A065, A066, A067, A068, A069, A070, A071, A072, A073, A074, A075, A076, A077, A078, A079, A080, A081, A082, A083, A084, A085, A086, A087, A088, A089, A090, A091, A092, A093, A094, A095, A096, A097, A098, A099, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135,

A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A175, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A191, A192, A193, A194, A195, 5 A196, A197, A198, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, B01, B02, B03, B04, B05, B06, B07, B08, B09, B10, B11, B12, B13, B14, B15, B16, B17, B18, B19, B20, B21, B22, B23, B24, B25, B26, B27, B28, B29, B30, B31, B32, B33, B34, B35, B36, B37, B38, B39, B40, B41, B42, B43, B44, B45, B46, B47, B48, B49, B50, B51, B52, B53, B54, B55, B56, B57, B58, B59, B60, B61, B62, B63, B64, B65, 10 B66, B67, B68, B69, C005, C006, C007, C009, C012, C013, C14, C15, C16, C17, C18, C20

Diabrotica balteata (Corn root worm):

A 24-well microtiter plate (MTP) with artificial diet was treated with test solutions at an application rate of 200 ppm (concentration in well 18 ppm) by pipetting. After drying, the MTP's were infested with L2 15 larvae (6-10 per well). After an incubation period of 5 days, samples were checked for larval mortality and growth regulation.

The following compounds gave at least 80% control of *Diabrotica balteata*:

A001, A002, A003, A004, A006, A010, A011, A013, A014, A015, A017, A018, A019, A020, A021, A022, A023, A024, A025, A026, A027, A028, A029, A030, A031, A032, A034, A035, A036, A037, 20 A038, A039, A040, A042, A047, A048, A049, A050, A051, A052, A053, A054, A055, A056, A057, A058, A059, A060, A061, A062, A063, A064, A065, A069, A074, A075, A076, A077, A078, A079, A080, A081, A082, A084, A085, A086, A087, A088, A089, A090, A091, A092, A093, A094, A095, A096, A097, A098, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A120, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, 25 A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A150, A157, A158, A159, A160, A166, A167, A168, A169, A170, A171, A176, A177, A178, A179, A180, A181, A182, A183, A184, A186, A187, A188, A189, A190, A191, A192, A193, A194, A195, A198, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, B01, B02, B03, B04, B05, B06, B09, B10, B11, B12, B13, B14, 30 B15, B16, B17, B18, B19, B20, B21, B22, B23, B24, B25, B26, B27, B28, B29, B30, B31, B32, B33, B34, B35, B36, B37, B38, B39, B40, B42, B43, B44, B45, B46, B49, B50, B55, B56, B57, B58, B59, B61, B63, B67, B68, B69, C005, C006, C007, C011, C012, C013, C14, C15, C16, C17, C18, C20

Thrips tabaci (Onion thrips):

35 Sunflower leaf discs were placed on agar in a 24-well microtiter plate and sprayed with test solutions at an application rate of 200 ppm. After drying, the leaf discs were infested with a thrip population of mixed ages. After an incubation period of 7 days, samples were checked for mortality.

The following compounds gave at least 80% control of *Thrips tabaci*:

A001, A002, A003, A004, A005, A007, A008, A013, A014, A015, A017, A019, A020, A021, A022, A023, A024, A025, A026, A027, A028, A029, A030, A031, A032, A034, A035, A036, A037, A038, A039, A040, A041, A042, A043, A047, A048, A049, A050, A051, A052, A053, A054, A055, A056, A057, A058, A060, A061, A062, A063, A064, A065, A066, A067, A068, A069, A074, A075, A076, 5 A077, A078, A079, A080, A081, A082, A083, A084, A085, A086, A087, A088, A089, A090, A091, A092, A093, A094, A095, A096, A097, A098, A099, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A120, A121, A122, A123, A124, A125, A126, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A151, A152, A153, A155, A157, A158, 10 A159, A160, A163, A166, A167, A168, A169, A170, A171, A175, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A191, A192, A193, A194, A195, A196, A197, A198, A200, A201, A202, A203, A204, A206, A207, A208, A209, A210, A211, A213, A215, A216, A217, A218, A219, A220, B02, B03, B04, B05, B06, B08, B09, B10, B11, B12, B13, B14, B15, B16, B17, B19, B20, B21, B22, B23, B24, B25, B26, B27, B28, B29, B30, B31, B32, B33, B34, 15 B35, B36, B37, B38, B39, B40, B41, B42, B43, B44, B45, B46, B47, B49, B50, B51, B53, B56, B58, B59, B61, B62, B63, B66, B68, B69, C006, C007, C008, C009, C010, C013, C14, C15, C16, C17, C18, C20

Tetranychus urticae (Two-spotted spider mite):

20 Bean leaf discs on agar in 24-well microtiter plates were sprayed with test solutions at an application rate of 200 ppm. After drying, the leaf discs are infested with mite populations of mixed ages. 8 days later, discs are checked for egg mortality, larval mortality, and adult mortality.

The following compounds gave at least 80% control of *Tetranychus urticae*:

A001, A002, A003, A004, A005, A006, A008, A012, A013, A014, A015, A017, A020, A021, A022, 25 A023, A024, A025, A026, A027, A028, A029, A030, A031, A032, A033, A034, A035, A036, A037, A038, A039, A040, A042, A047, A048, A049, A050, A051, A052, A053, A054, A055, A056, A057, A058, A059, A060, A061, A062, A063, A064, A065, A066, A067, A068, A069, A074, A075, A076, A077, A078, A079, A080, A081, A082, A083, A084, A085, A086, A087, A088, A089, A090, A092, A093, A094, A095, A096, A097, A098, A099, A101, A102, A103, A104, A105, A106, A107, A108, 30 A109, A110, A111, A112, A113, A114, A115, A116, A117, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A151, A152, A153, A155, A156, A157, A158, A159, A160, A162, A163, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, 35 A190, A191, A192, A193, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A213, A214, A215, A216, A217, A218, A219, A220, B01, B02, B03, B04, B05, B06, B09, B10, B11, B12, B13, B14, B15, B16, B17, B18, B19, B20, B21, B22, B23, B24, B25, B26, B27, B28, B29, B30, B31, B32, B33, B34, B35, B36, B37, B38, B39, B40,

B43, B44, B45, B46, B47, B48, B50, B51, B56, B57, B58, B59, B61, B62, B63, B64, B66, B67, B68, B69, C005, C006, C007, C008, C009, C010, C012, C013, C14, C15, C16, C17, C18, C20

Myzus persicae (sachet) (Green peach aphid) mixed population

- 5 Test compounds were applied by pipette into 24 well plates and mixed with Sucrose solution. The plates were closed with a stretched Parafilm. A plastic stencil with 24 holes is placed onto the plate and infested pea seedlings were placed directly on the Parafilm. The infested plate is closed with a gel blotting paper and another plastic stencil and then turned upside down. 5 days after infestation the samples were checked on mortality. Application rate: 12.5ppm.
- 10 The following compounds gave at least 80% control of *Myzus persicae* : A002, A004, A026, A027, A038, A064, A080, A088, A089, A090, A122, A124, A125, A132, A133, A139, A145, A178, A188, A190, A218, B06, B15, B16, B27, B31, B36, B39, B50, B58, C15, C16, C17, C18, C20

Euschistus heros (Neotropical brown stink bug)

- 15 (contact/feeding activity)
- 2 week old Soybean plants were sprayed in a turn table spray chamber with the diluted spray solution at an application rate of 50 ppm. After drying, 2 soybean seeds were added and plants were infested with 10 N-2 nymphs of the neotropical brown stink bug *Euschistus heros* in plastic test boxes. Boxes were incubated in a climate chamber at 25°C and 60 % RH. Evaluation is done 5 days after infestation on
- 20 mortality and growth effect.
- The following compound gave at least 80% control of *Euschistus heros*:
- A188, C15, C16, C17, C18, C20

Nilaparvata lugens (Brown plant hopper)

- 25 (larvicide, feeding/contact)
- Rice seedlings are treated with the diluted test solutions in a spray chamber an application rate of 200 ppm. After drying, plants are infested with 20 N₃ nymphs (2 replicates). 6-12 days after the treatment samples are checked for mortality. The following compound gave at least 80% control of *Nilaparvata lugens*:
- 30 A002, A026, A027, A038, A064, A080, A088, A090, A124, A125, A133, A139, A145, B04, B06, B27, B31, B36, B39, B50, B58, C15, C17, C18, C20

Anthonomus grandis (Cotton boll weevil)

- (adulticide, feeding/contact)
- 35 Cotton plants are treated with the diluted test solutions in a spray chamber at an application rate of 50ppm. After drying, plants are infested with 10 adult weevils (2 replicates). 5 days after the treatment samples are checked for mortality. The following compound gave at least 80% control of *Anthonomus grandis*:

WO 2013/026931

210


PCT/EP2012/066554

C15, C16, C17, C18, C20

CLAIMS

1. A compound of formula I

5




wherein

G¹ is oxygen;

R¹ is hydrogen;

10 R² is group P



L is a bond, methylene or ethylene;

one of A¹ and A² is S, SO or SO₂ and the other is -C(R⁴)R⁴;

R³ is hydrogen or methyl;

15 each R⁴ is independently hydrogen or methyl;

Y¹, Y² and Y³ are independently CH or nitrogen;

wherein no more than two of Y¹, Y² and Y³ are nitrogen and wherein Y² and Y³ are not both nitrogen;

R⁵ is hydrogen, halogen, cyano, nitro, NH₂, C₁-C₂alkyl, C₁-C₂haloalkyl, C₃-C₃cycloalkyl, C₁-

C₂halocycloalkyl, C₁-C₂alkoxy, C₁-C₂haloalkoxy;

20 X² is C-X⁶ or nitrogen;

X¹, X³ and X⁶ are independently hydrogen, halogen or trihalomethyl, wherein at least two of X¹, X³ and X⁶ are not hydrogen;

X⁴ is trifluoromethyl, difluoromethyl or chlorodifluoromethyl.

25 2. A compound according to claim 1, wherein R² is thietan-3-yl-, 1-oxo-thietan-3-yl-, 1,1-Dioxo-thietan-3-yl-, 2,2-dimethylthietan-3-yl-, 2,2-dimethyl-1-oxo-thietan-3-yl-, 2,2-dimethyl-1,1-Dioxo-thietan-3-yl-, 3-Methyl-thietan-3-yl-, 3-Methyl-1-oxo-thietan-3-yl-, 3-Methyl-1,1-Dioxo-thietan-3-yl-, thietan-3-ylmethyl-, 1-oxo-thietan-3-ylmethyl-, 1,1-Dioxo-thietan-3-ylmethyl-, thietan-2-ylmethyl-, (1-

oxothietan-2-yl)methyl-, (1,1-dioxothietan-2-yl)methyl-, 2-(thietan-3-yl)ethanyl, 2-(1,1-dioxothietan-3-yl)ethanyl, or 2-(1-oxothietan-3-yl)ethanyl.

3. A compound according to claim 1 or claim 2, wherein

5 Y¹ is CH, Y² is CH, Y³ is CH, or

Y¹ is N, Y² is CH, Y³ is CH, or

Y¹ is N, Y² is N, Y³ is CH, or

Y¹ is CH, Y² is N, Y³ is CH, or

Y¹ is CH, Y² is CH, Y³ is N.

10

4. A compound according to any one of claims 1 to 3, wherein R⁵ is hydrogen, chloro, bromo, fluoro, trifluoromethyl, methyl, ethyl, methoxy, nitro, trifluoromethoxy, cyano, or cyclopropyl.

5. A compound according to any one of claims 1 to 4, wherein

15 X¹ is chloro, X² is CH, X³ is chloro, or

X¹ is chloro, X² is C-F, X³ is hydrogen, or

X¹ is fluoro, X² is C-Cl, X³ is hydrogen, or

X¹ is chloro, X² is C-Cl, X³ is hydrogen, or

X¹ is chloro, X² is C-Br, X³ is chloro, or

20 X¹ is chloro, X² is C-F, X³ is chloro, or

X¹ is chloro, X² is C-Cl, X³ is chloro, or

X¹ is chloro, X² is C-I, X³ is chloro, or

X¹ is fluoro, X² is C-F, X³ is fluoro, or

X¹ is chloro, X² is CH, X³ is bromo, or

25 X¹ is chloro, X² is CH, X³ is fluoro, or

X¹ is chloro, X² is CH, X³ is trifluoromethyl, or

X¹ is chloro, X² is C-Cl, X³ is trifluoromethyl, or

X¹ is trifluoromethyl, X² is CH, X³ is trifluoromethyl, or

X¹ is trifluoromethyl, X² is C-Cl, X³ is trifluoromethyl, or

30 X¹ is trifluoromethyl, X² is CH, X³ is hydrogen, or


X¹ is chloro, X² is N, X³ is chloro, or

X¹ is trifluoromethyl, X² is N, X³ is trifluoromethyl.

6. A compound according to any one of claims 1 to 5, wherein X² is C-X⁶, Y¹, Y² and Y³ are C-H,

35 R³ is hydrogen and each R⁴ is hydrogen.


7. A compound of formula X1



wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I in any one of claims 1 to 6.

5


8. A compound of formula X2



wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I in any one of claims 1 to 6.


10

9. A compound of formula X3




15 wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I in any one of claims 1 to 6.

10. A compound of formula X4



wherein G^1 , R^1 , R^2 , R^5 , X^1 , X^2 , X^3 , X^4 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I in any one of claims 1 to 6.

5 11. A compound of formula X5



wherein G^1 , R^1 , R^2 , R^5 , Y^1 , Y^2 , and Y^3 are as defined for the compound of formula I in any one of claims
 10 1 to 6, and X^5 is chloro, bromo, iodo, amino, CHO, CN, OH, C(=O)OH, C(=NOH)H, C(=NOH)Cl, C(=NOH)NH₂, C(=O)CH₃, C(=NOH)CH₃, C(=O)CH₂Cl, or C(=O)CH₂Br.

12. A method of controlling insects, acarines, nematodes or molluscs which comprises applying to a pest, to a locus of a pest, or to a plant susceptible to attack by a pest an insecticidally, acaricidally,
 15 nematocidally or molluscicidally effective amount of a compound of formula (I) as defined in any one of claims 1 to 6.

13. An insecticidal, acaricidal, nematocidal or molluscicidal composition comprising an insecticidally, acaricidally, nematocidally or molluscicidally effective amount of a compound of formula (I) as defined in
 20 any one of claims 1 to 6.

14. An insecticidal, acaricidal, nematocidal or molluscicidal composition according to claim 13 comprising at least one additional compound having biological activity.

25 15. A combination product comprising a pesticidally effective amount of a component A and a pesticidally effective amount of component B, wherein component A is a compound of formula (I) as defined in any one of claims 1 to 6, and compound B is imidacloprid, enrofloxacin, praziquantel, pyrantel embonate, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, fipronil,

ivermectin, omeprazole, tiamulin, benazepril, milbemycin, cyromazine, thiamethoxam, pyriprole, deltamethrin, cefquinome, florfenicol, buserelin, ceftiofur, selamectin, carprofen, metaflumizone, moxidectin, methoprene (including S-methoprene), clorsulon, pyrantel, amitraz, triclabendazole, avermectin, abamectin, emamectin, eprinomectin, doramectin, selamectin, 5 nemadectin, albendazole, cambendazole, fenbendazole, flubendazole, mebendazole, oxfendazole, oxibendazole, parbendazole, tetramisole, levamisole, pyrantel pamoate, oxantel, morantel, triclabendazole, epsiprantel, fipronil, lufenuron, ecdysone or tebufenozide, wherein the combination product is for use in a method of therapeutic treatment.

- 10 16. A combination product comprising a pesticidally effective amount of a component A and a pesticidally effective amount of component B, wherein component A is a compound of formula (I) as defined in any one of claims 1 to 6, wherein component B is enrofloxacin, praziquantel, pyrantel embonate, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, omeprazole, tiamulin, benazepril, pyriprole, cefquinome, florfenicol, buserelin, ceftiofur, selamectin, carprofen, moxidectin, clorsulon, pyrantel, eprinomectin, doramectin, selamectin, 15 nemadectin, albendazole, cambendazole, fenbendazole, flubendazole, mebendazole, oxfendazole, oxibendazole, parbendazole, tetramisole, levamisole, pyrantel pamoate, oxantel, morantel, triclabendazole, epsiprantel, lufenuron or ecdysone.
- 20 17. A combination product according to claim 15 or claim 16, wherein component B is enrofloxacin, praziquantel, pyrantel embonate, febantel, penethamate, moloxicam, cefalexin, kanamycin, pimobendan, clenbuterol, omeprazole, tiamulin, benazepril, pyriprole, cefquinome, florfenicol, buserelin, ceftiofur, selamectin, carprofen, moxidectin, clorsulon or pyrantel.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/066554

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D331/04 C07D409/12 C07D413/12 A01N43/36 A01N43/80
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2010/020521 A1 (SYNGENTA PARTICIPATIONS AG [CH]; PITTERNA THOMAS [CH]; EL QACEMI MYRIE) 25 February 2010 (2010-02-25) Whole document, specially claim 1 -----	1-17
X	WO 2009/080250 A2 (SYNGENTA PARTICIPATIONS AG [CH]; RENOLD PETER [CH]; ZAMBACH WERNER [CH]) 2 July 2009 (2009-07-02) cited in the application Whole document, specially compounds in table A, page 63-65 and tests in page 67 -----	1-6, 12-17
X,P	WO 2011/154555 A1 (SYNGENTA PARTICIPATIONS AG [CH]; SYNGENTA LTD [GB]; EL QACEMI MYRIEM []) 15 December 2011 (2011-12-15) Compound in lines 13-14 in page 131 ----- -/--	9

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

25 October 2012

Date of mailing of the international search report

09/11/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Sahagún Krause, H

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/066554

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 2011/104089 A1 (SYNGENTA PARTICIPATIONS AG [CH]; SYNGENTA LTD [GB]; MULHOLLAND NICHOLA) 1 September 2011 (2011-09-01) Formula (IB) in claim 1; examples and formula (II) and intermediates and products in examples 10, 11 and 12 -----	1-6,9, 11-17
X,P	WO 2012/104331 A2 (SYNGENTA PARTICIPATIONS AG [CH]; SYNGENTA LTD [GB]; VOCK CHRISTOPH [CH]) 9 August 2012 (2012-08-09) compounds in claims 5 and 10 -----	1-6, 12-17
X,P	WO 2012/067235 A1 (NISSAN CHEMICAL IND LTD [JP]; MITA TAKESHI [JP]; MAIDURU YUKIHIRO [JP]) 24 May 2012 (2012-05-24) compounds 1-001 to 1-012 in p. 109, 4-001 in p. 110 -----	1-6, 12-17
X,P	WO 2011/104088 A1 (SYNGENTA PARTICIPATIONS AG [CH]; SYNGENTA LTD [GB]; DUTTON ANA CRISTIN) 1 September 2011 (2011-09-01) claim 1 and compounds in Table A, p. 8 -----	1-6, 12-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/066554

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010020521 A1	25-02-2010	AR 073115 A1 AU 2009284229 A1 CA 2733160 A1 CN 102131789 A CO 6341524 A2 EA 201100356 A1 EP 2331515 A1 JP 2012500782 A MA 32574 B1 TW 201021707 A US 2011166184 A1 UY 32070 A WO 2010020521 A1	13-10-2010 25-02-2010 25-02-2010 20-07-2011 21-11-2011 31-10-2011 15-06-2011 12-01-2012 01-08-2011 16-06-2010 07-07-2011 26-03-2010 25-02-2010
WO 2009080250 A2	02-07-2009	AR 069702 A1 AU 2008340691 A1 CA 2709246 A1 CN 101910161 A CO 6300905 A2 CR 11531 A DK 2238132 T3 DO P2010000196 A EA 201001008 A1 EC SP10010304 A EP 2238132 A2 EP 2518068 A2 JP 2011507813 A KR 20100112588 A MA 32749 B1 NZ 586083 A TW 200926983 A US 2010279999 A1 UY 31578 A1 WO 2009080250 A2	10-02-2010 02-07-2009 02-07-2009 08-12-2010 21-07-2011 13-09-2010 22-10-2012 15-09-2010 28-02-2011 30-07-2010 13-10-2010 31-10-2012 10-03-2011 19-10-2010 01-11-2011 29-06-2012 01-07-2009 04-11-2010 03-08-2009 02-07-2009
WO 2011154555 A1	15-12-2011	TW 201211057 A WO 2011154555 A1	16-03-2012 15-12-2011
WO 2011104089 A1	01-09-2011	AU 2011220041 A1 UY 33249 A WO 2011104089 A1	23-08-2012 30-09-2011 01-09-2011
WO 2012104331 A2	09-08-2012	NONE	
WO 2012067235 A1	24-05-2012	NONE	
WO 2011104088 A1	01-09-2011	NONE	