



- (51) **International Patent Classification:**
C07D 213/74 (2006.01) *A01N 43/40* (2006.01)
- (21) **International Application Number:**
PCT/EP2015/057088
- (22) **International Filing Date:**
31 March 2015 (31.03.2015)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
14164464.1 11 April 2014 (11.04.2014) EP
- (71) **Applicant:** SYNGENTA PARTICIPATIONS AG
[CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).
- (72) **Inventors:** HOFFMAN, Thomas James; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH). SULZER-MOSSE, Sarah; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH). NEBEL, Kurt; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH). CEDERBAUM, Fredrik; Syngenta Crop Protection, Münchwilen AG, Schaffhauserstrasse, CH-4332 Stein (CH).
- (74) **Agent:** SYNGENTA INTERNATIONAL AG; Intellectual Property, WRO 1008-Z1-26, Schwarzwaldallee 215, CH-4058 Basel (CH).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(54) **Title:** FUNGICIDAL N'-[2-METHYL-6-[2-ALKOXY-ETHOXY]-3-PYRIDYL]-N-ALKYL-FORMAMIDINE DERIVATIVES FOR USE IN AGRICULTURE



(57) **Abstract:** The invention relates to compounds of formula (I), wherein R^1 , R^2 , R^3 , R^4 , R^5 and R^6 are as defined in the claims. The invention further provides compositions which comprise these compounds and to their use in agriculture or horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.




FUNGICIDAL N'-[2-METHYL-6-[2-ALKOXY-ETHOXY]-3-PYRIDYL]-N-ALKYL-FORMAMIDINE
DERIVATIVES FOR USE IN AGRICULTURE

The present invention relates to novel microbiocidal, in particular fungicidal, pyridylamidine compounds. It further relates to intermediates used in the preparation of these compounds, to compositions which comprise these compounds, and to their use in agriculture or horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.

Certain pyridylamidines derivatives have been proposed in the literature as microbiocidal active ingredients in pesticides. For example, WO 00/46184 and WO 03/093224 disclose pyridylamidines which are useful as fungicides. However, the biological properties of these known compounds are not entirely satisfactory for controlling or preventing infestation of plants by phytopathogenic microorganisms, which is why there is a need to provide other compounds which have microbicidal properties.

The present invention relates to compounds of formula (I)



wherein

R^1 represents hydrogen, halogen, cyano, OH, NH_2 , C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, $NH(C_1$ - C_4 alkyl), $N(C_1$ - C_4 alkyl) $_2$, $CO(C_1$ - C_4 alkyl), $CO_2(C_1$ - C_4 alkyl), CO_2H , $CONH(C_1$ - C_4 alkyl), $CON(C_1$ - C_4 alkyl) $_2$, $SO_2NH(C_1$ - C_4 alkyl), $SO_2N(C_1$ - C_4 alkyl) $_2$, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkyl- C_1 - C_4 alkoxy or C_2 - C_4 alkynyl;

R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, R^7 or $-C_1$ - C_2 alkyl- R^7 , each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl and C_1 - C_4 haloalkoxy;

R^3 and R^4 independently of each other represent hydrogen, C_1 - C_4 alkyl or C_3 - C_6 cycloalkyl; or

R^3 and R^4 together with the nitrogen atom to which they are attached form a 3-6-membered saturated cyclic group;

R^5 represents H, C_1 - C_4 alkyl or C_1 - C_4 haloalkyl;

R^6 represents C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl or C_1 - C_4 haloalkoxy;

- 2 -

R⁷ represents a three- to ten-membered monocyclic or fused bicyclic ring system which can be aromatic, partially saturated or fully saturated and can contain 1 to 4 hetero atoms selected from the group consisting of nitrogen, oxygen and sulfur, it being possible for the three- to ten-membered ring system itself to be optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₄ haloalkyl, C₁-C₄ alkoxy and C₁-C₄ haloalkoxy;

and tautomers/isomers/enantiomers/salts and N-oxides of these compounds

Substituents at a nitrogen atom are always different from halogen. A hydroxy, mercapto or amino substituent is not to be placed on an α -carbon relative to a heteroatom of a core fragment.

Halogen, either as a lone substituent or in combination with another substituent (e.g. haloalkyl) is generally fluorine, chlorine, bromine or iodine, and usually fluorine, chlorine or bromine.

Each alkyl moiety (including the alkyl moiety of alkoxy, alkylthio, etc.) is a straight or branched chain and, depending on the number of carbon atoms it contains, is, for example, methyl, ethyl, *n*-propyl, *n*-butyl, *n*-pentyl, *n*-hexyl, *iso*-propyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *neo*-pentyl, *n*-heptyl or 1,3-dimethylbutyl, and usually methyl or ethyl.

The alkenyl and alkynyl groups can be mono- or di-unsaturated and examples thereof are derived from the above mentioned alkyl groups.

The alkenyl group is an unsaturated straight or branched chain having a carbon-carbon double bond and, depending on the number of carbon atoms it contains, is, for example ethenyl, 1-propenyl, 2-propenyl, 1-methyl-ethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 1-methyl-1-butenyl, 2-methyl-1-butenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl, 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-propenyl, 1,2-dimethyl-1-propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-2-propenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-1-pentenyl, 2-methyl-1-pentenyl, 3-methyl-1-pentenyl, 4-methyl-1-pentenyl, 1-methyl-2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl, 4-methyl-2-pentenyl, 1-methyl-3-pentenyl, 2-methyl-3-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl, 3-methyl-4-pentenyl, 4-methyl-4-pentenyl, 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl, 1,2-dimethyl-1-butenyl, 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl, 1,3-dimethyl-1-butenyl, 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl, and usually 2-propenyl, 1-methyl-2-propenyl, 2-butenyl, 2-methyl-2-propenyl.

- 3 -

The alkynyl group is an unsaturated straight or branched chain having a carbon-carbon triple bond and, depending on the number of carbon atoms it contains, is, for example ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-methyl-2-propynyl, 1-pentynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 3-methyl-1-butynyl, 1-methyl-2-butynyl, 1-methyl-3-butynyl, 2-methyl-3-butynyl, 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl, 3-methyl-1-pentynyl, 4-methyl-1-pentynyl, 1-methyl-2-pentynyl, 4-methyl-2-pentynyl, 1-methyl-3-pentynyl, 2-methyl-3-pentynyl, 1-methyl-4-pentynyl, 2-methyl-4-pentynyl, 3-methyl-4-pentynyl, 3,3,-dimethyl-1-butynyl, 1-ethyl-2-butynyl, 1,1-dimethyl-2-butynyl, 1-ethyl-3-butynyl, 2-ethyl-3-butynyl, 1,1-dimethyl-3-butynyl, 2,2-dimethyl-3-butynyl, 1,2-dimethyl-3-butynyl.

Haloalkyl moieties are alkyl moieties which are substituted by one or more of the same or different halogen atoms and are, for example, monofluoromethyl, difluoromethyl, trifluoromethyl, monochloromethyl, dichloromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, 2-fluoroethyl, 1,1-difluoroethyl, 1-fluoroethyl, 2-chloroethyl, pentafluoroethyl, 1,1-difluoro-2,2,2-trichloroethyl, 2,2,3,3-tetrafluoroethyl and 2,2,2-trichloroethyl, and typically trichloromethyl, difluorochloromethyl, difluoromethyl, trifluoromethyl and dichlorofluoromethyl.

Alkoxy is, for example, methoxy, ethoxy, propoxy, *iso*-propoxy, *n*-butoxy, *iso*-butoxy, *sec*-butoxy and *tert*-butoxy, and usually methoxy or ethoxy.

Haloalkoxy is, for example, fluoromethoxy, difluoromethoxy, trifluoromethoxy, 2,2,2-trifluoroethoxy, 1,1,2,2-tetrafluoroethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2,2-difluoroethoxy and 2,2,2-trichloroethoxy, and usually difluoromethoxy, 2-chloroethoxy and trifluoromethoxy.

Alkylthio is, for example, methylthio, ethylthio, propylthio, *iso*-propylthio, *n*-butylthio, *iso*-butylthio, *sec*-butylthio or *tert*-butylthio, and usually methylthio or ethylthio.

Alkylsulphonyl is, for example, methylsulphonyl, ethylsulphonyl, propylsulphonyl, *iso*-propylsulphonyl, *n*-butylsulphonyl, *iso*-butylsulphonyl, *sec*-butylsulphonyl or *tert*-butylsulphonyl, and usually methylsulphonyl or ethylsulphonyl.

Alkylsulphinyl is, for example, methylsulphinyl, ethylsulphinyl, propylsulphinyl, *iso*-propylsulphinyl, *n*-butylsulphinyl, *iso*-butylsulphinyl, *sec*-butylsulphinyl or *tert*-butylsulphinyl, and usually methylsulphinyl or ethylsulphinyl.

Cycloalkyl may be saturated or partially unsaturated, preferably fully saturated, and is, for example, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

Alkoxyalkyl is, for example, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, *n*-propoxymethyl, *n*-propoxyethyl, *iso*-propoxymethyl or *iso*-propoxyethyl.

Aryl includes phenyl, naphthyl, anthracyl, fluorenyl and indanyl, but is usually phenyl.

Carbocycle includes cycloalkyl groups and aryl groups.

- 4 -

Heterocycloalkyl is a non-aromatic ring that may be saturated or partially unsaturated, preferably fully saturated, containing carbon atoms as ring members and at least one heteroatom selected from O, S and N as ring members. Examples include oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, 1,3-dioxolanyl, 1,4-dioxanyl, aziridinyl, azetidiny,
5 pyrrolidinyl, piperidinyl, oxazinanyl, morpholinyl, thiomorpholinyl, imidazolidinyl, pyrazolidinyl and piperazinyl, preferably morpholinyl, pyrrolidinyl, piperdiny and piperazinyl, more preferably morpholinyl and pyrrolidinyl.

Heteroaryl is, for example, a monovalent monocyclic or bicyclic aromatic hydrocarbon radical. Examples of monocyclic groups include pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl,
10 pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, thiophenyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, and thiadiazolyl. Examples of bicyclic groups include quinolinyl, cinnolinyl, quinoxalinyl, benzimidazolyl, benzothiophenyl, and benzothiadiazolyl. Monocyclic heteroaryl groups are preferred, preferably pyridyl, pyrrolyl, imidazolyl and triazolyl, e.g. 1,2,4 triazolyl, pyridyl and imidazolyl being most preferred.

The terms "heterocycle" and "heterocyclic ring" are used interchangeably and are defined to include heterocycloalkyl and heteroaryl groups. Any reference herein to a heterocycle or heterocyclic ring preferably refers to the specific examples given under the definition of heteroaryl and heterocycloalkyl above, and are preferably morpholinyl, pyrrolidinyl, piperdiny, piperazinyl pyridyl, pyrrolyl, imidazolyl and triazolyl, e.g. 1,2,4 triazolyl,
20 more preferably morpholinyl, pyrrolidinyl, pyridyl and imidazolyl. No heterocycle contains adjacent oxygen atoms, adjacent sulphur atoms, or adjacent oxygen and sulphur atoms.

Where a moiety is indicated as being (optionally) substituted, e.g. alkyl, this includes those moieties where they are part of a larger group, e.g. the alkyl in the alkylthio group. The same applies, e.g. to the phenyl moiety in phenylthio etc. Where a moiety is indicated as being optionally substituted by one or more other groups, preferably there are one to five optional substituents, more preferably one to three optional substituents. Where a moiety is substituted by a cyclic group, e.g. aryl, heteroaryl, cycloalkyl, preferably there are no more than two such substituents, more preferably no more than one such substituent.

The following list provides definitions, including preferred definitions, for substituents R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 with reference to compounds of formula I. For any one of these substituents, any of the definitions given below may be combined with any definition of any other substituent given below or elsewhere in this document.

R^1 represents hydrogen, halogen, cyano, OH, NH_2 , C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, $NH(C_1$ - C_4 alkyl), $N(C_1$ - C_4 alkyl) $_2$, $CO(C_1$ - C_4 alkyl), $CO_2(C_1$ - C_4 alkyl), CO_2H , $CONH(C_1$ - C_4

- 5 -

alkyl), CON(C₁-C₄ alkyl)₂, SO₂NH(C₁-C₄ alkyl), SO₂N(C₁-C₄ alkyl)₂, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, C₁-C₄ alkyl-C₁-C₄ alkoxy or C₂-C₄ alkynyl

Preferably, R¹ represents hydrogen, halogen, cyano, OH, NH₂, C₁-C₄ alkyl, C₃-C₆ cycloalkyl, NH(C₁-C₄ alkyl), N(C₁-C₄ alkyl)₂, CO(C₁-C₄ alkyl), CO₂(C₁-C₄ alkyl), CO₂H,
 5 CONH(C₁-C₄ alkyl), CON(C₁-C₄ alkyl)₂, SO₂NH(C₁-C₄ alkyl), SO₂N(C₁-C₄ alkyl)₂, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, or C₂-C₄ alkynyl.

More preferably, R¹ represents hydrogen, halogen, cyano, OH, NH₂, methyl, ethyl, cyclopropyl, NH(C₁-C₂ alkyl), N(C₁-C₂ alkyl)₂, CO(C₁-C₂ alkyl), CO₂(C₁-C₂ alkyl), CO₂H,
 10 CONH(C₁-C₂ alkyl), CON(C₁-C₂ alkyl)₂, SO₂NH(C₁-C₂ alkyl), SO₂N(C₁-C₂ alkyl)₂, C₁-C₄ fluoroalkyl, C₁-C₄ alkoxy, C₁-C₂ haloalkoxy or C₂-C₄ alkynyl.

Even more preferably, R¹ represents hydrogen, halogen, cyano, OH, NH₂, methyl, ethyl, cyclopropyl, NHMe, NMe₂, COMe, CO₂Me, CO₂H, CONHMe, CONMe₂, SO₂NHMe, SO₂NMe₂, CHF₂, CF₃, OMe, OCHF₂ or acetylenyl.

Yet more preferably, R₁ represents hydrogen, halogen, cyano, methyl, ethyl,
 15 cyclopropyl, CHF₂, CF₃, OMe, or OCHF₂.

Most preferably, R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano.

R² represents C₃-C₆alkyl, C₃-C₆alkenyl, C₃-C₆alkynyl, R⁷ or -C₁-C₂alkyl-R⁷, each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl and C₁-C₄ haloalkoxy.

20 Preferably, R² represents C₃-C₆alkyl, C₃-C₆alkenyl, C₃-C₆alkynyl, C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, -C₁-C₂alkyl-C₃-C₆cycloalkyl, -C₁-C₂alkyl-C₃-C₆cycloalkenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₂ alkyl, C₁-C₂ haloalkyl and C₁-C₂ haloalkoxy.

More preferably, R² represents C₃-C₆alkyl, C₃-C₆alkenyl, C₃-C₆alkynyl, C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, -C₁-C₂alkyl-C₃-C₆cycloalkyl, -C₁-C₂alkyl-C₃-C₆cycloalkenyl which may be optionally substituted by one or more groups independently selected from the group
 25 consisting fluoro, C₁-C₂ alkyl, C₁-C₂ fluoroalkyl and C₁-C₂ fluoroalkoxy.

Yet more preferably, R² is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, 2-methylbutyl, 2,2-dimethylpropyl, 2-methylpentyl, 3-methylpentyl, 2,3-dimethylbutyl,
 30 2,2-dimethylbutyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl -CH₂-cyclopropyl, -CH₂-cyclobutyl, -CH₂-cyclopentyl, and -CH₂-cyclopentenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro, methyl and difluoromethoxy.

- 6 -

Yet more preferably again, R_2 is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro and difluoromethoxy.

Most preferably, R_2 is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or cyclopropyl.

5 In another group of compounds, R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, R^7 or $-C_1$ - C_2 alkyl- R^7 , each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 alkyl and C_1 - C_4 haloalkyl.

10 Preferably in this group of compounds, R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_2 alkyl and C_1 - C_2 haloalkyl.

15 More preferably in this group of compounds, R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkenyl which may be optionally substituted by one or more groups independently selected from the group consisting fluoro, C_1 - C_2 alkyl and C_1 - C_2 fluoroalkyl.

20 Yet more preferably in this group of compounds, R^2 is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, 2-methylbutyl, 2,2-dimethylpropyl, 2-methylpentyl, 3-methylpentyl, 2,3-dimethylbutyl, 2,2-dimethylbutyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl $-CH_2$ -cyclopropyl, $-CH_2$ -cyclobutyl, $-CH_2$ -cyclopentyl, and $-CH_2$ -cyclopentenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro and methyl.

Yet more preferably again in this group of compounds, R_2 is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more fluoro.

25 R^3 and R^4 independently of each other represent hydrogen, C_1 - C_4 alkyl or C_3 - C_6 cycloalkyl; or

R^3 and R^4 together with the nitrogen atom to which they are attached form a 3-6-membered saturated cyclic group;

Preferably, R^3 and R^4 independently of each other represent hydrogen, C_1 - C_3 alkyl or C_3 - C_5 cycloalkyl; or

30 R^3 and R^4 together with the nitrogen atom to which they are attached form a 4- or 5-membered saturated cyclic group.

More preferably, R^3 and R^4 independently of each other represent hydrogen, methyl, ethyl, isopropyl or cyclopropyl; or

35 R^3 and R^4 together with the nitrogen atom to which they are attached form a 4-membered saturated cyclic group.

- 7 -

In one group of compounds, R^3 is hydrogen or methyl;

R^4 is methyl or ethyl.

Preferably in this group of compounds, R^3 represents hydrogen or methyl;

R^4 is ethyl.

5 R^5 represents H, C_1 - C_4 alkyl or C_1 - C_4 haloalkyl.

Preferably, R^5 represents H or C_1 - C_4 alkyl.

More preferably, R^5 represents H or methyl

Most preferably, R^5 is hydrogen.

R^6 represents C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl or C_1 - C_4 haloalkoxy.

10 Preferably, R^6 represents C_1 - C_4 alkyl or C_1 - C_4 alkoxy.

More preferably, R^6 represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl, methoxy or ethoxy.

Even more preferably, R^6 represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, or cyclopentyl.



15 Yet more preferably, R^6 represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl or cyclopentyl.

Even yet more preferably, R^6 represents methyl, ethyl, n-propyl, iso-propyl, or cyclopropyl.

More preferably still, R^6 represents methyl, ethyl, n-propyl or iso-propyl.

20 Most preferably, R^6 is methyl.

When R^5 and R^6 are not the same, the compounds of formula (I) can occur in (at least) two enantiomeric forms: (Ia) and (Ib).



25 A racemic compound (I) is a 1:1 mixture of the compounds of formula (Ia) and (Ib). Other ratios of (Ia) and (Ib) are possible and part of the present invention. Examples of such ratios of (Ia) to (Ib) are 1:99, 2:98, 5:95, 10:90, 20:80, 30:70; 40:60, 45:55; 55:45; 60:40, 70:30, 80:20, 90:10, 95:5, 98:2, and 99:1.

- 8 -

In one embodiment of the invention, the weight ratio of (Ib) to (Ia) is weighted towards compound of formula (Ia), for example, the the ratio of (Ib) to (Ia) being 1:99, 2:98, 5:95, 10:90, 20:80, 30:70; 40:60 or 45:55. More preferably in this embodiment of the invention, the compound of formula (I) consists essentially of the compound of formula (Ia); even more
5 preferably, the compound of formula (I) is the compound of formula (Ia).

In another embodiment of the invention, the weight ratio of (Ia) to (Ib) is weighted towards compound of formula (Ib), for example, the the ratio of (Ia) to (Ib) being 1:99, 2:98, 5:95, 10:90, 20:80, 30:70; 40:60 or 45:55. More preferably in this embodiment of the invention, the compound of formula (I) consists essentially of the compound of formula (Ib);
10 even more preferably, the compound of formula (I) is the compound of formula (Ib).

R⁷ represents a three- to ten-membered monocyclic or fused bicyclic ring system which can be aromatic, partially saturated or fully saturated and can contain 1 to 4 hetero atoms selected from the group consisting of nitrogen, oxygen and sulfur, it being possible for the three- to ten-membered ring system itself to be optionally substituted by one or more groups
15 independently selected from the group consisting halogen, C₁-C₄ haloalkyl, C₁-C₄ alkoxy and C₁-C₄ haloalkoxy.

Preferably, R⁷ represents C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, phenyl, naphthyl, anthracyl, fluorenyl, indanyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, 1,3-dioxolanyl, 1,4-dioxanyl, aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, oxazinanyl, morpholinyl, thiomorpholinyl, imidazolidinyl, pyrazolidinyl, piperazinyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, thiophenyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, quinolinyl, cinnolinyl, quinoxalinyl, benzimidazolyl, benzothiophenyl, or benzothiadiazolyl, each of which is optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₄
20 haloalkyl, C₁-C₄ alkoxy and C₁-C₄ haloalkoxy.

More preferably, R⁷ represents C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, phenyl, naphthyl, anthracyl, fluorenyl, indanyl, morpholinyl, pyrrolidinyl, piperidinyl, piperazinyl, pyridyl, pyrrolyl, imidazolyl and triazolyl, each of which is optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₄ haloalkyl, C₁-C₄ alkoxy and
30 C₁-C₄ haloalkoxy.

Yet more preferably, R⁷ represents C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, phenyl, naphthyl, morpholinyl, pyrrolidinyl, piperidinyl, piperazinyl, pyridyl, pyrrolyl, imidazolyl and triazolyl, each of which is optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₄ haloalkyl, C₁-C₄ alkoxy and C₁-C₄
35 haloalkoxy.

- 9 -

Even more preferably, R^7 represents C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, phenyl, morpholinyl, pyrrolidinyl, piperdiny, piperazinyl, pyridyl, pyrrolyl, imidazolyl and triazolyl, each of which is optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy and C_1 - C_4 haloalkoxy.

5 More preferably again, R^7 represents C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, phenyl, morpholinyl, pyrrolidinyl, pyridyl and imidazolyl, each of which is optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy and C_1 - C_4 haloalkoxy.

10 Most preferably, R^7 represents C_3 - C_6 cycloalkyl or C_3 - C_5 cycloalkenyl, each of which is optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy and C_1 - C_4 haloalkoxy

In one group of compounds of formula (I), R^1 represents hydrogen, halogen, cyano, OH, NH_2 , C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, $NH(C_1$ - C_4 alkyl), $N(C_1$ - C_4 alkyl) $_2$, $CO(C_1$ - C_4 alkyl), $CO_2(C_1$ - C_4 alkyl), CO_2H , $CONH(C_1$ - C_4 alkyl), $CON(C_1$ - C_4 alkyl) $_2$, $SO_2NH(C_1$ - C_4 alkyl),
15 $SO_2N(C_1$ - C_4 alkyl) $_2$, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, or C_2 - C_4 alkynyl;

R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 haloalkyl and C_1 - C_4 haloalkoxy;

20 R^3 and R^4 independently of each other represent hydrogen, C_1 - C_4 alkyl or C_3 - C_6 cycloalkyl; or

R^3 and R^4 together with the nitrogen atom to which they are attached form a 3-6-membered saturated cyclic group;

R^5 represents H, C_1 - C_4 alkyl or C_1 - C_4 haloalkyl;

25 R^6 represents C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl or C_1 - C_4 haloalkoxy.

In another group of compounds of formula (I), R^1 represents hydrogen, halogen, cyano, OH, NH_2 , methyl, ethyl, cyclopropyl, $NH(C_1$ - C_2 alkyl), $N(C_1$ - C_2 alkyl) $_2$, $CO(C_1$ - C_2 alkyl), $CO_2(C_1$ - C_2 alkyl), CO_2H , $CONH(C_1$ - C_2 alkyl), $CON(C_1$ - C_2 alkyl) $_2$, $SO_2NH(C_1$ - C_2 alkyl), $SO_2N(C_1$ - C_2 alkyl) $_2$, C_1 - C_4 fluoroalkyl, C_1 - C_4 alkoxy, C_1 - C_2 haloalkoxy or C_2 - C_4 alkynyl;

30 R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_2 haloalkyl and C_1 - C_2 haloalkoxy;

35 R^3 and R^4 independently of each other represent hydrogen, C_1 - C_3 alkyl or C_3 - C_5 cycloalkyl; or

- 10 -

R^3 and R^4 together with the nitrogen atom to which they are attached form a 5-membered saturated cyclic group;

R^5 represents H or C_1 - C_4 alkyl;

R^6 represents C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl or C_1 - C_4 haloalkoxy.

5 In another group of compounds of formula (I), R^1 represents hydrogen, halogen, cyano, OH, NH_2 , methyl, ethyl, cyclopropyl, $NHMe$, NMe_2 , $COMe$, CO_2Me , CO_2H , $CONHMe$, $CONMe_2$, SO_2NHMe , SO_2NMe_2 , CHF_2 , CF_3 , OMe , $OCHF_2$ or acetylenyl;

R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, C_3 - C_6 cycloalkyl, C_3 - C_5 cycloalkenyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkyl, $-C_1$ - C_2 alkyl- C_3 - C_6 cycloalkenyl which may be
10 optionally substituted by one or more groups independently selected from the group consisting fluoro, C_1 - C_2 alkyl, C_1 - C_2 fluoroalkyl and C_1 - C_2 fluoroalkoxy;

R^3 and R^4 independently of each other represent hydrogen, methyl, ethyl, isopropyl or cyclopropyl; or

R^3 and R^4 together with the nitrogen atom to which they are attached form a 5-
15 membered saturated cyclic group;

R^5 represents H or methyl;

R^6 represents C_1 - C_4 alkyl or C_1 - C_4 alkoxy.

In another group of compounds of formula (I), R_1 represents hydrogen, halogen, cyano, methyl, ethyl, cyclopropyl, CHF_2 , CF_3 , OMe , or $OCHF_2$;

20 R_2 is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, 2-methylbutyl, 2,2-dimethylpropyl, 2-methylpentyl, 3-methylpentyl, 2,3-dimethylbutyl, 2,2-dimethylbutyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl $-CH_2$ -cyclopropyl, $-CH_2$ -cyclobutyl, $-CH_2$ -cyclopentyl, and $-CH_2$ -cyclopentenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro, methyl and
25 difluoromethoxy;

R^3 is hydrogen or methyl;

R^4 is methyl or ethyl;

R^5 represents H or methyl;

R^6 represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl,
30 methoxy or ethoxy.

In another group of compounds of formula (I), R^1 represents hydrogen, Cl, Br, methyl, CHF_2 or cyano;

R_2 is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro
35 and difluoromethoxy;

- 11 -

R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl,

5 methoxy or ethoxy.

In another group of compounds of formula (I), R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro and difluoromethoxy;

10

R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl or cyclopentyl.

15 In another group of compounds of formula (I), R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro and difluoromethoxy;

20 R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ represents methyl, ethyl, n-propyl or iso-propyl.

25 In another group of compounds of formula (I), R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro and difluoromethoxy;

R³ represents hydrogen or methyl;

30 R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ is methyl.

In another group of compounds of formula (I), R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

35 R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or cyclopropyl;

- 12 -

R^3 represents hydrogen or methyl;


R^4 is ethyl;

R^5 is hydrogen;

R^6 is methyl.

5 Tables 1 to 22: Compounds of formula (I)

The invention is further illustrated by making available the following individual compounds of formula (I) listed below in Tables 1 to 22.




Each of Tables 1 to 22, which follow Table A below, make available 47 compounds of
 10 the formula (I) in which R^2 , R^3 , R^4 , R^5 and R^6 are the substituents defined in Table A and R^1 is the substituent defined in the relevant Table 1 to 22. Thus Table 1 individualises 47 compounds of formula (I) wherein for each row of Table A, R^1 is as defined in Table 1. Similarly, Table 2 individualises 47 compounds of formula (I) wherein for each row of Table A, R^1 is as defined in Table 2; and so on for Tables 3 to 22.

15 Table A discloses 47 sets of meanings of the variables R^2 , R^3 , R^4 , R^5 and R^6 in a compound of formula I.

Tables 23 to 44: Compounds of formula (Ia)

The invention is further illustrated by making available the following individual compounds of formula (Ia) listed below in Tables 23 to 44.



(Ia)

20

Each of Tables 23 to 44, which follow Table A below, make available 47 compounds of the formula (Ia) in which R^2 , R^3 , R^4 , R^5 and R^6 are the substituents defined in Table A and R^1 is the substituent defined in the relevant Table 23 to 44. Thus Table 23 individualises 47 compounds of formula (Ia) wherein for each row of Table A, R^1 is as defined in Table 23.


- 13 -

Similarly, Table 24 individualises 47 compounds of formula (Ia) wherein for each row of Table A, R¹ is as defined in Table 24; and so on for Tables 25 to 44.

Table A discloses 47 sets of meanings of the variables R², R³, R⁴, R⁵ and R⁶ in a compound of formula Ia.

5 Tables 45 to 66: Compounds of formula (Ib)

The invention is further illustrated by making available the following individual compounds of formula (Ib) listed below in Tables 45 to 66.



Each of Tables 45 to 66, which follow Table A below, make available 47 compounds of the formula (Ib) in which R², R³, R⁴, R⁵ and R⁶ are the substituents defined in Table A and R¹ is the substituent defined in the relevant Table 45 to 66. Thus Table 45 individualises 47 compounds of formula (Ib) wherein for each row of Table A, R¹ is as defined in Table 45. Similarly, Table 46 individualises 47 compounds of formula (Ib) wherein for each row of Table A, R¹ is as defined in Table 46; and so on for Tables 47 to 66.

15 Table A discloses 47 sets of meanings of the variables R², R³, R⁴, R⁵ and R⁶ in a compound of formula Ib.

Table A


	R ²	R ³	R ⁴	R ⁵	R ⁶
A.1.001	n-propyl	ethyl	ethyl	H	methyl
A.1.002	n-propyl	methyl	cyclopropyl	H	methyl
A.1.003	n-propyl	methyl	isopropyl	H	methyl
A.1.004	n-propyl	-CH ₂ -CH ₂ -CH ₂ -CH ₂ -		H	methyl
A.1.005	n-propyl	methyl	methyl	H	methyl
A.1.006	n-propyl	ethyl	H	H	methyl
A.1.007	n-propyl	methyl	H	H	methyl
A.1.008	n-propyl	isopropyl	H	H	methyl
A.1.09	prop-2-enyl	ethyl	ethyl	H	methyl
A.1.010	prop-2-enyl	methyl	cyclopropyl	H	methyl
A.1.011	prop-2-enyl	methyl	isopropyl	H	methyl
A.1.012	prop-2-enyl	-CH ₂ -CH ₂ -CH ₂ -CH ₂ -		H	methyl
A.1.013	prop-2-enyl	methyl	methyl	H	methyl
A.1.014	prop-2-enyl	ethyl	H	H	methyl
A.1.015	prop-2-enyl	isopropyl	H	H	methyl
A.1.016	prop-2-enyl	ethyl	methyl	H	methyl
A.1.017	isopropyl	ethyl	methyl	H	methyl
A.1.018	isopropyl	ethyl	methyl	methyl	methyl
A.1.019	propargyl	ethyl	methyl	H	methyl
A.1.020	n-propyl	ethyl	methyl	H	methyl
A.1.021	prop-2-enyl	ethyl	methyl	methyl	methyl
A.1.022	n-propyl	ethyl	methyl	methyl	methyl
A.1.023	propargyl	ethyl	methyl	methyl	methyl
A.1.024	3,3-difluoropropyl	ethyl	methyl	H	methyl
A.1.025	3,3,3-trifluoropropyl	ethyl	methyl	methyl	methyl
A.1.026	3,3,3-trifluoropropyl	ethyl	methyl	H	methyl
A.1.027	3,3-difluoroprop-2-enyl	ethyl	methyl	H	methyl
A.1.028	n-propyl	ethyl	methyl	H	ethyl
A.1.029	prop-2-enyl	ethyl	methyl	H	ethyl
A.1.030	propargyl	ethyl	methyl	H	ethyl
A.1.031	n-propyl	ethyl	methyl	H	methoxymethyl
A.1.032	prop-2-enyl	ethyl	methyl	H	methoxymethyl
A.1.033	propargyl	ethyl	methyl	H	methoxymethyl
A.1.034	2,2-dimethyl-propyl	ethyl	methyl	H	methyl
A.1.035	2-methyl-butyl	ethyl	methyl	H	methyl
A.1.036	2,2-dimethyl-butyl	ethyl	methyl	H	methyl
A.1.037	cyclopropyl	ethyl	methyl	H	methyl
A.1.038	cyclobutyl	ethyl	methyl	H	methyl
A.1.039	cyclopentyl	ethyl	methyl	H	methyl

- 15 -


	R²	R³	R⁴	R⁵	R⁶
A.1.040	cyclopent-3-enyl	ethyl	methyl	H	methyl
A.1.041	3-methyl-but-2-enyl	ethyl	methyl	H	methyl
A.1.042	but-2-enyl	ethyl	methyl	H	methyl
A.1.043	cyclopropyl-methyl	ethyl	methyl	H	methyl
A.1.044	cyclobutyl-methyl	ethyl	methyl	H	methyl
A.1.045	cyclopentyl-methyl	ethyl	methyl	H	methyl
A.1.046	cyclopent-3-enyl-methyl	ethyl	methyl	H	methyl
A.1.047	n-propyl	-CH ₂ -CH ₂ -CH ₂ -		H	methyl

Tables 1, 23 and 45: Tables 1, 23 and 45 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is hydrogen and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A. For example;

compound 1.1.016 has the following structure:




compound 23.1.043 has the following structure:



compound 45.1.043 has the following structure:

- 16 -



Tables 2, 24 and 46: Tables 2, 24 and 46 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is chloro and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

Tables 3, 25 and 47: Tables 3, 25 and 47 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is bromo and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

Tables 4, 26 and 48: Tables 4, 26 and 48 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is iodo and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

Tables 5, 27 and 49: Tables 5, 27 and 49 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is methyl and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

Tables 6, 28 and 50: Tables 6, 28 and 50 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is ethyl and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

Tables 7, 29 and 51: Tables 7, 29 and 51 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is cyclopropyl and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

Tables 8, 30 and 52: Tables 8, 30 and 52 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is trifluoromethyl and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

- 17 -

Tables 9, 31 and 53: Tables 9, 31 and 53 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is difluoromethyl and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

5 Tables 10, 32 and 54: Tables 10, 32 and 54 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is ethynyl and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.


10 Tables 11, 33 and 55: Tables 11, 33 and 55 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is methoxy and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

15 Tables 12, 34 and 56: Tables 12, 34 and 56 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is difluoromethoxy and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

20 Tables 13, 35 and 57: Tables 13, 35 and 57 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is methylamino and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.

25 Tables 14, 36 and 58: Tables 14, 36 and 58 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is dimethylamino and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.


30 Tables 15, 37 and 59: Tables 15, 37 and 59 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is N-methylsulfonamide, wherein the broken line below indicates the point of attachment of the group R¹ to the remainder of the compound and each of the variables R², R³, R⁴, R⁵ and R⁶ has the specific meaning given in the corresponding row of Table A.



Tables 16, 38 and 60: Tables 16, 38 and 60 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R¹ is N,N-

- 18 -

dimethylsulfonamide, wherein the broken line below indicates the point of attachment of the group R^1 to the remainder of the compound and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.




5

Tables 17, 39 and 61: Tables 17, 39 and 61 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R^1 is fluoro and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.

10 Tables 18, 40 and 62: Tables 18, 40 and 62 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R^1 is acetyl and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.

Tables 19, 41 and 63: Tables 19, 41 and 63 disclose 47 compounds of formula (I), 47
15 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R^1 is methoxyformyl wherein the broken line below indicates the point of attachment of the group R^1 to the remainder of the compound and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.



20 Tables 20, 42 and 64: Tables 20, 42 and 64 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R^1 is hydroxy and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.

Tables 21, 43 and 65: Tables 21, 43 and 65 disclose 47 compounds of formula (I), 47
25 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R^1 is fluoromethyl and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.

Tables 22, 44 and 66: Tables 22, 44 and 66 disclose 47 compounds of formula (I), 47 compounds of formula (Ia) and 47 compounds of formula (Ib) respectively wherein R^1 is

- 19 -

cyano and each of the variables R^2 , R^3 , R^4 , R^5 and R^6 has the specific meaning given in the corresponding row of Table A.

It has now been found that the compounds of formula (I) according to the invention have, for practical purposes, a very advantageous spectrum of activities for protecting useful plants against diseases that are caused by phytopathogenic microorganisms, such as fungi, bacteria or viruses, in particular against diseases that are caused by fungi.

The invention therefore also relates to a method of controlling or preventing infestation of useful plants by phytopathogenic microorganisms, wherein a compound of formula (I) is applied as active ingredient to the plants, to parts thereof or the locus thereof.

The term "plants" refers to all physical parts of a plant, including seeds, seedlings, saplings, roots, tubers, stems, stalks, foliage, and fruits.

The term "locus" as used herein means fields in or on which plants are growing, or where seeds of cultivated plants are sown, or where seed will be placed into the soil. It includes soil, seeds, and seedlings, as well as established vegetation.

The compounds of formula I can be used in the agricultural sector and related fields of use e.g. as active ingredients for controlling plant pests or on non-living materials for control of spoilage microorganisms or organisms potentially harmful to man.

The compounds of formula (I) according to the invention are distinguished by excellent activity at low rates of application, by being well tolerated by plants and by being environmentally safe. They have very useful curative, preventive and systemic properties and are used for protecting numerous useful plants. The compounds of formula (I) can be used to inhibit or destroy the diseases that occur on plants or parts of plants of different crops of useful plants, while at the same time protecting also those parts of the plants that grow later from phytopathogenic micro-organisms.

It is possible to use compounds of formula I as fungicide. The term "fungicide" as used herein means a compound that controls, modifies, or prevents the growth of fungi. The term "fungicidally effective amount" means the quantity of such a compound or combination of such compounds that is capable of producing an effect on the growth of fungi. Controlling or modifying effects include all deviation from natural development, such as killing, retardation and the like, and prevention includes barrier or other defensive formation in or on a plant to prevent fungal infection.

A preferred method of applying a compound of formula (I) is foliar application. The frequency of application and the rate of application will depend on the risk of infestation by the corresponding pathogen. However, the compounds of formula (I) may also penetrate the plant through the roots via the soil (systemic action) by drenching the locus of the plant with a

- 20 -

liquid formulation, or by applying the compounds in solid form to the soil, e.g. in granular form (soil application). In crops of water rice such granulates can be applied to the flooded rice field.

5 It is also possible to use compounds of formula (I) as dressing agents for the treatment of plant propagation material, in particular of seeds and plant cuttings (e.g. rice), for the protection against fungal infections as well as against phytopathogenic fungi occurring in the soil.

10 The propagation material can be treated with a composition comprising a compound of formula I before planting: seed, for example, can be dressed before being sown. The active ingredients according to the invention can also be applied to grains (coating), either by impregnating the seeds in a liquid formulation or by coating them with a solid formulation. The composition can also be applied to the planting site when the propagation material is being planted, for example, to the seed furrow during sowing. The invention relates also to
15 such methods of treating plant propagation material and to the plant propagation material so treated.

The term "plant propagation material" denotes all generative parts of a plant, for example seeds or vegetative parts of plants such as cuttings and tubers. It includes seeds in the strict sense, as well as roots, fruits, tubers, bulbs, rhizomes, and parts of plants.

20 Furthermore the compounds of formula (I) according to the invention may be used for controlling fungi in related areas, for example in the protection of technical materials, including wood and wood related technical products, in food storage or in hygiene management.

25 In addition, the invention could be used to protect non-living materials from fungal attack, e.g. lumber, wall boards and paint.

The compounds of formula I are for example, effective against Fungi and fungal vectors of disease as well as phytopathogenic bacteria and viruses. These Fungi and fungal vectors of disease as well as phytopathogenic bacteria and viruses are for example:

30 Absidia corymbifera, Alternaria spp, Aphanomyces spp, Ascochyta spp, Aspergillus spp. including A. flavus, A. fumigatus, A. nidulans, A. niger, A. terreus, Aureobasidium spp. including A. pullulans, Blastomyces dermatitidis, Blumeria graminis, Bremia lactucae, Botryosphaeria spp. including B. dothidea, B. obtusa, Botrytis spp. including B. cinerea, Candida spp. including C. albicans, C. glabrata, C. krusei, C. lusitanae, C. parapsilosis, C. tropicalis, Cephalosporium fragrans, Ceratocystis spp, Cercospora spp. including C.
35 arachidicola, Cercosporidium personatum, Cladosporium spp, Claviceps purpurea,

- 21 -

Coccidioides immitis, Cochliobolus spp, Colletotrichum spp. including C. musae, Cryptococcus neoformans, Diaporthe spp, Didymella spp, Drechslera spp, Elsinoe spp, Epidermophyton spp, Erwinia amylovora, Erysiphe spp. including E. cichoracearum, Eutypa lata, Fusarium spp. including F. culmorum, F. graminearum, F. langsethiae, F. moniliforme, 5 F. oxysporum, F. proliferatum, F. subglutinans, F. solani, Gaeumannomyces graminis, Gibberella fujikuroi, Gloeodes pomigena, Gloeosporium musarum, Glomerella cingulate, Guignardia bidwellii, Gymnosporangium juniperi-virginianae, Helminthosporium spp, Hemileia spp, Histoplasma spp. including H. capsulatum, Laetisaria fuciformis, Leptographium lindbergi, Leveillula taurica, Lophodermium seditiosum, Microdochium nivale, 10 Microsporum spp, Monilinia spp, Mucor spp, Mycosphaerella spp. including M. graminicola, M. pomi, Oncobasidium theobromaeon, Ophiostoma piceae, Paracoccidioides spp, Penicillium spp. including P. digitatum, P. italicum, Petriellidium spp, Peronosclerospora spp. Including P. maydis, P. philippinensis and P. sorghi, Peronospora spp, Phaeosphaeria nodorum, Phakopsora pachyrhizi, Phellinus igniarius, Phialophora spp, Phoma spp, 15 Phomopsis viticola, Phytophthora spp. including P. infestans, Plasmopara spp. including P. halstedii, P. viticola, Pleospora spp., Podosphaera spp. including P. leucotricha, Polymyxa graminis, Polymyxa betae, Pseudocercospora herpotrichoides, Pseudomonas spp, Pseudoperonospora spp. including P. cubensis, P. humuli, Pseudopeziza tracheiphila, Puccinia Spp. including P. hordei, P. recondita, P. striiformis, P. tritici, Pyrenopeziza spp, 20 Pyrenophora spp, Pyricularia spp. including P. oryzae, Pythium spp. including P. ultimum, Ramularia spp, Rhizoctonia spp, Rhizomucor pusillus, Rhizopus arrhizus, Rhynchosporium spp, Scedosporium spp. including S. apiospermum and S. prolificans, Schizothyrium pomi, Sclerotinia spp, Sclerotium spp, Septoria spp, including S. nodorum, S. tritici, Sphaerotheca macularis, Sphaerotheca fusca (Sphaerotheca fuliginea), Sporothrix spp, Stagonospora 25 nodorum, Stemphylium spp, Stereum hirsutum, Thanatephorus cucumeris, Thielaviopsis basicola, Tilletia spp, Trichoderma spp. including T. harzianum, T. pseudokoningii, T. viride, Trichophyton spp, Typhula spp, Uncinula necator, Urocystis spp, Ustilago spp, Venturia spp. including V. inaequalis, Verticillium spp, and Xanthomonas spp.

Crops of useful plants in which the composition according to the invention can be used 30 include perennial and annual crops, such as berry plants for example blackberries, blueberries, cranberries, raspberries and strawberries; cereals for example barley, maize (corn), millet, oats, rice, rye, sorghum tritcale and wheat; fibre plants for example cotton, flax, hemp, jute and sisal; field crops for example sugar and fodder beet, coffee, hops, mustard, oilseed rape (canola), poppy, sugar cane, sunflower, tea and tobacco; fruit trees for example 35 apple, apricot, avocado, banana, cherry, citrus, nectarine, peach, pear and plum; grasses for

- 22 -

example Bermuda grass, bluegrass, bentgrass, centipede grass, fescue, ryegrass, St. Augustine grass and Zoysia grass; herbs such as basil, borage, chives, coriander, lavender, lovage, mint, oregano, parsley, rosemary, sage and thyme; legumes for example beans, lentils, peas and soya beans; nuts for example almond, cashew, ground nut, hazelnut, peanut, pecan, pistachio and walnut; palms for example oil palm; ornamentals for example flowers, shrubs and trees; other trees, for example cacao, coconut, olive and rubber; vegetables for example asparagus, aubergine, broccoli, cabbage, carrot, cucumber, garlic, lettuce, marrow, melon, okra, onion, pepper, potato, pumpkin, rhubarb, spinach and tomato; and vines for example grapes.

Crops are to be understood as being those which are naturally occurring, obtained by conventional methods of breeding, or obtained by genetic engineering. They include crops which contain so-called output traits (e.g. improved storage stability, higher nutritional value and improved flavour).

Crops are to be understood as also including those crops which have been rendered tolerant to herbicides like bromoxynil or classes of herbicides such as ALS-, EPSPS-, GS-, HPPD- and PPO-inhibitors. An example of a crop that has been rendered tolerant to imidazolinones, e.g. imazamox, by conventional methods of breeding is Clearfield® summer canola. Examples of crops that have been rendered tolerant to herbicides by genetic engineering methods include e.g. glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady®, Herculex I® and LibertyLink®.

Crops are also to be understood as being those which naturally are or have been rendered resistant to harmful insects. This includes plants transformed by the use of recombinant DNA techniques, for example, to be capable of synthesising one or more selectively acting toxins, such as are known, for example, from toxin-producing bacteria. Examples of toxins which can be expressed include δ -endotoxins, vegetative insecticidal proteins (Vip), insecticidal proteins of bacteria colonising nematodes, and toxins produced by scorpions, arachnids, wasps and fungi.

An example of a crop that has been modified to express the *Bacillus thuringiensis* toxin is the Bt maize KnockOut® (Syngenta Seeds). An example of a crop comprising more than one gene that codes for insecticidal resistance and thus expresses more than one toxin is VipCot® (Syngenta Seeds). Crops or seed material thereof can also be resistant to multiple types of pests (so-called stacked transgenic events when created by genetic modification). For example, a plant can have the ability to express an insecticidal protein while at the same

- 23 -

time being herbicide tolerant, for example Herculex I® (Dow AgroSciences, Pioneer Hi-Bred International).

The compounds of formula (I) can be used in unmodified form or, preferably, together with carriers and adjuvants conventionally employed in the art of formulation.

5 Therefore the invention also relates to compositions for controlling and protecting against phytopathogenic microorganisms, comprising a compound of formula (I) and an inert carrier, and to a method of controlling or preventing infestation of useful plants by phytopathogenic microorganisms, wherein a composition, comprising a compound of formula (I) as active ingredient and an inert carrier, is applied to the plants, to parts thereof or the
10 locus thereof.

To this end compounds of formula (I) and inert carriers are conveniently formulated in known manner to emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations e.g. in polymeric substances. As with the type of the compositions, the
15 methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. The compositions may also contain further adjuvants such as stabilizers, antifoams, viscosity regulators, binders or tackifiers as well as fertilizers, micronutrient donors or other formulations for obtaining special effects.

20 Suitable carriers and adjuvants (auxiliaries) can be solid or liquid and are substances useful in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilizers. Such carriers are for example described in WO 97/33890.

A formulation, i.e. a composition comprising the compound of formula (I) and, if
25 desired, a solid or liquid adjuvant, is prepared in a known manner, typically by intimately mixing and/or grinding the compound with extenders, for example solvents, solid carriers and, optionally, surface-active compounds (surfactants).

The agrochemical formulations will usually contain from 0.1 to 99% by weight, preferably from 0.1 to 95% by weight, of the compound of formula (I), 99.9 to 1% by weight, preferably 99.8 to 5% by weight, of a solid or liquid adjuvant, and from 0 to 25% by weight, preferably from 0.1 to 25% by weight, of a surfactant.
30

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.

- 24 -

Advantageous rates of application are normally from 5g to 2kg of active ingredient (a.i.) per hectare (ha), preferably from 10g to 1kg a.i./ha, most preferably from 20g to 600g a.i./ha. When used as seed drenching agent, convenient rates of application are from 10mg to 1g of active substance per kg of seeds. The rate of application for the desired action can be
5 determined by experiments. It depends for example on the type of action, the developmental stage of the useful plant, and on the application (location, timing, application method) and can, owing to these parameters, vary within wide limits.

Normally, in the management of a crop a grower would use one or more other agronomic chemicals in addition to the compound of the present invention. Examples of
10 agronomic chemicals include pesticides, such as acaricides, bactericides, fungicides, herbicides, insecticides, nematocides, as well as plant nutrients and plant fertilizers.

Accordingly, the present invention provides a composition comprising a compound of formula (I) according to the present invention together with one or more pesticides, plant nutrients or plant fertilizers. The combination may also encompass specific plant traits
15 incorporated into the plant using any means, for example conventional breeding or genetic modification. Such compositions may also contain one or more inert carriers as described above.

The invention also provides for the use of provides a composition comprising a compound of formula (I) according to the present invention together with one or more
20 pesticides, plant nutrients or plant fertilizers. The combination may also encompass specific plant traits incorporated into the plant using any means, for example conventional breeding or genetic modification.

Suitable examples of plant nutrients or plant fertilizers are calcium sulfate (CaSO_4), calcium nitrate ($\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$), calcium carbonate (CaCO_3), potassium nitrate (KNO_3),
25 magnesium sulfate (MgSO_4), potassium hydrogen phosphate (KH_2PO_4), manganese sulfate (MnSO_4), copper sulfate (CuSO_4), zinc sulfate (ZnSO_4), nickel chloride (NiCl_2), cobalt sulfate (CoSO_4), potassium hydroxide (KOH), sodium chloride (NaCl), boric acid (H_3BO_3) and metal salts thereof (Na_2MoO_4). The nutrients may be present in an amount of 5% to 50% by weight, preferably of 10% to 25% by weight or of 15% to 20% by weight each. Preferred additional
30 nutrients are urea ($(\text{NH}_2)_2\text{CO}$), melamine ($\text{C}_3\text{H}_6\text{N}_6$), potassium oxide (K_2O), and inorganic nitrates. The most preferred additional plant nutrient is potassium oxide. Where the preferred additional nutrient is urea, it is present in an amount of generally 1% to 20% by weight, preferably 2% to 10% by weight or of 3% to 7% by weight.

Suitable examples of pesticides are acycloamino acid fungicides, aliphatic nitrogen
35 fungicides, amide fungicides, anilide fungicides, antibiotic fungicides, aromatic fungicides,

- 25 -

arsenical fungicides, aryl phenyl ketone fungicides, benzamide fungicides, benzanilide fungicides, benzimidazole fungicides, benzothiazole fungicides, botanical fungicides, bridged diphenyl fungicides, carbamate fungicides, carbanilate fungicides, conazole fungicides, copper fungicides, dicarboximide fungicides, , dinitrophenol fungicides, dithiocarbamate
5 fungicides, dithiolane fungicides, furamide fungicides, furanilide fungicides, hydrazide fungicides, imidazole fungicides, mercury fungicides, morpholine fungicides, organophosphorous fungicides, organotin fungicides, oxathiin fungicides, oxazole fungicides, phenylsulfamide fungicides, polysulfide fungicides, pyrazole fungicides, pyridine fungicides, pyrimidine fungicides, pyrrole fungicides, quaternary ammonium fungicides, quinoline
10 fungicides, quinone fungicides, quinoxaline fungicides, strobilurin fungicides, sulfonanilide fungicides, thiadiazole fungicides, thiazole fungicides, thiazolidine fungicides, thiocarbamate fungicides, thiophene fungicides, triazine fungicides, triazole fungicides, triazolopyrimidine fungicides, urea fungicides, valinamide fungicides, zinc fungicides, Benzoylureas, carbamates, chloronicotinylns, diacylhydrazines, diamides, fiproles, macrolides, nitroimines,
15 nitromethylenes, organochlorines, organophosphates, organosilicons, organotins, phenylpyrazoles, phosphoric esters, pyrethroids, spinosyns, tetramic acid derivatives, tetronic acid derivatives, Antibiotic nematocides, avermectin nematocides, botanical nematocides, carbamate nematocides, oxime carbamate nematocides, organophosphorus nematocides, nematophagous fungi or bacteria, amide herbicides, anilide herbicides,
20 arsenical herbicides, arylalanine herbicides, aryloxyphenoxypropionic herbicides, benzofuranyl herbicides, benzoic acid herbicides, benzothiazole herbicides, benzoylcyclohexanedione herbicides, carbamate herbicides, carbanilate herbicides, chloroacetanilide herbicides, chlorotriazine herbicides, cyclohexene oxmie herbicides, cyclopropylisoxazole herbicides, dicarboximide herbicides, dinitroaniline herbicides,
25 dinitrophenol herbicides, diphenyl ether herbicides, dithiocarbamate herbicides, fluoroalkyltriazine herbicides, halogenated aliphatic herbicides, imidazolinone herbicides, inorganic herbicides, methoxytriazine herbicides, methylthiotriazine herbicides, nitrile herbicides, nitrophenyl ether herbicides, organophosphorous herbicides, oxadiazolone herbicides, oxazole herbicides, phenoxy herbicides, phenoxyacetic herbicides,
30 phenoxybutyric herbicides, phenoxypropionic herbicides, phenylenediamine herbicides, phenylurea herbicides, phthalic acid herbicides, picolinic acid herbicides, pyrazole herbicides, pyridazine herbicides, pyridazinone herbicides, pyridine herbicides, pyrimidinediamine herbicides, pyrimidinylloxybenzylamine herbicides, pyrimidinylsulfonylurea herbicides, quaternary ammonium herbicides, quinolinecarboxylic acid herbicides,
35 sulfonamide herbicides, sulfonanilide herbicides, sulfonylurea herbicides, thiadiazolylurea

- 26 -


herbicides, thioamide herbicides, thiocarbamate herbicides, thiocarbonate herbicides, thiourea herbicides, triazine herbicides, triazinone herbicides, triazinylsulfonylurea herbicides, triazole herbicides, triazolone herbicides, triazolopyrimidine herbicides, uracil herbicides, urea herbicides, microbials, plant extracts, pheromones, macrobials and other biologicals.

5 The following non-limiting Examples illustrate the above-described invention in greater detail without limiting it. Those skilled in the art will promptly recognise appropriate variations from the procedures both as to reactants and as to reaction conditions and techniques. All references mentioned herein are incorporated by reference in their entirety.

10 **SYNTHETIC EXAMPLES**

Using techniques analogous to those described in WO 12/146125 (pp.370-378) and further techniques known to the person skilled in the art, for example as found in WO 08/101682 (pp.22-33), compounds of formula (I) may be prepared.

Preparation of (2R)-1-propoxypropan-2-ol




To an ice-bath cooled solution of THF (400 mL) under inert atmosphere (Ar) and sodium hydride (12 g, 490 mmol, 5 equiv.) 1-propanol (40 mL, 490 mmol, 5 equiv) was added dropwise. The ice bath was removed and the reaction mixture was stirred at room temperature for 30 minutes then (2R)-2-methyloxirane (5.8 g, 99 mmol) was added dropwise and the reaction was stirred for 18h under heating at 50°C. After this time, GC-MS and NMR indicated that the starting material was consumed and the reaction mixture was allowed to reach room temperature before quenching with aqueous NH₄Cl solution and extracting with dichloromethane. The organic layer was dried over anhydrous Na₂SO₄ and filtered. The solvent was removed in vacuo (not dropping below 200 mbar) at 30°C and provided the title compound (4.4 g, 38% yield) as a yellow liquid.

¹H NMR (400 MHz, CDCl₃): δ (ppm) 4.00 – 3.87 (1H, m) 3.50 – 3.40 (m, 3H), 3.30 – 3.20 (m, 1H), 2.64 (d, 1H), 1.61 (m, 2H), 1.12 (d, 3H), 0.95 (t, 3H)

30 Preparation of N'-[5-bromo-2-methyl-6-[(1S)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine


- 27 -



To a stirring suspension of N'-(5-bromo-6-hydroxy-2-methyl-3-pyridyl)-N-ethyl-N-methyl-formamidinium (0.75 g, 2.8 mmol) in THF (15 mL), (2R)-1-propoxypropan-2-ol (0.36 g, 3 mmol, 1.1 equiv) and triphenylphosphine (0.80 g, 3 mmol, 1.1 equiv) were added at room temperature under inert atmosphere (Ar). To this mixture, DIAD (diisopropyl diazodicarboxylate) (0.60 mL, 3 mmol, 1.1 equiv) was added dropwise over 10 minutes while keeping the temperature below 40°C. The reaction mixture was stirred for 24h at room temperature. After this time, LC-MS indicated that the starting material had been nearly consumed and the reaction mixture was quenched with water (40 mL). The water phase was extracted with ethyl acetate (3x50 mL). The organic layer was dried over anhydrous Na₂SO₄ and filtered. The solvent was removed in vacuo to give a brown residue, which was purified by preparative reverse phase chromatography to afford the desired (0.10 g, 10% yield).

¹H NMR (400 MHz, CDCl₃): δ (ppm) 7.45 – 7.30 (broad s, 1H), 7.24 (s, 1H), 5.40 – 5.30 (m, 1H), 3.70 – 3.60 (m, 1H), 3.55 – 3.45 (m, 3H), 3.45 – 3.30 (broad m, 2H), 3.00 (s, 3H), 2.35 (s, 3H), 1.65 – 1.50 (m, 2H), 1.35 (m, 3H), 1.20 (m, 3H), 0.90 (t, 3H).

Preparation of N'-[5-bromo-2-methyl-6-[(1R)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidinium




To an ice-bath cooled solution of (2R)-1-propoxypropan-2-ol (0.103 g, 0.88 mmol, 1.2 equiv) in DMF (4 mL) under inert atmosphere (Ar), potassium tertbutoxide (0.25 g, 2.19 mmol, 3 equiv) and triphenylphosphine (0.14 g, 0.55 mmol, 1.5 equiv) was added. The reaction mixture was stirred for 20 minutes before N'-(5-bromo-6-fluoro-2-methyl-3-pyridyl)-N-ethyl-N-methyl-formamidinium (0.20 g, 0.73 mmol) was added. The reaction mixture was stirred for 4h at room temperature and was quenched with water upon completion. The water phase was extracted with ethyl acetate (2x50 mL). The organic layers were combined, washed with water (3x50 mL), dried over anhydrous Na₂SO₄, and filtered. The solvent was

- 28 -

removed in vacuo to give a brown residue, which was purified by preparative reverse phase chromatography to afford the desired compound (0.130 g, 13% yield).

^1H NMR (400 MHz, CDCl_3): δ (ppm) 7.45 – 7.30 (broad s, 1H), 7.24 (s, 1H), 5.40 – 5.30 (m, 1H), 3.70 – 3.60 (m, 1H), 3.55 – 3.45 (m, 3H), 3.50 – 3.30 (broad m, 2H), 3.00 (s, 3H), 2.35 (s, 3H), 1.65 – 1.50 (m, 2H), 1.35 (m, 3H), 1.20 (m, 3H), 0.90 (t, 3H).


Preparation of N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine



To a stirring suspension of N'-(5-bromo-6-hydroxy-2-methyl-3-pyridyl)-N-ethyl-N-methyl-formamidine (6.0 g, 22.05 mmol) in THF (30 mL), 1-propoxypropan-2-ol (3.53 mL, 26.46 mmol, 1.2 equiv) and triphenylphosphine (6.94 g, 26.46 mmol, 1.2 equiv) were added at room temperature under inert atmosphere (Ar). To this mixture, DIAD (diisopropyl diazodicarboxylate) (5.21 mL, 26.46 mmol, 1.2 equiv) was added dropwise over 10 minutes while keeping the temperature below 40°C. The reaction mixture was stirred for 1.5 h at room temperature. After this time, LC-MS indicated that the starting material had been consumed and the reaction mixture was concentrated in vacuo. Heptane was added to the residue and the mixture was cooled with an ice bath to recrystallize triphenylphosphine oxide. The brown residue was purified by combiflash column chromatography (silica gel, heptane/ethyl acetate, v/v = 90/10 to 4/1). Fractions containing the pure compound were collected and concentrated in vacuo to give the title compound (7.80 g, 95% yield) as a light yellow oil.

^1H NMR (400 MHz, CDCl_3): δ (ppm) 7.45 – 7.30 (broad s, 1H), 7.24 (s, 1H), 5.40 – 5.30 (m, 1H), 3.70 – 3.60 (m, 1H), 3.55 – 3.45 (m, 3H), 3.45 – 3.30 (broad m, 2H), 3.00 (s, 3H), 2.35 (s, 3H), 1.65 – 1.50 (m, 2H), 1.35 (m, 3H), 1.20 (m, 3H), 0.90 (t, 3H).

Preparation of N'-[5-cyano-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine




- 29 -

To a stirred solution of N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidinium (0.25 g, 0.67 mmol) in DMF (1 mL) under inert atmosphere (Ar), zinc cyanide (0.087 g, 0.74 mmol, 1.1 equiv) and tetrakis(triphenylphosphine)palladium (0.23 g, 0.20 mmol, 0.3 equiv) were added and the reaction mixture was stirred for 18h under heating at 120°C. After this time, TLC and LC-MS indicated that the starting material was consumed and the reaction mixture was allowed to reach room temperature before quenching with water. The water phase was extracted with ethyl acetate (2x30 mL). The organic layer was washed with brine (3x50 mL), dried over anhydrous Na₂SO₄ and filtered. The solvent was removed in vacuo to give a brown residue, which was purified by combiflash column chromatography (silica gel, heptane/ethyl acetate, v/v = 90/10 to 70/30). Fractions containing the pure compound were collected and concentrated in vacuo to give the title compound (0.207 g, 97% yield) as a colorless oil.

¹H NMR (400 MHz, CDCl₃): δ (ppm) 7.45 – 7.30 (broad s, 1H), 7.20 (s, 1H), 5.50 – 5.40 (m, 1H), 3.70 – 3.60 (m, 1H), 3.55 – 3.40 (m, 3H), 3.45 – 3.30 (broad m, 2H), 3.00 (s, 3H), 2.40 (s, 3H), 1.65 – 1.50 (m, 2H), 1.35 (m, 3H), 1.20 (m, 3H), 0.90 (t, 3H).


Preparation of 3-chloro-6-methyl-5-nitro-pyridin-2-ol



To an ice-bath cooled suspension of 6-methyl-5-nitro-pyridin-2-ol (0.50 g, 3.24 mmol) in acetonitrile (5 mL) under inert atmosphere (Ar), was added N-chlorosuccinimide (0.43 g, 3.24 mmol, 1 equiv) portionwise. The reaction mixture was stirred for 20h under heating at 67°C. At this time, LC-MS indicated that the starting material was consumed and the reaction mixture was cooled to 0°C and the precipitate was filtered to give the title compound (0.36g, 48% yield) as a beige-white solid.

¹H NMR (400 MHz, CD₃OD): δ (ppm) 8.5 (s, 1H), 2.7 (s, 3H).

Preparation of 5-chloro-2-methyl-3-nitro-6-(o-tolylmethoxy)pyridine




To a stirred solution of o-tolylmethanol (3.99 g, 32.1 mmol, 1.2 equiv) in THF (100 mL), 3-chloro-6-methyl-5-nitro-pyridin-2-ol (5.30g, 26.7 mmol) and triphenylphosphine (8.41 g,

- 30 -

32.1 mmol, 1.2 equiv) were added at room temperature under inert atmosphere (Ar). To this mixture, DIAD (diisopropyl diazodicarboxylate) (6.58 mL, 33.4 mmol, 1.25 equiv) was added dropwise over 10 minutes while keeping the temperature below 40°C. The reaction mixture was stirred for 16 h at room temperature. At this time, LC-MS indicated that the starting material was consumed and the reaction mixture was quenched with water (20 mL). A precipitate formed and was filtered and washed with a mixture of methanol/water (v/v = 5/1), suspended in toluene and concentrated in vacuo to give the title compound (5.28g, 47% yield) as a yellow solid.

¹H NMR (400 MHz, DMSO): δ (ppm) = 8.60 (s, 1H), 7.45 (d, 1H), 7.30 – 7.20 (m, 3H), 5.50 (s, 2H), 2.70 (s, 3H), 2.35 (s, 3H).


Preparation of 6-benzyloxy-5-chloro-2-methyl-pyridin-3-amine



A solution of 2-benzyloxy-3-chloro-6-methyl-5-nitropyridine (250 mg, 0.90 mmol), 10% platinum on carbon (12 mg, 0.062 mmol) in THF (5 mL) was placed under hydrogen (3 equiv., 2.70 mmol) pressure of 3 bar and the reaction was stirred for 18 h at 37°C. After this time, TLC indicated that the starting material has been consumed. The reaction mixture was filtered and the residue was washed with methanol. The organic layer was concentrated to give the title compound (0.216 g, 97% yield) which was used without further purification.

¹H NMR (400 MHz, CDCl₃): δ (ppm) 7.50 - 7.45 (m, 1H), 7.25 - 7.15 (m, 3H), 7.0 (s, 1H), 5.35 (s, 2H), 3.40 – 3.10 (broad s, 2H), 2.42 (s, 3H), 2.30 (s, 3H).

Preparation of N'-(5-chloro-6-hydroxy-2-methyl-3-pyridyl)-N-ethyl-N-methyl-formamidine




To a solution of N-ethyl-N-methyl-formamide (1.29 g, 14.76 mmol, 1.1 equiv) in dichloromethane (70 mL) was added phosphorus oxychloride (1.38 mL, 14.76 mmol, 1.1 equiv). The solution was stirred for 1.5 h at room temperature and then a solution of 5-chloro-2-methyl-6-(o-tolylmethoxy)pyridin-3-amine (3.52 g, 13.41 mmol) in dichloromethane (10 mL)

- 31 -

was added dropwise. After being stirred for 20 h at room temperature the solid was filtered and washed with dichloromethane. The residue was purified by combiflash column chromatography (silica gel, dichloromethane / methanol + 5% triethylamine v/v = 10/0 to 9/1). Fractions containing the compound were collected and concentrated in vacuo to give the title compound (2.52 g, 82 % yield) as a yellow solid.

¹H NMR (400 MHz, CD₃OD): δ (ppm) 7.65 – 7.50 (broad s, 1H), 3.50 – 3.30 (broad s, 1H), 3.0 (s, 2H), 2.25 (s, 3H), 1.35 (m, 3H), 1.25 (m, 3H).


Preparation of N'-(5-chloro-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl)-N-ethyl-N-methyl-formamidine



To a stirred solution of N'-(5-chloro-6-hydroxy-2-methyl-3-pyridyl)-N-ethyl-N-methyl-formamidine (0.25 g, 1.10 mmol) in THF (10 mL), 1-propoxypropan-2-ol (0.14 g, 1.21 mmol, 1.1 equiv) and triphenylphosphine (0.32 g, 1.21 mmol, 1.1 equiv) were added at room temperature under inert atmosphere (Ar). To this mixture, DIAD (diisopropyl diazodicarboxylate) (0.24 mL, 1.21 mmol, 1.1 equiv) was added dropwise over 10 minutes while keeping the temperature below 40°C. The reaction mixture was stirred for 16h under heating at 60°C. After this time, triphenylphosphine (0.15 g, 0.55 mmol, 0.5 equiv) and DIAD (diisopropyl diazodicarboxylate) (0.11 mL, 0.55 mmol, 0.5 equiv) were added again and the reaction mixture was further stirred for 9 h. The reaction mixture was allowed to reach room temperature before quenching with water (10 mL) and 2M NaOH aqueous solution (2 mL). The water phase was extracted with ethyl acetate (1x25 mL). The organic layers were combined, dried over anhydrous Na₂SO₄ and filtered. The solvent was removed in vacuo to give a brown residue, which was purified by combiflash column chromatography (silica gel, heptane/ethyl acetate, v/v = 9/1 to 1/1). Fractions containing the pure compound were collected and concentrated in vacuo to give the title compound (0.19 g, 53% yield) as a yellow oil.


¹H NMR (400 MHz, CDCl₃): δ (ppm) 7.45 – 7.30 (broad s, 1H), 7.05 (s, 1H), 5.40 – 5.30 (m, 1H), 3.70 – 3.60 (m, 1H), 3.55 – 3.40 (m, 3H), 3.45 – 3.30 (broad m, 2H), 3.00 (s, 3H), 2.35 (s, 3H), 1.65 – 1.50 (m, 2H), 1.35 (m, 3H), 1.20 (m, 3H), 0.85 (t, 3H).

- 32 -

Preparation of 5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-nitro-pyridine

To a stirred suspension of 3-bromo-6-methyl-5-nitro-pyridin-2-ol (0.23 g, 1 mmol) in THF (0.08 mL), 1-propoxypropan-2-ol (0.15 g, 1.2 mmol, 1.2 equiv) and triphenylphosphine (0.32 g, 1.2 mmol, 1.2 equiv) were added at room temperature under inert atmosphere (Ar). To this mixture, DIAD (diisopropyl diazodicarboxylate) (0.24 mL, 1.2 mmol, 1.2 equiv) was added dropwise over 10 minutes while keeping the temperature below 40°C. The reaction mixture was stirred for 12h under heating at 65°C. After this time, LC-MS still showed remaining starting material but the reaction mixture was allowed to reach room temperature and the solvent was removed in vacuo to give a brown residue, which was purified by combiflash column chromatography (silica gel, heptane/ triethylamine, v/v = 95/5). Fractions containing the pure compound were collected and concentrated in vacuo to give the title compound (0.20 g, 60% yield) as a beige oil.

¹H NMR (400 MHz, CDCl₃): δ (ppm) 8.45 (s, 1H), 5.50 – 5.40 (m, 1H), 3.65 – 3.60 (m, 1H), 3.65 – 3.50 (m, 1H), 3.50 – 3.35 (m, 2H), 2.70 (s, 3H), 1.50 (m, 2H), 1.30 (m, 3H), 0.80 (t, 3H).

Preparation of 5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)pyridin-3-amine


To a stirred suspension of 5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-nitro-pyridine (0.47 g, 1.41 mmol) in ethanol (10 mL) were added ammonium chloride (0.15 g, 2.82 mmol, 2 equiv), water (2.8 mL) and then iron (0.32 g, 5.64 mmol, 4 equiv). The reaction mixture was stirred for 4h under heating at 85°C. As reaction monitoring still showed a lot of remaining starting material, ammonium chloride (0.75 g, 1.41 mmol, 1 equiv) and iron (0.16 g, 2.82 mmol, 2 equiv) were added and reaction mixture was further stirred for 10h under heating at 85°C. After this time, LC-MS indicated that the starting material has been consumed and the reaction mixture was allowed to reach room temperature before filtering it over celite. The solvent was removed in vacuo and the residue was redissolved with ethyl acetate (15 mL). The organic phase was washed with a 2N aqueous NaOH solution (2x25

- 33 -

mL), dried over anhydrous Na_2SO_4 and filtered. The solvent was removed in vacuo to give a brown residue, which was purified by combiflash column chromatography (silica gel, heptane/ethyl acetate + 10% triethylamine, v/v = 10/0 to 1/1). Fractions containing the pure compound were collected and concentrated in vacuo to give the title compound (0.27 g, 63% yield) as an orange-brown oil.

^1H NMR (400 MHz, CDCl_3): δ (ppm) 7.10 (s, 1H), 5.25 – 5.15 (m, 1H), 3.65 – 3.60 (m, 1H), 3.55 – 3.50 (m, 3H), 3.50 – 3.35 (broad m, 2H), 2.20 (s, 3H), 1.50 (m, 2H), 1.25 (m, 3H), 0.80 (t, 3H).


Preparation of N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-formamidine



To a solution of N-ethylformamide (0.070 mL, 0.91 mmol, 1.1 equiv) in dichloromethane (6.6 mL) was added phosphorus oxychloride (0.085 mL, 0.91 mmol, 1.1 equiv). The solution was stirred for 1 h at room temperature. Then a solution of 5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)pyridin-3-amine (0.25 g, 0.83 mmol) in dichloromethane (3 mL) was added dropwise. The suspension was stirred for 2 h at room temperature then poured onto a mixture of 2N aqueous NaOH solution (25 mL) and ice. The aqueous layer was separated, extracted with dichloromethane (2x15 mL). The organic phase was washed with 2N aqueous NaOH solution (25 mL) and brine (25 mL), dried over anhydrous Na_2SO_4 and filtered. The solvent was removed in vacuo to give a dark yellow residue, which was purified by reverse phase preparative HPLC to give the title compound (0.09 g, 29% yield) as an orange-brown oil.

^1H NMR (400 MHz, CDCl_3): δ (ppm) 7.45 (s, 1H), 7.25 (s, 1H), 5.40 – 5.30 (m, 1H), 4.70 – 4.50 (broad s, 1H), 3.75 – 3.60 (m, 1H), 3.55–3.45 (m, 3H), 3.50 – 3.35 (broad m, 2H), 2.30 (s, 3H), 1.60 – 1.50 (m, 2H), 1.35 (m, 3H), 1.25 (m, 3H), 0.80 (t, 3H).

Preparation of N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-methyl-formamidine



To a solution of N-methyl N-ethylformamide (0.30 g, 2.37 mmol, 1.1 equiv, 80% wt.) in dichloromethane (4.0 mL) was added phosphorus oxychloride (0.20 mL, 2.2 mmol, 1.1 equiv). The solution was stirred for 1h at room temperature. Then a solution of 5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)pyridin-3-amine (0.60 g, 2.0 mmol) in dichloromethane (0.5 mL) was added dropwise. The suspension was stirred for 2h at room temperature. then poured onto a mixture of 2N aqueous NaOH solution (25 mL) and ice. The aqueous layer was separated, extracted with dichloromethane (2x15 mL). The organic phase was washed with 2N aqueous NaOH solution (25 mL) and brine (25 mL), dried over anhydrous Na₂SO₄ and filtered. The solvent was removed in vacuo to give a dark yellow residue, which was purified by reverse phase preparative HPLC to give the title compound (0.30 g, 40% yield) as an orange oil.

¹H NMR (400 MHz, CDCl₃): δ (ppm) 7.47 (broad s, 1H), 7.25 (s, 1H), 5.40 – 5.30 (m, 2H), 3.70 – 3.64 (m, 3H), 3.60-3.40 (m, 3H), 2.94 (broad s, 2H) 2.35 (d, 3H), 1.65 - 1.50 (m, 2H), 1.35 (m, 3H), 1.25 (m, 6H), 0.90 (t, 3H).

Table 67

This table gives analytical data for compounds of formula (I) prepared using techniques analogous to those above, as well as those described in WO 12/146125 (pp.370-378) and further techniques known to the person skilled in the art, for example as found in WO 08/101682 (pp.22-33).

Compound No.	Structural formula	LC-Method: R _t (min); MS-ESI (m/z; (M+H) ⁺);
67.001	5-[(E)-[ethyl(methyl)amino]methyleneamino]-6-methyl-2-(1-methyl-2-propoxy-ethoxy)pyridine-3-carboxamide	Method 1 0.9 min.; 337
67.002	N'-[5-bromo-2-methyl-6-[(1S)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 0.76 min.; 372
67.003	N'-[5-bromo-2-methyl-6-[(1R)-1-methyl-2-propoxy-ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 0.76 min.; 372

Compound No.	Structural formula	LC-Method: R _t (min); MS-ESI (m/z; (M+H) ⁺);
67.004	5-[(E)-[ethyl(methyl)amino]methyleneamino]-N,N,6-trimethyl-2-(1-methyl-2-propoxy-ethoxy)pyridine-3-carboxamide	Method 1 0.8 min.; 361
67.005	5-[(E)-[ethyl(methyl)amino]methyleneamino]-N,6-dimethyl-2-(1-methyl-2-propoxy-ethoxy)pyridine-3-carboxamide	Method 1 0.8 min.; 351
67.006	5-[(E)-[ethyl(methyl)amino]methyleneamino]-6-methyl-2-(1-methyl-2-propoxy-ethoxy)pyridine-3-carboxylic acid	Method 1 0.7 min.; 338
67.007	N'-[5-cyano-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 0.78 min.; 319
67.008	N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-isopropyl-N-methyl-formamidine	oil
67.009	N'-[5-(difluoromethyl)-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 2 13.04 min.; 342
67.010	N'-[5-(difluoromethoxy)-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 0.85 min.; 360
67.011	N'-[2,5-dimethyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 0.79 min.; 308
67.012	N-ethyl-N-methyl-N'-[2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]formamidine	Method 1 0.67 min.; 294
67.013	N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 0.76 min.; 372
67.014	N'-[6-(2-allyloxy-1-methyl-ethoxy)-5-chloro-2-methyl-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.08 min.; 326
67.015	N'-[5-chloro-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 1 1.11 min.; 328
67.016	N'-[6-[1-(allyloxymethyl)-2-methoxy-ethoxy]-5-bromo-2-methyl-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.02 min.; 400
67.017	N'-[6-(2-allyloxy-1-methyl-ethoxy)-5-cyano-2-methyl-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 0.96 min.; 317
67.018	N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-isopropyl-N-methyl-formamidine	oil

Compound No.	Structural formula	LC-Method: R _t (min); MS-ESI (m/z; (M+H) ⁺);
67.019	Methyl 5-[(E)-[ethyl(methyl)amino]methyleneamino]-6-methyl-2-(1-methyl-2-propoxy-ethoxy)pyridine-3-carboxylate	Method 1 0.80 min.; 352
67.020	N'-[5-cyano-2-methyl-6-[1-methyl-2-[3-(trifluoromethyl)phenoxy]ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.33 min.; 422
67.021	N'-[6-[2-(4-chlorophenoxy)-1-methyl-ethoxy]-5-cyano-2-methyl-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.29 min.; 387
67.022	N'-[5-cyano-2-methyl-6-(1-methyl-2-phenoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.21 min.; 353
67.023	N'-[5-chloro-2-methyl-6-[1-methyl-2-[3-(trifluoromethyl)phenoxy]ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.38 min.; 430
67.024	N'-[5-chloro-6-[2-(4-chlorophenoxy)-1-methyl-ethoxy]-2-methyl-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.38 min.; 396
67.025	N'-[5-chloro-2-methyl-6-(1-methyl-2-phenoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.23 min.; 362
67.026	N'-[5-bromo-6-[2-(4-chlorophenoxy)-1-methyl-ethoxy]-2-methyl-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.37 min.; 440
67.027	N'-[5-bromo-2-methyl-6-[1-methyl-2-[3-(trifluoromethyl)phenoxy]ethoxy]-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.42 min.; 474
67.028	N'-[5-bromo-2-methyl-6-(1-methyl-2-phenoxy-ethoxy)-3-pyridyl]-N-ethyl-N-methyl-formamidine	Method 3 1.25 min.; 406
67.029	N'-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]-N-cyclopropyl-N-methyl-formamidine	Method 1 0.77 min.; 371
67.030	1-(azetidin-1-yl)-N-[5-bromo-2-methyl-6-(1-methyl-2-propoxy-ethoxy)-3-pyridyl]methanimine	Method 1 0.76 min.; 385

Analytical Methods Used

Method 1: Mass spectra were recorded on a Mass Spectrometer from Waters (SQD or ZQ Single quadrupole mass spectrometer) equipped with an electrospray source (Polarity: positive or negative ions, Capillary: 3.00 kV, Cone range: 30-60 V, Extractor: 2.00 V, Source Temperature: 150°C, Desolvation Temperature: 350°C, Cone Gas Flow: 0 L/Hr, Desolvation

- 37 -

Gas Flow: 650 L/Hr, Mass range: 100 to 900 Da) and an Acquity UPLC from Waters: Binary pump, heated column compartment and diode-array detector. Solvent degasser, binary pump, heated column compartment and diode-array detector. Column: Waters UPLC HSS T3, 1.8 μ m, 30 x 2.1 mm, Temp: 60 °C, DAD Wavelength range (nm): 210 to 500, Solvent Gradient: A = water + 5% MeOH + 0.05 % HCOOH, B= Acetonitrile + 0.05 % HCOOH. Gradient: 0 min 0% B, 100%A; 1.2-1.5min 100% B; Flow (ml/min) 0.85.

Method 2: Mass spectra were recorded on a Mass Spectrometer from Shimadzu (SQD or ZQ Single quadrupole mass spectrometer) equipped with an electrospray source (Polarity: positive or negative ions, Capillary: 1.5 kV, Cone range: unknown, Extractor: 5.00 V, Source Temperature: 200°C, Desolvation Temperature: 250°C, Cone Gas Flow: 90 L/Hr, Desolvation Gas Flow: 90 L/Hr, Mass range: 50 to 900 Da) and an SPD-20A from LC from Shimadzu: Solvent degasser, binary pump, heated column compartment and ultraviolet detector. Column: Diamonsil C18 (2) 5 μ m 150*4.6 mm, Temp: 40 °C, SPD-20A Wavelength range (nm): 210 to 500, Solvent Gradient: A = water + 0.1 % F₃CCOOH, B= Acetonitrile + 0.1 % F₃CCOOH; Gradient: 0 min 10% B, 90% A; 15min 100% B; Flow 1.00 (ml/min)

Method 3: Mass spectra were recorded on a ZQ2000 Mass Spectrometer from Waters (Single quadrupole mass spectrometer) equipped with an electrospray source (Polarity: positive ions, Capillary (kV) 3.5, Cone (V) 60.00, Extractor (V) 3.00, Source Temperature (°C) 150, Desolvation Temperature (°C) 350, Cone Gas Flow (L/Hr) 50, Desolvation Gas Flow (L/Hr) 800, Mass range: 140 to 800 Da) DAD Wavelength range (nm): 210 to 400, Type of column: Waters ACQUITY UPLC HSS T3; Column length: 30 mm; Internal diameter of column: 2.1 mm; Particle Size: 1.8 micron; Temperature: 60°C. Solvent Gradient: A = water + 10% MeOH + 0.1 % HCOOH, B= Acetonitrile + 0.1 % HCOOH. Gradient: 0 min 0% B, 100% A; 2.5-2.8 min 100% B; 0% A; 3.0 min 100% A, 0% B; Flow (ml/min) 0.85.

Biological examples

Blumeria graminis f. sp. tritici (Erysiphe graminis f. sp. tritici) / wheat / leaf disc preventative (Powdery mildew on wheat)

Wheat leaf segments cv. Kanzler were placed on agar in a multiwell plate (24-well format) and sprayed with the formulated test compound diluted in water. The leaf disks were inoculated by shaking powdery mildew infected plants above the test plates 1 day after application. The inoculated leaf disks were incubated at 20°C and 60% rh under a light regime of 24 h darkness followed by 12 h light / 12 h darkness in a climate chamber and the activity of a compound was assessed as percent disease control compared to untreated when an appropriate level of disease damage appears on untreated check leaf segments (6 - 8 days after application).

- 38 -

The following compounds gave at 200 ppm give at least 80% disease control in this test when compared to untreated control leaf disks under the same conditions, which show extensive disease development:

Q.002, Q.003, Q.007, Q.008, Q.009, Q.010, Q.011, Q.013, Q.014, Q.015, Q.016,
 5 Q.017, Q.018, Q.020, Q.021, Q.022, Q.023, Q.024, Q.025, Q.026, Q.027, Q.028, Q.029,
 Q.030.

Puccinia recondita f. sp. tritici / wheat / leaf disc preventative (brown rust)

Wheat leaf segments cv. Kanzler were placed on agar in multiwell plates (24-well format) and sprayed with the formulated test compound diluted in water. The leaf disks were
 10 inoculated with a spore suspension of the fungus 1 day after application. The inoculated leaf segments were incubated at 19°C and 75% rh under a light regime of 12 h light / 12 h darkness in a climate cabinet and the activity of a compound was assessed as percent disease control compared to untreated when an appropriate level of disease damage appears in untreated check leaf segments (7 – 9 days after application).

15 The following compounds gave at 200 ppm gave at least 80% disease control in this test when compared to untreated control leaf disks under the same conditions, which show extensive disease development:

Q.002, Q.003, Q.007, Q.008, Q.009, Q.010, Q.011, Q.013, Q.014, Q.015, Q.016,
 Q.017, Q.018, Q.020, Q.021, Q.022, Q.023, Q.024, Q.025, Q.026, Q.028, Q.029, Q.030..

20 **Puccinia recondita f. sp. tritici / wheat / leaf disc curative (Brown rust)**

Wheat leaf segments cv. Kanzler are placed on agar in multiwell plates (24-well format). The leaf segments are inoculated with a spore suspension of the fungus. Plates were stored in darkness at 19°C and 75% rh. The formulated test compound diluted in water was applied 1 day after inoculation. The leaf segments were incubated at 19°C and 75% rh
 25 under a light regime of 12 h light / 12 h darkness in a climate cabinet and the activity of a compound was assessed as percent disease control compared to untreated when an appropriate level of disease damage appears in untreated check leaf segments (6 – 8 days after application).

30 The following compounds gave at 200 ppm gave at least 80% disease control in this test when compared to untreated control leaf disks under the same conditions, which show extensive disease development:

Q.002, Q.003, Q.007, Q.008, Q.009, Q.010, Q.011, Q.012, Q.013, Q.014, Q.015,
 Q.016, Q.017, Q.018, Q.019, Q.020, Q.021, Q.022, Q.023, Q.024, Q.025, Q.026, Q.027,
 Q.028, Q.029, Q.030..

35 **Phakopsora pachyrhizi on soybean, preventive treatment**

- 39 -

The compound activity was tested under 1 day preventive conditions. Soybean plants with a fully enfolded first trifoliolate leaf were sprayed with a track sprayer and 50 l/ha spray volume with the test compounds, either solo or in tankmix as shown in the table below.

1 day after application leaf discs were cut from the first trifoliolate leaf and placed in multiwell

5 plates on water-agar. 5 leaf discs per treatment where infested with spores of a triazole tolerant soybean rust strain. The multiwell plates where sealed and placed in an incubator 48 h in darkness and 12 h light/dark cycle afterwards. Rust infestation on leaf discs was evaluated visually 11 days after application and average activity calculated in relation to disease severity on untreated check leaf discs.


10 The following compounds gave at 200 ppm gave at least 80% disease control in this test when compared to untreated control leaf disks under the same conditions, which show extensive disease development:

Q.002, Q.003, Q.007, Q.008, Q.009, Q.010, Q.011, Q.012, Q.013, Q.014, Q.015,
Q.016, Q.017, Q.018, Q.019 Q.020, Q.023, Q.024, Q.025, Q.026, Q.027, Q.028, Q.029,
15 Q.030..

- 40 -

WHAT IS CLAIMED IS

1. A compound of formula (I)



5 wherein

R^1 represents hydrogen, halogen, cyano, OH, NH_2 , C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, $NH(C_1$ - C_4 alkyl), $N(C_1$ - C_4 alkyl) $_2$, $CO(C_1$ - C_4 alkyl), $CO_2(C_1$ - C_4 alkyl), CO_2H , $CONH(C_1$ - C_4 alkyl), $CON(C_1$ - C_4 alkyl) $_2$, $SO_2NH(C_1$ - C_4 alkyl), $SO_2N(C_1$ - C_4 alkyl) $_2$, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkyl- C_1 - C_4 alkoxy or C_2 - C_4 alkynyl;

10 R^2 represents C_3 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, R^7 or $-C_1$ - C_2 alkyl- R^7 , each of which may be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl and C_1 - C_4 haloalkoxy;

R^3 and R^4 independently of each other represent hydrogen, C_1 - C_4 alkyl or C_3 - C_6 cycloalkyl; or

15 R^3 and R^4 together with the nitrogen atom to which they are attached form a 3-6-membered saturated cyclic group;

R^5 represents H, C_1 - C_4 alkyl or C_1 - C_4 haloalkyl;

R^6 represents C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl or C_1 - C_4 haloalkoxy;

20 R^7 represents a three- to ten-membered monocyclic or fused bicyclic ring system which can be aromatic, partially saturated or fully saturated and can contain 1 to 4 hetero atoms selected from the group consisting of nitrogen, oxygen and sulfur, it being possible for the three- to ten-membered ring system itself to be optionally substituted by one or more groups independently selected from the group consisting halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy and C_1 - C_4 haloalkoxy;

25 and tautomers/isomers/enantiomers/salts and N-oxides of these compounds.

2. A compound of formula (I) according to claim 1 wherein

R^1 represents hydrogen, halogen, cyano, OH, NH_2 , C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, $NH(C_1$ - C_4 alkyl), $N(C_1$ - C_4 alkyl) $_2$, $CO(C_1$ - C_4 alkyl), $CO_2(C_1$ - C_4 alkyl), CO_2H , $CONH(C_1$ - C_4

- 41 -

alkyl), CON(C₁-C₄ alkyl)₂, SO₂NH(C₁-C₄ alkyl), SO₂N(C₁-C₄ alkyl)₂, C₁-C₄ haloalkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkoxy, or C₂-C₄ alkynyl;

R² represents C₃-C₆alkyl, C₃-C₆alkenyl, C₃-C₆alkynyl, C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, -C₁-C₂alkyl-C₃-C₆cycloalkyl, -C₁-C₂alkyl-C₃-C₆cycloalkenyl, each of which
5 may be optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₂ alkyl, C₁-C₂ haloalkyl, C₁-C₂ alkoxy and C₁-C₂ haloalkoxy;

R³ and R⁴ independently of each other represent hydrogen, C₁-C₄ alkyl or C₃-C₆ cycloalkyl; or

R³ and R⁴ together with the nitrogen atom to which they are attached form a 3-6-
10 membered saturated cyclic group;

R⁵ represents H, C₁-C₄ alkyl or C₁-C₄ haloalkyl;

R⁶ represents C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy.

3. A compound of formula (I) according to either claim 1 or claim 2 wherein

R¹ represents hydrogen, halogen, cyano, OH, NH₂, methyl, ethyl, cyclopropyl, NH(C₁-C₂ alkyl), N(C₁-C₂ alkyl)₂, CO(C₁-C₂ alkyl), CO₂(C₁-C₂ alkyl), CO₂H, CONH(C₁-C₂ alkyl), CON(C₁-C₂ alkyl)₂, SO₂NH(C₁-C₂ alkyl), SO₂N(C₁-C₂ alkyl)₂, C₁-C₄ fluoroalkyl, C₁-C₄ alkoxy, C₁-C₂ haloalkoxy or C₂-C₄ alkynyl;

R² represents C₃-C₆alkyl, C₃-C₆alkenyl, C₃-C₆alkynyl, C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, -C₁-C₂alkyl-C₃-C₆cycloalkyl, -C₁-C₂alkyl-C₃-C₆cycloalkenyl, each of which
20 may be optionally substituted by one or more groups independently selected from the group consisting halogen, C₁-C₂ alkyl, C₁-C₂ haloalkyl and C₁-C₂ haloalkoxy;

R³ and R⁴ independently of each other represent hydrogen, C₁-C₃ alkyl or C₃-C₅ cycloalkyl; or

R³ and R⁴ together with the nitrogen atom to which they are attached form a 5-
25 membered saturated cyclic group;

R⁵ represents H or C₁-C₄ alkyl;

R⁶ represents C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy.

4. A compound of formula (I) according to any preceding claim wherein

R¹ represents hydrogen, halogen, cyano, OH, NH₂, methyl, ethyl, cyclopropyl, NHMe, NMe₂, COMe, CO₂Me, CO₂H, CONHMe, CONMe₂, SO₂NHMe, SO₂NMe₂, CHF₂, CF₃, OMe, OCHF₂ or acetylenyl;

R² represents C₃-C₆alkyl, C₃-C₆alkenyl, C₃-C₆alkynyl, C₃-C₆cycloalkyl, C₃-C₅cycloalkenyl, -C₁-C₂alkyl-C₃-C₆cycloalkyl, -C₁-C₂alkyl-C₃-C₆cycloalkenyl which may be
35

- 42 -

optionally substituted by one or more groups independently selected from the group consisting fluoro, C₁-C₂ alkyl, C₁-C₂ fluoroalkyl and C₁-C₂ fluoroalkoxy;

R³ and R⁴ independently of each other represent hydrogen, methyl, ethyl, isopropyl or cyclopropyl; or

- 5 R³ and R⁴ together with the nitrogen atom to which they are attached form a 5-membered saturated cyclic group;

R⁵ represents H or methyl;

R⁶ represents C₁-C₄ alkyl or C₁-C₄ alkoxy.

- 10 5. A compound of formula (I) according to any preceding claim wherein

R₁ represents hydrogen, halogen, cyano, methyl, ethyl, cyclopropyl, CHF₂, CF₃, OMe, or OCHF₂;

- 15 R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, 2-methylbutyl, 2,2-dimethylpropyl, 2-methylpentyl, 3-methylpentyl, 2,3-dimethylbutyl, 2,2-dimethylbutyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl -CH₂-cyclopropyl, -CH₂-cyclobutyl, -CH₂-cyclopentyl, and -CH₂-cyclopentenyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro, methyl and difluoromethoxy;

R³ is hydrogen or methyl;

- 20 R⁴ is methyl or ethyl;

R⁵ represents H or methyl;

R⁶ represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl, methoxy or ethoxy.

- 25 6. A compound of formula (I) according any preceding claim wherein

R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro and difluoromethoxy;

- 30 R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl, cyclopentyl, methoxy or ethoxy.

- 43 -

7. A compound of formula (I) according any preceding claim wherein

R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro
5 and difluoromethoxy;

R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ represents methyl, ethyl, n-propyl, iso-propyl, cyclopropyl, cyclobutyl or cyclopentyl.

10

8. A compound of formula (I) according any preceding claim wherein

R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro
15 and difluoromethoxy;

R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ represents methyl, ethyl, n-propyl or iso-propyl.

20

9. A compound of formula (I) according any preceding claim wherein

R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, each of which may be optionally substituted by one or more groups independently selected from the group consisting fluoro
25 and difluoromethoxy;

R³ represents hydrogen or methyl;

R⁴ is ethyl;

R⁵ is hydrogen or methyl;

R⁶ is methyl.

30

10. A compound of formula (I) according any preceding claim wherein

R¹ represents hydrogen, Cl, Br, methyl, CHF₂ or cyano;

R₂ is n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or cyclopropyl;

R³ represents hydrogen or methyl;

35

R⁴ is ethyl;

- 44 -

R⁵ is hydrogen;

R⁶ is methyl.

5 11. A composition comprising a fungicidally effective amount of a compound of formula (I) as defined in any one of claims 1 to 10, optionally comprising at least one additional active ingredient.

10 12. A method of controlling or preventing phytopathogenic diseases on useful plants or on propagation material thereof, which comprises applying to the useful plants, the locus thereof or propagation material thereof a fungicidally effective amount of a compound of formula (I) as defined in any one of claims 1 to 10.

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2015/057088

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D213/74 A01N43/40
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2012/146125 A1 (SYNGENTA PARTICIPATIONS AG [CH]) 1 November 2012 (2012-11-01) cited in the application page 207; line A.1355 page 317; table 3; compound T3.1355 page 400; compound Q.429 pages 401-448; biological examples claims 1, 14, 15,16 ----- -/--	1-12



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

12 May 2015

Date of mailing of the international search report

20/05/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Cortés, José

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2015/057088

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2008/101682 A2 (SYNGENTA PARTICIPATIONS AG [CH]; SYNGENTA LTD [GB]) 28 August 2008 (2008-08-28) cited in the application 53; example P6 page 101; table P; compound P.28 pages 159 and 160; lines A.1.351 and A.1355 page 186; table 5; compounds T5.1.351 and T5.1.355 pages 263 to 266; biological examples claims 1,15,16 -----	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2015/057088

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2012146125 A1	01-11-2012	AR 086110 A1	20-11-2013
		US 2014335201 A1	13-11-2014
		UY 34039 A	30-11-2012
		WO 2012146125 A1	01-11-2012

WO 2008101682 A2	28-08-2008	BR PI0807791 A2	08-07-2014
		CA 2677949 A1	28-08-2008
		CN 101657423 A	24-02-2010
		CN 103396359 A	20-11-2013
		EP 2125734 A2	02-12-2009
		JP 2010519267 A	03-06-2010
		JP 2014051522 A	20-03-2014
		RU 2009135067 A	27-03-2011
		RU 2012147703 A	20-05-2014
		UA 102514 C2	25-07-2013
		US 2011046088 A1	24-02-2011
		US 2013338105 A1	19-12-2013
		WO 2008101682 A2	28-08-2008
