



- (51) **International Patent Classification:**
C07D 413/14 (2006.01) *A01N 43/80* (2006.01)
- (21) **International Application Number:**
PCT/EP2016/072588
- (22) **International Filing Date:**
22 September 2016 (22.09.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
15186540.9 23 September 2015 (23.09.2015) EP
- (71) **Applicant:** SYNGENTA PARTICIPATIONS AG
[CH/CH]; Schwarzwaldallee 215, 4058 Basel (CH).
- (72) **Inventors:** CASSAYRE, Jérôme, Yves; Syngenta Crop Protection AG, Breitenloh 5, 4333 Münchwilen (CH). STOLLER, André; Syngenta Crop Protection AG, Schaffhauserstrasse, 4332 Stein (CH). PITTERNA, Thomas; Syngenta Crop Protection AG, Schaffhauserstrasse, 4332 Stein (CH). EL QACEMI, Myriem; Syngenta Crop Protection AG, Schaffhauserstrasse, 4332 Stein (CH).
- (74) **Agent:** SYNGENTA INTERNATIONAL AG; Schwarzwaldallee 215, (WRO B8-Z1-30), 4058 Basel (CH).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

- (54) **Title:** ISOXAZOLINE-SUBSTITUTED BENZAMIDES AND ANALOGUES AS INSECTICIDES



- (57) **Abstract:** The present invention relates to compounds of formula (I) wherein A¹, A², A³, A⁴, R¹, R², R³, R⁴ and n are as defined in claim 1; or a tautomer, isomer, enantiomer, salt or N-oxide thereof; to intermediates for preparing compounds of formula (I), to compositions comprising them and to methods of using them to combat and control insect, acarine, nematode and mollusc pests.


ISOXAZOLINE-SUBSTITUTED BENZAMIDES AND ANALOGUES AS INSECTICIDES

The present invention relates to certain isoxazolidine derivatives, to processes and intermediates for preparing these derivatives, to insecticidal, acaricidal, nematocidal and molluscicidal compositions comprising these derivatives and to methods of using these derivatives to control insect, acarine, nematode and mollusc pests.

Certain isoxazoline derivatives with insecticidal properties are disclosed, for example, in WO2011067272.

It has now surprisingly been found that certain isoxazolidine derivatives have highly potent insecticidal properties.

The present invention provides, in a first aspect of the invention, compounds of formula (I)



wherein

A¹, A², A³ and A⁴ are independently of one another C-H, C-R⁵, or nitrogen;

R¹ is hydrogen, C₁-C₈alkyl, C₁-C₈alkylcarbonyl-, C₁-C₈alkoxy, C₁-C₈alkoxy-C₁-C₈alkyl or C₁-

C₈alkoxycarbonyl-;

R² is C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a}, C₁-C₈haloalkyl, C₁-C₈haloalkyl substituted by one to three R^{6a}, C₃-C₈cycloalkyl, C₃-C₈cycloalkyl substituted by one to three R^{6b}, C₃-C₈cycloalkyl where one carbon atom is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈alkyl, C₃-C₈cycloalkyl-C₁-C₈alkyl where one carbon atom in the cycloalkyl group is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-

C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a}, C₂-C₈haloalkenyl, C₂-C₈haloalkenyl substituted by one to three R^{6a}, C₂-C₈alkynyl, C₂-C₈haloalkynyl, phenyl, phenyl substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷, 5-6 membered heteroaryl-C₁-C₄alkyl, 5-6 membered heteroaryl-C₁-C₄alkyl wherein the heteroaryl moiety is

substituted by one to three R⁷, -N(R⁸)(R⁹), -OR¹⁰ or halogen;

R³ is C₁-C₈haloalkyl;

R⁴ is aryl, aryl substituted by one to three R⁷, heteroaryl or heteroaryl substituted by one to three R⁷;

R⁵ is independently halogen, cyano, nitro, C₁-C₈alkyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈haloalkenyl, C₂-C₈alkynyl, C₂-C₈haloalkynyl, C₁-C₈alkoxy, C₁-C₈haloalkoxy, or C₁-C₈alkoxycarbonyl-, or

two R⁵ on adjacent carbon atoms together form a -CH=CH-CH=CH- bridge or a -N=CH-CH=CH- bridge;

R^{6a} is independently cyano, nitro, amino, C₁-C₈alkylamino, N,N-C₁-C₈dialkylamino, hydroxy, C₁-C₈alkoxy, or C₁-C₈haloalkoxy;

R^{6b} is independently halogen, cyano, nitro, oxo, C₁-C₈alkyl, C₁-C₈haloalkyl, amino, C₁-C₈alkylamino, N,N-C₁-C₈dialkylamino, hydroxyl, C₁-C₈alkoxy, C₁-C₈haloalkoxy, phenyl, phenyl substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷;

R⁷ is independently halogen, cyano, nitro, C₁-C₈alkyl, C₁-C₈haloalkyl, C₁-C₈alkoxy, C₁-C₈haloalkoxy;

- 5 R⁸ and R⁹ are independently hydrogen, cyano, cyano-C₁-C₈alkyl, C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a}, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a}, C₁-C₈alkoxy, C₁-C₈haloalkoxy, C₁-C₈haloalkoxy substituted by one to three R^{6a}, C₁-C₈alkoxy substituted by one to three R^{6a}, C₁-C₈haloalkyl, C₁-C₈haloalkyl substituted by one to three R^{6a}, C₃-C₈cycloalkyl, C₃-C₈cycloalkyl substituted by one to three R^{6b}, C₃-C₈cycloalkyl where one carbon atom is replaced by O, S, S(O) or SO₂,
- 10 C₃-C₈cycloalkyl-C₁-C₈alkyl, C₃-C₈cycloalkyl-C₁-C₈alkyl where one carbon atom in the cycloalkyl group is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a}, C₂-C₈haloalkenyl, C₂-C₈haloalkenyl substituted by one to three R^{6a}, C₂-C₈alkynyl, C₂-C₈haloalkynyl, phenyl, phenyl substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered
- 15 heteroaryl substituted by one to three R⁷, 5-6 membered heteroaryl-C₁-C₄alkyl, 5-6 membered heteroaryl-C₁-C₄alkyl wherein the heteroaryl moiety is substituted by one to three R⁷, -S(O)R¹⁰, -S(O)₂R¹⁰, COR¹⁰, COOR¹⁰, or R⁸ and R⁹ together with the nitrogen atom can be linked through a C₃-C₈alkylene chain, a C₃-C₈alkylene chain substituted by one to three R^{6b} or a C₃-C₈alkylene chain, where one carbon atom is replaced by O, S, S(O) or SO₂;
- 20 R¹⁰ is hydrogen, cyano-C₁-C₈alkyl, C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a}, C₁-C₈haloalkyl, C₁-C₈haloalkyl substituted by one to three R^{6a}, C₃-C₈cycloalkyl, C₃-C₈cycloalkyl substituted by one to three R^{6b}, C₃-C₈cycloalkyl where one carbon atom is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈alkyl, C₃-C₈cycloalkyl-C₁-C₈alkyl where one carbon atom in the cycloalkyl group is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a},
- 25 C₂-C₈haloalkenyl, C₂-C₈haloalkenyl substituted by one to three R^{6a}, C₂-C₈alkynyl, C₂-C₈haloalkynyl, phenyl, phenyl substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷, 5-6 membered heteroaryl-C₁-C₄alkyl or 5-6 membered heteroaryl-C₁-C₄alkyl wherein the heteroaryl moiety is substituted by one to three R⁷;
- 30 n is 1 or 2;
- or a tautomer, isomer, enantiomer, salt or N-oxide thereof.

Compounds of formula (I) which have at least one basic centre can form, for example, acid addition salts, for example with strong inorganic acids such as mineral acids, for example perchloric acid, sulfuric acid, nitric acid, nitrous acid, a phosphorus acid or a hydrohalic acid, with strong organic

35 carboxylic acids, such as C₁-C₄alkanecarboxylic acids which are unsubstituted or substituted, for example by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic acid, malonic acid, succinic acid, maleic acid, fumaric acid or phthalic acid, such as hydroxycarboxylic acids, for example ascorbic acid, lactic acid, malic acid, tartaric acid or citric acid, or such as benzoic acid, or with organic sulfonic acids, such as C₁-C₄alkane- or arylsulfonic acids which are

40 unsubstituted or substituted, for example by halogen, for example methane- or p-toluenesulfonic acid.

Compounds of formula (I) which have at least one acidic group can form, for example, salts with bases, for example mineral salts such as alkali metal or alkaline earth metal salts, for example sodium,

potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower-alkylamine, for example ethyl-, diethyl-, triethyl- or dimethylpropylamine, or a mono-, di- or trihydroxy-lower-alkylamine, for example mono-, di- or triethanolamine.

5 The alkyl groups occurring in the definitions of the substituents can be straight-chain or branched and are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, pentyl, hexyl, nonyl, decyl and their branched isomers. Alkylsulfanyl, alkylsulfanyl, alkylsulfonyl, alkoxy, alkenyl and alkynyl radicals are derived from the alkyl radicals mentioned. The alkenyl and alkynyl groups can be mono- or polyunsaturated.

10 Halogen is generally fluorine, chlorine, bromine or iodine. This also applies, correspondingly, to halogen in combination with other meanings, such as haloalkyl or halophenyl.

Haloalkyl groups preferably have a chain length of from 1 to 6 carbon atoms. Haloalkyl is, for example, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 2-fluoroethyl, 2-chloroethyl, pentafluoroethyl, 1,1-difluoro-2,2,2-trichloroethyl, 2,2,3,3-
15 tetrafluoroethyl and 2,2,2-trichloroethyl.

Alkoxy is, for example, methoxy, ethoxy, propoxy, i-propoxy, n-butoxy, isobutoxy, sec-butoxy and tert-butoxy and also the isomeric pentyloxy and hexyloxy radicals.

Alkoxyalkyl groups preferably have a chain length of 1 to 6 carbon atoms.

Alkoxyalkyl is, for example, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, n-
20 propoxymethyl, n-propoxyethyl, isopropoxymethyl or isopropoxyethyl.

Alkoxyalkyl is for example methoxycarbonyl (which is C₁alkoxyalkyl), ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, n-butoxycarbonyl, tert-butoxycarbonyl, n-pentoxycarbonyl or hexoxycarbonyl.

The cycloalkyl groups preferably have from 3 to 6 ring carbon atoms, for example cyclopropyl,
25 cyclobutyl, cyclopentyl and cyclohexyl.

Heteroaryl groups are preferably 5-6 membered heteroaryl or are 5-6 membered heteroaryl substituted by one to three R⁷, where heteroaryl groups contain 1 to 3 hetero atoms selected from the group consisting of nitrogen, oxygen and sulfur, it not being possible for each ring system to contain more than 2 oxygen atoms and more than 2 sulfur atoms. Examples of monocyclic groups include pyridyl,
30 pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl (e.g. 1,2,4 triazolyl), furanyl, thiophenyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, tetrazolyl and thiadiazolyl.

The compounds of formula (I) according to the invention also include hydrates which may be formed during the salt formation.

Preferred values of A¹, A², A³, A⁴, R¹, R², R³, R⁴, R⁵, R^{6a}, R^{6b}, R⁷, R⁸, R⁹, and R¹⁰, in relation to
35 each compound of the present invention, including the intermediate compounds, are, in any combination (including combinations of preferred values with the original values) as set out below.

Preferably no more than two of A¹, A², A³ and A⁴ are nitrogen.

Preferably A¹ is C-H or C-R⁵; more preferably A¹ is C-R⁵.

Preferably A² is C-H or C-R⁵; more preferably A² is C-H.

40 Preferably A³ is C-H or N; more preferably A³ is C-H.

Preferably A⁴ is C-H or N; more preferably A⁴ is C-H.

Preferably R^1 is hydrogen, C_1 - C_8 alkyl, C_1 - C_8 alkylcarbonyl- or C_1 - C_8 alkoxycarbonyl-; more preferably hydrogen, methyl, ethyl, methylcarbonyl-, or methoxycarbonyl-; most preferably hydrogen, methyl or ethyl; especially hydrogen or methyl; more especially hydrogen.

Preferably R^2 is C_1 - C_8 alkyl, C_1 - C_8 alkyl substituted by one to three R^{6a} , C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, di- C_1 - C_8 alkylamino, $-N(R^8)(R^9)$, aryl, aryl substituted by one to three R^{6b} , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^{6b} or halogen; more preferably C_1 - C_4 alkyl, C_1 - C_4 alkyl substituted by one to three R^{6a} , C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_5 cycloalkyl, C_1 - C_4 haloalkyl, di- C_1 - C_4 alkylamino, $-N(R^8)(R^9)$, 1-3 halo-substituted phenyl, 5-6 membered heteroaryl or fluoro; most preferably methyl, ethyl, cyclopropyl, chloromethyl, fluoromethyl, difluoromethyl, fluoro, dimethylamino or methylamino.

Preferably R^3 is C_1 - C_4 haloalkyl; more preferably chlorodifluoromethyl or trifluoromethyl; most preferably trifluoromethyl.

Preferably R^4 is aryl, aryl substituted by one to three R^7 , 5-6 membered heteroaryl or 5-6 membered heteroaryl substituted by one to three R^7 ; more preferably R^4 is aryl or aryl substituted by one to three R^7 ; most preferably phenyl or phenyl substituted by one to three R^7 ; even more preferably R^4 is phenyl substituted by one to three R^7 ; especially R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; more especially R^4 is 3-chloro-5-trifluoromethyl-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 3,5-dichloro-4-fluoro-phenyl, or 3,4,5-trichloro-phenyl.

Preferably R^5 is independently halogen, cyano, nitro, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, or C_2 - C_8 alkenyl, or two R^5 on adjacent carbon atoms together form a $-CH=CH-CH=CH-$ bridge; more preferably halogen, cyano, nitro, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_3 - C_8 cycloalkyl, or C_1 - C_8 haloalkyl; even more preferably bromo, chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, cyclopropyl, or vinyl; yet even more preferably bromo, chloro, fluoro, cyclopropyl, trifluoromethyl, vinyl, or methyl; most preferably chloro, bromo, trifluoromethyl, fluoro, or methyl.

Preferably R^{6a} independently is cyano, halogen, nitro, C_1 - C_4 alkoxy, or C_1 - C_4 haloalkoxy; more preferably fluoro, cyano, nitro, methoxy, difluoromethoxy or trifluoromethoxy.

Preferably R^{6b} independently is halogen, cyano, nitro, C_1 - C_4 alkyl, or C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, or C_1 - C_4 haloalkoxy; more preferably bromo, chloro, fluoro, cyano, nitro methyl, ethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy; most preferably chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, methoxy, or trifluoromethoxy; especially chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, methoxy, or trifluoromethoxy; more especially bromo, fluoro, chloro, or trifluoromethyl.

Preferably R^7 is independently halogen, cyano, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy; more preferably, methyl, fluoro, chloro, bromo, trifluoromethyl, trifluoromethoxy, cyano or methoxy, even more preferably R^7 is Cl, Br, F, CF_3 , CH_3 or OCF_3 .

Preferably R^8 and R^9 are independently hydrogen, cyano- C_1 - C_8 alkyl, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, C_1 - C_8 alkoxyalkyl, C_1 - C_8 hydroxyalkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl- C_1 - C_4 alkyl or phenyl- C_1 - C_4 alkyl wherein the phenyl moiety is substituted by one to three R^7 , 5-6 membered heteroaryl- C_1 - C_4 alkyl or 5-6 membered heteroaryl- C_1 - C_4 alkyl wherein the heteroaryl moiety is substituted by one to three R^7 ; more preferably R^8 and R^9 are independently hydrogen, cyano- C_1 - C_8 alkyl-, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 alkoxyalkyl, C_1 - C_8 hydroxyalkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl- C_1 - C_4 alkyl or

phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl-C₁-C₄alkyl or 5-6 membered heteroaryl-C₁-C₄alkyl wherein the heteroaryl moiety is substituted by one to three R⁷; yet even more preferably R⁸ and R⁹ are independently hydrogen, C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄haloalkyl, C₁-C₄hydroxyalkyl, C₁-C₄alkoxy-C₁-C₄alkyl, phenyl-CH₂- or phenyl-CH₂- wherein the phenyl moiety is substituted by one to three R⁷, furanyl or furanyl substituted by one to three R⁷, thietanyl, oxetanyl, oxo-thietanyl, or dioxo-thietanyl; yet even more preferably R⁸ and R⁹ are independently hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclobutyl, oxetanyl, thietanyl, trifluoroethyl, difluoroethyl, allyl, propargyl, cyanomethyl, benzyl, benzyl substituted by one to three R⁷, or pyridine-methyl- or pyridine-methyl- substituted by one to three R⁷; especially R⁸ and R⁹ are independently hydrogen and methyl.

Preferably each R¹⁰ is independently hydrogen, C₁-C₈alkyl, C₁-C₈haloalkyl or C₃-C₈cycloalkyl, more preferably hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₃-C₆cycloalkyl, most preferably, hydrogen, methyl, ethyl or cyclopropyl.

Preferably n is 2.

In an embodiment E1 of formula (I), independent of other embodiments, A¹ is CR⁵ and A², A³ and A⁴ are each CH, wherein R⁵ is as defined in the first aspect of the invention.

In an embodiment E2 of formula (I), independent of other embodiments, R¹ is hydrogen, C₁-C₈alkyl, C₁-C₈alkylcarbonyl- or C₁-C₈alkoxycarbonyl-.

In an embodiment E3 of formula (I), independent of other embodiments, R² is C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a}, C₂-C₈alkenyl, C₂-C₈alkynyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, di-C₁-C₈alkylamino, -N(R⁶)(R⁹), aryl, aryl substituted by one to three R^{6b}, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^{6b}, wherein R^{6a} and R^{6b} are as defined in the first aspect of the invention or halogen.

In an embodiment E4 of formula (I), independent of other embodiments, R³ is C₁-C₄haloalkyl.

In an embodiment E5 of formula (I), independent of other embodiments, R⁴ is aryl, aryl substituted by one to three R⁷, 5-6 membered heteroaryl or 5-6 membered heteroaryl substituted by one to three R⁷, wherein R⁷ is as defined in the first aspect of the invention.

Embodiment E6 of formula (I) provides compounds of formula (I) wherein A¹ is CR⁵ and A², A³ and A⁴ are each CH; R¹ is hydrogen, methyl, ethyl, methylcarbonyl-, or methoxycarbonyl; R² is C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a}, C₂-C₈alkenyl, C₂-C₈alkynyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, di-C₁-C₈alkylamino, -N(R⁶)(R⁹), aryl, aryl substituted by one to three R^{6b}, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^{6b} or fluoro; R³ is C₁-C₄haloalkyl; R⁴ is aryl or aryl substituted by one to three R⁷; and n is 2; wherein R⁵ is halogen or C₁-C₈alkyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, or C₂-C₈alkenyl; R^{6a} is independently cyano, halogen, nitro, C₁-C₄alkoxy, or C₁-C₄haloalkoxy; and R^{6b} is independently halogen, cyano, nitro, C₁-C₄alkyl, or C₁-C₄haloalkyl, C₁-C₄alkoxy, or C₁-C₄haloalkoxy.

Embodiment E7 of formula (I) provides compounds of formula (I) wherein A¹ is CR⁵ and A², A³ and A⁴ are each CH; R¹ is hydrogen, methyl, ethyl, methylcarbonyl-, or methoxycarbonyl; R² is C₁-C₄alkyl, C₁-C₄alkyl substituted by one to three R^{6a}, C₂-C₄alkenyl, C₂-C₄alkynyl, C₃-C₅cycloalkyl, C₁-C₄haloalkyl, di-C₁-C₄alkylamino, -NH(C₁-C₄alkyl), fluoro, phenyl, or 5-6 membered heteroaryl; R³ is C₁-C₄haloalkyl; R⁴ is aryl or aryl substituted by one to three R⁷; and n is 2; wherein R⁵ is halogen, cyano, nitro, C₁-C₈alkyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, C₂-C₈alkenyl; R^{6a} is independently fluoro, cyano, nitro, methoxy,

difluoromethoxy or trifluoromethoxy; and R⁷ is independently bromo, chloro, fluoro, cyano, nitro methyl, ethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy.

Embodiment E8 of formula (I) provides compounds of formula (I) wherein A¹ is CR⁵ and A², A³ and A⁴ are each CH; R¹ is hydrogen; R² is C₁-C₄alkyl, C₁-C₄alkyl substituted by one to three R^{6a}, C₂-C₄alkenyl, C₂-C₄alkynyl, C₃-C₅cycloalkyl, C₁-C₄haloalkyl, di-C₁-C₄alkylamino, -NH(C₁-C₄alkyl), fluoro, 1-3 halo-substituted phenyl, or 5-6 membered heteroaryl; R³ is chlorodifluoromethyl or trifluoromethyl; R⁴ is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; and n is 2; wherein R⁵ is bromo, chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, cyclopropyl, or vinyl; R^{6a} is independently fluoro, cyano, nitro, methoxy, difluoromethoxy or trifluoromethoxy; and R^{6b} is independently bromo, chloro, fluoro, cyano, nitro methyl, ethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy.

Embodiment E9 of formula (I) provides compounds of formula (I) wherein A¹ is CR⁵ and A², A³ and A⁴ are each CH; R¹ is hydrogen; R² is C₁-C₄alkyl, C₁-C₄alkyl substituted by one to three R^{6a}, C₂-C₄alkenyl, C₂-C₄alkynyl, C₃-C₅cycloalkyl, C₁-C₄haloalkyl, di-C₁-C₄alkylamino, -NH(C₁-C₄alkyl), fluoro or 5-6 membered heteroaryl; R³ is chlorodifluoromethyl or trifluoromethyl; R⁴ is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; and n is 2; wherein R⁵ is bromo, chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, cyclopropyl, or vinyl; R^{6a} is independently fluoro, cyano, nitro, methoxy, difluoromethoxy or trifluoromethoxy; and R^{6b} is independently bromo, chloro, fluoro, cyano, nitro methyl, ethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy.

Embodiment E10 of formula (I) provides compounds of formula (I) wherein A¹ is CR⁵ and A², A³ and A⁴ are each CH; R¹ is hydrogen; R² is methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclobutyl, oxetanyl, thietanyl, chloromethyl, fluoromethyl, difluoromethyl, trifluoroethyl, difluoroethyl, allyl, propargyl, cyanomethyl, dimethylamino, methylamino or fluoro; R³ is chlorodifluoromethyl or trifluoromethyl; R⁴ is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; and n is 2; wherein R⁵ is bromo, chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, cyclopropyl, or vinyl; R^{6a} is independently fluoro, cyano, nitro, methoxy, difluoromethoxy or trifluoromethoxy; and R^{6b} is independently bromo, chloro, fluoro, cyano, nitro methyl, ethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy.

Embodiment E11 of formula (I) provides compounds of formula (I) wherein A², A³ and A⁴ are each CH; R¹ is hydrogen; R³ is chlorodifluoromethyl or trifluoromethyl; n is 2; A¹ is CR⁵, wherein R⁵ is chloro, bromo, trifluoromethyl, fluoro, or methyl; R² is methyl, ethyl, cyclopropyl, chloromethyl, fluoromethyl, difluoromethyl, iso-propyl, methylamino, dimethylamino, fluoro, cyclobutyl, 3,3,3-trifluoropropyl, or 2,2,2-trifluoroethyl, or methoxyethyl; and R⁴ is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R⁴ is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R⁴ is 3,5-

dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E12 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro or methyl; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E13 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro, bromo, trifluoromethyl, fluoro, or methyl; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is phenyl, pyridyl, mono-, di- or tri-substituted phenyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl, or mono- or di-substituted pyridyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl.

Embodiment E14 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro or methyl; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is 3,5-dichloro-4-fluorophenyl, 3,4,5-trichloro-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3-chloro-5-trifluoromethyl-phenyl.

Embodiment E15 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro; R^2 is methyl, ethyl, cyclopropyl, chloromethyl, fluoromethyl, difluoromethyl, iso-propyl, methylamino, dimethylamino, fluoro, cyclobutyl, 3,3,3-trifluoropropyl, or 2,2,2-trifluoroethyl, or methoxyethyl; and R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E16 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl,

3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E17 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is phenyl, pyridyl, mono-, di- or tri-substituted phenyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl, or mono- or di-substituted pyridyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl.

Embodiment E18 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is chloro; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is 3,5-dichloro-4-fluorophenyl, 3,4,5-trichloro-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3-chloro-5-trifluoromethyl-phenyl.

Embodiment E19 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is bromo; R^2 is methyl, ethyl, cyclopropyl, chloromethyl, fluoromethyl, difluoromethyl, iso-propyl, methylamino, dimethylamino, fluoro, cyclobutyl, 3,3,3-trifluoropropyl, or 2,2,2-trifluoroethyl, or methoxyethyl; and R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E20 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is bromo; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E21 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is bromo; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is phenyl, pyridyl, mono-, di- or tri-substituted phenyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl, or mono- or di-substituted pyridyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl.

Embodiment E22 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is bromo; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is 3,5-dichloro-4-fluorophenyl, 3,4,5-trichloro-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3-chloro-5-

5 trifluoromethyl-phenyl.

Embodiment E23 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is methyl; R^2 is methyl, ethyl, cyclopropyl, chloromethyl, fluoromethyl, difluoromethyl, iso-propyl, methylamino, dimethylamino, fluoro, cyclobutyl, 3,3,3-trifluoropropyl or 2,2,2-trifluoroethyl, or methoxyethyl; and R^4 is

10 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from

15 bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E24 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is methyl; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is

20 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from

25 bromo, chloro, fluoro or trifluoro; more preferably R^4 is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E25 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is methyl; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl

30 or fluoro; and R^4 is phenyl, pyridyl, mono-, di- or tri-substituted phenyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl, or mono- or di-substituted pyridyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl.

Embodiment E26 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is methyl; R^2 is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R^4 is

35 3,5-dichloro-4-fluorophenyl, 3,4,5-trichloro-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3-chloro-5-trifluoromethyl-phenyl.

Embodiment E27 of formula (I) provides compounds of formula (I) wherein A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^3 is chlorodifluoromethyl or trifluoromethyl; n is 2; A^1 is CR^5 , wherein R^5 is trifluoromethyl; R^2 is methyl, ethyl, cyclopropyl, chloromethyl, fluoromethyl, difluoromethyl, iso-propyl, methylamino, dimethylamino, fluoro, cyclobutyl, 3,3,3-trifluoropropyl or 2,2,2-trifluoroethyl, or methoxyethyl; and R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-

40 trifluoromethyl-phenyl.

trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R⁴ is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the

5 substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R⁴ is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E28 of formula (I) provides compounds of formula (I) wherein A², A³ and A⁴ are each CH; R¹ is hydrogen; R³ is trifluoromethyl; n is 2; A¹ is CR⁵, wherein R⁵ is trifluoromethyl; R² is methyl,

10 ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R⁴ is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro-phenyl or 3,4,5-trichloro-phenyl; especially R⁴ is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl,

15 3,5-dichloro-4-fluorophenyl, pyridyl, or 3-5-substituted pyridyl wherein the substituents are selected from bromo, chloro, fluoro or trifluoro; more preferably R⁴ is 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,5-bis-(trifluoromethyl)-phenyl, 4-bromo-3,5-dichlorophenyl, 3,4,5-trichloro-phenyl, 3,5-dichloro-4-fluorophenyl.

Embodiment E29 of formula (I) provides compounds of formula (I) wherein A², A³ and A⁴ are each CH; R¹ is hydrogen; R³ is chlorodifluoromethyl or trifluoromethyl; n is 2; A¹ is CR⁵, wherein R⁵ is

20 trifluoromethyl; R² is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R⁴ is phenyl, pyridyl, mono-, di- or tri-substituted phenyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl, or mono- or di-substituted pyridyl where the substituents are independently selected from chloro, bromo, fluoro, or trifluoromethyl.

Embodiment E30 of formula (I) provides compounds of formula (I) wherein A², A³ and A⁴ are each CH; R¹ is hydrogen; R³ is trifluoromethyl; n is 2; A¹ is CR⁵, wherein R⁵ is trifluoromethyl; R² is methyl, ethyl, cyclopropyl, methylamino, dimethylamino, chloromethyl, fluoromethyl, difluoromethyl or fluoro; and R⁴ is 3,5-dichloro-4-fluorophenyl, 3,4,5-trichloro-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3-chloro-5-trifluoromethyl-phenyl.

One preferred embodiment of formula (I) provides compounds of formula (I) wherein

A¹, A², A³ and A⁴ are independently of one another C-H, C-R⁵, or nitrogen;

R¹ is hydrogen;

R² is C₁-C₈alkyl or C₁-C₈alkyl substituted by one to three R^{6a}, C₁-C₈haloalkyl or C₁-C₈haloalkyl substituted by one to three R^{6a}, C₃-C₈cycloalkyl or C₃-C₈cycloalkyl substituted by one to three R^{6b}, phenyl, phenyl

35 substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷, -N(R⁸)(R⁹) or fluoro;

R³ is C₁-C₈haloalkyl;

R⁴ is phenyl or phenyl substituted by one to three R⁷;

40 R⁵ is independently halogen, cyano, nitro, C₁-C₈alkyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, C₂-C₈alkenyl;

R^{6a} is independently C₁-C₈alkoxy, or C₁-C₈haloalkoxy;

R^{6b} is independently oxo, amino, C_1 - C_8 alkylamino, N,N - C_1 - C_8 dialkylamino, C_1 - C_8 alkoxy, C_1 - C_8 haloalkoxy, phenyl, phenyl substituted by one to three R^7 , 5-6 membered heteroaryl or 5-6 membered heteroaryl substituted by one to three R^7 ;

R^7 is independently halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_1 - C_8 alkoxy, or C_1 - C_8 haloalkoxy;


5 R^8 and R^9 are independently hydrogen, cyano- C_1 - C_8 alkyl, C_1 - C_8 alkyl, C_1 - C_8 alkyl substituted by one to three R^{6a} , C_2 - C_8 alkenyl or C_2 - C_8 alkenyl substituted by one to three R^{6a} , phenyl- C_1 - C_4 alkyl, phenyl- C_1 - C_4 alkyl wherein the phenyl moiety is substituted by one to three R^7 , $-S(O)_2R^{10}$;

R^{10} is hydrogen, cyano- C_1 - C_8 alkyl, C_1 - C_8 alkyl or C_1 - C_8 alkyl substituted by one to three R^{6a} , phenyl, phenyl substituted by one to three R^7 , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to
10 three R^7 ;

n is 2;

or a tautomer, isomer, enantiomer, salt or N -oxide thereof.

Equally especially preferred compounds of formula (I) are represented by the compounds of formula (Ib)



15 wherein

R^a is hydrogen, halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, or C_1 - C_8 haloalkoxy;

R^b is hydrogen, halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, or C_1 - C_8 haloalkoxy;

R^c is hydrogen, halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, or C_1 - C_8 haloalkoxy;

20 R^5 is halogen, cyano, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, C_2 - C_8 alkenyl;

R^d is $S(O)_2-R^2$;

R^2 is C_1 - C_8 alkyl or C_1 - C_8 alkyl substituted by one to three R^{6a} , C_1 - C_8 haloalkyl or C_1 - C_8 haloalkyl substituted by one to three R^{6a} , C_3 - C_8 cycloalkyl or C_3 - C_8 cycloalkyl substituted by one to three R^{6b} , phenyl, phenyl substituted by one to three R^7 , phenyl- C_1 - C_4 alkyl, phenyl- C_1 - C_4 alkyl wherein the phenyl moiety is

25 substituted by one to three R^7 , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^7 , $-N(R^8)(R^9)$, halogen;

R^5 is independently halogen, cyano, nitro, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, C_2 - C_8 alkenyl;

R^{6a} is independently C_1 - C_8 alkoxy, or C_1 - C_8 haloalkoxy;

R^{6b} is independently oxo, amino, C_1 - C_8 alkylamino, N,N - C_1 - C_8 dialkylamino, C_1 - C_8 alkoxy, C_1 - C_8 haloalkoxy, phenyl, phenyl substituted by one to three R^7 , 5-6 membered heteroaryl or 5-6 membered heteroaryl substituted by one to three R^7 ;

R^7 is independently halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_1 - C_8 alkoxy, or C_1 - C_8 haloalkoxy;

R^8 and R^9 are independently hydrogen, cyano- C_1 - C_8 alkyl, C_1 - C_8 alkyl, C_1 - C_8 alkyl substituted by one to three R^{6a} , C_2 - C_8 alkenyl or C_2 - C_8 alkenyl substituted by one to three R^{6a} , phenyl- C_1 - C_4 alkyl, phenyl- C_1 -

35 C_4 alkyl wherein the phenyl moiety is substituted by one to three R^7 , $-S(O)_2R^{10}$;

R¹⁰ is hydrogen, cyano-C₁-C₈alkyl, C₁-C₈alkyl or C₁-C₈alkyl substituted by one to three R^{6a}, phenyl, phenyl substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷. Preferably at least one of the R^a, R^b and R^c is not hydrogen.

More preferably


- 5 R^a is hydrogen, halogen, cyano, C₁-C₈alkyl, C₁-C₈haloalkyl, or C₁-C₈haloalkoxy;
 R^b is hydrogen, halogen, cyano, C₁-C₈alkyl, C₁-C₈haloalkyl, or C₁-C₈haloalkoxy;
 R^c is hydrogen, halogen, cyano, C₁-C₈alkyl, C₁-C₈haloalkyl, or C₁-C₈haloalkoxy;
 R⁵ is halogen, cyano, C₁-C₈alkyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, C₂-C₈alkenyl;
 R^d is SO₂-CH₃, SO₂-4-chlorophenyl, SO₂-propyl, SO₂-CH₂Cl, SO₂-CH₂F, SO₂-CHF₂, SO₂-cyclopropyl,
 10 SO₂-CH₂CH₂-Cl, SO₂-CH₂CF₃, SO₂-2-thienyl, SO₂-N(CH₃)₂, SO₂-CH₂CH₂CF₃, SO₂-cyclobutyl, SO₂-2-propyl,
 SO₂-CH₂CH₂-O-CH₃, SO₂CH₂CH₃, SO₂F, SO₂-NH₂, SO₂-NH(CH₃), SO₂-N(CH₃)(CH₂CH₃), SO₂-N(CH₃)(C(=O)CH₃),
 SO₂-N(CH₃)C(=O)N(CH₃)₂, SO₂-N(CH₃)C(=O)OCH₂CH₃, SO₂-N(CH₃)C(=O)OCH₃,
 SO₂-N(CH₃)C(=O)(3-pyridyl), SO₂-N(CH₃)SO₂N(CH₃)₂, SO₂-N(CH₃)C(=O)Ph, SO₂-N(CH₃)CN, SO₂-N(CH₃)CH₂CHCH₂,
 15 SO₂-N(CH₃)CH₂COCH₂, SO₂-N(CH₃)CH₂CH₂CH₃, SO₂-N(CH₃)CH₂CH₂CH₂CH₃, SO₂-N(CH₃)CH₂Ph,
 SO₂-N(CH₃)CH₂CN, SO₂-N(CH₃)CH₂CO₂CH₂CH₃, SO₂-N(CH₃)CH₂CH₂OCH₃, SO₂-N(CH₂CH₃)CO₂C(CH₃)₃,
 SO₂-N(CH₃)SO₂CH₃, SO₂-NHCH₂CH₃, SO₂-N(CH₃)CH₂OCH₃, SO₂-NHC(CH₃)₃,
 SO₂-N(CH₂CH₃)₂, SO₂-N(CH₂)₄, SO₂-N(CH₃)C(CH₃)₃, SO₂-NHCH(CH₃)₂, SO₂-NHC(=O)CH₃, SO₂-NHSO₂CH₃,
 SO₂-N(CH₃)CH(CH₃)₂, SO₂-N(CH₂CH₃)CH(CH₃)₂ or SO₂-NHCH₂CHCH₂. Preferably at least one of the R^a, R^b and R^c is not hydrogen. More preferably at least two of the R^a, R^b and R^c are not
 20 hydrogen.

Equally more preferably

- R^a is H, Cl, Br, F, CF₃, CH₃ or OCF₃;
 R^b is H, Cl, Br, F, CF₃, CH₃ or OCF₃;
 R^c is H, Cl, Br, F, CF₃, CH₃ or OCF₃;
 25 R⁵ is CH₃, Cl, Br, CN, cyclopropyl, CF₃, CHCH₂;
 R^d is SO₂-CH₃, SO₂-4-chlorophenyl, SO₂-propyl, SO₂-CH₂Cl, SO₂-CH₂F, SO₂-CHF₂, SO₂-cyclopropyl,
 SO₂-CH₂CH₂-Cl, SO₂-CH₂CF₃, SO₂-2-thienyl, SO₂-N(CH₃)₂, SO₂-CH₂CH₂CF₃, SO₂-cyclobutyl, SO₂-2-propyl,
 SO₂-CH₂CH₂-O-CH₃, SO₂CH₂CH₃, SO₂F, SO₂-NH₂, SO₂-NH(CH₃), SO₂-N(CH₃)(CH₂CH₃), SO₂-N(CH₃)(C(=O)CH₃),
 SO₂-N(CH₃)C(=O)N(CH₃)₂, SO₂-N(CH₃)C(=O)OCH₂CH₃, SO₂-N(CH₃)C(=O)OCH₃,
 30 SO₂-N(CH₃)C(=O)(3-pyridyl), SO₂-N(CH₃)SO₂N(CH₃)₂, SO₂-N(CH₃)C(=O)Ph, SO₂-N(CH₃)CN, SO₂-N(CH₃)CH₂CHCH₂,
 SO₂-N(CH₃)CH₂COCH₂, SO₂-N(CH₃)CH₂CH₂CH₃, SO₂-N(CH₃)CH₂CH₂CH₂CH₃, SO₂-N(CH₃)CH₂Ph,
 SO₂-N(CH₃)CH₂CN, SO₂-N(CH₃)CH₂CO₂CH₂CH₃, SO₂-N(CH₃)CH₂CH₂OCH₃, SO₂-N(CH₂CH₃)CO₂C(CH₃)₃,
 SO₂-N(CH₃)SO₂CH₃, SO₂-NHCH₂CH₃, SO₂-N(CH₃)CH₂OCH₃, SO₂-NHC(CH₃)₃,
 SO₂-N(CH₂CH₃)₂, SO₂-N(CH₂)₄, SO₂-N(CH₃)C(CH₃)₃, SO₂-NHCH(CH₃)₂, SO₂-NHC(=O)CH₃, SO₂-NHSO₂CH₃,
 35 SO₂-N(CH₃)CH(CH₃)₂, SO₂-N(CH₂CH₃)CH(CH₃)₂ or SO₂-NHCH₂CHCH₂. Preferably at least one of the R^a, R^b and R^c is not hydrogen. More preferably at least two of the R^a, R^b and R^c are not hydrogen.


The present invention also provides intermediates useful for the preparation of compounds of formula (I). Certain intermediates are novel and as such form a further aspect of the invention. One

- 40 group of novel intermediates are compounds of formula (Int-I)




wherein A^1 , A^2 , A^3 , A^4 , R^1 , R^2 and n are as defined for a compound of formula (I) and X^B is a halogen, such as bromo, or X^B is cyano, formyl, $CH=N-OH$ or acetyl; or a salt or *N*-oxide thereof. The preferences for A^1 , A^2 , A^3 , A^4 , R^1 , R^2 and n are the same as the preferences set out for the corresponding substituents of a compound of formula (I).

Another group of novel intermediates are compounds of formula (Int-II)





wherein A^1 , A^2 , A^3 , A^4 , R^1 , R^2 and n are as defined for a compound of formula (I); X^C is CH_2 -halogen, wherein halogen is preferably bromo or chloro, $CH=C(R^3)R^4$ or $CH_2C(OH)(R^3)R^4$ wherein R^3 and R^4 are as defined for a compound of formula (I); or a salt or *N*-oxide thereof. The preferences for A^1 , A^2 , A^3 , A^4 , R^1 , R^2 and n are the same as the preferences set out for the corresponding substituents of a compound of formula (I).

Another group of novel intermediates are compounds of formula (Int-III)



wherein R^1 , R^2 and n are as defined for a compound of formula (I); or a salt or *N*-oxide thereof. The preferences for R^1 , R^2 and n are the same as the preferences set out for the corresponding substituents of a compound of formula (I).

Compounds of formula (I) include at least one chiral centre and may exist as compounds of formula (I*) or compounds of formula (I**):



Generally compounds of formula (I**) are more biologically active than compounds of formula (I*).

The invention includes mixtures of compounds (I*) and (I**) in any ratio e.g. in a molar ratio of 1:99 to


5 99:1, e.g. 10:1 to 1:10, e.g. a substantially 50:50 molar ratio. In an enantiomerically (or epimerically) enriched mixture of formula (I**), the molar proportion of compound (I**) compared to the total amount of both enantiomers (or epimers) is for example greater than 50%, e.g. at least 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or at least 99%. Likewise, in enantiomerically (or epimerically) enriched mixture of formula (I*), the molar proportion of the compound of formula (I*) compared to the total amount of both

10 enantiomers (or epimers) is for example greater than 50%, e.g. at least 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or at least 99%. Enantiomerically (or epimerically) enriched mixtures of formula (I**) are preferred.

Tables 1 to 96: Compounds of formula (Ia)

The invention is further illustrated by making available the following individual compounds of

15 formula (Ia) listed below in Tables 1 to 96.



Each of Tables 1 to 96, which follow the Table P below, make available 250 compounds of the formula (Ia) in which X₃, R³, X₁ and R⁵ are the substituents defined in Table P and R² and X₂ are the

20 substituents defined in the relevant Table 1 to 96. Thus Table 1 individualises 250 compounds of formula (Ia) wherein for each row of Table P, the R² and X₂ substituents are as defined in Table 1; similarly, Table 2 individualises 250 compounds of formula (Ia) wherein for each row of Table P, the R² and X₂ substituents are as defined in Table 2; and so on for Tables 3 to 96.

Each compound disclosed in Tables 1 to 96 represents a disclosure of a compound according to

25 the compound of formula (I*), and a disclosure according to the compound of formula (I**) as well as mixtures thereof.

Table P

	X ₃	R ³	X ₁	R ⁵
1	H	CF ₃	H	Me
2	Cl	CF ₃	H	Me
3	Br	CF ₃	H	Me
4	F	CF ₃	H	Me
5	CF ₃	CF ₃	H	Me
6	H	CF ₂ Cl	H	Me
7	Cl	CF ₂ Cl	H	Me
8	Br	CF ₂ Cl	H	Me
9	F	CF ₂ Cl	H	Me
10	CF ₃	CF ₂ Cl	H	Me
11	H	CF ₃	Cl	Me
12	Cl	CF ₃	Cl	Me
13	Br	CF ₃	Cl	Me
14	F	CF ₃	Cl	Me
15	CF ₃	CF ₃	Cl	Me
16	H	CF ₂ Cl	Cl	Me
17	Cl	CF ₂ Cl	Cl	Me
18	Br	CF ₂ Cl	Cl	Me
19	F	CF ₂ Cl	Cl	Me
20	CF ₃	CF ₂ Cl	Cl	Me
21	H	CF ₃	Br	Me
22	Cl	CF ₃	Br	Me
23	Br	CF ₃	Br	Me
24	F	CF ₃	Br	Me
25	CF ₃	CF ₃	Br	Me
26	H	CF ₂ Cl	Br	Me
27	Cl	CF ₂ Cl	Br	Me
28	Br	CF ₂ Cl	Br	Me
29	F	CF ₂ Cl	Br	Me
30	CF ₃	CF ₂ Cl	Br	Me
31	H	CF ₃	F	Me
32	Cl	CF ₃	F	Me
33	Br	CF ₃	F	Me
34	F	CF ₃	F	Me
35	CF ₃	CF ₃	F	Me
36	H	CF ₂ Cl	F	Me
37	Cl	CF ₂ Cl	F	Me
38	Br	CF ₂ Cl	F	Me

	X ₃	R ³	X ₁	R ⁵
39	F	CF ₂ Cl	F	Me
40	CF ₃	CF ₂ Cl	F	Me
41	H	CF ₃	CF ₃	Me
42	Cl	CF ₃	CF ₃	Me
43	Br	CF ₃	CF ₃	Me
44	F	CF ₃	CF ₃	Me
45	CF ₃	CF ₃	CF ₃	Me
46	H	CF ₂ Cl	CF ₃	Me
47	Cl	CF ₂ Cl	CF ₃	Me
48	Br	CF ₂ Cl	CF ₃	Me
49	F	CF ₂ Cl	CF ₃	Me
50	CF ₃	CF ₂ Cl	CF ₃	Me
51	H	CF ₃	H	Cl
52	Cl	CF ₃	H	Cl
53	Br	CF ₃	H	Cl
54	F	CF ₃	H	Cl
55	CF ₃	CF ₃	H	Cl
56	H	CF ₂ Cl	H	Cl
57	Cl	CF ₂ Cl	H	Cl
58	Br	CF ₂ Cl	H	Cl
59	F	CF ₂ Cl	H	Cl
60	CF ₃	CF ₂ Cl	H	Cl
61	H	CF ₃	Cl	Cl
62	Cl	CF ₃	Cl	Cl
63	Br	CF ₃	Cl	Cl
64	F	CF ₃	Cl	Cl
65	CF ₃	CF ₃	Cl	Cl
66	H	CF ₂ Cl	Cl	Cl
67	Cl	CF ₂ Cl	Cl	Cl
68	Br	CF ₂ Cl	Cl	Cl
69	F	CF ₂ Cl	Cl	Cl
70	CF ₃	CF ₂ Cl	Cl	Cl
71	H	CF ₃	Br	Cl
72	Cl	CF ₃	Br	Cl
73	Br	CF ₃	Br	Cl
74	F	CF ₃	Br	Cl
75	CF ₃	CF ₃	Br	Cl
76	H	CF ₂ Cl	Br	Cl
77	Cl	CF ₂ Cl	Br	Cl

	X ₃	R ³	X ₁	R ⁵
78	Br	CF ₂ Cl	Br	Cl
79	F	CF ₂ Cl	Br	Cl
80	CF ₃	CF ₂ Cl	Br	Cl
81	H	CF ₃	F	Cl
82	Cl	CF ₃	F	Cl
83	Br	CF ₃	F	Cl
84	F	CF ₃	F	Cl
85	CF ₃	CF ₃	F	Cl
86	H	CF ₂ Cl	F	Cl
87	Cl	CF ₂ Cl	F	Cl
88	Br	CF ₂ Cl	F	Cl
89	F	CF ₂ Cl	F	Cl
90	CF ₃	CF ₂ Cl	F	Cl
91	H	CF ₃	CF ₃	Cl
92	Cl	CF ₃	CF ₃	Cl
93	Br	CF ₃	CF ₃	Cl
94	F	CF ₃	CF ₃	Cl
95	CF ₃	CF ₃	CF ₃	Cl
96	H	CF ₂ Cl	CF ₃	Cl
97	Cl	CF ₂ Cl	CF ₃	Cl
98	Br	CF ₂ Cl	CF ₃	Cl
99	F	CF ₂ Cl	CF ₃	Cl
100	CF ₃	CF ₂ Cl	CF ₃	Cl
101	H	CF ₃	H	Br
102	Cl	CF ₃	H	Br
103	Br	CF ₃	H	Br
104	F	CF ₃	H	Br
105	CF ₃	CF ₃	H	Br
106	H	CF ₂ Cl	H	Br
107	Cl	CF ₂ Cl	H	Br
108	Br	CF ₂ Cl	H	Br
109	F	CF ₂ Cl	H	Br
110	CF ₃	CF ₂ Cl	H	Br
111	H	CF ₃	Cl	Br
112	Cl	CF ₃	Cl	Br
113	Br	CF ₃	Cl	Br
114	F	CF ₃	Cl	Br
115	CF ₃	CF ₃	Cl	Br
116	H	CF ₂ Cl	Cl	Br

	X ₃	R ³	X ₁	R ⁵
117	Cl	CF ₂ Cl	Cl	Br
118	Br	CF ₂ Cl	Cl	Br
119	F	CF ₂ Cl	Cl	Br
120	CF ₃	CF ₂ Cl	Cl	Br
121	H	CF ₃	Br	Br
122	Cl	CF ₃	Br	Br
123	Br	CF ₃	Br	Br
124	F	CF ₃	Br	Br
125	CF ₃	CF ₃	Br	Br
126	H	CF ₂ Cl	Br	Br
127	Cl	CF ₂ Cl	Br	Br
128	Br	CF ₂ Cl	Br	Br
129	F	CF ₂ Cl	Br	Br
130	CF ₃	CF ₂ Cl	Br	Br
131	H	CF ₃	F	Br
132	Cl	CF ₃	F	Br
133	Br	CF ₃	F	Br
134	F	CF ₃	F	Br
135	CF ₃	CF ₃	F	Br
136	H	CF ₂ Cl	F	Br
137	Cl	CF ₂ Cl	F	Br
138	Br	CF ₂ Cl	F	Br
139	F	CF ₂ Cl	F	Br
140	CF ₃	CF ₂ Cl	F	Br
141	H	CF ₃	CF ₃	Br
142	Cl	CF ₃	CF ₃	Br
143	Br	CF ₃	CF ₃	Br
144	F	CF ₃	CF ₃	Br
145	CF ₃	CF ₃	CF ₃	Br
146	H	CF ₂ Cl	CF ₃	Br
147	Cl	CF ₂ Cl	CF ₃	Br
148	Br	CF ₂ Cl	CF ₃	Br
149	F	CF ₂ Cl	CF ₃	Br
150	CF ₃	CF ₂ Cl	CF ₃	Br
151	H	CF ₃	H	CF ₃
152	Cl	CF ₃	H	CF ₃
153	Br	CF ₃	H	CF ₃
154	F	CF ₃	H	CF ₃
155	CF ₃	CF ₃	H	CF ₃

	X ₃	R ³	X ₁	R ⁵
156	H	CF ₂ Cl	H	CF ₃
157	Cl	CF ₂ Cl	H	CF ₃
158	Br	CF ₂ Cl	H	CF ₃
159	F	CF ₂ Cl	H	CF ₃
160	CF ₃	CF ₂ Cl	H	CF ₃
161	H	CF ₃	Cl	CF ₃
162	Cl	CF ₃	Cl	CF ₃
163	Br	CF ₃	Cl	CF ₃
164	F	CF ₃	Cl	CF ₃
165	CF ₃	CF ₃	Cl	CF ₃
166	H	CF ₂ Cl	Cl	CF ₃
167	Cl	CF ₂ Cl	Cl	CF ₃
168	Br	CF ₂ Cl	Cl	CF ₃
169	F	CF ₂ Cl	Cl	CF ₃
170	CF ₃	CF ₂ Cl	Cl	CF ₃
171	H	CF ₃	Br	CF ₃
172	Cl	CF ₃	Br	CF ₃
173	Br	CF ₃	Br	CF ₃
174	F	CF ₃	Br	CF ₃
175	CF ₃	CF ₃	Br	CF ₃
176	H	CF ₂ Cl	Br	CF ₃
177	Cl	CF ₂ Cl	Br	CF ₃
178	Br	CF ₂ Cl	Br	CF ₃
179	F	CF ₂ Cl	Br	CF ₃
180	CF ₃	CF ₂ Cl	Br	CF ₃
181	H	CF ₃	F	CF ₃
182	Cl	CF ₃	F	CF ₃
183	Br	CF ₃	F	CF ₃
184	F	CF ₃	F	CF ₃
185	CF ₃	CF ₃	F	CF ₃
186	H	CF ₂ Cl	F	CF ₃
187	Cl	CF ₂ Cl	F	CF ₃
188	Br	CF ₂ Cl	F	CF ₃
189	F	CF ₂ Cl	F	CF ₃
190	CF ₃	CF ₂ Cl	F	CF ₃
191	H	CF ₃	CF ₃	CF ₃
192	Cl	CF ₃	CF ₃	CF ₃
193	Br	CF ₃	CF ₃	CF ₃
194	F	CF ₃	CF ₃	CF ₃

	X ₃	R ³	X ₁	R ⁵
195	CF ₃	CF ₃	CF ₃	CF ₃
196	H	CF ₂ Cl	CF ₃	CF ₃
197	Cl	CF ₂ Cl	CF ₃	CF ₃
198	Br	CF ₂ Cl	CF ₃	CF ₃
199	F	CF ₂ Cl	CF ₃	CF ₃
200	CF ₃	CF ₂ Cl	CF ₃	CF ₃
201	H	CF ₃	H	F
202	Cl	CF ₃	H	F
203	Br	CF ₃	H	F
204	F	CF ₃	H	F
205	CF ₃	CF ₃	H	F
206	H	CF ₂ Cl	H	F
207	Cl	CF ₂ Cl	H	F
208	Br	CF ₂ Cl	H	F
209	F	CF ₂ Cl	H	F
210	CF ₃	CF ₂ Cl	H	F
211	H	CF ₃	Cl	F
212	Cl	CF ₃	Cl	F
213	Br	CF ₃	Cl	F
214	F	CF ₃	Cl	F
215	CF ₃	CF ₃	Cl	F
216	H	CF ₂ Cl	Cl	F
217	Cl	CF ₂ Cl	Cl	F
218	Br	CF ₂ Cl	Cl	F
219	F	CF ₂ Cl	Cl	F
220	CF ₃	CF ₂ Cl	Cl	F
221	H	CF ₃	Br	F
222	Cl	CF ₃	Br	F
223	Br	CF ₃	Br	F
224	F	CF ₃	Br	F
225	CF ₃	CF ₃	Br	F
226	H	CF ₂ Cl	Br	F
227	Cl	CF ₂ Cl	Br	F
228	Br	CF ₂ Cl	Br	F
229	F	CF ₂ Cl	Br	F
230	CF ₃	CF ₂ Cl	Br	F
231	H	CF ₃	F	F
232	Cl	CF ₃	F	F
233	Br	CF ₃	F	F

	X ₃	R ³	X ₁	R ⁵
234	F	CF ₃	F	F
235	CF ₃	CF ₃	F	F
236	H	CF ₂ Cl	F	F
237	Cl	CF ₂ Cl	F	F
238	Br	CF ₂ Cl	F	F
239	F	CF ₂ Cl	F	F
240	CF ₃	CF ₂ Cl	F	F
241	H	CF ₃	CF ₃	F
242	Cl	CF ₃	CF ₃	F
243	Br	CF ₃	CF ₃	F
244	F	CF ₃	CF ₃	F
245	CF ₃	CF ₃	CF ₃	F
246	H	CF ₂ Cl	CF ₃	F
247	Cl	CF ₂ Cl	CF ₃	F
248	Br	CF ₂ Cl	CF ₃	F
249	F	CF ₂ Cl	CF ₃	F
250	CF ₃	CF ₂ Cl	CF ₃	F

Table 1

Table 1 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is Me, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

5 Table 2

Table 2 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is Me, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 3

10 Table 3 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is Me, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 4

Table 4 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is Me, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 5

15 Table 5 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is Me, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 6

Table 6 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is Me, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

20 Table 7

Table 7 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is Et, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 8

25 Table 8 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is Et, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 9

Table 9 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is Et, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 10

30 Table 10 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is Et, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 11

Table 11 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is Et, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

35 Table 12

Table 12 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is Et, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 13

40 Table 13 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is Pr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 14

Table 14 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is Pr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 15

Table 15 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is Pr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 16

Table 16 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is Pr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 17

Table 17 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is Pr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 18

Table 18 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is Pr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 19

Table 19 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is N(Et)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 20

Table 20 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is N(Et)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 21

Table 21 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is N(Et)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 22

Table 22 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is N(Et)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 23

Table 23 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is N(Et)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 24

Table 24 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is N(Et)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 25

Table 25 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is iPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 26

Table 26 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is iPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 27

Table 27 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is iPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 28

Table 28 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is iPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 29

Table 29 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is iPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 30

Table 30 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is iPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 31

Table 31 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is N(Me)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 32

Table 32 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is N(Me)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 33

Table 33 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is N(Me)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 34

Table 34 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is N(Me)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 35

Table 35 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is N(Me)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 36

Table 36 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is N(Me)₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 37

Table 37 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is cBu, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 38

Table 38 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is cBu, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 39

Table 39 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is cBu, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 40

Table 40 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is cBu, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 41

Table 41 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is cBu, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 42

Table 42 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is cBu, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 43

Table 43 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is NHEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 44

Table 44 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is NHEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 45

Table 45 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is NHEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 46

Table 46 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is NHEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 47

Table 47 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is NHEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 48

Table 48 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is NHEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 49

Table 49 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is CH₂CF₃, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 50

Table 50 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is CH₂CF₃, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 51

Table 51 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is CH₂CF₃, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 52

Table 52 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is CH₂CF₃, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 53

Table 53 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is CH₂CF₃, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 54

Table 54 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is CH₂CF₃, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 55

Table 55 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is NHMe, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 56

Table 56 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is NHMe, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 57

Table 57 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is NHMe, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 58

Table 58 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is NHMe, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 59

10 Table 59 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is NHMe, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 60

Table 60 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is NHMe, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

15 Table 61

Table 61 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is NMeEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 62

20 Table 62 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is NMeEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 63

Table 63 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is NMeEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 64

25 Table 64 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is NMeEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 65

Table 65 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is NMeEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

30 Table 66

Table 66 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is NMeEt, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 67

35 Table 67 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is cPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 68

Table 68 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is cPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 69

40 Table 69 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is cPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 70

Table 70 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is cPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 71

Table 71 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is cPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 72

Table 72 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is cPr, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 73

Table 73 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is CH₂Cl, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 74

Table 74 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is CH₂Cl, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 75

Table 75 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is CH₂Cl, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 76

Table 76 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is CH₂Cl, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 77

Table 77 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is CH₂Cl, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 78

Table 78 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is CH₂Cl, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 79

Table 79 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is CH₂F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 80

Table 80 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is CH₂F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 81

Table 81 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is CH₂F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 82

Table 82 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is CH₂F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 83

Table 83 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is CH₂F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 84

Table 84 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is CH₂F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 85

Table 85 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is CHF₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 86

Table 86 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is CHF₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 87

Table 87 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is CHF₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 88

Table 88 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is CHF₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 89

Table 89 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is CHF₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 90

Table 90 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is CHF₂, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 91

Table 91 provides 250 compounds of formula (Ia) wherein X_2 is N, R^2 is F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 92

Table 92 provides 250 compounds of formula (Ia) wherein X_2 is CH, R^2 is F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 93

Table 93 provides 250 compounds of formula (Ia) wherein X_2 is C-F, R^2 is F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 94

Table 94 provides 250 compounds of formula (Ia) wherein X_2 is C-Br, R^2 is F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 95

Table 95 provides 250 compounds of formula (Ia) wherein X_2 is C-Cl, R^2 is F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Table 96

Table 96 provides 250 compounds of formula (Ia) wherein X_2 is C-CF₃, R^2 is F, and X_3 , R^3 , X_1 , R^5 are as defined in Table P.

Examples of compounds of formula (Int-I) made available are those where X^B is bromo, chloro, iodo, cyano, formyl, CH=NOH or acetyl, A^2 , A^3 and A^4 are each CH, R^1 is hydrogen, n is 2, A^1 is CR⁵, and wherein R^5 and R^2 each correspond to a substituent R^5 and R^2 respectively as defined in each of Tables 1 to 96 above in context of formula (Ia). So for example, Table 1 individualises a compound of formula


(Int-I) wherein X^B is bromo, chloro, iodo, cyano, formyl, $CH=NOH$ or acetyl, A_2 , A_3 and A_4 are each CH , R^1 is hydrogen, n is 2, A^1 is CR^5 , and wherein R^5 and R^2 are each as defined in Table 1; similarly, Table 2 individualises a compound of formula (Int-I) wherein X^B is bromo, chloro, iodo, cyano, formyl, $CH=NOH$ or acetyl, A^2 , A^3 and A^4 are each CH , R^1 is hydrogen, n is 2, A^1 is CR^5 , and wherein R^5 and R^2 are each as defined in Table 2; and so on for Tables 3 to 96.

Examples of compounds of formula (Int-II) made available are those where X^C is CH_2Cl , CH_2Br , $CH=C(CF_3)(3\text{-chloro-5-trifluoromethyl-phenyl})$, $CH=C(CF_3)(3\text{-bromo-5-trifluoromethyl-phenyl})$, $CH=C(CF_3)(3,5\text{-dichloro-4-fluoro-phenyl})$, $CH=C(CF_3)(3,4,5\text{-trichloro-phenyl})$, $CH_2C(OH)(CF_3)(3\text{-chloro-5-trifluoromethyl-phenyl})$, $CH_2C(OH)(CF_3)(3\text{-bromo-5-trifluoromethyl-phenyl})$, $CH_2C(OH)(CF_3)(3,5\text{-dichloro-4-fluoro-phenyl})$ or $CH_2C(OH)(CF_3)(3,4,5\text{-trichloro-phenyl})$, A_2 , A_3 and A_4 are each CH , R^1 is hydrogen, n is 2, A^1 is CR^5 , and wherein R^5 and R^2 each correspond to a substituent R^5 and R^2 respectively as defined in each of Tables 1 to 96 above in context of formula (Ia). So for example, Table 1 individualises a compound of formula (Int-II) wherein X^C is CH_2Cl , CH_2Br , $CH=C(CF_3)(3\text{-chloro-5-trifluoromethyl-phenyl})$, $CH=C(CF_3)(3\text{-bromo-5-trifluoromethyl-phenyl})$, $CH=C(CF_3)(3,5\text{-dichloro-4-fluoro-phenyl})$, $CH=C(CF_3)(3,4,5\text{-trichloro-phenyl})$, $CH_2C(OH)(CF_3)(3\text{-chloro-5-trifluoromethyl-phenyl})$, $CH_2C(OH)(CF_3)(3\text{-bromo-5-trifluoromethyl-phenyl})$, $CH_2C(OH)(CF_3)(3,5\text{-dichloro-4-fluoro-phenyl})$ or $CH_2C(OH)(CF_3)(3,4,5\text{-trichloro-phenyl})$, A_2 , A_3 and A_4 are each CH , R^1 is hydrogen, n is 2, A^1 is CR^5 , and wherein R^5 and R^2 are each as defined in Table 1; similarly, Table 2 individualises a compound of formula (Int-II) wherein X^C is CH_2Cl , CH_2Br , $CH=C(CF_3)(3\text{-chloro-5-trifluoromethyl-phenyl})$, $CH=C(CF_3)(3\text{-bromo-5-trifluoromethyl-phenyl})$, $CH=C(CF_3)(3,5\text{-dichloro-4-fluoro-phenyl})$, $CH=C(CF_3)(3,4,5\text{-trichloro-phenyl})$, $CH_2C(OH)(CF_3)(3\text{-chloro-5-trifluoromethyl-phenyl})$, $CH_2C(OH)(CF_3)(3\text{-bromo-5-trifluoromethyl-phenyl})$, $CH_2C(OH)(CF_3)(3,5\text{-dichloro-4-fluoro-phenyl})$ or $CH_2C(OH)(CF_3)(3,4,5\text{-trichloro-phenyl})$, A_2 , A_3 and A_4 are each CH , R^1 is hydrogen, n is 2, A^1 is CR^5 , and wherein R^5 and R^2 are each as defined in Table 2; and so on for Tables 3 to 96.

Examples of compounds of formula (Int-III) made available are those where R^1 is hydrogen, n is 2, and wherein R^2 corresponds to a substituent R^2 as defined in each of Tables 1 to 96 above in context of formula (Ia). So for example, Table 1 individualises a compound of formula (Int-III) wherein R^1 is hydrogen, n is 2, and R^2 is as defined in Table 1; similarly, Table 2 individualises a compound of formula (Int-III) wherein R^1 is hydrogen, n is 2, and wherein R^2 is as defined in Table 2; and so on for Tables 3 to 96.

The compounds of the invention may be made by a variety of methods as shown in Schemes 1 and 2.

Scheme 1



1) Compounds of formula (I) can be prepared by reacting a compound of formula (II) wherein R is OH, C₁-C₆alkoxy or Cl, F or Br, with an amine of formula (III) wherein R^G is S(O)_nR² or a protecting group, such as a Boc, as shown in **Scheme 1**. When R is OH such reactions are usually carried out in the presence of a coupling reagent, such as N,N'-dicyclohexylcarbodiimide ("DCC"), 1-ethyl-3-(3-dimethyl-amino-propyl)carbodiimide hydrochloride ("EDC") or bis(2-oxo-3-oxazolidinyl)phosphonic chloride ("BOP-Cl"), in the presence of a base, and optionally in the presence of a nucleophilic catalyst, such as hydroxybenzotriazole ("HOBT"). When R is Cl, such reactions are usually carried out in the presence of a base, and optionally in the presence of a nucleophilic catalyst. Alternatively, it is possible to conduct the reaction in a biphasic system comprising an organic solvent, preferably ethyl acetate, and an aqueous solvent, preferably a solution of sodium hydrogen carbonate. When R is C₁-C₆alkoxy it is sometimes possible to convert the ester directly to the amide by heating the ester and amine together in a thermal process. Suitable bases include pyridine, triethylamine, 4-(dimethylamino)-pyridine ("DMAP") or diisopropylethylamine (Hunig's base). Preferred solvents are *N,N*-dimethylacetamide, tetrahydrofuran, dioxane, 1,2-dimethoxyethane, ethyl acetate and toluene. The reaction is carried out at a temperature of from 0°C to 100°C, preferably from 15°C to 30°C, in particular at ambient temperature. Amines of formula (III) are either known in the literature or can be prepared using methods known to a person skilled in the art. Some of these methods are described in the preparation Examples.

2) Acid halides of formula (II), wherein R is Cl, F or Br, may be made from carboxylic acids of formula (II), wherein R is OH, under standard conditions, as described for example in WO2009080250.


3) Carboxylic acids of formula (II), wherein R is OH, may be formed from esters of formula (II), wherein R is C₁-C₆alkoxy as described for example in WO2009080250.

4) Compounds of formula (I) can be prepared by reacting a compound of formula (IV) wherein X^B is a leaving group, for example a halogen, such as bromo, with carbon monoxide and an amine of formula (III), in the presence of a catalyst, such as palladium(II) acetate or bis(triphenylphosphine)palladium(II) dichloride, optionally in the presence of a ligand, such as triphenylphosphine, and a base, such as sodium carbonate, pyridine, triethylamine, 4-(dimethylamino)-pyridine ("DMAP") or diisopropylethylamine (Hunig's base), in a solvent, such as water, *N,N*-dimethylformamide or tetrahydrofuran. The reaction is carried out

at a temperature of from 50°C to 200°C, preferably from 100°C to 150°C. The reaction is carried out at a pressure of from 50 to 200 bar, preferably from 100 to 150 bar.


5) Compounds of formula (IV) wherein X^B is a leaving group, for example a halogen, such as bromo, can be made by a various of methods, for example as described in WO2009080250.

5 Scheme 2



6) Alternatively, compounds of formula (I) can be prepared by various methods from an intermediate of formula (V) as shown in **Scheme 2** wherein X^B is a leaving group, for example a halogen, such as bromo, or X^B is cyano, formyl or acetyl according to similar methods to those described in
 10 WO2009080250. An intermediate of formula (V) can be prepared for example from an intermediate of formula (VI) as described in the same reference.

Scheme 3



15 7) Alternatively, compounds of formula (I) can be prepared by various methods from an intermediate of formula (VII) as shown in **Scheme 3** wherein X^C is $CH=C(R^3)R^4$, or $CH_2C(OH)(R^3)R^4$

wherein R^3 and R^4 are as defined for a compound of formula (I) according to similar methods to those described in WO2009080250.

8) Compounds of formula (VII) wherein X^C is $CH=C(R^3)R^4$, or $CH_2C(OH)(R^3)R^4$ can be prepared from a compound of formula (Va) or from a compound of formula (VII) wherein X^C is CH_2 -halogen using similar methods to those described in WO2009080250.

9) Compounds of formula (VII) wherein X^C is CH_2 -halogen, such as bromo or chloro, can be prepared by reacting a methyl ketone of formula (Va), with a halogenating agent, such as bromine or chlorine, in a solvent, such as acetic acid, at a temperature of from 0°C to 50°C , preferably from ambient temperature to 40°C .

10) Compounds of formula (III) are either known compounds or can be prepared by known methods to the person skilled in the art. Examples of such methods can be found in the Examples below.

The compounds of formula (I) according to the invention are preventively and/or curatively valuable active ingredients in the field of pest control, even at low rates of application, which have a very favorable biocidal spectrum and are well tolerated by warm-blooded species, fish and plants. The pests which may be combated and controlled by the use of the compounds of the invention include those pests associated with agriculture (which term includes the growing of crops for food and fiber products), horticulture and animal husbandry, companion animals, forestry and the storage of products of vegetable origin (such as fruit, grain and timber); those pests associated with the damage of man-made structures and the transmission of diseases of man and animals; and also nuisance pests (such as flies). The active ingredients according to the invention act against all or individual developmental stages of normally sensitive, but also resistant, animal pests, such as insects or representatives of the order Acarina. The insecticidal or acaricidal activity of the active ingredients according to the invention can manifest itself directly, i. e. in destruction of the pests, which takes place either immediately or only after some time has elapsed, for example during ecdysis, or indirectly, for example in a reduced oviposition and/or hatching rate, a good activity corresponding to a destruction rate (mortality) of at least 50 to 60%.

Examples of the abovementioned animal pests are:

from the order *Acarina*, for example,

Acalitus spp, *Aculus* spp, *Acaricalus* spp, *Aceria* spp, *Acarus* siro, *Amblyomma* spp., *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., *Bryobia* spp, *Calipitrimerus* spp., *Chorioptes* spp., *Dermanyssus* gallinae, *Dermatophagoides* spp, *Eotetranychus* spp, *Eriophyes* spp., *Hemitarsonemus* spp, *Hyalomma* spp., *Ixodes* spp., *Olygonychus* spp, *Ornithodoros* spp., *Polyphagotarsonus latus*, *Panonychus* spp., *Phyllocoptura oleivora*, *Phytonemus* spp, *Polyphagotarsonemus* spp, *Psoroptes* spp., *Rhipicephalus* spp., *Rhizoglyphus* spp., *Sarcoptes* spp., *Steneotarsonemus* spp, *Tarsonemus* spp. and *Tetranychus* spp.;

from the order *Anoplura*, for example,

Haematopinus spp., *Linognathus* spp., *Pediculus* spp., *Pemphigus* spp. and *Phylloxera* spp.;

from the order *Coleoptera*, for example,

Agriotes spp., *Amphimallon majale*, *Anomala orientalis*, *Anthonomus* spp., *Aphodius* spp, *Astylus atomaculatus*, *Ataenius* spp, *Atomaria linearis*, *Chaetocnema tibialis*, *Cerotoma* spp, *Conoderus* spp, *Cosmopolites* spp., *Cotinis nitida*, *Curculio* spp., *Cyclocephala* spp, *Dermestes* spp., *Diabrotica* spp., *Diloboderus abderus*, *Epilachna* spp., *Eremnus* spp., *Heteronychus arator*, *Hypothenemus hampei*, *Lagria vilosa*, *Leptinotarsa decemlineata*, *Lissorhoptrus* spp., *Liogenys* spp, *Maecolaspis* spp, *Maladera*

castanea, Megascelis spp, Melighetes aeneus, Melolontha spp., Myochrous armatus, Orycaephilus spp., Otorhynchus spp., Phyllophaga spp, Phlyctinus spp., Popillia spp., Psylliodes spp., Rhyssomatus aubtilis, Rhizopertha spp., Scarabeidae, Sitophilus spp., Sitotroga spp., Somaticus spp, Sphenophorus spp, Sternechus subsignatus, Tenebrio spp., Tribolium spp. and Trogoderma spp.;

5 from the order *Diptera*, for example,

Aedes spp., Anopheles spp, Antherigona soccata, Bactrocea oleae, Bibio hortulanus, Bradysia spp, Calliphora erythrocephala, Ceratitis spp., Chrysomyia spp., Culex spp., Cuterebra spp., Dacus spp., Delia spp, Drosophila melanogaster, Fannia spp., Gastrophilus spp., Geomyza tripunctata, Glossina spp., Hypoderma spp., Hyppobosca spp., Liriomyza spp., Lucilia spp., Melanagromyza spp., Musca spp.,
10 Oestrus spp., Orseolia spp., Oscinella frit, Pegomyia hyoscyami, Phorbia spp., Rhagoletis spp, Rivelia quadrifasciata, Scatella spp, Sciara spp., Stomoxys spp., Tabanus spp., Tannia spp. and Tipula spp.;

from the order *Hemiptera*, for example,

Acanthocoris scabator, Acrosternum spp, Adelphocoris lineolatus, Amblypelta nitida, Bathycoelia thalassina, Blissus spp, Cimex spp., Clavigralla tomentosicollis, Creontiades spp, Distantiella theobroma,
15 Dichelops furcatus, Dysdercus spp., Edessa spp, Euchistus spp., Eurydema pulchrum, Eurygaster spp., Halyomorpha halys, Horcias nobilellus, Leptocoris spp., Lygus spp, Margarodes spp, Murgantia histrionic, Neomegalotomus spp, Nesidiocoris tenuis, Nezara spp., Nysius simulans, Oebalus insularis, Piesma spp., Piezodorus spp, Rhodnius spp., Sahlbergella singularis, Scaptocoris castanea, Scotinophara spp. , Thyanta spp , Triatoma spp., Vatica illudens;

20 Acyrthosium pisum, Adalges spp, Agalliana ensigera, Agonoscena targionii, Aleurodicus spp, Aleurocanthus spp, Aleurolobus barodensis, Aleurothrix floccosus, Aleyrodes brassicae, Amarasca biguttula, Amritodus atkinsoni, Aonidiella spp., Aphididae, Aphis spp., Aspidiotus spp., Aulacorthum solani, Bactericera cockerelli, Bemisia spp, Brachycaudus spp, Brevicoryne brassicae, Cacopsylla spp, Cavariella aegopodii Scop., Ceroplaster spp., Chrysomphalus aonidium, Chrysomphalus dictyospermi,
25 Cicadella spp, Cofana spectra, Cryptomyzus spp, Cicadulina spp, Coccus hesperidum, Dalbulus maidis, Dialeurodes spp, Diaphorina citri, Diuraphis noxia, Dysaphis spp, Empoasca spp., Eriosoma larigerum, Erythroneura spp., Gascardia spp., Glycaspis brimblecombei, Hyadaphis pseudobrassicae, Hyalopterus spp, Hyperomyzus pallidus, Idioscopus clypealis, Jacobiasca lybica, Laodelphax spp., Lecanium corni, Lepidosaphes spp., Lopaphis erysimi, Lyogenys maidis, Macrosiphum spp., Mahanarva spp, Metcalfa pruinosa, Metopolophium dirhodum, Myndus crudus, Myzus spp., Neotoxoptera sp, Nephotettix spp.,
30 Nilaparvata spp., Nippolachnus piri Mats, Odonaspis ruthae, Oregma lanigera Zehnter, Parabemisia myricae, Paratrioza cockerelli, Parlatoria spp., Pemphigus spp., Peregrinus maidis, Perkinsiella spp, Phorodon humuli, Phylloxera spp, Planococcus spp., Pseudaulacaspis spp., Pseudococcus spp., Pseudatomoscelis seriatus, Psylla spp., Pulvinaria aethiopica, Quadraspidiotus spp., Quesada gigas,
35 Recilia dorsalis, Rhopalosiphum spp., Saissetia spp., Scaphoideus spp., Schizaphis spp., Sitobion spp., Sogatella furcifera, Spissistilus festinus, Tarophagus Proserpina, Toxoptera spp, Trialeurodes spp, Tridiscus sporoboli, Trionymus spp, Trioza erytrae , Unaspis citri, Zyginia flammigera, Zyginidia scutellaris, ;

from the order *Hymenoptera*, for example,

40 Acromyrmex, Arge spp, Atta spp., Cephus spp., Diprion spp., Diprionidae, Gilpinia polytoma, Hoplocampa spp., Lasius spp., Monomorium pharaonis, Neodiprion spp., Pogonomymex spp, Slenopsis invicta, Solenopsis spp. and Vespa spp.;

from the order *Isoptera*, for example,

Coptotermes spp, Cornitermes cumulans, Incisitermes spp, Macrotermes spp, Mastotermes spp, Microtermes spp, Reticulitermes spp.; Solenopsis geminate

from the order *Lepidoptera*, for example,

- 5 Acleris spp., Adoxophyes spp., Aegeria spp., Agrotis spp., Alabama argillaceae, Amylois spp., Anticarsia gemmatilis, Archips spp., Argyresthia spp, Argyrotaenia spp., Autographa spp., Bucculatrix thurberiella, Busseola fusca, Cadra cautella, Carposina nipponensis, Chilo spp., Choristoneura spp., Chrysoteuchia topiaria, Clysia ambiguella, Cnaphalocrocis spp., Cnephasia spp., Cochylis spp., Coleophora spp., Colias lesbia, Cosmophila flava, Crambus spp, Crocidolomia binotalis, Cryptophlebia
- 10 leucotreta, Cydalima perspectalis, Cydia spp., Diaphania perspectalis, Diatraea spp., Diparopsis castanea, Earias spp., Eldana saccharina, Ephestia spp., Epinotia spp, Estigmene acrea, Etiella zinckinella, Eucosma spp., Eupoecilia ambiguella, Euproctis spp., Euxoa spp., Feltia jaculiferia, Grapholita spp., Hedya nubiferana, Heliothis spp., Hellula undalis, Herpetogramma spp, Hyphantria cunea, Keiferia lycopersicella, Lasmopalpus lignosellus, Leucoptera scitella, Lithocolletis spp., Lobesia botrana,
- 15 Loxostege bifidalis, Lymantria spp., Lyonetia spp., Malacosoma spp., Mamestra brassicae, Manduca sexta, Mythimna spp, Noctua spp, Operophtera spp., Orniodes indica, Ostrinia nubilalis, Pammene spp., Pandemis spp., Panolis flammea, Papaipema nebris, Pectinophora gossypiella, Perileucoptera coffeella, Pseudaletia unipuncta, Phthorimaea operculella, Pieris rapae, Pieris spp., Plutella xylostella, Prays spp., Pseudoplusia spp, Rachiplusia nu, Richia albicosta, Scirpophaga spp., Sesamia spp., Sparganothis spp.,
- 20 Spodoptera spp., Sylepta derogate, Synanthedon spp., Thaumetopoea spp., Tortrix spp., Trichoplusia ni, Tuta absoluta, and Yponomeuta spp.;

from the order *Mallophaga*, for example,

Damalinae spp. and Trichodectes spp.;

from the order *Orthoptera*, for example,

- 25 Blatta spp., Blattella spp., Gryllotalpa spp., Leucophaea maderae, Locusta spp., Neocurtilla hexadactyla, Periplaneta spp. , Scapteriscus spp, and Schistocerca spp.;

from the order *Psocoptera*, for example,

Liposcelis spp.;

from the order *Siphonaptera*, for example,

- 30 Ceratophyllus spp., Ctenocephalides spp. and Xenopsylla cheopis;

from the order *Thysanoptera*, for example,

Calliothrips phaseoli, Frankliniella spp., Heliothrips spp, Hercinothrips spp., Parthenothrips spp, Scirtothrips aurantii, Sericothrips variabilis, Taeniothrips spp., Thrips spp;

from the order *Thysanura*, for example, Lepisma saccharina.

- 35 The active ingredients according to the invention can be used for controlling, i. e. containing or destroying, pests of the abovementioned type which occur in particular on plants, especially on useful plants and ornamentals in agriculture, in horticulture and in forests, or on organs, such as fruits, flowers, foliage, stalks, tubers or roots, of such plants, and in some cases even plant organs which are formed at a later point in time remain protected against these pests.

- 40 Suitable target crops are, in particular, cereals, such as wheat, barley, rye, oats, rice, maize or sorghum; beet, such as sugar or fodder beet; fruit, for example pomaceous fruit, stone fruit or soft fruit, such as apples, pears, plums, peaches, almonds, cherries or berries, for example strawberries,

raspberries or blackberries; leguminous crops, such as beans, lentils, peas or soya; oil crops, such as oilseed rape, mustard, poppies, olives, sunflowers, coconut, castor, cocoa or ground nuts; cucurbits, such as pumpkins, cucumbers or melons; fibre plants, such as cotton, flax, hemp or jute; citrus fruit, such as oranges, lemons, grapefruit or tangerines; vegetables, such as spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes or bell peppers; Lauraceae, such as avocado, Cinnamomum or camphor; and also tobacco, nuts, coffee, eggplants, sugarcane, tea, pepper, grapevines, hops, the plantain family and latex plants.

The compositions and/or methods of the present invention may be also used on any ornamental and/or vegetable crops, including flowers, shrubs, broad-leaved trees and evergreens.

For example the invention may be used on any of the following ornamental species: *Ageratum* spp., *Alonsoa* spp., *Anemone* spp., *Anisodonteia capensis*, *Anthemis* spp., *Antirrhinum* spp., *Aster* spp., *Begonia* spp. (e.g. *B. elatior*, *B. semperflorens*, *B. tuberosa*), *Bougainvillea* spp., *Brachycome* spp., *Brassica* spp. (ornamental), *Calceolaria* spp., *Capsicum annuum*, *Catharanthus roseus*, *Canna* spp., *Centaurea* spp., *Chrysanthemum* spp., *Cineraria* spp. (*C. maritime*), *Coreopsis* spp., *Crassula coccinea*, *Cuphea ignea*, *Dahlia* spp., *Delphinium* spp., *Dicentra spectabilis*, *Dorotheanthus* spp., *Eustoma grandiflorum*, *Forsythia* spp., *Fuchsia* spp., *Geranium gnaphalium*, *Gerbera* spp., *Gomphrena globosa*, *Heliotropium* spp., *Helianthus* spp., *Hibiscus* spp., *Hortensia* spp., *Hydrangea* spp., *Hypoestes phyllostachya*, *Impatiens* spp. (*I. Walleriana*), *Iresines* spp., *Kalanchoe* spp., *Lantana camara*, *Lavatera trimestris*, *Leonotis leonurus*, *Lilium* spp., *Mesembryanthemum* spp., *Mimulus* spp., *Monarda* spp., *Nemesia* spp., *Tagetes* spp., *Dianthus* spp. (carnation), *Canna* spp., *Oxalis* spp., *Bellis* spp., *Pelargonium* spp. (*P. peltatum*, *P. Zonale*), *Viola* spp. (pansy), *Petunia* spp., *Phlox* spp., *Plecthranthus* spp., *Poinsettia* spp., *Parthenocissus* spp. (*P. quinquefolia*, *P. tricuspidata*), *Primula* spp., *Ranunculus* spp., *Rhododendron* spp., *Rosa* spp. (rose), *Rudbeckia* spp., *Saintpaulia* spp., *Salvia* spp., *Scaevola aemola*, *Schizanthus wisetonensis*, *Sedum* spp., *Solanum* spp., *Surfinia* spp., *Tagetes* spp., *Nicotinia* spp., *Verbena* spp., *Zinnia* spp. and other bedding plants.

For example the invention may be used on any of the following vegetable species: *Allium* spp. (*A. sativum*, *A. cepa*, *A. oschaninii*, *A. Porrum*, *A. ascalonicum*, *A. fistulosum*), *Anthriscus cerefolium*, *Apium graveolus*, *Asparagus officinalis*, *Beta vulgaris*, *Brassica* spp. (*B. Oleracea*, *B. Pekinensis*, *B. rapa*), *Capsicum annuum*, *Cicer arietinum*, *Cichorium endivia*, *Cichorium* spp. (*C. intybus*, *C. endivia*), *Citrullus lanatus*, *Cucumis* spp. (*C. sativus*, *C. melo*), *Cucurbita* spp. (*C. pepo*, *C. maxima*), *Cyanara* spp. (*C. scolymus*, *C. cardunculus*), *Daucus carota*, *Foeniculum vulgare*, *Hypericum* spp., *Lactuca sativa*, *Lycopersicon* spp. (*L. esculentum*, *L. lycopersicum*), *Mentha* spp., *Ocimum basilicum*, *Petroselinum crispum*, *Phaseolus* spp. (*P. vulgaris*, *P. coccineus*), *Pisum sativum*, *Raphanus sativus*, *Rheum raphaniticum*, *Rosemarinus* spp., *Salvia* spp., *Scorzonera hispanica*, *Solanum melongena*, *Spinacea oleracea*, *Valerianella* spp. (*V. locusta*, *V. eriocarpa*) and *Vicia faba*.

Preferred ornamental species include African violet, *Begonia*, *Dahlia*, *Gerbera*, *Hydrangea*, *Verbena*, *Rosa*, *Kalanchoe*, *Poinsettia*, *Aster*, *Centaurea*, *Coreopsis*, *Delphinium*, *Monarda*, *Phlox*, *Rudbeckia*, *Sedum*, *Petunia*, *Viola*, *Impatiens*, *Geranium*, *Chrysanthemum*, *Ranunculus*, *Fuchsia*, *Salvia*, *Hortensia*, rosemary, sage, St. Johnswort, mint, sweet pepper, tomato and cucumber.

The active ingredients according to the invention are especially suitable for controlling Aphis craccivora, Diabrotica balteata, Heliothis virescens, Myzus persicae, Plutella xylostella and Spodoptera littoralis in cotton, vegetable, maize, rice and soya crops. The active ingredients according to the

invention are further especially suitable for controlling *Mamestra* (preferably in vegetables), *Cydia pomonella* (preferably in apples), *Empoasca* (preferably in vegetables, vineyards), *Leptinotarsa* (preferably in potatoes) and *Chilo suppressalis* (preferably in rice).

The active ingredients according to the invention are especially suitable for controlling *Aphis craccivora*, *Diabrotica balteata*, *Heliothis virescens*, *Myzus persicae*, *Plutella xylostella* and *Spodoptera littoralis* in cotton, vegetable, maize, rice and soya crops. The active ingredients according to the invention are further especially suitable for controlling *Mamestra* (preferably in vegetables), *Cydia pomonella* (preferably in apples), *Empoasca* (preferably in vegetables, vineyards), *Leptinotarsa* (preferably in potatoes) and *Chilo suppressalis* (preferably in rice).

10 In a further aspect, the invention may also relate to a method of controlling damage to plant and parts thereof by plant parasitic nematodes (Endoparasitic-, Semiendoparasitic- and Ectoparasitic nematodes), especially plant parasitic nematodes such as root knot nematodes, *Meloidogyne hapla*, *Meloidogyne incognita*, *Meloidogyne javanica*, *Meloidogyne arenaria* and other *Meloidogyne* species; cyst-forming nematodes, *Globodera rostochiensis* and other *Globodera* species; *Heterodera avenae*,
15 *Heterodera glycines*, *Heterodera schachtii*, *Heterodera trifolii*, and other *Heterodera* species; Seed gall nematodes, *Anguina* species; Stem and foliar nematodes, *Aphelenchoides* species; Sting nematodes, *Belonolaimus longicaudatus* and other *Belonolaimus* species; Pine nematodes, *Bursaphelenchus xylophilus* and other *Bursaphelenchus* species; Ring nematodes, *Criconema* species, *Criconemella* species, *Criconemoides* species, *Mesocriconema* species; Stem and bulb nematodes, *Ditylenchus*
20 *destructor*, *Ditylenchus dipsaci* and other *Ditylenchus* species; Awl nematodes, *Dolichodorus* species; Spiral nematodes, *Helicotylenchus multicinctus* and other *Helicotylenchus* species; Sheath and sheathoid nematodes, *Hemicycliophora* species and *Hemicriconemoides* species; *Hirshmanniella* species; Lance nematodes, *Hoploaimus* species; false rootknot nematodes, *Nacobbus* species; Needle nematodes, *Longidorus elongatus* and other *Longidorus* species; Pin nematodes, *Pratylenchus* species;
25 Lesion nematodes, *Pratylenchus neglectus*, *Pratylenchus penetrans*, *Pratylenchus curvatus*, *Pratylenchus goodeyi* and other *Pratylenchus* species; Burrowing nematodes, *Radopholus similis* and other *Radopholus* species; Reniform nematodes, *Rotylenchus robustus*, *Rotylenchus reniformis* and other *Rotylenchus* species; *Scutellonema* species; Stubby root nematodes, *Trichodorus primitivus* and other *Trichodorus* species, *Paratrichodorus* species; Stunt nematodes, *Tylenchorhynchus claytoni*,
30 *Tylenchorhynchus dubius* and other *Tylenchorhynchus* species; Citrus nematodes, *Tylenchulus* species; Dagger nematodes, *Xiphinema* species; and other plant parasitic nematode species, such as *Subanguina* spp., *Hypsoperine* spp., *Macroposthonia* spp., *Melinius* spp., *Punctodera* spp., and *Quinisulcius* spp..

The compounds of the invention may also have activity against the molluscs. Examples of which include, for example, *Ampullariidae*; *Arion* (*A. ater*, *A. circumscriptus*, *A. hortensis*, *A. rufus*);

35 *Bradybaenidae* (*Bradybaena fruticum*); *Cepaea* (*C. hortensis*, *C. Nemoralis*); *ochlodina*; *Deroceras* (*D. agrestis*, *D. empiricorum*, *D. laeve*, *D. reticulatum*); *Discus* (*D. rotundatus*); *Euomphalia*; *Galba* (*G. trunculata*); *Helicella* (*H. itala*, *H. obvia*); *Helicidae* *Helicigona arbustorum*); *Helicodiscus*; *Helix* (*H. aperta*); *Limax* (*L. cinereoniger*, *L. flavus*, *L. marginatus*, *L. maximus*, *L. tenellus*); *Lymnaea*; *Milax* (*M. gagates*, *M. marginatus*, *M. sowerbyi*); *Opeas*; *Pomacea* (*P. canaticulata*); *Vallonia* and *Zanitoides*.

40 Crops are to be understood as also including those crops which have been rendered tolerant to herbicides like bromoxynil or classes of herbicides such as ALS-, EPSPS-, GS-, HPPD- and PPO-inhibitors. An example of a crop that has been rendered tolerant to imidazolinones, e.g. imazamox, by

conventional methods of breeding is Clearfield® summer canola. Examples of crops that have been rendered tolerant to herbicides by genetic engineering methods include e.g. glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady®, Herculex I® and LibertyLink®.

5 Crops are also to be understood as being those which naturally are or have been rendered resistant to harmful insects. This includes plants transformed by the use of recombinant DNA techniques, for example, to be capable of synthesising one or more selectively acting toxins, such as are known, for example, from toxin-producing bacteria. Examples of toxins which can be expressed include δ -endotoxins, vegetative insecticidal proteins (Vip), insecticidal proteins of bacteria colonising nematodes, 10 and toxins produced by scorpions, arachnids, wasps and fungi.

An example of a crop that has been modified to express the *Bacillus thuringiensis* toxin is the Bt maize KnockOut® (Syngenta Seeds). An example of a crop comprising more than one gene that codes for insecticidal resistance and thus expresses more than one toxin is VipCot® (Syngenta Seeds). Crops or seed material thereof can also be resistant to multiple types of pests (so-called stacked transgenic 15 events when created by genetic modification). For example, a plant can have the ability to express an insecticidal protein while at the same time being herbicide tolerant, for example Herculex I® (Dow AgroSciences, Pioneer Hi-Bred International).

Further areas of use of the compositions according to the invention are the protection of stored goods and store rooms and the protection of raw materials, such as wood, textiles, floor coverings or 20 buildings, and also in the hygiene sector, especially the protection of humans, domestic animals and productive livestock against pests of the mentioned type.

The present invention also provides a method for controlling pests (such as mosquitoes and other disease vectors; see also http://www.who.int/malaria/vector_control/irs/en/). In one embodiment, the method for controlling pests comprises applying the compositions of the invention to the target pests, to 25 their locus or to a surface or substrate by brushing, rolling, spraying, spreading or dipping. By way of example, an IRS (indoor residual spraying) application of a surface such as a wall, ceiling or floor surface is contemplated by the method of the invention. In another embodiment, it is contemplated to apply such compositions to a substrate such as non-woven or a fabric material in the form of (or which can be used in the manufacture of) netting, clothing, bedding, curtains and tents.

30

In one embodiment, the method for controlling such pests comprises applying a pesticidally effective amount of the compositions of the invention to the target pests, to their locus, or to a surface or substrate so as to provide effective residual pesticidal activity on the surface or substrate. Such application may be made by brushing, rolling, spraying, spreading or dipping the pesticidal composition of 35 the invention. By way of example, an IRS application of a surface such as a wall, ceiling or floor surface is contemplated by the method of the invention so as to provide effective residual pesticidal activity on the surface. In another embodiment, it is contemplated to apply such compositions for residual control of pests on a substrate such as a fabric material in the form of (or which can be used in the manufacture of) netting, clothing, bedding, curtains and tents.

40 Substrates including non-woven, fabrics or netting to be treated may be made of natural fibres such as cotton, raffia, jute, flax, sisal, hessian, or wool, or synthetic fibres such as polyamide, polyester,

polypropylene, polyacrylonitrile or the like. The polyesters are particularly suitable. The methods of textile treatment are known, e.g. WO2008151984, WO2003034823, US5631072, WO2005064072, WO2006128870, EP1724392, WO2005113886 or WO2007090739.

Further areas of use of the compositions according to the invention are the field of tree

5 injection/trunk treatment for all ornamental trees as well all sort of fruit and nut trees.

In the field of tree injection/trunk treatment, the compounds according to the present invention are especially suitable against wood-boring insects from the order *Lepidoptera* as mentioned above and from the order *Coleoptera*, especially against woodborers listed in the following tables X and Y:

Table X. Examples of exotic woodborers of economic importance.

Family	Species	Host or Crop Infested
Buprestidae	<i>Agrilus planipennis</i>	Ash
Cerambycidae	<i>Anoplura glabripennis</i>	Hardwoods
Scolytidae	<i>Xylosandrus crassiusculus</i>	Hardwoods
	<i>X. mutilatus</i>	Hardwoods
	<i>Tomicus piniperda</i>	Conifers

10

Table Y. Examples of native woodborers of economic importance.

Family	Species	Host or Crop Infested
Buprestidae	<i>Agrilus anxius</i>	Birch
	<i>Agrilus politus</i>	Willow, Maple
	<i>Agrilus sayi</i>	Bayberry, Sweetfern
	<i>Agrilus vittaticollis</i>	Apple, Pear, Cranberry, Serviceberry, Hawthorn
	<i>Chrysobothris femorata</i>	Apple, Apricot, Beech, Boxelder, Cherry, Chestnut, Currant, Elm, Hawthorn, Hackberry, Hickory, Horsechestnut, Linden, Maple, Mountain-ash, Oak, Pecan, Pear, Peach, Persimmon, Plum, Poplar, Quince, Redbud, Serviceberry, Sycamore, Walnut, Willow
	<i>Texania campestris</i>	Basswood, Beech, Maple, Oak, Sycamore, Willow, Yellow-poplar
Cerambycidae	<i>Goes pulverulentus</i>	Beech, Elm, Nuttall, Willow, Black oak, Cherrybark oak, Water oak, Sycamore
	<i>Goes tigrinus</i>	Oak
	<i>Neoclytus acuminatus</i>	Ash, Hickory, Oak, Walnut, Birch, Beech, Maple, Eastern hophornbeam, Dogwood,

Family	Species	Host or Crop Infested
		Persimmon, Redbud, Holly, Hackberry, Black locust, Honeylocust, Yellow-poplar, Chestnut, Osage-orange, Sassafras, Lilac, Mountain-mahogany, Pear, Cherry, Plum, Peach, Apple, Elm, Basswood, Sweetgum
	<i>Neoptychodes trilineatus</i>	Fig, Alder, Mulberry, Willow, Netleaf hackberry
	<i>Oberea ocellata</i>	Sumac, Apple, Peach, Plum, Pear, Currant, Blackberry
	<i>Oberea tripunctata</i>	Dogwood, Viburnum, Elm, Sourwood, Blueberry, Rhododendron, Azalea, Laurel, Poplar, Willow, Mulberry
	<i>Oncideres cingulata</i>	Hickory, Pecan, Persimmon, Elm, Sourwood, Basswood, Honeylocust, Dogwood, Eucalyptus, Oak, Hackberry, Maple, Fruit trees
	<i>Saperda calcarata</i>	Poplar
	<i>Strophiona nitens</i>	Chestnut, Oak, Hickory, Walnut, Beech, Maple
Scolytidae	<i>Corthylus columbianus</i>	Maple, Oak, Yellow-poplar, Beech, Boxelder, Sycamore, Birch, Basswood, Chestnut, Elm
	<i>Dendroctonus frontalis</i>	Pine
	<i>Dryocoetes betulae</i>	Birch, Sweetgum, Wild cherry, Beech, Pear
	<i>Monarthrum fasciatum</i>	Oak, Maple, Birch, Chestnut, Sweetgum, Blackgum, Poplar, Hickory, Mimosa, Apple, Peach, Pine
	<i>Phloeotribus liminaris</i>	Peach, Cherry, Plum, Black cherry, Elm, Mulberry, Mountain-ash
	<i>Pseudopityophthorus pruinosis</i>	Oak, American beech, Black cherry, Chickasaw plum, Chestnut, Maple, Hickory, Hornbeam, Hophornbeam
Sesiidae	<i>Paranthrene simulans</i>	Oak, American chestnut
	<i>Sannina uroceriformis</i>	Persimmon
	<i>Synanthedon exitiosa</i>	Peach, Plum, Nectarine, Cherry,

Family	Species	Host or Crop Infested
		Apricot, Almond, Black cherry
	<i>Synanthedon pictipes</i>	Peach, Plum, Cherry, Beach, Black Cherry
	<i>Synanthedon rubrofascia</i>	Tupelo
	<i>Synanthedon scitula</i>	Dogwood, Pecan, Hickory, Oak, Chestnut, Beech, Birch, Black cherry, Elm, Mountain-ash, Viburnum, Willow, Apple, Loquat, Ninebark, Bayberry
	<i>Vitacea polistiformis</i>	Grape

The present invention may be also used to control any insect pests that may be present in turfgrass, including for example beetles, caterpillars, fire ants, ground pearls, millipedes, sow bugs, mites, mole crickets, scales, mealybugs ticks, spittlebugs, southern chinch bugs and white grubs. The present invention may be used to control insect pests at various stages of their life cycle, including eggs, larvae, nymphs and adults.

In particular, the present invention may be used to control insect pests that feed on the roots of turfgrass including white grubs (such as *Cyclocephala* spp. (e.g. masked chafer, *C. lurida*), *Rhizotrogus* spp. (e.g. European chafer, *R. majalis*), *Cotinus* spp. (e.g. Green June beetle, *C. nitida*), *Popillia* spp. (e.g. Japanese beetle, *P. japonica*), *Phyllophaga* spp. (e.g. May/June beetle), *Ataenius* spp. (e.g. Black turfgrass ataenius, *A. spretulus*), *Maladera* spp. (e.g. Asiatic garden beetle, *M. castanea*) and *Tomarus* spp.), ground pearls (*Margarodes* spp.), mole crickets (tawny, southern, and short-winged; *Scapteriscus* spp., *Gryllotalpa africana*) and leatherjackets (European crane fly, *Tipula* spp.).

The present invention may also be used to control insect pests of turfgrass that are thatch dwelling, including armyworms (such as fall armyworm *Spodoptera frugiperda*, and common armyworm *Pseudaletia unipuncta*), cutworms, billbugs (*Sphenophorus* spp., such as *S. venatus verstitus* and *S. parvulus*), and sod webworms (such as *Crambus* spp. and the tropical sod webworm, *Herpetogramma phaeopteralis*).

The present invention may also be used to control insect pests of turfgrass that live above the ground and feed on the turfgrass leaves, including chinch bugs (such as southern chinch bugs, *Blissus insularis*), Bermudagrass mite (*Eriophyes cynodoniensis*), rhodesgrass mealybug (*Antonina graminis*), two-lined spittlebug (*Prospapia bicincta*), leafhoppers, cutworms (*Noctuidae* family), and greenbugs.

The present invention may also be used to control other pests of turfgrass such as red imported fire ants (*Solenopsis invicta*) that create ant mounds in turf.

In the hygiene sector, the compositions according to the invention are active against ectoparasites such as hard ticks, soft ticks, mange mites, harvest mites, flies (biting and licking), parasitic fly larvae, lice, hair lice, bird lice and fleas.

Examples of such parasites are:

Of the order Anoplurida: Haematopinus spp., Linognathus spp., Pediculus spp. and Phtirus spp., Solenopotes spp..

Of the order Mallophagida: Trimenopon spp., Menopon spp., Trinoton spp., Bovicola spp., Werneckiella spp., Lepikentron spp., Damalina spp., Trichodectes spp. and Felicola spp..

Of the order Diptera and the suborders Nematocerina and Brachycerina, for example Aedes spp., Anopheles spp., Culex spp., Simulium spp., Eusimulium spp., Phlebotomus spp., Lutzomyia spp.,
 5 Culicoides spp., Chrysops spp., Hybomitra spp., Atylotus spp., Tabanus spp., Haematopota spp., Philipomyia spp., Braula spp., Musca spp., Hydrotaea spp., Stomoxys spp., Haematobia spp., Morellia spp., Fannia spp., Glossina spp., Calliphora spp., Lucilia spp., Chrysomyia spp., Wohlfahrtia spp., Sarcophaga spp., Oestrus spp., Hypoderma spp., Gasterophilus spp., Hippobosca spp., Lipoptena spp. and Melophagus spp..

10 Of the order Siphonaptera, for example Pulex spp., Ctenocephalides spp., Xenopsylla spp., Ceratophyllus spp..

Of the order Heteroptera, for example Cimex spp., Triatoma spp., Rhodnius spp., Panstrongylus spp..

15 Of the order Blattellidae, for example Blatta orientalis, Periplaneta americana, Blattella germanica and Supella spp..

Of the subclass Acarina (Acarida) and the orders Meta- and Mesostigmata, for example Argas spp., Ornithodoros spp., Otobius spp., Ixodes spp., Amblyomma spp., Boophilus spp., Dermacentor spp., Haemaphysalis spp., Hyalomma spp., Rhipicephalus spp., Dermanyssus spp., Raillietia spp., Pneumonyssus spp., Sternostoma spp. and Varroa spp..

20 Of the orders Actinoptera (Prostigmata) and Acaridida (Astigmata), for example Acarapis spp., Cheyletiella spp., Ornithocheyletia spp., Myobia spp., Psorergates spp., Demodex spp., Trombicula spp., Listrophorus spp., Acarus spp., Tyrophagus spp., Caloglyphus spp., Hypodectes spp., Pterolichus spp., Psoroptes spp., Chorioptes spp., Otodectes spp., Sarcoptes spp., Notoedres spp., Knemidocoptes spp., Cytodites spp. and Laminosioptes spp..

25 The compositions according to the invention are also suitable for protecting against insect infestation in the case of materials such as wood, textiles, plastics, adhesives, glues, paints, paper and card, leather, floor coverings and buildings.

The compositions according to the invention can be used, for example, against the following pests: beetles such as Hylotrupes bajulus, Chlorophorus pilosus, Anobium punctatum, Xestobium rufovillosum,
 30 Ptilinuspecticornis, Dendrobium pertinax, Ernobius mollis, Priobium carpini, Lyctus brunneus, Lyctus africanus, Lyctus planicollis, Lyctus linearis, Lyctus pubescens, Trogoxylon aequale, Minthesrugicollis, Xyleborus spec., Trypodendron spec., Apate monachus, Bostrychus capucinus, Heterobostrychus brunneus, Sinoxylon spec. and Dinoderus minutus, and also hymenoptera such as Sirex juvencus, Urocerus gigas, Urocerus gigas taignus and Urocerus augur, and termites such as Kaloterms flavicollis,
 35 Cryptotermes brevis, Heterotermes indicola, Reticulitermes flavipes, Reticulitermes santonensis, Reticulitermes lucifugus, Mastotermes darwiniensis, Zootermopsis nevadensis and Coptotermes formosanus, and bristletails such as Lepisma saccharina.

The compounds according to the invention can be used as pesticidal agents in unmodified form, but they are generally formulated into compositions in various ways using formulation adjuvants, such as
 40 carriers, solvents and surface-active substances. The formulations can be in various physical forms, e.g. in the form of dusting powders, gels, wettable powders, water-dispersible granules, water-dispersible tablets, effervescent pellets, emulsifiable concentrates, microemulsifiable concentrates, oil-in-water

emulsions, oil-flowables, aqueous dispersions, oily dispersions, suspo-emulsions, capsule suspensions, emulsifiable granules, soluble liquids, water-soluble concentrates (with water or a water-miscible organic solvent as carrier), impregnated polymer films or in other forms known e.g. from the Manual on

- 5 Second Revision (2010). Such formulations can either be used directly or diluted prior to use. The dilutions can be made, for example, with water, liquid fertilisers, micronutrients, biological organisms, oil or solvents.

The formulations can be prepared e.g. by mixing the active ingredient with the formulation adjuvants in order to obtain compositions in the form of finely divided solids, granules, solutions,

10 dispersions or emulsions. The active ingredients can also be formulated with other adjuvants, such as finely divided solids, mineral oils, oils of vegetable or animal origin, modified oils of vegetable or animal origin, organic solvents, water, surface-active substances or combinations thereof.

The active ingredients can also be contained in very fine microcapsules. Microcapsules contain the active ingredients in a porous carrier. This enables the active ingredients to be released into the

15 environment in controlled amounts (e.g. slow-release). Microcapsules usually have a diameter of from 0.1 to 500 microns. They contain active ingredients in an amount of about from 25 to 95 % by weight of the capsule weight. The active ingredients can be in the form of a monolithic solid, in the form of fine particles in solid or liquid dispersion or in the form of a suitable solution. The encapsulating membranes can comprise, for example, natural or synthetic rubbers, cellulose, styrene/butadiene copolymers,

20 polyacrylonitrile, polyacrylate, polyesters, polyamides, polyureas, polyurethane or chemically modified polymers and starch xanthates or other polymers that are known to the person skilled in the art. Alternatively, very fine microcapsules can be formed in which the active ingredient is contained in the form of finely divided particles in a solid matrix of base substance, but the microcapsules are not themselves encapsulated.

25 The formulation adjuvants that are suitable for the preparation of the compositions according to the invention are known *per se*. As liquid carriers there may be used: water, toluene, xylene, petroleum ether, vegetable oils, acetone, methyl ethyl ketone, cyclohexanone, acid anhydrides, acetonitrile, acetophenone, amyl acetate, 2-butanone, butylene carbonate, chlorobenzene, cyclohexane, cyclohexanol, alkyl esters of acetic acid, diacetone alcohol, 1,2-dichloropropane, diethanolamine, p-diethylbenzene, diethylene glycol,

30 diethylene glycol abietate, diethylene glycol butyl ether, diethylene glycol ethyl ether, diethylene glycol methyl ether, *N,N*-dimethylformamide, dimethyl sulfoxide, 1,4-dioxane, dipropylene glycol, dipropylene glycol methyl ether, dipropylene glycol dibenzoate, diproxitol, alkylpyrrolidone, ethyl acetate, 2-ethylhexanol, ethylene carbonate, 1,1,1-trichloroethane, 2-heptanone, alpha-pinene, d-limonene, ethyl lactate, ethylene glycol, ethylene glycol butyl ether, ethylene glycol methyl ether, gamma-butyrolactone, glycerol,

35 glycerol acetate, glycerol diacetate, glycerol triacetate, hexadecane, hexylene glycol, isoamyl acetate, isobornyl acetate, isooctane, isophorone, isopropylbenzene, isopropyl myristate, lactic acid, laurylamine, mesityl oxide, methoxypropanol, methyl isoamyl ketone, methyl isobutyl ketone, methyl laurate, methyl octanoate, methyl oleate, methylene chloride, m-xylene, *n*-hexane, *n*-octylamine, octadecanoic acid, octylamine acetate, oleic acid, oleylamine, o-xylene, phenol, polyethylene glycol, propionic acid, propyl

40 lactate, propylene carbonate, propylene glycol, propylene glycol methyl ether, p-xylene, toluene, triethyl phosphate, triethylene glycol, xylenesulfonic acid, paraffin, mineral oil, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol methyl ether, diethylene

glycol methyl ether, methanol, ethanol, isopropanol, and alcohols of higher molecular weight, such as amyl alcohol, tetrahydrofurfuryl alcohol, hexanol, octanol, ethylene glycol, propylene glycol, glycerol, *N*-methyl-2-pyrrolidone and the like.

Suitable solid carriers are, for example, talc, titanium dioxide, pyrophyllite clay, silica, attapulgite
5 clay, kieselguhr, limestone, calcium carbonate, bentonite, calcium montmorillonite, cottonseed husks, wheat flour, soybean flour, pumice, wood flour, ground walnut shells, lignin and similar substances.

A large number of surface-active substances can advantageously be used in both solid and liquid formulations, especially in those formulations which can be diluted with a carrier prior to use. Surface-active substances may be anionic, cationic, non-ionic or polymeric and they can be used as emulsifiers,
10 wetting agents or suspending agents or for other purposes. Typical surface-active substances include, for example, salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; salts of alkylarylsulfonates, such as calcium dodecylbenzenesulfonate; alkylphenol/alkylene oxide addition products, such as nonylphenol ethoxylate; alcohol/alkylene oxide addition products, such as tridecylalcohol ethoxylate; soaps, such as sodium stearate; salts of alkylnaphthalenesulfonates, such as sodium
15 dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl)sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryltrimethylammonium chloride, polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono- and di-alkylphosphate esters; and also further substances described e.g. in McCutcheon's Detergents and
20 Emulsifiers Annual, MC Publishing Corp., Ridgewood New Jersey (1981).

Further adjuvants that can be used in pesticidal formulations include crystallisation inhibitors, viscosity modifiers, suspending agents, dyes, anti-oxidants, foaming agents, light absorbers, mixing auxiliaries, antifoams, complexing agents, neutralising or pH-modifying substances and buffers, corrosion inhibitors, fragrances, wetting agents, take-up enhancers, micronutrients, plasticisers, glidants, lubricants,
25 dispersants, thickeners, antifreezes, microbicides, and liquid and solid fertilisers.

The compositions according to the invention can include an additive comprising an oil of vegetable or animal origin, a mineral oil, alkyl esters of such oils or mixtures of such oils and oil derivatives. The amount of oil additive in the composition according to the invention is generally from 0.01 to 10 %, based on the mixture to be applied. For example, the oil additive can be added to a spray tank in the desired
30 concentration after a spray mixture has been prepared. Preferred oil additives comprise mineral oils or an oil of vegetable origin, for example rapeseed oil, olive oil or sunflower oil, emulsified vegetable oil, alkyl esters of oils of vegetable origin, for example the methyl derivatives, or an oil of animal origin, such as fish oil or beef tallow. Preferred oil additives comprise alkyl esters of C₈-C₂₂ fatty acids, especially the methyl derivatives of C₁₂-C₁₈ fatty acids, for example the methyl esters of lauric acid, palmitic acid and
35 oleic acid (methyl laurate, methyl palmitate and methyl oleate, respectively). Many oil derivatives are known from the Compendium of Herbicide Adjuvants, 10th Edition, Southern Illinois University, 2010.

The inventive compositions generally comprise from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, of compounds of the present invention and from 1 to 99.9 % by weight of a formulation adjuvant which preferably includes from 0 to 25 % by weight of a surface-active substance. Whereas
40 commercial products may preferably be formulated as concentrates, the end user will normally employ dilute formulations.

The rates of application vary within wide limits and depend on the nature of the soil, the method of application, the crop plant, the pest to be controlled, the prevailing climatic conditions, and other factors governed by the method of application, the time of application and the target crop. As a general guideline compounds may be applied at a rate of from 1 to 2000 l/ha, especially from 10 to 1000 l/ha.

5 Preferred formulations can have the following compositions (weight %):

Emulsifiable concentrates:

active ingredient:	1 to 95 %, preferably 60 to 90 %
surface-active agent:	1 to 30 %, preferably 5 to 20 %
liquid carrier:	1 to 80 %, preferably 1 to 35 %

10

Dusts:

active ingredient:	0.1 to 10 %, preferably 0.1 to 5 %
solid carrier:	99.9 to 90 %, preferably 99.9 to 99 %

15 Suspension concentrates:

active ingredient:	5 to 75 %, preferably 10 to 50 %
water:	94 to 24 %, preferably 88 to 30 %
surface-active agent:	1 to 40 %, preferably 2 to 30 %

20 Wettable powders:

active ingredient:	0.5 to 90 %, preferably 1 to 80 %
surface-active agent:	0.5 to 20 %, preferably 1 to 15 %
solid carrier:	5 to 95 %, preferably 15 to 90 %

25 Granules:

active ingredient:	0.1 to 30 %, preferably 0.1 to 15 %
solid carrier:	99.5 to 70 %, preferably 97 to 85 %

The following Examples further illustrate, but do not limit, the invention.

30

<u>Wettable powders</u>	a)	b)	c)
active ingredients	25 %	50 %	75 %
sodium lignosulfonate	5 %	5 %	-
sodium lauryl sulfate	3 %	-	5 %
sodium diisobutyl naphthalenesulfonate	-	6 %	10 %
phenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	-
highly dispersed silicic acid	5 %	10 %	10 %
Kaolin	62 %	27 %	-

The combination is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording wettable powders that can be diluted with water to give suspensions of the desired concentration.

<u>Powders for dry seed treatment</u>	a)	b)	c)
active ingredients	25 %	50 %	75 %
light mineral oil	5 %	5 %	5 %
highly dispersed silicic acid	5 %	5 %	-
Kaolin	65 %	40 %	-
Talcum	-	-	20 %

The combination is thoroughly mixed with the adjuvants and the mixture is thoroughly ground in a suitable mill, affording powders that can be used directly for seed treatment.

<u>Emulsifiable concentrate</u>	
active ingredients	10 %
octylphenol polyethylene glycol ether (4-5 mol of ethylene oxide)	3 %
calcium dodecylbenzenesulfonate	3 %
castor oil polyglycol ether (35 mol of ethylene oxide)	4 %
Cyclohexanone	30 %
xylene mixture	50 %

Emulsions of any required dilution, which can be used in plant protection, can be obtained from this concentrate by dilution with water.

<u>Dusts</u>	a)	b)	c)
Active ingredients	5 %	6 %	4 %
Talcum	95 %	-	-
Kaolin	-	94 %	-
mineral filler	-	-	96 %

Ready-for-use dusts are obtained by mixing the combination with the carrier and grinding the mixture in a suitable mill. Such powders can also be used for dry dressings for seed.

<u>Extruder granules</u>	
Active ingredients	15 %
sodium lignosulfonate	2 %
carboxymethylcellulose	1 %
Kaolin	82 %

The combination is mixed and ground with the adjuvants, and the mixture is moistened with water. The mixture is extruded and then dried in a stream of air.

<u>Coated granules</u>	
Active ingredients	8 %
polyethylene glycol (mol. wt. 200)	3 %
Kaolin	89 %

The finely ground combination is uniformly applied, in a mixer, to the kaolin moistened with polyethylene glycol. Non-dusty coated granules are obtained in this manner.

Suspension concentrate

active ingredients	40 %
propylene glycol	10 %
nonylphenol polyethylene glycol ether (15 mol of ethylene oxide)	6 %
Sodium lignosulfonate	10 %
carboxymethylcellulose	1 %
silicone oil (in the form of a 75 % emulsion in water)	1 %
Water	32 %

The finely ground combination is intimately mixed with the adjuvants, giving a suspension concentrate

- 5 from which suspensions of any desired dilution can be obtained by dilution with water. Using such dilutions, living plants as well as plant propagation material can be treated and protected against infestation by microorganisms, by spraying, pouring or immersion.

Flowable concentrate for seed treatment

active ingredients	40 %
propylene glycol	5 %
copolymer butanol PO/EO	2 %
Tristyrenephenole with 10-20 moles EO	2 %
1,2-benzisothiazolin-3-one (in the form of a 20% solution in water)	0.5 %
monoazo-pigment calcium salt	5 %
Silicone oil (in the form of a 75 % emulsion in water)	0.2 %
Water	45.3 %

The finely ground combination is intimately mixed with the adjuvants, giving a suspension

- 10 concentrate from which suspensions of any desired dilution can be obtained by dilution with water. Using such dilutions, living plants as well as plant propagation material can be treated and protected against infestation by microorganisms, by spraying, pouring or immersion.

Slow Release Capsule Suspension

- 28 parts of the combination are mixed with 2 parts of an aromatic solvent and 7 parts of toluene
 15 diisocyanate/polymethylene-polyphenylisocyanate-mixture (8:1). This mixture is emulsified in a mixture of 1.2 parts of polyvinylalcohol, 0.05 parts of a defoamer and 51.6 parts of water until the desired particle size is achieved. To this emulsion a mixture of 2.8 parts 1,6-diaminohexane in 5.3 parts of water is added. The mixture is agitated until the polymerization reaction is completed. The obtained capsule suspension is stabilized by adding 0.25 parts of a thickener and 3 parts of a dispersing agent. The capsule
 20 suspension formulation contains 28% of the active ingredients. The medium capsule diameter is 8-15 microns. The resulting formulation is applied to seeds as an aqueous suspension in an apparatus suitable for that purpose.

- Formulation types include an emulsion concentrate (EC), a suspension concentrate (SC), a suspo-emulsion (SE), a capsule suspension (CS), a water dispersible granule (WG), an emulsifiable granule
 25 (EG), an emulsion, water in oil (EO), an emulsion, oil in water (EW), a micro-emulsion (ME), an oil dispersion (OD), an oil miscible flowable (OF), an oil miscible liquid (OL), a soluble concentrate (SL), an

ultra-low volume suspension (SU), an ultra-low volume liquid (UL), a technical concentrate (TK), a dispersible concentrate (DC), a wettable powder (WP), a soluble granule (SG) or any technically feasible formulation in combination with agriculturally acceptable adjuvants.

The activity of the compositions according to the invention can be broadened considerably, and adapted to prevailing circumstances, by adding other insecticidally, acaricidally and/or fungicidally active ingredients. The mixtures of the compounds of formula (I) with other insecticidally, acaricidally and/or fungicidally active ingredients may also have further surprising advantages which can also be described, in a wider sense, as synergistic activity. For example, better tolerance by plants, reduced phytotoxicity, insects can be controlled in their different development stages or better behaviour during their production, for example during grinding or mixing, during their storage or during their use.

Suitable additions to active ingredients here are, for example, representatives of the following classes of active ingredients: organophosphorus compounds, nitrophenol derivatives, thioureas, juvenile hormones, formamidines, benzophenone derivatives, ureas, pyrrole derivatives, carbamates, pyrethroids, chlorinated hydrocarbons, acylureas, pyridylmethyleamino derivatives, macrolides, neonicotinoids and *Bacillus thuringiensis* preparations.

The following mixtures of the compounds of formula (I) with active ingredients are preferred (the abbreviation "TX" means "one compound selected from the group consisting of the compounds described in Tables 1 to 96 and Table P of the present invention"):

an adjuvant selected from the group of substances consisting of petroleum oils (alternative name)
 (628) + TX,
 an acaricide selected from the group of substances consisting of 1,1-bis(4-chlorophenyl)-2-ethoxyethanol (IUPAC name) (910) + TX, 2,4-dichlorophenyl benzenesulfonate (IUPAC/Chemical Abstracts name) (1059) + TX, 2-fluoro-*N*-methyl-*N*-1-naphthylacetamide (IUPAC name) (1295) + TX, 4-chlorophenyl phenyl sulfone (IUPAC name) (981) + TX, abamectin (1) + TX, acequinocyl (3) + TX, acetoprole [CCN] + TX, acrinathrin (9) + TX, aldicarb (16) + TX, aldoxycarb (863) + TX, alpha-cypermethrin (202) + TX, amidithion (870) + TX, amidoflumet [CCN] + TX, amidothioate (872) + TX, amiton (875) + TX, amiton hydrogen oxalate (875) + TX, amitraz (24) + TX, aramite (881) + TX, arsenous oxide (882) + TX, AVI 382 (compound code) + TX, AZ 60541 (compound code) + TX, azinphos-ethyl (44) + TX, azinphos-methyl (45) + TX, azobenzene (IUPAC name) (888) + TX, azocyclotin (46) + TX, azotohate (889) + TX, benomyl (62) + TX, benoxafos (alternative name) [CCN] + TX, benzoximate (71) + TX, benzyl benzoate (IUPAC name) [CCN] + TX, bifenazate (74) + TX, bifenthrin (76) + TX, binapacryl (907) + TX, brofenvalerate (alternative name) + TX, bromocyclen (918) + TX, bromophos (920) + TX, bromophos-ethyl (921) + TX, bromopropylate (94) + TX, buprofezin (99) + TX, butocarboxim (103) + TX, butoxycarboxim (104) + TX, butylpyridaben (alternative name) + TX, calcium polysulfide (IUPAC name) (111) + TX, camphechlor (941) + TX, carbanolate (943) + TX, carbaryl (115) + TX, carbofuran (118) + TX, carbophenothion (947) + TX, CGA 50'439 (development code) (125) + TX, chinomethionat (126) + TX, chlordimeform (964) + TX, chlordimeform hydrochloride (964) + TX, chlorfenapyr (130) + TX, chlorfenethol (968) + TX, chlorfenson (970) + TX, chlorfensulfide (971) + TX, chlorfenvinphos (131) + TX, chlorobenzilate (975) + TX, chloromebuform (977) + TX, chloromethiuron (978) + TX, chloropropylate (983) + TX, chlorpyrifos (145) + TX, chlorpyrifos-methyl (146) + TX, chlorthiophos (994) + TX, cinerin I (696) + TX, cinerin II (696) + TX, cinerins (696) + TX, clofentezine (158) + TX,

closantel (alternative name) [CCN] + TX, coumaphos (174) + TX, crotamiton (alternative name) [CCN] + TX, crotoxyphos (1010) + TX, cufraneb (1013) + TX, cyanthoate (1020) + TX, cyflumetofen (CAS Reg. No.: 400882-07-7) + TX, cyhalothrin (196) + TX, cyhexatin (199) + TX, cypermethrin (201) + TX, DCPM (1032) + TX, DDT (219) + TX, demephion (1037) + TX, demephion-O (1037) + TX,
 5 demephion-S (1037) + TX, demeton (1038) + TX, demeton-methyl (224) + TX, demeton-O (1038) + TX, demeton-O-methyl (224) + TX, demeton-S (1038) + TX, demeton-S-methyl (224) + TX, demeton-S-methylsulfon (1039) + TX, diafenthiuron (226) + TX, dialifos (1042) + TX, diazinon (227) + TX, dichlofluanid (230) + TX, dichlorvos (236) + TX, dicliphos (alternative name) + TX, dicofol (242) + TX, dicrotophos (243) + TX, dienochlor (1071) + TX, dimefox (1081) + TX, dimethoate (262) + TX,
 10 dinactin (alternative name) (653) + TX, dinex (1089) + TX, dinex-diclexine (1089) + TX, dinobuton (269) + TX, dinocap (270) + TX, dinocap-4 [CCN] + TX, dinocap-6 [CCN] + TX, dinoocton (1090) + TX, dinopenton (1092) + TX, dinosulfon (1097) + TX, dinoterbon (1098) + TX, dioxathion (1102) + TX, diphenyl sulfone (IUPAC name) (1103) + TX, disulfiram (alternative name) [CCN] + TX, disulfoton (278) + TX, DNOC (282) + TX, dofenapyn (1113) + TX, doramectin (alternative name) [CCN] + TX,
 15 endosulfan (294) + TX, endothion (1121) + TX, EPN (297) + TX, eprinomectin (alternative name) [CCN] + TX, ethion (309) + TX, ethoate-methyl (1134) + TX, etoxazole (320) + TX, etrimfos (1142) + TX, fenazaflor (1147) + TX, fenazaquin (328) + TX, fenbutatin oxide (330) + TX, fenothiocarb (337) + TX, fenpropathrin (342) + TX, fenpyrad (alternative name) + TX, fenpyroximate (345) + TX, fenson (1157) + TX, fentrifanil (1161) + TX, fenvalerate (349) + TX, fipronil (354) + TX, fluacrypyrim (360) + TX,
 20 TX, fluazuron (1166) + TX, flubenzimine (1167) + TX, flucyclohexuron (366) + TX, flucythrinate (367) + TX, fluenetil (1169) + TX, flufenoxuron (370) + TX, flumethrin (372) + TX, fluorbenside (1174) + TX, fluvalinate (1184) + TX, FMC 1137 (development code) (1185) + TX, formetanate (405) + TX, formetanate hydrochloride (405) + TX, formothion (1192) + TX, formparanate (1193) + TX, gamma-HCH (430) + TX, glyodin (1205) + TX, halfenprox (424) + TX, heptenophos (432) + TX, hexadecyl
 25 cyclopropanecarboxylate (IUPAC/Chemical Abstracts name) (1216) + TX, hexythiazox (441) + TX, iodomethane (IUPAC name) (542) + TX, isocarbophos (alternative name) (473) + TX, isopropyl O-(methoxyaminothiophosphoryl)salicylate (IUPAC name) (473) + TX, ivermectin (alternative name) [CCN] + TX, jasmolin I (696) + TX, jasmolin II (696) + TX, jodfenphos (1248) + TX, lindane (430) + TX, lufenuron (490) + TX, malathion (492) + TX, malonoben (1254) + TX, mecarbam (502) + TX,
 30 mephosfolan (1261) + TX, mesulfen (alternative name) [CCN] + TX, methacrifos (1266) + TX, methamidophos (527) + TX, methidathion (529) + TX, methiocarb (530) + TX, methomyl (531) + TX, methyl bromide (537) + TX, metolcarb (550) + TX, mevinphos (556) + TX, mexacarbate (1290) + TX, milbemectin (557) + TX, milbemycin oxime (alternative name) [CCN] + TX, mipafox (1293) + TX, monocrotophos (561) + TX, morphothion (1300) + TX, moxidectin (alternative name) [CCN] + TX,
 35 naled (567) + TX, NC-184 (compound code) + TX, NC-512 (compound code) + TX, nifluridide (1309) + TX, nikkomycins (alternative name) [CCN] + TX, nitrilacarb (1313) + TX, nitrilacarb 1:1 zinc chloride complex (1313) + TX, NNI-0101 (compound code) + TX, NNI-0250 (compound code) + TX, omethoate (594) + TX, oxamyl (602) + TX, oxydeprofos (1324) + TX, oxydisulfoton (1325) + TX, pp'-DDT (219) + TX, parathion (615) + TX, permethrin (626) + TX, petroleum oils (alternative name) (628)
 40 + TX, phenkapton (1330) + TX, phenthoate (631) + TX, phorate (636) + TX, phosalone (637) + TX, phosfolan (1338) + TX, phosmet (638) + TX, phosphamidon (639) + TX, phoxim (642) + TX, pirimiphos-methyl (652) + TX, polychloroterpenes (traditional name) (1347) + TX, polynactins

(alternative name) (653) + TX, proclonol (1350) + TX, profenofos (662) + TX, promacyl (1354) + TX, propargite (671) + TX, propetamphos (673) + TX, propoxur (678) + TX, prothidathion (1360) + TX, prothoate (1362) + TX, pyrethrin I (696) + TX, pyrethrin II (696) + TX, pyrethrins (696) + TX, pyridaben (699) + TX, pyridaphenthion (701) + TX, pyrimidifen (706) + TX, pyrimitate (1370) + TX, quinalphos (711) + TX, quintiofos (1381) + TX, R-1492 (development code) (1382) + TX, RA-17 (development code) (1383) + TX, rotenone (722) + TX, schradan (1389) + TX, sebufos (alternative name) + TX, selamectin (alternative name) [CCN] + TX, SI-0009 (compound code) + TX, sophamide (1402) + TX, spirodiclofen (738) + TX, spiromesifen (739) + TX, SSI-121 (development code) (1404) + TX, sulfiram (alternative name) [CCN] + TX, sulfluramid (750) + TX, sulfotep (753) + TX, sulfur (754) + TX, SZI-121 (development code) (757) + TX, tau-fluvalinate (398) + TX, tebufenpyrad (763) + TX, TEPP (1417) + TX, terbam (alternative name) + TX, tetrachlorvinphos (777) + TX, tetradifon (786) + TX, tetranactin (alternative name) (653) + TX, tetrasul (1425) + TX, thiafenox (alternative name) + TX, thiocarboxime (1431) + TX, thiofanox (800) + TX, thiometon (801) + TX, thioquinox (1436) + TX, thuringiensin (alternative name) [CCN] + TX, triamiphos (1441) + TX, triarathene (1443) + TX, triazophos (820) + TX, triazuron (alternative name) + TX, trichlorfon (824) + TX, trifenofos (1455) + TX, trinactin (alternative name) (653) + TX, vamidothion (847) + TX, vaniliprole [CCN] and YI-5302 (compound code) + TX,

an algicide selected from the group of substances consisting of bethoxazin [CCN] + TX, copper dioctanoate (IUPAC name) (170) + TX, copper sulfate (172) + TX, cybutryne [CCN] + TX, dichlone (1052) + TX, dichlorophen (232) + TX, endothal (295) + TX, fentin (347) + TX, hydrated lime [CCN] + TX, nabam (566) + TX, quinoxamine (714) + TX, quinonamid (1379) + TX, simazine (730) + TX, triphenyltin acetate (IUPAC name) (347) and triphenyltin hydroxide (IUPAC name) (347) + TX,

an anthelmintic selected from the group of substances consisting of abamectin (1) + TX, crufomate (1011) + TX, doramectin (alternative name) [CCN] + TX, emamectin (291) + TX, emamectin benzoate (291) + TX, eprinomectin (alternative name) [CCN] + TX, ivermectin (alternative name) [CCN] + TX, milbemycin oxime (alternative name) [CCN] + TX, moxidectin (alternative name) [CCN] + TX, piperazine [CCN] + TX, selamectin (alternative name) [CCN] + TX, spinosad (737) and thiophanate (1435) + TX,

an avicide selected from the group of substances consisting of chloralose (127) + TX, endrin (1122) + TX, fenthion (346) + TX, pyridin-4-amine (IUPAC name) (23) and strychnine (745) + TX,

a bactericide selected from the group of substances consisting of 1-hydroxy-1*H*-pyridine-2-thione (IUPAC name) (1222) + TX, 4-(quinoxalin-2-ylamino)benzenesulfonamide (IUPAC name) (748) + TX, 8-hydroxyquinoline sulfate (446) + TX, bronopol (97) + TX, copper dioctanoate (IUPAC name) (170) + TX, copper hydroxide (IUPAC name) (169) + TX, cresol [CCN] + TX, dichlorophen (232) + TX, dipyrithione (1105) + TX, dodicin (1112) + TX, fenaminosulf (1144) + TX, formaldehyde (404) + TX, hydrargaphen (alternative name) [CCN] + TX, kasugamycin (483) + TX, kasugamycin hydrochloride hydrate (483) + TX, nickel bis(dimethyldithiocarbamate) (IUPAC name) (1308) + TX, nitrapyrin (580) + TX, octhiline (590) + TX, oxolinic acid (606) + TX, oxytetracycline (611) + TX, potassium hydroxyquinoline sulfate (446) + TX, probenazole (658) + TX, streptomycin (744) + TX, streptomycin sesquisulfate (744) + TX, tecloftalam (766) + TX, and thiomersal (alternative name) [CCN] + TX,

a biological agent selected from the group of substances consisting of *Adoxophyes orana* GV (alternative name) (12) + TX, *Agrobacterium radiobacter* (alternative name) (13) + TX, *Amblyseius* spp.

(alternative name) (19) + TX, *Anagrapha falcifera* NPV (alternative name) (28) + TX, *Anagrus atomus* (alternative name) (29) + TX, *Aphelinus abdominalis* (alternative name) (33) + TX, *Aphidius colemani* (alternative name) (34) + TX, *Aphidoletes aphidimyza* (alternative name) (35) + TX, *Autographa californica* NPV (alternative name) (38) + TX, *Bacillus firmus* (alternative name) (48) + TX, *Bacillus*
 5 *sphaericus* Neide (scientific name) (49) + TX, *Bacillus thuringiensis* Berliner (scientific name) (51) + TX, *Bacillus thuringiensis* subsp. *aizawai* (scientific name) (51) + TX, *Bacillus thuringiensis* subsp. *israelensis* (scientific name) (51) + TX, *Bacillus thuringiensis* subsp. *japonensis* (scientific name) (51) + TX, *Bacillus thuringiensis* subsp. *kurstaki* (scientific name) (51) + TX, *Bacillus thuringiensis* subsp. *tenebrionis* (scientific name) (51) + TX, *Beauveria bassiana* (alternative name) (53) + TX, *Beauveria*
 10 *brongniartii* (alternative name) (54) + TX, *Chrysoperla carnea* (alternative name) (151) + TX, *Cryptolaemus montrouzieri* (alternative name) (178) + TX, *Cydia pomonella* GV (alternative name) (191) + TX, *Dacnusa sibirica* (alternative name) (212) + TX, *Diglyphus isaea* (alternative name) (254) + TX, *Encarsia formosa* (scientific name) (293) + TX, *Eretmocerus eremicus* (alternative name) (300) + TX, *Helicoverpa zea* NPV (alternative name) (431) + TX, *Heterorhabditis bacteriophora* and *H. megidis*
 15 (alternative name) (433) + TX, *Hippodamia convergens* (alternative name) (442) + TX, *Leptomastix dactylopii* (alternative name) (488) + TX, *Macrolophus caliginosus* (alternative name) (491) + TX, *Mamestra brassicae* NPV (alternative name) (494) + TX, *Metaphycus helvolus* (alternative name) (522) + TX, *Metarhizium anisopliae* var. *acridum* (scientific name) (523) + TX, *Metarhizium anisopliae* var. *anisopliae* (scientific name) (523) + TX, *Neodiprion sertifer* NPV and *N. lecontei* NPV (alternative name)
 20 (575) + TX, *Orius* spp. (alternative name) (596) + TX, *Paecilomyces fumosoroseus* (alternative name) (613) + TX, *Phytoseiulus persimilis* (alternative name) (644) + TX, *Spodoptera exigua* multicapsid nuclear polyhedrosis virus (scientific name) (741) + TX, *Steinernema bibionis* (alternative name) (742) + TX, *Steinernema carpocapsae* (alternative name) (742) + TX, *Steinernema feltiae* (alternative name) (742) + TX, *Steinernema glaseri* (alternative name) (742) + TX, *Steinernema riobrave* (alternative
 25 name) (742) + TX, *Steinernema riobrans* (alternative name) (742) + TX, *Steinernema scapterisci* (alternative name) (742) + TX, *Steinernema* spp. (alternative name) (742) + TX, *Trichogramma* spp. (alternative name) (826) + TX, *Typhlodromus occidentalis* (alternative name) (844) and *Verticillium lecanii* (alternative name) (848) + TX,

a soil sterilant selected from the group of substances consisting of iodomethane (IUPAC name)
 30 (542) and methyl bromide (537) + TX,

a chemosterilant selected from the group of substances consisting of apholate [CCN] + TX, bisazir (alternative name) [CCN] + TX, busulfan (alternative name) [CCN] + TX, diflubenzuron (250) + TX, dimatif (alternative name) [CCN] + TX, hemel [CCN] + TX, hempa [CCN] + TX, metepa [CCN] + TX, methiotepa [CCN] + TX, methyl apholate [CCN] + TX, morzid [CCN] + TX, penfluron (alternative
 35 name) [CCN] + TX, tepa [CCN] + TX, thiohempa (alternative name) [CCN] + TX, thiotepa (alternative name) [CCN] + TX, tretamine (alternative name) [CCN] and uredepa (alternative name) [CCN] + TX,

an insect pheromone selected from the group of substances consisting of (*E*)-dec-5-en-1-yl acetate with (*E*)-dec-5-en-1-ol (IUPAC name) (222) + TX, (*E*)-tridec-4-en-1-yl acetate (IUPAC name) (829) + TX, (*E*)-6-methylhept-2-en-4-ol (IUPAC name) (541) + TX, (*E,Z*)-tetradeca-4,10-dien-1-yl acetate (IUPAC
 40 name) (779) + TX, (*Z*)-dodec-7-en-1-yl acetate (IUPAC name) (285) + TX, (*Z*)-hexadec-11-enal (IUPAC name) (436) + TX, (*Z*)-hexadec-11-en-1-yl acetate (IUPAC name) (437) + TX, (*Z*)-hexadec-13-en-11-yn-1-yl acetate (IUPAC name) (438) + TX, (*Z*)-icos-13-en-10-one (IUPAC name) (448) + TX, (*Z*)-

- tetradec-7-en-1-al (IUPAC name) (782) + TX, (Z)-tetradec-9-en-1-ol (IUPAC name) (783) + TX, (Z)-tetradec-9-en-1-yl acetate (IUPAC name) (784) + TX, (7E,9Z)-dodeca-7,9-dien-1-yl acetate (IUPAC name) (283) + TX, (9Z,11E)-tetradeca-9,11-dien-1-yl acetate (IUPAC name) (780) + TX, (9Z,12E)-tetradeca-9,12-dien-1-yl acetate (IUPAC name) (781) + TX, 14-methyloctadec-1-ene (IUPAC name) (545) + TX, 4-methylnonan-5-ol with 4-methylnonan-5-one (IUPAC name) (544) + TX, alpha-multistriatin (alternative name) [CCN] + TX, brevicomin (alternative name) [CCN] + TX, codlure (alternative name) [CCN] + TX, codlemone (alternative name) (167) + TX, cuelure (alternative name) (179) + TX, disparlure (277) + TX, dodec-8-en-1-yl acetate (IUPAC name) (286) + TX, dodec-9-en-1-yl acetate (IUPAC name) (287) + TX, dodeca-8 + TX, 10-dien-1-yl acetate (IUPAC name) (284) + TX, dominicalure (alternative name) [CCN] + TX, ethyl 4-methyloctanoate (IUPAC name) (317) + TX, eugenol (alternative name) [CCN] + TX, frontalinal (alternative name) [CCN] + TX, gossyplure (alternative name) (420) + TX, grandlure (421) + TX, grandlure I (alternative name) (421) + TX, grandlure II (alternative name) (421) + TX, grandlure III (alternative name) (421) + TX, grandlure IV (alternative name) (421) + TX, hexalure [CCN] + TX, ipsdienol (alternative name) [CCN] + TX, ipsenol (alternative name) [CCN] + TX, japonilure (alternative name) (481) + TX, lineatin (alternative name) [CCN] + TX, litlure (alternative name) [CCN] + TX, looplure (alternative name) [CCN] + TX, medlure [CCN] + TX, megatomoic acid (alternative name) [CCN] + TX, methyl eugenol (alternative name) (540) + TX, muscalure (563) + TX, octadeca-2,13-dien-1-yl acetate (IUPAC name) (588) + TX, octadeca-3,13-dien-1-yl acetate (IUPAC name) (589) + TX, orfuralure (alternative name) [CCN] + TX, oryctalure (alternative name) (317) + TX, ostramone (alternative name) [CCN] + TX, siglure [CCN] + TX, sordidin (alternative name) (736) + TX, sulcatol (alternative name) [CCN] + TX, tetradec-11-en-1-yl acetate (IUPAC name) (785) + TX, trimedlure (839) + TX, trimedlure A (alternative name) (839) + TX, trimedlure B₁ (alternative name) (839) + TX, trimedlure B₂ (alternative name) (839) + TX, trimedlure C (alternative name) (839) and trunc-call (alternative name) [CCN] + TX,
- an insect repellent selected from the group of substances consisting of 2-(octylthio)ethanol (IUPAC name) (591) + TX, butopyronoxyl (933) + TX, butoxy(polypropylene glycol) (936) + TX, dibutyl adipate (IUPAC name) (1046) + TX, dibutyl phthalate (1047) + TX, dibutyl succinate (IUPAC name) (1048) + TX, diethyltoluamide [CCN] + TX, dimethyl carbate [CCN] + TX, dimethyl phthalate [CCN] + TX, ethyl hexanediol (1137) + TX, hexamide [CCN] + TX, methoquin-butyl (1276) + TX, methylneodecanamide [CCN] + TX, oxamate [CCN] and picaridin [CCN] + TX,
- an insecticide selected from the group of substances consisting of 1-dichloro-1-nitroethane (IUPAC/Chemical Abstracts name) (1058) + TX, 1,1-dichloro-2,2-bis(4-ethylphenyl)ethane (IUPAC name) (1056), + TX, 1,2-dichloropropane (IUPAC/Chemical Abstracts name) (1062) + TX, 1,2-dichloropropane with 1,3-dichloropropene (IUPAC name) (1063) + TX, 1-bromo-2-chloroethane (IUPAC/Chemical Abstracts name) (916) + TX, 2,2,2-trichloro-1-(3,4-dichlorophenyl)ethyl acetate (IUPAC name) (1451) + TX, 2,2-dichlorovinyl 2-ethylsulfinyethyl methyl phosphate (IUPAC name) (1066) + TX, 2-(1,3-dithiolan-2-yl)phenyl dimethylcarbamate (IUPAC/ Chemical Abstracts name) (1109) + TX, 2-(2-butoxyethoxy)ethyl thiocyanate (IUPAC/Chemical Abstracts name) (935) + TX, 2-(4,5-dimethyl-1,3-dioxolan-2-yl)phenyl methylcarbamate (IUPAC/ Chemical Abstracts name) (1084) + TX, 2-(4-chloro-3,5-xylyloxy)ethanol (IUPAC name) (986) + TX, 2-chlorovinyl diethyl phosphate (IUPAC name) (984) + TX, 2-imidazolidone (IUPAC name) (1225) + TX, 2-isovalerylindan-1,3-dione (IUPAC name) (1246) + TX, 2-methyl(prop-2-ynyl)aminophenyl methylcarbamate (IUPAC name) (1284) + TX, 2-thiocyanatoethyl

laurate (IUPAC name) (1433) + TX, 3-bromo-1-chloroprop-1-ene (IUPAC name) (917) + TX, 3-methyl-1-phenylpyrazol-5-yl dimethylcarbamate (IUPAC name) (1283) + TX, 4-methyl(prop-2-ynyl)amino-3,5-xylyl methylcarbamate (IUPAC name) (1285) + TX, 5,5-dimethyl-3-oxocyclohex-1-enyl dimethylcarbamate (IUPAC name) (1085) + TX, abamectin (1) + TX, acephate (2) + TX, acetamiprid (4) + TX, acethion
 5 (alternative name) [CCN] + TX, acetoprole [CCN] + TX, acrinathrin (9) + TX, acrylonitrile (IUPAC name) (861) + TX, alanycarb (15) + TX, aldicarb (16) + TX, aldoxycarb (863) + TX, aldrin (864) + TX, allethrin (17) + TX, allosamidin (alternative name) [CCN] + TX, allyxycarb (866) + TX, alpha-cypermethrin (202) + TX, alpha-ecdysone (alternative name) [CCN] + TX, aluminium phosphide (640) + TX, amidithion (870) + TX, amidothioate (872) + TX, aminocarb (873) + TX, amiton (875) + TX,
 10 amiton hydrogen oxalate (875) + TX, amitraz (24) + TX, anabasine (877) + TX, athidathion (883) + TX, AVI 382 (compound code) + TX, AZ 60541 (compound code) + TX, azadirachtin (alternative name) (41) + TX, azamethiphos (42) + TX, azinphos-ethyl (44) + TX, azinphos-methyl (45) + TX, azothoate (889) + TX, *Bacillus thuringiensis* delta endotoxins (alternative name) (52) + TX, barium hexafluorosilicate (alternative name) [CCN] + TX, barium polysulfide (IUPAC/Chemical Abstracts name)
 15 (892) + TX, barthrin [CCN] + TX, Bayer 22/190 (development code) (893) + TX, Bayer 22408 (development code) (894) + TX, bendiocarb (58) + TX, benfuracarb (60) + TX, bensultap (66) + TX, beta-cyfluthrin (194) + TX, beta-cypermethrin (203) + TX, bifenthrin (76) + TX, bioallethrin (78) + TX, bioallethrin S-cyclopentenyl isomer (alternative name) (79) + TX, bioethanomethrin [CCN] + TX, biopermethrin (908) + TX, bioresmethrin (80) + TX, bis(2-chloroethyl) ether (IUPAC name) (909) + TX,
 20 bistrifluron (83) + TX, borax (86) + TX, brofenvalerate (alternative name) + TX, bromfenvinfos (914) + TX, bromocyclen (918) + TX, bromo-DDT (alternative name) [CCN] + TX, bromophos (920) + TX, bromophos-ethyl (921) + TX, bufencarb (924) + TX, buprofezin (99) + TX, butacarb (926) + TX, butathiofos (927) + TX, butocarboxim (103) + TX, butonate (932) + TX, butoxycarboxim (104) + TX, butylpyridaben (alternative name) + TX, cadusafos (109) + TX, calcium arsenate [CCN] + TX, calcium
 25 cyanide (444) + TX, calcium polysulfide (IUPAC name) (111) + TX, camphechlor (941) + TX, carbanolate (943) + TX, carbaryl (115) + TX, carbofuran (118) + TX, carbon disulfide (IUPAC/Chemical Abstracts name) (945) + TX, carbon tetrachloride (IUPAC name) (946) + TX, carbophenothion (947) + TX, carbosulfan (119) + TX, cartap (123) + TX, cartap hydrochloride (123) + TX, cevadine (alternative name) (725) + TX, chlorbicyclen (960) + TX, chlordane (128) + TX,
 30 chlordacone (963) + TX, chlordimeform (964) + TX, chlordimeform hydrochloride (964) + TX, chlorethoxyfos (129) + TX, chlorfenapyr (130) + TX, chlorfenvinphos (131) + TX, chlorfluazuron (132) + TX, chlormephos (136) + TX, chloroform [CCN] + TX, chloropicrin (141) + TX, chlorphoxim (989) + TX, chlorprazophos (990) + TX, chlorpyrifos (145) + TX, chlorpyrifos-methyl (146) + TX, chlorthiophos (994) + TX, chromafenozide (150) + TX, cinerin I (696) + TX, cinerin II (696) + TX,
 35 cinerins (696) + TX, cis-resmethrin (alternative name) + TX, cismethrin (80) + TX, clocythrion (alternative name) + TX, cloethocarb (999) + TX, closantel (alternative name) [CCN] + TX, clothianidin (165) + TX, copper acetoarsenite [CCN] + TX, copper arsenate [CCN] + TX, copper oleate [CCN] + TX, coumaphos (174) + TX, coumithoate (1006) + TX, crotamiton (alternative name) [CCN] + TX, crotoxyphos (1010) + TX, cruformate (1011) + TX, cryolite (alternative name) (177) + TX, CS 708
 40 (development code) (1012) + TX, cyanofenphos (1019) + TX, cyanophos (184) + TX, cyanthoate (1020) + TX, cyclethrin [CCN] + TX, cycloprothrin (188) + TX, cyfluthrin (193) + TX, cyhalothrin (196) + TX, cypermethrin (201) + TX, cyphenothrin (206) + TX, cyromazine (209) + TX, cythioate

(alternative name) [CCN] + TX, *d*-limonene (alternative name) [CCN] + TX, *d*-tetramethrin (alternative name) (788) + TX, DAEP (1031) + TX, dazomet (216) + TX, DDT (219) + TX, decarbofuran (1034) + TX, deltamethrin (223) + TX, demephion (1037) + TX, demephion-O (1037) + TX, demephion-S (1037) + TX, demeton (1038) + TX, demeton-methyl (224) + TX, demeton-O (1038) + TX, demeton-O-methyl (224) + TX, demeton-S (1038) + TX, demeton-S-methyl (224) + TX, demeton-S-methylsulphon (1039) + TX, diafenthion (226) + TX, dialifos (1042) + TX, diamidafos (1044) + TX, diazinon (227) + TX, dicapthon (1050) + TX, dichlofenthion (1051) + TX, dichlorvos (236) + TX, dicliphos (alternative name) + TX, dicresyl (alternative name) [CCN] + TX, dicrotophos (243) + TX, dicyclanil (244) + TX, dieldrin (1070) + TX, diethyl 5-methylpyrazol-3-yl phosphate (IUPAC name) (1076) + TX, diflubenzuron (250) + TX, dilor (alternative name) [CCN] + TX, dimefluthrin [CCN] + TX, dimefox (1081) + TX, dimetan (1085) + TX, dimethoate (262) + TX, dimethrin (1083) + TX, dimethylvinphos (265) + TX, dimetilan (1086) + TX, dinex (1089) + TX, dinex-diclexine (1089) + TX, dinoprop (1093) + TX, dinosam (1094) + TX, dinoseb (1095) + TX, dinotefuran (271) + TX, diofenolan (1099) + TX, dioxabenzofos (1100) + TX, dioxacarb (1101) + TX, dioxathion (1102) + TX, disulfoton (278) + TX, dithicrofos (1108) + TX, DNOC (282) + TX, doramectin (alternative name) [CCN] + TX, DSP (1115) + TX, ecdysterone (alternative name) [CCN] + TX, EI 1642 (development code) (1118) + TX, emamectin (291) + TX, emamectin benzoate (291) + TX, EMPC (1120) + TX, empenthrin (292) + TX, endosulfan (294) + TX, endothion (1121) + TX, endrin (1122) + TX, EPBP (1123) + TX, EPN (297) + TX, epofenonane (1124) + TX, eprinomectin (alternative name) [CCN] + TX, esfenvalerate (302) + TX, etaphos (alternative name) [CCN] + TX, ethiofencarb (308) + TX, ethion (309) + TX, ethiprole (310) + TX, ethoate-methyl (1134) + TX, ethoprophos (312) + TX, ethyl formate (IUPAC name) [CCN] + TX, ethyl-DDD (alternative name) (1056) + TX, ethylene dibromide (316) + TX, ethylene dichloride (chemical name) (1136) + TX, ethylene oxide [CCN] + TX, etofenprox (319) + TX, etrimfos (1142) + TX, EXD (1143) + TX, famphur (323) + TX, fenamiphos (326) + TX, fenazaflor (1147) + TX, fenchlorphos (1148) + TX, fenethacarb (1149) + TX, fenfluthrin (1150) + TX, fenitrothion (335) + TX, fenobucarb (336) + TX, fenoxacrim (1153) + TX, fenoxycarb (340) + TX, fenpirithrin (1155) + TX, fenpropathrin (342) + TX, fenpyrad (alternative name) + TX, fensulfothion (1158) + TX, fenthion (346) + TX, fenthion-ethyl [CCN] + TX, fenvalerate (349) + TX, fipronil (354) + TX, flonicamid (358) + TX, flubendiamide (CAS. Reg. No.: 272451-65-7) + TX, flucufuron (1168) + TX, flucycloxuron (366) + TX, flucythrinate (367) + TX, fluenetil (1169) + TX, flufenerim [CCN] + TX, flufenoxuron (370) + TX, flufenprox (1171) + TX, flumethrin (372) + TX, fluvalinate (1184) + TX, FMC 1137 (development code) (1185) + TX, fonofos (1191) + TX, formetanate (405) + TX, formetanate hydrochloride (405) + TX, formothion (1192) + TX, formparanate (1193) + TX, fosmethilan (1194) + TX, fospirate (1195) + TX, fosthiazate (408) + TX, fosthietan (1196) + TX, furathiocarb (412) + TX, furethrin (1200) + TX, gamma-cyhalothrin (197) + TX, gamma-HCH (430) + TX, guazatine (422) + TX, guazatine acetates (422) + TX, GY-81 (development code) (423) + TX, halfenprox (424) + TX, halofenozide (425) + TX, HCH (430) + TX, HEOD (1070) + TX, heptachlor (1211) + TX, heptenophos (432) + TX, heterophos [CCN] + TX, hexaflumuron (439) + TX, HHN (864) + TX, hydramethylnon (443) + TX, hydrogen cyanide (444) + TX, hydroprene (445) + TX, hyquincarb (1223) + TX, imidacloprid (458) + TX, imiprothrin (460) + TX, indoxacarb (465) + TX, iodomethane (IUPAC name) (542) + TX, IPSP (1229) + TX, isazofos (1231) + TX, isobenzan (1232) + TX, isocarbophos (alternative name) (473) + TX, isodrin (1235) + TX, isofenphos (1236) + TX, isolane (1237) + TX,

isoprocab (472) + TX, isopropyl O-(methoxyaminothiophosphoryl)salicylate (IUPAC name) (473) + TX, isoprothiolane (474) + TX, isothioate (1244) + TX, isoxathion (480) + TX, ivermectin (alternative name) [CCN] + TX, jasmolin I (696) + TX, jasmolin II (696) + TX, jodfenphos (1248) + TX, juvenile hormone I (alternative name) [CCN] + TX, juvenile hormone II (alternative name) [CCN] + TX, juvenile hormone III (alternative name) [CCN] + TX, kelevan (1249) + TX, kinoprene (484) + TX, lambda-cyhalothrin (198) + TX, lead arsenate [CCN] + TX, lepimectin (CCN) + TX, leptophos (1250) + TX, lindane (430) + TX, lirimfos (1251) + TX, lufenuron (490) + TX, lythidathion (1253) + TX, *m*-cumenyl methylcarbamate (IUPAC name) (1014) + TX, magnesium phosphide (IUPAC name) (640) + TX, malathion (492) + TX, malonoben (1254) + TX, mazidox (1255) + TX, mecarbam (502) + TX, mecarphon (1258) + TX, menazon (1260) + TX, mephosfolan (1261) + TX, mercurous chloride (513) + TX, mesulfenfos (1263) + TX, metaflumizone (CCN) + TX, metam (519) + TX, metam-potassium (alternative name) (519) + TX, metam-sodium (519) + TX, methacrifos (1266) + TX, methamidophos (527) + TX, methanesulfonyl fluoride (IUPAC/Chemical Abstracts name) (1268) + TX, methidathion (529) + TX, methiocarb (530) + TX, methocrotophos (1273) + TX, methomyl (531) + TX, methoprene (532) + TX, methoquin-butyl (1276) + TX, methothrin (alternative name) (533) + TX, methoxychlor (534) + TX, methoxyfenozide (535) + TX, methyl bromide (537) + TX, methyl isothiocyanate (543) + TX, methylchloroform (alternative name) [CCN] + TX, methylene chloride [CCN] + TX, metofluthrin [CCN] + TX, metolcarb (550) + TX, metoxadiazone (1288) + TX, mevinphos (556) + TX, mexacarbate (1290) + TX, milbemectin (557) + TX, milbemycin oxime (alternative name) [CCN] + TX, mipafox (1293) + TX, mirex (1294) + TX, monocrotophos (561) + TX, morphothion (1300) + TX, moxidectin (alternative name) [CCN] + TX, naftalofos (alternative name) [CCN] + TX, naled (567) + TX, naphthalene (IUPAC/Chemical Abstracts name) (1303) + TX, NC-170 (development code) (1306) + TX, NC-184 (compound code) + TX, nicotine (578) + TX, nicotine sulfate (578) + TX, nifluridide (1309) + TX, nitenpyram (579) + TX, nithiazine (1311) + TX, nitrilacarb (1313) + TX, nitrilacarb 1:1 zinc chloride complex (1313) + TX, NNI-0101 (compound code) + TX, NNI-0250 (compound code) + TX, nornicotine (traditional name) (1319) + TX, novaluron (585) + TX, noviflumuron (586) + TX, O-5-dichloro-4-iodophenyl O-ethyl ethylphosphonothioate (IUPAC name) (1057) + TX, O,O-diethyl O-4-methyl-2-oxo-2*H*-chromen-7-yl phosphorothioate (IUPAC name) (1074) + TX, O,O-diethyl O-6-methyl-2-propylpyrimidin-4-yl phosphorothioate (IUPAC name) (1075) + TX, O,O,O',O'-tetrapropyl dithiopyrophosphate (IUPAC name) (1424) + TX, oleic acid (IUPAC name) (593) + TX, omethoate (594) + TX, oxamyl (602) + TX, oxydemeton-methyl (609) + TX, oxydeprofos (1324) + TX, oxydisulfoton (1325) + TX, pp'-DDT (219) + TX, para-dichlorobenzene [CCN] + TX, parathion (615) + TX, parathion-methyl (616) + TX, penfluron (alternative name) [CCN] + TX, pentachlorophenol (623) + TX, pentachlorophenyl laurate (IUPAC name) (623) + TX, permethrin (626) + TX, petroleum oils (alternative name) (628) + TX, PH 60-38 (development code) (1328) + TX, phenkapton (1330) + TX, phenothrin (630) + TX, phenthoate (631) + TX, phorate (636) + TX, phosalone (637) + TX, phosfolan (1338) + TX, phosmet (638) + TX, phosnichlor (1339) + TX, phosphamidon (639) + TX, phosphine (IUPAC name) (640) + TX, phoxim (642) + TX, phoxim-methyl (1340) + TX, pirimetaphos (1344) + TX, pirimicarb (651) + TX, pirimiphos-ethyl (1345) + TX, pirimiphos-methyl (652) + TX, polychlorodicyclopentadiene isomers (IUPAC name) (1346) + TX, polychloroterpenes (traditional name) (1347) + TX, potassium arsenite [CCN] + TX, potassium thiocyanate [CCN] + TX, prallethrin (655) + TX, precocene I (alternative name) [CCN] + TX, precocene II (alternative name) [CCN] + TX,

precocene III (alternative name) [CCN] + TX, primidophos (1349) + TX, profenofos (662) + TX,
 profluthrin [CCN] + TX, promacyl (1354) + TX, promecarb (1355) + TX, propaphos (1356) + TX,
 propetamphos (673) + TX, propoxur (678) + TX, prothidathion (1360) + TX, prothiofos (686) + TX,
 prothoate (1362) + TX, protrifenbute [CCN] + TX, pymetrozine (688) + TX, pyraclofos (689) + TX,
 5 pyrazophos (693) + TX, pyresmethrin (1367) + TX, pyrethrin I (696) + TX, pyrethrin II (696) + TX,
 pyrethrins (696) + TX, pyridaben (699) + TX, pyridalyl (700) + TX, pyridaphenthion (701) + TX,
 pyrimidifen (706) + TX, pyrimitate (1370) + TX, pyriproxyfen (708) + TX, quassia (alternative name)
 [CCN] + TX, quinalphos (711) + TX, quinalphos-methyl (1376) + TX, quinothion (1380) + TX,
 quintiofos (1381) + TX, R-1492 (development code) (1382) + TX, rafoxanide (alternative name) [CCN]
 10 + TX, resmethrin (719) + TX, rotenone (722) + TX, RU 15525 (development code) (723) + TX, RU
 25475 (development code) (1386) + TX, ryania (alternative name) (1387) + TX, ryanodine (traditional
 name) (1387) + TX, sabadilla (alternative name) (725) + TX, schradan (1389) + TX, sebufos
 (alternative name) + TX, selamectin (alternative name) [CCN] + TX, SI-0009 (compound code) + TX,
 SI-0205 (compound code) + TX, SI-0404 (compound code) + TX, SI-0405 (compound code) + TX,
 15 silafluofen (728) + TX, SN 72129 (development code) (1397) + TX, sodium arsenite [CCN] + TX,
 sodium cyanide (444) + TX, sodium fluoride (IUPAC/Chemical Abstracts name) (1399) + TX, sodium
 hexafluorosilicate (1400) + TX, sodium pentachlorophenoxide (623) + TX, sodium selenate (IUPAC
 name) (1401) + TX, sodium thiocyanate [CCN] + TX, sophamide (1402) + TX, spinosad (737) + TX,
 spiromesifen (739) + TX, spirotetmat (CCN) + TX, sulcofuron (746) + TX, sulcofuron-sodium (746) +
 20 TX, sulfluramid (750) + TX, sulfotep (753) + TX, sulfuryl fluoride (756) + TX, sulprofos (1408) + TX,
 tar oils (alternative name) (758) + TX, tau-fluvalinate (398) + TX, tazimcarb (1412) + TX, TDE (1414)
 + TX, tebufenozide (762) + TX, tebufenpyrad (763) + TX, tebupirimfos (764) + TX, teflubenzuron
 (768) + TX, tefluthrin (769) + TX, temephos (770) + TX, TEPP (1417) + TX, terallethrin (1418) + TX,
 terbam (alternative name) + TX, terbufos (773) + TX, tetrachloroethane [CCN] + TX, tetrachlorvinphos
 25 (777) + TX, tetramethrin (787) + TX, theta-cypermethrin (204) + TX, thiaclopid (791) + TX, thiafenox
 (alternative name) + TX, thiamethoxam (792) + TX, thicrofos (1428) + TX, thiocarboxime (1431) + TX,
 thiocyclam (798) + TX, thiocyclam hydrogen oxalate (798) + TX, thiodicarb (799) + TX, thiofanox
 (800) + TX, thiometon (801) + TX, thionazin (1434) + TX, thiosultap (803) + TX, thiosultap-sodium
 (803) + TX, thuringiensin (alternative name) [CCN] + TX, tolfenpyrad (809) + TX, tralomethrin (812) +
 30 TX, transluthrin (813) + TX, transpermethrin (1440) + TX, triamiphos (1441) + TX, triazamate (818)
 + TX, triazophos (820) + TX, triazuron (alternative name) + TX, trichlorfon (824) + TX,
 trichlormetaphos-3 (alternative name) [CCN] + TX, trichloronat (1452) + TX, trifenofos (1455) + TX,
 triflumuron (835) + TX, trimethacarb (840) + TX, triprene (1459) + TX, vamidothion (847) + TX,
 vaniliprole [CCN] + TX, veratridine (alternative name) (725) + TX, veratrine (alternative name) (725) +
 35 TX, XMC (853) + TX, xylylcarb (854) + TX, YI-5302 (compound code) + TX, zeta-cypermethrin (205)
 + TX, zetamethrin (alternative name) + TX, zinc phosphide (640) + TX, zolaprofos (1469) and ZXI
 8901 (development code) (858) + TX, cyantraniliprole [736994-63-19 + TX, chlorantraniliprole [500008-
 45-7] + TX, cyenopyrafen [560121-52-0] + TX, cyflumetofen [400882-07-7] + TX, pyrifluquinazon
 [337458-27-2] + TX, spinetoram [187166-40-1 + 187166-15-0] + TX, spirotetramat [203313-25-1] + TX,
 40 sulfoxaflo [946578-00-3] + TX, flufiprole [704886-18-0] + TX, meperfluthrin [915288-13-0] + TX,
 tetramethylfluthrin [84937-88-2] + TX, triflumezopyrim (disclosed in WO 2012/092115) + TX,
 fluxametamide (WO 2007/026965) + TX, epsilon-metofluthrin [240494-71-7] + TX, epsilon-momfluorothrin

[1065124-65-3] + TX, fluazaindolizine [1254304-22-7] + TX, chloroprallethrin [399572-87-3] + TX, fluxametamide [928783-29-3] + TX, cyhalodiamide [1262605-53-7] + TX, tioxazafen [330459-31-9] + TX, broflanilide [1207727-04-5] + TX, flufiprole [704886-18-0] + TX, cyclaniliprole [1031756-98-5] + TX, tetraniliprole [1229654-66-3] + TX, guadi pyr (described in WO2010/060231) + TX, cycloxaprid (described in WO2005/077934) + TX,

a molluscicide selected from the group of substances consisting of bis(tributyltin) oxide (IUPAC name) (913) + TX, bromoacetamide [CCN] + TX, calcium arsenate [CCN] + TX, cloethocarb (999) + TX, copper acetoarsenite [CCN] + TX, copper sulfate (172) + TX, fentin (347) + TX, ferric phosphate (IUPAC name) (352) + TX, metaldehyde (518) + TX, methiocarb (530) + TX, niclosamide (576) + TX, niclosamide-olamine (576) + TX, pentachlorophenol (623) + TX, sodium pentachlorophenoxide (623) + TX, tazimcarb (1412) + TX, thiodicarb (799) + TX, tributyltin oxide (913) + TX, trifenmorph (1454) + TX, trimethacarb (840) + TX, triphenyltin acetate (IUPAC name) (347) and triphenyltin hydroxide (IUPAC name) (347) + TX, pyriprole [394730-71-3] + TX,

a nematicide selected from the group of substances consisting of AKD-3088 (compound code) + TX, 1,2-dibromo-3-chloropropane (IUPAC/Chemical Abstracts name) (1045) + TX, 1,2-dichloropropane (IUPAC/ Chemical Abstracts name) (1062) + TX, 1,2-dichloropropane with 1,3-dichloropropene (IUPAC name) (1063) + TX, 1,3-dichloropropene (233) + TX, 3,4-dichlorotetrahydrothiophene 1,1-dioxide (IUPAC/Chemical Abstracts name) (1065) + TX, 3-(4-chlorophenyl)-5-methylrhodanine (IUPAC name) (980) + TX, 5-methyl-6-thioxo-1,3,5-thiadiazinan-3-ylacetic acid (IUPAC name) (1286) + TX, 6-isopentenylaminopurine (alternative name) (210) + TX, abamectin (1) + TX, acetoprole [CCN] + TX, alanycarb (15) + TX, aldicarb (16) + TX, aldoxycarb (863) + TX, AZ 60541 (compound code) + TX, benclonthiaz [CCN] + TX, benomyl (62) + TX, butylpyridaben (alternative name) + TX, cadusafos (109) + TX, carbofuran (118) + TX, carbon disulfide (945) + TX, carbosulfan (119) + TX, chloropicrin (141) + TX, chlorpyrifos (145) + TX, cloethocarb (999) + TX, cytokinins (alternative name) (210) + TX, dazomet (216) + TX, DBCP (1045) + TX, DCIP (218) + TX, diamidafos (1044) + TX, dichlofenthion (1051) + TX, dicliphos (alternative name) + TX, dimethoate (262) + TX, doramectin (alternative name) [CCN] + TX, emamectin (291) + TX, emamectin benzoate (291) + TX, eprinomectin (alternative name) [CCN] + TX, ethoprophos (312) + TX, ethylene dibromide (316) + TX, fenamiphos (326) + TX, fenpyrad (alternative name) + TX, fensulfthion (1158) + TX, fosthiazate (408) + TX, fosthietan (1196) + TX, furfural (alternative name) [CCN] + TX, GY-81 (development code) (423) + TX, heterophos [CCN] + TX, iodomethane (IUPAC name) (542) + TX, isamidofos (1230) + TX, isazofos (1231) + TX, ivermectin (alternative name) [CCN] + TX, kinetin (alternative name) (210) + TX, mecarphon (1258) + TX, metam (519) + TX, metam-potassium (alternative name) (519) + TX, metam-sodium (519) + TX, methyl bromide (537) + TX, methyl isothiocyanate (543) + TX, milbemycin oxime (alternative name) [CCN] + TX, moxidectin (alternative name) [CCN] + TX, *Myrothecium verrucaria* composition (alternative name) (565) + TX, NC-184 (compound code) + TX, oxamyl (602) + TX, phorate (636) + TX, phosphamidon (639) + TX, phosphocarb [CCN] + TX, sebufos (alternative name) + TX, selamectin (alternative name) [CCN] + TX, spinosad (737) + TX, terbam (alternative name) + TX, terbufos (773) + TX, tetrachlorothiophene (IUPAC/ Chemical Abstracts name) (1422) + TX, thiafenox (alternative name) + TX, thionazin (1434) + TX, triazophos (820) + TX, triazuron (alternative name) + TX, xyleneols [CCN] + TX, YI-5302 (compound code) and zeatin (alternative name) (210) + TX, fluensulfone [318290-98-1] + TX,

a nitrification inhibitor selected from the group of substances consisting of potassium ethylxanthate [CCN] and nitrapyrin (580) + TX,

a plant activator selected from the group of substances consisting of acibenzolar (6) + TX, acibenzolar-S-methyl (6) + TX, probenazole (658) and *Reynoutria sachalinensis* extract (alternative name) (720) + TX,

a rodenticide selected from the group of substances consisting of 2-isovalerylindan-1,3-dione (IUPAC name) (1246) + TX, 4-(quinoxalin-2-ylamino)benzenesulfonamide (IUPAC name) (748) + TX, alpha-chlorohydrin [CCN] + TX, aluminium phosphide (640) + TX, antu (880) + TX, arsenous oxide (882) + TX, barium carbonate (891) + TX, bithiosemi (912) + TX, brodifacoum (89) + TX, bromadiolone (91) + TX, bromethalin (92) + TX, calcium cyanide (444) + TX, chloralose (127) + TX, chlorophacinone (140) + TX, cholecalciferol (alternative name) (850) + TX, coumachlor (1004) + TX, coumafuryl (1005) + TX, coumatetralyl (175) + TX, crimidine (1009) + TX, difenacoum (246) + TX, difethialone (249) + TX, diphacinone (273) + TX, ergocalciferol (301) + TX, flocoumafen (357) + TX, fluoroacetamide (379) + TX, flupropadine (1183) + TX, flupropadine hydrochloride (1183) + TX, gamma-HCH (430) + TX, HCH (430) + TX, hydrogen cyanide (444) + TX, iodomethane (IUPAC name) (542) + TX, lindane (430) + TX, magnesium phosphide (IUPAC name) (640) + TX, methyl bromide (537) + TX, norbormide (1318) + TX, phosacetim (1336) + TX, phosphine (IUPAC name) (640) + TX, phosphorus [CCN] + TX, pindone (1341) + TX, potassium arsenite [CCN] + TX, pyrinuron (1371) + TX, scilliroside (1390) + TX, sodium arsenite [CCN] + TX, sodium cyanide (444) + TX, sodium fluoroacetate (735) + TX, strychnine (745) + TX, thallium sulfate [CCN] + TX, warfarin (851) and zinc phosphide (640) + TX,

a synergist selected from the group of substances consisting of 2-(2-butoxyethoxy)ethyl piperonylate (IUPAC name) (934) + TX, 5-(1,3-benzodioxol-5-yl)-3-hexylcyclohex-2-enone (IUPAC name) (903) + TX, farnesol with nerolidol (alternative name) (324) + TX, MB-599 (development code) (498) + TX, MGK 264 (development code) (296) + TX, piperonyl butoxide (649) + TX, piprotal (1343) + TX, propyl isomer (1358) + TX, S421 (development code) (724) + TX, sesamex (1393) + TX, sesasmolin (1394) and sulfoxide (1406) + TX,

an animal repellent selected from the group of substances consisting of anthraquinone (32) + TX, chloralose (127) + TX, copper naphthenate [CCN] + TX, copper oxychloride (171) + TX, diazinon (227) + TX, dicyclopentadiene (chemical name) (1069) + TX, guazatine (422) + TX, guazatine acetates (422) + TX, methiocarb (530) + TX, pyridin-4-amine (IUPAC name) (23) + TX, thiram (804) + TX, trimethacarb (840) + TX, zinc naphthenate [CCN] and ziram (856) + TX,

a virucide selected from the group of substances consisting of imanin (alternative name) [CCN] and ribavirin (alternative name) [CCN] + TX,

a wound protectant selected from the group of substances consisting of mercuric oxide (512) + TX, octhilinone (590) and thiophanate-methyl (802) + TX,

and biologically active compounds selected from the group consisting of azaconazole (60207-31-0) + TX, bitertanol [70585-36-3] + TX, bromuconazole [116255-48-2] + TX, cyproconazole [94361-06-5] + TX, difenoconazole [119446-68-3] + TX, diniconazole [83657-24-3] + TX, epoxiconazole [106325-08-0] + TX, fenbuconazole [114369-43-6] + TX, fluquinconazole [136426-54-5] + TX, flusilazole [85509-19-9] + TX, flutriafol [76674-21-0] + TX, hexaconazole [79983-71-4] + TX, imazalil [35554-44-0] + TX, imibenconazole [86598-92-7] + TX, ipconazole [125225-28-7] + TX, metconazole [125116-

23-6] + TX, myclobutanil [88671-89-0] + TX, pefurazoate [101903-30-4] + TX, penconazole [66246-88-6] + TX, prothioconazole [178928-70-6] + TX, pyrifenoxy [88283-41-4] + TX, prochloraz [67747-09-5] + TX, propiconazole [60207-90-1] + TX, simeconazole [149508-90-7] + TX, tebuconazole [107534-96-3] + TX, tetraconazole [112281-77-3] + TX, triadimefon [43121-43-3] + TX, triadimenol [55219-65-3] + TX, triflumizole [99387-89-0] + TX, triticonazole [131983-72-7] + TX, ancymidol [12771-68-5] + TX, fenarimol [60168-88-9] + TX, nuarimol [63284-71-9] + TX, bupirimate [41483-43-6] + TX, dimethirimol [5221-53-4] + TX, ethirimol [23947-60-6] + TX, dodemorph [1593-77-7] + TX, fenpropidine [67306-00-7] + TX, fenpropimorph [67564-91-4] + TX, spiroxamine [118134-30-8] + TX, tridemorph [81412-43-3] + TX, cyprodinil [121552-61-2] + TX, mepanipyrim [110235-47-7] + TX, pyrimethanil [53112-28-0] + TX, fenpiclonil [74738-17-3] + TX, fludioxonil [131341-86-1] + TX, benalaxyl [71626-11-4] + TX, furalaxyl [57646-30-7] + TX, metalaxyl [57837-19-1] + TX, R-metalaxyl [70630-17-0] + TX, ofurace [58810-48-3] + TX, oxadixyl [77732-09-3] + TX, benomyl [17804-35-2] + TX, carbendazim [10605-21-7] + TX, debacarb [62732-91-6] + TX, fuberidazole [3878-19-1] + TX, thiabendazole [148-79-8] + TX, chlozolinate [84332-86-5] + TX, dichlozoline [24201-58-9] + TX, iprodione [36734-19-7] + TX, myclozoline [54864-61-8] + TX, procymidone [32809-16-8] + TX, vinclozoline [50471-44-8] + TX, boscalid [188425-85-6] + TX, carboxin [5234-68-4] + TX, fenfuram [24691-80-3] + TX, flutolanil [66332-96-5] + TX, mepronil [55814-41-0] + TX, oxycarboxin [5259-88-1] + TX, penthiopyrad [183675-82-3] + TX, thifluzamide [130000-40-7] + TX, guazatine [108173-90-6] + TX, dodine [2439-10-3] [112-65-2] (free base) + TX, iminoctadine [13516-27-3] + TX, azoxystrobin [131860-33-8] + TX, dimoxystrobin [149961-52-4] + TX, enestroburin {Proc. BCPC, Int. Congr., Glasgow, 2003, 1, 93} + TX, fluoxastrobin [361377-29-9] + TX, kresoxim-methyl [143390-89-0] + TX, metominostrobin [133408-50-1] + TX, trifloxystrobin [141517-21-7] + TX, orysastrobin [248593-16-0] + TX, picoxystrobin [117428-22-5] + TX, pyraclostrobin [175013-18-0] + TX, ferbam [14484-64-1] + TX, mancozeb [8018-01-7] + TX, maneb [12427-38-2] + TX, metiram [9006-42-2] + TX, propineb [12071-83-9] + TX, thiram [137-26-8] + TX, zineb [12122-67-7] + TX, ziram [137-30-4] + TX, captafol [2425-06-1] + TX, captan [133-06-2] + TX, dichlofluanid [1085-98-9] + TX, fluoroimide [41205-21-4] + TX, folpet [133-07-3] + TX, tolylfluanid [731-27-1] + TX, bordeaux mixture [8011-63-0] + TX, copperhydroxid [20427-59-2] + TX, copperoxychlorid [1332-40-7] + TX, coppersulfat [7758-98-7] + TX, copperoxid [1317-39-1] + TX, mancopper [53988-93-5] + TX, oxine-copper [10380-28-6] + TX, dinocap [131-72-6] + TX, nitrothal-isopropyl [10552-74-6] + TX, edifenphos [17109-49-8] + TX, iprobenphos [26087-47-8] + TX, isoprothiolane [50512-35-1] + TX, phosdiphen [36519-00-3] + TX, pyrazophos [13457-18-6] + TX, tolclofos-methyl [57018-04-9] + TX, acibenzolar-S-methyl [135158-54-2] + TX, anilazine [101-05-3] + TX, benthiavalicarb [413615-35-7] + TX, blasticidin-S [2079-00-7] + TX, chinomethionat [2439-01-2] + TX, chloroneb [2675-77-6] + TX, chlorothalonil [1897-45-6] + TX, cyflufenamid [180409-60-3] + TX, cymoxanil [57966-95-7] + TX, dichlone [117-80-6] + TX, diclocymet [139920-32-4] + TX, diclomezine [62865-36-5] + TX, dicloran [99-30-9] + TX, diethofencarb [87130-20-9] + TX, dimethomorph [110488-70-5] + TX, SYP-LI90 (Flumorph) [211867-47-9] + TX, dithianon [3347-22-6] + TX, ethaboxam [162650-77-3] + TX, etridiazole [2593-15-9] + TX, famoxadone [131807-57-3] + TX, fenamidone [161326-34-7] + TX, fenoxanil [115852-48-7] + TX, fentin [668-34-8] + TX, ferimzone [89269-64-7] + TX, fluazinam [79622-59-6] + TX, fluopicolide [239110-15-7] + TX, flusulfamide [106917-52-6] + TX, fenhexamid [126833-17-8] + TX, fosetyl-aluminium [39148-24-8] + TX, hymexazol [10004-44-1] + TX, iprovalicarb [140923-17-7] + TX, IKF-916 (Cyazofamid) [120116-

88-3] + TX, kasugamycin [6980-18-3] + TX, methasulfocarb [66952-49-6] + TX, metrafenone [220899-03-6] + TX, pencycuron [66063-05-6] + TX, phthalide [27355-22-2] + TX, polyoxins [11113-80-7] + TX, probenazole [27605-76-1] + TX, propamocarb [25606-41-1] + TX, proquinazid [189278-12-4] + TX, pyroquilon [57369-32-1] + TX, quinoxifen [124495-18-7] + TX, quintozone [82-68-8] + TX, sulfur
 5 [7704-34-9] + TX, tiadinil [223580-51-6] + TX, triazoxide [72459-58-6] + TX, tricyclazole [41814-78-2] + TX, triforine [26644-46-2] + TX, validamycin [37248-47-8] + TX, zoxamide (RH7281) [156052-68-5] + TX, mandipropamid [374726-62-2] + TX, isopyrazam [881685-58-1] + TX, sedaxane [874967-67-6] + TX, 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxylic acid (9-dichloromethylene-1,2,3,4-tetrahydro-1,4-methano-naphthalen-5-yl)-amide (disclosed in WO 2007/048556) + TX, 3-difluoromethyl-1-methyl-1H-
 10 pyrazole-4-carboxylic acid (3',4',5'-trifluoro-biphenyl-2-yl)-amide (disclosed in WO 2006/087343) + TX, [(3S,4R,4aR,6S,6aS,12R,12aS,12bS)-3-[(cyclopropylcarbonyl)oxy]-1,3,4,4a,5,6,6a,12,12a,12b-decahydro-6,12-dihydroxy-4,6a,12b-trimethyl-11-oxo-9-(3-pyridinyl)-2H,11Hnaphtho[2,1-b]pyrano[3,4-e]pyran-4-yl)methyl-cyclopropanecarboxylate [915972-17-7] + TX, 1,3,5-trimethyl-N-(2-methyl-1-oxopropyl)-N-[3-(2-methylpropyl)-4-[2,2,2-trifluoro-1-methoxy-1-(trifluoromethyl)ethyl]phenyl]-1H-pyrazole-
 15 4-carboxamide [926914-55-8] + TX; lancotrione [1486617-21-3] + TX; florypyrauxifen [943832-81-3] + TX; ipfentrifluconazole [1417782-08-1] + TX; mefentrifluconazole [1417782-03-6] + TX; quinofumelin [861647-84-9] + TX; chloroprallethrin [399572-87-3] + TX; cyhalodiamide [1262605-53-7] + TX; fluazaindolizine [1254304-22-7] + TX; fluxametamide [928783-29-3] + TX; epsilon-metofluthrin [240494-71-7] + TX; epsilon-momfluorothrin [1065124-65-3] + TX; pydiflumetofen [1228284-64-7] + TX; kappa-bifenthrin
 20 [439680-76-9] + TX; broflanilide [1207727-04-5] + TX; dicloromezotiaz [1263629-39-5] + TX; dipymetitrone [16114-35-5] + TX; pyraziflumid [942515-63-1] + TX; and kappa-tefluthrin [391634-71-2] + TX; and

microbials including: *Acinetobacter lwoffii* + TX, *Acremonium alternatum* + TX + TX, *Acremonium cephalosporium* + TX + TX, *Acremonium diospyri* + TX, *Acremonium obclavatum* + TX, *Adoxophyes*
 25 *orana granulovirus* (AdoxGV) (Capex®) + TX, *Agrobacterium radiobacter* strain K84 (Galltrol-A®) + TX, *Alternaria alternate* + TX, *Alternaria cassia* + TX, *Alternaria destruens* (Smolder®) + TX, *Ampelomyces quisqualis* (AQ10®) + TX, *Aspergillus flavus* AF36 (AF36®) + TX, *Aspergillus flavus* NRRL 21882 (Aflaguard®) + TX, *Aspergillus* spp. + TX, *Aureobasidium pullulans* + TX, *Azospirillum* + TX, (MicroAZ® + TX, TAZO B®) + TX, *Azotobacter* + TX, *Azotobacter chroococcum* (Azotomeal®) + TX, *Azotobacter* cysts
 30 (Bionatural Blooming Blossoms®) + TX, *Bacillus amyloliquefaciens* + TX, *Bacillus cereus* + TX, *Bacillus chitinosporus* strain CM-1 + TX, *Bacillus chitinosporus* strain AQ746 + TX, *Bacillus licheniformis* strain HB-2 (Biostart™ Rhizoboost®) + TX, *Bacillus licheniformis* strain 3086 (EcoGuard® + TX, Green Releaf®) + TX, *Bacillus circulans* + TX, *Bacillus firmus* (BioSafe® + TX, BioNem-WP® + TX, VOTIVO®) + TX, *Bacillus firmus* strain I-1582 + TX, *Bacillus macerans* + TX, *Bacillus marismortui* + TX, *Bacillus*
 35 *megaterium* + TX, *Bacillus mycoides* strain AQ726 + TX, *Bacillus papillae* (Milky Spore Powder®) + TX, *Bacillus pumilus* spp. + TX, *Bacillus pumilus* strain GB34 (Yield Shield®) + TX, *Bacillus pumilus* strain AQ717 + TX, *Bacillus pumilus* strain QST 2808 (Sonata® + TX, Ballad Plus®) + TX, *Bacillus spahericus* (VectoLex®) + TX, *Bacillus* spp. + TX, *Bacillus* spp. strain AQ175 + TX, *Bacillus* spp. strain AQ177 + TX, *Bacillus* spp. strain AQ178 + TX, *Bacillus subtilis* strain QST 713 (CEASE® + TX, Serenade® + TX,
 40 Rhapsody®) + TX, *Bacillus subtilis* strain QST 714 (JAZZ®) + TX, *Bacillus subtilis* strain AQ153 + TX, *Bacillus subtilis* strain AQ743 + TX, *Bacillus subtilis* strain QST3002 + TX, *Bacillus subtilis* strain QST3004 + TX, *Bacillus subtilis* var. *amyloliquefaciens* strain FZB24 (Taegro® + TX, Rhizopro®) + TX,

Bacillus thuringiensis Cry 2Ae + TX, *Bacillus thuringiensis* Cry1Ab + TX, *Bacillus thuringiensis* aizawai GC 91 (Agree®) + TX, *Bacillus thuringiensis israelensis* (BMP123® + TX, Aquabac® + TX, VectoBac®) + TX, *Bacillus thuringiensis kurstaki* (Javelin® + TX, Deliver® + TX, CryMax® + TX, Bonide® + TX, Scutella WP® + TX, Turilav WP® + TX, Astuto® + TX, Dipel WP® + TX, Biobit® + TX, Foray®) + TX, *Bacillus*
 5 *thuringiensis kurstaki* BMP 123 (Baritone®) + TX, *Bacillus thuringiensis kurstaki* HD-1 (Bioprotec-CAF / 3P®) + TX, *Bacillus thuringiensis* strain BD#32 + TX, *Bacillus thuringiensis* strain AQ52 + TX, *Bacillus thuringiensis* var. *aizawai* (XenTari® + TX, DiPel®) + TX, bacteria spp. (GROWMEND® + TX, GROWSWEET® + TX, Shootup®) + TX, bacteriophage of *Clavipacter michiganensis* (AgriPhage®) + TX, Bakflor® + TX, *Beauveria bassiana* (Beaugenic® + TX, Brocaril WP®) + TX, *Beauveria bassiana* GHA
 10 (Mycotrol ES® + TX, Mycotrol O® + TX, BotaniGuard®) + TX, *Beauveria brongniartii* (Engerlingspilz® + TX, Schweizer Beauveria® + TX, Melocont®) + TX, *Beauveria* spp. + TX, *Botrytis cineria* + TX, *Bradyrhizobium japonicum* (TerraMax®) + TX, *Brevibacillus brevis* + TX, *Bacillus thuringiensis tenebrionis* (Novodor®) + TX, BtBooster + TX, *Burkholderia cepacia* (Deny® + TX, Intercept® + TX, Blue Circle®) + TX, *Burkholderia gladii* + TX, *Burkholderia gladioli* + TX, *Burkholderia* spp. + TX, Canadian thistle fungus
 15 (CBH Canadian Bioherbicide®) + TX, *Candida butyri* + TX, *Candida famata* + TX, *Candida fructus* + TX, *Candida glabrata* + TX, *Candida guilliermondii* + TX, *Candida melibiosica* + TX, *Candida oleophila* strain O + TX, *Candida parapsilosis* + TX, *Candida pelliculosa* + TX, *Candida pulcherrima* + TX, *Candida reukaufii* + TX, *Candida saitoana* (Bio-Coat® + TX, Biocure®) + TX, *Candida sake* + TX, *Candida* spp. + TX, *Candida tenius* + TX, *Cedecea dravisae* + TX, *Cellulomonas flavigena* + TX, *Chaetomium cochliodes*
 20 (Nova-Cide®) + TX, *Chaetomium globosum* (Nova-Cide®) + TX, *Chromobacterium subtsugae* strain PRAA4-1T (Grandevo®) + TX, *Cladosporium cladosporioides* + TX, *Cladosporium oxysporum* + TX, *Cladosporium chlorocephalum* + TX, *Cladosporium* spp. + TX, *Cladosporium tenuissimum* + TX, *Clonostachys rosea* (EndoFine®) + TX, *Colletotrichum acutatum* + TX, *Coniothyrium minitans* (Cotans WG®) + TX, *Coniothyrium* spp. + TX, *Cryptococcus albidus* (YIELDPLUS®) + TX, *Cryptococcus*
 25 *humicola* + TX, *Cryptococcus infirmo-minutus* + TX, *Cryptococcus laurentii* + TX, *Cryptophlebia leucotreta granulovirus* (Cryptex®) + TX, *Cupriavidus campinensis* + TX, *Cydia pomonella granulovirus* (CYD-X®) + TX, *Cydia pomonella granulovirus* (Madex® + TX, Madex Plus® + TX, Madex Max/ Carpovirusine®) + TX, *Cylindrobasidium laeve* (Stumpout®) + TX, *Cylindrocladium* + TX, *Debaryomyces hansenii* + TX, *Drechslera hawaiiensis* + TX, *Enterobacter cloacae* + TX, *Enterobacteriaceae* + TX,
 30 *Entomophthora virulenta* (Vektor®) + TX, *Epicoccum nigrum* + TX, *Epicoccum purpurascens* + TX, *Epicoccum* spp. + TX, *Filobasidium floriforme* + TX, *Fusarium acuminatum* + TX, *Fusarium chlamydosporum* + TX, *Fusarium oxysporum* (Fusaclean® / Biofox C®) + TX, *Fusarium proliferatum* + TX, *Fusarium* spp. + TX, *Galactomyces geotrichum* + TX, *Gliocladium catenulatum* (Primastop® + TX, Prestop®) + TX, *Gliocladium roseum* + TX, *Gliocladium* spp. (SoilGard®) + TX, *Gliocladium virens*
 35 (Soilgard®) + TX, *Granulovirus* (Granupom®) + TX, *Halobacillus halophilus* + TX, *Halobacillus litoralis* + TX, *Halobacillus trueperi* + TX, *Halomonas* spp. + TX, *Halomonas subglaciescola* + TX, *Halovibrio variabilis* + TX, *Hanseniaspora uvarum* + TX, *Helicoverpa armigera nucleopolyhedrovirus* (Helicovex®) + TX, *Helicoverpa zea nuclear polyhedrosis virus* (Gemstar®) + TX, Isoflavone – formononetin (Myconate®) + TX, *Kloeckera apiculata* + TX, *Kloeckera* spp. + TX, *Lagenidium giganteum* (Laginex®) +
 40 TX, *Lecanicillium longisporum* (Vertiblast®) + TX, *Lecanicillium muscarium* (Vertikil®) + TX, *Lymantria Dispar nucleopolyhedrosis virus* (Disparvirus®) + TX, *Marinococcus halophilus* + TX, *Meira geulakonigii* + TX, *Metarhizium anisopliae* (Met52®) + TX, *Metarhizium anisopliae* (Destruxin WP®) + TX,

- Metschnikowia fruticola* (Shemer®) + TX, *Metschnikowia pulcherrima* + TX, *Microdochium dimerum* (Antibot®) + TX, *Micromonospora coerulea* + TX, *Microsphaeropsis ochracea* + TX, *Muscudor albus* 620 (Muscudor®) + TX, *Muscudor roseus* strain A3-5 + TX, *Mycorrhizae* spp. (AMykor®) + TX, Root Maximizer®) + TX, *Myrothecium verrucaria* strain AARC-0255 (DiTera®) + TX, BROS PLUS® + TX,
- 5 *Ophiostoma piliferum* strain D97 (Sylvanex®) + TX, *Paecilomyces farinosus* + TX, *Paecilomyces fumosoroseus* (PFR-97® + TX, PreFeRal®) + TX, *Paecilomyces linacinus* (Biostat WP®) + TX, *Paecilomyces lilacinus* strain 251 (MeloCon WG®) + TX, *Paenibacillus polymyxa* + TX, *Pantoea agglomerans* (BlightBan C9-1®) + TX, *Pantoea* spp. + TX, *Pasteuria* spp. (Econem®) + TX, *Pasteuria nishizawae* + TX, *Penicillium aurantiogriseum* + TX, *Penicillium billai* (Jumpstart® + TX, TagTeam®) +
- 10 TX, *Penicillium brevicompactum* + TX, *Penicillium frequentans* + TX, *Penicillium griseofulvum* + TX, *Penicillium purpurogenum* + TX, *Penicillium* spp. + TX, *Penicillium viridicatum* + TX, *Phlebiopsis gigantea* (Rotstop®) + TX, phosphate solubilizing bacteria (Phosphomeal®) + TX, *Phytophthora cryptogea* + TX, *Phytophthora palmivora* (Devine®) + TX, *Pichia anomala* + TX, *Pichia guillemontii* + TX, *Pichia membranaefaciens* + TX, *Pichia onychis* + TX, *Pichia stipites* + TX, *Pseudomonas aeruginosa* +
- 15 TX, *Pseudomonas aureofaciens* (Spot-Less Biofungicide®) + TX, *Pseudomonas cepacia* + TX, *Pseudomonas chlororaphis* (AtEze®) + TX, *Pseudomonas corrugate* + TX, *Pseudomonas fluorescens* strain A506 (BlightBan A506®) + TX, *Pseudomonas putida* + TX, *Pseudomonas reactans* + TX, *Pseudomonas* spp. + TX, *Pseudomonas syringae* (Bio-Save®) + TX, *Pseudomonas viridiflava* + TX, *Pseudomonas fluorescens* (Zequanox®) + TX, *Pseudozyma flocculosa* strain PF-A22 UL (Sporodex L®) +
- 20 TX, *Puccinia canaliculata* + TX, *Puccinia thlaspeos* (Wood Warrior®) + TX, *Pythium paroecandrum* + TX, *Pythium oligandrum* (Polygandron® + TX, Polyversum®) + TX, *Pythium periplocum* + TX, *Rhanella aquatilis* + TX, *Rhanella* spp. + TX, *Rhizobia* (Dormal® + TX, Vault®) + TX, *Rhizoctonia* + TX, *Rhodococcus globerulus* strain AQ719 + TX, *Rhodospiridium diobovatum* + TX, *Rhodospiridium toruloides* + TX, *Rhodotorula* spp. + TX, *Rhodotorula glutinis* + TX, *Rhodotorula graminis* + TX,
- 25 *Rhodotorula mucilagnosa* + TX, *Rhodotorula rubra* + TX, *Saccharomyces cerevisiae* + TX, *Salinococcus roseus* + TX, *Sclerotinia minor* + TX, *Sclerotinia minor* (SARRITOR®) + TX, *Scytalidium* spp. + TX, *Scytalidium uredinicola* + TX, *Spodoptera exigua nuclear polyhedrosis virus* (Spod-X® + TX, Spexit®) + TX, *Serratia marcescens* + TX, *Serratia plymuthica* + TX, *Serratia* spp. + TX, *Sordaria fimicola* + TX, *Spodoptera littoralis nucleopolyhedrovirus* (Littovir®) + TX, *Sporobolomyces roseus* + TX,
- 30 *Stenotrophomonas maltophilia* + TX, *Streptomyces ahngroscopicus* + TX, *Streptomyces albaduncus* + TX, *Streptomyces exfoliates* + TX, *Streptomyces galbus* + TX, *Streptomyces griseoplanus* + TX, *Streptomyces griseoviridis* (Mycostop®) + TX, *Streptomyces lydicus* (Actinovate®) + TX, *Streptomyces lydicus* WYEC-108 (ActinoGrow®) + TX, *Streptomyces violaceus* + TX, *Tilletiopsis minor* + TX, *Tilletiopsis* spp. + TX, *Trichoderma asperellum* (T34 Biocontrol®) + TX, *Trichoderma gamsii* (Tenet®) +
- 35 TX, *Trichoderma atroviride* (Plantmate®) + TX, *Trichoderma hamatum* TH 382 + TX, *Trichoderma harzianum rifai* (Mycostar®) + TX, *Trichoderma harzianum* T-22 (Trianum-P® + TX, PlantShield HC® + TX, RootShield® + TX, Trianum-G®) + TX, *Trichoderma harzianum* T-39 (Trichodex®) + TX, *Trichoderma inhamatum* + TX, *Trichoderma koningii* + TX, *Trichoderma* spp. LC 52 (Sentinel®) + TX, *Trichoderma lignorum* + TX, *Trichoderma longibrachiatum* + TX, *Trichoderma polysporum* (Binab T®) +
- 40 TX, *Trichoderma taxi* + TX, *Trichoderma virens* + TX, *Trichoderma virens* (formerly *Gliocladium virens* GL-21) (SoilGuard®) + TX, *Trichoderma viride* + TX, *Trichoderma viride* strain ICC 080 (Remedier®) + TX, *Trichosporon pullulans* + TX, *Trichosporon* spp. + TX, *Trichothecium* spp. + TX, *Trichothecium*

roseum + TX, *Typhula phacorrhiza* strain 94670 + TX, *Typhula phacorrhiza* strain 94671 + TX, *Ulocladium atrum* + TX, *Ulocladium oudemansii* (Botry-Zen®) + TX, *Ustilago maydis* + TX, various bacteria and supplementary micronutrients (Natural II®) + TX, various fungi (Millennium Microbes®) + TX, *Verticillium chlamydosporium* + TX, *Verticillium lecanii* (Mycotal® + TX, Vertalec®) + TX, Vip3Aa20
 5 (VIPtera®) + TX, *Virgibacillus marismortui* + TX, *Xanthomonas campestris* pv. *Poae* (Camperico®) + TX, *Xenorhabdus bovienii* + TX, *Xenorhabdus nematophilus*; and

Plant extracts including: pine oil (Retenol®) + TX, azadirachtin (Plasma Neem Oil® + TX, AzaGuard® + TX, MeemAzal® + TX, Molt-X® + TX, Botanical IGR (Neemazad® + TX, Neemix®) + TX, canola oil (Lilly Miller Vegol®) + TX, *Chenopodium ambrosioides* near *ambrosioides* (Requiem®) + TX,
 10 *Chrysanthemum* extract (Crisant®) + TX, extract of neem oil (Trilogy®) + TX, essentials oils of *Labiatae* (Botania®) + TX, extracts of clove rosemary peppermint and thyme oil (Garden insect killer®) + TX, Glycinebetaine (Greenstim®) + TX, garlic + TX, lemongrass oil (GreenMatch®) + TX, neem oil + TX, *Nepeta cataria* (Catnip oil) + TX, *Nepeta catarina* + TX, nicotine + TX, oregano oil (MossBuster®) + TX, *Pedaliaceae* oil (Nematon®) + TX, pyrethrum + TX, *Quillaja saponaria* (NemaQ®) + TX, *Reynoutria*
 15 *sachalinensis* (Regalia® + TX, Sakalia®) + TX, rotenone (Eco Roten®) + TX, *Rutaceae* plant extract (Soleo®) + TX, soybean oil (Ortho ecosense®) + TX, tea tree oil (Timorex Gold®) + TX, thymus oil + TX, AGNIQUE® MMF + TX, BugOil® + TX, mixture of rosemary sesame peppermint thyme and cinnamon extracts (EF 300®) + TX, mixture of clove rosemary and peppermint extract (EF 400®) + TX, mixture of clove peppermint garlic oil and mint (Soil Shot®) + TX, kaolin (Screen®) + TX, storage glucan of brown
 20 algae (Laminarin®); and

pheromones including: blackheaded fireworm pheromone (3M Sprayable Blackheaded Fireworm Pheromone®) + TX, Codling Moth Pheromone (Paramount dispenser-(CM)/ Isomate C-Plus®) + TX, Grape Berry Moth Pheromone (3M MEC-GBM Sprayable Pheromone®) + TX, Leafroller pheromone (3M MEC – LR Sprayable Pheromone®) + TX, Muscamone (Snip7 Fly Bait® + TX, Starbar Premium Fly
 25 Bait®) + TX, Oriental Fruit Moth Pheromone (3M oriental fruit moth sprayable pheromone®) + TX, Peachtree Borer Pheromone (Isomate-P®) + TX, Tomato Pinworm Pheromone (3M Sprayable pheromone®) + TX, Entostat powder (extract from palm tree) (Exosex CM®) + TX, (E + TX,Z + TX,Z)-3 + TX,8 + TX,11 Tetradecatrienyl acetate + TX, (Z + TX,Z + TX,E)-7 + TX,11 + TX,13-Hexadecatrienal + TX, (E + TX,Z)-7 + TX,9-Dodecadien-1-yl acetate + TX, 2-Methyl-1-butanol + TX, Calcium acetate + TX,
 30 Scenturion® + TX, Biolure® + TX, Check-Mate® + TX, Lavandulyl senecioate; and

Macrobiales including: *Aphelinus abdominalis* + TX, *Aphidius ervi* (Aphelinus-System®) + TX, *Acerophagus papaya* + TX, *Adalia bipunctata* (Adalia-System®) + TX, *Adalia bipunctata* (Adaline®) + TX, *Adalia bipunctata* (Aphidalia®) + TX, *Ageniaspis citricola* + TX, *Ageniaspis fuscicollis* + TX, *Amblyseius andersoni* (Anderline® + TX, Andersoni-System®) + TX, *Amblyseius californicus* (Amblyline® + TX,
 35 Spical®) + TX, *Amblyseius cucumeris* (Thripex® + TX, Bugline cucumeris®) + TX, *Amblyseius fallacis* (Fallacis®) + TX, *Amblyseius swirskii* (Bugline swirskii® + TX, Swirskii-Mite®) + TX, *Amblyseius womersleyi* (WomerMite®) + TX, *Amitus hesperidum* + TX, *Anagrus atomus* + TX, *Anagrus fusciventris* + TX, *Anagrus kamali* + TX, *Anagrus loecki* + TX, *Anagrus pseudococci* (Citripar®) + TX, *Anicetus benefices* + TX, *Anisopteromalus calandrae* + TX, *Anthocoris nemoralis* (Anthocoris-System®) + TX,
 40 *Aphelinus abdominalis* (Apheline® + TX, Aphiline®) + TX, *Aphelinus asychis* + TX, *Aphidius colemani* (Aphipar®) + TX, *Aphidius ervi* (Ervipar®) + TX, *Aphidius gifuensis* + TX, *Aphidius matricariae* (Aphipar-M®) + TX, *Aphidoletes aphidimyza* (Aphidend®) + TX, *Aphidoletes aphidimyza* (Aphidoline®) + TX,

- Aphytis lingnanensis* + TX, *Aphytis melinus* + TX, *Aprostocetus hagenowii* + TX, *Atheta coriaria* (Staphyline®) + TX, *Bombus* spp. + TX, *Bombus terrestris* (Natupol Beehive®) + TX, *Bombus terrestris* (Beeline® + TX, Tripol®) + TX, *Cephalonomia stephanoderis* + TX, *Chilocorus nigritus* + TX, *Chrysoperla carnea* (Chrysoline®) + TX, *Chrysoperla carnea* (Chrysopa®) + TX, *Chrysoperla rufilabris* + TX,
- 5 *Cirrospilus ingenuus* + TX, *Cirrospilus quadristriatus* + TX, *Citrostichus phyllocnistoides* + TX, *Closterocerus chamaeleon* + TX, *Closterocerus* spp. + TX, *Coccidoxenoides perminutus* (Planopar®) + TX, *Coccophagus cowperi* + TX, *Coccophagus lycimnia* + TX, *Cotesia flavipes* + TX, *Cotesia plutellae* + TX, *Cryptolaemus montrouzieri* (Cryptobug® + TX, Cryptoline®) + TX, *Cybocephalus nipponicus* + TX, *Dacnusa sibirica* + TX, *Dacnusa sibirica* (Minusa®) + TX, *Diglyphus isaea* (Diminex®) + TX, *Delphastus*
- 10 *catalinae* (Delphastus®) + TX, *Delphastus pusillus* + TX, *Diachasmimorpha krausii* + TX, *Diachasmimorpha longicaudata* + TX, *Diaparsis jucunda* + TX, *Diaphorencyrtus aligarhensis* + TX, *Diglyphus isaea* + TX, *Diglyphus isaea* (Miglyphus® + TX, Digline®) + TX, *Dacnusa sibirica* (DacDigline® + TX, Minex®) + TX, *Diversinervus* spp. + TX, *Encarsia citrina* + TX, *Encarsia formosa* (Encarsia max® + TX, Encarline® + TX, En-Strip®) + TX, *Eretmoceris eremicus* (Enermix®) + TX, *Encarsia guadeloupae* +
- 15 TX, *Encarsia haitiensis* + TX, *Episyrphus balteatus* (Syrphidend®) + TX, *Eretmoceris siphonini* + TX, *Eretmoceris californicus* + TX, *Eretmoceris eremicus* (Ercal® + TX, Eretline e®) + TX, *Eretmoceris eremicus* (Bemimix®) + TX, *Eretmoceris hayati* + TX, *Eretmoceris mundus* (Bemipar® + TX, Eretline m®) + TX, *Eretmoceris siphonini* + TX, *Exochomus quadripustulatus* + TX, *Feltiella acarisuga* (Spidend®) + TX, *Feltiella acarisuga* (Feltiline®) + TX, *Fopius arisanus* + TX, *Fopius ceratitivorus* + TX,
- 20 Formononetin (Wirless Beehome®) + TX, *Franklinothrips vespiformis* (Vespop®) + TX, *Galendromus occidentalis* + TX, *Goniozus legneri* + TX, *Habrobracon hebetor* + TX, *Harmonia axyridis* (HarmoBeetle®) + TX, *Heterorhabditis* spp. (Lawn Patrol®) + TX, *Heterorhabditis bacteriophora* (NemaShield HB® + TX, Nemaseek® + TX, Terranem-Nam® + TX, Terranem® + TX, Larvanem® + TX, B-Green® + TX, NemAttack® + TX, Nematop®) + TX, *Heterorhabditis megidis* (Nemasys H® + TX, BioNem H® + TX,
- 25 Exhibitline hm® + TX, Larvanem-M®) + TX, *Hippodamia convergens* + TX, *Hypoaspis aculeifer* (Aculeifer-System® + TX, Entomite-A®) + TX, *Hypoaspis miles* (Hypoline m® + TX, Entomite-M®) + TX, *Lbalia leucospoides* + TX, *Lecanoideus floccissimus* + TX, *Lemophagus errabundus* + TX, *Leptomastidea abnormis* + TX, *Leptomastix dactylopii* (Leptopar®) + TX, *Leptomastix epona* + TX, *Lindorus lophanthae* + TX, *Lipolexis oregmae* + TX, *Lucilia caesar* (NatuFly®) + TX, *Lysiphlebus testaceipes* + TX,
- 30 *Macrolophus caliginosus* (Mirical-N® + TX, Macroline c® + TX, Mirical®) + TX, *Mesoseiulus longipes* + TX, *Metaphycus flavus* + TX, *Metaphycus lounsburyi* + TX, *Micromus angulatus* (Milacewing®) + TX, *Microterys flavus* + TX, *Muscidifurax raptorellus* and *Spalangia cameroni* (Biopar®) + TX, *Neodryinus typhlocybae* + TX, *Neoseiulus californicus* + TX, *Neoseiulus cucumeris* (THRYPEX®) + TX, *Neoseiulus fallacis* + TX, *Nesideocoris tenuis* (NesidioBug® + TX, Nesibug®) + TX, *Ophyra aenescens* (Biofly®) +
- 35 TX, *Orius insidiosus* (Thrip-or-I® + TX, Oriline i®) + TX, *Orius laevigatus* (Thrip-or-L® + TX, Oriline l®) + TX, *Orius majusculus* (Oriline m®) + TX, *Orius strigicollis* (Thrip-or-S®) + TX, *Pauesia juniperorum* + TX, *Pediobius foveolatus* + TX, *Phasmarhabditis hermaphrodita* (Nemaslug®) + TX, *Phymastichus coffea* + TX, *Phytoseiulus macropilus* + TX, *Phytoseiulus persimilis* (Spidex® + TX, Phytoline p®) + TX, *Podisus maculiventris* (Podisus®) + TX, *Pseudacteon curvatus* + TX, *Pseudacteon obtusus* + TX, *Pseudacteon*
- 40 *tricusps* + TX, *Pseudaphycus maculipennis* + TX, *Pseudleptomastix mexicana* + TX, *Psyllaephagus pilosus* + TX, *Psytalia concolor* (complex) + TX, *Quadrastichus* spp. + TX, *Rhyzobius lophanthae* + TX, *Rodolia cardinalis* + TX, *Rumina decollate* + TX, *Semiellacher petiolatus* + TX, *Sitobion avenae*

(Ervibank®) + TX, *Steinernema carpocapsae* (Nematac C® + TX, Millenium® + TX, BioNem C® + TX, NemAttack® + TX, Nemastar® + TX, Capsanem®) + TX, *Steinernema feltiae* (NemaShield® + TX, Nemasys F® + TX, BioNem F® + TX, Steinernema-System® + TX, NemAttack® + TX, Nemaplus® + TX, Exhibitline sf® + TX, Scia-rid® + TX, Entonem®) + TX, *Steinernema kraussei* (Nemasys L® + TX, BioNem L® + TX, Exhibitline srb®) + TX, *Steinernema riobrave* (BioVector® + TX, BioVektor®) + TX, *Steinernema scapterisci* (Nematac S®) + TX, *Steinernema* spp. + TX, *Steinernematid* spp. (Guardian Nematodes®) + TX, *Stethorus punctillum* (Stethorus®) + TX, *Tamarixia radiata* + TX, *Tetrastichus setifer* + TX, *Thripobius semiluteus* + TX, *Torymus sinensis* + TX, *Trichogramma brassicae* (Tricholine b®) + TX, *Trichogramma brassicae* (Tricho-Strip®) + TX, *Trichogramma evanescens* + TX, *Trichogramma minutum* + TX, *Trichogramma ostrinae* + TX, *Trichogramma platneri* + TX, *Trichogramma pretiosum* + TX, *Xanthopimpla stemmator*, and

other biologicals including: abscisic acid + TX, bioSea® + TX, *Chondrostereum purpureum* (Chontrol Paste®) + TX, *Colletotrichum gloeosporioides* (Collego®) + TX, Copper Octanoate (Cueva®) + TX, Delta traps (Trapline d®) + TX, *Erwinia amylovora* (Harpin) (ProAct® + TX, Ni-HIBIT Gold CST®) + TX, Ferri-phosphate (Ferramol®) + TX, Funnel traps (Trapline y®) + TX, Gallex® + TX, Grower's Secret® + TX, Homo-brassonolide + TX, Iron Phosphate (Lilly Miller Worry Free Ferramol Slug & Snail Bait®) + TX, MCP hail trap (Trapline f®) + TX, *Microctonus hyperodae* + TX, *Mycoleptodiscus terrestris* (Des-X®) + TX, BioGain® + TX, Aminomite® + TX, Zenox® + TX, Pheromone trap (Thripline ams®) + TX, potassium bicarbonate (MilStop®) + TX, potassium salts of fatty acids (Sanova®) + TX, potassium silicate solution (Sil-Matrix®) + TX, potassium iodide + potassiumthiocyanate (Enzicur®) + TX, SuffOil-X® + TX, Spider venom + TX, *Nosema locustae* (Semaspore Organic Grasshopper Control®) + TX, Sticky traps (Trapline YF® + TX, Rebell Amarillo®) + TX and Traps (Takitraps y + b®) + TX.

The references in brackets behind the active ingredients, e.g. [3878-19-1] refer to the Chemical Abstracts Registry number. The above described mixing partners are known. Where the active ingredients are included in "The Pesticide Manual" [The Pesticide Manual - A World Compendium; Thirteenth Edition; Editor: C. D. S. Tomlin; The British Crop Protection Council], they are described therein under the entry number given in round brackets hereinabove for the particular compound; for example, the compound "abamectin" is described under entry number (1). Where "[CCN]" is added hereinabove to the particular compound, the compound in question is included in the "Compendium of Pesticide Common Names", which is accessible on the internet [A. Wood; Compendium of Pesticide Common Names, Copyright © 1995-2004]; for example, the compound "acetoprole" is described under the internet address <http://www.alanwood.net/pesticides/acetoprole.html>.

Most of the active ingredients described above are referred to hereinabove by a so-called "common name", the relevant "ISO common name" or another "common name" being used in individual cases. If the designation is not a "common name", the nature of the designation used instead is given in round brackets for the particular compound; in that case, the IUPAC name, the IUPAC/Chemical Abstracts name, a "chemical name", a "traditional name", a "compound name" or a "development code" is used or, if neither one of those designations nor a "common name" is used, an "alternative name" is employed. "CAS Reg. No" means the Chemical Abstracts Registry Number.

The active ingredient mixture of the compounds of formula (I) selected from Tables 1 to 96 and Table A of the present invention with active ingredients described above comprises a compound selected from Tables 1 to 96 and Table A and an active ingredient as described above preferably in a mixing ratio

of from 100:1 to 1:6000, especially from 50:1 to 1:50, more especially in a ratio of from 20:1 to 1:20, even more especially from 10:1 to 1:10, very especially from 5:1 and 1:5, special preference being given to a ratio of from 2:1 to 1:2, and a ratio of from 4:1 to 2:1 being likewise preferred, above all in a ratio of 1:1, or 5:1, or 5:2, or 5:3, or 5:4, or 4:1, or 4:2, or 4:3, or 3:1, or 3:2, or 2:1, or 1:5, or 2:5, or 3:5, or 4:5, or 1:4, or 2:4, or 3:4, or 1:3, or 2:3, or 1:2, or 1:600, or 1:300, or 1:150, or 1:35, or 2:35, or 4:35, or 1:75, or 2:75, or 4:75, or 1:6000, or 1:3000, or 1:1500, or 1:350, or 2:350, or 4:350, or 1:750, or 2:750, or 4:750. Those mixing ratios are by weight.

The mixtures as described above can be used in a method for controlling pests, which comprises applying a composition comprising a mixture as described above to the pests or their environment, with the exception of a method for treatment of the human or animal body by surgery or therapy and diagnostic methods practised on the human or animal body.

The mixtures comprising a compound of formula (I) selected from Tables 1 to 96 and Table P and one or more active ingredients as described above can be applied, for example, in a single "ready-mix" form, in a combined spray mixture composed from separate formulations of the single active ingredient components, such as a "tank-mix", and in a combined use of the single active ingredients when applied in a sequential manner, i.e. one after the other with a reasonably short period, such as a few hours or days. The order of applying the compounds of formula (I) selected from Tables 1 to 96 and Table P and the active ingredients as described above is not essential for working the present invention.

The compositions according to the invention can also comprise further solid or liquid auxiliaries, such as stabilizers, for example unepoxidized or epoxidized vegetable oils (for example epoxidized coconut oil, rapeseed oil or soya oil), antifoams, for example silicone oil, preservatives, viscosity regulators, binders and/or tackifiers, fertilizers or other active ingredients for achieving specific effects, for example bactericides, fungicides, nematocides, plant activators, molluscicides or herbicides.

The compositions according to the invention are prepared in a manner known per se, in the absence of auxiliaries for example by grinding, screening and/or compressing a solid active ingredient and in the presence of at least one auxiliary for example by intimately mixing and/or grinding the active ingredient with the auxiliary (auxiliaries). These processes for the preparation of the compositions and the use of the compounds I for the preparation of these compositions are also a subject of the invention.

The application methods for the compositions, that is the methods of controlling pests of the abovementioned type, such as spraying, atomizing, dusting, brushing on, dressing, scattering or pouring - which are to be selected to suit the intended aims of the prevailing circumstances - and the use of the compositions for controlling pests of the abovementioned type are other subjects of the invention. Typical rates of concentration are between 0.1 and 1000 ppm, preferably between 0.1 and 500 ppm, of active ingredient. The rate of application per hectare is generally 1 to 2000 g of active ingredient per hectare, in particular 10 to 1000 g/ha, preferably 10 to 600 g/ha.

A preferred method of application in the field of crop protection is application to the foliage of the plants (foliar application), it being possible to select frequency and rate of application to match the danger of infestation with the pest in question. Alternatively, the active ingredient can reach the plants via the root system (systemic action), by drenching the locus of the plants with a liquid composition or by incorporating the active ingredient in solid form into the locus of the plants, for example into the soil, for example in the form of granules (soil application). In the case of paddy rice crops, such granules can be metered into the flooded paddy-field.

The compounds of the invention and compositions thereof are also be suitable for the protection of plant propagation material, for example seeds, such as fruit, tubers or kernels, or nursery plants, against pests of the abovementioned type. The propagation material can be treated with the compound prior to planting, for example seed can be treated prior to sowing. Alternatively, the compound can be applied to
5 seed kernels (coating), either by soaking the kernels in a liquid composition or by applying a layer of a solid composition. It is also possible to apply the compositions when the propagation material is planted to the site of application, for example into the seed furrow during drilling. These treatment methods for plant propagation material and the plant propagation material thus treated are further subjects of the invention. Typical treatment rates would depend on the plant and pest/fungi to be controlled and are
10 generally between 1 to 200 grams per 100 kg of seeds, preferably between 5 to 150 grams per 100 kg of seeds, such as between 10 to 100 grams per 100 kg of seeds.

The term seed embraces seeds and plant propagules of all kinds including but not limited to true seeds, seed pieces, suckers, corns, bulbs, fruit, tubers, grains, rhizomes, cuttings, cut shoots and the like and means in a preferred embodiment true seeds.

15 The present invention also comprises seeds coated or treated with or containing a compound of formula (I). The term "coated or treated with and/or containing" generally signifies that the active ingredient is for the most part on the surface of the seed at the time of application, although a greater or lesser part of the ingredient may penetrate into the seed material, depending on the method of application. When the said seed product is (re)planted, it may absorb the active ingredient. In an
20 embodiment, the present invention makes available a plant propagation material adhered thereto with a compound of formula (I). Further, it is hereby made available, a composition comprising a plant propagation material treated with a compound of formula (I).

Seed treatment comprises all suitable seed treatment techniques known in the art, such as seed dressing, seed coating, seed dusting, seed soaking and seed pelleting. The seed treatment application of
25 the compound formula (I) can be carried out by any known methods, such as spraying or by dusting the seeds before sowing or during the sowing/planting of the seeds.

The invention further relates to a method for controlling pests, which comprises applying a composition according to the invention to the pests or their environment preferably with the exception of a method for treatment of the human or animal body by surgery or therapy and diagnostic methods
30 practised on the human or animal body.

The invention further relates to a method for the protection of plant propagation material from the attack by pests, which comprises treating the propagation material or the site, where the propagation material is planted, with a composition comprising a compound according to this invention or with a compound according to this invention. The invention further relates to a plant propagation material treated
35 with the pesticidal composition comprising a compound according to this invention or with a compound according to this invention.

The Examples which follow serve to illustrate the invention. The compounds of the invention can be distinguished from known compounds by virtue of greater efficacy at low application rates, which can be verified by the person skilled in the art using the experimental procedures outlined in the Examples,
40 using lower application rates if necessary, for example 50 ppm, 12.5 ppm, 6 ppm, 3 ppm, 1.5 ppm or 0.8 ppm. Temperatures are given in degrees Celsius; mixing ratios of solvents are given in parts by volume.

Preparatory Examples:

"Mp" means melting point in °C. Free radicals represent methyl groups. ¹H NMR measurements were recorded on a Bruker 400MHz spectrometer, chemical shifts are given in ppm relevant to a TMS standard. Spectra measured in deuterated solvents as indicated.

The following LC-MS methods were used to characterize the compounds:

5

Method A

MS	Spectra were recorded on a Mass Spectrometer from Waters (SQD or SQDII Single quadrupole mass spectrometer) equipped with an electrospray source (Polarity: positive or negative ions, Capillary: 3.00 kV, Cone range: 30 V, Extractor: 2.00 V, Source Temperature: 150°C, Desolvation Temperature: 350°C, Cone Gas Flow: 0 L/Hr, Desolvation Gas Flow: 650 L/Hr, Mass range: 100 to 900 Da).
LC	Acquity UPLC from Waters: Binary pump, heated column compartment and diode-array detector. Solvent degasser, binary pump, heated column compartment and diode-array detector. Column: Waters UPLC HSS T3 , 1.8 µm, 30 x 2.1 mm, Temp: 60 °C, DAD Wavelength range (nm): 210 to 500, Solvent Gradient: A = water + 5% MeOH + 0.05 % HCOOH, B= Acetonitrile + 0.05 % HCOOH: gradient: 0 min 0% B, 100%A; 1.2-1.5min 100% B; Flow (ml/min) 0.85.

Method B

MS	Spectra were recorded on a Mass Spectrometer from Waters (SQD or SQDII Single quadrupole mass spectrometer) equipped with an electrospray source (Polarity: positive or negative ions, Capillary: 3.00 kV, Cone range: 45 V, Extractor: 2.00 V, Source Temperature: 150°C, Desolvation Temperature: 350°C, Cone Gas Flow: 0 L/Hr, Desolvation Gas Flow: 650 L/Hr, Mass range: 100 to 900 Da).
LC	Acquity UPLC from Waters: Binary pump, heated column compartment and diode-array detector. Solvent degasser, binary pump, heated column compartment and diode-array detector. Column: Waters UPLC HSS T3 , 1.8 µm, 30 x 2.1 mm, Temp: 60 °C, DAD Wavelength range (nm): 210 to 500, Solvent Gradient: A = water + 5% MeOH + 0.05 % HCOOH, B= Acetonitrile + 0.05 % HCOOH: gradient: 0 min 0% B, 100%A; 1.2-1.5min 100% B; Flow (ml/min) 0.85.

10 Method C

MS	Spectra were recorded on a Mass Spectrometer from Waters (SQD, SQDII Single quadrupole mass spectrometer) equipped with an electrospray source (Polarity: positive and negative ions), Capillary: 3.00 kV, Cone range: 30V, Extractor: 2.00 V, Source Temperature: 150°C, Desolvation Temperature: 350°C, Cone Gas Flow: 50 l/h, Desolvation Gas Flow: 650 l/h, Mass range: 100 to 900 Da)
LC	Acquity UPLC from Waters: Binary pump, heated column compartment, diode-array detector and ELSD detector. Column: Waters UPLC HSS T3, 1.8 µm, 30 x 2.1 mm, Temp: 60 °C, DAD Wavelength range (nm): 210 to 500, Solvent Gradient: A = water + 5% MeOH + 0.05 % HCOOH, B= Acetonitrile + 0.05 % HCOOH, gradient: 10-100% B


	in 2.7 min; Flow (ml/min) 0.85
--	--------------------------------

Method D

MS	Spectra were recorded on a Waters SQD2 Mass Spectrometer (Single quadrupole mass spectrometer) Ionisation method: Electrospray, Polarity: positive ions Capillary (kV) 3.50, Cone (V) 30.00, Extractor (V) 3.00, Source Temperature (°C) 150, Desolvation Temperature (°C) 400 Cone Gas Flow (L/Hr) 60, Desolvation Gas Flow (L/Hr) 700, Mass range: 140 to 800 Da; DAD Wavelength range (nm): 210 to 400																				
LC	Method Waters ACQUITY UPLC with the following HPLC gradient conditions (Solvent A: Water/Methanol 9:1,0.1% formic acid and Solvent B: Acetonitrile,0.1% formic acid) <table><tr><td>Time (minutes)</td><td>A (%)</td><td>B (%)</td><td>Flow rate (ml/min)</td></tr><tr><td>0</td><td>100</td><td>0</td><td>0.75</td></tr><tr><td>2.5</td><td>0</td><td>100</td><td>0.75</td></tr><tr><td>2.8</td><td>0</td><td>100</td><td>0.75</td></tr><tr><td>3.0</td><td>100</td><td>0</td><td>0.75</td></tr></table> Type of column: Waters ACQUITY UPLC HSS T3; Column length: 30 mm; Internal diameter of column: 2.1 mm; Particle Size: 1.8 micron; Temperature: 60°C.	Time (minutes)	A (%)	B (%)	Flow rate (ml/min)	0	100	0	0.75	2.5	0	100	0.75	2.8	0	100	0.75	3.0	100	0	0.75
Time (minutes)	A (%)	B (%)	Flow rate (ml/min)																		
0	100	0	0.75																		
2.5	0	100	0.75																		
2.8	0	100	0.75																		
3.0	100	0	0.75																		

Example 1: N-(2-cyclobutylsulfonylisoxazolidin-4-yl)-4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide (example A014 in table A)

Step A: tert-butyl 4-[[4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate



10

A solution of tert-butyl 4-aminoisoxazolidine-2-carboxylate (preparation described in WO11/067272) (0.377 g) in tetrahydrofuran (5 ml) was treated with triethylamine (0.307 g) and stirred at 20°C. An amount of 4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl chloride (0.911 g) (the preparation of 4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid is described in WO13/026695) dissolved in tetrahydrofuran (5 ml) was slowly added.

The mixture was stirred for one hour then it was diluted with dichloromethane and aqueous solution of sodium bicarbonate. The phases were separated and the aqueous layer was extracted with dichloromethane. The organic phases were combined, dried over sodium sulfate and evaporated to yield a crude product as a yellow solid. Flash chromatography over silica gel with ethyl acetate-heptane mixture (1:1) gave the title compound as a slightly yellow solid. ¹H-NMR (CDCl₃, 400 MHz, δ in ppm):

20

7.58 (d, 2H), 7.54 – 7.46 (m, 2H), 7.40 (d, 1H), 6.25 (d, 1H), 5.02 (m, 1H), 4.12 – 3.95 (m, 4H), 3.72 – 3.57 (m, 2H), 2.47 (s, 3H), 1.49 (s, 9H).

Step B: 4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-isoxazolidin-2-ium-4-yl-

5 2-methyl-benzamide; 2,2,2-trifluoroacetate




A solution of tert-butyl 4-[4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate (1.00 g) in dichloromethane (8.5 ml) was treated with trifluoroacetic acid (0.945 g) and stirred at 20°C for 16 hours. The reaction was followed by TLC analysis
10 on silica gel, eluting with methyl alcohol-ethyl acetate (5/95).

The reaction mixture was concentrated under reduced pressure and the residue was submitted to flash chromatography over silica gel, eluting with methyl alcohol-ethyl acetate 0:10 to 1:9. After evaporation of the selected fractions, the title compound was obtained as a yellow powder.

1H-NMR (CDCl_3 , 400 MHz, δ in ppm): 8.40 - 8.10 (brs, 2H), 7.58 (d, 2H), 7.52 – 7.45 (m, 3H), 5.17 (m,
15 1H), 4.30 (m, 1H), 4.20 (m, 1H), 4.08 (d, 1H), 3.70 (m, 2H), 3.60 (m, 1H), 2.40 (s, 3H).

Step C: N-(2-cyclobutylsulfonylisoxazolidin-4-yl)-4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide




20

A solution of 4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-isoxazolidin-2-ium-4-yl-2-methyl-benzamide; 2,2,2-trifluoroacetate (described in step B) (0.150 g,) in tetrahydrofuran (5.0 ml), under argon atmosphere was treated with pyridine (0.038 g) and cyclubutane sulfonyl chloride (0.187
25 g). The reaction mixture was stirred for 18 hours at 20°C. The reaction was followed by TLC (silica gel, Ethyl acetate/Heptane 1:1). The reaction mixture was evaporated under reduced pressure and the residue was submitted to flash column chromatography over silica gel, eluting with Ethyl acetate/Heptane 1:1. The selected fractions were evaporated to yield the title compound (example A014 in table A) as a colourless solid.

30 1H-NMR (CDCl_3 , 400 MHz, δ in ppm): 7.60 (d, 2H), 7.52 – 7.40 (m, 3H), 6.97 (d, 1H), 5.32 (m, 1H), 4.45 (t, 1H), 4.27 (dd, 1H), 4.10 – 3.95 (m, 3H), 3.73 - 3.63 (m, 2H), 2.49 (s, 3H), 1.30 – 1.15 (m, 2H), 0.75 (d, 2H), 0.45 (m, 2H).

Example 2: 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide (example A001 in table A)

Step A: tert-butyl 4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate




Following similar conditions as described in step A in example 1, but starting with the stereoenriched acid 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid (described in WO 2013026931), tert-butyl 4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate was prepared as a mixture of diastereomers.

¹HNMR (CDCl₃, 400 MHz): 7.59 (s, 1H), 7.58 (s, 1H), 7.53 - 7.47 (m, 2H), 7.38 (d, J = 8 Hz, 1H), 6.24 (d, J = 8 Hz, 1H), 5.05 - 4.97 (m, 1H), 4.12 - 4.01 (m, 3H), 3.97 (dd, J₁ = 10 Hz, J₂ = 3 Hz, 1H), 3.68 (d, J = 17 Hz, 1H), 3.60 (dd, J₁ = 12 Hz, J₂ = 3 Hz, 1H), 2.46 (s, 3H), 1.48 (s, 9H).

Steps B and C: 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide

Similarly to steps B and C of example 1, but using methanesulfonyl chloride instead of cyclobutanesulfonyl chloride, the title compound was obtained as a mixture of diastereomers (compound A001 in table A) (LC-MS data in the table).

Example 3: Single diastereomers 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide (example A013 in Table A) and 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4S)-2-methylsulfonylisoxazolidin-4-yl]benzamide (example A012 in table A)



The intermediate, tert-butyl 4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate obtained in step A of preparation example 2 was submitted to chiral resolution by preparative HPLC chromatography using the conditions outlined hereafter.

Analytical HPLC method:

HPLC: Waters UPLC – HClass, DAD Detector Waters UPLC

Column: Daicel CHIRALPAK® IB, 3µm, 0.46cm x 10cm

Mobile phase: Hept/EtOH 90/10

5 Flow rate: 1.0 ml/min

Detection: 265 nm

Sample concentration: 1mg/mL in DCM/iPrOH 50/50

Injection: 2µl

10 Preparative HPLC method:

Autopurification System from Waters: 2767 sample Manager, 2489 UV/Visible Detector, 2545 Quaternary Gradient Module.

Column: Daicel CHIRALPAK® IB, 5µm, 1.0 cm x 25cm

Mobile phase: Hept/EtOH 90/10

15 Flow rate: 10 ml/min

Detection: UV 260 nm

Sample concentration: 50mg/mL in (4ml Heptane/2ml EtOH)


Injection: 250µl - 500µl

Results:

First eluting enantiomer	Second eluting enantiomer
Retention time (min) ~ 6.68	Retention time (min) ~ 10.79
Chemical purity (area% at 265 nm) 99	Chemical purity (area% at 265 nm) 99
Enantiomeric excess (%) > 99	Enantiomeric excess (%) > 99


20

The intermediate with the elution time of 6.68 min is tert-butyl (4R)-4-[[4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate



while the intermediate with the elution time of 10.79 min is tert-butyl (4S)-4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-carboxylate.

25



Each diastereomer was submitted to the synthetic steps B and C, as described for the preparation

30 example 2 to yield, respectively:


4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide (example A013 in Table A) with a retention time of 21.71 min under the conditions described hereafter and

- 5 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4S)-2-methylsulfonylisoxazolidin-4-yl]benzamide (example A012 in table A) with a retention time of 17.61 min under the same conditions.

Analytical method:

- 10 HPLC: Waters UPLC - HClass
DAD Detector Waters UPLC
Column: Daicel CHIRALPAK® IB, 3µm, 0.46cm x 10cm
Mobile phase: Heptane/EtOH 90/10
Flow rate: 1.0 ml/min
15 Detection: 265nm
Sample concentration: 1 mg/mL in DCM/iPr 50/50
Injection: 2µL

- Example 4: 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfonyl)isoxazolidin-4-yl]-2-methyl-benzamide (example A010 in table A)
- 20



- Following exactly the same synthetic sequence as in synthesis example 2, but using dimethylsulfonyl chloride instead of methanesulfonyl chloride, the title compound was prepared as a mixture of
- 25 diastereomers. LC-MS data is indicated in table A.

The mixture was submitted to preparative separation by chiral HPLC under the conditions described hereunder.

Analytical HPLC method:

- 30 HPLC: Waters UPLC – HClass, DAD Detector Waters UPLC
Column: Daicel CHIRALPAK® IA, 3µm, 0.46cm x 10cm
Mobile phase: TBME/EtOH 98/02
Flow rate: 1.0 ml/min
Detection: 265 nm
35 Sample concentration: 1mg/mL in DCM/iPrOH 50/50
Injection: 2µl

Preparative HPLC method:

Autopurification System from Waters: 2767 sample Manager, 2489 UV/Visible Detector, 2545 Quaternary Gradient Module.

Column: Daicel CHIRALPAK® IA, 5µm, 1.0 cm x 25cm

Mobile phase: TBME/EtOH 98/02

5 Flow rate: 10 ml/min

Detection: UV 260 nm

Sample concentration: 25mg/mL in dichloromethane, filtered

Injection: 100µl - 200µl


First eluting enantiomer (compound A143 in table A)	Second eluting enantiomer (compound A034 in table A)
Retention time (min) ~ 3.6	Retention time (min) ~ 4.77
Chemical purity (area% at 266 nm) 99	Chemical purity (area% at 266 nm) 99
Enantiomeric excess (%) > 99	Enantiomeric excess (%) > 99

10

Example 5: 2-chloro-N-[(4R)-2-(cyclopropylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide (compound A154 in table A)

Step A: methyl 2-[(4S)-4-hydroxyisoxazolidine-2-carbonyl]benzoate


15




In a sulfonation flask with mechanical stirrer, condenser, thermometer, under inert atmosphere, a suspension of N-hydroxyphthalimide (40.0 g) and (R)-(-)-epichlorhydrin (25.0 g) in anhydrous 1,4-dioxane (240 mL), was treated with triethylamine (3.42 mL). The suspension turned immediately orange. The resulting mixture was then heated to 55°C and stirred at that temperature for 6 days. To the resulting dark red solution were added methanol (240 mL) and triethylamine (34.2 mL) and the stirring was continued for 3 hours at the same temperature.

The reaction mixture was then concentrated under vacuum to yield a dark red oil which was treated with a little dichloromethane and saturated aqueous sodium bicarbonate solution. The mixture was extracted twice with dichloromethane. The organic phase was dried over sodium sulfate and evaporated under reduced pressure. The residue was submitted to silica gel chromatography, using methanol/dichloromethane 5:95 as eluent. The selected fractions were evaporated and the residue was triturated with ethyl acetate. A white suspension formed slowly. Methyl 2-[(4S)-4-hydroxyisoxazolidine-2-carbonyl]benzoate was isolated by filtration.

¹HNMR (CDCl₃, 400 MHz): 7.99 (d, J = 8 Hz, 1H), 7.62 (t, J = 8 Hz, 1H), 7.51-7.42 (m, 2H), 4.81-4.72 (m, 1H), 4.35-3.63 (m, 4H), 3.92 (s, 3H).


Step B: (4S)-isoxazolidin-2-ium-4-ol chloride

- 5 Methyl 2-[(4S)-4-hydroxyisoxazolidine-2-carbonyl]benzoate (264 g) was placed in a flask equipped with a mechanical stirrer, a condenser, a thermometer and an inert gas inlet and outlet through a bubbler. A 4 molar anhydrous solution of hydrochloric acid in dioxane (1.313 L) was added. The resulting solution was stirred at 80°C for 20 hours. Care was taken to trap the hydrochloric gas leaving the apparatus. The reaction mixture was then cooled to 0°C and the solid that was formed was isolated by careful
- 10 filtration. After rinsing with dry dioxane and drying, (4S)-isoxazolidin-2-ium-4-ol chloride was isolated as colorless powder. The compound was analyzed by NMR.
- ¹HNMR (DMSO-d₆, 400 MHz): 13.1-12.1 (br s, 2H), 6.2-5.2 (br s, 1H), 4.85-4.77 (m, 1H), 4.08 (d, 1H), 4.04 (dd, 1H), 3.49 (dd, 1H), 3.33 (d, 1H).

15 Step C: tert-butyl (4S)-4-hydroxyisoxazolidine-2-carboxylate

- A suspension of (4S)-isoxazolidin-2-ium-4-ol chloride (67.8 g) in dry tetrahydrofuran (300 mL) under inert atmosphere was stirred at 0°C. A solution of triethylamine (82 g) in tetrahydrofuran (100 mL) was added
- 20 over a period of 10 minutes. A solution di-tert-butyl dicarbonate (141 g) in tetrahydrofuran (100 mL) was then added to the reaction mixture. After the removal of the ice bath, the reaction mixture was stirred at room temperature until the starting material was reacted.
- The progression of the reaction is monitored by TLC analysis of aliquots eluted on a silica gel plate with a mixture of ethyl acetate/heptane 1:1. The plate can be developed with iodine vapour or ninhydrine.
- 25 The reaction mixture was filtered and the precipitate washed with tetrahydrofuran. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column using first a gradient heptane/ethyl acetate 3:1 to 1:3.
- The tert-butyl (4S)-4-hydroxyisoxazolidine-2-carboxylate is isolated as a pale yellow oil, characterized by its NMR spectrum.
- 30 ¹HNMR (CDCl₃, 400 MHz): 4.78-4.74 (m, 1H), 3.96-3.94 (m, 2H), 3.72-3.68 (m, 2H), 2.36 (broad d, 1H), 1.48 (s, 9H).

Step D: tert-butyl (4S)-4-methylsulfonyloxyisoxazolidine-2-carboxylate




To a solution of tert-butyl (4S)-4-hydroxyisoxazolidine-2-carboxylate (5.00 g) in dichloromethane (30 mL) was added triethylamine (7.52 mL). The reaction flask was cooled in an ice bath and a solution of methanesulfonyl chloride (3.13 mL) in dichloromethane (10 mL) was slowly added in such a way to keep the temperature below 20°C. The resulting orange colored suspension was stirred for 16 hours at 20°C and resulted in a brown suspension.

The reaction mixture was then washed with 1M aqueous hydrochloric acid (30 mL) and the aqueous phase was extracted with dichloromethane. The combined organic extracts were washed with 1M aqueous sodium hydroxide (30 mL). The aqueous phase was extracted with dichloromethane and the combined organic extracts were dried over sodium sulfate and concentrated under vacuo to yield crude tert-butyl (4S)-4-methylsulfonyloxyisoxazolidine-2-carboxylate that was pure enough to be used in the following step.

¹HNMR (CDCl₃, 400 MHz): 5.53-5.47 (m, 1H), 4.20-4.15 (broad d, J=10 Hz, 1H), 4.11-4.05 (dd, J₁=10 Hz, J₂= 5 Hz, 1H), 4.03-3.97 (broad d, J=14 Hz, 1H), 3.89-3.82 (dd, J₁=14 Hz, J₂= 5 Hz, 1H), 3.07 (s, 3H), 1.50 (s, 9H).

Step E: tert-butyl (4R)-4-azidoisoxazolidine-2-carboxylate



Sodium azide (21.5 g) was suspended in 170 mL of dimethylformamide (dried over molecular sieves) in a 500 mL flask fitted with a mechanical stirrer, a thermometer, a condenser, an argon inlet and outlet with a bubbler and a dropping funnel. The reaction flask was placed in an oil bath and heated under stirring at 60°C. The reaction was performed under argon atmosphere and behind a safety shield. Special care was given to the absence of chlorinated solvent traces in the starting mesylate for safety reasons. A solution of tert-butyl (4S)-4-methylsulfonyloxyisoxazolidine-2-carboxylate (described above) (68 g) in dry dimethylformamide (170 mL) was added under stirring. No exotherm was observed. After the addition was complete, the reaction mixture was stirred at the same temperature for 22 hours. The progression of the reaction can be followed by TLC analysis (silica gel plate, eluent ethyl acetate- heptane 1:1, R_f starting material 0.3, R_f product 0.6).


The reaction mixture was then let cool down to room temperature and transferred in a mixture of water (1 L) and ethyl acetate. The resulting mixture was extracted three times with ethyl acetate. The combined organic extracts were washed three times with water, then with brine before being dried over sodium

sulfate. After removal of the solvent under vacuum, the crude tert-butyl (4R)-4-azidoisoxazolidine-2-carboxylate was obtained as a light orange oil. It was used without purification for the following step.

¹HNMR (CDCl₃, 400 MHz): 4.47-4.42 (m, 1H), 4.07-4.01 (dd, 1H), 3.97-3.92 (dd, 1H), 3.80-3.77 (d, 2H), 1.51 (s, 9H).

5

Step F: tert-butyl (4R)-4-aminoisoxazolidine-2-carboxylate



A solution of tert-butyl (4R)-4-azidoisoxazolidine-2-carboxylate (56 g) in tetrahydrofuran (1300 mL) was placed in an inertized autoclave equipped with a mechanical stirrer. Palladium on carbon (2.50 g, 5% catalyst loading) was introduced and the reactor was closed. After evacuation of the reactor's atmosphere hydrogen was introduced (this operation was repeated twice). The applied pressure didn't exceed 3 atm. The reaction was performed under strong agitation, at room temperature. The progression was checked by analysis of aliquots of the reaction mixture (after removal of hydrogen under vacuum and filling the reactor with argon). After 8 hours, another portion of catalyst (1.25 g) was introduced, following a safe protocol and the hydrogenation was continued for two more hours, after which the conversion was complete. After inertizing the reactor, the reaction mixture was filtered over a short path of celite and the filtrate was evaporated on the rotary evaporator at 60°C. The crude tert-butyl (4R)-4-aminoisoxazolidine-2-carboxylate was obtained as a pale brown oil.

¹HNMR (CDCl₃, 400 MHz): 4.00-3.89 (m, 2H), 3.87-3.80 (dd, 1H), 3.70-3.65 (dd, 1H), 3.38-3.33 (dd, 1H), 1.48 (s, 9H).

Enantiomeric purity of tert-butyl (4R)-4-aminoisoxazolidine-2-carboxylate: Chiral analysis of the product was performed on a chiral HPLC, using a racemic compound as reference.

HPLC: Waters UPLC - HClass

DAD Detector Waters UPLC

Column: Daicel CHIRALPAK® IF, 3μm, 0.46cm x 10cm

Mobile phase: Heptane/EtOH/DEA 80/20/0.1%

Flow rate: 1.0 ml/min

Detection: 220 nm

Sample concentration: 1 mg/mL in iPrOH 100%


Injection: 2μL

Under these conditions, the enantiomers have a retention time of 5.99 min and 9.69 min respectively.

The product obtained as described above showed only the peak of the short retention time, the other

isomer being not detected.


Step G: tert-butyl (4R)-4-[[2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]]benzoyl]amino]isoxazolidine-2-carboxylate



This step was performed in an analogous way as described in previous examples.

5

Step H: 2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-isoxazolidin-4-yl]benzamide



10

This step was performed in an analogous way as described in previous examples.

Step I: N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide (compound A154 in table A)

15




A solution of 2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-isoxazolidin-4-yl]benzamide (0.35 g) in tetrahydrofuran (5.5 mL) under nitrogen atmosphere, was treated with cyclopropylsulfamoyl chloride (0.161 g) (obtained in two steps by adding one equivalent of chlorosulfonic acid to three equivalents of cyclopropylamine in dichloromethane at 0°C, filtering and treating the solid residue with PCl₅ in toluene at 75°C for 2 hours, the product being in solution can be recovered by decanting and evaporation of the solvent.) followed by pyridine (0.065 g). After one hour stirring at 20°C, the reaction mixture was evaporated and the residue was submitted to chromatography over silica gel to yield the title compound that was characterized by NMR.

¹HNMR (CDCl₃, 400 MHz): 7.75-7.65 (m, 2H), 7.64 – 7.48 (m, 1H), 7.50 (s, 2H), 7.42 (s, 1H), 7.20 (d, 1H), 5.45-5.35 (m, 1H), 5.11 (s, 1H), 4.48 (t, 1H), 4.21-4.15 (m, 1H), 4.05 (d, 1H), 3.94-3.86 (m, 1H), 3.85-3.76 (m, 1H), 3.68 (d, 1H), 2.73-2.67 (m, 1H), 0.90-0.65 (m, 4H).

25

Example 6: N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide (compound A037 in table A)

Step A: [(4S)-2-(dimethylsulfamoyl)isoxazolidin-4-yl] methanesulfonate



5


To a suspension of (4S)-isoxazolidin-2-ium-4-ol chloride (synthesis described in step B of example 5) (5.00 g) in tetrahydrofuran (50 mL) under inert atmosphere was added triethylamine (19.6 mL). The mixture was stirred at 20°C for 15 minutes, then N,N-dimethylsulfamoyl chloride (4.75 mL) was added. The reaction mixture was then heated to 60°C and stirred for 20 hours.

- 10 The resulting suspension was then cooled down to 0°C and methanesulfonyl chloride (3.42 mL) was added dropwise, so that the temperature was kept under 30°C. The mixture was then stirred at 20°C for 2 hours. The reaction mixture was treated with water (150mL) and extracted three times with ethyl acetate. The combined organic phases were washed with a saturated aqueous solution of sodium bicarbonate, followed by brine and dried over sodium sulfate. After removal of the solvent, the pale brown oil was
- 15 crystallized from ethyl acetate / hexanes to yield off-white crystals of the title product that was characterized by NMR.

¹HNMR (CDCl₃, 400 MHz): 5.67-5.58 (m, 1H), 4.50 (dd, J₁ = 10.4 Hz, J₂ = 6.2 Hz, 1H), 4.24 (dd, J₁ = 10.4 Hz, J₂ = 2.6 Hz, 1H), 4.15 (dd, J₁ = 14.3 Hz, J₂ = 6.2 Hz, 1H), 3.71 (dd, J₁ = 14.3 Hz, J₂ = 2.6 Hz, 1H), 3,10 (s, 3H), 2.96 (s, 3H).

20

Steps B then C: (4R)-4-amino-N,N-dimethyl-isoxazolidine-2-sulfonamide



The title compound was obtained in an analogous way as described in steps E and F of synthesis example 5. It was characterized by NMR analysis.

- 25 ¹HNMR (CDCl₃, 400 MHz): 4.25 (t, 1H), 4.07-3.99 (m, 1H), 3.79 (dd, 1H), 3.73 (dd, 1H), 3.25 (dd, 1H), 2.89 (s, 6H).

Step D: N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide

In an analogous way as described in previous synthesis examples, the acid chloride obtained from the stereo enriched 2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoic acid was reacted with the (4R)-4-amino-N,N-dimethyl-isoxazolidine-2-sulfonamide described above to yield the title compound. The LC-MS analysis (conditions in Table A) indicated a retention time of 1.23


5 minute and [M-H]⁻ m/z peaks: 627, 629, 631, 633.

¹HNMR (CDCl₃, 400 MHz): 7.65 (s, 2H), 7.54-7.42 (m, 3H), 7.00 (br. d, 1H), 5.36-5.27 (m, 1H), 4.45 (t, 1H), 4.15 (dd, 1H), 4.07 (d, 1H), 3.87-3.80 (m, 1H), 3.78 - 3.62 (m, 2H), 2.97 (s, 6H), 2.47 (s, 3H).

Example 7: Synthesis of 4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-

10 ethylsulfonylisoxazolidin-4-yl]-2-methyl-benzamide (Compound A121 in table A)

Step A: [(4S)-2-ethylsulfonylisoxazolidin-4-yl] ethanesulfonate




To a stirred suspension of (4S)-isoxazolidin-2-ium-4-ol chloride (synthesis described in step B of example 15 5) (2.50 g) in dichloromethane (100 mL) at 0°C was added triethylamine (11.2 mL), whereby the reaction mixture became a yellow solution. Ethanesulfonyl chloride (6.53 g) was then added in such a way that the temperature didn't exceed 25°C. A suspension formed and the reaction mixture was stirred at 0°C for 30 min, then for 1 hour at 20°C.

The reaction mixture was then washed with hydrochloric acid aqueous solution (1 N), followed by water 20 and brine. After drying of the organic phase over sodium sulfate and removal of the solvent, the dark brown oil was purified by chromatography over a silica gel, eluting with a mixture of heptane and ethyl acetate. The title compound was isolated as a brown oil and characterized by NMR.

¹HNMR (CDCl₃, 400 MHz): 5.65-5.59 (m, 1H), 4.47 (dd, 1H), 4.37 (dd, 1H), 4.21 (dd, 1H), 3.72 (dd, 1H), 3.42-3.34 (m, 2H), 3.22 (q, 2H), 1.47-1.38 (m, 6H).

25

Step B: (4R)-4-azido-2-ethylsulfonyl-isoxazolidine




In a similar way as described above, but with heating at 90°C, the title compound was obtained as a pale yellow oil.

¹HNMR (CDCl₃, 400 MHz): 4.63-4.55 (m, 1H), 4.41 (dd, 1H), 4.21 (dd, 1H), 4.04 (dd, 1H), 3.45-3.20 (m, 3H), 1.44 (t, 3H).

5

Step C: (4R)-2-ethylsulfonylisoxazolidin-4-amine




In a similar way as described above, the title compound was obtained as a colorless oil.

¹HNMR (CDCl₃, 400 MHz): 4.31 (dd, 1H), 4.14-4.06 (m, 1H), 3.90 (dd, 1H), 3.82 (dd, 1H), 3.38-3.25 (m, 3H), 1.60 (br. s, 2H), 1.42 (t, 3H).

10

By analogy, the compound (4R)-2-methylsulfonylisoxazolidin-4-amine, hereafter, was also prepared



as a light yellow oil.

¹HNMR (CDCl₃, 400 MHz): 4.34 (dd, 1H), 4.13-4.07 (m, 1H), 3.88 (dd, 1H), 3.84 (dd, 1H), 3.34 (dd, 1H), 3.12 (s, 3H), 1.56 (br. s, 2H).

15

Step D: 4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-2-methyl-benzamide

20

After reaction of the acid chloride obtained from the stereo-enriched 4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid with the (4R)-2-ethylsulfonylisoxazolidin-4-amine, in a similar way as described in the previous synthetic examples, the title compound was obtained.


¹HNMR (CDCl₃, 400 MHz): 7.53-7.47 (m, 4H), 7.46 – 7.41 (m, 2H), 6.95 (d, 1H), 5.36-5.28 (m, 1H), 4.50-4.43 (m, 1H), 4.18-4.13 (m, 1H), 4.08 (d, 1H), 3.98 (d, 1H), 3.72 - 3.64 (m, 2H), 3.45-3.23 (m, 2H), 2.47 (s, 3H), 1.43 (t, 3H).

25

Example 8 : 4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide

30


Step A: tert-butyl 4-[(4-acetyl-2-methyl-benzoyl)amino]isoxazolidine-2-carboxylate



Treatment of 4-acetyl-2-methyl-benzoyl chloride with tert-butyl 4-aminoisoxazolidine-2-carboxylate in an analogous way as for step A of example 1 leads to the title product as a viscous oil.

¹HNMR (CDCl₃, 400 MHz): 7.76 (s, 1H), 7.72 (d, 2H), 7.39 (d, 1H), 6.58 (br. d, 1H), 5.05-4.96 (m, 1H),
 5 4.10-4.02 (m, 2H), 3.97 (dd, 1H), 3.60 (dd, 1H), 2.58 (s, 3H), 2.45 (s, 3H), 1.47 (s, 9H).


Step B: 4-acetyl-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide



Proceeding in an analogous way as in step B and step C of the example 1, but using methanesulfonyl
 10 chloride instead of cyclobutanesulfonyl chloride, the title compound was obtained as colorless crystals
 with a melting point of 153-155°C.

¹HNMR (CDCl₃, 400 MHz): 7.80-7.73 (m, 2H), 7.44 (d, 1H), 6.94 (br. d, 1H), 5.37-5.29 (m, 1H), 4.48 (t,
 1H), 4.18 (dd, 1H), 3.96 (d, 1H), 3.71 (dd, 1H), 3.17 (s, 3H), 2.60 (s, 3H), 2.48 (s, 3H).

15 Step C: 4-[(E-Z)-3-(3,5-dichlorophenyl)-4,4,4-trifluoro-but-2-enoyl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide




4-Acetyl-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide (3.00 g), 1-(3,5-dichlorophenyl)-2,2,2-trifluoro-ethanone (3.35 g) and potassium carbonate (3.18 g) were mixed in tetrahydrofuran (50 mL). The
 20 suspension was stirred at 65°C for 44 hours.

The resulting mixture was diluted with water (200 mL) and 2N hydrochloric acid (25 mL), then extracted twice with ethyl acetate. The organic phase was washed with water, brine and dried over sodium sulfate. After evaporation of the solvent, the crude product was submitted to chromatography over silica gel.

¹HNMR (CDCl₃, 400 MHz): 7.68-7.63 (m, 2H), 7.45 (d, 1H), 7.35 (d, 1H), 7.14 (s, 2H), 6.90 (br.d, 1H),
5 5.38-5.28 (m, 1H), 4.50 (t, 1H), 4.20 (dd, 1H), 4.01-3.95 (m, 1H), 3.68 (dd, 1H), 3.18 (s, 3H), 2.48 (s, 3H).

Step D : 4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide



10


To a stirred solution of 4-[(E-Z)-3-(3,5-dichlorophenyl)-4,4,4-trifluoro-but-2-enoyl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide (3.15 g) in chloroform (57 mL) stirred at -20°C under nitrogen atmosphere, was added a solution of cesium hydroxide hydrate (2.89 g) in water (3 mL), followed by a 50% aqueous solution of hydroxylamine (0.76 g) and and (R)-[1-(9-anthrylmethyl)-5-vinyl-quinuclidin-1-ium-2-yl]-(6-methoxy-4-quinolyl)methanol chloride (0.63 g). (0.2 equiv.). After 3 hours at that temperature,
15 the bath was removed and the mixture was stirred at 20°C for 15 hours.

The reaction mixture was diluted with dichloromethane (50 ml), then washed with diluted HCl. As the phase separation was difficult, the mixture was treated with aqueous sodium hydroxide to pH 11. The organic layer was dried over sodium sulfate and concentrated. The residue was submitted to silica gel

20 chromatography to yield the title product that was characterized by NMR.

¹HNMR (CDCl₃, 400 MHz): 7.54-7.47 (m, 4H), 7.46 – 7.39 (m, 2H), 6.91 (br.d, 1H), 5.35-5.27 (m, 1H), 4.50-4.48 (t, 1H), 4.20-4.14 (m, 1H), 4.14-4.04 (m, 1H), 3.95 (d, 1H), 3.75 - 3.66 (m, 2H), 3.17 (s, 3H), 2.47 (s, 3H)


25 Example 9: Method for preparing compounds of the invention from a carboxylic acid



To a solution of an acid of formula RCOOH (45 μmol), for example 4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoic acid in the case of Compound No. A040 of Table AX in dimethylacetamide (0.3 ml) was added successively a solution of 2-methylsulfonylisoxazolidin-4-amine
30 (30 μmol) in dimethylacetamide (0.25 ml), diisopropylethylamine (Hunig's base) (0.6 ml), and a solution of bis(2-oxo-3-oxazolidinyl)phosphonic chloride ("BOP-Cl") (15.3mg) in dimethylacetamide (0.25 ml). The reaction mixture was stirred at 50°C for 16 hours. Then the reaction mixture was evaporated to dryness.

The remaining mixture was dissolved with methanol / dimethylacetamide (4:1) (0.7 ml) and purified by preparative HPLC. This method was used to prepare a number of compounds (Compound Nos. A040 and A045 to A066 of Table A) in parallel.

- 5 Example 10: Resolution of 4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-benzamide (examples A037 and A112 in table A)



The mixture of isomers of N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-

- 10 trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide was submitted to preparative separation by chiral HPLC under the conditions described hereunder.

Analytical HPLC method:

HPLC: Waters UPLC – HClass, DAD Detector Waters UPLC

Column: Daicel CHIRALPAK® IF, 3μm, 0.46cm x 10cm

- 15 Mobile phase: Hept/EtOAc 70/30

Flow rate: 1.0 ml/min

Detection: 265 nm

Sample concentration: 1mg/mL in DCM/iPrOH 50/50

Injection: 2μl

20

Preparative HPLC method:

Autopurification System from Waters: 2767 sample Manager, 2489 UV/Visible Detector, 2545 Quaternary Gradient Module.

Column: Daicel CHIRALPAK® IF, 5μm, 1.0 cm x 25cm

- 25 Mobile phase: Hept/EtOAc 70/30

Flow rate: 10 ml/min

Detection: UV 260 nm

Sample concentration: 166 mg/mL in EtOAc/Hept, filtered

Injection: 30μl - 150μl


30

First eluting enantiomer (compound A112 in table A)	Second eluting enantiomer (compound A037 in table A)
Retention time (min) ~ 5.45	Retention time (min) ~ 8.86
Chemical purity (area% at 265 nm) 99	Chemical purity (area% at 265 nm) 99

Enantiomeric excess (%) > 99

Enantiomeric excess (%) > 99

Example 11: (4R)-4-[[4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidine-2-sulfonyl fluoride (compound A198 in table A)



To a solution of 1.000 g of the compound 4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-isoxazolidin-4-yl]-2-methyl-benzamide (obtained by analogy as in step G of example 5) in tetrahydrofuran (15 ml), was added triethylamine (0.207 g). A slow stream of sulfonyl fluoride was bubbled through the mixture for one hour, then the mixture was stirred overnight at 20°C. The reaction mixture was then concentrated under vacuum. The residue was dissolved in ethyl acetate and washed with water, dried over sodium sulfate and the crude product was purified by chromatography over silica gel eluting with a mixture of ethyl acetate and heptane.

The title compound was characterized by LC-MS (see table A) and NMR.

¹HNMR (CDCl₃, 400 MHz): 7.50 (s, 2H), 7.48 – 7.39 (m, 3H), 7.38 – 7.32 (m, 1H), 6.75 – 6.60 (m, 1H), 5.28-5.18 (m, 1H), 4.47-4.40 (m, 1H), 4.20 (d, 1H), 4.18-4.05 (m, 2H), 3.84 (d, 1H), 3.71 (d, 1H), 2.40 (s, 3H)

Table A: Compounds according to the invention

Compound number	chemical name	LC-MS Method	RT (min)	[M+H] ⁺	[M-H] ⁻
A1	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	A	1.14	584 / 586 / 588	
A2	N-[2-(4-chlorophenyl)sulfonylisoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide	A	1.26	678 / 680 / 682 / 684	
A3	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-propylsulfonylisoxazolidin-4-yl)benzamide	A	1.21	612 / 614 / 616	
A4	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-	A	1.18	598 / 600 / 602	

	ethylsulfonylisoxazolidin-4-yl)-2-methylbenzamide				
A5	N-[2-(chloromethylsulfonyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.18	618 / 620 / 622 / 624	
A6	N-(2-cyclopropylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.18	610 / 612 / 614	
A7	N-[2-(2-chloroethylsulfonyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.17	632 / 634 / 636 / 638	
A8	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[2-(2,2,2-trifluoroethylsulfonyl)isoxazolidin-4-yl]benzamide	A	1.2	652 / 654 / 656	
A9	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[2-(2-thienylsulfonyl)isoxazolidin-4-yl]benzamide	A	1.19	652 / 654 / 656	
A10	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.19	613 / 615 / 617	
A11	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[2-(3,3,3-trifluoropropylsulfonyl)isoxazolidin-4-yl]benzamide	A	1.22	666 / 668 / 670	
A12	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4S)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.14	584 / 586 / 588	
A13	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.14	584 / 586 / 588	
A14	N-(2-cyclobutylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.22	624 / 626 / 628	

A15	2-chloro-N-(2-methylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.17	620 / 622 / 624 / 626	
A16	4-[(5S)-5-[3-bromo-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	A	1.14	644 / 646	
A17	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-isopropylsulfonylisoxazolidin-4-yl)-2-methylbenzamide	A	1.21	612 / 614 / 616	
A18	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(2-methoxyethylsulfonyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.17	628 / 630 / 632	
A19	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	A	1.15		602 / 604 / 606 / 608
A20	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	A	1.14	600 / 602	
A21	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.18	633 / 635 / 637	
A22	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.17	629 / 631	
A23	4-[(5S)-5-[3-bromo-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.18	673 / 675	
A24	2-chloro-N-(2-cyclopropylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.18	630 / 632 / 634 / 636	
A25	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-ethylsulfonylisoxazolidin-4-yl)benzamide	A	1.2	618 / 620 / 622 / 624	
A26	2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.18		598 / 600 / 602

A27	N-(2-ethylsulfonylisoxazolidin-4-yl)-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.2	614 / 616 / 618 / 620	
A28	2-chloro-N-(2-ethylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.22	634 / 636 / 638 / 640 / 642	
A29	N-(2-cyclopropylsulfonylisoxazolidin-4-yl)-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.21	626 / 628 / 630 / 632	
A30	2-chloro-N-(2-cyclopropylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.22	646 / 648 / 650 / 652	
A31	N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.22	629 / 631 / 633 / 635	
A32	2-chloro-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.23	649 / 651 / 653 / 655	
A33	4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	A	1.14		564 / 565 / 568
A34	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4S)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.19	613 / 615 / 617	
A35	2-methyl-N-(2-sulfamoylisoxazolidin-4-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.13	601 / 603 / 605	
A36	2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.18	600 / 602 / 604	
A37	N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.23		627 / 629 / 631 / 633
A38	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.14	566 / 568	
A39	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.19	595 / 597	

A40	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.89	566.13	
A41	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	B	1.17		593 / 595
A42	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-ethylsulfonylisoxazolidin-4-yl)-2-methylbenzamide	B	1.17		612 / 614
A43	2-bromo-N-(2-ethylsulfonylisoxazolidin-4-yl)-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	B	1.22		676 / 678 / 680
A44	2-bromo-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-ethylsulfonylisoxazolidin-4-yl)benzamide	B	1.19		660 / 662 / 664
A45	4-[5-(4-bromo-3,5-dichloro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.98	644.01	
A46	4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)naphthalene-1-carboxamide	D	1.97	602.11	
A47	2-cyano-4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.9	577.09	
A48	2-bromo-4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.89	630	
A49	2-chloro-4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.88	586.05	
A50	4-[5-(3,5-dichloro-4-cyano-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.76	591.09	
A51	4-[5-(3-chloro-5-methyl-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.81	546.16	
A52	2-cyclopropyl-4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.93	592.13	

A53	2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)-4-[5-[4-(trifluoromethoxy)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	D	1.79	582.16	
A54	4-[5-(2-fluoro-4-pyridyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.44	517.15	
A55	4-[5-(4-bromophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.74	576.1	
A56	8-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)quinoline-5-carboxamide	D	1.93	603.09	
A57	4-[5-[4-chloro-3,5-bis(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	2	668.1	
A58	4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)-2-(trifluoromethyl)benzamide	D	1.93	620.07	
A59	4-[5-[3,5-bis(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.91	634.15	
A60	2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)-4-[5-(trifluoromethyl)-5-[3-(trifluoromethyl)phenyl]-4H-isoxazol-3-yl]benzamide	D	1.74	566.17	
A61	2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)-4-[5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	D	1.97	600.06	
A62	4-[5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.89	600.12	
A63	4-[5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.89	584.11	
A64	4-[5-[chloro(difluoro)methyl]-5-(3,5-dichlorophenyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)benzamide	D	1.91	582.08	
A65	4-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)-2-vinyl-benzamide	D	1.91	578.12	
A66	4-[5-(3,4-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-	D	1.83	566.13	

	methylsulfonylisoxazolidin-4-yl)benzamide				
A67	2-methyl-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.17		615 / 617 / 619
A68	2-methyl-N-[(4R)-2-sulfamoylisoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.13		599 / 601 / 603
A69	2-methyl-N-[2-(methylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.18		613 / 615 / 617
A70	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.14	599 / 601 / 603	
A71	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-sulfamoylisoxazolidin-4-yl)benzamide	A	1.1	585 / 587 / 589	
A72	N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.25		641 / 643 / 645
A73	N-[2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.25		641 / 643 / 645
A74	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-[dimethylcarbamoyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-benzamide	A	1.16	670 / 672 / 674	
A75	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.14		598 / 600
A76	ethyl N-[4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidin-2-yl]sulfonyl-N-methyl-carbamate	A	1.21	671 / 673	
A77	methyl N-[4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidin-2-yl]sulfonyl-N-methyl-carbamate	A	1.18		655 / 657 / 659

A78	N-[4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidin-2-yl]sulfonyl-N-methyl-pyridine-3-carboxamide	A	1.18	704 / 706 / 708	
A79	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-[dimethylsulfamoyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-benzamide	A	1.2		704 / 706 / 708
A80	N-[2-[benzoyl(methyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide	A	1.24		701 / 703 / 705
A81	N-[2-[cyano(methyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide	A	1.18		622 / 624 / 626
A82	N-[2-[allyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.27		653 / 655 / 656
A83	2-methyl-N-[2-[methyl(oxiran-2-ylmethyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.21		669 / 671 / 674
A84	2-methyl-N-[2-[methyl(propyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.28		655 / 657 / 659
A85	N-[2-[butyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.31		669 / 671 / 673
A86	N-[2-[benzyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.31		703 / 706 / 707
A87	N-[2-[cyanomethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.2		652 / 654 / 656
A88	methyl 2-[methyl-4-[[2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoyl]amino]isoxazolidin-2-yl]sulfonyl-amino]acetate	A	1.21		*[M+HCOO]- 721 / 723 / 725

A89	ethyl 2-[methyl-4-[[2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoyl]amino]isoxazolidin-2-yl]sulfonyl-amino]acetate	A	1.24		699 / 701 / 703
A90	N-[2-[2-methoxyethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.23		671 / 673 / 676
A91	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.19	629 / 631	
A92	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[2-[methyl(methylsulfonyl)sulfamoyl]isoxazolidin-4-yl]benzamide	A	1.17	677 / 679 / 701	
A93	tert-butyl N-[4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidin-2-yl]sulfonyl-N-ethyl-carbamate	A	1.28		712 / 714 / 716
A94	tert-butyl N-allyl-N-[4-[[4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzoyl]amino]isoxazolidin-2-yl]sulfonyl-carbamate	A	1.28		723 / 725 / 727
A95	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(ethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.15	613 / 615 / 617	
A96	N-[2-[methoxymethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.22		657 / 659
A97	N-[2-(tert-butylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.19		641 / 643
A98	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[2-(diethylsulfamoyl)isoxazolidin-4-yl]-2-methylbenzamide	A	1.23		639 / 641 / 643

A99	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-pyrrolidin-1-ylsulfonylisoxazolidin-4-yl)benzamide	A	1.2		639 / 641 / 643
A100	N-[2-[tert-butyl(methyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide	A	1.25		655 / 657 / 659
A101	N-[2-(isopropylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.2		641 / 643 / 645
A102	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[2-(methylsulfonylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.01		661 / 663 / 665
A103	N-[2-[isopropyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.26		655 / 657 / 659
A104	N-[2-[ethyl(isopropyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.29	671 / 673 / 675	
A105	N-[2-(allylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-benzamide	A	1.15	625 / 627 / 629	
A106	4-[[2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoyl]amino]isoxazolidine-2-sulfonyl fluoride	A	1.25		602 / 604 / 606
A107	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.24	633 / 635 / 637	
A108	2-chloro-4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.23		647 / 649
A109	2-chloro-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.23	649 / 651 / 653 / 655	
A110	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.19		613 / 615 / 617

A111	N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(trifluoromethyl)-5-[3-(trifluoromethyl)phenyl]-4H-isoxazol-3-yl]benzamide	A	1.12	595	
A112	N-[(4S)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	*	*	*	*
A113	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.21	615 / 617 / 619	613 / 615 / 617
A114	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.18	601 / 603 / 605	599 / 601 / 603
A115	2-chloro-4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.19	635 / 637 / 639	633 / 635 / 637
A116	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.19	586 / 588 / 590	
A117	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]benzamide	A	1.22	600 / 602 / 604	
A118	N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.22	614 / 616 / 618	
A119	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-2-methylbenzamide	A	1.18	598 / 600 / 602	
A120	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-2-methylbenzamide	A	1.18		612 / 614 / 616
A121	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-2-methylbenzamide	A	1.17	580 / 582 / 584	
A122	2-chloro-4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-	A	1.18	634 / 636	

	4-yl]benzamide				
A123	2-chloro-4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.15	620 / 622	
A124	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]benzamide	C	2.05	633 / 635 / 637	631 / 633 / 635
A125	2-chloro-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	C	2.13	649 / 651 / 653 / 655	647 / 649 / 651 / 653
A126	2-chloro-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	C	2.07	635 / 637 / 639 / 641	633 / 635 / 637
A127	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.27	629 / 631 / 633	627 / 629 / 631
A128	2-chloro-4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.21	649 / 651 / 653	647 / 649
A129	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]benzamide	A	1.27	629 / 631 / 633	627 / 629 / 631
A130	2-chloro-4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]benzamide	A	1.27	663 / 665	661 / 663 / 665
A131	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]benzamide	C	2.18	647 / 649 / 651	645 / 647 / 649
A132	2-chloro-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	C	2.26	663 / 665 / 667	661 / 663 / 665 / 667

A133	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-benzamide	A	1.2		611 / 613 / 615
A134	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-benzamide	A	1.2		627 / 629
A135	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-benzamide	A	1.2		593 / 595 / 597
A136	N-[(4R)-2-(ethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.24		627 / 629 / 631
A137	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.19		597 / 599 / 601
A138	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.18		613 / 615
A139	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-(methylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.18		579 / 581 / 583
A140	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-benzamide	A	1.26		625 / 627 / 629
A141	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-benzamide	A	1.26		641 / 643
A142	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-benzamide	A	1.26		607 / 609
A143	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)isoxazolidin-4-yl]-2-methyl-	A	1.23	613 / 615	

	benzamide				
A144	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]benzamide	A	1.16	604 / 606 / 608 / 610	
A145	2-chloro-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.19	620 / 622 / 624	
A146	2-chloro-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]benzamide	A	1.23	618 / 620 / 622	
A147	2-chloro-N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.27		632 / 634 / 636
A148	N-[(4R)-2-(azetidin-1-ylsulfonyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.26		623 / 625
A149	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-(2,2,2-trifluoroethylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.25		665 / 667 / 669
A150	4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-[(4R)-2-[methyl(2,2,2-trifluoroethyl)sulfamoyl]isoxazolidin-4-yl]benzamide	A	1.25		679 / 681 / 683
A151	tert-butyl N-[(4R)-4-[[2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoyl]amino]isoxazolidin-2-yl]sulfonylcarbamate	A	1.19		685 / 687 / 687
A152	tert-butyl N-[(4R)-4-[[2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzoyl]amino]isoxazolidin-2-yl]sulfonyl-N-prop-2-ynyl-carbamate	A	1.28		723 / 725
A153	2-chloro-N-[(4R)-2-[cyclopropyl(methyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.23	641 / 643 / 645	

A154	2-chloro-N-[(4R)-2-(cyclopropylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.17	627 / 629 / 631	
A155	N-[(4R)-2-(cyclopropylsulfamoyl)isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.16	607 / 609. 611	
A156	N-[(4R)-2-[cyclopropyl(methyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methylbenzamide	A	1.23	621 / 623	
A157	2-chloro-N-[(4R)-2-[cyclopropyl(ethyl)sulfamoyl]isoxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.26	655 / 657 / 659	
A158	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-(prop-2-ynylsulfamoyl)isoxazolidin-4-yl]benzamide	A	1.15	625 / 627 / 629	
A159	2-methyl-N-[(4R)-2-methylsulfonylisoxazolidin-4-yl]-4-[(5S)-5-(trifluoromethyl)-5-[3-(trifluoromethyl)phenyl]-4H-isoxazol-3-yl]benzamide	A	1.07		564
A160	N-[(4R)-2-ethylsulfonylisoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(trifluoromethyl)-5-[3-(trifluoromethyl)phenyl]-4H-isoxazol-3-yl]benzamide	A	1.11	580	
A161	2-chloro-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-[(4R)-2-[ethyl(prop-2-ynyl)sulfamoyl]isoxazolidin-4-yl]benzamide	A	1.23	653 / 655 / 657 / 659	
A162	N-[2-[acetyl(methyl)sulfamoyl]isoxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,5-dichloro-4-fluoro-phenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]benzamide	A	1.2	641 / 643	
A163	N-[(4R)-2-(cyanomethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.17	609, 611	
A164	N-[(4R)-2-(chloromethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-	A	1.21	618, 620, 622	

	(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide				
A165	N-[(4R)-2-(cyanomethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.16		589, 591, 593
A166	N-[(4R)-2-(chloromethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.19	600, 602, 604	
A167	N-[(4R)-2-[(1-cyanocyclopropyl)sulfamoyl]-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.15	632, 634	
A168	N-[(4R)-2-[(1-cyanocyclopropyl)sulfamoyl]-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.16	650, 652	
A169	N-[(4S)-2-(dimethylsulfamoyl)-1,2-oxazolidin-4-yl]-4-[(5S)-5-[3-fluoro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.19	613	
A170	4-[(5S)-5-[3-fluoro-4-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.13	584	
A171	N-[(4S)-2-(dimethylsulfamoyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-[3-fluoro-4-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.18	613	
A172	4-[(5S)-5-[3-fluoro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-methylsulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.14	584	
A173	N-[(4S)-2-(bromomethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.20	644, 646, 648	
A174	2-[[[(4S)-4-[[4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidin-2-	A	1.10	610, 612, 614	

	yl)sulfonyl]acetic acid				
A175	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-ethenylsulfonyl-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.17	578, 580	
A176	methyl 2-[[[(4R)-4-[[4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidin-2-yl)sulfonyl]acetate	A	1.16	624, 628, 630	
A177	N-[(4R)-2-(2-bromoethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.22		656, 658, 660
A178	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-sulfamoyl-1,2-oxazolidin-4-yl]benzamide	A	1.09	567, 569, 571	
A179	N-[(4R)-2-[(1-cyanocyclopropyl)-methylsulfamoyl]-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.20	645, 647, 649	
A180	4-[(5R)-5-[3-fluoro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-(methylsulfamoyl)-1,2-oxazolidin-4-yl]benzamide	A	1.10	599	
A181	4-[(5S)-5-[4-fluoro-3-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(methylsulfamoyl)-1,2-oxazolidin-4-yl]benzamide	A	1.09	599	
A182	4-[(5S)-5-[3-fluoro-4-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(methylsulfamoyl)-1,2-oxazolidin-4-yl]benzamide	A	1.10	599	
A183	4-[(5S)-5-[4-fluoro-3-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.10	584	
A184	N-[(4S)-2-(dimethylsulfamoyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-[4-fluoro-3-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.15	613	

A185	4-[(5R)-5-[3-chloro-5-(difluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-methylsulfonyl-1,2-oxazolidin-4-yl]benzamide	B	1.09	582, 584	
A186	4-[(5R)-5-[3-chloro-5-(difluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-(dimethylsulfamoyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	B	1.14	611, 613	
A187	N-[(4S)-2-[(1-cyanocyclopropyl)-methylsulfamoyl]-1,2-oxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.22	664, 666	
A188	4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-morpholin-4-ylsulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.20	655, 657, 659	
A189	N-[(4R)-2-(bromomethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.20	662, 664, 666	
A190	methyl 2-[[[(4S)-4-[[4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidin-2-yl]sulfonyl]acetate	A	1.17	642, 644, 646	
A191	N-[(4R)-2-(2-bromoethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.23	676, 678, 680	
A192	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-(2,2,2-trifluoroethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.22	652, 654, 656	
A193	4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-ethenylsulfonyl-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.18	596, 598, 600	
A194	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(2,2,2-trifluoroethylsulfamoyl)-1,2-oxazolidin-4-yl]benzamide	A	1.19	649, 651	

A195	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(2,2,2-trifluoroethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.22	634, 636, 638	
A196	4-[(5S)-5-[3,5-dichloro-4-(difluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-(dimethylsulfamoyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.19	645, 647, 649	
A197	4-[(5S)-5-[3,5-dichloro-4-(difluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-methylsulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.13	616, 618, 620	
A198	(4R)-4-[[4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidine-2-sulfonyl fluoride	A	1.23	570, 572, 574	
A199	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-(3,3,3-trifluoropropylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.24	666, 668, 670	
A200	2-[[[(4S)-4-[[4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidin-2-yl]sulfonyl]acetic acid	A	1.10	628, 630	
A201	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-[2-(methylamino)-2-oxoethyl]sulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.10	623, 625	
A202	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-[2-(dimethylamino)-2-oxoethyl]sulfonyl-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.12	637, 639, 641	
A203	4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-[2-(hydroxyamino)-2-oxoethyl]sulfonyl-1,2-oxazolidin-4-yl]-2-methylbenzamide	C	1.79	643, 645, 647	
A204	4-[(5S)-5-[3,5-dichloro-4-(difluoromethoxy)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-methylsulfonyl-1,2-oxazolidin-4-yl]benzamide	A	1.13	632, 634, 636	

A205	4-[(5S)-5-[3,5-dichloro-4-(difluoromethoxy)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-(dimethylsulfamoyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.18	661, 663, 665	
A206	4-[(5R)-5-[3,5-dichloro-4-(difluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-(methylsulfamoyl)-1,2-oxazolidin-4-yl]benzamide	A	1.12	631, 633, 635	
A207	4-[(5R)-5-[3,5-dichloro-4-(difluoromethoxy)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-(methylsulfamoyl)-1,2-oxazolidin-4-yl]benzamide	A	1.12	647, 649, 651	
A208	4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-sulfamoyl-1,2-oxazolidin-4-yl]benzamide	A	1.10	585, 587, 589	
A209	tert-butyl N-[[[(4R)-4-[[4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidin-2-yl]sulfonyl]carbamate	A	1.21		683, 685, 687
A210	4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-(fluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.17	583, 586	
A211	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-(fluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.19	602, 604	
A212	2-methyl-N-[(4S)-2-methylsulfonyl-1,2-oxazolidin-4-yl]-4-[(5R)-5-[3-methyl-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]benzamide	C	1.91	580	
A213	(4S)-4-[[4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidine-2-sulfonyl fluoride	A	1.22	588, 590, 592	
A214	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(2,2,2-trifluoroethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.23	668, 670	
A215	2-methyl-N-[(4R)-2-methylsulfonyl-1,2-oxazolidin-4-yl]-4-[(5S)-5-(trifluoromethyl)-5-[6-	C	1.70	567	

	(trifluoromethyl)pyridin-2-yl]-4H-1,2-oxazol-3-yl]benzamide				
A216	N-[(4R)-2-(fluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methyl-4-[(5R)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]benzamide	A	1.23	618, 620, 622	
A217	N-[(4R)-2-(chloromethylsulfonyl)-1,2-oxazolidin-4-yl]-4-[(5R)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.20	634, 636	
A218	4-[(5R)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-(fluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.19	618, 620	
A219	(4R)-4-[[4-[(5R)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidine-2-sulfonyl fluoride	A	1.23	604, 606	
A220	N-[(4R)-2-(chloromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methyl-4-[(5R)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]benzamide	A	1.25	634, 636, 638	
A221	2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-(2,2,2-trifluoroethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.27	668, 670, 672	
A222	2-methyl-N-[(4R)-2-methylsulfonyl-1,2-oxazolidin-4-yl]-4-[5-(trifluoromethyl)-5-[4-(trifluoromethyl)pyridin-2-yl]-4H-1,2-oxazol-3-yl]benzamide	C	1.71	567	
A223	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(trifluoromethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.26	638, 640, 642	
A224	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4S)-2-(trifluoromethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.26	606, 608, 610	
A225	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-(2,2-difluoroethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.20	634, 636, 638	

A226	4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4R)-2-(difluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.21	620, 622, 624	
A227	4-[(5R)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methyl-N-[(4R)-2-(trichloromethylsulfonyl)-1,2-oxazolidin-4-yl]benzamide	A	1.31	654, 656, 658, 660, 662	
A228	N-[(4S)-2-cyclopropylsulfonyl-1,2-oxazolidin-4-yl]-4-[(5S)-5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.19	610, 612, 614	
A229	N-[(4S)-2-cyclopropylsulfonyl-1,2-oxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]benzamide	A	1.24	626, 628, 630	
A230	N-[(4S)-2-cyclopropylsulfonyl-1,2-oxazolidin-4-yl]-4-[(5S)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzamide	A	1.19	592, 594	
A231	N-[(4R)-2-(difluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methyl-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]benzamide	A	1.24	636, 638, 640	
A232	4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-(difluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.20	602, 604, 606	
A233	4-[(5S)-5-[3-chloro-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-N-[(4S)-2-(difluoromethylsulfonyl)-1,2-oxazolidin-4-yl]-2-methylbenzamide	A	1.20	636, 638	
A234	tert-butyl N-[[[(4R)-4-[[4-[(5R)-5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]-2-methylbenzoyl]amino]-1,2-oxazolidin-2-yl]sulfonyl]carbamate	A	1.20		665, 667, 669
A235	2-methyl-N-[(4S)-2-(1-methylcyclopropyl)sulfonyl-1,2-oxazolidin-4-yl]-4-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4H-1,2-oxazol-3-yl]benzamide	A	1.27	640, 642, 644	
A236	6-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-4-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)pyridine-3-	D	1.83	567.1	

	carboxamide				
A237	6-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-2-methyl-N-(2-methylsulfonylisoxazolidin-4-yl)pyridine-3-carboxamide	D	1.86	567.12	
A238	5-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)pyridine-2-carboxamide	D	1.85	553.09	
A239	1-[5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4H-isoxazol-3-yl]-N-(2-methylsulfonylisoxazolidin-4-yl)isoquinoline-4-carboxamide	D	2	603.11	

Biological examples

These Examples illustrate the pesticidal/insecticidal properties of compounds of formula (I).

Tests were performed as follows:

5

Diabrotica balteata (Corn root worm)

Maize sprouts placed onto an agar layer in 24-well microtiter plates were treated with aqueous test solutions prepared from 10'000 ppm DMSO stock solutions by spraying. After drying, the plates were infested with L2 larvae (6 to 10 per well). The samples were assessed for mortality and growth inhibition in comparison to untreated samples 4 days after infestation.

10

The following compounds gave an effect of at least 80% in at least one of the two categories (mortality or growth inhibition) at an application rate of 200 ppm:

A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17, A18, A19, A20, A21, A22, A23, A24, A25, A27, A28, A29, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43, A44, A45, A46, A48, A49, A50, A51, A52, A53, A54, A55, A57, A58, A59, A60, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, A98, A99, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A176, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A192, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, A221, A222, A223, A224, A225, A226, A227, A228, A229, A230, A231, A232, A233, A236, A237, A239.

25

Euschistus heros (Neotropical Brown Stink Bug)

Soybean leaves on agar in 24-well microtiter plates were sprayed with aqueous test solutions prepared from 10'000 ppm DMSO stock solutions. After drying the leaves were infested with N2 nymphs. The samples were assessed for mortality and growth inhibition in comparison to untreated samples 5 days after infestation.

- 5 The following compounds gave an effect of at least 80% in at least one of the two categories (mortality or growth inhibition) at an application rate of 200 ppm:

A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A17, A18, A19, A20, A26, A27, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43, A44, A45, A46, A48, A49, A50, A52, A54, A57, A58, A59, A60, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, A98, A99, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A172, A173, A174, A176, A178, A179, A180, A181, A183, A184, A185, A186, A187, A188, A189, A190, A192, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, A221, A223, A224, A225, A226, A227, A228, A229, A230, A231, A232, A233, A236, A239

20 **Myzus persicae (Green peach aphid):Feeding/Contact activity**

Sunflower leaf discs were placed onto agar in a 24-well microtiter plate and sprayed with aqueous test solutions prepared from 10'000 ppm DMSO stock solutions. After drying, the leaf discs were infested with an aphid population of mixed ages. The samples were assessed for mortality 6 days after infestation.

The following compounds resulted in at least 80% mortality at an application rate of 200 ppm:

25 A1, A3, A4, A5, A6, A7, A10, A11, A12, A13, A14, A15, A16, A17, A18, A19, A20, A21, A22, A23, A24, A25, A27, A28, A29, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43, A44, A45, A50, A59, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A90, A91, A93, A95, A96, A97, A98, A99, A100, A101, A103, A104, A105, A106, A107, A108, A109, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A161, A162, A164, A166, A168, A169, A172, A173, A174, A178, A180, A181, A183, A184, A185, A186, A187, A188, A189, A192, A196, A197, A198, A200, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, A223, A226, A228, A229, A230, A231, A232, A233.

Plutella xylostella (Diamond back moth)

24-well microtiter plates with artificial diet were treated with aqueous test solutions prepared from 10'000 ppm DMSO stock solutions by pipetting. After drying, the plates were infested with L2 larvae (10 to 15 per well). The samples were assessed for mortality and growth inhibition in comparison to untreated samples 5 days after infestation.

The following compounds gave an effect of at least 80% in at least one of the two categories (mortality or growth inhibition) at an application rate of 200 ppm:

A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17, A18, A19, A20, A21, A22, A23, A24, A25, A27, A28, A29, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43, A44, A45, A46, A48, A49, A50, A51, A52, A53, A54, A55, A56, A57, A58, A59, A60, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, A98, A99, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A192, A193, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, A221, A222, A223, A224, A225, A226, A227, A228, A229, A230, A231, A232, A233, A236, A237, A239.

Spodoptera littoralis (Egyptian cotton leaf worm)

Cotton leaf discs were placed onto agar in 24-well microtiter plates and sprayed with aqueous test

solutions prepared from 10'000 ppm DMSO stock solutions. After drying the leaf discs were infested with five L1 larvae. The samples were assessed for mortality, anti-feeding effect, and growth inhibition in comparison to untreated samples 3 days after infestation. Control of *Spodoptera littoralis* by a test sample is given when at least one of the categories mortality, anti-feedant effect, and growth inhibition is higher than the untreated sample.

The following compounds resulted in at least 80% control at an application rate of 200 ppm:

A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17, A18, A19, A20, A21, A22, A23, A24, A25, A27, A28, A29, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43, A44, A45, A46, A47, A48, A49, A50, A51, A52, A53, A54, A55, A56, A57, A58, A59, A60, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, A98, A99, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A175, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A192, A193, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, A221, A222, A223, A224, A225, A226, A227, A228, A229, A230, A231, A232, A233, A236, A237, A238, A239

40

Spodoptera littoralis (Egyptian cotton leaf worm)

Test compounds were applied by pipette from 10'000 ppm DMSO stock solutions into 24-well plates and mixed with agar. Lettuce seeds were placed onto the agar and the multi well plate was closed by another plate which contained also agar. After 7 days the compound was absorbed by the roots and the lettuce grew into the lid plate. The lettuce leaves were then cut off into the lid plate. Spodoptera eggs were

5 pipetted through a plastic stencil onto a humid gel blotting paper and the lid plate was closed with it. The samples were assessed for mortality, anti-feedant effect and growth inhibition in comparison to untreated samples 6 days after infestation.

The following compounds gave an effect of at least 80% in at least one of the three categories (mortality, anti-feeding, or growth inhibition) at a test rate of 12.5 ppm:

10 A1, A5, A6, A8, A10, A11, A12, A13, A16, A19, A20, A21, A22, A23, A24, A25, A27, A28, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A44, A45, A48, A52, A57, A59, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A72, A73, A75, A76, A78, A81, A82, A87, A90, A91, A95, A97, A98, A105, A107, A108, A109, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A125, A126, A127, A128, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140,

15 A141, A142, A143, A144, A145, A146, A147, A155, A158, A159, A160, A236.

Tetranychus urticae (Two-spotted spider mite):Feeding/contact activity

Bean leaf discs on agar in 24-well microtiter plates were sprayed with aqueous test solutions prepared from 10'000 ppm DMSO stock solutions. After drying the leaf discs were infested with a mite population of

20 mixed ages. The samples were assessed for mortality on mixed population (mobile stages) 8 days after infestation.

The following compounds resulted in at least 80% mortality at an application rate of 200 ppm:

A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17, A18, A19, A20, A21, A22, A23, A24, A25, A27, A28, A29, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43,

25 A44, A45, A46, A48, A49, A50, A51, A52, A53, A54, A55, A57, A58, A59, A60, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, A98, A99, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135,

30 A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, A170, A171, A172, A173, A174, A176, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A192, A193, A194, A195, A196, A197, A198, A199, A200, A201, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218,

35 A219, A220, A221, A222, A223, A224, A225, A226, A227, A228, A229, A230, A231, A232, A233, A236.

Thrips tabaci (Onion thrips) Feeding/Contact activity

Sunflower leaf discs were placed on agar in 24-well microtiter plates and sprayed with aqueous test solutions prepared from 10'000 ppm DMSO stock solutions. After drying the leaf discs were infested with


40 a thrips population of mixed ages. The samples were assessed for mortality 6 days after infestation.

The following compounds resulted in at least 80% mortality at an application rate of 200 ppm:

A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17, A18, A19, A20, A21, A22, A23, A24, A25, A27, A28, A29, A30, A31, A32, A33, A34, A35, A36, A37, A38, A39, A40, A41, A42, A43, A44, A45, A46, A48, A49, A50, A51, A52, A55, A57, A58, A59, A60, A61, A62, A63, A64, A65, A66, A67, A68, A69, A70, A71, A72, A73, A74, A75, A76, A77, A78, A79, A80, A81, A82, A83, A84, A85, A86, A87, 5 A88, A89, A90, A91, A92, A93, A94, A95, A96, A97, A98, A99, A100, A101, A102, A103, A104, A105, A106, A107, A108, A109, A110, A111, A112, A113, A114, A115, A116, A117, A118, A119, A120, A121, A122, A123, A124, A125, A126, A127, A128, A129, A130, A131, A132, A133, A134, A135, A136, A137, A138, A139, A140, A141, A142, A143, A144, A145, A146, A147, A148, A149, A150, A151, A152, A153, A154, A155, A156, A157, A158, A159, A160, A161, A162, A163, A164, A165, A166, A167, A168, A169, 10 A170, A171, A172, A173, A174, A176, A177, A178, A179, A180, A181, A182, A183, A184, A185, A186, A187, A188, A189, A190, A192, A193, A194, A195, A196, A197, A198, A199, A200, A202, A203, A204, A205, A206, A207, A208, A209, A210, A211, A212, A213, A214, A215, A216, A217, A218, A219, A220, A221, A223, A224, A225, A226, A227, A228, A229, A230, A231, A232, A233, A236, A239.

CLAIMS

1. A compound of formula (I)



5 wherein

A^1 , A^2 , A^3 and A^4 are independently of one another C-H, C- R^5 , or nitrogen;

R^1 is hydrogen, C₁-C₈alkyl, C₁-C₈alkylcarbonyl-, C₁-C₈alkoxy, C₁-C₈alkoxy-C₁-C₈alkyl or C₁-C₈alkoxycarbonyl-;

R^2 is C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a} , C₁-C₈haloalkyl, C₁-C₈haloalkyl substituted by one to three R^{6a} , C₃-C₈cycloalkyl, C₃-C₈cycloalkyl substituted by one to three R^{6b} , C₃-C₈cycloalkyl where one carbon atom is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈alkyl, C₃-C₈cycloalkyl-C₁-C₈alkyl where one carbon atom in the cycloalkyl group is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a} , C₂-C₈haloalkenyl, C₂-C₈haloalkenyl substituted by one to three R^{6a} , C₂-C₈alkynyl, C₂-C₈haloalkynyl, phenyl, phenyl substituted by one to three R^7 , phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R^7 , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^7 , 5-6 membered heteroaryl-C₁-C₄alkyl, 5-6 membered heteroaryl-C₁-C₄alkyl wherein the heteroaryl moiety is substituted by one to three R^7 , -N(R^8)(R^9), halogen or -OR¹⁰;

R^3 is C₁-C₈haloalkyl;

R^4 is aryl, aryl substituted by one to three R^7 , heteroaryl or heteroaryl substituted by one to three R^7 ;

R^5 is independently halogen, cyano, nitro, C₁-C₈alkyl, C₃-C₈cycloalkyl, C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈haloalkenyl, C₂-C₈alkynyl, C₂-C₈haloalkynyl, C₁-C₈alkoxy, C₁-C₈haloalkoxy, or C₁-C₈alkoxycarbonyl-, or two R^5 on adjacent carbon atoms together form a -CH=CH-CH=CH- bridge or a -N=CH-CH=CH- bridge;

R^{6a} is independently cyano, nitro, amino, C₁-C₈alkylamino, N,N-C₁-C₈dialkylamino, hydroxy, C₁-C₈alkoxy,

25 or C₁-C₈haloalkoxy;

R^{6b} is independently halogen, cyano, nitro, oxo, C₁-C₈alkyl, C₁-C₈haloalkyl, amino, C₁-C₈alkylamino, N,N-C₁-C₈dialkylamino, hydroxyl, C₁-C₈alkoxy, C₁-C₈haloalkoxy, phenyl, phenyl substituted by one to three R^7 , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^7 ;

R^7 is independently halogen, cyano, nitro, C₁-C₈alkyl, C₁-C₈haloalkyl, C₁-C₈alkoxy, C₁-C₈haloalkoxy;

30 R^8 and R^9 are independently hydrogen, cyano, cyano-C₁-C₈alkyl, C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a} , C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a} , C₁-C₈alkoxy, C₁-C₈haloalkoxy, C₁-C₈haloalkoxy substituted by one to three R^{6a} , C₁-C₈alkoxy substituted by one to three R^{6a} , C₁-C₈haloalkyl, C₁-C₈haloalkyl substituted by one to three R^{6a} , C₃-C₈cycloalkyl, C₃-C₈cycloalkyl substituted by one to three R^{6b} , C₃-C₈cycloalkyl where one carbon atom is replaced by O, S, S(O) or SO₂,

C₃-C₈cycloalkyl-C₁-C₈alkyl, C₃-C₈cycloalkyl-C₁-C₈alkyl where one carbon atom in the cycloalkyl group is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a}, C₂-C₈haloalkenyl, C₂-C₈haloalkenyl substituted by one to three R^{6a}, C₂-C₈alkynyl, C₂-C₈haloalkynyl, phenyl, phenyl substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl

5 wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷, 5-6 membered heteroaryl-C₁-C₄alkyl, 5-6 membered heteroaryl-C₁-C₄alkyl wherein the heteroaryl moiety is substituted by one to three R⁷, -S(O)R¹⁰, -S(O)₂R¹⁰, COR¹⁰, COOR¹⁰, or R⁸ and R⁹ together with the nitrogen atom can be linked through a C₃-C₈alkylene chain, a C₃-C₈alkylene chain substituted by one to three R^{6b} or a C₃-C₈alkylene chain, where one carbon atom is
10 replaced by O, S, S(O) or SO₂;

R¹⁰ is hydrogen, cyano-C₁-C₈alkyl, C₁-C₈alkyl, C₁-C₈alkyl substituted by one to three R^{6a}, C₁-C₈haloalkyl, C₁-C₈haloalkyl substituted by one to three R^{6a}, C₃-C₈cycloalkyl, C₃-C₈cycloalkyl substituted by one to three R^{6b}, C₃-C₈cycloalkyl where one carbon atom is replaced by O, S, S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈alkyl, C₃-C₈cycloalkyl-C₁-C₈alkyl where one carbon atom in the cycloalkyl group is replaced by O, S,
15 S(O) or SO₂, C₃-C₈cycloalkyl-C₁-C₈haloalkyl, C₂-C₈alkenyl, C₂-C₈alkenyl substituted by one to three R^{6a}, C₂-C₈haloalkenyl, C₂-C₈haloalkenyl substituted by one to three R^{6a}, C₂-C₈alkynyl, C₂-C₈haloalkynyl, phenyl, phenyl substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷, 5-6 membered heteroaryl-C₁-C₄alkyl or 5-6 membered heteroaryl-C₁-C₄alkyl wherein
20 the heteroaryl moiety is substituted by one to three R⁷;

n is 1 or 2;

or a tautomer, isomer, enantiomer, salt or *N*-oxide thereof.

2. The compound according to claim 1, wherein A¹ is CR⁵ and A², A³ and A⁴ are each CH, wherein
25 R⁵ is as defined in claim 1.

3. The compound according to either claim 1 or claim 2, wherein R¹ is hydrogen, C₁-C₈alkyl, C₁-C₈alkylcarbonyl- or C₁-C₈alkoxycarbonyl-.

30 4. The compound according to any one of claims 1 to 3, wherein R² is C₁-C₈alkyl or C₁-C₈alkyl substituted by one to three R^{6a}, C₁-C₈haloalkyl or C₁-C₈haloalkyl substituted by one to three R^{6a}, C₃-C₈cycloalkyl or C₃-C₈cycloalkyl substituted by one to three R^{6b}, phenyl, phenyl substituted by one to three R⁷, phenyl-C₁-C₄alkyl, phenyl-C₁-C₄alkyl wherein the phenyl moiety is substituted by one to three R⁷, 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R⁷, halogen or -N(R⁸)(R⁹)
35 wherein R⁸, R⁹, R^{6a} and R^{6b} are as defined in claim 1.

5. The compound according to any one of claims 1 to 4, wherein R³ is C₁-C₄haloalkyl.

6. The compound according to any one of claims 1 to 5, wherein R⁴ is phenyl or phenyl substituted
40 by one to three R⁷; wherein R⁷ is independently halogen, cyano, C₁-C₈alkyl, C₁-C₈haloalkyl, C₁-C₈alkoxy, or C₁-C₈haloalkoxy.


7. The compound according to any one of claims 1 to 6, wherein A^1 is CR^5 and A^2 , A^3 and A^4 are each CH; R^1 is hydrogen, methyl, ethyl, methylcarbonyl-, or methoxycarbonyl; R^2 is C_1 - C_8 alkyl, C_1 - C_8 alkyl substituted by one to three R^{6a} , C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, di- C_1 - C_8 alkylamino, C_1 - C_4 alkylamino, fluoro, aryl, aryl substituted by one to three R^{6b} , 5-6 membered heteroaryl or 5-6 membered heteroaryl substituted by one to three R^{6b} ; R^3 is C_1 - C_4 haloalkyl; R^4 is aryl or aryl substituted by one to three R^{6b} ; and n is 2; wherein R^5 is halogen or C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, or C_2 - C_8 alkenyl; R^{6a} is independently cyano, halogen, nitro, C_1 - C_4 alkoxy, or C_1 - C_4 haloalkoxy; and R^{6b} is independently halogen, cyano, nitro, C_1 - C_4 alkyl, or C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy, or C_1 - C_4 haloalkoxy.

10

8. The compound according to any one of claims 1 to 7, wherein A^1 is CR^5 and A^2 , A^3 and A^4 are each CH; R^1 is hydrogen; R^2 is C_1 - C_4 alkyl, C_1 - C_4 alkyl substituted by one to three R^{6a} , C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_3 - C_5 cycloalkyl, C_1 - C_4 haloalkyl, di- C_1 - C_4 alkylamino, C_1 - C_4 alkylamino, fluoro, 1-3 halo-substituted phenyl, or 5-6 membered heteroaryl; R^3 is chlorodifluoromethyl or trifluoromethyl; R^4 is 3,5-bis-(trifluoromethyl)-phenyl, 3-chloro-5-trifluoromethyl-phenyl, 3-bromo-5-trifluoromethyl-phenyl, 3,5-dibromo-phenyl, 3,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3-trifluoromethyl-phenyl, 4-bromo-3,5-dichlorophenyl, 3,5-dichloro-4-fluoro phenyl or 3,4,5-trichloro-phenyl; and n is 2; wherein R^5 is bromo, chloro, fluoro, cyano, nitro, methyl, ethyl, trifluoromethyl, cyclopropyl, or vinyl; R^{6a} is independently fluoro, cyano, nitro, methoxy, difluoromethoxy or trifluoromethoxy; and R^{6b} is independently bromo, chloro, fluoro, cyano, nitro methyl, ethyl, trifluoromethyl, methoxy, difluoromethoxy, trifluoromethoxy.

15

9. A compound of formula (I) according to claim 1 represented by the compounds of formula (Ib)



wherein

25 R^a is hydrogen, halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, or C_1 - C_8 haloalkoxy;
 R^b is hydrogen, halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, or C_1 - C_8 haloalkoxy;
 R^c is hydrogen, halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, or C_1 - C_8 haloalkoxy;
 R^5 is halogen, cyano, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, C_2 - C_8 alkenyl;
 R^d is $S(O)_2$ - R^2 ;
 30 R^2 is C_1 - C_8 alkyl or C_1 - C_8 alkyl substituted by one to three R^{6a} , C_1 - C_8 haloalkyl or C_1 - C_8 haloalkyl substituted by one to three R^{6a} , C_3 - C_8 cycloalkyl or C_3 - C_8 cycloalkyl substituted by one to three R^{6b} , phenyl, phenyl substituted by one to three R^7 , phenyl- C_1 - C_4 alkyl, phenyl- C_1 - C_4 alkyl wherein the phenyl moiety is substituted by one to three R^7 , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^7 , $-N(R^8)(R^9)$, C_1 - C_4 alkylamino, fluoro;
 35 R^5 is independently halogen, cyano, nitro, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, C_1 - C_8 haloalkyl, C_2 - C_8 alkenyl;

R^{6a} is independently C_1 - C_8 alkoxy, or C_1 - C_8 haloalkoxy;


R^{6b} is independently oxo, amino, C_1 - C_8 alkylamino, N,N - C_1 - C_8 dialkylamino, C_1 - C_8 alkoxy, C_1 - C_8 haloalkoxy, phenyl, phenyl substituted by one to three R^7 , 5-6 membered heteroaryl or 5-6 membered heteroaryl substituted by one to three R^7 ;

5 R^7 is independently halogen, cyano, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, C_1 - C_8 alkoxy, or C_1 - C_8 haloalkoxy;


R^8 and R^9 are independently hydrogen, cyano- C_1 - C_8 alkyl, C_1 - C_8 alkyl, C_1 - C_8 alkyl substituted by one to three R^{6a} , C_2 - C_8 alkenyl or C_2 - C_8 alkenyl substituted by one to three R^{6a} , phenyl- C_1 - C_4 alkyl, phenyl- C_1 - C_4 alkyl wherein the phenyl moiety is substituted by one to three R^7 , $-S(O)_2R^{10}$;

10 R^{10} is hydrogen, cyano- C_1 - C_8 alkyl, C_1 - C_8 alkyl or C_1 - C_8 alkyl substituted by one to three R^{6a} , phenyl, phenyl substituted by one to three R^7 , 5-6 membered heteroaryl, 5-6 membered heteroaryl substituted by one to three R^7 .


10. A compound of formula (Int-I)



15 wherein A^1 , A^2 , A^3 , A^4 , R^1 and R^2 , are as defined for a compound of formula (I) according to any one of claims 1 to 9 and X^B is a leaving group, or X^B is cyano, formyl, $CH=N-OH$ or acetyl; or a tautomer, isomer, enantiomer, salt or N -oxide thereof; or
a compound of formula (Int-II)



20 wherein A^1 , A^2 , A^3 , A^4 , R^1 and R^2 , are as defined for a compound of formula (I) according to any one of claims 1 to 9 and X^C is CH_2 -halogen, $CH=C(R^3)R^4$, or $CH_2C(OH)(R^3)R^4$ wherein R^3 and R^4 are as defined for a compound of formula (I) according to any one of claims 1 to 9; or a tautomer, isomer, enantiomer, salt or N -oxide thereof; or
a compound of formula (Int-III)



wherein R^1 and R^2 , are as defined for a compound of formula (I) according to any one of claims 1 to 9; or a tautomer, isomer, enantiomer, salt or *N*-oxide thereof.

- 5 11. A method of combating and/or controlling an invertebrate animal pest which comprises applying to the pest, to a locus of the pest, or to a plant susceptible to attack by the pest a pesticidally effective amount of a compound of formula (I) as defined in any one of claims 1 to 9.
12. A pesticidal composition, which comprises at least one compound of formula (I) according to any of
- 10 the claims 1 to 9 or where appropriate, a tautomer thereof, in each case in free form or in agrochemically utilizable salt form, as active ingredient.
13. A method for controlling pests, which comprises applying a composition according to claim 12 to the pests or their environment with the exception of a method for treatment of the human or animal body by
- 15 surgery or therapy and diagnostic methods practised on the human or animal body.
14. A method for the protection of plant propagation material from the attack by pests, which comprises treating the propagation material or the site, where the propagation material is planted, with a composition according to claim 12.
- 20 15. Plant propagation material treated with the pesticidal composition described in claim 12.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2016/072588

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D413/14 A01N43/80
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2011/067272 A1 (SYNGENTA PARTICIPATIONS AG [CH]; CASSAYRE JEROME YVES [CH]; RENOLD PET) 9 June 2011 (2011-06-09) cited in the application claim 1	1-15
X	----- US 2011/144334 A1 (MITA TAKESHI [JP] ET AL) 16 June 2011 (2011-06-16) claim 1	1-15
X	----- US 2007/066617 A1 (MITA TAKESHI [JP] ET AL) 22 March 2007 (2007-03-22) claim 1	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

10 November 2016

Date of mailing of the international search report

22/11/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Panday, Narendra

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2016/072588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011067272	A1	09-06-2011	AR 079224 A1 04-01-2012
		AU 2010326766 A1 07-06-2012	
		BR 112012013085 A2 08-09-2015	
		CA 2780522 A1 09-06-2011	
		CN 102639529 A 15-08-2012	
		CO 6541622 A2 16-10-2012	
		CR 20120299 A 04-07-2012	
		EA 201200826 A1 30-01-2013	
		EC SP12011940 A 31-07-2012	
		EP 2507230 A1 10-10-2012	
		JP 5980119 B2 31-08-2016	
		JP 2013512274 A 11-04-2013	
		KR 20120102066 A 17-09-2012	
		MA 33782 B1 01-11-2012	
		MY 157849 A 29-07-2016	
		NZ 600036 A 31-05-2013	
		TW 201130417 A 16-09-2011	
		US 2012238517 A1 20-09-2012	
		US 2014228577 A1 14-08-2014	
		UY 33070 A 29-07-2011	
		WO 2011067272 A1 09-06-2011	
		ZA 201203587 B 29-10-2014	
US 2011144334	A1	16-06-2011	AU 2006285613 A1 08-03-2007
			BR PI0617076 A2 04-08-2009
			CA 2621228 A1 08-03-2007
			EP 1932836 A1 18-06-2008
			ES 2443690 T3 20-02-2014
			JP 5293921 B2 18-09-2013
			JP 5594490 B2 24-09-2014
			JP 2013040184 A 28-02-2013
			KR 20080049091 A 03-06-2008
			TW 200804315 A 16-01-2008
			US 7951828 B1 31-05-2011
			US 2011144334 A1 16-06-2011
			US 2014135496 A1 15-05-2014
			WO 2007026965 A1 08-03-2007
US 2007066617	A1	22-03-2007	AU 2005219788 A1 15-09-2005
			AU 2010212450 A1 09-09-2010
			BR PI0508140 A 24-07-2007
			CA 2558848 A1 15-09-2005
			CN 101768129 A 07-07-2010
			CN 102532048 A 04-07-2012
			CN 102558082 A 11-07-2012
			DK 1731512 T3 05-01-2015
			EA 200601639 A1 27-02-2007
			EA 200900197 A1 30-06-2009
			EP 1731512 A1 13-12-2006
			ES 2526614 T3 13-01-2015
			HR P20141241 T1 13-02-2015
			IL 177949 A 28-02-2013
			JP 4883296 B2 22-02-2012
			JP 5321856 B2 23-10-2013
			JP 2012006944 A 12-01-2012
			KR 20070010145 A 22-01-2007
			KR 20110132638 A 08-12-2011
			LU 92672 I2 02-11-2015

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2016/072588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		NL 300721 I1	29-12-2015
		PT 1731512 E	24-11-2014
		SI 1731512 T1	30-01-2015
		TW I355896 B	11-01-2012
		US 2007066617 A1	22-03-2007
		US 2009312330 A1	17-12-2009
		US 2011251398 A1	13-10-2011
		US 2012172615 A1	05-07-2012
		US 2013296559 A1	07-11-2013
		US 2015105398 A1	16-04-2015
		WO 2005085216 A1	15-09-2005
