



(51) International Patent Classification:

C07D 487/04 (2006.01) C07K 1/00 (2006.01)
C07K 7/64 (2006.01) A61K 38/12 (2006.01)
A61K 47/62 (2017.01) A61P 35/00 (2006.01)

Published:

— with international search report (Art. 21(3))

(21) International Application Number:

PCT/EP2018/071268

(22) International Filing Date:

06 August 2018 (06.08.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17185182.7 07 August 2017 (07.08.2017) EP

(71) Applicant: HEIDELBERG PHARMA GMBH [DE/DE];
Schriesheimer Strasse 101, 68526 Ladenburg (DE).

(72) Inventors: LUTZ, Christian; Am Feldrain 36, 69469
Weinheim (DE). SIMON, Werner; In der Liewiese
29, 55595 Hüffelsheim (DE). WERNER-SIMON, Su-
sanne; In der Liewiese 29, 55595 Hüffelsheim (DE).
MÜLLER, Christoph; Mumbacherstr. 38, 69488 Birke-
nau (DE). HECHLER, Thorsten; Mühlbergweg 24, 64686
Lautertal (DE). KULKE, Michael; Egellstr. 17, 67071
Ludwigshafen am Rhein (DE).

(74) Agent: WALLINGER RICKER SCHLOTTER TOST-
MANN; Patent- und Rechtsanwälte, Partnerschaft mbB,
Zweibrückenstraße 5-7, 80331 Munich (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(54) Title: NOVEL METHOD FOR SYNTHESIZING AMANITINS

(57) Abstract: The invention relates to novel methods for synthesizing amanitin derivatives having a hydroxy group attached to the central tryptophan moiety. The invention furthermore relates to novel amanitin derivatives having a hydroxy group attached to position 4', 5' or 7' of the central tryptophan moiety, novel conjugates of such amanitin derivatives, and pharmaceutical compositions comprising such conjugates.

NOVEL METHOD FOR SYNTHESIZING AMANITINS

FIELD OF THE INVENTION

[001] The invention relates to novel methods for synthesizing amanitin derivatives having a hydroxy group attached to the central tryptophan moiety. The invention furthermore relates to novel amanitin derivatives having a hydroxy group attached to position 4', 5' or 7' of the central tryptophan moiety, novel conjugates of such amanitin derivatives, and pharmaceutical compositions comprising such conjugates.

BACKGROUND OF THE INVENTION

[002] Amatoxins are cyclic peptides composed of 8 amino acids that are found in *Amanita phalloides* mushrooms (see **Fig. 1**). Amatoxins specifically inhibit the DNA-dependent RNA polymerase II of mammalian cells, and thereby also the transcription and protein biosynthesis of the affected cells. Inhibition of transcription in a cell causes stop of growth and proliferation. Though not covalently bound, the complex between amanitin and RNA-polymerase II is very tight ($K_D = 3$ nM). Dissociation of amanitin from the enzyme is a very slow process, thus making recovery of an affected cell unlikely. When the inhibition of transcription lasts too long, the cell will undergo programmed cell death (apoptosis).

[003] The use of amatoxins as cytotoxic moieties for tumour therapy had already been explored in 1981 by coupling an anti-Thy 1.2 antibody to α -amanitin using a linker attached to the indole ring of Trp (amino acid 4; see **Fig. 1**) via diazotation (Davis & Preston, Science 213 (1981) 1385-1388). Davis & Preston identified the site of attachment as position 7'. Morris & Venton demonstrated as well that substitution at position 7' results in a derivative, which maintains cytotoxic activity (Morris & Venton, Int. J. Peptide Protein Res. 21 (1983) 419-430).

[004] Patent application EP 1 859 811 A1 (published November 28, 2007) described conjugates, in which the γ C-atom of amatoxin amino acid 1 of β -amanitin was directly coupled, i.e. without a linker structure, to albumin or to monoclonal antibody HEA125, OKT3, or PA-1. Furthermore, the inhibitory effect of these conjugates on the proliferation of breast cancer cells (MCF-7), Burkitt's lymphoma cells (Raji) and T-lymphoma cells (Jurkat) was shown. The use of linkers was suggested, including linkers comprising elements such as amide, ester, ether, thioether, disulfide, urea, thiourea, hydrocarbon moieties and the like, but no such constructs were actually shown, and no more details, such as attachment sites on the amatoxins, were provided.

[005] Patent applications WO 2010/115629 and WO 2010/115630 (both published October 14, 2010) describe conjugates, where antibodies, such as anti-EpCAM antibodies such as humanized antibody huHEA125, are coupled to amatoxins via (i) the γ C-atom of amatoxin amino acid 1, (ii) the 6' C-atom of amatoxin amino acid 4, or (iii) via the δ C-atom of amatoxin amino acid 3, in each case either directly or via a linker between the antibody and the amatoxins. The suggested linkers comprise elements such as amide, ester, ether, thioether, disulfide, urea, thiourea, hydrocarbon moieties and the like. Furthermore, the inhibitory effects of these conjugates on the proliferation of breast cancer cells (cell line MCF-7), pancreatic carcinoma (cell line Capan-1), colon cancer (cell line Colo205), and cholangiocarcinoma (cell line OZ) were shown.

[006] Patent application WO 2012/119787 describes that target-binding moieties can be attached to amatoxins via linkers at additional attachment sites on tryptophan amino acid 4, namely positions 1'-N, without interference with the interaction of such amatoxins with their target, the DNA-dependent RNA polymerase II of mammalian cells.

[007] Patent application WO 2014/009025 describes the total synthesis of amanitin derivatives using a novel synthon for γ,δ -dihydroxyisoleucine as one of the starting materials. Furthermore, patent application PCT/EP2016/078984 [published as WO 2017/089607] describes the total synthesis of derivatives of γ - and ϵ -amanitin using readily available (2S,3R,4S)-L-4-hydroxyisoleucine as one of the starting

materials. However, these approaches, and all other fully or partially synthetic approaches pursued so far, are incorporating the central tryptophan moiety into the amatoxin core by using the method according to Savige-Fontana. In this method, cis-2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole ("Hpi") is incorporated into a linear amanitin precursor structure. In an acid-catalyzed Hpi-cysteine coupling reaction, the amanitin ring system with central tryptophan moiety is formed.

[008] Hpi, however, does not possess any functional moiety attached to the phenyl ring of Hpi's indole moiety. Thus, the amanitin products resulting from the method according to Savige-Fontana contain a central tryptophan moiety without any further substituents. Naturally occurring α -, β -, γ , and ϵ -amanitins, however, contain a central 6'-hydroxy-tryptophan moiety, and that 6'-hydroxy group has successfully been used as functional group for the functionalization of amanitins, e.g. by attaching targeting moieties, either directly or via linkers (see, for example, WO 2010/115629, WO 2010/115630, and WO 2012/041504). Thus, in the case of synthetic amanitins, functionalization had to be done via (i) the γ C-atom of amatoxin amino acid 1, or (ii) via the δ C-atom of amatoxin amino acid 3, as described above, or via the nitrogen atom of the tryptophan moiety, as described in WO 2012/119787.

[009] Hpi can be obtained by reacting tryptophan with peracetic acid or photochemically with singlet oxygen. However, no derivatives with substituents attached to the phenyl ring of Hpi's indole moiety have been described so far.

[0010] While the use of fully synthetic routes to amatoxins may offer an option for the supply of larger quantities of amatoxins required for therapeutic uses, and may offer the construction of a variety of novel amatoxin variants by using appropriate starting materials as building blocks, the approaches pursued in the past had been limited by the fact that the native structure α -, β -, γ , and/or ϵ -amanitin could not yet be obtained, since the 6'-hydroxy moiety attached to the core tryptophan moiety in these amanitins could not be incorporated. Thus, options for functionalizing synthetic amanitins have been rather limited so far. Furthermore, it would be highly desirable to expand the range of options available for the functionalization, since factors such as steric hindrance and reactivity might have a strong impact on the

reactivity, biological activity and/or stability of synthetic amanitins and of conjugates thereof. However, stability and efficacy of conjugates comprising highly toxic amanitins are of utmost importance for the envisaged use as therapeutic molecules for administration to human beings.


OBJECT OF THE INVENTION

[0011] Thus, there was still a great need for a cost efficient and robust way of synthesizing amatoxins with a hydroxy group attached to the phenyl ring of the central tryptophan moiety. In particular, there is a strong need for identifying starting materials that could be used in the established Savige-Fontana reaction and that are set up in a way that they can cause the incorporation of such hydroxy groups.

SUMMARY OF THE INVENTION

[0012] The present invention is based on the unexpected observation that variants of Hpi can be synthesized that permit the introduction of hydroxyl groups during the synthesis of amanitin derivatives.

[0013] Thus, in one aspect the present invention relates to a hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole according to Formula I



wherein R1 is selected from alkyl, aryl, heteroaryl, substituted alkyl, substituted aryl, and substituted heteroaryl;

P₁ is hydrogen or a protecting group, particularly a protecting group selected from Boc, PhCH₂OCO-, CH₂=CHCH₂O-CO-, and trityl;

P₂ is hydrogen or a protecting group, particularly a protecting group, particularly a protecting group selected from Boc, PhCH₂OCO-, CH₂=CHCH₂O-CO-, and trityl; and R₂ is selected from OH, OR₁, and a polypeptide chain consisting of 1-7 amino acid residues.

[0014] In a second aspect, the present invention relates to a method for the synthesis of a linear precursor comprising eight amino acid residues of an amanitin derivative comprising a hydroxylated tryptophan moiety, comprising the step of using a hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole of any one of the present invention in the peptide synthesis of said precursor.

[0015] In a third aspect, the invention relates to a method for the synthesis of an amanitin derivative comprising a hydroxylated tryptophan moiety, comprising the steps of.

- (i) causing or allowing the formation of a bond between the cysteine residue and the tryptophan moiety of the linear precursor of the present invention; and
- (ii) causing or allowing the formation of said amanitin derivative by reacting the N-terminus of the linear precursor of the present invention with the C-terminus of said precursor.

[0016] In a fourth aspect, the invention relates to an amanitin derivative comprising a hydroxylated tryptophan moiety, which is selected from (i) S-desoxy-4'-hydroxy-amanin, 4'-hydroxy-amanin, S-desoxy-5'-hydroxy-amanin, 5'-hydroxy-amanin, S-desoxy-7'-hydroxy-amanin, 7'-hydroxy-amanin, (ii) S-desoxy-4'-hydroxy-amaninamide, 4'-hydroxy-amaninamide, S-desoxy-5'-hydroxy-amaninamide, 5'-hydroxy-amaninamide, S-desoxy-7'-hydroxy-amaninamide, and 7'-hydroxy-amaninamide, (iii) a derivative of the amanitin according to (i), wherein the free carboxylic acid moiety of amino acid 1 is converted to a carboxylic ester –C(=O)OR₁ or to a moiety –C(=O)NH-OR₁, wherein R₁ is selected from alkyl, aryl, heteroaryl, substituted alkyl, substituted aryl, and substituted heteroaryl.

[0017] In a fifth aspect, the invention relates to a conjugate comprising (a) amanitin derivative comprising a hydroxylated tryptophan moiety of the present invention; (b) a target-binding moiety; and (c) optionally a linker linking said amanitin derivative and said target-binding moiety.

[0018] In a sixth aspect, the invention relates to a pharmaceutical composition comprising the amanitin of the present invention or the conjugate of the present invention.

[0019] In a seventh aspect, the invention relates to an amanitin derivative of the present invention, the conjugate of the present invention, or the pharmaceutical composition of the present invention for use in the treatment of cancer in a patient, particularly wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, cholangiocarcinoma, colorectal cancer, lung cancer, prostate cancer, ovarian cancer, prostate cancer, stomach cancer, kidney cancer, malignant melanoma, leukemia, and malignant lymphoma.

[0020] In an eighth aspect, the invention relates to a construct comprising (a) an amanitin derivative of the present invention; and (b) a linker moiety carrying a reactive group Y for linking said amanitin derivative to a target-binding moiety.

BRIEF DESCRIPTION OF THE DRAWING

[0021] **Fig. 1** shows the structural formulae of different amatoxins. The numbers in bold type (1 to 8) designate the standard numbering of the eight amino acids forming the amatoxin. The standard designations of the atoms in amino acids 1, 3 and 4 are also shown (Greek letters α to γ , Greek letters α to δ , and numbers from 1' to 7', respectively).

[0022] **Fig. 2** shows the structure of compound S-deoxy- α -amanitin (HDP 30.0735) and of α -amanitin.

[0023] **Fig. 3** shows that Hpi (HDP 30.0079) is generated in a one-step synthesis starting from Boc-L-tryptophan.

[0024] **Fig. 4** shows that compound S-deoxy- α -amanitin (HDP 30.0735) is generated by solid-phase peptide synthesis incorporating Hpi derivative HDP 30.2555.

[0025] **Fig. 5** shows the cytotoxicity of S-deoxy- α -amanitin (HDP 30.0735) and S-desoxy-5'-hydroxy-amaninamide HDP 30.2548 on HEK293 cells and on HEK293 OATP1B3 cells.

[0026] **Fig. 6** shows a construct based on S-deoxy- α -amanitin (HDP 30.0735) with a cleavable linker attached to the 6'-hydroxy group and a terminal maleimide group as an example of a reactive group Y for linking said construct to a target-binding moiety.

[0027] **Fig. 7** shows an alternative construct based on S-deoxy- α -amanitin (HDP 30.0735) with a cleavable linker attached to the carboxyl group of amino acid residue 1 and a terminal maleimide group as an example of a reactive group Y for linking said construct to a target-binding moiety.

[0028] **Fig. 8** shows that hydroxylated Hpi derivative HDP 30.2536 is generated in a multi-step synthesis starting from 5'-hydroxy-L-tryptophan.

[0029] **Fig. 9** shows that compound the amanitin precursor HDP 30.2544 is generated by solid-phase peptide synthesis incorporating Hpi derivative HDP 30.2536.

[0030] **Fig. 10** shows that compound the amanitin derivative HDP 30.2546 is generated by ring closure from the amanitin precursor HDP 30.2544.

[0031] **Fig. 11** shows that the amanitin derivative HDP 30.2548 (S-desoxy-5'-hydroxy-amaninamide) is generated by removal of protecting groups from amanitin derivative HDP 30.2546.

[0032] **Fig. 12** shows a construct based on amanitin derivative HDP 30.2548 (S-desoxy-5'-hydroxy-amaninamide) with a cleavable linker attached to the 5'-hydroxy group and a terminal maleimide group as an example of a reactive group Y for linking said construct to a target-binding moiety.

[0033] **Fig. 13** shows the cytotoxicity of HDP 30.2347, HDP 30.2371 and HDP 30.2602 ADCs targeting A) HER-2/neu on SKBR-3 cells (HER-2/neu+++) and JIMT-1 cells (HER-2/neu+), B) PSMA on LnCap cells (PSMA+++) and 22rv1 cells (PSMA++) and C) CD19 on Raji cells (CD19+++) and Nalm-6 cells (CD19++) in comparison to HDP 30.1699 ADCs (containing the same cleavable linker as HDP 30.2347, HDP 30.2371 and HDP 30.2602, but alpha-amanitin instead of the above described Amanitin derivatives)

DETAILED DESCRIPTION OF THE INVENTION

[0034] Before the present invention is described in detail below, it is to be understood that this invention is not limited to the particular methodology, protocols and reagents described herein as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art.

[0035] Particularly, the terms used herein are defined as described in "A multilingual glossary of biotechnological terms: (IUPAC Recommendations)", Leuenberger, H.G.W, Nagel, B. and Kölbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland).


[0036] Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer, composition or step or group of integers or steps, while any additional integer, composition or step or group of integers, compositions or steps may optionally be present as well, including embodiments, where no additional integer, composition or step or group of integers, compositions or steps are present. In such latter embodiments, the term "comprising" is used coterminous with "consisting of".

[0037] Several documents are cited throughout the text of this specification. Each of the documents cited herein (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, GenBank Accession Number sequence submissions etc.), whether supra or infra, is hereby incorporated by reference in its entirety to the extent possible under the respective patent law. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

[0038] The present invention will now be further described. In the following passages different aspects of the invention are defined in more detail. Each aspect so defined may be combined with any other aspect or aspects unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous.

[0039] The present invention is based on the unexpected observation that variants of Hpi can be synthesized that permit the introduction of hydroxyl groups during the synthesis of amanitin derivatives

[0040] Thus, in one aspect the present invention relates to a hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole according to Formula I



wherein R1 is selected from alkyl, aryl, heteroaryl, substituted alkyl, substituted aryl, and substituted heteroaryl;

P₁ is hydrogen or a protecting group;

P₂ is hydrogen or a protecting group; and

R2 is selected from OH, OR1, and a polypeptide chain consisting of 1-7 amino acid residues.

[0041] In the context of the present invention, the term “protecting group” refers to a group that is attached to a nitrogen atom in positions 1 or 8 of the central hexahydropyrrolo[2,3-b]indole moiety in order to block the nitrogen atom from reacting with other reactants used to synthesize and/or to further functionalize compounds according to Formula I.. One of ordinary skill in the art is well familiar with the different protecting groups that are available in the art and that can be attached to the corresponding nitrogen atom when needed to protect the nitrogen atom, and that can be cleaved off subsequently, when N-protection is no longer needed. In particular embodiments, the N-protection uses an N-acylating reagent. Thus, in such embodiments, P₁ and/or P₂ are acyl groups. In particular other embodiments, the N-protection uses an N-alkylating reagent. Thus, in such embodiments, P₁ and/or P₂ is an alkyl group.

[0042] In particular embodiments, the protecting group P₁ or P₂, when present, is independently selected from Boc, PhCH₂OCO-, CH₂=CHCH₂O-CO-, and trityl.

[0043] In a particular embodiment, the hydroxy group in position 3a and the hydrogen atom in position 8a are in *cis*-configuration with respect to the functional

group attached to position 2. In another particular embodiment, the hydroxy group in position 3a and the hydrogen atom in position 8a are in *trans*-configuration with respect to the functional group attached to position 2. In a particular embodiment, the amino-substituted derivative according to the present invention is a mixture of compounds with *cis*- and *trans*-configuration.

[0044] In a particular embodiment, the substituent R1-C(=O)-O- is attached to position 4 in Formula I.

[0045] In a particular embodiment, the substituent R1-C(=O)-O- is attached to position 5 in Formula I.

[0046] In a particular embodiment, the substituent R1-C(=O)-O- is attached to position 6 in Formula I.

[0047] In a particular embodiment, the substituent R1-C(=O)-O- is attached to position 7 in Formula I.

[0048] In a second aspect, the present invention relates to a method for the synthesis of a linear precursor comprising eight amino acid residues of an amanitin derivative comprising a hydroxylated tryptophan moiety, comprising the step of using a hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole of the present invention in the peptide synthesis of said precursor.

[0049] In a third aspect, the invention relates to a method for the synthesis of an amanitin derivative comprising a hydroxylated tryptophan moiety, comprising the steps of.

- (i) causing or allowing the formation of a bond between the cysteine residue and the tryptophan moiety of the linear precursor of the present invention; and
- (ii) causing or allowing the formation of said amanitin derivative by reacting the N-terminus of the linear precursor of the present invention with the C-terminus of said precursor.

[0050] In additional aspects, the invention relates to individual amanitin precursors synthesized as shown in the examples, in particular compounds HDP 30.2569, HDP 30.2572 and the solid phase based intermediates synthesized according to [00145].

[0051] In a particular embodiment, the method of the present invention further comprises the oxidation of the sulfur atom of the cysteine moiety to form a sulfoxide or a sulfone, particularly a sulfoxide.

[0052] In a fourth aspect, the invention relates to an amanitin derivative comprising a hydroxylated tryptophan moiety, which is selected from (i) S-desoxy-4'-hydroxy-amanin, 4'-hydroxy-amanin, S-desoxy-5'-hydroxy-amanin, 5'-hydroxy-amanin, S-desoxy-7'-hydroxy-amanin, 7'-hydroxy-amanin, (ii) S-desoxy-4'-hydroxy-amaninamide, 4'-hydroxy-amaninamide, S-desoxy-5'-hydroxy-amaninamide, 5'-hydroxy-amaninamide, S-desoxy-7'-hydroxy-amaninamide, and 7'-hydroxy-amaninamide, (iii) a derivative of the amanitin according to (i), wherein the free carboxylic acid moiety of amino acid 1 is converted to a carboxylic ester – C(=O)OR1 or to a moiety –C(=O)NH-OR1, wherein R1 is selected from alkyl, aryl, heteroaryl, substituted alkyl, substituted aryl, and substituted heteroaryl.

[0053] In a particular embodiment, the amanitin derivative of the present invention is selected from S-desoxy-5'-hydroxy-amanin, 5'-hydroxy-amanin, S-desoxy-5'-hydroxy-amaninamide, and 5'-hydroxy-amaninamide

[0054] In a fifth aspect, the invention relates to a conjugate comprising (a) amanitin derivative comprising a hydroxylated tryptophan moiety of the present invention; (b) a target-binding moiety; and (c) optionally a linker linking said amanitin derivative and said target-binding moiety.

[0055] In a sixth aspect, the invention relates to a pharmaceutical composition comprising the amanitin of the present invention or the conjugate of the present invention.

[0056] In a seventh aspect, the invention relates to an amanitin derivative of the present invention, the conjugate of the present invention, or the pharmaceutical composition of the present invention for use in the treatment of cancer in a patient, particularly wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, cholangiocarcinoma, colorectal cancer, lung cancer, prostate cancer, ovarian cancer, prostate cancer, stomach cancer, kidney cancer, malignant melanoma, leukemia, and malignant lymphoma.

[0057] In an eighth aspect, the invention relates to a construct comprising (a) an amanitin derivative of the present invention; and (b) a linker moiety carrying a reactive group Y for linking said amanitin derivative to a target-binding moiety.

[0058] In the context of the present invention, the term “amanitin” refers to a particular group of amatoxins. In the context of the present invention the term “amatoxin” includes all cyclic peptides composed of 8 amino acids as isolated from the genus *Amanita* and described in Wieland, T. and Faulstich H. (Wieland T, Faulstich H., CRC Crit Rev Biochem. 5 (1978) 185-260). In the context of the present invention, the term “amanitins” refers to bicyclic structure that are based on an aspartic acid or asparagine residue in position 1, a proline residue, particularly a hydroxyproline residue in position 2, an isoleucine, hydroxyisoleucine or dihydroxyisoleucine in position 3, a hydroxytryptophan residue in position 4, glycine residues in positions 5 and 7, an isoleucine residue in position 6, and a cysteine residue in position 8, particularly a derivative of cysteine that is oxidized to a sulfoxide or sulfone derivative (for the numbering and representative examples of amanitins, see Figure 1), and furthermore includes all chemical derivatives thereof; further all semisynthetic analogues thereof; further all synthetic analogues thereof built from building blocks according to the master structure of the natural compounds (cyclic, 8 amino acids), further all synthetic or semisynthetic analogues containing non-hydroxylated amino acids instead of the hydroxylated amino acids (provided that there is at least one hydroxy group present at the phenyl ring of the tryptophan moiety), further all synthetic or semisynthetic analogues, in each case wherein any such derivative or analogue is functionally active by inhibiting mammalian RNA polymerase II.

[0059] Thus, in the context of the present invention, the term "eight amino acid residues of an amanitin derivative" refers to the specific amino acids that form the bicyclic amanitin polypeptide structure.

[0060] Functionally, amatoxins are defined as peptides or depsipeptides that inhibit mammalian RNA polymerase II. Preferred amatoxins are those with a functional group (e.g. a carboxylic group or carboxylic acid derivative such as a carboxamide or hydroxamic acid, an amino group, a hydroxy group, a thiol or a thiol-capturing group) that can be reacted with linker molecules or target-binding moieties as defined above. Amatoxins which are particularly suitable for the conjugates of the present invention are di-deoxy variants of α -amanitin, β -amanitin, γ -amanitin, ϵ -amanitin, amanullin, or amanullinic acid, or mono-deoxy variants of amanin, amaninamide, γ -amanin, or γ -amaninamide as shown in Fig. 1 as well as salts, chemical derivatives, semisynthetic analogues, and synthetic analogues thereof.

[0061] In a particular embodiment, the hydroxy-substituted derivative, the amanitin derivative comprising a hydroxylated tryptophan moiety, and/or the conjugate of the present invention have/has a purity greater than 90%, particularly greater than 95%, more particularly greater than 98%, or even more than 99%.

[0062] In the context of the present invention, the term "purity" refers to the total amount of, e.g. conjugates being present. A purity of greater than 90%, for example, means that in 1 mg of a composition comprising a conjugate of the present invention, there are more than 90%, i.e. more than 900 μ g, of such conjugate. The remaining part, i.e. the impurities may include unreacted starting material and other reactants, solvents, cleavage products and/or side products.

[0063] In a particular embodiment, a composition comprising the hydroxy-substituted derivative, the amanitin derivative comprising a hydroxylated tryptophan moiety, and/or the conjugate of the present invention comprises more than 100 mg, in particular more than 500 mg, and more particularly more than 1 g of such hydroxy-substituted derivative, amanitin derivative comprising a hydroxylated tryptophan

moiety, and/or conjugate. Thus, trace amount of, e.g. a conjugate of the present invention that arguably may be present in complex preparations of conjugates of the prior art are explicitly excluded.

[0064] The term “target-binding moiety”, as used herein, refers to any molecule or part of a molecule that can specifically bind to a target molecule or target epitope. Preferred target-binding moieties in the context of the present application are (i) antibodies or antigen-binding fragments thereof; (ii) antibody-like proteins; and (iii) nucleic acid aptamers. “Target-binding moieties” suitable for use in the present invention typically have a molecular mass of 40 000 Da (40 kDa) or more.

[0065] As used herein, a first compound (e.g. an antibody) is considered to “specifically bind” to a second compound (e.g. an antigen, such as a target protein), if it has a dissociation constant K_D to said second compound of 100 μM or less, particularly 50 μM or less, particularly 30 μM or less, particularly 20 μM or less, particularly 10 μM or less, particularly 5 μM or less, more particularly 1 μM or less, more particularly 900 nM or less, more particularly 800 nM or less, more particularly 700 nM or less, more particularly 600 nM or less, more particularly 500 nM or less, more particularly 400 nM or less, more particularly 300 nM or less, more particularly 200 nM or less, even more particularly 100 nM or less, even more particularly 90 nM or less, even more particularly 80 nM or less, even more particularly 70 nM or less, even more particularly 60 nM or less, even more particularly 50 nM or less, even more particularly 40 nM or less, even more particularly 30 nM or less, even more particularly 20 nM or less, and even more particularly 10 nM or less.

[0066] In the context of the present application the terms “target molecule” and “target epitope”, respectively, refers to an antigen and an epitope of an antigen, respectively, that is specifically bound by a target-binding moiety. Particularly the target molecule is a tumour-associated antigen, in particular an antigen or an epitope which is present on the surface of one or more tumour cell types in an increased concentration and/or in a different steric configuration as compared to the surface of non-tumour cells. Particularly, said antigen or epitope is present on the surface of one or more tumour cell types, but not on the surface of non-tumour cells. In particular embodiments, the target-binding moiety specifically binds to an epitope of

an antigen selected from: PSMA, CD19, CD269, sialyl Lewis^a, HER-2/neu and epithelial cell adhesion molecule (EpCAM). In other embodiments, said antigen or epitope is preferentially expressed on cells involved in autoimmune diseases. In particular such embodiments, the target-binding moiety specifically binds to an epitope of the IL-6 receptor (IL-6R).

[0067] The term “antibody or antigen binding fragment thereof”, as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e. molecules that contain an antigen-binding site that immunospecifically binds an antigen. Thus, the term “antigen-binding fragments thereof” refers to a fragment of an antibody comprising at least a functional antigen-binding domain. Also comprised are immunoglobulin-like proteins that are selected through techniques including, for example, phage display to specifically bind to a target molecule, e.g. to a target protein selected from: PSMA, CD19, CD269, sialyl Lewis^a, HER-2/neu and EpCAM. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. “Antibodies and antigen-binding fragments thereof” suitable for use in the present invention include, but are not limited to, polyclonal, monoclonal, monovalent, bispecific, heteroconjugate, multispecific, human, humanized (in particular CDR-grafted), deimmunized, or chimeric antibodies, single chain antibodies (e.g. scFv), Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, diabodies or tetrabodies (Holliger P. et al., Proc Natl Acad Sci U S A. 90 (1993) 6444-8), nanobodies, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above.

[0068] In some embodiments the antigen-binding fragments are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (dsFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable domain(s) alone or in combination with the entirety or a portion of the

following: hinge region, CL, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments also comprising any combination of variable domain(s) with a hinge region, CL, CH1, CH2, and CH3 domains.

[0069] Antibodies usable in the invention may be from any animal origin including birds and mammals. Particularly, the antibodies are from human, rodent (e.g. mouse, rat, guinea pig, or rabbit), chicken, pig, sheep, goat, camel, cow, horse, donkey, cat, or dog origin. It is particularly preferred that the antibodies are of human or murine origin. As used herein, “human antibodies” include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described for example in U.S. Patent No. 5,939,598 by Kucherlapati & Jakobovits.

[0070] The term “antibody-like protein” refers to a protein that has been engineered (e.g. by mutagenesis of loops) to specifically bind to a target molecule. Typically, such an antibody-like protein comprises at least one variable peptide loop attached at both ends to a protein scaffold. This double structural constraint greatly increases the binding affinity of the antibody-like protein to levels comparable to that of an antibody. The length of the variable peptide loop typically consists of 10 to 20 amino acids. The scaffold protein may be any protein having good solubility properties. Particularly, the scaffold protein is a small globular protein. Antibody-like proteins include without limitation affibodies, anticalins, and designed ankyrin repeat proteins (for review see: Binz et al., Nat Biotechnol. 2005, 1257-68). Antibody-like proteins can be derived from large libraries of mutants, e.g. be panned from large phage display libraries and can be isolated in analogy to regular antibodies. Also, antibody-like binding proteins can be obtained by combinatorial mutagenesis of surface-exposed residues in globular proteins.

[0071] The term “nucleic acid aptamer” refers to a nucleic acid molecule that has been engineered through repeated rounds of *in vitro* selection or SELEX (systematic evolution of ligands by exponential enrichment) to bind to a target molecule (for a review see: Brody and Gold, J Biotechnol. 74 (2000) 5-13). The

nucleic acid aptamer may be a DNA or RNA molecule. The aptamers may contain modifications, e.g. modified nucleotides such as 2'-fluorine-substituted pyrimidines.

[0072] A "linker" in the context of the present invention refers to a structure that is connecting two components, each being attached to one end of the linker. In the case of the linker being a bond, a direct linkage of amatoxin to the antibody may decrease the ability of the amatoxin to interact with RNA polymerase II. In particular embodiments, the linker increases the distance between two components and alleviates steric interference between these components, such as in the present case between the antibody and the amatoxin. In particular embodiments, the linker has a continuous chain of between 1 and 30 atoms (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 atoms) in its backbone, i.e. the length of the linker is defined as the shortest connection as measured by the number of atoms or bonds between the amatoxin moiety and the antibody, wherein one side of the linker backbone has been reacted with the amatoxin and, the other side is available for reaction, or has been reacted, with an antibody. In the context of the present invention, a linker particularly is a C₁₋₂₀-alkylene, C₁₋₂₀-heteroalkylene, C₂₋₂₀-alkenylene, C₂₋₂₀-heteroalkenylene, C₂₋₂₀-alkynylene, C₂₋₂₀-heteroalkynylene, cycloalkylene, heterocycloalkylene, arylene, heteroarylene, aralkylene, or a heteroaralkylene group, optionally substituted. The linker may contain one or more structural elements such as carboxamide, ester, ether, thioether, disulfide, urea, thiourea, hydrocarbon moieties and the like. The linker may also contain combinations of two or more of these structural elements. Each one of these structural elements may be present in the linker more than once, e.g. twice, three times, four times, five times, or six times. In some embodiments the linker may comprise a disulfide bond. It is understood that the linker has to be attached either in a single step or in two or more subsequent steps to the amatoxin and the antibody. To that end the linker to be will carry two groups, particularly at a proximal and distal end, which can (i) form a covalent bond to a group present in one of the components to be linked, particularly an activated group on an amatoxin or the target binding-peptide or (ii) which is or can be activated to form a covalent bond with a group on an amatoxin. Accordingly, it is preferred that chemical groups are at the distal and proximal end of the linker, which are the result of such a coupling reaction, e.g. an ester, an ether, a urethane, a peptide bond etc.

[0073] In particular embodiments, the linker L is a linear chain of between 1 and 20 atoms independently selected from C, O, N and S, particularly between 2 and 18 atoms, more particularly between 5 and 16 atoms, and even more particularly between 6 and 15 atoms. In particular embodiments, at least 60% of the atoms in the linear chain are C atoms. In particular embodiments, the atoms in the linear chain are linked by single bonds.

[0074] In particular embodiments, the linker L is an alkylene, heteroalkylene, alkenylene, heteroalkenylene, alkynylene, heteroalkynylene, cycloalkylene, heterocycloalkylene, arylene, heteroarylene, aralkylene, or a heteroaralkylene group, comprising from 1 to 4 heteroatoms selected from N, O, and S, wherein said linker is optionally substituted.

[0075] The term “alkylene” refers to a bivalent straight chain saturated hydrocarbon groups having from 1 to 20 carbon atoms, including groups having from 1 to 10 carbon atoms. In certain embodiments, alkylene groups may be lower alkylene groups. The term “lower alkylene” refers to alkylene groups having from 1 to 6 carbon atoms, and in certain embodiments from 1 to 5 or 1 to 4 carbon atoms. Examples of alkylene groups include, but are not limited to, methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{CH}_2-$), n-propylene, n-butylene, n-pentylene, and n-hexylene.

[0076] The term “alkenylene” refers to bivalent straight chain groups having 2 to 20 carbon atoms, wherein at least one of the carbon-carbon bonds is a double bond, while other bonds may be single bonds or further double bonds. The term “alkynylene” herein refers to groups having 2 to 20 carbon atoms, wherein at least one of the carbon-carbon bonds is a triple bond, while other bonds may be single, double or further triple bonds. Examples of alkenylene groups include ethenylene ($-\text{CH}=\text{CH}-$), 1-propenylene, 2-propenylene, 1-butenylene, 2-butenylene, 3-butenylene, and the like. Examples of alkynylene groups include ethynylene, 1-propynylene, 2-propynylene, and so forth.

[0077] As used herein, “cycloalkylene” is intended to refer to a bivalent ring being part of any stable monocyclic or polycyclic system, where such ring has between 3 and 12 carbon atoms, but no heteroatom, and where such ring is fully saturated, and the term “cycloalkenylene” is intended to refer to a bivalent ring being part of any stable monocyclic or polycyclic system, where such ring has between 3 and 12 carbon atoms, but no heteroatom, and where such ring is at least partially unsaturated (but excluding any arylene ring). Examples of cycloalkylenes include, but are not limited to, cyclopropylene, cyclobutylene, cyclopentylene, cyclohexylene, and cycloheptylene. Examples of cycloalkenylenes include, but are not limited to, cyclopentenylene and cyclohexenylene.

[0078] As used herein, the terms “heterocycloalkylene” and “heterocycloalkenylene” are intended to refer to a bivalent ring being part of any stable monocyclic or polycyclic ring system, where such ring has between 3 and about 12 atoms, and where such ring consists of carbon atoms and at least one heteroatom, particularly at least one heteroatom independently selected from the group consisting of N, O and S, with heterocycloalkylene referring to such a ring that is fully saturated, and heterocycloalkenylene referring to a ring that is at least partially unsaturated (but excluding any arylene or heteroarylene ring).

[0079] The term “arylene” is intended to mean a bivalent ring or ring system being part of any stable monocyclic or polycyclic system, where such ring or ring system has between 3 and 20 carbon atoms, but has no heteroatom, which ring or ring system consists of an aromatic moiety as defined by the “ $4n+2$ ” π electron rule, including phenylene.

[0080] As used herein, the term “heteroarylene” refers to a bivalent ring or ring system being part of any stable mono- or polycyclic system, where such ring or ring system has between 3 and 20 atoms, which ring or ring system consists of an aromatic moiety as defined by the “ $4n+2$ ” π electron rule and contains carbon atoms and one or more nitrogen, sulfur, and/or oxygen heteroatoms.

[0081] In the context of the present invention, the term “substituted” is intended to indicate that one or more hydrogens present in the backbone of a linker is replaced with a selection from the indicated group(s), provided that the indicated atom's normal valency, or that of the appropriate atom of the group that is substituted, is not exceeded, and that the substitution results in a stable compound. The term “optionally substituted” is intended to mean that the linker is either unsubstituted or substituted, as defined herein, with one or more substituents, as defined herein. When a substituent is a keto (or oxo, i.e. =O) group, a thio or imino group or the like, then two hydrogens on the linker backbone atom are replaced. Exemplary substituents include, for example, alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, acyl, aroyl, heteroaroyl, carboxyl, alkoxy, aryloxy, acyloxy, aroyloxy, heteroaroyloxy, alkoxycarbonyl, halogen, (thio)ester, cyano, phosphoryl, amino, imino, (thio)amido, sulfhydryl, alkylthio, acylthio, sulfonyl, a sulfate, a sulfonate, a sulfamoyl, a sulfonamido, nitro, azido, haloalkyl, including perfluoroalkyl (such as trifluoromethyl), haloalkoxy, alkylsulfanyl, alkylsulfinyl, alkylsulfonyl, alkylsulfonylamino, arylsulfonylamino, phosphoryl, phosphate, phosphonate, phosphinate, alkylcarboxy, alkylcarboxamide, oxo, hydroxy, mercapto, amino (optionally mono- or di-substituted, e.g. by alkyl, aryl, or heteroaryl), imino, carboxamide, carbamoyl (optionally mono- or di-substituted, e.g. by alkyl, aryl, or heteroaryl), amidino, aminosulfonyl, acylamino, aroylamino, (thio)ureido, (arylthio)ureido, alkyl(thio)ureido, cycloalkyl(thio)ureido, aryloxy, aralkoxy, or -O(CH₂)_n-OH, -O(CH₂)_n-NH₂, -O(CH₂)_nCOOH, -(CH₂)_nCOOH, -C(O)O(CH₂)_nR, -(CH₂)_nN(H)C(O)OR, or -N(R)S(O)₂R wherein n is 1-4 and R is independently selected from hydrogen, -alkyl, -alkenyl, -alkynyl, -cycloalkyl, -cycloalkenyl, -(C-linked-heterocycloalkyl), -(C-linked-heterocycloalkenyl), -aryl, and -heteroaryl, with multiple degrees of substitution being allowed. It will be understood by those skilled in the art that substituents, such as heterocycloalkyl, aryl, heteroaryl, alkyl, etc., or functional groups such as -OH, -NHR etc., can themselves be substituted, if appropriate. It will also be understood by those skilled in the art that the substituted moieties themselves can be substituted as well when appropriate.

[0082] In particular embodiments, the linker L comprises a moiety selected from one of the following moieties: a disulfide (-S-S-), an ether (-O-), a thioether (-S-), an

amine (-NH-), an ester (-O-C(=O)- or -C(=O)-O-), a carboxamide (-NH-C(=O)- or -C(=O)-NH-), a urethane (-NH-C(=O)-O- or -O-C(=O)-NH-), and a urea moiety (-NH-C(=O)-NH-).

[0083] In particular embodiments of the present invention, the linker L comprises a number of m groups selected from the list of: alkylene, alkenylene, alkynylene, cycloalkylene, heteroalkylene, heteroalkenylene, heteroalkynylene, heterocycloalkylene, arylene, heteroarylene, aralkylene, and a heteroaralkylene group, wherein each group may optionally be independently substituted, the linker further comprises a number of n moieties independently selected from one of the following moieties: a disulfide (-S-S-), an ether (-O-), a thioether (-S-), an amine (-NH-), an ester (-O-C(=O)- or -C(=O)-O-), a carboxamide (-NH-C(=O)- or -C(=O)-NH-), a urethane (-NH-C(=O)-O- or -O-C(=O)-NH-), and a urea moiety (-NH-C(=O)-NH-), wherein $m = n + 1$. In particular embodiments, m is 2 and n is 1, or m is 3 and n is 2. In particular embodiments, the linker comprises 2 or 3 unsubstituted alkylene groups, and 1 or 2, respectively, disulfide, ether, thioether, amine, ester, carboxamide, urethane or urea moieties linking the unsubstituted alkylene groups.

[0084] In a particular embodiment, the linker L does not comprise a heteroarylene group.

[0085] In particular embodiments, the C atoms in the linear chain are independently part of optionally substituted methylene groups (-CH₂-). In particular such embodiments, the optional substituents are independently selected from halogen and C₁₋₆-alkyl, particularly methyl.

[0086] In particular embodiments, the linker L is a stable linker.

[0087] In the context of the present invention, the term “stable linker” refers to a linker that is stable (i) in the presence of enzymes, and (ii) in an intracellular reducing environment.

[0088] In particular embodiments, the stable linker does not contain (i) an enzyme-cleavable substructure, and/or (ii) a disulfide group. In particular such embodiments,

the linker has a length of up to 12 atoms, particularly from 2 to 10, more particularly from 4 to 9, and most particularly from 6 to 8 atoms.


[0089] In particular other embodiments, the linker is a cleavable linker.

[0090] In the context of the present invention, the term “cleavable linker” refers to a linker that is (i) cleavable by an enzyme, or (ii) a reducible linker. In particular embodiments, the term only refers to a linker that is cleavable by an enzyme (not to a reducible linker).


[0091] In the context of the present invention, the term “linker that is cleavable ... by an enzyme” refers to a linker that can be cleaved by an enzyme, particularly by a lysosomal peptidase, such as Cathepsin B, resulting in the intracellular release of the toxin cargo conjugated to the targeting antibody after internalization (see Dubowchik et al., Bioconjug Chem. 13 (2002) 855-69). In particular embodiments, the cleavable linker comprises a dipeptide selected from: Phe-Lys, Val-Lys, Phe-Ala, Val-Ala, Phe-Cit and Val-Cit, particularly wherein the cleavable linker further comprises a p-aminobenzyl (PAB) spacer between the dipeptides and the amatoxin.

[0092] In particular such embodiments, the cleavable linker comprises a structure $L^1-L^*-L^2$, wherein L^* is p-aminobenzyl dipeptide moiety, L^1 is a part of the linker that connects L^* to the amatoxin, in particular, wherein L^1 is connected to L^* via a $-NH-$ or a $-O-$ group, particularly a $-C(=O)-NH-$, a $-C(=O)-NH-O-$ or a $-C(=O)-O-$ group, and wherein L^2 is a part of the linker that connects L^* to the target-binding moiety, in particular wherein L^2 is connected to L^* via a $-(CH_2)_m-$ moiety, with m being an integer selected from 1 to 8, in particular from 1 to 5, or via a $-(CH_2CH_2O)_n-$ moiety, with n being an integer selected from 1 to 3, in particular from 1 to 2. For example, in the case of the cleavable linker comprising the dipeptide Val-Ala, the structure of $L^1-L^*-L^2$ is as follows:





[0093] In particular other such embodiments, L^* comprises the dipeptide Val-Lys and has the following structure:




[0094] In particular embodiments, the linker L^1 is a linear chain of between 1 and 4 atoms independently selected from C, O, N and S, particularly between 1 and 3 atoms, more particularly between 1 and 2 atoms, and even more just 1 atom. In particular embodiments, at least 50% of the atoms in the linear chain are C atoms. In particular embodiments, the atoms in the linear chain are linked by single bonds.

In a particular embodiment, L^1 is a $-NH-$ or a $-O-$ group that is part of the amatoxin. In particular embodiments, L^1 is a $-O-$ group originating from a hydroxy group attached to position 4', 5', 6' or 7' of the central tryptophan moiety. In particular embodiments, L^1 is a $-O-$ group originating from the hydroxyl group being part of the carboxylic acid group of amino acid residue 1 of an amanin derivative according to the present invention. In particular embodiments, L^1 is a $-NH-$ group originating from the amino group being part of the carboxamide group of amino acid residue 1 of an amanineamide derivative according to the present invention. In particular embodiments, L^1 is a $-O-$ group originating from a hydroxyl group being part of amino

acid residue 3 of an amanin or amaninamide derivative according to the present invention.

[0095] In the context of the present invention, the term “reducible linker” refers to a linker that can be cleaved in the intracellular reducing environment, particularly a linker that contains a disulfide groups, resulting in the intracellular release of the toxin cargo conjugated to the target-binding moiety after internalization by the intracellular reducing environment (see Shen et al., J. Biol. Chem. 260 (1985) 10905–10908). In particular embodiments, the reducible linker comprises a moiety



wherein R1 to R4 are independently selected from H and methyl.

[0096] In particular such embodiments, such cleavable linker has a length of up to 20 atoms, particularly from 6 to 18, more particularly from 8 to 16, and most particularly from 10 to 15 atoms. In particular such embodiments, the part of the linker linking the amatoxin according to the present invention and the cleavable disulfide group is a linear chain of 3 or 4 C atoms, particularly 3 C atoms. In particular embodiments, the 3 or 4 C atoms in the linear chain are linked by single bonds. In particular embodiments, the linker is an n-propylene group.

[0097] In particular embodiments, said linker is present and is connected on one side to the hydroxyl group attached to the phenyl ring of the central tryptophan moiety, i. e. to a 4', 5' or 7' hydroxy substituent.

[0098] In particular other embodiments, said linker is present and is connected on one side to a position in the amanitin derivative of the present invention, wherein said position is selected from

- (i) in the case of S-desoxy-4'-hydroxy-amaninamide, 4'-hydroxy-amaninamide, S-desoxy-5'-hydroxy-amaninamide, 5'-hydroxy-amaninamide, S-desoxy-7'-hydroxy-amaninamide, and 7'-hydroxy-amaninamide, the nitrogen

atom of the carboxamide group at the γ C-atom of amatoxin amino acid 1 (amide linkage);

(ii) in the case of S-desoxy-4'-hydroxy-amanin, 4'-hydroxy-amanin, S-desoxy-5'-hydroxy-amanin, 5'-hydroxy-amanin, S-desoxy-7'-hydroxy-amanin, 7'-hydroxy-amanin,, the oxygen atom of the acid group at the γ C-atom of amatoxin amino acid 1 (ester linkage);

(iii) in the case of a derivative of the amanitin of the present invention, wherein the free carboxylic acid moiety of amino acid 1 has been converted to a moiety $-C(=O)NH-OR_1$, the oxygen atom of the hydroxamic acid group at the γ C-atom of amatoxin amino acid 1;


(iv) the oxygen atom of the hydroxy group at the δ C-atom of amatoxin amino acid 3, particularly via an ester linkage, an ether linkage or a urethane linkage; or

(v) the ring nitrogen of amino acid 4.

[0099] Coupling of the linker to the target-binding moiety can be achieved by a variety of methods well known to one of ordinary skill in the art, particularly in the art of antibody-drug conjugates (ADCs).

[00100] In particular embodiments, said linker is connected to the target-binding moiety via a urea moiety (...-linker-NH-C(=O)-NH-target-binding moiety). In particular such embodiments, the urea moiety results from a reaction of a primary amine originally present in the target-binding moiety, such as an amino group of a lysine side chain, with a carbamic acid derivative ...-linker-NH-C(O)-Z, wherein Z is a leaving group that can be replaced by a primary amine.

[00101] In particular other embodiments, said linker is present and is connected to the target-binding moiety via a thioether moiety (...-linker-S-target-binding moiety). Thus, in such embodiments, the present invention relates to a conjugate of generic formula:



wherein Amanitin is an amanitin derivative according to the present invention, L is a linker, X^* is a moiety resulting from coupling of a thiol group to a thiol-reactive group,

S is the sulphur atom of said thiol group, particularly the thiol group of a cysteine amino acid residue, and Tbm is a target-binding moiety, particularly an antibody or a functional antibody fragment comprising said cysteine amino acid residue. In particular embodiments, said cysteine amino acid residue (i) is located in an antibody domain selected from CL, CH1, CH2, and CH3; (ii) is located at a position, where the germline sequence exhibiting the closest homology to the sequence of said antibody domain contains an amino acid residue different from cysteine; and (iii) is located a position that is solvent-exposed.

[00102] In the context of the present invention, the term “thiol-reactive group” refers to a group that selectively reacts with the thiol group of, for example, a free cysteine of an antibody, particularly in a pH value in the range between 6.0 and 8.0, more particularly in a pH value in the range between 6.5 and 7.5. In particular, the term “selectively” means that less than 10% of the coupling reactions of a molecule comprising a thiol-reactive group with an antibody comprising at least one free cysteine residue are coupling reactions with non-cysteine residues of the antibody, such as lysine residues, particularly less than 5%, more particularly less than 2%. In particular embodiments, the thiol-reactive group is selected from bromoacetamide, iodoacetamide, maleimide, a maleimide having a leaving group in position 3, in particular a leaving group selected from –Br, and substituted thiol (see, for example, US 9,295,729), a 1,2-dihydropyridazine-3,6-dione having a leaving group in position 4, in particular a leaving group selected from –Br, and substituted thiol (see, for example, US 9,295,729), methylsulfonyl benzothiazole, methylsulfonyl phenyltetrazole, methylsulfonyl phenyloxadiazole (see Toda et al., *Angew. Chem. Int. Ed. Engl.*, 52 (2013) 12592-6), a 3-arylpropionitrile (see Kolodych et al, *Bioconjugate Chem.* 2015, 26, 197–200), and 5-nitro-pyridin-2-yl-disulfide (...-L-S-S-(5-nitro-pyridine-2-yl)).

[00103] In particular embodiments, said position or functional group, which is on one side connected to the linker and which can directly or indirectly be connected to a position or functional group present in a target-binding moiety is a moiety that can react with two thiol groups present in one target-binding moiety or in two target-binding moieties. In particular embodiments, the thiol-reactive groups is a maleimide having two leaving groups in positions 3 and 4, in particular selected from 3,4-

dibromomaleimide, 3,4-bis(arylthio)-maleimide, in particular 3,4-diphenylthio-maleimide, and 3,4-bis(heteroarylthio)-maleimide, in particular 3,4-bis(2-pyridinyl-sulfanyl)-maleimide, and. In particular other embodiments, the thiol-reactive groups is a 1,2-dihydropyridazine-3,6-dione having two leaving groups in positions 4 and 5, in particular selected from 4,5-bromo-1,2-dihydropyridazine-3,6-dione, 4,5-bis(arylthio)-1,2-dihydropyridazine-3,6-dione, in particular 4,5-diphenylthio-1,2-dihydropyridazine-3,6-dione, and 4,5-bis(heteroarylthio)-1,2-dihydropyridazine-3,6-dione, in particular 4,5-bis(2-pyridinyl-sulfanyl)-1,2-dihydropyridazine-3,6-dione.

[00104] In particular embodiments, the moiety resulting from coupling of a thiol group to a thiol-reactive group is selected from: thiol-substituted acetamide; thiol-substituted succinimide; thiol-substituted succinamic acid; thiol-substituted heteroaryl, particularly thiol-substituted benzothiazole, thiol-substituted phenyltetrazole and thiol-substituted phenyloxadiazole; and a disulphide, wherein one sulphur atom is derived from a cysteine residue of the antibody. In particular embodiments, the moiety resulting from coupling of a thiol group to a thiol-reactive group is a thiol-substituted succinimide.

[00105] In particular embodiments, the linker L in the moiety L-X*-S present in the generic formula of section [00101], is selected from the following group of moieties:


- (Amanitin side) $-(\text{CH}_2)_2\text{-S-S-(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_3\text{-S-S-(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_2\text{-S-S-(CH}_2)_3\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_3\text{-S-S-(CH}_2)_3\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_4\text{-S-S-(CH}_2)_4\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_2\text{-CMe}_2\text{-S-S-(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_2\text{-S-S-CMe}_2\text{-(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-(\text{CH}_2)_3\text{-S-S-}$ (Tbm side);
- (Amanitin side) $-\text{CH}_2\text{-C}_6\text{H}_4\text{-NH-Cit-Val-CO(CH}_2)_5\text{-X-S-}$ (Tbm side)
- (Amanitin side) $-\text{CH}_2\text{-C}_6\text{H}_4\text{-NH-Ala-Val-CO(CH}_2)_5\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-\text{CH}_2\text{-C}_6\text{H}_4\text{-NH-Ala-Val-CO(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-\text{CH}_2\text{-C}_6\text{H}_4\text{-NH-Ala-Phe-CO(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-\text{CH}_2\text{-C}_6\text{H}_4\text{-NH-Lys-Phe-CO(CH}_2)_2\text{-X-S-}$ (Tbm side);
- (Amanitin side) $-\text{CH}_2\text{-C}_6\text{H}_4\text{-NH-Cit-Phe-CO(CH}_2)_2\text{-X-S-}$ (Tbm side);

(Amanitin side) $-\text{CH}_2-\text{C}_6\text{H}_4-\text{NH}-\text{Val}-\text{Val}-\text{CO}(\text{CH}_2)_2-\text{X}-\text{S}-$ (Tbm side);
 (Amanitin side) $-\text{CH}_2-\text{C}_6\text{H}_4-\text{NH}-\text{Ile}-\text{Val}-\text{CO}(\text{CH}_2)_2-\text{X}-\text{S}-$ (Tbm side);
 (Amanitin side) $-\text{CH}_2-\text{C}_6\text{H}_4-\text{NH}-\text{His}-\text{Val}-\text{CO}(\text{CH}_2)_2-\text{X}-\text{S}-$ (Tbm side);
 (Amanitin side) $-\text{CH}_2-\text{C}_6\text{H}_4-\text{NH}-\text{Met}-\text{Val}-\text{CO}(\text{CH}_2)_2-\text{X}-\text{S}-$ (Tbm side);
 (Amanitin side) $-\text{CH}_2-\text{C}_6\text{H}_4-\text{NH}-\text{Asn}-\text{Lys}-\text{CO}(\text{CH}_2)_2-\text{X}-\text{S}-$ (Tbm side); and

wherein $-\text{NH}-$ and $-\text{CO}-$ flanking the dipeptide sequences represent amino and carbonyl moieties of the linker forming amide bonds to the carboxy- and the amino-terminus of the dipeptide, respectively.

[00106] In the context of the present invention, the term “a moiety resulting from coupling of a thiol group to a thiol-reactive group” refers to a structure that results from (i) the nucleophilic substitution of a leaving group Y present in a thiol-reactive group by the sulphur atom of a cysteine residue, for example a bromo acetamide group, a iodo acetamide, a 4,6-dichloro-1,3,5-triazin-2-ylamino group, an alkylsulfone or a heteroarylsulfone; (ii) the addition of the HS-group of a cysteine residue to an activated double bond of a thiol-reactive group, for example maleimide, or (iii) an disulfide exchange of an activated disulfide or methanethiosulfonate with the sulphur atom of a cysteine residue, for example with pyridine-2-thiol, 5-nitropyridine-2-thiol or methanesulfinat as leaving group; or (iv) any other chemical reaction that results in a stable bond between the sulphur atom of a cysteine residue and a reactive moiety being part of the thiol-reactive group.


[00107] The primary moiety resulting from coupling of thiol group may be optionally further derivatized, e.g. the succinimidyl thioether resulting from a maleimide can be hydrolysed to succinamic acid thioethers of the following generic structures



[00108] In particular other embodiments, site-specific coupling can be achieved by reducing a disulfide bridge present in the target-binding moiety, and by reacting the


two cysteine residues with a bridging moiety X* present in an Amanitin – L – X* construct (see Badescu et al. Bridging disulfides for stable and defined antibody drug conjugates. *Bioconjugate Chemistry*. 25 (2014) 1124-1136).

[00109] In a similar embodiment, site-specific coupling can be achieved by reducing a disulfide bridge present in the target-binding moiety, and by reacting the two cysteine residues with a bridging moiety X* present in a Amanitin – L – X* construct, particularly wherein X* is




(see Bryden et al., *Bioconjug Chem*, 25 (2014) 611-617; Schumacher et al., *Org Biomol Chem*, 2014, 12, 7261-7269)

[00110] In a particular other embodiment, coupling is achieved by regiospecific coupling of an amino group present in the linker to a glutamine residue present in the target-binding moiety via a transaminase, particularly by coupling to glutamine Q295 of an antibody.



[00111] In a particular embodiment, coupling is achieved by site-specific conjugation to target-binding moieties comprising N-glycan side chains. In particular, the N-glycan side chain can be degraded enzymatically, followed by transglycosylation with an azido-galactose. Using click chemistry, such modified target-binding moiety can be coupled to appropriately modified constructs Amanitin – L – X*, wherein X* is, for example, a dibenzo-cyclooctyne (DIBO) or an analogous moiety comprising a C-C triple bond. For example, a construct Amanitin –L–NH₂ can be coupled to DIBO-SE



DIBO-SE

by nucleophilic substitution of the hydroxy succinimide moiety. The resulting DIBO-modified linker construct can then be coupled to the azido derivative mentioned above. In an alternative embodiment, the target-binding moiety can be modified by incorporation of a non-natural amino acid that permits click-chemistry, in particular by incorporation of a para-azidomethyl-L-phenylalanine (pAMF).


[00112] In particular embodiments, the linker L in $-L-X^*$ is a linear chain of at least 5, particularly at least 10, more particularly between 10 and 20 atoms independently selected from C, O, N and S, particularly between 10 and 18 atoms, more particularly between 10 and 16 atoms, and even more particularly between 10 and 15 atoms. In particular embodiments, at least 60% of the atoms in the linear chain are C atoms. In particular embodiments, the atoms in the linear chain are linked by single bonds.

[00113] In alternative embodiments, the position or functional group, which can directly or indirectly be connected to a position or functional group present in a target-binding moiety, is not an ethynyl group, or, more generally, is not an alkynyl group, or is not a group that can be reacted with an 1,3 dipole in a 1,3-dipolar cycloaddition (click chemistry).

[00114] In particular other embodiments, site-specific coupling of a Amanitin $-L-X^*$ construct to a target-binding moiety can be achieved by incorporation of a non-natural amino acid comprising a keto group, in particular p-acetylphenylalanine (pAcPhe), into the target-binding moiety, and by reacting such modified target-

binding moiety with an Amanitin – L – X* construct, wherein X* is a hydroxylamine moiety.

[00115] In a further embodiment, a formyl group can be introduced by formylglycine generating enzyme (FGE), which is highly selective for a cysteine group in a recognition sequence CxPxR to generate an aldehyde tag. Such aldehyde tag can be reacted with an appropriate group X* present in an Amanitin – L – X* construct, in particular wherein X* is



(see Agarwal et al., Bioconjugate Chem 24 (2013) 846–851).

[00116] In a second aspect, the present invention relates to a pharmaceutical composition comprising the conjugate of the present invention.

[00117] In a third aspect, the present invention relates to a conjugate of the present invention for use in the treatment of cancer in a patient, particularly wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, cholangiocarcinoma, colorectal cancer, lung cancer, prostate cancer, ovarian cancer, prostate cancer, stomach cancer, kidney cancer, malignant melanoma, leukemia, and malignant lymphoma.

[00118] As used herein, "treat", "treating" or "treatment" of a disease or disorder means accomplishing one or more of the following: (a) reducing the severity of the disorder; (b) limiting or preventing development of symptoms characteristic of the disorder(s) being treated; (c) inhibiting worsening of symptoms characteristic of the disorder(s) being treated; (d) limiting or preventing recurrence of the disorder(s) in patients that have previously had the disorder(s); and (e) limiting or preventing recurrence of symptoms in patients that were previously symptomatic for the disorder(s).

[00119] As used herein, the treatment may comprise administering a conjugate or a pharmaceutical composition according to the present invention to a patient, wherein "administering" includes in vivo administration, as well as administration directly to tissue ex vivo, such as vein grafts.

[00120] In particular embodiments, a therapeutically effective amount of the conjugate of the present invention is used.

[00121] A "therapeutically effective amount" is an amount of a therapeutic agent sufficient to achieve the intended purpose. The effective amount of a given therapeutic agent will vary with factors such as the nature of the agent, the route of administration, the size and species of the animal to receive the therapeutic agent, and the purpose of the administration. The effective amount in each individual case may be determined empirically by a skilled artisan according to established methods in the art.

[00122] In another aspect the present invention relates to pharmaceutical composition comprising an amanitin derivative according to the present invention, or a conjugate of the present invention of an amanitin derivative with a target-binding moiety, and further comprising one or more pharmaceutically acceptable diluents, carriers, excipients, fillers, binders, lubricants, glidants, disintegrants, adsorbents; and/or preservatives.

[00123] "Pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans.

[00124] In particular embodiments, the pharmaceutical composition is used in the form of a systemically administered medicament. This includes parenterals, which comprise among others injectables and infusions. Injectables are formulated either in the form of ampoules or as so called ready-for-use injectables, e.g. ready-to-use syringes or single-use syringes and aside from this in puncturable flasks for

multiple withdrawal. The administration of injectables can be in the form of subcutaneous (s.c.), intramuscular (i.m.), intravenous (i.v.) or intracutaneous (i.c.) application. In particular, it is possible to produce the respectively suitable injection formulations as a suspension of crystals, solutions, nanoparticulate or a colloid dispersed systems like, e.g. hydrosols.

[00125] Injectable formulations can further be produced as concentrates, which can be dissolved or dispersed with aqueous isotonic diluents. The infusion can also be prepared in form of isotonic solutions, fatty emulsions, liposomal formulations and micro-emulsions. Similar to injectables, infusion formulations can also be prepared in the form of concentrates for dilution. Injectable formulations can also be applied in the form of permanent infusions both in in-patient and ambulant therapy, e.g. by way of mini-pumps.

[00126] It is possible to add to parenteral drug formulations, for example, albumin, plasma, expander, surface-active substances, organic diluents, pH-influencing substances, complexing substances or polymeric substances, in particular as substances to influence the adsorption of the target-binding moiety toxin conjugates of the invention to proteins or polymers or they can also be added with the aim to reduce the adsorption of the target-binding moiety toxin conjugates of the invention to materials like injection instruments or packaging-materials, for example, plastic or glass.

[00127] The amanitin derivatives of the present invention comprising a target-binding moiety can be bound to microcarriers or nanoparticles in parenterals like, for example, to finely dispersed particles based on poly(meth)acrylates, polylactates, polyglycolates, polyamino acids or polyether urethanes. Parenteral formulations can also be modified as depot preparations, e.g. based on the "multiple unit principle", if the target-binding moiety toxin conjugates of the invention are introduced in finely dispersed, dispersed and suspended form, respectively, or as a suspension of crystals in the medicament or based on the "single unit principle" if the target-binding moiety toxin conjugate of the invention is enclosed in a formulation, e.g. in a tablet or a rod which is subsequently implanted. These implants or depot medicaments in

single unit and multiple unit formulations often consist of so called biodegradable polymers like e.g. polyesters of lactic acid and glycolic acid, polyether urethanes, polyamino acids, poly(meth)acrylates or polysaccharides.

[00128] Adjuvants and carriers added during the production of the pharmaceutical compositions of the present invention formulated as parenterals are particularly *aqua sterilisata* (sterilized water), pH value influencing substances like, e.g. organic or inorganic acids or bases as well as salts thereof, buffering substances for adjusting pH values, substances for isotonization like e.g. sodium chloride, sodium hydrogen carbonate, glucose and fructose, tensides and surfactants, respectively, and emulsifiers like, e.g. partial esters of fatty acids of polyoxyethylene sorbitans (for example, Tween[®]) or, e.g. fatty acid esters of polyoxyethylenes (for example, Cremophor[®]), fatty oils like, e.g. peanut oil, soybean oil or castor oil, synthetic esters of fatty acids like, e.g. ethyl oleate, isopropyl myristate and neutral oil (for example, Miglyol[®]) as well as polymeric adjuvants like, e.g. gelatine, dextran, polyvinylpyrrolidone, additives which increase the solubility of organic solvents like, e.g. propylene glycol, ethanol, N,N-dimethylacetamide, propylene glycol or complex forming substances like, e.g. citrate and urea, preservatives like, e.g. benzoic acid hydroxypropyl ester and methyl ester, benzyl alcohol, antioxidants like e.g. sodium sulfite and stabilizers like e.g. EDTA.


[00129] When formulating the pharmaceutical compositions of the present invention as suspensions in a preferred embodiment thickening agents to prevent the setting of the target-binding moiety toxin conjugates of the invention or, tensides and polyelectrolytes to assure the resuspendability of sediments and/or complex forming agents like, for example, EDTA are added. It is also possible to achieve complexes of the active ingredient with various polymers. Examples of such polymers are polyethylene glycol, polystyrene, carboxymethyl cellulose, Pluronic[®] or polyethylene glycol sorbit fatty acid ester. The target-binding moiety toxin conjugates of the invention can also be incorporated in liquid formulations in the form of inclusion compounds e.g. with cyclodextrins. In particular embodiments dispersing agents can be added as further adjuvants. For the production of lyophilisates scaffolding agents

like mannite, dextran, saccharose, human albumin, lactose, PVP or varieties of gelatine can be used.

EXAMPLES


[00130] In the following, the invention is explained in more detail by non-limiting examples:

A. Total synthesis of S-desoxy- α -amanitin HDP 30.0735



HDP 30.0735

1. Preparation of N-(tert-butoxycarbonyl)-L-6-acetoxy-tryptophan HDP 30.2550



HDP 30.2550


[00131] 590.0 mg (2.68 mmol) 6-Hydroxy-L-tryptophan (CAS: 13567-14-1) was suspended in a mixture of 30 ml 1,4-dioxane/water 1:1 (v,v). Under argon 2.68 ml (2.68 mmol) 1N NaOH were added at once at ambient temperature. The resulting yellow solution was then treated with 574.6 μ l (2.68 mmol) Boc anhydride (Boc_2O) and stirred for 24 h at room temperature. The solution was acidified with 1N hydrochloric acid to pH 2.4 and extracted 3 times with 25 ml ethylacetate. The combined ethylacetate extracts were washed with saturated NaCl solution and dried over MgSO_4 . Filtration and evaporation to dryness gave 785.0 mg crude material. The crude N-Boc-6-hydroxy-L-tryptophan was dissolved in 4.91 ml (4.91mmol) 1N NaOH and treated with 463.2 ml (500.3mg, 4.90mmol) acetic anhydride. The reaction mixture was stirred for 3 h under argon and acidified with 5% citric acid. The aqueous phase was extracted three times with 25 ml ethylacetate, washed with saturated NaCl and dried over MgSO_4 . Filtration and evaporation gave 635 mg of a crude solid.

[00132] The crude product was purified by flash chromatography on a 330 g silica gel column (detection wave length 254 nm) with a gradient of CH_2Cl_2 +1% acetic acid to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (15:1) +1% acetic acid and gave after co evaporation with toluene 564.4 mg (56 % yield) of a white powder.

MS (ESI⁺)found: 361.08 [M-H]⁺;

calc.: 362.15 ($\text{C}_{18}\text{H}_{22}\text{N}_2\text{O}_6$)

2. Preparation of cis,trans-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-6-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole cis-HDP 30.2555 and trans HDP 30.2555 (cis,trans-6-Acetoxy-Hpi)



Photooxygenation

[00133] The photo-oxygenation was carried out with a 400 W high-pressure sodium vapor lamp (Sirius X400 lamp 230 V, 400 W; 55000 lumen at a distance of 1.3 m) or alternatively with a tungsten-halogen lamp (500 W). A filter solution ($\text{CuCl}_2\text{-CaCO}_3$) cutting off light with $\lambda < 490$ nm is used for a tungsten-halogen lamp.

[00134] Methylene blue or Rose Bengal is used as a dye sensitizer.

[00135] The reaction was carried out in a 500 ml cylindrical reaction vessel with heat exchange jacket made of DURAN[®] borosilicate glass, flat bottom and flat laboratory flange (DN) with two connectors with GL 18 thread. The distance from lamp to reaction vessel, was 15 cm and the reaction temperature was in a range of 3-4°C.

[00136] The final product was purified on a Teledyne ISCO Flash chromatography system with a 330 g Silica Redi Sept Flash column (Teledyne ISCO cat. 69-2203-330). Solvents CH₂Cl₂, CH₃OH, CH₃COOH were standard HPLC or BP grade.

[00137] Dry oxygen (99.5% purity) was bubbled through the reaction mixture with a rate of 2-4 l per minute.

[00138] 943.0 mg (2.60 mmol) N-(tert-butoxycarbonyl)-L-6-acetoxy-tryptophan HDP **30.2550** and 100 mg Rose Bengal were dissolved in 500 ml methanol and cooled to 3°C by using a Huber cryostat with glycol/water as cooling media. The reaction solution was irradiated with the 400 W high-pressure sodium vapor lamp. During the irradiation a slow stream of oxygen was bubbled through the reaction solution. After 5 hours irradiation, oxygenation and cooling was stopped and the reaction media was treated with 10 ml of dimethyl sulfide. The mixture was stirred for 2 hours and evaporated to dryness by using a rotary evaporator with a water bath temperature of 35°C. The dark red residue was dried further in high vacuum to a crystalline solid of 1.20 g. The crude product was purified on a 330 g silica gel column (detection wave length 254nm) with a gradient of CH₂Cl₂ + 5% acetic acid to CH₂Cl₂/MeOH (30:1) + 5% acetic acid. 380 mg cis-HDP 30.2555 and 290 mg trans-HDP 30.2555 were eluted and co evaporated with toluene. After lyophilisation in tert-butanol both isomers were obtained as off-white powders.

cis-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-6-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (**cis-HDP 30.2555**)

380 mg cis-HDP 30.2555 yield: 39%

¹H-NMR (400 MHz, CD₃OD, δ = ppm)

δ = 1.22, 1.44, 1.54 [s, 9H, C(CH₃)₃]; 2.23 (s, 3H, OCOCH₃); 2.46-2.63 (m, 2H, CH₂); 4.14-4.29 (m, 1H, 2-H); 5.35 (s, 1H, 8a-H); 6.39-6.46 (m, 2H, 7-H, 5-H); 7.20- 7.24 (m, 1H, 4-H)

¹³C-NMR (100MHz, CD₃OD, δ=ppm)

δ = 20.93, 28.45, 31.12, 42.80, 61.12, 69.44, 82.21, 85.82, 87.93, 104.97, 112.98, 124.84, 129.42, 151.51, 154.04, 155.97, 171.34, 175.79

MS (ESI⁺) found: 378.92 [MH]⁺; calc.: 378.14 (C₁₈H₂₂N₂O₇)

MS (ESI⁺) found: 401.17 [M+Na]⁺; calc.: 401.14 (C₁₈H₂₂N₂NaO₇)

UV/VIS (CH₃OH): $\lambda_{\max}$ = 296 nm, 239 nm, 215 nm

$\lambda_{\min}$ = 266 nm, 227 nm

trans-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-6-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (**trans-HDP 30.2555**)

290 mg trans-HDP 30.2555 yield: 30%

¹H-NMR (400MHz, CD₃OD, δ =ppm)

δ = 1.22, 1.45, 1.54 [s, 9H, C(CH₃)₃]; 2.22 (s, 3H, OCOCH₃); 2.55-2.73 (m, 2H, CH₂); 4.51-4.57 (m, 1H, 2-H); 5.21-5.24 (s, 1H, 8a-H); 6.36-6.41 (m, 2H, 7-H, 5-H); 7.17-7.18 (m, 1H, 4-H)

¹³C-NMR (100MHz, CD₃OD, δ =ppm)

δ = 20.95, 28.50, 31.12, 42.47, 60.97, 69.44, 82.06, 84.84, 87.54, 104.74, 112.67, 125.03, 128.70, 152.31, 154.22, 156.00, 171.23, 174.67

MS (ESI⁺) found: 379.00 [MH]⁺; calc.: 378.14 (C₁₈H₂₂N₂O₇)

MS (ESI⁺) found: 401.17 [M+Na]⁺; calc.: 401.14 (C₁₈H₂₂N₂NaO₇)


MS (ESI⁺) found: 779.00 [2M+Na]⁺; calc.: 779.28 (C₃₆H₄₄N₄Na₂O₁₄)

UV/VIS (CH₃OH): $\lambda_{\max}$ = 299 nm, 241 nm, 215 nm

$\lambda_{\min}$ = 268 nm, 228 nm

3. Preparation of S-desoxy- α -amanitin HDP 30.0735


Step 1: HDP 30.0013



[00139] FmocHypOH (10.0 g, 28.3 mmol) was suspended in 100 ml 80% MeOH and Cs_2CO_3 (4.6 g, 14.1 mmol) was added. The suspension was stirred at 50°C for 30 minutes until complete dissolution. The reaction mixture was concentrated to dryness and redissolved in 100 ml DMF. Allylbromide (1.6 ml, 3.6 g, 29.7 mmol) was added dropwise and the reaction was stirred over night at room temperature. DMF was distilled off and the residue dissolved in *tert*-butylmethyl ether. Precipitates were filtered and the clear solution was absorbed on Celite prior column chromatography. The compound was purified on 220 g silicagel with n-hexane/ethylacetate gradient.

Yield: 11.5 g, 100%


Step 2: HDP 30.0400



[00140] HDP 30.0013 (5.0 g, 14.1 mmol), pyridinium 4-toluenesulfonate (1.33 g, 5.3mmol) were added to a suspension of 1,3-dihydro-2H-pyran-2-yl-methoxymethyl resin (5.0 g, 1.0 mmol/g THP-resin) in 40 ml dichloroethane. The reaction was stirred at 80°C overnight. After cooling the resin was filtered and extensively washed with dichloroethane, dimethylformamide, acetonitrile, dichloromethane and tert-butylmethyl ether.

[00141] Loading was 0.62 mmol/g (determined by UV-spectroscopy of the fluorenylmethyl group after deprotection)

Step 3: HDP 30.2569 (Solid Phase Synthesis)



Resin pre-treatment:

[00142] HDP 30.0400 (0.31 g, 0.25 mmol) was treated with N,N-dimethylbarbituric acid (241 mg, 1.55 mmol) and $Pd(PPh_3)_4$ (35 mg, 0.03 mmol). The resin was shaken over night at room temperature. Thereafter the resin was extensively washed with dichloromethane, DMF, acetonitrile, dichloromethane and tert-butylmethyl ether and dried under reduced pressure.

Coupling Procedure:

[00143] All reactants and reagents were dissolved in dichloromethane/DMF (1:1, v/v). HDP 30.0477 [see WO 2014/009025] (102 mg, 0.30 mmol) was dissolved in 6.0ml dichloromethane/N,N-dimethylformamide and treated with 4.0 ml of a 0.2 N solution PyBOP/ HOBt and 2 ml DIEA (40% in DMF). After addition of 2.0 ml N,N-dimethylformamide, the reaction was heated to 50°C for 8 minutes by microwave

irradiation (20 W, CEM microwave reactor) and was washed with N,N-dimethylformamide after coupling.

Fmoc-Deprotection:

[00144] Deprotection was performed by addition of 6.0 ml 20% piperidine in N,N-dimethylformamide at 50°C for 10 minutes. The resin was washed with N,N-dimethylformamide (no deprotection after coupling of the final amino acid).

[00145] All other amino acids were coupled following the above protocol, weightings are shown below:

(0.102 g, 0.30mmol	1.5eq	HDP 30.0477	MW: 339.6, see above)
0.72 g, 1.2 mmol	5.0 eq	FmocAsn(Trt)OH	MW: 599.7
0.71 g, 1.2 mmol	5.0 eq	FmocCys(OTrt)OH	MW: 586.7
0.36 g, 1.2 mmol	5.0 eq	FmocGlyOH	MW: 297.3
0.36 g, 1.2 mmol	5.0 eq	FmocIleOH	MW: 353.4
0.36 g, 1.2 mmol	5.0 eq	FmocGlyOH	MW: 297.3
0.114 g, 0.30 mmol	1.5 eq	HDP 30.2555	MW: 378.4

[00146] After completion, the resin was finally transferred into a syringe with bottom frit, washed with DCM and dried under reduced pressure.

Resin Release and B-Ring Formation

[00147] A solution of 5 ml TFA, 5 ml DCM plus 10% MeOH was aspirated to the resin and shaken for 15 min at ambient temperature. The solution was dispensed into a 50 ml reaction flask and the resin washed with TFA/DCM 1:1 plus 10% MeOH once and poured into the same flask. The reaction flask was stirred for 16 h. Triisopropylsilane (0.5 ml) was added and the reaction concentrated in vacuum. The residue was dissolved in 500 µl MeOH and the peptide precipitated in 50 ml ice-cold TBME. After centrifugation the supernatant was decanted and the precipitate washed once with 50 ml TBME and dried under reduced pressure.


[00148] The precipitate was solubilized in 2 ml methanol and purified by preparative reverse phase column chromatography. Methanol was distilled off under reduced pressure and the remaining aqueous phase was freeze dried.

Yield: 136.5 mg, 62.5%

MS (ESI+) found: 1047.4 [M+H]⁺; calc.: 1047.4 (C₄₅H₆₃N₁₀O₁₇S)

HPLC: 91.9area%

Step 4: Cyclisation (A-Ring Formation, HDP 30.2572)




[00149] The above freeze dried monocyclic intermediate (136 mg, 130 µmol) was dissolved in 16 ml DMF and treated with diphenylphosphoryl azide (DPPA, 131 µl, 1300 µmol, 10 eq) and diisopropylethylamine (DIEA, 162 µl, 1300 µmol, 10 eq). The reaction was stirred for 16 h and quenched with 500 µl water upon completion. Conversion was monitored by HPLC. The mixture was concentrated by reduced pressure, re-dissolved in 1 ml methanol and purified by preparative reverse phase column chromatography.

Yield: 55.3 mg, 41%

MS (ESI+) found: 1029.33 [M+Na]⁺; calc.: 1030.10 (C₄₅H₆₁N₁₀O₁₆S)

HPLC: 99.2area%

Step 5: Acetate-Deprotection (HDP 30.0735)


[00150] HDP 30.2572 (55.3 mg, 53.7 μ mol) was dissolved in a 7 N methanolic NH_3 solution (3.0 ml) and stirred. Conversion was checked by HPLC/MS. After completion (6 – 8 h) the reaction was concentrated in vacuum, re-suspended in 100 μ l MeOH and purified by prep-HPLC.

Yield: 14.1 mg, 29%

HPLC: 100 %

MS (ESI+) found: 903.3 $[M+H]^+$; calc.: 902.9 ($C_{39}H_{54}N_{10}O_{13}S$)
found: 925.33 $[M+Na]^+$

4. Preparation of 6'-((3-Maleidopropanamido)-Val-Ala-PAB)-S-deoxy- α -amanitin (HDP 30.2371)

Step 1: HDP 30.2364

[00151] S-Desoxy- α -amanitin **HDP 30.0735** (30 mg, 33.2 μ mol) and **HDP 30.1690** (WO 2016/142049, 78 mg, 133 μ mol = 4.0 eq.) were dissolved in 1500 μ l dry dimethylacetamide (DMA). A 0.2 M cesium hydrogencarbonate solution in water (199 μ l, 1.2 eq.) was added in one portion and the mixture is stirred at room temperature. After 1.5 and 4 h additional portions 99 μ l (0.6 eq.) of cesium carbonate solution were added.


[00152] After 18 h the solvent evaporated by high vacuum.

[00153] The residue was dissolved in 400 μ l of methanol and added dropwise to ice cooled MTBE (10 ml). After standing at 0°C for 10 min, the resulted precipitate was isolated by centrifugation (4 min, 4000 $\times$ g). The supernatant was discharged and the pellet was resuspended in additional MTBE (10 ml) and centrifugation was repeated. The vacuum dried crude product was dissolved in 400 μ l of methanol and purified by prep. HPLC on a Phenomenex Luna-C₁₈(2), 10 μ m column (250 $\times$ 21.2 mm) with a gradient 5% to 100% methanol in water. Product fractions were combined and reduced to 23 mg (70%) product as amorphous solid.

MS (ESI+) [M+Na]⁺ found: 1430.58; calc.: 1430.65 (C₆₅H₉₇N₁₃NaO₁₈SSi)

[00154] By evaporation of the early eluting peak 5 mg (17%) starting material were recovered.

Step 2: HDP 30.2366




[00155] HDP 30.2364 (9.47 mg, 6.72 μ mol) was dissolved in 750 μ l trifluoroacetic acid (TFA). After 2 min the volatiles are removed *in vacuo* and the

residue was dissolved in 750 µl water and 3% ammonia is added drop wise until a pH of 10 was reached and precipitation occurred.

[00156] The solution was freeze-dried and purified on prep. HPLC subsequently (Phenomenex Luna-C₁₈(2), 10 µm column 250 x 21.2 mm, gradient of 5-100% of methanol (0.05% TFA) in water (0.05% TFA) to give 7.72 mg (89% based on TFA salt) product.

MS (ESI⁺) [MH]⁺ found: 1178.42; calc.: 1178.53 (C₅₄H₇₆N₁₃O₁₅S)

Step 3 HDP 30.2371



HDP 30.2366 (4.01mg, 3.10 µmol) was dissolved in 500 µl dry DMF.


[00157] 3-(Maleimido)propionic acid N-hydroxysuccinimide ester (BMPS, 1.65 mg, 2 eq.), dissolved in 100 µl DMF followed by 2.1 µl DIPEA were added.

[00158] After stirring for 1h the reaction mixture was dropped into 10 ml of ice-cooled methyl-*tert*-butyl ether. The tube was kept on ice for 10 min and centrifuged at 4000 × g. The supernatant was removed and the pellet was washed with 10 ml of fresh methyl-*tert*-butyl ether.

[00159] The vacuum-dried pellet was purified by RP-18 HPLC with a gradient 5-100% methanol in water. The pure fractions were evaporated to dryness and lyophilized from 1 ml of *tert*-butanol/water to give 2.70 mg (65%) product as a colorless powder.


MS (ESI+) [MH] ⁺	found: 1329.3;	calc.: 1329.6 (C ₆₁ H ₈₁ N ₁₄ O ₁₈ S)
[M+Na] ⁺	found: 1351.5;	calc.: 1351.5 (C ₆₁ H ₈₀ N ₁₄ NaO ₁₈ S)

B. Total synthesis of S-desoxy-5'-hydroxy-amaninamide HDP 30.2548



HDP 30.2548

1. Preparation of N-(tert-butoxycarbonyl)-L-5-acetoxy-tryptophan HDP 30.2531



HDP 30.2531

[00160] 800.0 mg (3.63 mmol) 5-Hydroxy-L-tryptophan (CAS: 4350-09-8) were suspended in a mixture of 40 ml 1,4-dioxane/water 1:1 (v:v). Under argon 3.64 ml (3.64 mmol) 1N NaOH were added at once at ambient temperature. The resulting

yellow solution was then treated with 779.0 μ l (794.0 mg, 3.63 mmol) Boc anhydride (Boc_2O) and stirred for 24 hours at room temperature. The solution was acidified with 1N hydrochloric acid to pH 2.4 and extracted 3 times with 35 ml ethylacetate. The combined ethylacetate extracts were washed with saturated NaCl solution and dried over MgSO_4 . Filtration and evaporation to dryness gave 1.26 g crude material. The crude N-Boc-6-hydroxy-L-tryptophan was dissolved in 7.87 ml (7.87 mmol) 1 N NaOH and treated with 743.0 μ l (802.4 mg, 7.86 mmol) acetanhydride. The reaction mixture was stirred for 3 hours under argon and acidified with 5% citric acid. The aqueous phase was extracted three times with 25 ml ethylacetate, washed with saturated NaCl and dried over MgSO_4 . Filtration and evaporation.

[00161] The crude solid was purified by flash chromatography on a 330 g silica gel column (detection wave length 254 nm) with a gradient of CH_2Cl_2 +1% acetic acid to $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (15:1) +1% acetic acid and gave after co evaporation with toluene 1020.0mg (78 % yield) of a white powder.


MS (ESI⁺) found: 361.08[M-H]⁺;

calc.: 362.15 ($\text{C}_{18}\text{H}_{22}\text{N}_2\text{O}_6$)

¹H-NMR (400 MHz, CD_3OD , δ = ppm)

δ = 1.39 (s, 9H, $\text{C}(\text{CH}_3)_3$); 2.27 (s, 3H, OCOCH_3); 3.08-3.31 (m, 2H, CH_2); 4.39-4.41 (m, 1H, 2-H); 6.81-6.83; 7.13; 7.24-7.26 (m, 2H, 6-H, 7-H); 7.30- 7.32 (1H, 4-H)

2. Preparation of cis,trans-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-5-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole cis-HDP 30.2536 and trans HDP 30.2536 (cis,trans-5-Acetoxy-Hpi)



Photooxygenation

[00162] The photo-oxygenation was carried out with a 400 W high-pressure sodium vapor lamp (Sirius X400 lamp 230 V, 400 W; 55,000 lumen at a distance of 1.3 m) or alternatively with a tungsten-halogen lamp (500 W). A filter solution ($\text{CuCl}_2\text{-CaCO}_3$) cutting off light with $\lambda < 490 \text{ nm}$ is used for a tungsten-halogen lamp.

[00163] Methylene blue or rose Bengal is used as a dye sensitizer.

The reaction was carried out in a 500 ml cylindrical reaction vessel with heat exchange jacket made of DURAN[®] borosilicate glass, flat bottom and flat laboratory flange (DN) with two connectors with GL 18 thread. The distance from lamp to reaction vessel, was 15 cm and the reaction temperature was in a range of 3-4°C.

[00164] The final product was purified on a Teledyne ISCO Flash chromatography system with a 330 g Silica Redi Sept Flash column (Teledyne ISCO cat. 69-2203-330). Solvents CH_2Cl_2 , CH_3OH , CH_3COOH were standard HPLC or BP grade.

Dry oxygen (99.5% purity) was bubbled through the reaction mixture with a rate of 2-4 l per minute.

[00165] 1.20 g (3.17mmol) N-(tert-butoxycarbonyl)-L-5-acetoxy-tryptophan **HDP 30.2531** and 100 mg Rose Bengal were dissolved in 500 ml methanol and cooled to 3°C by using a Huber cryostat with glycol/water as cooling media. The reaction solution was irradiated with the 400 W high-pressure sodium vapor lamp. During the irradiation a slow stream of oxygen was bubbled through the reaction solution. After 4 hours irradiation, oxygenation and cooling was stopped and the reaction media was treated with 20 ml of dimethyl sulfide. The mixture was stirred for 2 hours and evaporated to dryness by using a rotary evaporator with a water bath temperature of 35°C. The dark red residue was dried further in high vacuum to a crystalline solid. The crude product was purified on a 330 g silica gel column (detection wave length 254 nm) with a gradient of CH₂Cl₂ +5% acetic acid to CH₂Cl₂/MeOH (30:1) + 5% acetic acid.

[00166] 178 mg **cis-HDP 30.2536** and 132 mg **trans-HDP 30.2536** were eluted and co evaporated with toluene. After lyophilisation in tert-butanol both isomers were obtained as off-white powders.

cis-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-5-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (**cis-HDP 30.2536**)

178 mg cis-HDP 30.2536 yield: 15%

¹H-NMR (400MHz, CD₃OD, δ=ppm)

δ = 1.44, 1.54 (s, 9H, C(CH₃)₃); 2.23 (s, 3H, OCOCH₃); 2.45-2.62 (m, 2H, CH₂); 4.18-4.33 (m, 1H, 2-H); 5.37 (s, 1H, 8a-H); 6.63-6.67; 6.82-6.84; (m, 2H, 7-H, 6-H); 6.96- 6.98 (m, 1H, 4-H)

¹³C-NMR (100MHz, CD₃OD, δ = ppm)

δ = 20.86, 28.46, 31.12, 42.81, 61.21, 82.14, 85.60, 111.49, 117.83, 123.98, 132.77, 144.99, 148.03, 156.07, 172.01, 175.89

MS (ESI⁺) found: 379.00 [MH]⁺; calc.: 378.14 (C₁₈H₂₂N₂O₇)

MS (ESI⁺) found: 401.17 [M+Na]⁺; calc.: 401.14 (C₁₈H₂₂N₂NaO₇)

MS (ESI⁻) found: 377.17 [M-H]⁻; calc.: 378.14 (C₁₈H₂₂N₂O₇)

UV/VIS (CH₃OH): $\lambda_{\max}$ = 299nm, 242nm, 207nm

$\lambda_{\min}$ = 270nm, 222nm

trans-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-5-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole (**trans-HDP 30.2536**)

132mg trans-HDP 30.2536 yield: 11%

¹H-NMR (400MHz, CD₃OD, δ = ppm)

δ = 1.47, 1.54 [s, 9H, C(CH₃)₃]; 2.21 (s, 3H, OCOCH₃); 2.50-2.70 (m, 2H, CH₂); 4.51-4.56 (m, 1H, 2-H); 5.22-5.26 (s, 1H, 8a-H); 6.58-6.63; 6.80-6.81 (m, 6-H, 7-H); 6.92 (s, 1H, 4-H)

¹³C-NMR (100MHz, CD₃OD, δ = ppm)

δ = 20.88, 28.50, 31.12, 42.60, 61.11, 82.05, 85.38, 111.26, 117.88, 124.07, 131.94, 144.66, 148.95, 156.11, 172.01, 174.81

MS (ESI⁺) found: 379.00 [MH]⁺; calc.: 378.14 (C₁₈H₂₂N₂O₇)

MS (ESI⁺) found: 401.17 [M+Na]⁺; calc.: 401.14 (C₁₈H₂₂N₂NaO₇)


MS (ESI⁻) found: 377.17 [M-H]⁻; calc.: 378.14 (C₁₈H₂₂N₂O₇)

UV/VIS (CH₃OH): $\lambda_{\max}$ = 302nm, 244nm, 208nm

$\lambda_{\min}$ = 272nm, 224nm

3. Preparation of S-desoxy-5'-hydroxy-amaninamide HDP 30.2548


Step 1: HDP 30.0013



[00167] FmocHypOH (10.0 g, 28.3 mmol) was suspended in 100 ml 80% MeOH and Cs_2CO_3 (4.6 g, 14.1 mmol) was added. The suspension was stirred at 50°C for 30 minutes until complete dissolution. The reaction mixture was concentrated to dryness and redissolved in 100 ml DMF. Allylbromide (1.6 ml, 3.6 g, 29.7 mmol) was added dropwise and the reaction was stirred over night at room temperature. DMF was distilled off and the residue dissolved in *tert*-butylmethyl ether. Precipitates were filtered and the clear solution was absorbed on Celite prior column chromatography. The compound was purified on 220 g Silicagel with n-hexane/ethylacetate gradient.

Yield: 11.5 g, 100%


Step 2: HDP 30.0400



[00168] **HDP 30.0013** (5.0 g, 14.1 mmol), pyridinium 4-toluenesulfonate (1.33 g, 5.3 mmol) were added to a suspension of 1,3-dihydro-2H-pyran-2-yl-methoxymethyl resin (5.0 g, 1.0 mmol/g THP-resin) in 40 ml dichloroethane. The reaction was stirred at 80°C overnight. After cooling the resin was filtered and extensively washed with dichloroethane, dimethylformamide, acetonitrile, dichloromethane and tert-butylmethylether.

[00169] Loading was 0.62 mmol/g (determined by UV-spectroscopy of the fluorenylmethyl group after deprotection)

Step 3: HDP 30.2544 (Solid Phase Synthesis)



Resin pre-treatment:

[00170] **HDP 30.0400** (0.31 g, 0.25 mmol) was treated with N,N-dimethylbarbituric acid (241 mg, 1.55 mmol) and Pd(PPh₃)₄ (35 mg, 0.03 mmol). The resin was shaken over night at room temperature. Thereafter the resin was extensively washed with dichloromethane, DMF, acetonitrile, dichloromethane and tert-butylmethyl ether and dried under reduced pressure.

Coupling Procedure:

[00171] All reactants and reagents were dissolved in dichloromethane/DMF (1:1, v/v). **HDP 30.0477** [see WO 2014/009025] (102mg, 0.30mmol) was dissolved in 6.0 ml dichloromethane/N,N-dimethylformamide and treated with 4.0 ml of a 0.2 N solution PyBOP/HOBt and 2 ml DIEA (40% in DMF). After addition of 2.0 ml N,N-dimethylformamide, the reaction was heated to 50°C for 8 minutes by microwave

irradiation (20 W, CEM microwave reactor) and was washed with N,N-dimethylformamide after coupling.

Fmoc-Deprotection:

[00172] Deprotection was performed by addition of 6.0 ml 20% piperidine in N,N-dimethylformamide at 50°C for 10 minutes. The resin was washed with N,N-dimethylformamide (no deprotection after coupling of the final amino acid).

[00173] All other amino acids were coupled following the above protocol, weightings are shown below:

(0.102 g, 0.30 mmol above)	1.5 eq	HDP 30.0477	MW: 339.6, see above)
0.72 g, 1.2 mmol	5.0 eq	FmocAsn(Trt)OH	MW: 599.7
0.71 g, 1.2 mmol	5.0 eq	FmocCys(OTrt)OH	MW: 586.7
0.36 g, 1.2 mmol	5.0 eq	FmocGlyOH	MW: 297.3
0.36 g, 1.2 mmol	5.0 eq	FmocIleOH	MW: 353.4
0.36 g, 1.2 mmol	5.0 eq	FmocGlyOH	MW: 297.3
0.114 g, 0.30 mmol	1.5 eq	HDP 30.2536	MW: 378.4

[00174] After completion, the resin was finally transferred into a syringe with bottom frit, washed with DCM and dried under reduced pressure.

Resin Release and B-Ring Formation

[00175] A solution of 5 ml TFA, 5 ml DCM plus 10% MeOH was aspirated to the resin and shaken for 15 min at ambient temperature. The solution was dispensed into a 50 ml reaction flask and the resin washed with TFA/DCM 1:1 plus 10% MeOH once and poured into the same flask. The reaction flask was stirred for 16 h. Triisopropylsilane (0.5 ml) was added and the reaction concentrated in vacuum. The residue was dissolved in 500 µl MeOH and the peptide precipitated in 50 ml ice-cold TBME. After centrifugation the supernatant was decanted and the precipitate washed once with 50 ml TBME and dried under reduced pressure.


[00176] The precipitate was solubilized in 2 ml methanol and purified by preparative reverse phase column chromatography. Methanol was distilled off under reduced pressure and the remaining aqueous phase was freeze dried.

Yield: 75.1 mg, 35.9%

MS (ESI⁺) found: 1047.4 [M+H]⁺; calc.: 1047.4 (C₄₅H₆₃N₁₀O₁₇S)

HPLC: 99.3area%

Step 4: Cyclisation (A-Ring Formation, HDP 30.2546)




[00177] The above freeze dried monocyclic intermediate (49.4 mg, 47.2 μmol) was dissolved in 3 ml DMF and treated with diphenylphosphoryl azide (DPPA, 13 μl, 237 μmol, 5 eq) and diisopropylethylamine (DIEA, 40 μl, 237 μmol, 5 eq). The reaction was stirred for 16 h and quenched with 500 μl water upon completion. Conversion was monitored by HPLC. The mixture was concentrated by reduced pressure, re-dissolved in 1ml methanol and purified by preparative reverse phase column chromatography.

Yield: 32.5 mg, 67.9%

MS (ESI⁺) found: [M+H]⁺1029.33 [MH]⁺ 1051.3 [M+Na]⁺; calc.: 1029.39; 1051.42 (C₄₅H₆₁N₁₀O₁₆S; C₄₅H₆₀N₁₀O₁₆SNa)

HPLC: 88.3area%

Step 5: Acetate-Deprotection (HDP 30.2548)

[00178] **HDP 30.2546** (23.4 mg, 22.7 μ mol) was dissolved in a 7 N methanolic NH_3 solution (2.3 ml) and stirred. Conversion was checked by HPLC/MS. After completion (6 – 8h) the reaction was concentrated in vacuum, re-suspended in 100 μ l MeOH and purified by prep-HPLC.


Yield: 12.5 mg, 53.3%

HPLC: 99 %

MS (ESI⁺) found: 903.4 [M+H]⁺; calc.: 902.9 ($C_{39}H_{54}N_{10}O_{13}S$)
 found: 925.4 [M+Na]⁺

5. Preparation of 5'-((3-Maleidopropanamido)-Val-Ala-*p*-amininobenzyloxy)-S-deoxy-amaninamide (HDP 30.2602)

Step 1: HDP 30.2563




[00179] S-desoxy-5'-hydroxy-amaninamide **HDP 30.2548** (9.66 mg, 10.7 μ mol) and **HDP 30.1690** (WO 2016/142049, 50.2 mg, 85.6 μ mol = 8.0 eq.) were dissolved in 500 μ l dry dimethylacetamide (DMA). A 1.0 M cesium hydrogen carbonate solution in water (17.12 μ l, 1.6 eq.) was added in one portion and the mixture is stirred at room temperature.

[00180] After 1.5 h additional portions 17.12 μ l (1.6 eq.) of cesium carbonate solution were added.

[00181] After 8 h the reaction mixture was neutralized with acetic acid, passed through a centrifugal filter (0.2 μ m) and purified by HPLC on a Phenomenex Luna-C₁₈(2), 10 μ m column (250 x 21.2 mm) with a gradient (5-100%) of acetonitrile in water. Product fractions were combined and reduced to 4.00 mg (26%) product as amorphous solid.

MS (ESI⁺) [M+Na]⁺ found: 1430.58; calc.: 1430.65 (C₆₅H₉₇N₁₃NaO₁₈SSi)

Step 2: HDP 30.2378




[00182] **HDP 30.2563** (7.53 mg, 5.34 μ mol) was dissolved in 500 μ l of a trifluoroacetic acid/water/triisopropylsilane 95:5:5 mixture. After 5 min the volatiles were removed *in vacuo* and the residue was dissolved in 1000 μ l water and 3% ammonia was added drop wise until a pH of 10 was reached.

[00183] The solution was freeze-dried and purified on prep. HPLC subsequently (Phenomenex Luna- C_{18} (2), 10 μ m column 250 x 21.2 mm, gradient of 20-30% in 16 min of acetonitrile in water (0.05% TFA) to give 3.95 mg (57% based on TFA salt) product.

MS (ESI⁺) [M+H]⁺ found: 1178.50; calc.: 1178.53 ($C_{54}H_{76}N_{13}O_{15}S$)

Step 3 HDP 30.2602




[00184] **HDP 30.2578** (3.95 mg, 3.06 μ mol) was dissolved in 400 μ l dry DMF.

[00185] 3-(Maleimido)propionic acid N-hydroxysuccinimide ester (BMPS, 1.63 mg, 2 eq.), dissolved in 81.4 μ l DMF followed by 2.1 μ l (2 eq.) DIPEA were added.


[00186] After stirring for 2 h the reaction mixture was passed through a centrifugal filter (0.2 μm) and purified by RP-18 HPLC with a gradient 5-70% acetonitrile in water+ 0.05% TFA. The pure fractions were evaporated to dryness and lyophilized from 2 ml of acetonitrile/water 1:1 to give 2.73 mg (67%) product as a colorless powder.

MS (ESI+) $[\text{M}+\text{H}]^+$ found: 1329.33; calc.: 1329.56 ($\text{C}_{61}\text{H}_{81}\text{N}_{14}\text{O}_{18}\text{S}$)


C. Total synthesis of S-desoxy-4'-hydroxy-amaninamide



- Schematic view of the synthesis of cis,trans-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-4-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole; cis- and trans (cis,trans-4-Acetoxy-Hpi)**




2. Schematic view of the synthesis of S-desoxy-4'-hydroxy-amaninamide



D. Total synthesis of S-desoxy-7'-hydroxy-amaninamide


1. Synthesis of cis,trans-1-(tert-butoxycarbonyl)-2-carboxy-3a-hydroxy-4-acetoxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole; (cis,trans-4-acetoxy-Hpi)

[00187] The synthesis of cis,trans-7-acetoxy-Hpi is done in analogy to the synthesis of cis,trans-4-acetoxy-Hpi starting from commercially available 7-hydroxy-L-tryptophan instead of 4-hydroxy-L-tryptophan.



2. Schematic view of the synthesis of S-desoxy-7'-hydroxy-amaninamide

[00188] The synthesis of S-desoxy-7'-hydroxy-amaninamide is done in analogy to the synthesis of S-desoxy-4'-hydroxy-amaninamide using cis,trans-7-acetoxy-Hpi instead of cis,trans-4-acetoxy-Hpi.



6. *In vitro* cytotoxicity of S-desoxy- α -amanitin HDP 30.0735 and S-desoxy-5'-hydroxy-amaninamide HDP 30.2548

BrdU Cell Proliferation Assay on HEK293 and HEK293 OATP1B3 cells.

[00189] HEK293-OATP1B3 cell culture plates were coated with poly-D-lysine.

Coating with Poly-D-Lysine:

- 5mg Poly D-Lysine in 50ml sterile water
- 50 μ l in each well of a 96 well plate
- Incubation for 1h at RT

- Washing of wells twice with 200µl sterile water
- Drying for at least for 2h (RT)

[00190] Black 96-well plates with clear bottom with 2.0×10^3 HEK293 and HEK293 OATP1B3 cells/well and 90 µl growth medium per well including 10% FCS and supplements were prepared. Controls: „Blank“ was set up with 100 µl medium without cells, „Background“ and „100%“ were set up with cells in 100 µl medium.

- Incubation for 24h at 37°C and 5% CO₂.

Dilution scheme of HDP 30.0735, HDP 30.2548 and alpha-amanitin:

[00191] The stock solutions were diluted 1:1000 (1:10 and 1:100 dilution) in medium:

10µl Amanitin derivative stock solution (1.0×10^{-2} M) + 90 µl PBS =
100 µl 1.0×10^{-3} M
2 µl dilution + 198 µl medium = 200 µl 1×10^{-5} M

Further dilutions:

A: 200 µl 1.0×10^{-5} M
B: 80 µl growth medium + 20 µl solution A (1:5 dilution)
C: 80 µl growth medium + 20 µl solution B (1:5 dilution)
D: 80 µl growth medium + 20 µl solution C (1:5 dilution)
E: 80 µl growth medium + 20 µl solution D (1:5 dilution)
F: 80 µl growth medium + 20 µl solution E (1:5 dilution)
G: 80 µl growth medium + 20 µl solution F (1:5 dilution)
H: 80 µl growth medium + 20 µl solution G (1:5 dilution)

[00192] 10 µl of each solution were added to well triplicates.

Final volume: 100 µl/well

Final dose: starting 1×10^{-6} M; 1:5 dilution series.

- Incubation for 96h at 37°C and 5% CO₂.
- After 96 h: Roche Cell proliferation assay, luminescent according to manufacturer instructions.
- EC₅₀-concentrations were determined with Graphpad Prism 4.0 data analysis software.

7. : Synthesis of HDP 30.2347, HDP 2371 and 30.2602 conjugates

Example: Synthesis of T-D265C-30.2371:

[00193] 10 mg Thiomab T-D265C in PBS buffer were used for conjugation to HDP 30.2371.

Adjust antibody solution to 1 mM EDTA:

2 ml antibody solution (10.0 mg) + 20 µl 100 mM EDTA, pH 8.0

Amount antibody: 10 mg = 6.9×10^{-8} mol

[00194] Uncapping of cysteines by reaction of antibody with 40 eq. TCEP:

- 2 ml antibody solution (6.9×10^{-8} mol) + 55.2 µl 50 mM TCEP solution (2.76×10^{-6} mol)
- Incubate for 3 h at 37°C on a shaker.
- Two consecutive dialyses at 4°C in 2.0 l 1x PBS, 1 mM EDTA, pH 7.4 in a Slide-A-Lyzer Dialysis Cassette 20,000 MWCO, first dialysis ca. 4 h, second dialysis overnight

[00195] Oxidation by reaction of antibody with 20 eq. dehydroascorbic acid (dhAA):

- ca. 2 ml antibody solution (6.9×10^{-8} mol) + 27.6 µl fresh 50 mM dhAA solution (1.38×10^{-6} mol)
- Incubate for 3 h at RT on a shaker.

[00196] Conjugation with amanitin using 6 eq. HDP 30.2371 and quenching with 25 eq. N-acetyl-L-cysteine:

Solubilize 2 mg HDP 30.2371 in 200 µl DMSO = 10 µg/µl

- ca. 2 ml antibody solution (= 9.5 mg; 6.53×10^{-8} mol) + **52.1 µl** HDP 30.2371 (=520.8 µg; 3.92×10^{-7} mol).

- Incubate 1 h at RT.
- Quench by addition of 16.3 µl 100 mM N-acetyl-L-cysteine (1.63×10^{-6} mol).
- Incubate 15 min at RT (or overnight at 4°C).
- Purify reaction mix with 1x PD-10 columns equilibrated with 1 x PBS, pH 7.4. Identify protein-containing fractions with Bradford reagent on parafilm and bring protein-containing fractions together.
- Dialysis of antibody solution at 4°C overnight in 2.0 l PBS, pH 7.4 and Slide-A-Lyzer Dialysis Cassettes 20,000 MWCO.

Determination of DAR by LC-ESI-MS-analysis.

[00197] Adjust protein concentration to 5.0 mg/ml (3.4×10^{-5} M) and bring to sterile conditions by filtration. Store at 4°C.

[00198] ADCs with a different antibody or with a different Amanitin derivative were produced accordingly. The molecular amount of antibody was calculated according to the MW of the respective antibody. The amounts of linker toxin, TCEP, dhAA, N-acetyl-L-cysteine were adjusted accordingly to reach the respective equivalents.

8. *In vitro* cytotoxicity of HDP 30.2347, HDP 2371 and 30.2602 conjugates

BrdU Cell Proliferation Assay on SKBR-3, JIMT-1, LnCap, and 22rv1 cells:

[00199] The assay was performed as described above (6.) with the following changes:

[00200] Cell culture plates were not coated with poly-D-lysine.

- Dilution scheme of ADCs:
- The stock solutions were diluted to 1.0×10^{-6} M in growth medium

Further dilutions:

A: 100 µl 1.0×10^{-6} M

B: 80 µl growth medium + 20 µl solution A (1:5 dilution)

C: 80 µl growth medium + 20 µl solution B (1:5 dilution)

D: 80 µl growth medium + 20 µl solution C (1:5 dilution)

E: 80 µl growth medium + 20 µl solution D (1:5 dilution)

F: 80 µl growth medium + 20 µl solution E (1:5 dilution)

G: 80 µl growth medium + 20 µl solution F (1:5 dilution)

H: 80 µl growth medium + 20 µl solution G (1:5 dilution)

- 10 µl of each solution were added to well triplicates. Final volume: 100 µl/well.
Final dose: starting 1×10^{-7} M; 1:5 dilution series

WST-I assay on Raji cells and Nalm-6 cells:

- Transparent F-bottom 96-well plates with 2.0×10^3 cells/well and 90 µl of the respective growth medium per well including 10% FCS and supplements were prepared. Controls: “Blank” was set up with 100 µl medium without cells; “cells only” was set up with cells in 100 µl medium.
- Incubation for 24 h at 37°C and 5% CO₂.
- Dilution scheme of ADCs:
- The stock solutions were diluted to 1.0×10^{-6} M in growth medium

Further dilutions:

A: 100 µl 1.0×10^{-6} M

B: 80 µl growth medium + 20 µl solution A (1:5 dilution)

C: 80 µl growth medium + 20 µl solution B (1:5 dilution)

D: 80 µl growth medium + 20 µl solution C (1:5 dilution)

E: 80 µl growth medium + 20 µl solution D (1:5 dilution)

F: 80 µl growth medium + 20 µl solution E (1:5 dilution)


G: 80 µl growth medium + 20 µl solution F (1:5 dilution)

H: 80 µl growth medium + 20 µl solution G (1:5 dilution)

- 10 µl of each solution were added to well triplicates. Final volume: 100 µl/well.
Final dose: starting 1 x 10⁻⁷ M; 1:5 dilution series
- Incubation for 96 h at 37°C and 5% CO₂.
- After 96 h: Roche WST-1 Cell Proliferation assay according to manufacturer instructions.
- EC₅₀-concentrations were determined with Graphpad Prism 4.0 data analysis software.

CLAIMS

1. A hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole according to Formula I



wherein R1 is selected from alkyl, aryl, heteroaryl, substituted alkyl, substituted aryl, and substituted heteroaryl;

P₁ is hydrogen or a protecting group;

P₂ is hydrogen or a protecting group, particularly a protecting group; and

R2 is selected from OH, OR1, and a polypeptide chain consisting of 1-7 amino acid residues.

2. The hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole of claim 1, wherein the protecting group, when present, is independently selected from Boc, PhCH₂OCO-, CH₂=CHCH₂O-CO-, and trityl.
3. The hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole of claim 1 or 2, wherein the substituent R1-C(=O)-O- is attached to position 5 in Formula I.
4. The hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole of claim 1 or 2, wherein the substituent R1-C(=O)-O- is attached to position 6 in Formula I.
5. A method for the synthesis of a linear precursor comprising eight amino acid residues of an amanitin derivative comprising a hydroxylated tryptophan


- moiety, comprising the step of using a hydroxy-substituted derivative of 2-carboxy-3a-hydroxy-1,2,3,3a,8,8a-hexahydropyrrolo[2,3-b]indole of any one of claims 1 to 5 in the peptide synthesis of said precursor.
6. A method for the synthesis of an amanitin derivative comprising a hydroxylated tryptophan moiety, comprising the steps of.
 - (i) causing or allowing the formation of a bond between the cysteine residue and the tryptophan moiety of the linear precursor of claim 5; and
 - (ii) causing or allowing the formation of said amanitin derivative by reacting the N-terminus of the linear precursor of claim 5 with the C-terminus of said precursor.
 7. The method of claim 6, further comprising the oxidation of the sulfur atom of the cysteine moiety to form a sulfoxide or a sulfone, particularly a sulfoxide.
 8. An amanitin derivative comprising a hydroxylated tryptophan moiety, which is selected from (i) S-desoxy-4'-hydroxy-amanin, 4'-hydroxy-amanin, S-desoxy-5'-hydroxy-amanin, 5'-hydroxy-amanin, S-desoxy-7'-hydroxy-amanin, 7'-hydroxy-amanin, (ii) S-desoxy-4'-hydroxy-amaninamide, 4'-hydroxy-amaninamide, S-desoxy-5'-hydroxy-amaninamide, 5'-hydroxy-amaninamide, S-desoxy-7'-hydroxy-amaninamide, and 7'-hydroxy-amaninamide, (iii) a derivative of the amanitin according to (i), wherein the free carboxylic acid moiety of amino acid 1 is converted to a carboxylic ester $-C(=O)OR_1$ or to a moiety $-C(=O)NH-OR_1$, wherein R_1 is selected from alkyl, aryl, heteroaryl, substituted alkyl, substituted aryl, and substituted heteroaryl.
 9. The amanitin derivative of claim 8, which is selected from S-desoxy-5'-hydroxy-amanin, 5'-hydroxy-amanin, S-desoxy-5'-hydroxy-amaninamide, and 5'-hydroxy-amaninamide
 10. A conjugate comprising (a) amanitin derivative comprising a hydroxylated tryptophan moiety of claim 8 or 9; (b) a target-binding moiety; and (c)

optionally a linker linking said amanitin derivative and said target-binding moiety.

11. A pharmaceutical composition comprising the amanitin derivative of claim 8 or 9 or the conjugate of claim 10.
12. The amanitin derivative of claim 8 or 9, the conjugate of claim 10, or the pharmaceutical composition of claim 11 for use in the treatment of cancer in a patient, particularly wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, cholangiocarcinoma, colorectal cancer, lung cancer, prostate cancer, ovarian cancer, prostate cancer, stomach cancer, kidney cancer, malignant melanoma, leukemia, and malignant lymphoma.
13. A construct comprising (a) an amanitin derivative of claim 8 or 9; and (b) a linker moiety carrying a reactive group Y for linking said amanitin derivative to a target-binding moiety.

FIGURES

Figure 1:



	R ₁	R ₂	R ₃	R ₄
α-amanitin	OH	OH	NH ₂	OH
β-amanitin	OH	OH	OH	OH
γ-amanitin	H	OH	NH ₂	OH
ε-amanitin	H	OH	OH	OH
amanin	OH	OH	OH	H
amaninamide	OH	OH	NH ₂	H
amanullin	H	H	NH ₂	OH
amanullinic acid	H	H	OH	OH
γ-amanin	H	OH	OH	H
γ-amaninamide	H	OH	NH ₂	H





Figure 2:**HDP 30.0735** **α -Amanitin****Figure 3:****Boc-L-tryptophan****HDP 30.0079**



Figure 4:**Figure 5:**


Figure 6:**HDP 30.2371**


Figure 7:**HDP 30.2347**




Figure 8:**Figure 9:**

Figure 10:



HDP 30.2544

HDP 30.2546

Figure 11:





Figure 12:



HDP 30.2602

Figure 13, A):




Figure 13, B)





Figure 13, C)



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/071268

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D487/04 C07K7/64 A61K47/62 C07K1/00 A61K38/12 A61P35/00 ADD. According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D C07K A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MASAKO NAKAGAWA ET AL: "DYE-SENSITIZED PHOTO-OXYGENATION OF TRYPTOPHAN", TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 41, no. 11, 1 January 1985 (1985-01-01), pages 2125-2132, XP002548899, ISSN: 0040-4020, DOI: 10.1016/S0040-4020(01)96583-4 compounds 16, 18b and 18c -----	1
Y	US 2016/220687 A1 (ALHAMDAN OMAR ABDULAZIZ H [SA]) 4 August 2016 (2016-08-04) paragraphs [0045], [-0046], [0077] - [0078] ----- -/-	8-13
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 3 September 2018		Date of mailing of the international search report 12/09/2018
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Schuemacher, Anne

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/071268

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 2 497 499 A1 (HEIDELBERG PHARMA GMBH [DE]) 12 September 2012 (2012-09-12) examples; claim 1; figure 1 -----	8-13
Y	WO 2014/043403 A1 (AGENSYS INC [US]; MENDELSON BRIAN ALAN [US]; MOON SUNG JU [US]) 20 March 2014 (2014-03-20) claims -----	8-13
Y	WO 2012/041504 A1 (HEIDELBERG PHARMA GMBH [DE]; ANDERL JAN [DE]; SIMON WERNER [DE]; MUELL) 5 April 2012 (2012-04-05) examples; claim 2; figure 1 -----	8-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/071268

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2016220687	A1	04-08-2016	NONE
EP 2497499	A1	12-09-2012	AU 2012224848 A1 12-09-2013
		BR 112013023015 A2 13-12-2016	
		CA 2829421 A1 13-09-2012	
		CN 103442736 A 11-12-2013	
		EP 2497499 A1 12-09-2012	
		EP 2683409 A1 15-01-2014	
		ES 2532855 T3 01-04-2015	
		HK 1192839 A1 08-04-2016	
		IL 228191 A 29-12-2016	
		JP 6105492 B2 29-03-2017	
		JP 2014508767 A 10-04-2014	
		KR 20140059163 A 15-05-2014	
		MX 344009 B 02-12-2016	
		NZ 614548 A 24-04-2015	
		RU 2013145314 A 20-04-2015	
		SG 193392 A1 30-10-2013	
		UA 110370 C2 25-12-2015	
		US 2014294865 A1 02-10-2014	
		US 2018185509 A1 05-07-2018	
		WO 2012119787 A1 13-09-2012	
		ZA 201307526 B 27-07-2016	
WO 2014043403	A1	20-03-2014	US 2015218220 A1 06-08-2015
			WO 2014043403 A1 20-03-2014
WO 2012041504	A1	05-04-2012	AU 2011307195 A1 11-04-2013
		BR 112013007577 A2 02-08-2016	
		CA 2812460 A1 05-04-2012	
		CN 103153345 A 12-06-2013	
		EP 2436398 A1 04-04-2012	
		EP 2621536 A1 07-08-2013	
		ES 2402254 T3 30-04-2013	
		ES 2651745 T3 29-01-2018	
		IL 225200 A 29-09-2016	
		JP 5977241 B2 24-08-2016	
		JP 2013544231 A 12-12-2013	
		KR 20130129193 A 27-11-2013	
		MX 344010 B 02-12-2016	
		NZ 608142 A 31-10-2014	
		RU 2013119960 A 10-11-2014	
		SG 189189 A1 31-05-2013	
		UA 111341 C2 25-04-2016	
		US 2013259880 A1 03-10-2013	
		WO 2012041504 A1 05-04-2012	