

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

28 February 2019 (28.02.2019)



(10) International Publication Number

WO 2019/038412 A1

(51) International Patent Classification:

A61K 9/00 (2006.01)

A61K 47/10 (2017.01)

A61K 38/00 (2006.01)

(21) International Application Number:

PCT/EP2018/072835

(22) International Filing Date:

24 August 2018 (24.08.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17187676.6

24 August 2017 (24.08.2017)

EP

(71) Applicant: **NOVO NORDISK A/S** [DK/DK]; Novo Allé, 2880 Bagsværd (DK).

(72) Inventors: **HORN MØLLER, Eva**; Novo Allé, 2880 Bagsværd (DK). **DUELUND SØRENSEN, Michael**; Novo Allé, 2880 Bagsværd (DK). **LUNDQVIST, Joakim**; Novo Allé, 2880 Bagsværd (DK).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: GLP-1 COMPOSITIONS AND USES THEREOF

(57) Abstract: The present invention relates to pharmaceutical compositions of the GLP-1 peptide semaglutide comprising no more than 0.01 % (w/w) phenol, their preparation, kits comprising such compositions as well as uses thereof.



WO 2019/038412 A1

GLP-1 COMPOSITIONS AND USES THEREOF

The present invention relates to the field of pharmaceutical compositions comprising the GLP-1 peptide semaglutide.

BACKGROUND

5 GLP-1 peptides are known to be prone to develop lack of stability in liquid solutions, for example lack of physical stability. Thus, liquid pharmaceutical compositions comprising GLP-1 peptides with even better stability are desired. Such improved stability may be physical stability and/or chemical stability.

SUMMARY

10 In some embodiments the invention relates to liquid pharmaceutical compositions comprising semaglutide and no more than 0.01 %(w/w) phenol. In some embodiments the invention relates to kits comprising the pharmaceutical composition as defined herein. In some embodiments the invention relates to the pharmaceutical composition as defined herein for use in medicine.

DESCRIPTION

15 The present invention relates to liquid pharmaceutical compositions comprising the GLP-1 peptide semaglutide and no more than 0.01 %(w/w) phenol. Surprisingly, the present inventors found that such compositions have improved chemical and/or physical stability. In some embodiments the composition comprises no phenol. In some embodiments the
20 composition comprises 0.01-10 mg/ml semaglutide. In some embodiments the composition has a pH in the range of 6.0-10.0, such as pH 7.0-7.8.

 In some embodiments the composition of the invention is a liquid pharmaceutical composition comprising semaglutide and no more than 0.01 %(w/w) phenol, wherein said composition

- 25 a. is for parenteral administration;
- b. is an aqueous solution comprising at least 60 %w/w water; or
- c. further comprises one or more pharmaceutically acceptable excipients selected from the group consisting of a buffer or an isotonic agent.

 In some embodiments the composition of the invention is a liquid pharmaceutical
30 composition comprising semaglutide, no more than 0.01 %(w/w) phenol, and optionally one or more pharmaceutically acceptable excipients, wherein the formulation is for parenteral administration, such as subcutaneous administration.

In some embodiments the composition of the invention is a liquid pharmaceutical composition comprising semaglutide, no more than 0.01 % (w/w) phenol, at least 60 % w/w water, and optionally one or more pharmaceutically acceptable excipients.

In some embodiments the term “**stability**” as used herein refers to stability of semaglutide in a liquid pharmaceutical composition. In some embodiments stability is chemical stability of the GLP-1 peptide (e.g. determined by HPLC, such as Assay (I) herein), and optionally physical stability of the GLP-1 peptide (e.g. determined by Thioflavine T assay, such as Assay (II) herein).

In some embodiments the term “**chemical stability**” in relation to semaglutide as used herein refers to the covalent bonds of the semaglutide compound being substantially intact. In some embodiments chemical stability of a GLP-1 peptide is determined by HPLC, such as Assay (I) herein. In some embodiments a composition possess chemical stability if its covalent bonds are intact in at least 80% (w/v) of said GLP-1 peptides after storage for 3 months at 25°C. In some embodiments chemical stability of semaglutide is determined by Assay (IV) herein.

In some embodiments the term “**physical stability**” in relation to semaglutide as used herein refers to semaglutide forming substantially no aggregates, e.g. in the form of fibril formation. In some embodiments physical stability is determined by Thioflavine T assay, such as Assay (II) herein.

In some embodiments the composition of the present invention is a stable pharmaceutical composition. The term “stable pharmaceutical composition” when used herein refers to a pharmaceutical composition, e.g. a solution or suspension, comprising GLP-1 peptide, and which composition following storage comprises at least 80% (w/v) of said GLP-1 peptide (e.g. after quiescent storage for 3 months at 25°C). Storage conditions for stability testing may be 2-8°C, such as 5°C, or at least 2.5 years at 5°C. Alternatively, storage conditions for stability testing may be at least 4 weeks, such as 6 weeks or 3 months, optionally at 30°C. The conditions of storage for this stable pharmaceutical composition may be at 5°C for 1 or 2 years. The conditions of storage for this stable pharmaceutical composition may be at 5°C for 3 years. Alternatively, the conditions of this storage may be at 25°C for 24 hours or 1 week. In yet another alternative, the conditions of this storage may be room temperature for two months, such as up to two months.

In some embodiments, chemical stability of the GLP-1 peptide requires at least 80% (w/v), such as at least 90% (w/v) or at least 95% (w/v), of said GLP-1 peptide remaining with its covalent bonds intact at the end of the storage period. In some embodiments chemical stability of the GLP-1 peptide requires at least 95% (w/v), such as at least 97% (w/v)

or at least 99%(w/v), of said GLP-1 peptide remaining with its covalent bonds intact at the end of the storage period.

The composition of the invention comprises no more than 0.01 %(w/w) phenol. In some embodiments the composition comprises substantially no phenol.

5 **Pharmaceutical compositions**

The terms "pharmaceutical composition" and "composition" are used interchangeably herein and refer to pharmaceutical compositions suitable for administration to a subject in need thereof.

In some embodiments the composition comprises 0.01-100 mg/ml semaglutide. In
10 some embodiments the composition comprises 0.1-50 mg/ml, such as 0.5-25 mg/ml or 1-15 mg/ml, semaglutide. In some embodiments the composition comprises 0.1-10 mg/ml, such as 0.5-5 mg/ml or 1-2 mg/ml, semaglutide. In some embodiments the composition comprises 0.01-10 mg/ml, such as 0.01-5 mg/ml, semaglutide. In some embodiments the composition comprises no more than 9 mg/ml, such as no more than 8 mg/ml or no more than 7 mg/ml,
15 semaglutide. In some embodiments the composition comprises no more than 6 mg/ml, such as no more than 5 mg/ml or no more than 4 mg/ml, semaglutide. In some embodiments the composition comprises no more than 3 mg/ml, such as no more than 2 mg/ml or no more than 1 mg/ml, semaglutide. In some embodiments the composition comprises at least 0.01 mg/ml, such as at least 0.02 mg/ml or at least 0.05 mg/ml, semaglutide. In some
20 embodiments the composition comprises 1.34 mg/ml semaglutide.

In some embodiments the composition of the invention has a pH in the range of 3-10, such as pH 6-10 or 6-9. In some embodiments the composition of the invention has a pH in the range of pH 6.5-8.5, such as pH 7.0-7.8.

In some embodiments the composition of the invention comprises one or more
25 pharmaceutically acceptable excipients.

In some embodiments the composition of the invention comprises an isotonic agent, such as propylene glycol. In some embodiments the isotonic agent is propylene glycol or sodium chloride.

In some embodiments the composition of the invention comprises a buffer, such as
30 phosphate buffer, TRIS, citrate, or no buffer. In some embodiments the phosphate buffer is a sodium phosphate buffer, such as disodium hydrogen phosphate.

In some embodiments the composition of the invention comprises no preservative.

The composition of the invention is in the form of a liquid pharmaceutical composition. In some embodiments the liquid pharmaceutical composition is a solution or a

suspension. In some embodiments the composition of the invention is in the form of a solution, such as an aqueous solution. In some embodiments the term "aqueous solution" as used herein refers to a solution comprising at least 60 %w/w water. In some embodiments the aqueous solution comprises 60-99 %w/w water. In some embodiments the aqueous solution comprises at least 75 %w/w water, such as at least 80 %w/w water or at least 85 %w/w water. In some embodiments the aqueous solution comprises at least 90 %w/w water, such as at least 92 %w/w water or at least 94 %w/w water.

Semaglutide

The GLP-1 peptide semaglutide may be prepared as described in WO2006/097537, Example 4. Semaglutide is also known as N^{6,26}-{18-[N-(17-carboxyheptadecanoyl)-L-γ-glutamyl]-10-oxo-3,6,12,15-tetraoxa-9,18-diazaoctadecanoyl}-[8-(2-amino-2-propanoic acid),34-L-arginine]human glucagon-like peptide 1(7-37), see WHO Drug Information Vol. 24, No. 1, 2010. In some embodiments semaglutide may be present in the composition in its fully or partly ionised form; for example one or more carboxylic acid groups (-COOH) may be deprotonated into the carboxylate group (-COO⁻) and/or one or more amino groups (-NH₂) may be protonated into the -NH₃⁺ group. In some embodiments semaglutide is added to the composition in the form of a salt.

Administration and kits

The composition of the invention is for parenteral administration. In some embodiments the composition is for subcutaneous administration.

In some embodiments the composition of the invention is for administration once weekly. In some embodiments the composition of the invention is for administration once daily, once every second or once every third day.

In some embodiments the invention relates to a kit comprising the pharmaceutical composition as defined herein and instructions for use. In some embodiments the instructions for use comprise the package insert of a drug.

In some embodiments the invention relates to a kit comprising the pharmaceutical composition as defined herein and an injection device. In some embodiments the injection device is selected from the group consisting of a durable pen and a prefilled pen. Examples of durable pens are NovoPen® 4 or NovoPen® 5 (both from Novo Nordisk A/S, Denmark). An example of a prefilled pen is FlexPen® (Novo Nordisk A/S, Denmark).

Indications

In some embodiments the compositions of the invention are for use in medicine. In some embodiments the composition of the invention may be used for the following medical treatments:

- 5 (i) prevention and/or treatment of all forms of diabetes, such as hyperglycaemia, type 2 diabetes, impaired glucose tolerance, type 1 diabetes, non-insulin dependent diabetes, MODY (maturity onset diabetes of the young), gestational diabetes, and/or for reduction of HbA1c;
- (ii) delaying or preventing diabetic disease progression, such as progression in type 10 2 diabetes, delaying the progression of impaired glucose tolerance (IGT) to insulin requiring type 2 diabetes, and/or delaying the progression of non-insulin requiring type 2 diabetes to insulin requiring type 2 diabetes;
- (iii) prevention and/or treatment of eating disorders, such as obesity, e.g. by decreasing food intake, reducing body weight, suppressing appetite, inducing satiety; treating 15 or preventing binge eating disorder, bulimia nervosa, and/or obesity induced by administration of an antipsychotic or a steroid; reduction of gastric motility; and/or delaying gastric emptying.

In some embodiments the indication is (i). In some embodiments the indication is (ii). In a still further particular aspect the indication is (iii). In some embodiments the indication 20 is type 2 diabetes and/or obesity.

In some embodiments the method or use comprises prevention, treatment, reduction and/or induction in one or more diseases or conditions defined herein. In some embodiments the indication is (i) and (iii). In some embodiments the indication is (ii) and (iii). In some embodiments the invention comprises administration of an effective amount of a 25 GLP-1 peptide. In some embodiments the invention relates to administration of an effective amount of a GLP-1 peptide.

Generally, all subjects suffering from obesity are also considered to be suffering from overweight. In some embodiments the invention relates to a method for treatment or prevention of obesity. In some embodiments the invention relates to use of the composition 30 for treatment or prevention of obesity. In some embodiments the subject suffering from obesity is human, such as an adult human or a paediatric human (including infants, children, and adolescents). Body mass index (BMI) is a measure of body fat based on height and weight. The formula for calculation is $BMI = \text{weight in kilograms} / \text{height in meters}^2$. A human subject suffering from obesity may have a BMI of ≥ 30 ; this subject may also be referred to as 35 obese. In some embodiments the human subject suffering from obesity may have a BMI of

≥35 or a BMI in the range of ≥30 to <40. In some embodiments the obesity is severe obesity or morbid obesity, wherein the human subject may have a BMI of ≥40.

In some embodiments the invention relates to a method for treatment or prevention of overweight, optionally in the presence of at least one weight-related comorbidity. In some
5 embodiments the invention relates to use of the composition for treatment or prevention of overweight, optionally in the presence of at least one weight-related comorbidity. In some embodiments the subject suffering from overweight is human, such as an adult human or a paediatric human (including infants, children, and adolescents). In some embodiments a human subject suffering from overweight may have a BMI of ≥25, such as a BMI of ≥27. In
10 some embodiments a human subject suffering from overweight has a BMI in the range of 25 to <30 or in the range of 27 to <30. In some embodiments the weight-related comorbidity is selected from the group consisting of hypertension, diabetes (such as type 2 diabetes), dyslipidaemia, high cholesterol, and obstructive sleep apnoea.

In some embodiments the invention relates to a method for reduction of body
15 weight. In some embodiments the invention relates to use of the composition for reduction of body weight. A human to be subjected to reduction of body weight according to the present invention may have a BMI of ≥25, such as a BMI of ≥27 or a BMI of ≥30. In some embodiments the human to be subjected to reduction of body weight according to the present invention may have a BMI of ≥35 or a BMI of ≥40. The term “reduction of body
20 weight” may include treatment or prevention of obesity and/or overweight.

In some embodiments, as used herein, specific values given in relation to numbers or intervals may be understood as the specific value or as about the specific value (e.g. plus or minus 10 percent of the specific value).

25 **Embodiments of the invention**

The following are non-limiting embodiments of the invention:

1. A liquid pharmaceutical composition comprising semaglutide and no more than 0.01 %w/w phenol.
- 30 2. A liquid pharmaceutical composition comprising semaglutide and substantially no phenol.
3. The composition according to claim 1 or 2, wherein said composition does not comprise phenol.
4. The composition according to any one of the preceding claims, wherein said composition is an aqueous solution comprising at least 60 %w/w water, such as at least 70 %w/w
35 water or at least 80 %w/w water.

5. The composition according to any one of the preceding claims, wherein the concentration of semaglutide is 0.5-10 mg/ml of said composition.
6. The composition according to any one of the preceding claims, wherein said semaglutide is in the form of a pharmaceutically acceptable salt.
- 5 7. The composition according to any one of the preceding claims, wherein said composition comprises one or more pharmaceutically acceptable excipients.
8. The composition according to any one of the preceding claims, wherein said composition comprises one or more agents for adjusting pH, such as HCl, NaOH, or acetate.
9. The composition according to any one of the preceding claims, wherein said composition
10 comprises a buffer and/or an isotonic agent.
10. The composition according to any one of the preceding claims, wherein said buffer is present in a concentration of 0.01-50 mM of said composition.
11. The composition according to any one of the preceding claims, wherein said buffer is a phosphate buffer.
- 15 12. The composition according to any one of the preceding claims, wherein said phosphate buffer is selected from the group consisting of sodium dihydrogen phosphate, disodium hydrogen phosphate, and sodium phosphate.
13. The composition according to any one of the preceding claims, wherein said isotonic agent is present in a concentration from 8 mg/ml to 50 mg/ml, such as 14 mg/ml to 30
20 mg/ml, of said composition.
14. The composition according to any one of the preceding claims, wherein said isotonic is propylene glycol.
15. The composition according to any one of the preceding claims, wherein said composition comprises no preservative.
- 25 16. The composition according to any one of the preceding claims, wherein said composition has a pH in the range of 6.0-10.0.
17. The composition according to any one of the preceding claims, wherein said composition is for parenteral administration.
18. The composition according to any one of the preceding claims, wherein said composition
30 is for subcutaneous administration.
19. A kit comprising the pharmaceutical composition as defined in any one of the preceding claims and instructions for use.
20. A kit comprising the pharmaceutical composition as defined in any one of the preceding
35 claims and an injection device for administration of said composition to a subject, wherein

said injection device is selected from the group consisting of a durable pen and a prefilled pen.

21. A pharmaceutical composition as defined in any one of the preceding claims for use in
5 medicine.
22. The pharmaceutical composition for use as defined in any one of the preceding claims for use in the treatment of diabetes or obesity.
23. A method for the prevention or treatment of diabetes or obesity, wherein the
pharmaceutical composition as defined in any one of the preceding claims is
10 administered to a subject in need thereof.

EXAMPLES

General methods and characterisation

15 *Preparation of semaglutide compositions:*

Unless otherwise noted, compositions of semaglutide were prepared by dissolving buffer (e.g. disodiumhydrogenphosphate dihydrate), isotonic agent (e.g. propylene glycol) and optionally preservative (phenol) in water. Semaglutide was dissolved therein, pH was adjusted to 7.4 using sodium hydroxide and/or hydrochloric acid, and the composition was
20 finally sterilised by filtration through a 0.22 µm sterile filter.

Preparation of liraglutide compositions:

Unless otherwise noted, compositions of liraglutide were prepared from Solution 1 and Solution 2: Solution 1 was prepared by dissolving buffer (disodiumhydrogenphosphate dihydrate), isotonic agent (mannitol), and optionally preservative (phenol) in water. Solution 2
25 was prepared by dissolving liraglutide while stirring slowly. Solution 1 and Solution 2 were mixed, pH was adjusted to 8.15 using sodium hydroxide and/or hydrochloric acid, and the composition was finally sterilised by filtration through a 0.22 µm sterile filter.

30 *Assay (I): Determination of High Molecular Weight Proteins (HMWP) content of semaglutide compositions*

Determination of HMWP content was performed using size exclusion chromatography (SE-HPLC) using a Waters Insulin HMWP column with a mobile phase of

sodium chloride, sodium phosphate, phosphoric acid and isopropanol, isocratic elution and detection at 280 nm. Content of HMWP is given in % as the combined area of chromatographic peaks eluting earlier than the semaglutide monomer peak (i.e. HMWP peaks), relative to the total area of HMWP and semaglutide monomer peaks.

5

Assay (II): Physical stability of semaglutide compositions assessed via ThT

The purpose of this assay is to assess the physical stability of a GLP-1 peptide in aqueous solution.

Low physical stability of a peptide or protein may lead to amyloid fibril formation.

10 Fibrils are structurally well-ordered, filamentous macromolecular structures formed by aggregation of soluble proteins and dominated by beta-sheet structure. Mature fibrils are insoluble and are resistant to degradation. For the sake of drug product quality and patient safety, it is desirable to minimize and control fibrillation events in pharmaceutical compositions of therapeutic peptides and proteins. Protein aggregation, including fibrillation, 15 can be assessed by visual inspection of a sample. Fibrillation can be assessed by the use of Thioflavine T (ThT), a small molecule indicator probe with a high specificity for fibrils. ThT has a distinct fluorescence signature when binding to fibrils compared to ThT in solution [Naiki et al. (1989) Anal. Biochem. 177, 244-249; LeVine (1999) Methods. Enzymol. 309, 274-284].

20 Formation of a partially folded intermediate of the peptide is suggested as a general initiating mechanism for fibrillation. A small amount of these intermediates nucleates to form a template onto which further intermediates may assembly and the fibrillation proceeds. The lag-time corresponds to the interval in which a critical amount of nuclei is generated and the apparent rate constant is the rate with which the fibril itself is formed. The lag-time described 25 in a ThT assay performed on a plate reader is therefore considered indicative of the fibrillation tendency of a peptide composition in solution.

Before performing the assay, ThT was added to the samples from a stock solution in H₂O to a final concentration of 20 µM in samples. Sample aliquots of 200 µl of the composition comprising the GLP-1 peptide were placed in a 96 well microtiter plate (optical 30 0.4 mL black Thermo Scientific Nunc) with a glass bead (2.8-3.2 mm, Whitehouse Scientific) placed in each well. Usually, eight replica of each sample were placed on the plate. The plate was sealed with sealing tape (Thermo Scientific Nunc).

Incubation at given temperature, shaking and measurement of the ThT fluorescence emission were performed in a BMG FLUOStar Omega or a BMG FLUOStar Optima. The 35 plate was incubated at 40°C with double orbital shaking at 300 rpm with an amplitude of 2

mm. Fluorescence measurement was performed using excitation through a 450 nm filter and measurement of emission through a 480 nm filter. The plate was measured every 20 minutes for a desired period of time. Between each measurement, the plate was shaken and heated as described.

5 The threshold value was determined as the highest ThT fluorescence (in relative fluorescence units (RFU)) measured on the plate at time 1 h 13 min, plus 100 RFU. The threshold value was then used to calculate the lag time using the "time to threshold" method in the BMG FLUOstar software.

10 *Assay (III): Determination of purity of liraglutide*

 Determination of purity was performed using high performance liquid chromatography (HPLC) using a Waters XTerra™ MS C18 column with a gradient elution of two mobile phases, where one mobile phase was an aqueous ammonium phosphate buffer (pH 8)/acetonitrile mixture and the other mobile phase was acetonitrile in water. Detection
15 was performed at 215 nm.

Assay (IV): Determination of sum of impurities of semaglutide

 Determination of purity of semaglutide is performed using reversed phase high performance liquid chromatography (RP-HPLC) using a Kinetex C18 column with an isocratic
20 elution followed by a gradient elution of two mobile phases, where one mobile phase was an aqueous phosphate buffer/acetonitrile mixture and the other mobile phase was an aqueous acetonitrile/isopropanol mixture. Detection was performed at 210 nm. Purity of semaglutide is given as sum of impurities in % as the combined area of all chromatographic peaks relative to semaglutide monomer peaks.

25

Example 1: Semaglutide

 Compositions comprising semaglutide were tested in this experiment. The tested compositions contained semaglutide (as specified in Table 1), propylene glycol (14 mg/ml), disodiumhydrogenphosphate dihydrate (1.42 mg/ml), and optionally phenol (5.5 mg/ml) (as
30 specified in Table 1), at pH 7.4 in an aqueous solution. These compositions were prepared as described herein in the section General Methods of Preparation. Chemical stability as expressed by HMWP was determined by Assay (I) described herein at start of the experiment and after storage at 25°C, 30°C or at 37°C. Physical stability as expressed by Thioflavin T (ThT) assay was determined by Assay (II) described herein.

The results are given in Tables 2 and 3. Surprisingly, these results show that physical and chemical stability of semaglutide were improved in compositions without phenol relative to those with phenol. Results shown in Table 3 are an average of 8 samples tested.

5

Table 1. Compositions tested in Example 1

Composition no.	Description
1	Semaglutide 1 mg/ml, with phenol
2	Semaglutide 1 mg/ml, without phenol
3	Semaglutide 1.34 mg/ml, with phenol
4	Semaglutide 1.34 mg/ml, without phenol
5	Semaglutide 0.5 mg/ml, without phenol
6	Semaglutide 0.5 mg/ml, with phenol
7	Semaglutide 1.0 mg/ml, without phenol
8	Semaglutide 1.0 mg/ml, with phenol
9	Semaglutide 2.0 mg/ml, without phenol
10	Semaglutide 2.0 mg/ml, with phenol

Table 2. Chemical stability of semaglutide compositions, as expressed by content of high molecular weight proteins (HMWP), following storage at different temperatures. A lower HMWP concentration corresponds to a better chemical stability.

10

Composition no.	HMWP (%)			
	0 months	25°C 6 months	30°C 3 months	37°C 3 months
1	0.1	2.0	1.9	4.1
2 (no phenol)	0.1	0.3	0.3	0.5
3	0.1	1.9	1.8	3.9
4 (no phenol)	0.1	0.3	0.4	0.6

Table 3. Physical stability of semaglutide compositions as expressed by Thioflavin T (ThT) assay. A longer lag time corresponds to a better physical stability.

Composition no.	Lag time (hours)
5 (no phenol)	> 117

6	19
7 (no phenol)	> 117
8	35
9 (no phenol)	> 117
10	35

Example 2 (Reference): Liraglutide

The results of Example 1 are also surprising in view of the fact that the GLP-1 compound liraglutide – contrary to semaglutide – is less chemically stable in a composition without phenol. These results are shown in Table 5.

The results in Table 5 were obtained as follows: Compositions comprising liraglutide were tested. The tested compositions contained liraglutide (as specified in Table 4), mannitol (36.9 mg/ml), disodium hydrogen phosphate (1.42 mg/ml), and optionally phenol (as specified in Table 4), at pH 7.4 in an aqueous solution. These compositions were prepared as described herein in the section General Methods of Preparation. Chemical stability as expressed by purity was determined by Assay (III) described herein at start of the experiment and after storage at 25°C or at 37°C.

Table 4. Compositions tested in Example 2

Composition no.	Description
11	Liraglutide (3 mg/ml), without phenol (pH 7.4)
12	Liraglutide (3 mg/ml), phenol (0.04 mg/ml) (pH 7.4)
13	Liraglutide (3 mg/ml), phenol (0.16 mg/ml) (pH 7.4)
14	Liraglutide (3 mg/ml), phenol (0.8 mg/ml) (pH 7.4)
15	Liraglutide (3 mg/ml), phenol (2.5 mg/ml) (pH 7.4)

Table 5. Chemical stability, as expressed by purity, of compositions comprising liraglutide following storage at different temperatures. A higher purity corresponds to a better chemical stability.

Composition no.	Purity (%)		
	0 months	3 months at 25°C	3 months at 37°C
11 (no phenol)	98	88	72
12	98	93	80

13	98	94	81
14	97	95	83
15	98	95	84

Example 3: Semaglutide - additional experiments

Compositions comprising semaglutide were tested in this experiment. The tested compositions contained semaglutide, isotonic agent (propylene glycol (14 mg/ml) or sodium chloride (6.3 mg/ml)), optionally buffer (disodiumhydrogenphosphate dihydrate (1.42 mg/ml) or trisodiumcitrate dihydrate (2.35 mg/ml)), and optionally phenol (5.5 mg/ml or 0.1 mg/ml), at pH 7.0, 7.4 or 7.8 in an aqueous solution; details of each composition tested is shown in Table 6. The compositions were prepared as described herein in the section General Methods of Preparation. Chemical stability as expressed by HMWP was determined by Assay (I) and as expressed by sum of impurities was determined by Assay (IV) described herein at start of the experiment and after storage at 30°C. Physical stability as expressed by Thioflavin T (ThT) assay was determined by Assay (II) described herein.

The results are given in Table 7 and 8. In line with the results of Example 1, these results show that physical stability and chemical stability of semaglutide were improved in compositions without or with low phenol concentration relative to those with phenol at 5.5 mg/ml. The results show that physical stability and chemical stability of semaglutide were also improved in compositions without phenol comprising either the buffer trisodiumcitrate dihydrate or no buffer or isotonic agent sodium chloride, relative to those with phenol. Chemical and physical stability were improved for compositions with 0.1 mg/ml phenol relative to compositions with 5.5 mg/ml phenol and similar to compositions with no phenol. This was demonstrated for compositions with pH 7.0-7.8 and semaglutide concentration 0.1-10 mg/ml.

Table 6. Compositions tested in Example 3

Comp. No.	Content of composition				
	Semaglutide (mg/ml)	Phenol (mg/ml)	Buffer	Isotonic agent	pH
1	0.5	0	Phos*	PG**	7.0
2	0.5	0.1	Phos	PG	7.0
3	0.5	5.5	Phos	PG	7.0

4	0.5	0	Phos	PG	7.4
5	0.5	0.1	Phos	PG	7.4
6	0.5	5.5	Phos	PG	7.4
7	0.5	0	Phos	PG	7.8
8	0.5	0.1	Phos	PG	7.8
9	0.5	5.5	Phos	PG	7.8
10	10	0	Phos	PG	7.0
11	10	0.1	Phos	PG	7.0
12	10	5.5	Phos	PG	7.0
13	10	0	Phos	PG	7.4
14	10	0.1	Phos	PG	7.4
15	10	5.5	Phos	PG	7.4
16	10	0	Phos	PG	7.8
17	10	0.1	Phos	PG	7.8
18	10	5.5	Phos	PG	7.8
19	0.1	0	Phos	PG	7.4
20	0.1	5.5	Phos	PG	7.4
21	0.5	0	Phos	Citrate***	7.4
22	0.5	5.5	Phos	Citrate	7.4
23	0.5	0	Phos	None [#]	7.4
24	0.5	5.5	Phos	None	7.4
25	0.5	0	NaCl ^{##}	PG	7.4
26	0.5	5.5	NaCl	PG	7.4

*) Phos: Disodiumhydrogenphosphate dihydrate, 1.42 mg/ml. **) PG: Propylene glycol, 14 mg/ml. ***) Citrate: Trisodiumcitrate dihydrate, 2.35 mg/ml. [#]) None: No pharmaceutical excipients added in the form of a buffer. ^{##}) NaCl: Sodium chloride, 6.3 mg/ml.

- 5 **Table 7.** Chemical stability of semaglutide compositions, as expressed by content of high molecular weight proteins (HMWP) and sum of impurities, following storage at 30°C temperature. A lower HMWP concentration and sum of impurities concentration corresponds to a better chemical stability.

Composition No.	DS (mg/ml)	Phenol (mg/ml)	Chemical Stability			
			HMWP (%)		Sum of impurities (%)	
			0 months	30°C	0 months	30°C

				3 months		3 months
1 (pH 7.0)	0.5	0	0.1	0.3	3.1	7.0
2 (pH 7.0)	0.5	0.1	0.1	0.3	3.2	7.2
3 (pH 7.0)	0.5	5.5	0.1	1.4	3.2	7.8
4 (pH 7.4)	0.5	0	0.1	0.3	3.1	6.7
5 (pH 7.4)	0.5	0.1	0.1	0.3	3.2	6.6
6 (pH 7.4)	0.5	5.5	0.1	2.4	3.2	8.4
7 (pH 7.8)	0.5	0	0.1	0.2	3.1	6.5
8 (pH 7.8)	0.5	0.1	0.1	0.3	3.2	6.6
9 (pH 7.8)	0.5	5.5	0.1	4.8	3.1	10.6
10 (pH 7.0)	10	0	0.1	1.4	3.1	8.4
11 (pH 7.0)	10	0.1	0.1	0.7	3.1	7.7
12 (pH 7.0)	10	5.5	0.1	N/A ¹	3.0	N/A ¹
13 (pH 7.4)	10	0	0.1	0.9	3.1	7.8
14 (pH 7.4)	10	0.1	0.1	0.7	3.1	6.9
15 (pH 7.4)	10	5.5	0.1	0.8	3.0	6.9
16 (pH 7.8)	10	0	0.1	0.9	3.0	6.6
17 (pH 7.8)	10	0.1	0.1	0.6	3.1	6.8
18 (pH 7.8)	10	5.5	0.1	1.0	3.1	6.9
19 (low DS)	0.1	0	0.1	0.2	3.5	7.7
20 (low DS)	0.1	5.5	0.1	4.7	3.7	11.4
21 (citrate)	0.5	0	0.1	0.2	3.1	6.2
22 (citrate)	0.5	5.5	0.1	2.2	3.2	7.7
23 (no buffer)	0.5	0	0.1	0.2	3.2	6.9
24 (no buffer)	0.5	5.5	0.1	2.3	3.2	9.3
25 (NaCl)	0.5	0	0.1	0.3	3.1	6.4
26 (NaCl)	0.5	5.5	0.1	3.4	3.2	8.9

DS: Semaglutide. ¹Not physically stable > 1 month at 30°C.

Table 8. Physical stability of semaglutide compositions as expressed by Thioflavin T (ThT) assay. A longer lag time corresponds to a better physical stability.

Composition No.	Semaglutide (mg/ml)	Phenol (mg/ml)	Lag time (hours)
1 (pH 7.0)	0.5	0	42

2 (pH 7.0)	0.5	0.1	63
3 (pH 7.0)	0.5	5.5	5
4 (pH 7.4)	0.5	0	>117
5 (pH 7.4)	0.5	0.1	>117
6 (pH 7.4)	0.5	5.5	87
7 (pH 7.8)	0.5	0	>117
8 (pH 7.8)	0.5	0.1	>117
9 (pH 7.8)	0.5	5.5	>117
10 (pH 7.0)	10	0	117
11 (pH 7.0)	10	0.1	>117
12 (pH 7.0)	10	5.5	25
13 (pH 7.4)	10	0	>117
14 (pH 7.4)	10	0.1	>117
15 (pH 7.4)	10	5.5	>117
16 (pH 7.8)	10	0	>117
17 (pH 7.8)	10	0.1	>117
18 (pH 7.8)	10	5.5	>117
19 (low DS)	0.1	0	>117
20 (low DS)	0.1	5.5	>117
21 (citrate)	0.5	0	>117
22 (citrate)	0.5	5.5	>117
23 (no buffer)	0.5	0	>117
24 (no buffer)	0.5	5.5	4
25 (NaCl)	0.5	0	>117
26 (NaCl)	0.5	5.5	8

Results are an average of 8 samples tested. DS: Semaglutide.

While certain features of the invention have been illustrated and described herein,
5 many modifications, substitutions, changes, and equivalents will now occur to those of
ordinary skill in the art. It is, therefore, to be understood that the appended claims are
intended to cover all such modifications and changes as fall within the true spirit of the
invention.

CLAIMS

1. A liquid pharmaceutical composition comprising semaglutide and no more than 0.01
%(w/w) phenol, wherein said composition
 - a. is for parenteral administration;
 - 5 b. is an aqueous solution comprising at least 60 %w/w water; or
 - c. further comprises one or more pharmaceutically acceptable excipients selected
from the group consisting of a buffer or an isotonic agent.
2. The composition according to claim 1, wherein said composition comprises no phenol.
3. The composition according to any one of the preceding claims, wherein said composition
10 is an aqueous solution comprising at least 60 %w/w water.
4. The composition according to any one of the preceding claims, wherein the concentration
of semaglutide is 0.5-10 mg/ml or 0.01-5 mg/ml of said composition.
5. The composition according to any one of the preceding claims, wherein said composition
comprises one or more pharmaceutically acceptable excipients.
- 15 6. The composition according to claim 5, wherein said one or more pharmaceutically
acceptable excipients are selected from the group consisting of a buffer or an isotonic
agent.
7. The composition according to any one of the preceding claims, wherein said composition
comprises no preservative.
- 20 8. The composition according to any one of the preceding claims, wherein said composition
has a pH in the range of 6.0-10.0, such as pH 7.0-7.8.
9. The composition according to any one of the preceding claims, wherein said composition
is for parenteral administration.
10. The composition according to claim 9, wherein said parenteral administration is
25 subcutaneous administration.
11. A kit comprising a liquid pharmaceutical composition comprising semaglutide and no
more than 0.01 %(w/w) and instructions for use.
12. A kit comprising a liquid pharmaceutical composition comprising semaglutide and no
more than 0.01 %(w/w) phenol and an injection device for administration of said
30 composition to a subject, wherein said injection device is selected from the group
consisting of a durable pen and a prefilled pen.
13. The kit according to claim 11 or 12, wherein said pharmaceutical composition is as
defined in any one of claims 1-10.
14. A pharmaceutical composition as defined in any one of the preceding claims for use in
35 medicine.

15. The pharmaceutical composition as defined in any one of the preceding claims for use in the treatment of diabetes or obesity.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/072835

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K9/00 A61K38/00 A61K47/10 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, EMBASE, WPI Data, BIOSIS, FSTA		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2014/060472 A1 (NOVO NORDISK AS [DK]) 24 April 2014 (2014-04-24) examples 1,2 table 2	1-15
A	----- WO 2006/097537 A2 (NOVO NORDISK AS [DK]; LAU JESPER [DK]; DOERWALD FLORENCIO ZARAGOZA [DK]) 21 September 2006 (2006-09-21) cited in the application page 63, line 10 - line 15 ----- -/--	1-15
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
14 September 2018		11/10/2018
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer
		Schüle, Stefanie

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/072835

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Basf: "BASF Chemical Emergency Medical Guidelines", 1 January 2016 (2016-01-01), XP055427004, Retrieved from the Internet: URL:https://www.basf.com/documents/corp/en/sustainability/employees-and-society/employees/occupational-medicine/medical-guidelines/Phenol_B_BASF_medGuidelines_E014.pdf [retrieved on 2017-11-20] the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/072835

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014060472 A1	24-04-2014	CN 104717972 A EP 2908844 A1 JP 2016500682 A US 2015273069 A1 WO 2014060472 A1	17-06-2015 26-08-2015 14-01-2016 01-10-2015 24-04-2014
WO 2006097537 A2	21-09-2006	AT 476446 T AU 2006224536 A1 BR PI0607762 A2 CA 2601784 A1 CN 101133082 A CN 104017062 A CN 104402989 A DK 1863839 T3 DK 2322546 T3 EP 1863839 A2 EP 2322546 A1 ES 2350051 T3 ES 2557313 T3 HU E028194 T2 HU S1800019 I1 IL 184051 A JP 4585037 B2 JP 5209463 B2 JP 2008533105 A JP 2010116407 A JP 2013063984 A KR 20070120089 A PT 1863839 E PT 2322546 E TW I362392 B TW 200942255 A US 2009156478 A1 US 2012295847 A1 WO 2006097537 A2 ZA 200707261 B	15-08-2010 21-09-2006 23-03-2010 21-09-2006 27-02-2008 03-09-2014 11-03-2015 22-11-2010 14-12-2015 12-12-2007 18-05-2011 17-01-2011 25-01-2016 28-12-2016 28-06-2018 31-08-2015 24-11-2010 12-06-2013 21-08-2008 27-05-2010 11-04-2013 21-12-2007 10-11-2010 22-01-2016 21-04-2012 16-10-2009 18-06-2009 22-11-2012 21-09-2006 28-05-2008