



(51) International Patent Classification:

C07K 16/28 (2006.01) A61K 39/00 (2006.01)

(21) International Application Number:

PCT/EP2019/064942

(22) International Filing Date:

07 June 2019 (07.06.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

18176888.8	09 June 2018 (09.06.2018)	EP
18176889.6	09 June 2018 (09.06.2018)	EP
19159321.9	26 February 2019 (26.02.2019)	EP

(71) Applicant: **BOEHRINGER INGELHEIM INTERNATIONAL GMBH** [DE/DE]; Binger Strasse 173, 55216 INGELHEIM AM RHEIN (DE).


(72) Inventors: **HIPP, Susanne**; Boehringer Ingelheim International GmbH, Corporate Patents, Binger Strasse 173, 55216 INGELHEIM AM RHEIN (DE). **ADAM, Paul**; Boehringer Ingelheim International GmbH, Corporate Patents, Binger Strasse 173, 55216 INGELHEIM

AM RHEIN (DE). **DZIEGELEWSKI, Michael**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **GANESAN, Rajkumar**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **GORMAN, Philip Nicholas**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **GUPTA, Priyanka**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **GUPTA, Pankaj**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **SCHEER, Justin**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US). **VOYNOV, Vladimir H.**; c/o VP, IP, Legal, Boehringer Ingelheim USA Corp. 900 Ridgebury Road, P.O. Box 368, Ridgefield, Connecticut 06877-0368 (US).

(74) Agent: **SIMON, Elke Anna Maria** et al.; Boehringer Ingelheim International GmbH, Corporate Patents Binger Strasse 173, 55216 INGELHEIM AM RHEIN (DE).

(54) Title: DLL3-CD3 BISPECIFIC ANTIBODIES

Fig. 1



(57) Abstract: The present disclosure relates to DLL3-CD3 bispecific antibodies, characterized by their sequences. The application further relates to uses of the antibodies and their compositions, in particular for therapeutic purposes in the field of cancer diseases.



(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

DLL3-CD3 BISPECIFIC ANTIBODIES

BACKGROUND OF THE INVENTION

5 TECHNICAL FIELD

The present invention relates to multi-specific binding proteins comprising a first antigen binding unit specific for DLL3 and a second antigen binding unit specific for CD3. The invention also relates to nucleic acids encoding such binding proteins, to methods for preparing such binding proteins; host cells expressing or capable of expressing such
10 binding proteins, compositions comprising such binding proteins and to uses of such binding proteins or such compositions, in particular for therapeutic purposes in the field of cancer diseases.

BACKGROUND INFORMATION

15 DLL3 is a type I transmembrane protein that belongs to the Notch ligand family. DLL3 is primarily involved in somitogenesis, where it predominantly is localized in the Golgi apparatus and acts as an inhibitor of Notch signalling, in contrast to the other DLL family members, DLL1 and DLL4, which are localized on the cell surface. Only in cancer cells, where DLL3 is overexpressed, some DLL3 molecules escape to the cell surface (Dylla,
20 Molecular & Cellular Oncology 2016, Vol. 3, No. 2). DLL3 has been identified as a tumor –specific target by using several methods, e.g. LC/MS analysis of tumor tissues (WO 2014/125273), RNA sequencing (WO 2017/021349), and immunohistochemistry (Saunders et al., Sci Transl Med. 2015 Aug 26;7(302):302ra136; Saunders et al., AACR 2017; WO 2013/126746) analysis.

25 DLL3 expression has been described in neuroendocrine tumors such as large cell neuroendocrine carcinoma (LCNEC), small cell lung cancer (SCLC), small cell bladder cancer, neuroendocrine prostate cancer, neuroendocrine pancreatic cancer and glioblastoma (Saunders et al., Sci Transl Med. 2015 Aug 26;7(302):302ra136; Saunders et al., AACR 2017).

30 SCLC represents an indication of extremely high medical need. SCLC accounts for 13% of Lung Cancer diagnoses and has a worse prognosis than NSCLC. Conventional treatments

for these cancers mainly include chemotherapy, radiotherapy, surgery or combinations thereof, there are no targeted therapies available yet. While the initial response rate to chemotherapy is high, many patients quickly relapse with chemo-resistant disease, for which there are no treatment options available. Although there have been improvements in
5 treating these cancers over the last years, overall survival rates for these tumor types remain unchanged, thus, there is an unmet medical need for more targeted and potent therapies.

One approach for a targeted therapy is provided by antibody drug conjugates (ADCs). However, for DLL3 this strategy is not preferred due to low cell surface expression of
10 DLL3 and high rate of resistance to chemotherapy. As the majority of patients relapse after chemotherapy treatment and due to the low expression of DLL3 on the cell surface, targeting DLL3 with ADCs might have limitations. In addition ADC approaches often have off-target toxicities caused by free drug as a result of linker instability or degradation.

There have been attempts to combine DLL3 targeting with targeting of other proteins. For
15 example, WO2017/021349 describes a bispecific antibody construct combining binding to human DLL3 on the surface of a target cell and binding to human CD3 on the surface of a T cell. It is however unproven whether such approaches will be successful and to date, no targeted therapies for SCLC and glioblastoma as well as other DLL3 expressing tumors are available.

20 In view of the poor outlook for such cancer patients, there is a need to identify more efficacious therapies, particularly efficacious therapies with improved tolerability. Thus, it is an object of the invention to provide pharmacologically active agents, compositions and/or methods of treatment that provide certain advantages compared to the agents, compositions and/or methods currently used and/or known in the art. These advantages
25 include in vivo efficacy, improved therapeutic and pharmacological properties, increased specificity, improved safety profile, less side effects, reduced immunogenicity, and other advantageous properties such as improved ease of preparation or reduced costs of goods, higher stability/longer half-life, need for less frequent administration regimens especially as compared to candidate drugs already known in the art.

30

BRIEF SUMMARY OF THE INVENTION

The present invention is based on a bispecific T cell engaging approach employing multi-specific binding proteins with a binding arm to CD3 on T cells and a binding arm to DLL3 on the cell surface of tumor cells. Through simultaneously binding to T cells and tumor
5 cells, the T cell engagers force formation of a cytolytic synapse between the two cells and so, redirect the T cell activity selectively to the targeted tumor cells.

In one aspect, the invention provides a multi-specific binding protein comprising a first antigen binding unit specifically binding to DLL3 and a second antigen binding unit specifically binding to CD3, said first antigen binding unit specifically binding to DLL3
10 selected from the group consisting of i) to xviii):

- i) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3);
- 15 ii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3);
- 20 iii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3);
- 25 iv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3);
- 30 v) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1), SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3);

- vi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:31 (CDR1), SEQ ID NO:32 (CDR2) and SEQ ID NO:33 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1), SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3);
- 5 vii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:133 (CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:136 (CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3);
- 10 viii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:139 (CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:142 (CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3);
- 15 ix) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:145 (CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:148 (CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3);
- 20 x) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:151 (CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:154 (CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3);
- 25 xi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:157 (CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:160 (CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3);
- 30 xii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:163 (CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy chain CDRs comprising the amino acid sequences of

SEQ ID NO:166 (CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3);

xiii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3);

xiv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3);

xv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3);

xvi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3);

xvii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194 (CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3); and

xviii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3).

In some embodiments of the binding protein of the invention, the first antigen binding unit specifically binding to DLL3 comprises a first light chain variable domain and a first heavy chain variable domain and is selected from the group consisting of i) to xviii):

- 5 i) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:37 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:38;
- ii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequences of SEQ ID NO:39 and heavy chain variable domain
10 comprising the amino acid sequences of SEQ ID NO:40;
- iii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:42;
- iv) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:43 and heavy chain variable domain
15 comprising the amino acid sequence of SEQ ID NO:44;
- v) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:45 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:46;
- 20 vi) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:47 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:48;
- vii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:205 and heavy chain variable domain
25 comprising the amino acid sequence of SEQ ID NO:206;
- viii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:207 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:208;

- ix) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:209 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:210;
- 5 x) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:211 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:212;
- xi) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:213 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:214;
- 10 xii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:215 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:216;
- xiii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:217 and heavy chain variable domain
15 comprising the amino acid sequence of SEQ ID NO:218;
- xiv) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:219 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:220;
- xv) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:221 and heavy chain variable domain
20 comprising the amino acid sequence of SEQ ID NO:222;
- xvi) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:223 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:224;
- 25 xvii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:225 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:226; and
- xviii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:227 and heavy chain variable domain
30 comprising the amino acid sequence of SEQ ID NO:228.

In some embodiments of the binding protein of the invention, the second antigen binding unit specifically binding to CD3 is selected from the group consisting of i) to iii):

- i) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3);
- ii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3); and
- iii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:96 (CDR1), SEQ ID NO:97 (CDR2) and SEQ ID NO:98 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:99 (CDR1), SEQ ID NO:100 (CDR2) and SEQ ID NO:101 (CDR3).

In some embodiments of the binding protein of the invention, the second antigen binding unit specifically binding to CD3 comprises a second light chain variable domain and a second heavy chain variable domain and is selected from the group consisting of:

- i) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:68;
- ii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70; and
- iii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:102 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:103.

In some embodiments of the invention, the first antigen binding unit specifically binding to DLL3 comprises from its N- to C-terminus: a first light chain variable domain, a first light chain constant domain, a first peptide linker, a first heavy chain variable domain and a first heavy chain constant CH1 domain; and the second antigen binding unit specifically binding to CD3 comprises from its N- to C-terminus: a second light chain variable domain, a second light chain constant domain, a second peptide linker, a second heavy chain

variable domain and a second heavy chain constant CH1 domain. In some embodiments of the invention, the first and/or second peptide linker comprises 26 to 42 amino acids, preferably any one of 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, more preferably 38 amino acids. In some embodiments of the invention, the first linker
5 and/or second linker is a Gly-Ser linker, preferably comprising the amino acid sequence of SEQ ID NO:89, more preferably said first and second peptide linker comprise the same sequence.

In some embodiments, the binding protein of the invention further comprises a first and a second Fc domain, said first Fc domain covalently linked to said first antigen binding unit,
10 and said second Fc domain covalently linked to said second antigen binding unit.

In some embodiments of the invention,

- i) the first Fc domain comprises a tyrosine (Y) at position 366 [T366Y], and the second Fc domain comprises a threonine (T) at position 407 [Y407T], or
- ii) the first Fc domain comprises a tryptophan (W) at position 366 [T366W], and the
15 second Fc domain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V], or
- iii) the second Fc domain comprises a tyrosine (Y) at position 366 [T366Y], and the first Fc domain comprises a threonine (T) at position 407 [Y407T], or
- iv) the second Fc domain comprises a tryptophan (W) at position 366 [T366W], and
20 the first Fc domain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V],

preferably the first or the second Fc domain further comprises an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F]. In some embodiments, the first and/or second Fc domain further comprises an alanine at position 234 [L234A] and at
25 position 235 [L235A]. In some embodiments, the first light chain constant domain and the second light chain constant domain comprise a human kappa or lambda domain.

In preferred embodiments, the binding protein of the invention comprises a first polypeptide chain specifically binding to DLL3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75,
30 SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO: 242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO: 245, SEQ ID NO: 246, SEQ ID NO: 247, SEQ ID NO: 248, SEQ ID NO: 249, SEQ ID NO: 250, SEQ ID NO: 251, and SEQ ID NO: 252

and a second polypeptide chain specifically binding to CD3 comprising the amino acid sequence of SEQ ID NO:79, SEQ ID NO:80 or SEQ ID NO:105.

In a further aspect, the invention provides an isolated nucleic acid molecule i) encoding a first antigen binding unit and/or a second antigen binding unit of a protein of the invention, optionally further encoding a first and/or a second Fc domain, or ii) encoding a first and/or a second polypeptide chain of a protein of the invention. In further aspects provided herein are expression vectors comprising the nucleic acid molecule of the invention, host cells transfected with such expression vectors, and methods of manufacturing a protein of the invention.

10 In a further aspect of the invention, provided herein is a multi-specific binding protein comprising a first polypeptide chain specifically binding to DLL3 and a second polypeptide chain specifically binding to CD3, where the first polypeptide chain comprises a first light chain, a first linker, and a first heavy chain and the second polypeptide chain comprises a second light chain, a second linker, and a second heavy chain, preferably the
15 C-terminus of said first light chain is covalently bound to the N-terminus of said first heavy chain via said first peptide linker and the C-terminus of said second light chain is covalently bound to the N-terminus of said second heavy chain via said second peptide linker.

In some embodiments of the binding protein of the invention, the first polypeptide chain specifically binding to DLL3 comprises a light chain variable domain and heavy chain
20 variable domain comprising CDR sequences and/or VH/VL sequences as defined for the antigen binding units of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17, or DLL3#18 described herein. In some embodiments, the
25 second polypeptide chain specifically binding to CD3 comprises a light chain variable and heavy chain variable domain comprising CDR sequences and/or VH/VL sequences as defined for the antigen binding units of CD3#1, CD3#2, or CD3#3 described herein.

Further aspects, embodiments, uses and methods involving the binding proteins of the invention will become clear from the following detailed description of the invention and
30 from the appended claims.

The invention provides for novel binding proteins that allow a more efficient treatment of DLL3 expressing cancers, such as SCLC, glioblastoma or a DLL3 expressing neuroendocrine tumor.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Schematic representation of a bispecific binding protein of the invention

Figure 2: Schematic representation of the DLL3 full length protein and DLL3 domain constructs expressed on HEK293 cells for epitope domain mapping. For the DLL3 domain constructs, the transmembrane and the intracellular domains are derived from EpCAM.

10 **Figure 3A:** DLL3 epitope domain mapping of seven exemplary DLL3/CD3 binding proteins. Exemplary DLL3/CD3 binding proteins recognizing the membrane proximal peptide, EGF4, EGF1 and DSL domains of DLL3. The y-axis depicts counts, the x-axis depicts PE-A (phycoerythrin signal area).

Figure 3B: DLL3 epitope domain mapping of six exemplary DLL3/CD3 binding proteins. Exemplary DLL3/CD3 binding proteins recognizing the EGF1, EGF3, EGF4 or EGF6 of
15 DLL3. The y-axis depicts counts, the x-axis depicts PE-A (phycoerythrin signal).

Figure 3C: DLL3 epitope domain mapping of six exemplary DLL3/CD3 binding proteins. Exemplary DLL3/CD3 binding proteins recognizing the DSL domains or neither DSL, EGF, nor membrane proximal peptide of DLL3. The y-axis depicts counts, the x-axis
20 depicts PE-A (phycoerythrin signal).

Figure 4: Binding of seven exemplary DLL3/CD3 binding proteins to cell lines expressing human and cyno DLL3. The y-axis depicts counts, the x-axis depicts PE-A (phycoerythrin signal area).

Figure 5: Binding of seven exemplary DLL3/CD3 binding proteins to cell lines expressing
25 human DLL1 and DLL4, no binding to human DLL1 and DLL4 detected. The y-axis depicts counts, the x-axis depicts PE-A (phycoerythrin signal area).

Figure 6A: Binding of seven exemplary DLL3/CD3 binding proteins to SCLC cell lines and human T cells by flow cytometry analysis.

Figure 6 B: Binding of exemplary DLL3/CD3 binding proteins (directed to a peptide that
30 is neither the DSL nor the EGF1-6 nor the membrane proximal peptide) to SHP77 cells.

Figure 6 C: Binding of exemplary DLL3/CD3 binding proteins (directed against the DSL domain) to SHP77 cells.

Figure 6 D: Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF1 domain) to SHP77 cells.

5 **Figure 6 E:** Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF3 domain) to SHP77 cells.

Figure 6 F: Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF4 domain) to SHP77 cells.

10 **Figure 6 G:** Binding of exemplary DLL3/CD3 binding proteins (directed against the DSL domain) to SHP77 cells.

Figure 6 H: Binding of exemplary DLL3/CD3 binding proteins (directed against the membrane proximal peptide domain) to SHP77 cells.

Figure 6 I: Binding of exemplary DLL3/CD3 binding proteins (directed to a peptide that is neither the DSL nor the EGF1-6 nor the membrane proximal peptide) to NCI-H82 cells.

15 **Figure 6 J:** Binding of exemplary DLL3/CD3 binding proteins (directed against the DSL domain) to NCI-H82 cells.

Figure 6 K: Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF1 domain) to NCI-H82 cells.

20 **Figure 6 L:** Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF3 domain) to NCI-H82 cells.

Figure 6 M: Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF4 domain) to NCI-H82 cells.

Figure 6 N: Binding of exemplary DLL3/CD3 binding proteins (directed against the DSL domain) to NCI-H82 cells.

25 **Figure 6 O:** Binding of exemplary DLL3/CD3 binding proteins (directed against the membrane proximal peptide domain) to NCI-H82 cells.

Figure 6 P: Binding of exemplary DLL3/CD3 binding proteins (directed to a peptide that is neither the DSL nor the EGF1-6 nor the membrane proximal peptide) to T cells.

Figure 6 Q: Binding of exemplary DLL3/CD3 binding proteins (directed against the DSL domain) to T cells.

Figure 6 R: Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF1 domain) to T cells.

5 **Figure 6 S:** Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF3 domain) to T cells.

Figure 6 T: Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF4 domain) to T cells.

10 **Figure 6 U:** Binding of exemplary DLL3/CD3 binding proteins (directed against the EGF6 domain) to T cells.

Figure 6 V: Binding of exemplary DLL3/CD3 binding proteins (directed against the membrane proximal peptide domain) to T cells.

Figure 7 A: Potency in lysing cells of seven DLL3/CD3 binding proteins redirecting non-stimulated PBMCs towards human SCLC cell lines.

15 **Figure 7 B:** Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed to a peptide that is neither the DSL nor the EGF1-6 nor the membrane proximal peptide) redirecting non-stimulated T cells towards human SHP77 cells.

Figure 7 C: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the DSL domain) redirecting non-stimulated T cells towards human SHP77 cells.

20 **Figure 7 D:** Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the EGF1 domain) redirecting non-stimulated T cells towards human SHP77 cells.

Figure 7 E: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the EGF3 domain) redirecting non-stimulated T cells towards human SHP77 cells.

25 **Figure 7 F:** Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the EGF4 domain) redirecting non-stimulated T cells towards human SHP77 cells.

Figure 7 G: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the EGF6 domain) redirecting non-stimulated T cells towards human SHP77 cells.

30 **Figure 7 H:** Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the membrane proximal peptide) redirecting non-stimulated T cells towards human SHP77 cells.

Figure 7 I: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed to a peptide that is neither the DSL nor the EGF1-6 nor the membrane proximal peptide) redirecting non-stimulated T cells towards human NCI-H82 cells.

Figure 7 J: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed
5 against the DSL domain) redirecting non-stimulated T cells towards human NCI-H82 cells.

Figure 7 K: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the EGF1 domain redirecting non-stimulated T cells towards human NCI-H82 cells.

Figure 7 L: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed
10 against the EGF3 domain) redirecting non-stimulated T cells towards human NCI-H82 cells.

Figure 7 M: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the EGF4 domain) redirecting non-stimulated T cells towards human NCI-H82 cells.

Figure 7 N: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed
15 against the EGF6 domain) redirecting non-stimulated T cells towards human NCI-H82 cells.

Figure 7 O: Potency in lysing cells of exemplary DLL3/CD3 binding proteins (directed against the membrane proximal peptide redirecting non-stimulated T cells towards human
20 NCI-H82 cells.

Figure 8: *In vitro* internalization assay with two exemplary DLL3/CD3 binding proteins. DLL3/CD3 binding proteins were pre-incubated with DLL3-positive SHP77 cells for 2 and 4 hours before co-incubation for 48 hours with human PBMCs.

Figure 9: Pharmacokinetics of two exemplary DLL3/CD3 binding proteins. Mean
25 pharmacokinetic profiles of DLL3#3/CD3#1 (open squares) and DLL3#3/CD3#2 (closed squares) in male C57BL/6 mice following a single 1 mg/kg intravenous dose.

Figure 10: Anti-tumor activity of two exemplary DLL3/CD3 binding proteins. The y-axis depicts median tumor volumes and the x-axis depicts time.

Figure 11: Potency in lysing cells of an exemplary DLL3/CD3 binding protein redirecting
30 non-stimulated PBMCs towards human SHP77 and RKO-E6 cells.

Figure 12: Potency in activation of T cells in presence of SHP77 cells of an exemplary DLL3/CD3 binding protein

Figure 13: Potency in degranulation of T cells in presence of SHP77 cells of an exemplary DLL3/CD3 binding protein

5 **Figure 14A:** Potency secretion of Interferon gamma by PBMCsin presence of SHP77 cells of an exemplary DLL3/CD3 binding protein

Figure 14B: Potency secretion of MCP-1 by PBMCsin presence of SHP77 cells of an exemplary DLL3/CD3 binding protein

10 **Figure 15:** Potency in proliferation of T cells in presence of SHP77 cells of an exemplary DLL3/CD3 binding protein

Figure 16: Potency in lysing cells of an exemplary DLL3/CD3 binding protein redirecting non-stimulated pan T cells and naïve T cells towards human SHP77 cells

Figure 17: Potency in lysing cells of an exemplary DLL3/CD3 binding protein redirecting non-stimulated CD4⁺ and CD4⁺ central memory T cells towards human SHP77 cells

15 **Figure 18:** Potency in lysing cells of an exemplary DLL3/CD3 binding protein redirecting non-stimulated CD4⁺ and CD4⁺ effector memory T cells towards human SHP77 cells

Figure 19: Potency in lysing cells of an exemplary DLL3/CD3 binding protein redirecting non-stimulated CD8⁺ and CD8⁺CD45RA⁺ effector memory T cells towards human SHP77 cells

20 **Figure 20:** Potency in lysing cells of an exemplary DLL3/CD3 binding protein redirecting non-stimulated CD8⁺ and CD8⁺ memory T cells towards human SHP77

Figure 21: T cell infiltration in SHP77 xenograft tumor tissue with an exemplary DLL3/CD3 binding protein.

25 **Figure 22:** Binding of three exemplary anti-DLL3 antibodies to recombinant DLL3 protein in ELISA assay. The y-axis depicts counts, the x-axis depicts concentration of DLL3 protein.

Figure 23: Binding of six exemplary anti-DLL3 antibodies to DLL3-positive SCLC cell lines determined by flow cytometry. The y-axis depicts counts, the x-axis depicts concentration of anti-DLL3 antibody.

30 **Figure 24:** Representative staining of a SCLC tissue sample with 5 exemplary anti-DLL3 antibodies. Anti-DLL3 antibodies DLL3#1 and DLL3#5 showed that the tumor cells exhibit a punctate and/or diffuse cytoplasmic and/or membranous DLL3 staining. Images

have been electronically generated by LEICA SCN400 automated scanner with a magnification of 10x.

Figure 25: Representative staining of cell pellets from SCLC cell lines with different DLL3 expression levels. Anti-DLL3 antibody DLL3#5

5

DETAILED DESCRIPTION OF THE INVENTION

USED TERMS AND DEFINITIONS

The above and other aspects and embodiments of the invention will become clear from the further description herein, in which:

10 Unless indicated or defined otherwise, all terms used have their usual meaning in the art, which will be clear to the skilled person. Reference is for example made to the standard handbooks, such as Sambrook et al, "Molecular Cloning: A Laboratory Manual" (2nd Ed.), Vols. 1-3, Cold Spring Harbor Laboratory Press (1989); Lewin, "Genes IV", Oxford University Press, New York, (1990), and Roitt *et al.*, "Immunology" (2nd Ed.), Gower
15 Medical Publishing, London, New York (1989), as well as to the general background art cited herein. Furthermore, unless indicated otherwise, all methods, steps, techniques and manipulations that are not specifically described in detail can be performed and have been performed in a manner known *per se*, as will be clear to the skilled person. Reference is for example again made to the standard handbooks, to the general background art referred to
20 above and to the further references cited therein.

When used herein the term “comprising” and variations thereof such as “comprises” and “comprise” can be substituted with the term “containing” or “including” or “having.”

The term "sequence" as used herein (for example in terms like "heavy/light chain sequence", "antibody sequence", "variable domain sequence", "constant domain sequence"
25 or "protein sequence"), should generally be understood to include both the relevant amino acid sequence as well as nucleic acid sequences or nucleotide sequences encoding the same, unless the context requires a more limited interpretation.

An “antigen binding unit” as used herein refers to a polypeptide capable of binding to its specific target or antigen and comprising the minimal structural requirements derived from
30 an antibody (typically present in an antibody) which allow for target binding. Thus, an antigen binding unit comprises at least the presence of three light chain and three heavy

chain CDR sequences, preferably at least a light chain variable domain and a heavy chain variable domain.

The generalized structure of an antibody or immunoglobulin is well known to those of skill in the art. These molecules are heterotetrameric glycoproteins, typically of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains and are typically referred to as full length antibodies. Each light chain is covalently linked to a heavy chain by one disulfide bond to form a heterodimer, and the heterotetrameric molecule is formed through a covalent disulfide linkage between the two identical heavy chains of the heterodimers. Although the light and heavy chains are linked together by one disulfide bond, the number of disulfide linkages between the two heavy chains varies by immunoglobulin isotype. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at the N-terminus a variable domain (VH), followed by three or four (in case of IgE) constant domains (CH1, CH2, CH3, and CH4), as well as a hinge region between CH1 and CH2. Each light chain has two domains, an N-terminal variable domain (VL) and a C-terminal constant domain (CL). The VL domain associates non-covalently with the VH domain, whereas the CL domain is commonly covalently linked to the CH1 domain via a disulfide bond. Particular amino acid residues are believed to form an interface between the light and heavy chain variable domains (Chothia et al., 1985, J. Mol. Biol. 186:651-663). Variable domains are also referred to herein as variable regions or Fv and denote the part that confers specificity to an antibody for the antigen by carrying the antigen-binding site.

The "light chain variable domain" (or "light chain variable region") and "heavy chain variable domain" (or "heavy chain variable region") as used herein have the same general structure and each domain essentially consists of four framework (FR) regions whose sequences are widely conserved, which are referred to in the art and hereinbelow as "framework region 1" or "FR1"; as "framework region 2" or "FR2"; as "framework region 3" or "FR3"; and as "framework region 4" or "FR4", respectively; which framework regions are interrupted by three hypervariable regions, HVRs (or CDRs), which are referred to in the art and herein below as "complementarity determining region 1" or "CDR1"; as "complementarity determining region 2" or "CDR2"; and as "complementarity determining region 3" or "CDR3", respectively. Thus, the general structure or sequence of an immunoglobulin variable domain can be indicated as follows: FR1 - CDR1 - FR2 - CDR2 - FR3 - CDR3 - FR4. The framework regions adopt a beta-sheet conformation and

the CDRs may form loops connecting the beta-sheet structure. The CDRs in each chain are held in their three-dimensional structure by the framework regions and form together with the CDRs from the other chain the antigen binding site.

Within the context of this invention, reference to CDR's is based on the definition of CCG, also referred to as IMGT (Lefranc MP, Pommié C, Ruiz M, Giudicelli V, Foulquier E, Truong L, Thouvenin-Contet V, Lefranc G. "IMGT unique numbering for immunoglobulin and T cell receptor variable domains and Ig superfamily V-like domains." Dev Comp Immunol. 2003 Jan;27(1):55-77; Giudicelli V, Brochet X, Lefranc MP. "IMGT/V-QUEST: IMGT standardized analysis of the immunoglobulin (IG) and T cell receptor (TR) nucleotide sequences". Cold Spring Harb Protoc. 2011;2011(6):695-715. An alternative definition of CDRs known in the art is based on Chothia (Chothia and Lesk, J. Mol. Biol. 1987, 196: 901-917), together with Kabat (E.A. Kabat, T.T. Wu, H. Bilofsky, M. Reid-Miller and H. Perry, Sequence of Proteins of Immunological Interest, National Institutes of Health, Bethesda (1983)).

The term "constant domains" or "constant region" as used within the current application denotes the sum of the domains of an antibody other than the variable region. Such constant domains and regions are well known in the state of the art and e.g. described by Kabat et al. ("Sequence of proteins of immunological interest", US Public Health Services, NIH Bethesda, MD, Publication No. 91).

The "Fc part" or "Fc domain" of an antibody is not involved directly in binding of an antibody to an antigen, but exhibits various effector functions. An "Fc part/domain of an antibody" is a term well known to the skilled artisan and defined on the basis of papain cleavage of antibodies. Depending on the amino acid sequence of the constant region of their heavy chains, antibodies or immunoglobulins are divided in the classes: IgA, IgD, IgE, IgG and IgM. According to the heavy chain constant regions the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ respectively. Several of these may be further divided into subclasses (isotypes), e.g. IgG1, IgG2, IgG3, and IgG4, IgA1, and IgA2. The Fc part of an antibody is directly involved in ADCC (antibody dependent cell-mediated cytotoxicity) and CDC (complement-dependent cytotoxicity) based on complement activation, C1q binding and Fc receptor binding. Complement activation (CDC) is initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. While the influence of an antibody on the complement system is dependent on certain conditions,

binding to Clq is caused by defined binding sites in the Fc part. Such binding sites are e.g. L234, L235, D270, N297, E318, K320, K322, P331 and P329 (numbering according to EU numbering (Edelman et al, Proc Natl Acad Sci U S A. 1969 May;63(1):78-85)). Most crucial among these residues in mediating Clq and Fcγ receptor binding in IgG1 are
5 L234 and L235 (Hezareh et al., J. Virology 75 (2001) 12161- 12168, Shields et al (2001) JBC, 276 (9): 6591–6604). Antibodies of subclass IgG1 and IgG3 usually show complement activation and Clq and C3 binding, whereas IgG2 and IgG4 do not activate the complement system and do not bind Clq and C3.

The term “antibody” or “antibody molecule” (used synonymously herein) encompasses a
10 monoclonal antibody, a polyclonal antibody, a human antibody, a humanized antibody, a chimeric antibody, multispecific antibodies (e.g., bispecific antibodies), a fragment of an antibody, in particular a Fv, Fab, Fab’, or F(ab’)2 fragment, a single chain antibody, in particular a single chain variable fragment (scFv), a single chain Fab fragment (scFab), a Small Modular Immunopharmaceutical (SMIP), a domain antibody, a nanobody, a
15 diabody. The antibody may have an effector function, such as ADCC or CDC, that is usually mediated by the Fc part (antibody constant region) of the antibody, or it may have no effector function, e.g. by lacking a Fc part or having a blocked, masked Fc part, in essence a Fc part that is not or insufficiently recognized by immune cells or immune system components, like the complement system.

20 Monoclonal antibodies (mAb) are monospecific antibodies that are identical in amino acid sequence. They may be produced by hybridoma technology from a hybrid cell line (called hybridoma) representing a clone of a fusion of a specific antibody-producing B cell with a myeloma (B cell cancer) cell (Kohler G, Milstein C. Continuous cultures of fused cells secreting antibody of predefined specificity. Nature 1975;256:495-7.). Alternatively,
25 monoclonal antibodies may be produced by recombinant expression in host cells (Norderhaug L, Olafsen T, Michaelsen TE, Sandlie I. (May 1997). "Versatile vectors for transient and stable expression of recombinant antibody molecules in mammalian cells." J Immunol Methods 204 (1): 77–87; see also below). A “recombinant antibody” or “recombinant binding protein” is an antibody or binding protein which has been produced
30 by a recombinantly engineered host cell. It is optionally isolated or purified.

Full length antibodies can be treated with enzymes such as papain or pepsin to generate useful antibody fragments. Papain digestion is used to produce two identical antigen-

binding antibody fragments called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment. The Fab fragment also contains the constant domain of the light chain and the CH1 domain of the heavy chain. Pepsin treatment yields a F(ab')₂ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

5 Fab' fragments differ from Fab fragments by the presence of additional residues including one or more cysteines from the antibody hinge region at the C-terminus of the CH1 domain. F(ab')₂ antibody fragments are pairs of Fab' fragments linked by cysteine residues in the hinge region. Other chemical couplings of antibody fragments are also known.

"Fv" fragment contains a complete antigen-recognition and binding site consisting of a
10 dimer of one heavy and one light chain variable domain in tight, non-covalent association. In this configuration, the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the VH-VL dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody.

A "single-chain Fv" or "scFv" antibody fragment is a single chain Fv variant comprising
15 the VH and VL domains of an antibody where the domains are present in a single polypeptide chain. The single chain Fv is capable of recognizing and binding an antigen. The scFv polypeptide may optionally also contain a polypeptide linker positioned between the VH and VL domains in order to facilitate formation of a desired three-dimensional structure for antigen binding by the scFv (see, e.g., Pluckthun, 1994, In The Pharmacology
20 of monoclonal Antibodies, Vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315).

A "single-chain Fab" or "scFab" antibody fragment is a single chain Fab variant comprising the VL, CL, VH and CH1 domains of an antibody where the domains are present in a single polypeptide chain. The single chain Fab is capable of recognizing and
25 binding an antigen. The scFab polypeptide may optionally also contain a polypeptide linker positioned between the CL and VH domains (Hust et al (2007) BMC Biotechnology).

For application in man, it is often desirable to reduce immunogenicity of therapeutic molecules, such as antibodies or binding proteins comprising an antigen binding unit as
30 described herein, originally derived from other species, like mouse. This can be done by construction of chimeric antibodies/binding proteins, or by a process called "humanization". In this context, a "chimeric antibody"; or "chimeric antigen binding unit"

is understood to be an antibody or an antigen binding unit comprising a sequence part (e.g. a variable domain) derived from one species (e.g. mouse) fused to a sequence part (e.g. the constant domains) derived from a different species (e.g. human). In this context, a “humanized antibody”, “a humanized binding protein” or a “humanized antigen binding unit” is an antibody, a protein or antigen binding unit comprising a variable domain originally derived from a non-human species, wherein certain amino acids have been mutated to make the overall sequence of that variable domain more closely resemble a sequence of a human variable domain. Methods of humanization of antibodies are well-known in the art (Billetta R, Lobuglio AF. “Chimeric antibodies”. Int Rev Immunol. 1993;10(2-3):165-76; Riechmann L, Clark M, Waldmann H, Winter G (1988). "Reshaping human antibodies for therapy". Nature: 332:323).

An “optimized antibody” or an “optimized antigen binding unit or protein” is a specific type of humanized antibody or humanized antigen binding unit/protein which includes an immunoglobulin amino acid sequence variant, or fragment thereof, which is capable of binding to a predetermined antigen and which comprises one or more FRs having substantially the amino acid sequence of a human immunoglobulin and one or more CDRs having substantially the amino acid sequence of a non-human immunoglobulin. This non-human amino acid sequence often referred to as an "import" sequence is typically taken from an "import" antibody domain, particularly a variable domain. In general, an optimized antibody includes at least the CDRs (or HVLs) of a non-human antibody or derived from a non-human antibody, inserted between the FRs of a human heavy or light chain variable domain. It will be understood that certain mouse FR residues may be important to the function of the optimized antibodies and therefore certain of the human germline sequence heavy and light chain variable domains residues are modified to be the same as those of the corresponding mouse sequence. During this process undesired amino acids may also be removed or changed, for example to avoid deamidation, undesirable charges or lipophilicity or non-specific binding. An “optimized antibody”, an “optimized antibody fragment” or “optimized” may sometimes be referred to as “humanized antibody”, “humanized antibody fragment” or “humanized”, or as “sequence-optimized”.

Furthermore, technologies have been developed for creating antibodies or VH/VL domains based on sequences derived from the human genome, for example by phage display or use of transgenic animals (WWW. Ablexis.com/technology-alivamab.php; WO 90/05144; D. Marks, H.R. Hoogenboom, T.P. Bonnert, J. McCafferty, A.D. Griffiths and G. Winter

(1991) "By-passing immunisation. Human antibodies from V-gene libraries displayed on phage." J.Mol.Biol., 222, 581-597; Knappik et al., J. Mol. Biol. 296: 57-86, 2000; S. Carmen and L. Jermutus, "Concepts in antibody phage display". Briefings in Functional Genomics and Proteomics 2002 1(2):189-203; Lonberg N, Huszar D. "Human antibodies from transgenic mice". Int Rev Immunol. 1995;13(1):65-93.; Brüggemann M, Taussig MJ. "Production of human antibody repertoires in transgenic mice". Curr Opin Biotechnol. 1997 Aug;8(4):455-8.). Such antibodies or antigen binding units or VH/VL domains are "human antibodies," "human antigen binding units," or "human VH/VL domains" in the context of the present invention.

10 The term "human antibody", "human antigen binding unit", or "human VH/VL domain" as used herein, is intended to include antibodies, antigen binding units or VH/VL domains having variable (and constant, if applicable) regions derived from human germline immunoglobulin sequences. The human antibodies, antigen binding units, proteins or VH/VL domains of the present technology may include amino acid residues not encoded
15 by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo). However, the term "human antibody", "human antigen binding unit", or "human VH/VL domain" as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another (mammalian species), such as a mouse, rat or rabbit, have been grafted
20 onto human framework sequences. Thus, as used herein, the term "human antibody", "human antigen binding unit", or "human VH/VL domain" refer to an antibody, antigen binding unit or VH/VL domain in which substantially every part of the protein (e.g., CDR, framework, CL, CH domains (e.g., CH1, CH2, CH3), hinge, VL, VH) is substantially non-immunogenic in humans, with only minor sequence changes or variations. Such changes or
25 variations optionally and preferably retain or reduce the immunogenicity in humans or other species relative to non-modified antibodies or antigen binding units.

Thus, a human antibody, human antigen binding unit or human VH/VL domain is distinct from e.g., a chimeric or humanized antibody. It is pointed out that a human antibody, human antigen binding unit or human VH/VL domain can be produced by a non-human
30 animal or prokaryotic or eukaryotic cell that is capable of expressing functionally rearranged human immunoglobulin (e.g., heavy chain and/or light chain) genes.

The term “monomer” refers to a homogenous form of an antibody or a multispecific protein as described herein. For example, for a full-length antibody, monomer means a monomeric antibody having two identical heavy chains and two identical light chains. In the context of the present invention, a monomer means a protein of the present invention having a single antigen binding unit specific for DLL3, and a single antigen binding unit specific for CD3 as described herein. For example, a monomer of a binding protein described herein may have two chains, a first chain comprising a single chain Fab with a first antigen binding unit and optionally a first Fc domain and a second chain comprising a single chain Fab with a second antigen binding unit and optionally a second Fc domain.

An epitope is a region of an antigen that is bound by an antibody or antigen binding moiety (e.g. the antigen binding unit of the proteins described herein). The term “epitope” includes any polypeptide determinant capable of specific binding to an antibody or antigen binding moiety. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, glycan side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and/or specific charge characteristics. Conformational and non-conformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

An antigen binding molecule/protein (such as an immunoglobulin, an antibody, an antigen binding unit, or a fragment of such antigen binding molecule/protein) that can “bind”, “bind to”, “specifically bind”, or “specifically bind to”, that “has affinity for”, “is specific for” and/or that “has specificity for” a certain epitope, antigen or protein (or for at least one part, fragment or epitope thereof) is said to be “against” or “directed against” said epitope, antigen or protein or is a “binding” molecule/protein with respect to such epitope, antigen or protein.

As used herein, the terms “binding” and “specific binding” refer to the binding of the antibody or antigen binding moiety (such as an immunoglobulin, an antibody, an antigen binding unit, or a fragment of such antigen binding molecule/protein) to an epitope of the antigen in an in vitro assay, preferably in a plasmon resonance assay ((Malmqvist M., “Surface plasmon resonance for detection and measurement of antibody-antigen affinity and kinetics.”, Curr Opin Immunol. 1993 Apr;5(2):282-6.)) with purified wild-type antigen. Antibody affinity can also be measured using kinetic exclusion assay (KinExA)

technology (Darling, R.J., and Brault P-A., "Kinetic exclusion assay technology: Characterization of Molecular Interactions." ASSAY and Drug Development Technologies. 2004, Dec 2(6): 647-657).

Generally, the term "*specificity*" refers to the number of different types of antigens or
5 epitopes to which a particular antigen binding molecule/protein (such as an immunoglobulin, an antibody, an antigen binding unit, or a fragment of such antigen binding molecule/protein) can bind. The specificity of an antigen-binding molecule/protein can be determined based on its affinity and/or avidity. The affinity, represented by the equilibrium constant for the dissociation of an antigen with an antigen-binding protein
10 (K_D), is a measure for the binding strength between an epitope and an antigen-binding site on the antigen-binding molecule/protein: the lesser the value of the K_D , the stronger the binding strength between an epitope and the antigen-binding molecule/protein (alternatively, the affinity can also be expressed as the affinity constant (K_A), which is $1/K_D$). As will be clear to the skilled person (for example on the basis of the further
15 disclosure herein), affinity can be determined in a manner known *per se*, depending on the specific antigen of interest. Avidity is the measure of the strength of binding between an antigen-binding molecule/protein (such as an immunoglobulin, an antibody, an antigen binding unit, or fragment of such antigen binding molecule/protein) and the pertinent antigen. Avidity is related to both the affinity between an epitope and its antigen binding
20 site on the antigen-binding molecule/protein and the number of pertinent binding sites present on the antigen-binding molecule/protein.

The term "isolated," as used herein, refers to material that is removed from its original or native environment (e.g. the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not
25 isolated, but the same polynucleotide or polypeptide, separated by human intervention from some or all of the co-existing materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of the environment in which it is found in nature. For example, a nucleic acid,
30 protein/polypeptide molecule is considered to be "(in) essentially isolated (form)" - when compared to its native biological source and/or the reaction medium or cultivation medium from which it has been obtained - when it has been separated from at least one other component with which it is usually associated in said source or medium, such as another

nucleic acid, another protein/polypeptide, another biological component or macromolecule or at least one contaminant, impurity or minor component. In particular, a nucleic acid or protein/polypeptide molecule is considered "essentially isolated" when it has been purified at least 2-fold, in particular at least 10-fold, more in particular at least 100-fold, and up to 1000-fold or more. A nucleic acid or protein/polypeptide molecule that is "in essentially isolated form" is preferably essentially homogeneous, as determined using a suitable technique, such as a suitable chromatographical technique, e.g., polyacrylamide-gelelectrophoresis.

As used herein, the terms "identical" or "percent identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence. To determine the percent identity, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., $\% \text{ identity} = \frac{\# \text{ of identical positions}}{\text{total \# of positions (e.g., overlapping positions)}} \times 100$). In some embodiments, the two sequences that are compared are the same length after gaps are introduced within the sequences, as appropriate (e.g., excluding additional sequence extending beyond the sequences being compared). For example, when variable region sequences are compared, the leader and/or constant domain sequences are not considered. For sequence comparisons between two sequences, a "corresponding" CDR refers to a CDR in the same location in both sequences (e.g., CDR-H1 of each sequence).

The determination of percent identity or percent similarity between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. USA 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. USA 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al., 1990, J. Mol.

Biol. 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12, to obtain nucleotide sequences homologous to a nucleic acid encoding a protein of interest. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3, to obtain amino acid sequences
5 homologous to a protein of interest. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-Blast can be used to perform an iterated search which detects distant relationships between molecules (Id.). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g.,
10 XBLAST and NBLAST) can be used. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight
15 residue table, a gap length penalty of 12, and a gap penalty of 4 can be used. Additional algorithms for sequence analysis are known in the art and include ADVANCE and ADAM as described in Torellis and Robotti, 1994, Comput. Appl. Biosci. 10:3-5; and FASTA described in Pearson and Lipman, 1988, Proc. Natl. Acad. Sci. USA 85:2444-8. Within FASTA, ktup is a control option that sets the sensitivity and speed of the search. If ktup=2,
20 similar regions in the two sequences being compared are found by looking at pairs of aligned residues; if ktup=1, single aligned amino acids are examined. ktup can be set to 2 or 1 for protein sequences, or from 1 to 6 for DNA sequences. The default if ktup is not specified is 2 for proteins and 6 for DNA. Alternatively, protein sequence alignment may be carried out using the CLUSTAL W algorithm, as described by Higgins et al., 1996,
25 Methods Enzymol. 266:383-402.

The term "covalently linked" as used herein means either a direct covalent bond between residues, or an indirect association where two residues are not directly bonded but are both covalently bonded to an intermediate molecule or domain, e.g. an intermediate domain of an immunoglobulin.

30 The terms "compete" or "cross-compete" are used interchangeably herein to refer to the ability of an antibody molecule to interfere with binding of an antibody molecule, e.g., an anti-DLL3 antibody molecule of the invention, to a target, e.g., human DLL3. The interference with binding can be direct or indirect (e.g., through an allosteric modulation of

the antibody molecule or the target). The extent to which an antibody molecule is able to interfere with the binding of another antibody molecule to the target, and therefore whether it can be said to compete, can be determined using a competition binding assay, for example, a FACS assay, an ELISA or BIACORE assay. In some embodiments, a competition binding assay is a quantitative competition assay. In some embodiments, a first anti-DLL3 antibody molecule is said to compete for binding to the target with a second anti- DLL3 antibody molecule when the binding of the first antibody molecule to the target is reduced by 10% or more, e.g., 20% or more, 30% or more, 40% or more, 50% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, 99% or more in a competition binding assay (e.g., a competition assay described herein).

It is to be inferred without explicit recitation and unless otherwise intended, that when the present technology relates to a polypeptide, protein, polynucleotide or antibody, an equivalent or a biologically equivalent of such is intended within the scope of the present technology.

As used herein, the term "biological equivalent thereof" is intended to be synonymous with "equivalent thereof" when referring to a reference protein, antibody, polypeptide, polynucleotide or nucleic acid, and intends those having minimal homology while still maintaining desired structure or functionality. Unless specifically recited herein, it is contemplated that any nucleic acid, polynucleotide, polypeptide, protein or antibody mentioned herein also includes equivalents thereof. For example, an equivalent intends at least about 80 % homology or identity and alternatively, at least about 85 %, or alternatively at least about 90 %, or alternatively at least about 95 %, or alternatively 98 % percent homology or identity and exhibits substantially equivalent biological activity to the reference protein, polypeptide, antibody or nucleic acid.

As used herein, the term "detectable label" refers to a molecule, compound or composition that can produce a detectable signal, i.e. physical or chemical signal including a colorimetric, fluorescent, electrical, radioactive and chemiluminescent signal, which can be measured by visual or instrumental methods, and which indicates the presence and/or quantity/concentration of the label in a sample.

A "detectable signal" can be generated by various mechanism including absorption, emission and/or scattering of a photon (including radio frequency, microwave frequency,

infrared frequency, visible frequency and ultra-violet frequency photons) and includes but is not limited to colorimetric, fluorescent, electrical, radioactive and chemiluminescent signals.

As used herein, "expression" refers to the process by which polynucleotides are transcribed into mRNA and/or the process by which the transcribed mRNA is subsequently being translated into peptides, polypeptides, or proteins. If the polynucleotide is derived from genomic DNA, expression may include splicing of the mRNA in an eukaryotic cell. The expression level of a gene may be determined by measuring the amount of mRNA or protein in a cell or tissue sample.

As used herein the term "biological sample" means a sample material derived from or contacted by living cells. The term is intended to include tissues, cells and biological fluids isolated from a subject. As used herein, the term "tissue sample" shall refer to a cellular sample that preserves the cross-sectional spatial relationship between the cells as they existed within the subject from which the sample was obtained. Biological samples can also be obtained from biopsies of internal organs or from cancers.

"Histochemistry" and cytochemistry" are techniques used to identify a molecule within the context of intact cells by labeling the samples with an agent that binds specifically to the molecule in a manner than can be visualized on a microscope. "Immunohistochemistry" and "immunocytochemistry" are types of histochemistry and cytochemistry that use antibodies to label the molecules.

MULTI-SPECIFIC BINDING PROTEINS OF THE INVENTION

The present invention provides multi-specific binding proteins comprising at least one antigen binding unit specifically binding to DLL3 (a first antigen binding unit), and at least one antigen binding unit specifically binding to CD3 (a second antigen binding unit). Such (multi-specific) binding proteins are also referred to herein as (multi-specific) binding molecules or DLL3/CD3 binding proteins or DLL3/CD3 binding molecules.

The inventors have surprisingly found that multi-specific binding proteins of the invention induce selective lysis of DLL3-positive SCLC cell lines in the presence of T cells and are already active at low effector to target cell ratios. Importantly, the binding proteins of the

invention do not cause T cell activation, T cell proliferation, and Cytokine secretion in the absence of DLL3-positive cells or lysis of DLL3-negative cells.

For avoidance of doubt, DLL3 as used herein refers to human DLL3 of UniProt Q9NYJ7 and the nucleic acid sequence encoding that protein. CD3 as used herein refers to human
5 CD3epsilon (UniProt P07766) and CD3 gamma (Uniprot: P09693) complexes (human CD3 ϵ gamma complexes).

Targeting DLL3 with a bispecific T cell engaging approach is expected to provide advantages over an ADC approach, as redirecting T cells is not influenced by resistance to chemotherapy and low expression levels on the cell surface are less critical for this mode
10 of action. T cell engagers are multi-specific binding proteins with binding arms to CD3 on T cells and binding arms to an antigen on the cell surface of tumor cells. Through simultaneously binding to T cells and tumor cells, the T cell engagers force formation of a cytolytic synapse between the two cells and so, redirect the T cell activity selectively to the targeted tumor cells.

15 In one aspect, the multi-specific binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3 and a second antigen binding unit specifically binding to CD3, wherein said first binding unit is selected from the group consisting of i) to xviii):

- 20 i) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3) (antigen binding unit DLL3#1);
- 25 ii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3) (antigen binding unit DLL3#2);
- 30 iii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid

sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3) (antigen binding unit DLL3#3);

iv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3) (antigen binding unit DLL3#4);

v) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1), SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3) (antigen binding unit DLL3#5);

vi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:31 (CDR1), SEQ ID NO:32 (CDR2) and SEQ ID NO:33 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1), SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3) (antigen binding unit DLL3#6);

vii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:133 (CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:136 (CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3) (antigen binding unit DLL3#7);

viii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:139 (CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:142 (CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3) (antigen binding unit DLL3#8);

ix) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:145 (CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:148 (CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3) (antigen binding unit DLL3#9);

- 5 x) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:151 (CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:154 (CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3) (antigen binding unit DLL3#10);
- 10 xi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:157 (CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:160 (CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3) (antigen binding unit DLL3#11);
- 15 xii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:163 (CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:166 (CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3) (antigen binding unit DLL3#12);
- 20 xiii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3) (antigen binding unit DLL3#13);
- 25 xiv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3) (antigen binding unit DLL3#14);
- 30 xv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3) (antigen binding unit DLL3#15);
- xvi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and

SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3) (antigen binding unit DLL3#16);

5 xvii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194(CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3) (antigen binding unit DLL3#17); and

10 xviii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3) (antigen binding unit DLL3#18).

15 Preferably, the first antigen binding unit of the binding protein of the invention is any one of i) to iii) as defined by the CDR sequences above.

For the avoidance of doubt, each of the specific embodiments listed herein can also be considered independent aspects of the invention.

In some embodiments of the binding protein of the invention, said second antigen binding unit specifically binding to CD3 is selected from the group consisting of i) to iii):

20 i) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3) (antigen binding unit CD3#1);

25 ii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3) (antigen binding unit CD3#2); and

30 iii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:96 (CDR1), SEQ ID NO:97 (CDR2) and SEQ ID

NO:98 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:99 (CDR1), SEQ ID NO:100 (CDR2) and SEQ ID NO:101 (CDR3) (antigen binding unit CD3#3).

The first antigen binding units i) to xviii) as outlined above are termed DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 and DLL3#18, respectively and the second antigen binding units i) to iii) as outlined above are termed CD3#1, CD3#2 and CD3#3, respectively. Provided herein is a sequence table which readily allows identification of individual amino acid sequences to specific antigen binding units and full length binding proteins of the present invention. A summary is provided in Table 1 in Example 2.

The terms “first” and “second” with respect to antigen binding units in general, as used herein, is solely intended to indicate that these units are two different units (as they bind to different target antigens). Thus, these terms shall not be understood to refer to the exact order or sequence of the units within the binding protein of the invention.

In some embodiments, the binding protein of the invention comprises a first antigen binding unit selected from the group consisting of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 and DLL3#18 as defined by the respective CDR sequences described above and a second antigen binding unit of CD3#1 as defined by the respective CDR sequences described above. In preferred embodiments, the binding protein of the invention comprises a first antigen binding unit selected from the group consisting of DLL3#1, DLL3#2, and DLL3#3 as defined by the respective CDR sequences described above and a second antigen binding unit of CD3#1 as defined by the respective CDR sequences described above.

In some embodiments, the binding protein of the invention comprises a first antigen binding unit selected from the group consisting of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 and DLL3#18 as defined by the respective CDR sequences described above and a second antigen binding unit of CD3#2 as defined by the respective CDR sequences described above. In preferred embodiments, the binding protein of the invention comprises a first antigen binding unit selected from the

group consisting of DLL3#1, DLL3#2, and DLL3#3 as defined by the respective CDR sequences described above and a second antigen binding unit of CD3#2 as defined by the respective CDR sequences described above.

In some embodiments, the binding protein of the invention comprises a first antigen binding unit selected from the group consisting of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 and DLL3#18 as defined by the respective CDR sequences described above and a second antigen binding unit of CD3#3 as defined by the respective CDR sequences described above. In preferred embodiments, the binding protein of the invention comprises a first antigen binding unit selected from the group consisting of DLL3#1, DLL3#2, and DLL3#3 as defined by the respective CDR sequences described above and a second antigen binding unit of CD3#3 as defined by the respective CDR sequences described above.

In one preferred embodiment, the binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3) and a second antigen binding unit specifically binding to CD3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3) and a second antigen binding unit specifically binding to CD3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3) and a second antigen binding unit specifically binding to CD3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3) and a second antigen binding unit specifically binding to CD3 comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3) and a second antigen binding unit specifically binding to CD3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises a first antigen binding unit specifically binding to DLL3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3) and a second antigen

binding unit specifically binding to CD3, comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3).

- 5 In addition to the CDR sequences as set out herein, the antigen binding units of the binding proteins of the invention include immunoglobulin framework region (FR) sequences. These sequences are preferably not immunogenic in humans, and are therefore preferably human or humanized FR sequences. Suitable human or humanized FR sequences are known in the art. Specifically preferred FR sequences can be taken from the embodiments
10 shown herein, disclosing the complete antigen binding units and thereby CDR sequences as well as FR sequences.

In preferred embodiments of the binding proteins of the invention, the first and the second binding unit each comprise a light chain variable domain and a heavy chain variable domain derived from an antibody molecule, said light/heavy chain variable domains
15 defined by the CDR sequences of any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 for the first antigen binding unit and said light/heavy chain variable domains defined by the CDR sequences of any one of CD3#1, CD3#2 or CD3#3 for the second antigen binding unit. In some
20 embodiments of the binding protein of the invention, the VH and/or VL domain of the antigen binding units of any one or more of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17, DLL3#18, CD3#1, CD3#2 or CD3#3 is a human or humanized VH and/or VL domain.

- 25 In preferred embodiments of the invention, the light/heavy chain variable domains of the first antigen binding unit are further defined as follows
- i) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:37 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:38 (antigen binding unit DLL3#1); or
 - 30 ii) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:39 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:40 (antigen binding unit DLL3#2); or

- iii) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:42 (antigen binding unit DLL3#3); or
- iv) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:43 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:44 (antigen binding unit DLL3#4); or
- v) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:45 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:46 (antigen binding unit DLL3#5); or
- vi) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:47 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:48 (antigen binding unit DLL3#6); or
- vii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:205 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:206 (antigen binding unit DLL3#7); or
- viii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:207 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:208 (antigen binding unit DLL3#8); or
- ix) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:209 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:210 (antigen binding unit DLL3#9); or
- x) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:211 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:212 (antigen binding unit DLL3#10); or
- xi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:213 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:214 (antigen binding unit DLL3#11); or
- xii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:215 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:216 (antigen binding unit DLL3#12); or
- xiii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:217 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:218 (antigen binding unit DLL3#13); or

- xiv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:219 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:220 (antigen binding unit DLL3#14); or
- 5 xv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:221 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:222 (antigen binding unit DLL3#15); or
- xvi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:223 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:224 (antigen binding unit DLL3#16); or
- 10 xvii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:225 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:226 (antigen binding unit DLL3#17); or
- xviii) comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:227 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:228 (antigen binding unit DLL3#18).
- 15

Preferably, the first antigen binding unit of the binding protein of the invention is any one of i) to iii) as defined by the VL and VH sequences above.

In preferred embodiments of the invention, the light/heavy chain variable domains of the second antigen binding unit are further defined as follows

- 20 i) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:68 (antigen binding unit CD3#1); or
- ii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70 (antigen binding unit CD3#2); or
- 25 iii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:102 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:103 (antigen binding unit CD3#3).

In some embodiments, the binding protein of the invention comprises a combination of a first and a second antigen binding unit selected from the group consisting of
 30 DLL3#1/CD3#1, DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#4/CD3#1, DLL3#5/CD3#1, DLL3#6/CD3#1, DLL3#7/CD3#1, DLL3#8/CD3#1, DLL3#9/CD3#1, DLL3#10/CD3#1,

DLL3#11/CD3#1, DLL3#12/CD3#1, DLL3#13/CD3#1, DLL3#14/CD3#1,
 DLL3#15/CD3#1, DLL3#16/CD3#1, DLL3#17/CD3#1, DLL3#18/CD3#1,
 DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#4/CD3#2, DLL3#5/CD3#2,
 DLL3#6/CD3#2, DLL3#7/CD3#2, DLL3#8/CD3#2, DLL3#9/CD3#2, DLL3#10/CD3#2,
 5 DLL3#11/CD3#2, DLL3#12/CD3#2, DLL3#13/CD3#2, DLL3#14/CD3#2,
 DLL3#15/CD3#2, DLL3#16/CD3#2, DLL3#17/CD3#2,
 DLL3#18/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#3, DLL3#3/CD3#3, DLL3#4/CD3#3,
 DLL3#5/CD3#3, DLL3#6/CD3#3, DLL3#7/CD3#3, DLL3#8/CD3#3, DLL3#9/CD3#3,
 DLL3#10/CD3#3, DLL3#11/CD3#3, DLL3#12/CD3#3, DLL3#13/CD3#3,
 10 DLL3#14/CD3#3, DLL3#15/CD3#3, DLL3#16/CD3#3, DLL3#17/CD3#3,
 DLL3#18/CD3#3, the first and second antigen binding unit being defined by the CDR
 and/or VH and VL sequences of the antigen binding units as described above. In preferred
 embodiments, the binding protein of the invention comprises a combination of a first and a
 second antigen binding unit selected from the group consisting of DLL3#1/CD3#1,
 15 DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2,
 DLL3#1/CD3#3, DLL3#2/CD3#3, and DLL3#3/CD3#3, the first and second antigen
 binding unit being defined by the CDR and/or VH and VL sequences of the antigen
 binding units as described above.

In one preferred embodiment, the binding protein of the invention comprises (i) a first
 20 antigen binding unit specifically binding to DLL3 comprising a light chain variable domain
 comprising the amino acid sequences of SEQ ID NO:37 and a heavy chain variable domain
 comprising the amino acid sequences of SEQ ID NO:38 and (ii) a second antigen binding
 unit specifically binding to CD3 comprising a light chain variable domain comprising the
 amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the
 25 amino acid sequence of SEQ ID NO:68.

In one preferred embodiment, the binding protein of the invention comprises (i) a first
 antigen binding unit specifically binding to DLL3 comprising a light chain variable domain
 comprising the amino acid sequences of SEQ ID NO:39 and a heavy chain variable domain
 comprising the amino acid sequences of SEQ ID NO:40 and (ii) a second antigen binding
 30 unit specifically binding to CD3 comprising a light chain variable domain comprising the
 amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the
 amino acid sequence of SEQ ID NO:68.

In one preferred embodiment, the binding protein of the invention comprises (i) a first antigen binding unit specifically binding to DLL3 comprising a light chain variable domain comprising the amino acid sequences of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:42 and (ii) a second antigen binding unit specifically binding to CD3 comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:68.

In one preferred embodiment, the binding protein of the invention comprises (i) a first antigen binding unit specifically binding to DLL3 comprising a light chain variable domain comprising the amino acid sequences of SEQ ID NO:37 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:38 and (ii) a second antigen binding unit specifically binding to CD3 comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70.

In one preferred embodiment, the binding protein of the invention comprises (i) a first antigen binding unit specifically binding to DLL3 comprising a light chain variable domain comprising the amino acid sequences of SEQ ID NO:39 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:40 and (ii) a second antigen binding unit specifically binding to CD3 comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70.

In one preferred embodiment, the binding protein of the invention comprises (i) a first antigen binding unit specifically binding to DLL3 comprising a light chain variable domain comprising the amino acid sequences of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:42 and a second antigen binding unit specifically binding to CD3 comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70.

In some embodiments, the binding protein of the invention comprises i) a first antigen binding unit specifically binding to DLL3 (e.g. any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 as defined

by the respective CDR or VH/VL sequences above) which comprises a first light chain variable domain covalently linked, either directly or indirectly, to a first heavy chain variable domain with a first peptide linker and/or ii) a second antigen binding unit specifically binding to CD3 (e.g. any one of CD3#1, CD3#2, or CD3#3 as defined by the
 5 respective CDR or VH/VL sequences above) which comprises a second light chain variable domain covalently linked, either directly or indirectly, to a second heavy chain variable domain with a second peptide linker.

In some embodiments of the binding proteins of the invention, the first and/or the second antigen binding unit further comprises a CL and a CH1 domain like in a light/heavy Fab
 10 domain of a conventional antibody molecule, thus said first binding unit comprises a) a VL domain (e.g., defined by the light chain CDR (LCCDR) or VL sequences of any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18) covalently linked (directly or indirectly bound) to a first CL domain and b) a
 15 VH domain (e.g., defined by the heavy chain CDR (HCCDR) or VH sequences of any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18) covalently linked (directly or indirectly bound) to a first CH1 domain and/or said second antigen binding unit comprises a) a VL domain (e.g., defined by the LCCDR
 20 or VL sequences of any one of CD3#1, CD3#2 or CD3#3) covalently linked (directly or indirectly bound) to a second CL domain and b) a VH domain (e.g., defined by HCCDR or VH sequences of any one of CD3#1, CD3#2 or CD3#3) covalently linked (directly or indirectly bound) to a second CH1 domain.

In the context of the present invention, a CL domain is the constant domain of an antibody
 25 light chain, for example a kappa (κ) or a lambda (λ) light chain. An example of a constant region of a kappa light chain is shown in SEQ ID NO:87. An example of a constant region of a lambda light chain is shown in SEQ ID NO:88. In some embodiments, the first and the second CL domain are the same, e.g. the first and the second CL domain are both a kappa light chain constant domain or the first and the second CL domain are both a lambda light
 30 chain constant domain. In some embodiments, the first and the second CL domain are different, e.g., the first CL domain is a constant kappa domain and the second CL domain is a constant lambda domain or vice versa.

In the context of the present invention, a CH1 domain is the first constant domain of an antibody heavy chain. An example of a constant CH1 domain is shown in SEQ ID NO:253.

In preferred embodiments of the binding proteins of the invention, the first antigen binding unit (e.g., any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 defined by the CDR and/or VH/VL sequences as outlined above) of the binding proteins of the invention comprises from N- to C-terminus: a first light chain variable domain, a first CL domain, a first linker peptide, a first VH domain and a first CH1 domain, and/or the second binding unit (e.g., any one of CD3#1, CD3#2 or CD3#3 defined by the CDR and/or VH/VL sequences as outlined above) of the binding proteins of the invention comprises from N- to C-terminus: a second light chain variable domain, a second CL domain, a second linker peptide, a second VH domain and a second CH1 domain. In this embodiment, the first and/or the second binding unit have the structure of a single chain Fab. For both, the first and/or the second antigen binding unit, when forming a single chain Fab, the order can be reversed such that from N- to C-terminus the antigen binding unit comprises: VH-CH1-[linker peptide]-VL-CL. In some embodiments of the protein of the invention when the first and/or second antigen binding unit comprise a Fab or a single chain Fab, the constant domains can be of the same type (e.g. both CL domains are kappa or lambda light chain constant domains) or of different types (the first CL domain is a kappa and the second CL domain is a lambda light chain constant domain or vice versa).

In one aspect, a linker used in a binding protein of the present invention comprises 26 to 42 amino acids, for example 30 to 40 amino acids. In a further aspect, a linker used in a protein of the present invention comprises 34 to 40 amino acids, for example 36 to 39 amino acids, for example 38 amino acids. In some embodiments, the linker comprises a sequence of any one of SEQ ID NOs: 89, 90, 91, 92, 93, 94, or 95, preferably SEQ ID NO: 89.

The linker sequence may be a naturally occurring sequence or a non-naturally occurring sequence. If used for therapeutic purposes, the linker is preferably non-immunogenic in the subject to which the binding protein of the invention is administered.

One useful group of linker sequences are linkers derived from the hinge region of heavy chain antibodies as described in WO1996/34103 and WO1994/04678. Other examples are poly-alanine linker sequences such as Ala-Ala-Ala.

Further preferred examples of linker sequences are Gly/Ser linkers of different length such as (glyxsery)_z linkers, including e.g. (gly4ser)₃, (gly4ser)₅, (gly4ser)₇, (gly3ser)₃,
 5 (gly3ser)₅, (gly3ser)₇, (gly3ser2)₃, (gly3ser2)₅, and (gly3ser2)₇ or a linker of any one of SEQ ID Nos: 89 to 95 .

In some embodiments of the binding proteins of the invention, the VL domain of the first antigen binding unit (e.g., defined by the light chain CDR (LCCDR) or VL sequences of
 10 DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18) is covalently linked (e.g., directly bound) via a first Gly/Ser linker (e.g., Gly/Ser linker of any one of 26 to 42 amino acids, 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, preferably 38 amino acids) to the VH domain of the first
 15 antigen binding unit (e.g., defined by the heavy chain CDR (HCCDR) or VH sequences of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18); and the VL domain of the second antigen binding unit (e.g., defined by the light chain CDR (LCCDR) or VL sequences of CD3#1, CD3#2 or CD3#3) is covalently
 20 linked (e.g. directly bound) via a second Gly/Ser linker (e.g., Gly/Ser linker of any one of 26 to 42 amino acids, 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, preferably 38 amino acids) to the VH domain of the second antigen binding unit (e.g., defined by the heavy chain CDR (HCCDR) or VH sequences of CD3#1, CD3#2 or CD3#3). More preferably, the first and the second linker are the same. Even more
 25 preferably, the first and the second linker each comprise the amino acid sequence of SEQ ID NO:89.

In some embodiments, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit specific for DLL3 and comprising the sequence selected from the group consisting of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51,
 30 SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, and SEQ ID NO:240 and

a second single chain Fab of SEQ ID NO:71 forming a second antigen binding unit specific for CD3. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:49 and a second single chain Fab comprising the sequence of SEQ ID NO:71 forming a second antigen binding unit. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:50 and a second single chain Fab comprising the sequence of SEQ ID NO:71 forming a second antigen binding unit. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:51 and a second single chain Fab comprising the sequence of SEQ ID NO:71 forming a second antigen binding unit.

In some embodiments, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit specific for DLL3 and comprising the sequence selected from the group consisting of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, and SEQ ID NO:240 and a second single chain Fab of SEQ ID NO:72 forming a second antigen binding unit specific for CD3. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:49 and a second single chain Fab comprising the sequence of SEQ ID NO:72 forming a second antigen binding unit. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:50 and a second single chain Fab comprising the sequence of SEQ ID NO:72 forming a second antigen binding unit. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:51 and a second single chain Fab comprising the sequence of SEQ ID NO:72 forming a second antigen binding unit.

In some embodiments, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit specific for DLL3 and comprising the sequence selected from the group consisting of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51,

SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54 SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, and SEQ ID NO:240 and a second single chain Fab of SEQ ID NO:104 forming a second antigen binding unit

5 specific for CD3. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:49 and a second single chain Fab comprising the sequence of SEQ ID NO:104 forming a second antigen binding unit. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen

10 binding unit comprising the sequence of SEQ ID NO:50 and a second single chain Fab comprising the sequence of SEQ ID NO:104 forming a second antigen binding unit. In a preferred embodiment, the binding protein of the invention comprises a first single chain Fab forming a first antigen binding unit comprising the sequence of SEQ ID NO:51 and a second single chain Fab comprising the sequence of SEQ ID NO:104 forming a second

15 antigen binding unit.

In some embodiments, the first antigen binding unit (e.g., DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 as defined by the CDR and/or VH/VL sequences outlined above) and/or the second antigen binding

20 unit (e.g. CD3#1, CD3#2 or CD3#3) comprises a VL domain covalently linked (e.g. directly bound) to a CL domain and a VH domain linked to a CH1 domain (i.e. together forming a Fab fragment derived from an antibody), and said CH1 domain is further covalently linked (e.g. directly bound) to an Fc domain thereby forming an arm of a conventional Y shaped antibody molecule with one light and one heavy chain. In some

25 embodiments, the first and the second antigen binding unit each form a Fab fragment, i.e. a first and a second Fab fragment, which is covalently linked (e.g. directly bound) to a first and a second Fc domain, respectively, thereby forming a conventional heterotetrameric bispecific and bivalent antibody molecule.

In preferred embodiments, the first antigen binding unit (e.g., any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 as defined by the CDR and/or VH/VL sequences outlined above) and/or the second antigen binding unit (e.g. any one of CD3#1, CD3#2 or CD3#3) comprise a single chain Fab, i.e.

30

an antibody light chain (VL-CL) covalently linked to the VH-CH1 domain of a heavy chain via a peptide linker (e.g., Gly/Ser linker of any one of 26 to 42 amino acids, 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, preferably 38 amino acids, even more preferably a linker of SEQ ID NO:89). In preferred embodiments, the binding protein of the invention comprises a first polypeptide chain comprising a first single chain Fab specifically binding to DLL3 (e.g., any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 as defined by the respective CDR or VH/VL sequences above, preferably the first antigen binding unit is DLL3#1, DLL3#2, DLL3#3) and a first Fc domain (this polypeptide chain herein referred to also as "DLL3 chain") and a second polypeptide chain comprising a second single chain Fab specifically binding to CD3 (e.g. any one of CD3#1 CD3#2 or CD3#3 as defined by the respective CDR or VH/VL sequences above) and a second Fc domain (this polypeptide chain herein referred to herein also as "CD3 chain"). In some embodiments, the first and the second Fc domain are the same. In preferred embodiments, the first and the second Fc domains are different. The resulting binding proteins of the invention bear a full Fc and have two independent binding sites, a first binding unit for DLL3 and a second binding unit for CD3. In some embodiments, the first antigen binding unit consists of a single polypeptide chain (a DLL3 chain). In some embodiments, the second antigen binding unit consists of a single polypeptide chain (a CD3 chain). In some embodiments, both the first and the second antigen binding unit consist of a single polypeptide chain, respectively.

In some embodiments, the first antigen binding unit of the protein of the invention is formed by a first polypeptide chain (a DLL3 chain) and the second antigen binding unit is formed by a second polypeptide chain (a CD3 chain). Thus, in some embodiments, the binding protein of the invention comprises two different polypeptide chains, each comprising an antigen binding unit with different specificity, the polypeptide chains covalently linked to each other, either via disulfide bonds or potentially via a peptide linker. In some embodiments, the binding protein of the invention is a bispecific, bivalent heterodimeric protein comprising two polypeptide chains, one polypeptide chain (a first polypeptide chain or DLL3 chain) comprising an antigen binding unit specifically binding to DLL3 (e.g. any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 defined by the respective CDR and/or

VH/VL sequences) and another polypeptide chain (a second polypeptide chain or CD3 chain) comprising an antigen binding unit specifically binding to CD3 (e.g. any one of CD3#1, CD3#2 or CD3#3).

In the context of the present invention, an Fc domain is for example derived from the heavy chain of an IgG, for example an IgG₁, IgG₂ or IgG₄. For example, an Fc domain of the present invention is a Fc domain of a heavy chain of an IgG₁ or IgG₄ and comprises a hinge region and two constant domains (C_{H2} and C_{H3}). Examples of Fc domains (including a hinge region) are shown in SEQ ID NOs:81 and 84.

The numbering of the amino acids in the amino acid chains of a protein of the present invention is herein according to the EU numbering system (Edelman, Cunningham et al. 1969), unless otherwise specified. This means that the amino acid numbers indicated herein correspond to the positions in a heavy chain of the corresponding sub-type (e.g. IgG₁ or IgG₄), according to the EU numbering system, unless otherwise specified.

In some embodiments, the first Fc domain and the second Fc domain in a protein of the present invention each comprises one or more amino acid changes which reduce the formation of homodimers of the first or second polypeptide chains instead of heterodimers of a first and a second polypeptide chain. Through these changes, a "protrusion" is generated in one of the Fc domains by replacing one or more, small amino acid side chains from the interface of one of the heavy chains with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size are created on the interface of the other Fc domain by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers, in particular homodimers of the Fc domain with the "protrusion" (see for example Ridgway et al. Protein Eng, 1996, 9(7): p. 617-21; Atwell et al, JMB, 1997, 270, 26-35). In some embodiments, such amino acid changes are a tyrosine (Y) at position 366 [T366Y] of the first Fc domain and a threonine (T) at position 407 [Y407T] of the second Fc domain. In some embodiments, the first Fc domain comprises a serine (S) at position 366 [T366S] and the second Fc domain comprises a tryptophan (W) at position 366 [T366W], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V]. In preferred embodiments, the first Fc domain comprises a tryptophan (W) at position 366 [T366W] and the second Fc domain comprises a serine (S) at position 366 [T366S], an alanine (A) at

position 368 [L368A] and a valine (V) at position 407 [Y407V]. For example, position 366 of the Fc domain according to EU numbering, corresponding to the amino acid position 146 in the human IgG1 Fc sequence of SEQ ID NO:81, is changed from T at position 146 in SEQ ID NO:81 to W at position 146 in SEQ ID NO:82; and positions 366, 368 and 407 according to EU numbering, corresponding to the amino acid positions 146, 148 and 187, respectively, in SEQ ID NO:81, are changed from T, L and Y at these positions in SEQ ID NO:81 to S, A and V at these positions in SEQ ID NO:83. In any of these embodiments, the amino acid changes described for the first Fc domain may be located in the second Fc domain and the respective amino acid changes for the second Fc domain may be located in the first Fc domain. In other words, the term “first” and “second” can be exchanged in these embodiments. In some embodiments, such a Fc domain is an Fc domain derived from the heavy chain of an IgG₁ or IgG₄.

In some embodiments, the first Fc domain comprises a cysteine (C) at position 354 [S354C] in addition to the tryptophan (W) at position 366 [T366W] and the second Fc domain comprises a cysteine (C) at position 349 [Y349C] in addition to the serine (S) at position 366 [T366S], the alanine (A) at position 368 [L368A] and the valine (V) at position 407 [Y407V]. In one aspect, such Fc domain is an Fc domain derived from the heavy chain of an IgG₄.

In some embodiments, the first Fc domain or the second Fc domain in a binding protein of the present invention further comprises one or more amino acid changes which reduce the binding of the Fc domain to protein A. In some embodiments, such amino acid changes are an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F] of one of the Fc domains. Both changes are derived from the sequence of human IgG3 (IgG3 does not bind to protein A). These two mutations are located in the CH3 domain and are incorporated in one of the Fc domains to reduce binding to Protein A (see for example Jendeberg et al. J Immunol Methods, 1997. 201(1): p. 25-34). These two changes facilitate the removal of homodimers of heavy chains comprising these changes during protein purification.

In some embodiments, in a binding protein of the present invention, the Fc domain, which comprises a threonine (T) at position 407 [Y407T], further comprises an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F]. In this case, the other heavy chain comprises a tyrosine (Y) at position 366 [T366Y], but does not include the

two changes at positions 435 and 436. Alternatively, in some embodiments, in a protein of the present invention, the Fc domain, which comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V], further comprises an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F]. In this case, the other Fc domain comprises a tryptophan (W) at position 366 [T366W], but does not include the two changes at positions 435 and 436. Thus, the Fc domain comprising the amino acid change resulting in a “cavity” as described above also comprises the amino acid changes, which reduce binding to Protein A. Homodimers comprising this Fc domain are removed through reduced binding to Protein A. The production of homodimers of the other Fc domain, which comprises the “protrusion”, is reduced by the presence of the “protrusion”.

In some embodiments, the Fc domain of a protein of the present invention may or may not further comprises YTE mutations (M252Y/S254T/T256E, EU numbering (Dall'Acqua, Kiener et al. 2006)). These mutations have been shown to improve the pharmacokinetic properties of Fc domains through preferential enhancement of binding affinity for neonatal FcRn receptor at pH 6.0.

In some embodiments, the first and/or the second Fc domain of the present invention derived from an IgG1 also includes the “KO” mutations (L234A, L235A). In a further aspect, the first and/or the second Fc domain of the present invention derived from an IgG4 also includes the Pro hinge mutation (S228P).

In preferred embodiments of the invention, the binding protein comprises i) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:73 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#1/CD3#1), or ii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:74 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#2/CD3#1), or iii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:75 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#3/CD3#1), or iv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:76 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#4/CD3#1), or v) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:77 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79

(DLL3#5/CD3#1), or vi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:78 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#6/CD3#1); or vii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:241 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#7/CD3#1); or viii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:242 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#8/CD3#1); or ix) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:243 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#9/CD3#1); or x) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:244 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#10/CD3#1); or xi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:245 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#11/CD3#1); or xii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:246 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#12/CD3#1); or xiii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:247 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#13/CD3#1); or xiv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:248 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#14/CD3#1); or xv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:249 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#15/CD3#1), or xvi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:250 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#16/CD3#1); or xvii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:251 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#17/CD3#1); or xviii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:252 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:79 (DLL3#18/CD3#1). Preferably, the first and second polypeptide chain are linked via one or more disulfide bonds and form an antibody like structure (Figure 1) similar to a conventional Y-shaped antibody molecule.

In another preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:73, SEQ ID NO:74, and SEQ ID NO:75, and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:79.

5 In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:73 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:79.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:74 and a second
10 polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:79.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:75 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:79.

In preferred embodiments of the invention, the binding protein comprises i) a first
15 polypeptide chain comprising the amino acid sequence of SEQ ID NO:73 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#1/CD3#2), or ii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:74 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#2/CD3#2), or iii) a first polypeptide chain comprising the amino
20 acid sequence of SEQ ID NO:75 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#3/CD3#2), or iv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:76 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#4/CD3#2), or v) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:77 and a second
25 polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#5/CD3#2), or vi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:78 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#6/CD3#2); or vii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:241 and a second polypeptide chain comprising the amino
30 acid sequence of SEQ ID NO:80 (DLL3#7/CD3#2); or viii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:242 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#8/CD3#2); or ix) a first

polypeptide chain comprising the amino acid sequence of SEQ ID NO:243 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#9/CD3#2); or x) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:244 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#10/CD3#2); or xi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:245 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#11/CD3#2); or xii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:246 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#12/CD3#2); or xiii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:247 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#13/CD3#2); or xiv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:248 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#14/CD3#2); or xv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:249 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#15/CD3#2), or xvi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:250 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#16/CD3#2); or xvii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:251 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#17/CD3#2); or xviii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:252 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:80 (DLL3#18/CD3#2). Preferably, the first and second polypeptide chain are linked via one or more disulfide bonds and form an antibody like structure (Figure 1) similar to a conventional Y-shaped antibody molecule.

In another preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:73, SEQ ID NO:74, and SEQ ID NO:75, and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:80.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:73 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:80.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:74 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:80.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:75 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:80.

In preferred embodiments of the invention, the binding protein comprises i) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:73 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#1/CD3#3), or ii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:74 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#2/CD3#3), or iii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:75 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#3/CD3#3), or iv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:76 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#4/CD3#3), or v) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:77 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#5/CD3#3), or vi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:78 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#6/CD3#3); or vii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:241 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#7/CD3#3); or viii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:242 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#8/CD3#3); or ix) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:243 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#9/CD3#3); or x) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:244 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#10/CD3#3); or xi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:245 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#11/CD3#3); or xii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:246 and a second polypeptide chain

comprising the amino acid sequence of SEQ ID NO:105 (DLL3#12/CD3#3); or xiii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:247 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#13/CD3#3); or xiv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:248 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#14/CD3#3); or xv) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:249 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#15/CD3#3), or xvi) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:250 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#16/CD3#3); or xvii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:251 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#17/CD3#3); or xviii) a first polypeptide chain comprising the amino acid sequence of SEQ ID NO:252 and a second polypeptide chain comprising the amino acid sequence of SEQ ID NO:105 (DLL3#18/CD3#3). Preferably, the first and second polypeptide chain are linked via one or more disulfide bonds and form an antibody like structure (Figure 1) similar to a conventional Y-shaped antibody molecule.

In another preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:73, SEQ ID NO:74, and SEQ ID NO:75, and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:105.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:73 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:105.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:74 and a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:105.

In one preferred embodiment, the binding protein comprises a first polypeptide chain specific for DLL3 comprising an amino acid sequence of SEQ ID NO:75 and a second

polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:105.

For all of the above embodiments it shall be understood that, by using the term "comprising", it is intended to also include an embodiment in which the respective domain
5 or molecule "consists of" the amino acid sequence as indicated.

In a further aspect, the present invention provides a protein comprising a first polypeptide chain specifically binding to DLL3 and a second polypeptide chain specifically binding to CD3, wherein the first chain comprises a first light chain covalently linked (e.g. directly bound) to a first linker, which is itself covalently linked (e.g. directly bound) to a first
10 heavy chain, and wherein the second chain specifically binding to CD3 comprises a second light chain covalently linked (e.g. directly bound) to a second linker, which is itself covalently linked (e.g. directly bound) to a second heavy chain.

In some embodiments, starting from its N-terminus, the first polypeptide chain comprises a first light chain variable region specifically binding to DLL3, a first light chain constant
15 region, a first linker, a first heavy chain variable region specific for DLL3 and a first heavy chain constant region. In some embodiments, starting from its N-terminus, the second polypeptide chain comprises a second light chain variable region specifically binding to CD3, a second light chain constant region, a second linker, a second heavy chain variable region specific for CD3 and a second heavy chain constant region.

20 The resulting proteins bear a full Fc, which is marginally larger than an IgG and has two independent binding sites (e.g., each binding site being monovalent for the respective antigen), a first binding site for DLL3 and a second binding site for CD3. Preferably, the first and second polypeptide chain are linked via one or more disulfide bonds. As such, the proteins of the invention are antibody-like structures, having the Y shaped structure of a
25 conventional full length antibody (see Figure 1). This bispecific format greatly reduces heterogeneity after expression and purification (e.g. by avoiding mispairing of light and heavy variable domains with different binding specificities), while maintaining the functional properties of the binding moieties within a structure less likely to generate unwanted immunogenic reactions. This also enables good expression of heterodimeric
30 proteins, e.g. in mammalian cells.

In some embodiments of the protein of the invention, the first polypeptide chain specifically binding to DLL3 comprises a first light chain variable domain and a first heavy

chain variable domain, which comprise CDR sequences selected from the group consisting of i) to xviii):

- i) light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and a heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3);
- ii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3);
- iii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3);
- iv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3);
- v) light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1), SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3); and
- vi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:31 (CDR1), SEQ ID NO:32 (CDR2) and SEQ ID NO:33 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1), SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3);
- vii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:133 (CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:136 (CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3);

- viii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:139 (CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:142 (CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3);
- 5 ix) light chain CDRs comprising the amino acid sequences of SEQ ID NO:145 (CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:148 (CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3);
- 10 x) light chain CDRs comprising the amino acid sequences of SEQ ID NO:151 (CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:154 (CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3);
- 15 xi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:157 (CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:160 (CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3);
- 20 xii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:163 (CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:166 (CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3);
- 25 xiii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3);
- 30 xiv) comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3);
- 30 xv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain

CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3);

xvi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3);

xvii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194 (CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3); and

xviii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3).

The respective light/heavy chain variable domains defined by these CDR sequences are termed DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17, and DLL3#18, respectively. Preferably, the CDR sequences are selected from the group consisting of i) to iii) (DLL3#1, DLL3#2, DLL3#3) as defined above.

In preferred embodiments of the binding protein of the invention, said second polypeptide chain specifically binding to CD3 comprises a second light chain variable domain and second heavy chain variable domain, which comprises CDR sequences selected from the group consisting of:

i) light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3);

ii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3); and

iii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:96 (CDR1), SEQ ID NO:97 (CDR2) and SEQ ID NO:98 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:99 (CDR1), SEQ ID NO:100 (CDR2) and SEQ ID NO:101 (CDR3).

5 The respective light/heavy chain variable domains defined by these CDR sequences are termed CD3#1, CD3#2 and CD3#3, respectively.

In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a first light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:1
10 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and a first heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3); and (ii) a second polypeptide chain specifically binding to CD3, comprising a second light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:55
15 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and a second heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a first light chain variable
20 domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and a first heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3); and (ii) a second polypeptide chain specifically binding to CD3, comprising a second light chain variable
25 domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and a second heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises (i) a first
30 polypeptide chain specifically binding to DLL3, comprising a first light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and a first heavy chain

variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3); and (ii) a second polypeptide chain specifically binding to CD3, comprising a second light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and a second heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a first light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and a first heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3); and (ii) a second polypeptide chain specifically binding to CD3, comprising a second light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and a second heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a first light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and a first heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3); and (ii) a second polypeptide chain specifically binding to CD3, comprising a second light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and a second heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3).

In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a first light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:13

(CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and a first heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3); and (ii) a second polypeptide chain specifically binding to CD3, comprising a second light chain variable domain with light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and a second heavy chain variable domain with heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3).

In preferred embodiments of the protein of the invention, said first polypeptide chain specifically binding to DLL3 comprises a light chain variable domain (a first light chain variable domain) and a heavy chain variable domain (a first heavy chain variable domain) selected from the group consisting of i) to xviii):

- i) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:37 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:38 (DLL3#1);
- ii) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:39 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:40 (DLL3#2);
- iii) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:42 (DLL3#3);
- iv) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:43 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:44 (DLL3#4);
- v) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:45 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:46 (DLL3#5);
- vi) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:47 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:48 (DLL3#6);

- vii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:205 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:206 (DLL3#7);
- 5 viii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:207 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:208 (DLL3#8);
- ix) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:209 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:210 (DLL3#9);
- 10 x) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:211 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:212 (DLL3#10);
- xi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:213 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:214 (DLL3#11);
- 15 xii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:215 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:216 (DLL3#12);
- xiii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:217 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:218 (DLL3#13);
- 20 xiv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:219 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:220 (DLL3#14);
- 25 xv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:221 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:222 (DLL3#15);
- xvi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:223 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:224 (DLL3#16);
- 30

xvii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:225 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:226 (DLL3#17); and

xviii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:227 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:228 (DLL3#18).

Preferably, the light chain variable and heavy chain variable domain sequences are selected from the group consisting of i) to iii) (DLL3#1, DLL3#2, DLL3#3) as defined above.

In preferred embodiments of the protein of the invention, said second polypeptide chain specifically binding to CD3 comprises a light chain variable domain (a second light chain variable domain) and a heavy chain variable domain (a second heavy chain variable domain) selected from the group consisting of:

- i) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:68 (CD3#1);
- ii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70 (CD3#2); and
- iii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:102 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:103 (CD3#3).

In some embodiments, the binding protein of the invention comprises a first and a second polypeptide chain comprising CDR and/or VH and VL sequences of the light/heavy chain variable domains selected from the list consisting of DLL3#1/CD3#1, DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#4/CD3#1, DLL3#5/CD3#1, DLL3#6/CD3#1, DLL3#7/CD3#1, DLL3#8/CD3#1, DLL3#9/CD3#1, DLL3#10/CD3#1, DLL3#11/CD3#1, DLL3#12/CD3#1, DLL3#13/CD3#1, DLL3#14/CD3#1, DLL3#15/CD3#1, DLL3#16/CD3#1, DLL3#17/CD3#1, DLL3#18/CD3#1, DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#4/CD3#2, DLL3#5/CD3#2, DLL3#6/CD3#2, DLL3#7/CD3#2, DLL3#8/CD3#2, DLL3#9/CD3#2, DLL3#10/CD3#2, DLL3#11/CD3#2, DLL3#12/CD3#2, DLL3#13/CD3#2, DLL3#14/CD3#2, DLL3#15/CD3#2, DLL3#16/CD3#2, DLL3#17/CD3#2, DLL3#18/CD3#2, DLL3#1/CD3#3,

- DLL3#2/CD3#3, DLL3#3/CD3#3, DLL3#4/CD3#3, DLL3#5/CD3#3 and DLL3#6/CD3#3, DLL3#7/CD3#3, DLL3#8/CD3#3, DLL3#9/CD3#3, DLL3#10/CD3#3, DLL3#11/CD3#3, DLL3#12/CD3#3, DLL3#13/CD3#3, DLL3#14/CD3#3, DLL3#15/CD3#3, DLL3#16/CD3#3, DLL3#17/CD3#3, DLL3#18/CD3#3. In preferred
- 5 embodiments, the binding protein of the invention comprises a first and a second polypeptide chain comprising CDR and/or VH and VL sequences of the light/heavy chain variable domains selected from the list consisting of DLL3#1/CD3#1, DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#3, and DLL3#3/CD3#3.
- 10 In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a light chain variable domain of SEQ ID NO:37 and heavy chain variable domain of SEQ ID NO:38; and (ii) a second polypeptide chain specifically binding to CD3, comprising a light chain variable domain of SEQ ID NO:67 and a heavy chain variable domain of SEQ ID NO:68.
- 15 In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a light chain variable domain of SEQ ID NO:39 and heavy chain variable domain of SEQ ID NO:40; and (ii) a second polypeptide chain specifically binding to CD3, comprising a light chain variable domain of SEQ ID NO:67 and a heavy chain variable domain of SEQ ID NO:68.
- 20 In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a light chain variable domain of SEQ ID NO:41 and heavy chain variable domain of SEQ ID NO:42; and (ii) a second polypeptide chain specifically binding to CD3, comprising a light chain variable domain of SEQ ID NO:67 and a heavy chain variable domain of SEQ ID NO:68.
- 25 In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a light chain variable domain of SEQ ID NO:37 and heavy chain variable domain of SEQ ID NO:38; and (ii) a second polypeptide chain specifically binding to CD3, comprising a light chain variable domain of SEQ ID NO:69 and a heavy chain variable domain of SEQ ID NO:70.
- 30 In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a light chain variable domain of SEQ ID NO:39 and heavy chain variable domain of SEQ ID NO:40; and (ii) a second

polypeptide chain specifically binding to CD3, comprising a light chain variable domain of SEQ ID NO:69 and a heavy chain variable domain of SEQ ID NO:70.

In one preferred embodiment, the binding protein of the invention comprises (i) a first polypeptide chain specifically binding to DLL3, comprising a light chain variable domain of SEQ ID NO:41 and heavy chain variable domain of SEQ ID NO:42; and (ii) a second polypeptide chain specifically binding to CD3, comprising a light chain variable domain of SEQ ID NO:69 and a heavy chain variable domain of SEQ ID NO:70.

In some embodiments of the invention, the first and/or the second linker peptide comprise a linker as described above, e.g. a linker derived from a hinge region, a poly-alanine linker or a Gly/Ser linker, wherein the linker comprises 26 to 42 amino acids, for example any one of 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, preferably 38 amino acids.

In preferred embodiments, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of any one of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, or SEQ ID NO:240 and the second polypeptide chain specific for CD3 comprises the amino acid sequence of SEQ ID NO:71.

In some embodiments, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of any one of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, or SEQ ID NO:240 and the second polypeptide chain specific for CD3 comprises the amino acid sequence of SEQ ID NO:72.

In some embodiments, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of any one of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, or SEQ ID NO:240 and the

second polypeptide chain specific for CD3 comprises the amino acid sequence of SEQ ID NO:104.

In one preferred embodiment, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of SEQ ID NO:49 and the second polypeptide chain specific for CD3
5 comprises the amino acid sequence of SEQ ID NO:71.

In one preferred embodiment, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of SEQ ID NO:50 and the second polypeptide chain specific for CD3 comprises the amino acid sequence of SEQ ID NO:71.

In one preferred embodiment, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of SEQ ID NO:51 and the second polypeptide chain specific for CD3
10 comprises the amino acid sequence of SEQ ID NO:71.

In one preferred embodiment, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of SEQ ID NO:49 and the second polypeptide chain specific for CD3 comprises the amino acid sequence of SEQ ID NO:72.

15 In one preferred embodiment, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of SEQ ID NO:50 and the second polypeptide chain specific for CD3 comprises the amino acid sequence of SEQ ID NO:72.

In one preferred embodiment, the first polypeptide chain specific for DLL3 comprises the amino acid sequence of SEQ ID NO:51 and the second polypeptide chain specific for CD3
20 comprises the amino acid sequence of SEQ ID NO:72.

In some embodiments of binding protein of the invention, the first and second polypeptide chain comprises an Fc domain derived from the heavy chain of an IgG, for example an IgG1, IgG2 or IgG4. For example, an Fc domain of the present invention is a Fc domain of a heavy chain of an IgG1 or IgG4 and comprises a hinge region and two constant domains
25 (CH2 and CH3). Examples of Fc domains of human IgGs are shown in SEQ ID NO:81 and 84.

In some embodiments of the binding protein of the invention, the heavy chain comprises one or more amino acid changes. For example, such amino acid changes are a tyrosine (Y) at position 366 [T366Y] of the first heavy chain and a threonine (T) at position 407
30 [Y407T] of the second heavy chain. In some embodiments, the first heavy chain comprises a serine (S) at position 366 [T366S] and the second heavy chain comprises a tryptophan

(W) at position 366 [T366W], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V]. In preferred embodiments, the first heavy chain comprises a tryptophan (W) at position 366 [T366W] and the second heavy chain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V]. For example, position 366 of the Fc domain according to EU numbering, corresponding to the amino acid position 146 in the human IgG1 Fc sequence of SEQ ID NO:81, is changed from T at position 146 in SEQ ID NO:81 to W at position 146 in SEQ ID NO:82; and positions 366, 368 and 407 according to EU numbering, corresponding to the amino acid positions 146, 148 and 187, respectively, in SEQ ID NO:81, are changed from T, L and Y at these positions in SEQ ID NO:81 to S, A and V at these positions in SEQ ID NO:83. In any of these embodiments, the amino acid changes described for the first heavy chain may be located in the second heavy chain and the respective amino acid changes for the second heavy chain may be located in the first heavy chain. In other words, the term “first” and “second” can be exchanged in these embodiments. In some embodiments, the heavy chain is derived from the heavy chain of an IgG₁ or IgG₄.

In some embodiments, the first heavy chain or the second heavy chain in a protein of the present invention further comprises one or more amino acid changes which reduce the binding of the heavy chain to protein A. In some embodiments, such amino acid changes are an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F] of one of the heavy chains.

In some embodiments, in a protein of the present invention, the heavy chain, which comprises a threonine (T) at position 407 [Y407T], further comprises an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F]. In this case, the other heavy chain comprises a tyrosine (Y) at position 366 [T366Y], but does not include the two changes at positions 435 and 436. Alternatively, in some embodiments, in a protein of the present invention, the heavy chain, which comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V], further comprises an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F]. In this case, the other heavy chain comprises a tryptophan (W) at position 366 [T366W], but does not include the two changes at positions 435 and 436. Thus, the heavy chain comprising the amino acid change resulting in a “cavity” as described above also comprises the amino acid changes, which reduce binding to Protein A. Homodimers

comprising these heavy chains are removed through reduced binding to Protein A. The production of homodimers of the other heavy chain, which comprises the “protrusion”, is reduced by the presence of the “protrusion”.

In some embodiments, the heavy chain of a protein of the present invention may or may not further comprise YTE mutations (M252Y/S254T/T256E, EU numbering (Dall'Acqua, Kiener et al. 2006)). These mutations have been shown to improve the pharmacokinetic properties of heavy chain through preferential enhancement of binding affinity for neonatal FcRn receptor at pH 6.0.

In some embodiments, the first and/or the second heavy chain of the present invention derived from an IgG1 also includes the “KO” mutations (L234A, L235A). In a further aspect, the first and/or the second heavy chain of the present invention derived from an IgG4 also includes the Pro hinge mutation (S228P).

In a further aspect, the proteins of the invention comprise a first antigen binding unit or polypeptide chain specific for DLL3 with an affinity of preferably ≤ 10 nM, more preferably ≤ 1 nM, even more preferable ≤ 0.1 nM, even more preferably ≤ 0.01 nM to human and cynomolgus monkey DLL3. The affinity can be measured in a SPR (BIAcore) assay using recombinant DLL3-protein, as described, e.g. in the examples or other methods that are well known for the skilled person. The proteins comprise a second antigen binding unit or polypeptide chain with an affinity of preferably ≤ 500 nM, more preferably ≤ 100 nM, even more preferably ≤ 10 nM to human and cynomolgus monkey CD3 ϵ complex.

In a further aspect, the DLL3/CD3 binding proteins of the invention do not bind to DLL3-negative cells and do not cross-react with the human and DLL3 paralogues DLL1 and DLL4, as shown in Example 5.

In a further aspect, the DLL3/CD3 binding proteins of the present invention comprise a first antigen binding unit or a first polypeptide chain specifically binding to the membrane proximal peptide of the DLL3 protein. In a further aspect, the proteins of the invention display weak binding to DLL3 protein expressing cells, e.g. no saturation of protein binding to the cell surface is achieved up to a concentration of 100 nM (see e.g. Figure 6).

In a further aspect, the DLL3/CD3 binding proteins of the present invention are capable of mediating cytotoxicity against tumor cells by providing optimal steric conditions for the formation of a cytolytic synapse between a tumor cell expressing DLL3 and a T cell, in

order to redirect the T cell activity selectively to the targeted tumor cells, leading to tumor cell lysis.

Various methods can be used to measure the cytotoxicity mediated by the DLL3/CD3 binding proteins of the present invention. For example, cytotoxicity can be measured using the method described in example 10. Effector cells can be e.g. stimulated or unstimulated (human or cynomolgus monkey) T cells or their subsets (e.g. CD4, CD8) or unstimulated (human or cynomolgus monkey) peripheral blood mononuclear cells (PBMCs). The target cells should express at least the extracellular domain of (human or cynomolgus monkey) DLL3 and can be cells with endogenous (natural) DLL3 expression, such as human small cell lung carcinoma cell lines SHP77, NCI-H82, alternatively also recombinant cells that express either the full-length DLL3 or the extracellular domain of DLL3. The effector to target cell ratio (E:T) is usually about 10:1 but can vary. Cytotoxic activity of DLL3/CD3 binding molecules can be determined e.g. in a LDH-release assay after 48 or 72 hours of incubation. Modifications in incubation time and read-out used for determination of cytotoxicity are possible and known to the skilled person. Read-out systems for cytotoxicity can comprise MTT/MTS assays, ATP-based assays, FACS-based assays, 51-Chromium release assays, sulforhodamine B (SRB) assays, colorimetric (WST) assays, clonogenic assays, ECIS technology and bioluminescent assays.

The cytotoxic activity mediated by DLL3/CD3 binding proteins of the present invention is preferably measured in a cell-based cytotoxicity assay. The cytotoxicity is represented by the EC_{90} values measured in the cytotoxicity assay. The skilled person is aware that an EC_{90} can be expected to be lower when purified T cells are used as effector cells, compared with PBMCs, the skilled person is also aware that the EC_{90} can be even lower when stimulated T cells are used. It can furthermore expected that the EC_{90} values are lower when the target cells express a high number of DLL3 on the cell surface compared to cell expressing a low number of DLL3 molecules on the cell surface. The EC_{90} of the DLL3/CD3 binding protein is preferably ≤ 10 nM, more preferably ≤ 5 nM and even more preferably ≤ 1 nM.

Preferably, the multi-specific binding proteins of the invention do not induce/mediate lysis of DLL3 negative cells. The term “do not induce/mediate lysis” of DLL3-negative cells means that an DLL3/CD3 binding molecule does not induce or mediate lysis of more than 30%, preferably not more than 20%, more preferably not more than 10% and particular not

more than 5% or DLL3-negative cells, whereas lysis of the DLL3-positive lung carcinoma cell line is set to be 100%. This usually applies for concentrations of the binding protein of up to 1000 nM.

Preferably, the DLL3/CD3 binding proteins of the invention are not internalized by the targeted cells. The rate of internalization can be assayed e.g. as described in Example 11. Preferably, the internalization rate (e.g. measured as a decrease in cytotoxicity) is $\leq 50\%$ after a 4 hour pre-incubation of the DLL3/CD3 binding proteins with the target cells, more preferably $\leq 40\%$ and even more preferably $\leq 30\%$.

Furthermore, the DLL3/CD3 binding proteins of the invention are shown to be stable with a monomer content above 95% (e.g., at least 98%, see example 14), have favorable pharmacokinetic properties and good downstream manufacturability and are further expected to have good bio-distribution (see e.g., example 12). The proteins of the present invention furthermore have a favorable immunogenicity profile (see example 16) and have good stability in-vitro and in-vivo (see e.g., examples 12 and 15). Furthermore, the DLL3/CD3 binding proteins of the invention (e.g., DLL3#3/CD3#1, DLL3#3/CD3#2) show favorable efficacy in a humanized *in vivo* xenograft mouse model. DLL3/CD3 binding proteins induced strong tumor regression starting already after the first dose of DLL3/CD3 binding proteins. Furthermore the DLL3/CD3 binding proteins of the invention induce tumor regression at very low doses of 0.25 mg/kg administered once weekly (q7d), further supporting their therapeutic applicability. In particular, the DLL3/CD3 binding proteins of the invention induce selective T cell proliferation, T cell activation, T cell degranulation and cytokine secretion (see example 18) only in the presence of DLL3-positive target cells and not in the presence of DLL3-negative target cells, and further significantly increase T cell infiltration into tumor tissue (see example 20). Furthermore example 19 demonstrates that DLL3/CD3 binding proteins mediate CD4⁺ as well as CD8⁺ T cell redirected lysis. In particular, naïve T cells as well as CD4⁺ effector memory, CD4⁺ central memory, CD8⁺CD45RA⁺ effector and CD8⁺ memory cells contribute to the T cell redirected lysis of DLL3-expression tumor cells.

A further aspect of the present invention provides isolated nucleic acid molecules encoding the first and/or the second antigen binding unit of a multi-specific binding protein of the invention. In some embodiments, the nucleic acid molecules further encode a first and/or a second Fc domain as described herein, the first and/or second Fc domain linked to the 3'

end of the nucleic acid molecule encoding the first and/or second antigen binding unit, respectively. In some embodiments, the nucleic acid molecule encodes i) a first polypeptide chain comprising a first single chain Fab specific for DLL3 (e.g. any one of DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, 5 DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 and DLL3#18), and optionally a first Fc domain and/or ii) a second polypeptide chain comprising a second single chain Fab specific for CD3 (e.g., any one of CD3#1, CD3#2 and CD3#3) and optionally a second Fc domain.

Preferably the nucleic acid molecule comprises a nucleotide sequence encoding a first 10 single chain Fab of any one of SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:229, SEQ ID NO:230, SEQ ID NO:231, SEQ ID NO:232, SEQ ID NO:233, SEQ ID NO:234, SEQ ID NO:235, SEQ ID NO:236, SEQ ID NO:237, SEQ ID NO:238, SEQ ID NO:239, or SEQ ID NO:240 and/or a second single chain Fab of SEQ ID NO:71, SEQ ID NO:72 or SEQ ID NO:104. In some 15 embodiments, the nucleic acid molecule comprises a nucleotide sequence encoding a first polypeptide chain of any one of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO: 242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO: 245, SEQ ID NO: 246, SEQ ID NO: 247, SEQ ID NO: 248, SEQ ID NO: 249, SEQ ID NO: 250, SEQ ID NO: 251, or SEQ ID NO: 252 20 and/or a second polypeptide chain specific for CD3 comprising the amino acid sequence of SEQ ID NO:79, SEQ ID NO:80 or SEQ ID NO:105.

A further aspect of the invention provides an expression vector containing a DNA molecule comprising the nucleotide sequence encoding the first and/or second antigen binding domain (e.g. a first and/or second single chain Fab of the invention). Preferably 25 the expression vector comprises, in addition, a nucleic acid molecule, preferably a DNA molecule, encoding a first and/or second Fc domain, linked to the nucleic acid molecule, preferably the DNA molecule, encoding the first and/or second antigen binding domain (e.g. first and/or second single chain Fab, respectively). As such, the expression vector comprises a nucleotide sequence encoding a polypeptide chain comprising a first single 30 chain Fab linked to a first Fc domain and/or a nucleotide sequence encoding a polypeptide chain comprising a second single chain Fab linked to a second Fc domain.

In a preferred embodiment, the expression vector contains a DNA molecule comprising the nucleotide sequence encoding the first and/or the second polypeptide chain of the invention. In a preferred embodiment, the expression vector comprises the nucleotide sequence encoding a first polypeptide chain of in any one of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO: 242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO: 245, SEQ ID NO: 246, SEQ ID NO: 247, SEQ ID NO: 248, SEQ ID NO: 249, SEQ ID NO: 250, SEQ ID NO: 251, or SEQ ID NO: 252 and/or a second polypeptide chain comprising SEQ ID NO:79.

In further preferred embodiments, the expression vector comprises the nucleotide sequence encoding a first polypeptide chain of any one of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO: 242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO: 245, SEQ ID NO: 246, SEQ ID NO: 247, SEQ ID NO: 248, SEQ ID NO: 249, SEQ ID NO: 250, SEQ ID NO: 251, or SEQ ID NO: 252 and/or a second polypeptide chain comprising SEQ ID NO:80.

In further preferred embodiments, the expression vector comprises the nucleotide sequence encoding a first polypeptide chain of any one of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO: 242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO: 245, SEQ ID NO: 246, SEQ ID NO: 247, SEQ ID NO: 248, SEQ ID NO: 249, SEQ ID NO: 250, SEQ ID NO: 251, or SEQ ID NO: 252 and/or a second polypeptide chain comprising SEQ ID NO:105.

In a specifically preferred embodiment, two expression vectors may be used, one of them for expression of the first polypeptide chain specific for DLL3, the other one for expression of the second polypeptide chain specific for CD3, which two expression vectors may then both be transfected into a host cell for recombinant protein expression.

Preferably, the expression vector will be a vector comprising said nucleic acid molecule or molecules, operably linked to at least one regulatory sequence, wherein such regulatory sequence may be a promoter, enhancer, or terminator sequence, and most preferably a heterologous promoter, enhancer, or terminator sequence.

In another aspect, the invention relates to a host cell having an expression vector encoding a first polypeptide chain specific for DLL3 of the invention and an expression vector encoding a second polypeptide chain specific for CD3 of the invention.

According to a particularly preferred embodiment, said host cells are eukaryotic cells such as mammalian cells. In another embodiment, such host cells are bacterial cells. Other useful cells are yeast cells or other fungal cells.

Suitable mammalian cells include for example CHO cells, BHK cells, HeLa cells, COS
5 cells, and the like. However, amphibian cells, insect cells, plant cells, and any other cells used in the art for the expression of heterologous proteins can be used as well.

ANTI-DLL3 ANTIBODIES

DLL3 is normally expressed on intracellular membranes, including those of the Golgi
10 apparatus in foetal brain, and plays a key role in somitogenesis in the paraxial mesoderm (Gefferis et al., J Cell Biol. 2007;178:465; Chapman et al., Hum. Mol. Genet. 2011;20:905–916). Marked induction of DLL3 expression in some tumor types including SCLC, LCNEC and glioblastoma results in localization to the cell surface: this together with the absence of detectable cell surface DLL3 in non-malignant cells opens a new window of
15 opportunity for tumour-cell specific therapy and for the use of anti-DLL3 antibodies as diagnostic and prognostic tools.

Several anti-DLL3 antibodies (such as LS-c167440, Lifespan; AP21739PU-N, Acris; LS-C148700, LSBio) are commercially available. However, these antibodies do not perform well in immunohistochemistry (IHC), FACS and ELISA assays and fail to specifically
20 bind to DLL3 protein expressed on cells, particularly tumor cells. Anti-DLL3 antibodies have also been reported, e.g., in WO2007111733 and suggested for diagnostic use in glioma patients, but no data of IHC assays are shown. Thus, the use of anti-DLL3 antibodies as reliable diagnostic tools to accurately measure DLL3 expression in patients, e.g. in tumor cells/tissues and/or to assess the efficacy of DLL3 targeted therapies remains
25 challenging.

Thus, there is a need to identify alternative anti-DLL3 antibodies, which can be used for accurate and specific detection of DLL3 protein expression in various assays such as FACS, ELISA, immunoprecipitation, Western blotting, ELISA, radioimmunoassay, flow cytometry, IHC and immunometric assays in any kind of biological sample and can be
30 used as reliable diagnostic reagents.

Therefore, a further aspect of the invention provides anti-DLL3 antibody molecules comprising

- 5 i) light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3); or
- 10 ii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3); or
- iii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3); or
- 15 iv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3); or
- 20 v) light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1), SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3); or
- 25 vi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:31 (CDR1), SEQ ID NO:32 (CDR2) and SEQ ID NO:33 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1), SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3); or
- 30 vii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:133 (CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:136 (CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3); or

- viii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:139 (CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:142 (CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3); or
- 5 ix) light chain CDRs comprising the amino acid sequences of SEQ ID NO:145 (CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:148 (CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3); or
- 10 x) light chain CDRs comprising the amino acid sequences of SEQ ID NO:151 (CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:154 (CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3); or
- 15 xi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:157 (CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:160 (CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3); or
- 20 xii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:163 (CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:166 (CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3); or
- 25 xiii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3); or
- 30 xiv) comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3); or
- 30 xv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3); or

- xvi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3); or
- 5 xvii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194(CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3); or
- 10 xviii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3).

The antibodies i) to xviii) as outlined above are termed DLL3#1, DLL3#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, 15 DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17 or DLL3#18 respectively. Provided herein is a sequence table which readily allows identification of individual amino acid sequences to specific antibodies of the present invention.

In preferred embodiments of the invention, the antibody molecule comprises CDR sequences as defined in i) or v) above corresponding to DLL3#1 or DLL3#5.

20 In some embodiments, the anti-DLL3 antibody of the invention is a chimeric, a humanized or a human antibody molecule. In some embodiments, the antibody molecule is a monoclonal antibody Fab, F(ab)2, Fv or scFv. . In some embodiments, the anti-DLL3 antibody molecule of the invention comprises a heavy chain constant region selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant regions. In 25 some embodiments, the light chain constant region of the anti-DLL3 antibody molecule of the invention is kappa or lambda.

In some embodiments, the anti-DLL3 antibody of the invention has a heavy chain variable domain comprising an amino acid sequence at least 85% identical to any one of SEQ ID NOs:38, 40, 42, 44, 46, 48, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, and 228. 30 Preferably, the antibody molecule has a heavy chain variable domain comprising an amino acid sequence of SEQ ID NOs:38, 40, 42, 44, 46, 48, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, or 228.

In some embodiments, the anti-DLL3 antibody molecule has a light chain variable domain comprising an amino acid sequence at least 85% identical to any one of SEQ ID NOs:37, 39, 41, 43, 45, 47, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, and 227.

Preferably, the antibody molecule has a light chain variable domain comprising an amino acid sequence of SEQ ID NOs:37, 39, 41, 43, 45, 47, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, or 227.

Methods of calculating amino acid sequence identities are well known in the art and further discussed herein in the Definitions section of the specification.

In some embodiments, the anti-DLL3 antibody molecule has i) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO:37 (DLL3#1), or ii) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 39 (DLL3#2); or iii) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:42 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO:41 (DLL3#3), or iv) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:44 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO:43 (DLL3#4); or v) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:46 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO:45 (DLL3#5); or vi) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:48 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO:47 (DLL3#6); or vii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:205 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:206 (DLL3#7); or viii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:207 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:208 (DLL3#8); or ix) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:209 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:210 (DLL3#9); or x) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:211 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:212 (DLL3#10); or xi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:213 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:214 (DLL3#11); or xii) a light chain

variable domain comprising the amino acid sequence of SEQ ID NO:215 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:216 (DLL3#12); or
xiii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:217 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:218
5 (DLL3#13); or xiv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:219 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:220 (DLL3#14); or xv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:221 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:222 (DLL3#15); or xvi) a light chain variable domain
10 comprising the amino acid sequence of SEQ ID NO:223 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:224 (DLL3#16); or xvii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:225 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:226 (DLL3#17); or xviii) a light chain variable domain comprising the amino acid sequence of SEQ ID
15 NO:227 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:228 (DLL3#18).

In some embodiments, the anti-DLL3 antibody of the invention is a mouse monoclonal antibody. In the context of this invention a mouse monoclonal antibody includes an antibody where the VH and VL are obtained from immunization of mice with human
20 DLL3 protein, subsequent selection of suitable VH and VL sequences binding with certain affinity to human DLL3, and then further joining such VH and VL sequences to constant domains which are derived from mouse (e.g. from mouse IgG2a) by recombinant techniques; and which are produced by recombinant expression in host cells. Further encompassed by the invention are chimeric antibodies, e.g., comprising variable and
25 constant regions from different species. In some embodiments, the antibody molecule of the invention is a chimeric antibody comprising VH and VL domains derived from mouse as described above and further comprising constant domains derived from another species such as human, rabbit, rat, goat, donkey. In some embodiments, the chimeric antibody comprises VH and VL domains derived from mouse and further humanized or sequence
30 optimized as defined above and further comprises constant domains derived from another species. In some embodiments, the chimeric antibody comprises VH and VL domains derived from a transgenic animal (e.g. a mouse) comprising human IgG sequences, thus comprises human VH and VL sequences, and further comprises constant domains derived

from another species. In any of the embodiments of chimeric antibodies as outlined above, the heavy chain constant region is a mouse, human, rabbit, rat, goat or donkey heavy chain region.

In some embodiments, the anti-DLL3 antibody molecule of the invention has a constant domain selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant domains. In a preferred embodiment, the anti-DLL3 antibody has a constant domain of IgG2a, preferably comprising the sequence of SEQ ID NO:254. In some embodiments, the anti-DLL3 antibody molecule has a light chain constant domain which is kappa or lambda, preferably the light chain constant domain is a kappa light chain constant domain, preferably comprising the sequence of SEQ ID NO:255.

In some embodiments, the anti-DLL3 antibody molecule is capable of binding to human and cynomolgus monkey DLL3 with a dissociation constant (KD) of preferably ≤ 10 nM, more preferably ≤ 1 nM, even more preferable ≤ 0.1 nM, even more preferably ≤ 0.01 nM. The affinity (KD value) can be measured in a SPR (BIAcore) assay using recombinant DLL3-protein, as described, e.g. in the examples or other methods that are well known for the skilled person.

In some embodiments, the anti-DLL3 antibody molecule does not bind to mouse DLL3.

In some embodiments, the anti-DLL3 antibody molecule is capable of detecting DLL3 expression (e.g. cytoplasmic and surface protein expression) on tissue samples (e.g., paraffin embedded/formalin-fixed tissue samples) such as tumor tissue samples, or cultured cell lines. Optionally, the paraffin embedded/formalin fixed cell culture or tissue samples are further treated with an epitope retriever such as Proteinase K.

In some embodiments, the anti-DLL3 antibody molecule of the invention binds to the same epitope as any one of the DLL3#1, DLL2#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17, or DLL3#18 antibody. In some embodiments, the anti-DLL3 antibody molecule of the invention cross-competes for binding to human DLL3 with any one of the DLL3#1, DLL2#2, DLL3#3, DLL3#4, DLL3#5, DLL3#6, DLL3#7, DLL3#8, DLL3#9, DLL3#10, DLL3#11, DLL3#12, DLL3#13, DLL3#14, DLL3#15, DLL3#16, DLL3#17, or DLL3#18 antibody molecules.

Another aspect of the present invention provides isolated nucleic acid molecules encoding the heavy chain variable domain and/or the light chain variable domain of an anti-DLL3 antibody molecule of the invention.

Preferably the nucleic acid molecule comprises a nucleotide sequence encoding the heavy chain variable domain of any one of SEQ ID NOs:38, 40, 42, 44, 46, 48, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, or 228. Preferably the nucleic acid molecule comprises a nucleotide sequence encoding the light chain variable domain of any one of SEQ ID NOs: 37, 39, 41, 43, 45, 47, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, or 227.

A further aspect of the invention provides an expression vector containing a DNA molecule comprising the nucleotide sequence encoding the heavy chain variable domain and/or the light chain variable domain of an anti-DLL3 antibody molecule of the invention.

Preferably the expression vector comprises, in addition, a nucleic acid molecule, preferably a DNA molecule, encoding the constant domains of a heavy chain and/or the constant domain of a light chain, respectively, linked to the nucleic acid molecule, preferably the DNA molecule, encoding the heavy chain variable domain and/or the light chain variable domain, respectively.

In a specifically preferred embodiment, two expression vectors may be used, one of them for expression of the heavy chain, the other one for expression of the light chain, which two expression vectors may then both be transfected into a host cell for recombinant protein expression.

Preferably, the expression vector will be a vector comprising said nucleic acid molecule or molecules, operably linked to at least one regulatory sequence, wherein such regulatory sequence may be a promoter, enhancer, or terminator sequence, and most preferably a heterologous promotor, enhancer, or terminator sequence.

In another aspect, the invention relates to a host cell having an expression vector encoding a heavy chain of an anti-DLL3 antibody molecule of the invention and an expression vector encoding a light chain of an anti-DLL3 antibody molecule of the invention.

According to a particularly preferred embodiment, said host cells are eukaryotic cells such as mammalian cells. In another embodiment, such host cells are bacterial cells. Other useful cells are yeast cells or other fungal cells.

Suitable mammalian cells include for example CHO cells, BHK cells, HeLa cells, COS cells, and the like. However, amphibian cells, insect cells, plant cells, and any other cells used in the art for the expression of heterologous proteins can be used as well.

5 METHODS OF MANUFACTURE AND PURIFICATION

The invention further provides methods of manufacturing a multi-specific binding protein of the invention, such methods generally comprising the steps of:

- culturing host cells comprising an expression vector comprising a nucleic acid encoding a binding protein of the invention under conditions that allow formation of the binding
- 10 protein of the invention; and,
- recovering the binding protein expressed by the host cells from the culture; and
- optionally further purifying and/or modifying and/or formulating the binding protein of the invention.

The invention further provides methods of manufacturing an anti-DLL3 antibody of the

15 invention, such methods generally comprising the steps of:

- culturing host cells comprising an expression vector comprising a nucleic acid encoding an antibody molecule of the invention under conditions that allow formation of the antibody molecule; and,
- recovering the antibody molecule expressed by the host cells from the culture; and
- 20 - optionally further purifying and/or modifying and/or formulating the antibody molecule of the invention.

A nucleic acid of the invention can e.g. be a DNA molecule comprising coding sequences as well as regulatory sequences and optionally natural or artificial introns, or can be a cDNA molecule. It may have its original codons or may have an optimized codon usage

25 that has been specifically adapted for expression in the intended host cell or host organism. According to one embodiment of the invention, the nucleic acid of the invention is in essentially isolated form, as defined above.

The nucleic acids of the invention may be prepared or obtained in a manner known per se (e.g. by automated DNA synthesis and/or recombinant DNA technology), based on the

30 information on the amino acid sequences for the proteins of the invention given herein.

The nucleic acid of the invention will typically be incorporated into an expression vector, i.e. a vector that can provide for expression of the protein when transfected into a suitable host cell or other expression system.

For manufacturing the binding proteins or antibodies of the invention, the skilled artisan
5 may choose from a great variety of expression systems well known in the art, e.g. those reviewed by Kipriyanow and Le Gall, 2004.

Expression vectors include plasmids, retroviruses, cosmids, EBV derived episomes, and the like. The expression vector and expression control sequences are selected to be compatible with the host cell. The nucleotide sequence encoding the first antigen binding
10 unit (e.g. the DLL3 specific single chain Fab or the full length DLL3 chain of the binding protein of the invention) and the nucleotide sequence encoding the second antigen binding unit (e.g. CD3 specific single chain Fab or the full length CD3 chain of the binding protein of the invention) of the DLL3/CD3 binding protein can be inserted into separate vectors. In certain embodiments, both DNA sequences are inserted into the same expression vector.
15 The nucleotide sequence encoding the light chain of a DLL3 antibody and the nucleotide sequence encoding the heavy chain of a DLL3 antibody can be inserted into separate vectors. In certain embodiments, both DNA sequences are inserted into the same expression vector.

Convenient vectors are those that encode a functionally complete human CH (constant
20 heavy) immunoglobulin sequence, with appropriate restriction sites engineered so that any antigen binding unit such as a single chain Fab sequence or any heavy/light chain variable domain can be easily inserted and expressed, as described above. For the antibody heavy chain, it can be, without limitation, any IgG isotype (IgG1, IgG2, IgG3, IgG4) or other immunoglobulins, including allelic variants.

The recombinant expression vector may also encode a signal peptide that facilitates
25 secretion of the full length CD3 or DLL3 chain from a host cell or of the light/heavy chain of an anti-DLL3 antibody. The DNA encoding the protein chain may be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the mature full length chain DNA. The signal peptide may be an immunoglobulin signal peptide or a
30 heterologous peptide from a non-immunoglobulin protein. Alternatively, the DNA sequence encoding the full length chains of the protein of the invention may already contain a signal peptide sequence.

In addition to the DLL3/CD3 chain encoding DNA sequences or the heavy/light chain of a DLL3 antibody encoding DNA sequences, the recombinant expression vectors typically carries regulatory sequences, optionally heterologous regulatory sequences, including promoters, enhancers, termination and polyadenylation signals and other expression control elements that control the expression of the protein chains in a host cell. Examples for promoter sequences (exemplified for expression in mammalian cells) are promoters and/or enhancers derived from CMV (such as the CMV Simian Virus 40 (SV40) promoter/enhancer), adenovirus, (e. g., the adenovirus major late promoter (AdMLP)), polyoma and strong mammalian promoters such as native immunoglobulin and actin promoters. Examples for polyadenylation signals are BGH polyA, SV40 late or early polyA; alternatively, 3'UTRs of immunoglobulin genes etc. can be used.

The recombinant expression vectors may also carry sequences that regulate replication of the vector in host cells (e. g. origins of replication) and selectable marker genes. Nucleic acid molecules encoding the full length chain with the first antigen binding unit (single chain Fab and Fc domain) or an antigen-binding portion thereof and/or the full length chain with the second antigen binding unit (single chain Fab and Fc domain) or an antigen-binding portion thereof, and vectors comprising these DNA molecules can be introduced into host cells, e.g. bacterial cells or higher eukaryotic cells, e.g. mammalian cells, according to transfection methods well known in the art, including liposome-mediated transfection, polycation-mediated transfection, protoplast fusion, microinjections, calcium phosphate precipitation, electroporation or transfer by viral vectors.

Preferably, the DNA molecules encoding the DLL3 and CD3 chain of the protein of the invention are present on two expression vectors which are co-transfected into the host cell, preferably a mammalian cell.

Mammalian cell lines available as hosts for expression are well known in the art and include, inter alia, Chinese hamster ovary (CHO) cells, NS0, SP2/0 cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human carcinoma cells (e. g., Hep G2 and A-549 cells), 3T3 cells or the derivatives/progenies of any such cell line. Other mammalian cells, including but not limited to human, mice, rat, monkey and rodent cells lines, or other eukaryotic cells, including but not limited to yeast, insect and plant cells, or prokaryotic cells such as bacteria may be used.

The proteins of the invention are produced by culturing the host cells for a period of time sufficient to allow for expression of the protein in the host cells. Protein molecules are preferably recovered from the culture medium as a secreted polypeptide or it can be recovered from host cell lysates if for example expressed without a secretory signal. It is
5 necessary to purify the protein molecules using standard protein purification methods used for recombinant proteins and host cell proteins in a way that substantially homogenous preparations of the protein are obtained. By way of example, state-of-the art purification methods useful for obtaining protein molecules of the invention include, as a first step, removal of cells and/or particulate cell debris from the culture medium or lysate. The
10 protein is then purified from contaminant soluble proteins, polypeptides and nucleic acids, for example, by fractionation on immunoaffinity or ion-exchange columns, ethanol precipitation, reverse phase HPLC, Sephadex chromatography, chromatography on silica or on a cation exchange resin. As a final step in the process for obtaining a protein molecule preparation, the purified protein molecule may be dried, e.g. lyophilized, as
15 described below for therapeutic applications.

The present invention relates to binding proteins that have binding specificities for at least two different targets. In relation to the present invention, the binding molecules are derived from antibodies. Techniques for making binding molecules include, but are not limited to, recombinant co-expression of two immunoglobulin chains having different
20 specificities (see Milstein and Cuello, *Nature* 305: 537 (1983)), WO 93/08829, and Traunecker et al., *EMBO J.* 10: 3655 (1991)), and "knob-in-hole" engineering (see, e.g., U.S. Patent No. 5,731, 168; Atwell et al, *JMB*, 1997, 270, 26-35). Binding proteins of the invention may also be made by engineering electrostatic steering effects for making antibody Fc-heterodimeric molecules (WO 2009/089004A1); cross- linking two or more
25 antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan et al., *Science*, 229: 81 (1985)); using leucine zippers to produce bi-specific proteins (see, e.g., Kostelny et al., *Immunol.*, 148(5): 1547-1553 (1992)); using "diabody" technology for making bispecific antibody fragments (see, e.g., Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444- 6448 (1993)); and using single-chain Fv (sFv) dimers (see, e.g. Gruber et al., /.
30 *Immunol.*, 152:5368 (1994)); and preparing trispecific antibodies as described, e.g., in Tutt et al. /. *Immunol.* 147: 60 (1991).

The compositions (e.g., multi-specific binding proteins and anti-DLL3 antibodies) and methods disclosed herein encompass polypeptides and nucleic acids having the sequences

specified, or sequences substantially identical or similar thereto, e.g., sequences at least 85%, 90%, 95% identical or higher to the sequence specified. In the context of an amino acid sequence, the term "substantially identical" is used herein to refer to a first amino acid sequence that contains a sufficient or minimum number of amino acid residues that are i) identical to, or ii) conservative substitutions of aligned amino acid residues in a second amino acid sequence such that the first and second amino acid sequences can have a common structural domain and/or common functional activity. For example, amino acid sequences that contain a common structural domain have at least about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a reference sequence, e.g., a sequence provided herein. In the context of nucleotide sequence, the term "substantially identical" is used herein to refer to a first nucleic acid sequence that contains a sufficient or minimum number of nucleotides that are identical to aligned nucleotides in a second nucleic acid sequence such that the first and second nucleotide sequences encode a polypeptide having common functional activity, or encode a common structural polypeptide domain or a common functional polypeptide activity, for example, nucleotide sequences having at least about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a reference sequence.

The nucleic acid molecules of the invention include, but are not limited to, the DNA molecules encoding the polypeptide sequences shown in the sequence listing. Also, the present invention also relates to nucleic acid molecules that hybridize to the DNA molecules encoding the polypeptide sequences shown in the sequence listing under high stringency binding and washing conditions, as defined in WO 2007/042309. Preferred molecules (from an mRNA perspective) are those that have at least 75% or 80% (preferably at least 85%, more preferably at least 90% and most preferably at least 95%) homology or sequence identity with one of the DNA molecules described herein. By way of example, in view of expressing the antibodies in eukaryotic cells, the DNA sequences shown in the sequence listing have been designed to match codon usage in eukaryotic cells. If it is desired to express the antibodies in *E. coli*, these sequences can be changed to match *E. coli* codon usage. Variants of DNA molecules of the invention can be constructed in several different ways, as described e.g. in WO 2007/042309.

The proteins of the invention may have a modified N-terminal sequence, e.g. a deletion of one or more of the N-terminal amino acids, or an exchange of e.g. the first, N-terminal amino acid (e.g. glutamate to alanine), to optimize the molecule for being expressed by

using certain expression systems (such as specific vectors or host cells), or for being expressed as inclusion bodies or in soluble form, or for being secreted into the medium or the periplasmic space or for being contained within the cell, or for yielding a more homogenous product. The polypeptides of the invention may have a modified C-terminal
5 sequence, such as an additional alanine, and/or further amino acid exchanges in the C-terminal part or at other defined positions within any of the framework regions, as explained e.g. in WO2012/175741, WO2011/075861, or WO2013/024059, in order to e.g. further enhance stability or reduce immunogenicity of such polypeptides.

For the avoidance of doubt, all of the embodiments relating to pharmaceutical
10 compositions, kits, treatment methods, medical uses, combinations, methods of administration and dosages as described herein are contemplated for any of the multi-specific binding proteins described herein, either alone or in combination with further therapeutic agents (as specified in more detail below).

15 PHARMACEUTICAL COMPOSITIONS; METHODS OF ADMINISTRATION;
DOSAGES

The invention further relates to pharmaceutical compositions for the treatment of a disease (as specified in more detail below), wherein such compositions comprise at least one multi-specific binding protein of the invention. The invention further encompasses methods of
20 treating a disease (as specified in more detail below) using at least one multi-specific protein of the invention or pharmaceutical composition as set out below, and further encompasses the preparation of a medicament for the treatment of such disease by using such binding protein of the invention or pharmaceutical composition.

The binding proteins of the invention (e.g., any one of DLL3#1/CD3#1, DLL3#2/CD3#1,
25 DLL3#3/CD3#1, DLL3#4/CD3#1, DLL3#5/CD3#1, DLL3#6/CD3#1, DLL3#7/CD3#1, DLL3#8/CD3#1, DLL3#9/CD3#1, DLL3#10/CD3#1, DLL3#11/CD3#1, DLL3#12/CD3#1, DLL3#13/CD3#1, DLL3#14/CD3#1, DLL3#15/CD3#1, DLL3#16/CD3#1, DLL3#17/CD3#1, DLL3#18/CD3#1, DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#4/CD3#2, DLL3#5/CD3#2, DLL3#6/CD3#2,
30 DLL3#7/CD3#2, DLL3#8/CD3#2, DLL3#9/CD3#2, DLL3#10/CD3#2, DLL3#11/CD3#2, DLL3#12/CD3#2, DLL3#13/CD3#2, DLL3#14/CD3#2, DLL3#15/CD3#2, DLL3#16/CD3#2, DLL3#17/CD3#2, DLL3#18/CD3#2, DLL3#1/CD3#3,

DLL3#2/CD3#3, DLL3#3/CD3#3, DLL3#4/CD3#3, DLL3#5/CD3#3 and
DLL3#6/CD3#3, DLL3#7/CD3#3, DLL3#8/CD3#3, DLL3#9/CD3#3, DLL3#10/CD3#3,
DLL3#11/CD3#3, DLL3#12/CD3#3, DLL3#13/CD3#3, DLL3#14/CD3#3,
DLL3#15/CD3#3, DLL3#16/CD3#3, DLL3#17/CD3#3, DLL3#18/CD3#3, preferably any
5 one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1,
DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3)
and/or the compositions comprising the same can be administered to a patient in need
thereof in any suitable manner, depending on the specific pharmaceutical formulation or
composition to be used. Thus, the binding proteins of the invention and/or the
10 compositions comprising the same can for example be administered intravenously (i.v.),
subcutaneously (s.c.), intramuscularly (i.m.), intraperitoneally (i.p.), transdermally, orally,
sublingually (e.g. in the form of a sublingual tablet, spray or drop placed under the tongue
and adsorbed through the mucus membranes into the capillary network under the tongue),
(intra-)nasally (e.g. in the form of a nasal spray and/or as an aerosol), topically, by means
15 of a suppository, by inhalation, or any other suitable manner in an effective amount or
dose. The binding protein can be administered by infusion, bolus or injection. In preferred
embodiments, the administration is by intravenous infusion or subcutaneous injection.

The binding proteins of the invention and/or the compositions comprising the same are
administered according to a regimen of treatment that is suitable for treating and/or
20 alleviating the disease, disorder or condition to be treated or alleviated. The clinician will
generally be able to determine a suitable treatment regimen, depending on factors such as
the disease, disorder or condition to be treated or alleviated, the severity of the disease, the
severity of the symptoms thereof, the specific binding protein of the invention to be used,
the specific route of administration and pharmaceutical formulation or composition to be
25 used, the age, gender, weight, diet, general condition of the patient, and similar factors well
known to the clinician. Generally, the treatment regimen will comprise the administration
of one or more binding proteins of the invention, or of one or more compositions
comprising the same, in therapeutically effective amounts or doses.

Generally, for the treatment and/or alleviation of the diseases, disorders and conditions
30 mentioned herein and depending on the specific disease, disorder or condition to be
treated, the potency of the specific binding protein of the invention to be used, the specific
route of administration and the specific pharmaceutical formulation or composition used,
the binding proteins of the invention will generally be administered in an amount between

0.005 and 20.0 mg per kilogram of body weight and dose, preferably between 0.05 and 10.0 mg/kg/dose, either continuously (e.g. by infusion) or more preferably as single doses (such as e.g. twice a week, weekly, or monthly doses; cf. below), but can significantly vary, especially, depending on the before-mentioned parameters. Thus, in some cases it
 5 may be sufficient to use less than the minimum dose given above, whereas in other cases the upper limit may have to be exceeded. When administering large amounts it may be advisable to divide them up into a number of smaller doses spread over the day.

Depending on the specific binding protein of the invention and its specific pharmacokinetic and other properties, it may be administered daily, every second, third, fourth, fifth or sixth
 10 day, weekly, monthly, and the like. An administration regimen could include long-term, weekly treatment. By "long-term" is meant at least two weeks and preferably months, or years of duration.

The efficacy of the multi-specific protein of the invention, and of compositions comprising the same, can be tested using any suitable in vitro assay, cell-based assay, in vivo assay
 15 and/or animal model known per se, or any combination thereof, depending on the specific disease involved. Suitable assays and animal models will be clear to the skilled person, and for example include the assays and animal models used in the Examples below.

FORMULATIONS

20 For pharmaceutical use, the binding proteins of the invention may be formulated as a pharmaceutical preparation comprising (i) at least one binding protein of the invention (e.g., any one of DLL3#1/CD3#1, DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#4/CD3#1, DLL3#5/CD3#1, DLL3#6/CD3#1, DLL3#7/CD3#1, DLL3#8/CD3#1, DLL3#9/CD3#1, DLL3#10/CD3#1, DLL3#11/CD3#1, DLL3#12/CD3#1, DLL3#13/CD3#1,
 25 DLL3#14/CD3#1, DLL3#15/CD3#1, DLL3#16/CD3#1, DLL3#17/CD3#1, DLL3#18/CD3#1, DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#4/CD3#2, DLL3#5/CD3#2, DLL3#6/CD3#2, DLL3#7/CD3#2, DLL3#8/CD3#2, DLL3#9/CD3#2, DLL3#10/CD3#2, DLL3#11/CD3#2, DLL3#12/CD3#2, DLL3#13/CD3#2, DLL3#14/CD3#2, DLL3#15/CD3#2, DLL3#16/CD3#2, DLL3#17/CD3#2,
 30 DLL3#18/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#3, DLL3#3/CD3#3, DLL3#4/CD3#3, DLL3#5/CD3#3 and DLL3#6/CD3#3, DLL3#7/CD3#3, DLL3#8/CD3#3, DLL3#9/CD3#3, DLL3#10/CD3#3, DLL3#11/CD3#3, DLL3#12/CD3#3,

DLL3#13/CD3#3, DLL3#14/CD3#3, DLL3#15/CD3#3, DLL3#16/CD3#3, DLL3#17/CD3#3, DLL3#18/CD3#3, preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) and (ii) at least one pharmaceutically acceptable carrier, diluent, excipient, adjuvant, and/or stabilizer, and (iii) optionally one or more further pharmacologically active polypeptides and/or compounds. By "pharmaceutically acceptable" is meant that the respective material does not show any biological or otherwise undesirable effects when administered to an individual and does not interact in a deleterious manner with any of the other components of the pharmaceutical composition (such as e.g. the pharmaceutically active ingredient) in which it is contained. Specific examples can be found in standard handbooks, such as e.g. Remington's Pharmaceutical Sciences, 18th Ed., Mack Publishing Company, USA (1990). For example, the binding proteins of the invention may be formulated and administered in any manner known per se for conventional antibodies and antibody fragments and other pharmaceutically active proteins. Thus, according to a further embodiment, the invention relates to a pharmaceutical composition or preparation that contains at least one binding protein of the invention and at least one pharmaceutically acceptable carrier, diluent, excipient, adjuvant and/or stabilizer, and optionally one or more further pharmacologically active substances, in the form of lyophilized or otherwise dried formulations or aqueous or non-aqueous solutions or suspensions.

Pharmaceutical preparations for parenteral administration, such as intravenous, intramuscular, subcutaneous injection or intravenous infusion may for example be sterile solutions, suspensions, dispersions, emulsions, or powders which comprise the active ingredient and which are suitable, optionally after a further dissolution or dilution step, for infusion or injection. Suitable carriers or diluents for such preparations for example include, without limitation, sterile water and pharmaceutically acceptable aqueous buffers and solutions such as physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's solution; water oils; glycerol; ethanol; glycols such as propylene glycol, as well as mineral oils, animal oils and vegetable oils, for example peanut oil, soybean oil, as well as suitable mixtures thereof.

Solutions of the binding proteins of the invention may also contain a preservative to prevent the growth of microorganisms, such as antibacterial and antifungal agents, for example, p-hydroxybenzoates, parabens, chlorobutanol, phenol, sorbic acid, thiomersal,

(alkali metal salts of) ethylenediamine tetraacetic acid, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Optionally, emulsifiers and/or dispersants may be used. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. Other agents delaying absorption, for example, aluminum monostearate and gelatin, may also be added. The solutions may be filled into injection vials, ampoules, infusion bottles, and the like.

In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

Usually, aqueous solutions or suspensions will be preferred. Generally, suitable formulations for therapeutic proteins such as the binding proteins of the invention are buffered protein solutions, such as solutions including the protein in a suitable concentration (such as from 0.001 to 400 mg/ml, preferably from 0.005 to 200 mg/ml, more preferably 0.01 to 200 mg/ml, more preferably 1.0 - 100 mg/ml, such as 1.0 mg/ml (i.v. administration) or 100 mg/ml (s.c. administration) and an aqueous buffer such as:

- phosphate buffered saline, pH 7.4,
- other phosphate buffers, pH 6.2 to 8.2,
- acetate buffers, pH 3.2 to 7.5, preferably pH 4.8 to 5.5
- histidine buffers, pH 5.5 to 7.0,
- succinate buffers, pH 3.2 to 6.6, and
- citrate buffers, pH 2.1 to 6.2,

and, optionally, salts (e.g. NaCl) and/or sugars (such as e.g. sucrose and trehalose) and/or other polyalcohols (such as e.g. mannitol and glycerol) for providing isotonicity of the solution.

In addition, other agents such as a detergent, e.g. 0.02 % Tween-20 or Tween-80, may be included in such solutions. Formulations for subcutaneous application may include significantly higher concentrations of the antibody of the invention, such as up to

100 mg/ml or even above 100 mg/ml. However, it will be clear to the person skilled in the art that the ingredients and the amounts thereof as given above do only represent one, preferred option. Alternatives and variations thereof will be immediately apparent to the skilled person, or can easily be conceived starting from the above disclosure. The above
 5 described formulations can optionally be provided as lyophilized formulation that is to be reconstituted in a solution, e.g. in water for injection (WFI).

According to a further aspect of the invention, a binding protein of the invention may be used in combination with a device useful for the administration of protein, such as a syringe, injector pen, micropump, or other device.

10

METHOD OF TREATMENT

A further aspect of the invention provides a method of treating cancer comprising administering to a patient in need thereof a therapeutically effective amount of the binding protein of the invention.

15 A further aspect of the invention provides a binding protein of the invention for use in a method of treating cancer.

A further aspect of the invention is the use of the binding protein of the invention for preparing a pharmaceutical composition for treating cancer.

For the avoidance of doubt, the medical use aspects of the invention may comprise any of
 20 the specific binding proteins of the invention as described above (e.g., any one of DLL3#1/CD3#1, DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#4/CD3#1, DLL3#5/CD3#1, DLL3#6/CD3#1, DLL3#7/CD3#1, DLL3#8/CD3#1, DLL3#9/CD3#1, DLL3#10/CD3#1, DLL3#11/CD3#1, DLL3#12/CD3#1, DLL3#13/CD3#1, DLL3#14/CD3#1, DLL3#15/CD3#1, DLL3#16/CD3#1, DLL3#17/CD3#1, DLL3#18/CD3#1,
 25 DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#4/CD3#2, DLL3#5/CD3#2, DLL3#6/CD3#2, DLL3#7/CD3#2, DLL3#8/CD3#2, DLL3#9/CD3#2, DLL3#10/CD3#2, DLL3#11/CD3#2, DLL3#12/CD3#2, DLL3#13/CD3#2, DLL3#14/CD3#2, DLL3#15/CD3#2, DLL3#16/CD3#2, DLL3#17/CD3#2, DLL3#18/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#3, DLL3#3/CD3#3, DLL3#4/CD3#3, DLL3#5/CD3#3
 30 and DLL3#6/CD3#3, DLL3#7/CD3#3, DLL3#8/CD3#3, DLL3#9/CD3#3, DLL3#10/CD3#3, DLL3#11/CD3#3, DLL3#12/CD3#3, DLL3#13/CD3#3, DLL3#14/CD3#3, DLL3#15/CD3#3, DLL3#16/CD3#3, DLL3#17/CD3#3,

DLL3#18/CD3#3, preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3).

As used herein, the term "cancer" is meant to include all types of cancerous growths or
5 oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness.

Exemplary cancers whose growth can be inhibited using the multi-specific binding proteins described herein are any DLL3 expressing tumors, preferably SCLC, LCNEC, glioma, glioblastoma, melanoma, or other DLL3 expressing neuroendocrine tumors (e.g.,
10 neuroendocrine prostate cancer or neuroendocrine pancreatic cancer, small cell bladder cancer).

Neuroendocrine tumors (NETs) arise from the dispersed endocrine system and can occur in many different areas of the body. Traditionally, NETs have been classified by their anatomic site of origin and are typically highly aggressive. They are most often located in
15 the gastrointestinal tract, pancreas or the lungs (small cell lung carcinoma and large cell neuroendocrine carcinoma) as well as in kidneys, or in the genitourinary tract (bladder, prostate, ovary, cervix, and endometrium). NETs include certain tumors of the gastrointestinal tract and of the pancreatic islet cells, certain thymus and lung tumors, and medullary carcinoma of the parafollicular cells of the thyroid.

20 Additional cancers whose growth can be inhibited using the multi-specific binding proteins described herein are pseudo neuroendocrine tumors (pNETs) that share certain genotypic, phenotypic or biochemical characteristics with traditionally defined neuroendocrine tumors.

In some embodiments, the following cancers, tumors, and other proliferative diseases may
25 be treated with multi-specific binding proteins of the invention: lung cancer; preferably SCLC, NSCLC or LCNEC; breast; cervical; colon; colorectal; endometrial; head and neck; liver (hepatoblastoma or hepatocellular carcinoma); ovarian; pancreatic; prostate; skin; gastric; testis; thyroid; adrenal; renal; bladder; uterine; esophageal; urothelial cancer; brain tumor; lymphoma; Ewing sarcoma; and other neuroendocrine and small blue round cell
30 tumors.

In a preferred embodiment of the invention the cancer is small cell lung cancer (SCLC) or glioblastoma.

All cancers, tumors, neoplasms, etc., mentioned above which are characterized by their specific location/origin in the body are meant to include both the primary tumors and the metastatic tumors derived therefrom.

It is possible that a patient is more likely to respond to treatment with a binding protein of the invention (as described herein) if that patient has a cancer which is characterized by having a high expression of DLL3. Thus, in some embodiments, the cancer to be treated with the binding proteins of the invention is a cancer with high expression of DLL3, e.g., DLL3 expression is higher than the average expression in cancer cells of a population of patients suffering from the same type of a DLL3 expressing cancer.

The binding proteins of the invention may be used in therapeutic regimens in the context of first line, second line, or any further line treatments and maintenance treatment.

The binding proteins of the invention may be used for the prevention, short-term or long-term treatment of the above-mentioned diseases, optionally also in combination with radiotherapy, one or more additional therapeutic agents and/or surgery.

In preferred embodiments, the protein of the invention is used for the treatment of cancer in combination with a PD-1 antagonist, such as an anti-PD-1 antibody or an anti-PDL-1 antibody. Preferably said anti-PD-1 antibody is selected from the group consisting of pembrolizumab, nivolumab, pidilizumab, PD1-1, PD1-2, PD1-3, PD1-4, and PD1-5 as described herein (as defined by the sequences in Table A below) and in WO2017/198741 (incorporated herein by reference). Preferably said anti-PDL-1 antibody is selected from the group consisting of atezolizumab, avelumab and durvalumab. In particular preferred embodiments, the binding protein of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, or DLL3#3/CD3#3) is used for the treatment of cancer in combination with PD1-1. In particular preferred embodiments, the binding protein of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, or DLL3#3/CD3#3) is used for the treatment of cancer in combination with PD1-2. In particular preferred embodiments, the binding protein of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is used for the treatment of cancer in combination with PD1-3. In particular preferred embodiments, the

- binding protein of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is used for the treatment of cancer in combination with PD1-4. In particular preferred embodiments, the binding protein of the invention
- 5 (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is used for the treatment of cancer in combination with PD1-5.

Table A: Amino acid sequences and SEQ ID NOs of heavy chain and light chain sequences of anti-PD1 antibodies PD1-1, PD1-2, PD1-3, PD1-4, PD1-5.

SEQ ID Number:	Brief description of sequence	Sequence
SEQ ID NO:256	PD1-1 HC	EVMLVESGGGLVQPGGSLRLSCTASGFTFSASAMSWV RQAPGKGLEWVAYISGGGGDTYYSSSVKGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCARHSNVNYYAMDY WGQGTLLTVSSASTKGPSVFPLAPCSRSTSESTAALGC LVKDYPPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGKTYTCNVDPKPSNTKVDKRVESKYG PPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNST YRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIS KAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVD KSRWQEGNVFSCSVMEALHNHYTQKSLSLSLG
SEQ ID NO:257	PD1-1 LC	EIVLTQSPATLSLSPGERATMSCRASENIDTSGISFMNW YQQKPGQAPKLLIYVASNQGSIGIPARFSGSGSTDFTLT ISRLEPEDFAVYYCQQSKEVPWTFGQGTKLEIKRTVAA PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID	PD1-2 HC	EVMLVESGGGLVQPGGSLRLSCTASGFTFSASAMSWV

SEQ ID Number:	Brief description of sequence	Sequence
NO:258		RQAPGKGLEWVAYISGGGGDTYYSSSVKGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCARHSNPNYAMDY WGQGTLTVTVSSASTKGPSVFPLAPCSRSTSESTAALGC LVKDYPPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGKTYTCNVDPHKPSNTKVDKRVESKYG PPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNST YRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIS KAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVD KSRWQEGNVFSCSVMHEALHNHYTQKSLSLGLG
SEQ ID NO:259	PD1-2 LC	EIVLTQSPATLSLSPGERATMSCRASENIDTSGISFMNW YQQKPGQAPKLLIYVASNQSGIPARFSGSGGTDFLT ISRLEPEDFAVYYCQQSKEVPWTFGGGTKLEIKRTVAA PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO:260	PD1-3 HC	EVMLVESGGGLVQPGGSLRLSCTASGFTFSKSAMSWV RQAPGKGLEWVAYISGGGGDTYYSSSVKGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCARHSNPVNYAMDY WGQGTLTVTVSSASTKGPSVFPLAPCSRSTSESTAALGC LVKDYPPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGKTYTCNVDPHKPSNTKVDKRVESKYG PPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNST YRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIS KAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVD KSRWQEGNVFSCSVMHEALHNHYTQKSLSLGLG

SEQ ID Number:	Brief description of sequence	Sequence
SEQ ID NO:261	PD1-3 LC	EIVLTQSPATLSLSPGERATMSCRASENIDVSGISFMNW YQQKPGQAPKLLIYVASNQGSVIPARFSGSGSGTDFTLT ISRLEPEDFAVYYCQQSKEVPWTFGQGTKLEIKRTVAA PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO:262	PD1-4 HC	EVMLVESGGGLVQPGGSLRLSCTASGFTFSKSAMSWV RQAPGKGLEWVAYISGGGGDTYYSSSVKGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCARHSNVNYYAMDY WGQGTLLTVSSASTKGPSVFPLAPCSRSTSESTAALGC LVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTKTYTCNVDHKPSNTKVDKRVESKYG PPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNST YRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIS KAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVD KSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLG
SEQ ID NO:263	PD1-4 LC	EIVLTQSPATLSLSPGERATMSCRASENIDVSGISFMNW YQQKPGQAPKLLIYVASNQGSVIPARFSGSGSGTDFTLT ISRLEPEDFAVYYCQQSKEVPWTFGQGTKLEIKRTVAA PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO:264	PD1-5 HC	EVMLVESGGGLVQPGGSLRLSCTASGFTFSKSAMSWV RQAPGKGLEWVAYISGGGGDTYYSSSVKGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCARHSNVNYYAMDY WGQGTLLTVSSASTKGPSVFPLAPCSRSTSESTAALGC LVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL

SEQ ID Number:	Brief description of sequence	Sequence
		SSVVTVPSSSLGKTYTCNVDPKPSNTKVDKRVESKYG PPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNST YRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIS KAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVD KSRWQEGNVFSCSVMHENHHTQKSLSLSLG
SEQ ID NO:265	PD1-5 LC	EIVLTQSPATLSLSPGERATMSCRASENIDVSGISFMNW YQQKPGQAPKLLIYVASNQGGIPARFSGSGGTDFTLT ISRLEPEDFAVYYCQQSKEVPWTFGQGTKLEIKRTVAA PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC

According to these preferred embodiments and any other of the aspects of the present invention, antibodies PD1-1, PD1-2, PD1-3, PD1-4 and PD1-5 are antibody molecules as disclosed in WO2017/198741, and are defined by the sequences as shown in Table A
5 above.

Accordingly, PD1-1 has a heavy chain comprising the amino acid sequence of SEQ ID NO:256 and a light chain comprising the amino acid sequence of SEQ ID NO:257;

PD1-2 has a heavy chain comprising the amino acid sequence of SEQ ID NO:258 and a light chain comprising the amino acid sequence of SEQ ID NO:259;

10 PD1-3 has a heavy chain comprising the amino acid sequence of SEQ ID NO:260 and a light chain comprising the amino acid sequence of SEQ ID NO:261;

PD1-4 has a heavy chain comprising the amino acid sequence of SEQ ID NO:262 and a light chain comprising the amino acid sequence of SEQ ID NO:263; and

PD1-5 has a heavy chain comprising the amino acid sequence of SEQ ID NO:264 and a
15 light chain comprising the amino acid sequence of SEQ ID NO:265.

The above also includes the use of the binding proteins of the invention in various methods of treating the above diseases by administering a therapeutically effective dose to a patient in need thereof, as well as the use of these binding proteins for the manufacture of medicaments for the treatment of such diseases, as well as pharmaceutical compositions including such binding proteins of the invention, as well as the preparation and/or manufacture of medicaments including such binding proteins of the invention, and the like.

COMBINATIONS WITH OTHER ACTIVE SUBSTANCES OR TREATMENTS

A binding protein of the invention may be used on its own or in combination with one or more additional therapeutic agents, in particular in combination with a chemotherapeutic agent like DNA damaging agents, a therapeutically active compound that inhibits angiogenesis, a signal transduction pathway inhibitor, an EGFR inhibitor, an immune modulator, an immune checkpoint inhibitor, a mitotic checkpoint inhibitor or a hormonal therapy agent.

The additional therapeutic agent may be administered simultaneously with, optionally as a component of the same pharmaceutical preparation, or before or after administration of the DLL3/CD3 binding protein.

Cytostatic and/or cytotoxic active substances which may be administered in combination with binding molecules of the invention include, without being restricted thereto, hormones, hormone analogues and antihormones, aromatase inhibitors, LHRH agonists and antagonists, inhibitors of growth factors (growth factors such as for example platelet derived growth factor (PDGF), fibroblast growth factor (FGF), vascular endothelial growth factor (VEGF), epidermal growth factor (EGF), insuline-like growth factors (IGF), human epidermal growth factor (HER, e.g. HER2, HER3, HER4) and hepatocyte growth factor (HGF)), inhibitors are for example (anti-)growth factor antibodies, (anti-)growth factor receptor antibodies and tyrosine kinase inhibitors, such as for example cetuximab, gefitinib, afatinib, nintedanib, imatinib, lapatinib, bosutinib and trastuzumab; antimetabolites (e.g. antifolates such as methotrexate, raltitrexed, pyrimidine analogues such as 5-fluorouracil (5-FU), gemcitabine, irinotecan, doxorubicin, TAS-102, capecitabine and gemcitabine, purine and adenosine analogues such as mercaptopurine, thioguanine, cladribine and pentostatin, cytarabine (ara C), fludarabine); antitumor antibiotics (e.g. anthracyclins); platinum derivatives (e.g. cisplatin, oxaliplatin,

carboplatin); alkylation agents (e.g. estramustin, meclorethamine, melphalan, chlorambucil, busulphan, dacarbazine, cyclophosphamide, ifosfamide, temozolomide, nitrosoureas such as for example carmustin and lomustin, thiotepa); antimitotic agents (e.g. Vinca alkaloids such as for example vinblastine, vindesine, vinorelbine and vincristine; and
 5 taxanes such as paclitaxel, docetaxel); angiogenesis inhibitors, including bevacizumab, ramucirumab and aflibercept, tubulin inhibitors; DNA synthesis inhibitors, PARP inhibitors, topoisomerase inhibitors (e.g. epipodophyllotoxins such as for example etoposide and etopophos, teniposide, amsacrin, topotecan, irinotecan, mitoxantrone), serine/threonine kinase inhibitors (e.g. PDK1 inhibitors, Raf inhibitors, A-Raf inhibitors,
 10 B-Raf inhibitors, C-Raf inhibitors, mTOR inhibitors, mTORC1/2 inhibitors, PI3K inhibitors, PI3K α inhibitors, dual mTOR/PI3K inhibitors, STK33 inhibitors, AKT inhibitors, PLK1 inhibitors (such as volasertib), inhibitors of CDKs, including CDK9 inhibitors, Aurora kinase inhibitors), tyrosine kinase inhibitors (e.g. PTK2/FAK inhibitors), protein-protein interaction inhibitors, MEK inhibitors, ERK inhibitors, FLT3 inhibitors,
 15 BRD4 inhibitors, IGF-1R inhibitors, Bcl-xL inhibitors, Bcl-2 inhibitors, Bcl-2/Bcl-xL inhibitors, ErbB receptor inhibitors, BCR-ABL inhibitors, ABL inhibitors, Src inhibitors, rapamycin analogs (e.g. everolimus, temsirolimus, ridaforolimus, sirolimus), androgen synthesis inhibitors, androgen receptor inhibitors, DNMT inhibitors, HDAC inhibitors, ANG1/2 inhibitors, CYP17 inhibitors, radiopharmaceuticals, immunotherapeutic agents
 20 such as immune checkpoint inhibitors (e.g. CTLA4, PD1, PD-L1, LAG3, and TIM3 binding molecules / immunoglobulins, such as ipilimumab, nivolumab, pembrolizumab) and various chemotherapeutic agents such as amifostin, anagrelid, clodronate, filgrastin, interferon, interferon alpha, leucovorin, rituximab, procarbazine, levamisole, mesna, mitotane, pamidronate and porfimer; proteasome inhibitors (such as Bortezomib); Smac
 25 and BH3 mimetics; agents restoring p53 functionality including mdm2-p53 antagonist; inhibitors of the Wnt/beta-catenin signaling pathway; and/or cyclin-dependent kinase 9 inhibitors.

Particularly preferred are treatments with the binding molecules of the invention in combination with one or more immunotherapeutic agents, including anti-PD-1 and anti-
 30 PD-L1 agents and anti LAG3 agents: Exemplary anti-PD1 agents include but are not limited to anti-PD-1 antibody PDR-001, pembrolizumab, nivolumab, pidilizumab and PD1-1, PD1-2, PD1-3, PD1-4 and PD1-5 as disclosed herein (Table A) and in WO2017/198741. Exemplary anti-PDL-1 agents include but are not limited to

atezolizumab, avelumab and durvalumab. In preferred embodiments, the binding molecule of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is combined with PD1-1. In preferred embodiments, 5 the binding molecule of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is combined with PD1-2. In preferred embodiments, the binding molecule of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, 10 DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is combined with PD1-3. In preferred embodiments, the binding molecule of the invention (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is combined with PD1-4. In preferred embodiments, the binding molecule of the invention 15 (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) is combined with PD1-5.

In certain embodiments, the additional therapeutic agent may be a further immunotherapeutic agent, such as modulators of: TIM-1, TIM-3, TIM-4, PD-L2, LAG3, 20 CTLA-4, Galectin 9, Galectin-1, CD69, CD113, GPR56, CD48, GARP, CAECAM-1, BTLA, TIGIT, CD160, LAIR1, 2B4, CEACAM, CD39, TGF β , IL-10, Fas ligand, ICOS, B7 family (B7-1, B7-2, B7-H1 (PDL-1), B7-DC (PD-L2), B7-H2 (ICOS-L), B7-H3, B7-H4, B7-H5 (VISTA), or B7-H6).

In some embodiments, the additional immunotherapeutic agent is a member of the TNF 25 family of molecules that bind to cognate TNF receptor family members, which include CD40 and CD40L, OX-40, OX-40L, CD70, CD27L, CD30, CD30L, 4-1BBL, CD137, GITR, TRAIL/Apo2-L, TRAILR1/DR4, TRAILR2/DR5, TRAILR3, TRAILR4, OPG, RANK, RANKL, TWEAKR/Fn14, TWEAK, BAFFR, EDAR, XEDAR, TACI, APRIL, BCMA, LIGHT, DcR3, HVEM, VEGI/TLIA, TRAMP/DR3, EDAR, EDA1, XEDAR, 30 EDA2, TNFR1, Lymphotoxin α /TNF β , TNFR2, TNF α , LT β R, Lymphotoxin α 1 β 2, FAS, FASL, RELT, DR6, TROY, NGFR.

In some embodiments, the additional therapeutic agent is a SMAC mimetic. SMAC

mimetics are compounds which bind to Inhibitor of Apoptosis Proteins (IAPs) and have immune modulatory function and mediate the induction of systemic cytokines (*e.g.* IL-6, TNF α *etc.*) and chemokines (*e.g.* MCP-1) when administered to animals or humans. In some embodiments, the SMAC mimetic is (i) LCL161, *i.e.* compound A in example 1 of
5 WO 2008/016893 (page 28/29; [122]), or a pharmaceutically acceptable salt thereof; (ii) the SMAC mimetic known as Debio-1143, or a pharmaceutically acceptable salt thereof; (iii) the SMAC mimetic known as birinapant, or a pharmaceutically acceptable salt thereof; (iv) the SMAC mimetic known as ASTX-660, or a pharmaceutically acceptable salt thereof; or (v) the SMAC mimetic known as CUDC-427, or a pharmaceutically acceptable
10 salt thereof.

In some embodiments, the additional immunotherapeutic agent is selected from (i) antagonists of cytokines that inhibit T cell activation (*e.g.*, IL-6, IL-10, TGF-B, VEGF; "immunosuppressive cytokines") and/or (ii) agonists of cytokines that stimulate T cell activation and/or cytokines such as IL2, for stimulating an immune response, *e.g.*, for
15 treating proliferative diseases, such as cancer.

In some embodiments, the additional immunotherapeutic agent is an agonist of a protein that stimulates T cell activation, such as CD28, GITRL, OX40L, CD27, and CD28H.

In some embodiments, the additional therapeutic agent is an oncolytic virus including but not limited to an oncolytic virus derived from vaccinia virus, adenovirus, (AdV), herpes
20 simplex virus (HSV1 or HSV2), reovirus, myxoma virus (MYXV), poliovirus, vesicular stomatitis virus (VSV), Maraba virus, varicella virus, measles virus (MV), or Newcastle disease virus (NDV).

DIAGNOSTIC USES

25 The anti-DLL3 antibody molecules of the invention are useful in diagnostic and prognostic methods and can be employed for labelling, localizing, or identifying cells or tissues expressing DLL3 (*e.g.* in ELISA assays, FACS analysis, immunohistology or the like) by attaching a dye, a drug or another molecule with binding specificity for a different antigen. For example, a detectable label can be conjugated to the anti-DLL3 antibody molecule of
30 the invention or a secondary reagent binding specifically to the antibody molecule of the invention and being conjugated with a detectable label (*e.g.* a secondary antibody) can be used in such diagnostic methods. In some embodiments, DLL3 specific antibodies

specifically bind to DLL3 expressed in cells, either in the cytoplasm and/or at the cell surface, and are used for localizing and/or identifying such cells. In some embodiments, the DLL3 specific antibodies provided herein are used for identifying cells or tissues expressing DLL3 (e.g. tumor cells).

- 5 Antibodies of the invention are selected such that they have a high level of epitope binding specificity and high binding affinity to the DLL3 polypeptide. In general, the higher the binding affinity of an antibody, the more stringent wash conditions can be performed in an immunoassay to remove nonspecifically bound material without removing the target polypeptide.
- 10 Thus, further provided herein are methods of detecting DLL3 in a biological sample. Specifically, in one aspect the invention provides a method of detecting DLL3 in a sample (e.g. a tissue sample, a tumor tissue sample) comprising
- (a) contacting the sample with an antibody of the invention; and
 - (b) detecting anti-DLL3 antibody bound to DLL3.
- 15 A biological sample can be obtained/isolated from any tissue (including biopsies), cell or body fluid of a subject. In a preferred embodiment, the sample is a tissue sample, preferably a tumor tissue sample. In a specific embodiment, the sample is a fixed (tumor) tissue sample, preferably a formalin-fixed, paraffin-embedded (FFPE) tissue sample, more preferably a formalin-fixed, paraffin-embedded (FFPE) tumor tissue sample.
- 20 The samples are typically divided into several portions and affixed to a medium for microscopic analysis, such as a microscope slide. Where the sample is a tissue sample, the several portions may be tissue sections. In some embodiments, serial sections are taken from FFPE tissue samples. In some embodiments, serial sections are taken from a plurality of different sites of a FFPE block, which can be done to capture both intra-section
- 25 heterogeneity and intra-block heterogeneity. In some embodiments, serial sections are taken from a plurality of different biopsy samples taken from different locations in the same tumor, which can be done to capture both intra-section heterogeneity and intra-tumor heterogeneity. Typically, tissue samples first undergo deparaffinization, antigen retrieval (e.g. with Proteinase K) before antigen detection. The tissue sample can be
- 30 obtained/isolated from a variety of tissues (e.g. from a biopsy), specifically tissues obtained from tumors including but not limit to SCLC, LCNEC, glioma, glioblastoma,

melanoma, or other neuroendocrine tumors (e.g., neuroendocrine prostate cancer or neuroendocrine pancreatic cancer, small cell bladder cancer).

Further exemplary tumors from which a (tissue) sample can be obtained/isolated include lung cancer; preferably SCLC, NSCLC or LCNEC; breast; cervical; colon; colorectal; endometrial; head and neck; liver (hepatoblastoma or hepatocellular carcinoma); ovarian; pancreatic; prostate; skin; gastric; testis; thyroid; adrenal; renal; bladder; uterine; esophageal; urothelial cancer; brain tumor; lymphoma; Ewing sarcoma; and other neuroendocrine (including pseudo-endocrine tumors) and small blue round cell tumors.

Exemplary methods of DLL3 detection in a sample are immunocytochemistry (ICC), immunohistochemistry (IHC), Western Blotting, Flow cytometry and/or ELISA.

In a preferred embodiment, DLL3 is detected using IHC.

In a specific aspect of the invention, the method of detecting DLL3 in a tissue sample comprises

- a) contacting a tissue sample (e.g. a tumor tissue such as SCLC, glioblastoma or neuroendocrine tumors), preferably said tissue sample is a fixed tissue sample (e.g., formalin-fixed and paraffin embedded), with an antibody of the invention (e.g. DLL3#1 or DLL3#5)
- b) permitting formation of antibody-antigen complexes in the sample, and
- c) detecting the anti-DLL3 antibody bound to DLL3.

In some embodiments of the methods of the invention, the anti-DLL3 antibody is detected by a detectable signal, preferably, the detectable signal is generated by a detectable label. In a preferred embodiment of the method of the invention, the detectable signal is detected in an IHC assay. In some embodiments, the detectable signal is detected in an ELISA assay, by flow cytometry or by Western blot. In some embodiments, the methods described herein can be used to detect DLL3 present on the cell surface of a cell (e.g. a tumor cell), for example using flow cytometry. In some embodiments, the methods described herein can be used to detect the presence of DLL3 in a tissue sample (e.g. tumor tissue sample) using ELISA, Western blot or IHC.

In some embodiments, the detectable label is directly conjugated to the anti-DLL3 antibody and thus is deposited on the sample upon binding of the anti-DLL3 antibody to DLL3, this is generally referred to as a direct labeling method). Direct labeling methods

are often more directly quantifiable, but often suffer from a lack of sensitivity. In other embodiments, deposition of the detectable label is effected by the use of a secondary detection reagent binding specifically to the antibody molecule of the invention and being conjugated with a detectable label (e.g. a secondary antibody), this is generally referred to as an indirect labeling method. In some embodiments, the specific secondary detection reagent may be a species-specific secondary antibody, an anti-hapten antibody binding to a hapten-conjugated anti-DLL3 antibody, or a biotin-binding protein bound to a biotinylated anti-DLL3 antibody).

Particular examples of detectable labels that can be conjugated either to the anti-DLL3 antibody of the invention or to a secondary detection reagent, include chromogenic, fluorescent, phosphorescent, luminescent and radioactive molecules and materials, catalysts (such as enzymes) that convert one substance into another substance to provide a detectable difference (such as by converting a colorless substance into a colored substance or vice versa, or by producing a precipitate or increasing sample turbidity), haptens that can be detected through antibody-hapten binding interactions using additional detectably labeled antibody conjugates, and paramagnetic and magnetic molecules or materials..

For example, the detectable label can be an enzyme such as horseradish peroxidase (HRP), alkaline phosphatase (AP), acid phosphatase, glucose oxidase, β -galactosidase, β -glucuronidase, and β -lactamase; a fluorophore such as fluoresceins, luminophores, coumarins, BODIPY dyes, resorufins, and rhodamines; nanoparticles such as quantum dots (U.S. Pat. Nos. 6,815,064, 6,682,596 and 6,649,138); metal chelates such as DOTA and DPTA chelates of radioactive or paramagnetic metal ions like Gd $<3+>$; and liposomes, for example, liposomes containing trapped fluorescent molecules.

Where the detectable label includes an enzyme, a detectable substrate such as a chromogen, a fluorogenic compound, or a luminogenic compound can be used in combination with the enzyme to generate a detectable signal. Particular examples of chromogenic compounds include diaminobenzidine (DAB), 4-nitrophenylphosphate (pNPP), fast red, bromochloroindolyl phosphate (BCIP), nitro blue tetrazolium (NBT), BCIP/NBT, fast red, AP Orange, AP blue, tetramethylbenzidine (TMB), 2,2'- azino-di-[3-ethylbenzothiazoline sulphonate] (ABTS), o-dianisidine, 4- chloronaphthol (4-CN), nitrophenyl- -D-galactopyranoside (ONPG), o- phenylenediamine (OPD), 5-bromo-4-chloro-3-indolyl- -galactopyranoside (X-Gal), methylumbelliferyl- -D-galactopyranoside (MU-Gal), p-nitrophenyl- a-D-galactopyranoside (PNP), 5-bromo-4-chloro-3-indolyl- -D-

glucuronide (X-Gluc), 3-amino-9-ethyl carbazol (AEC), fuchsin, iodonitrotetrazolium (INT), tetrazolium blue and tetrazolium violet.

In some examples, the detectable moiety is a fluorophore, which belongs to several common chemical classes including coumarins, fluoresceins (or fluorescein derivatives and
5 analogs), rhodamines, resorufins, luminophores and cyanines.

In other embodiments, the detectable moiety is a molecule detectable via brightfield microscopy, such as dyes including diaminobenzidine (DAB), 4-(dimethylamino) azobenzene-4'- sulfonamide (DABSYL), tetramethylrhodamine (DISCOVERY Purple), N,N'-biscarboxypentyl-5,5'-disulfonato-indo- dicarbocyanine (Cy5), and Rhodamine 110
10 (Rhodamine).

Haptens are small molecules that are specifically bound by antibodies, although by themselves they will not elicit an immune response in an animal and must first be attached to a larger carrier molecule such as a protein to generate an immune response. Examples of haptens include di-nitrophenyl, biotin, digoxigenin, and fluorescein.

15 Further provided herein in one aspect are methods of diagnosing or identifying a tumor as DLL3 expressing tumor (expressing DLL3 in the cytoplasm and/or the cell surface), comprising detecting DLL3 in a sample (e.g. a tumor tissue sample) of the subject using the anti-DLL3 antibodies (e.g. DLL3#1 or DLL3#5) of the invention (e.g. using IHC on a tumor tissue sample). In a further aspect, provided herein are methods of selecting a DLL3
20 targeted therapy for a subject, comprising detecting DLL3 in a sample (e.g. a tumor tissue sample) of the subject using the anti-DLL3 antibodies (e.g. DLL3#1 or DLL3#5) of the invention (e.g. using IHC on a tumor tissue sample). In a further aspect, provided herein are methods of monitoring the therapeutic effect (e.g. of a DLL3-targeted therapy) in a subject, comprising detecting DLL3 in a sample (e.g. a tumor tissue sample) of the subject
25 using the anti-DLL3 antibodies (e.g. DLL3#1 or DLL3#5) of the invention (e.g. using IHC on a tumor tissue sample).

The anti-DLL3 antibody molecules of the invention can also be used for targeting cells. In some embodiments, the DLL3 specific antibodies provided herein are used for delivering a drug or cytotoxic agent to a target cell (e.g. a tumor cell expressing DLL3) by attaching
30 such drug or cytotoxic agent to said DLL3 antibody, thereby, for example, killing said target cell.

KITS

The invention also encompasses kits comprising at least a multi-specific binding protein of the invention (e.g., any one of DLL3#1/CD3#1, DLL3#2/CD3#1, DLL3#3/CD3#1, DLL3#4/CD3#1, DLL3#5/CD3#1, DLL3#6/CD3#1, DLL3#7/CD3#1, DLL3#8/CD3#1, DLL3#9/CD3#1, DLL3#10/CD3#1, DLL3#11/CD3#1, DLL3#12/CD3#1, DLL3#13/CD3#1, DLL3#14/CD3#1, DLL3#15/CD3#1, DLL3#16/CD3#1, DLL3#17/CD3#1, DLL3#18/CD3#1, DLL3#1/CD3#2, DLL3#2/CD3#2, DLL3#3/CD3#2, DLL3#4/CD3#2, DLL3#5/CD3#2, DLL3#6/CD3#2, DLL3#7/CD3#2, DLL3#8/CD3#2, DLL3#9/CD3#2, DLL3#10/CD3#2, DLL3#11/CD3#2, DLL3#12/CD3#2, DLL3#13/CD3#2, DLL3#14/CD3#2, DLL3#15/CD3#2, DLL3#16/CD3#2, DLL3#17/CD3#2, DLL3#18/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#3, DLL3#3/CD3#3, DLL3#4/CD3#3, DLL3#5/CD3#3 and DLL3#6/CD3#3, DLL3#7/CD3#3, DLL3#8/CD3#3, DLL3#9/CD3#3, DLL3#10/CD3#3, DLL3#11/CD3#3, DLL3#12/CD3#3, DLL3#13/CD3#3, DLL3#14/CD3#3, DLL3#15/CD3#3, DLL3#16/CD3#3, DLL3#17/CD3#3, DLL3#18/CD3#3, preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) and optionally one or more other components selected from the group consisting of other drugs used for the treatment of the diseases and disorders as described above.

In one embodiment, the kit includes a composition containing an effective amount of a binding protein of the invention in unit dosage form.

The invention also encompasses kits comprising at least a multi-specific binding protein of the invention, and one or more other components selected from the group consisting of other drugs used for the treatment of the diseases and disorders as described above.

In one embodiment, the kit includes a composition containing an effective amount of a multi-specific binding protein of the invention in unit dosage form (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3). In a further embodiment the kit includes both a composition containing an effective amount of a multi-specific binding protein of the invention in unit dosage form (preferably any one of DLL3#1/CD3#1, DLL3#1/CD3#2, DLL3#1/CD3#3, DLL3#2/CD3#1, DLL3#2/CD3#2, DLL3#2/CD3#3, DLL3#3/CD3#1, DLL3#3/CD3#2, DLL3#3/CD3#3) and a composition

containing an effective amount of a PD-1 antagonist in unit dosage form, such as an anti PD-1 antibody, most preferably PD1-1, PD1-2, PD1-3, PD1-4, and PD1-5 as described herein (e.g. Table A) and in WO2017/198741.

In some embodiments, the kit comprises a sterile container which contains such a composition; such containers can be boxes, ampoules, bottles, vials, tubes, bags, pouches, blister-packs, or other suitable container forms known in the art. Such containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding medicaments. Further, the kit may comprise the pharmaceutical composition in a first container with the binding protein of the invention in lyophilized form and a second container with a pharmaceutically acceptable diluent (e.g., sterile water) for injection. The pharmaceutically acceptable diluent can be used for reconstitution or dilution of the binding protein.

If desired, a multi-specific binding protein of the invention, is provided together with instructions for administering the multi-specific binding proteins to a subject having cancer. The instructions will generally include information about the use of the composition for the treatment or prevention of a cancer. In other embodiments, the instructions include at least one of the following: description of the therapeutic agent; dosage schedule and administration for treatment or prevention of cancer or symptoms thereof; precautions; warnings; indications; counter-indications; overdose information; adverse reactions; animal pharmacology; clinical studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder supplied in or with the container.

As set forth herein, the present disclosure further provides diagnostic methods for determining the expression of DLL3 in a biological sample. In one aspect, the present disclosure provides kits for performing these methods as well as instructions for carrying out the methods of the present disclosure such as collecting tissue and/or performing the detection assay, and/or analyzing the results.

The kit comprises, or alternatively consists essentially of, or yet further consists of, a DLL3 antibody composition of the invention (e.g., monoclonal antibodies) and instructions for use. The kits are useful for detecting the presence of DLL3 polypeptides in a biological sample e.g., any body fluid or biopsy samples of body tissue. In some embodiments, the kit

can comprise: one or more DLL3 antibodies capable of binding a DLL3 polypeptide in a sample (e.g., anti-DLL3 antibody DLL3#1 or DLL3#5); means for detecting the DLL3 polypeptide in the sample; and optionally means for comparing the signal/amount of the DLL3 polypeptide in the sample with a control. One or more of the anti-DLL3 antibodies
5 of the invention may be labeled. The kit components, (e.g., reagents) can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect the DLL3 polypeptides. In certain embodiments, the kit comprises an anti-DLL3 antibody of the invention (a first antibody) and a second, different antibody which binds to either the DLL3 polypeptide or the first antibody and is conjugated to a detectable label.

10 The kit can also comprise, e.g., a buffering agent, a preservative or a protein-stabilizing agent. The kit can further comprise components necessary for detecting the detectable-label, e.g., an enzyme or a substrate. The kit can also contain a control sample or a series of control samples, which can be assayed and compared to the test sample. Each component of the kit can be enclosed within an individual container and all of the various containers
15 can be within a single package, along with instructions for interpreting the results of the assays performed using the kit. The kits of the present disclosure may contain a written product on or in the kit container. The written product describes how to use the reagents contained in the kit.

As amenable, these suggested kit components may be packaged in a manner customary for
20 use by those of skill in the art. For example, these suggested kit components may be provided in solution or as a liquid dispersion or the like.

EXAMPLES

25 The following examples illustrate the invention. These examples should not be construed as to limit the scope of this invention.

Example 1: Design and construction of CD3/DLL3 binding proteins

The present inventors have developed multi-specific binding proteins that bind DLL3 and
30 CD3 and that induce T-cell activation leading to lysis of DLL3-expressing tumor cells. The molecular design used has an IgG antibody scaffold and an IgG-like structure. It features knob-in-hole technology in the Fc for hetero-dimerization of the Knob (anti-DLL3) and

Hole (anti-CD3) arms. In addition, the binding protein has flexible peptide sequences between the light and the corresponding heavy chain in each arm. Thus, the binding protein comprises two arms, one binding to CD3, the other one binding to DLL3, each arm comprising a single chain Fab and an Fc region.

- 5 Preferably the binding molecule is bispecific and bivalent (monovalent for each of the two targets).

Preparation of binding domains that recognize DLL3 and CD3 using high throughput V gene recovery from hybridomas and cultured single B cells.

- 10 To obtain anti-DLL3 binders, hybridomas or single B cells derived from DLL3 immunized wild-type and AlivaMab humanized mice (Ablexis, San Francisco, CA, USA: AlivaMab transgenic mouse platform with human immunoglobulin loci) were cultured in vitro. Supernatants were screened for reactivity against recombinant human DLL3, by AlphaLISA (PerkinElmer, Waltham, MA, USA), and against SHP77 cells (ATCC®, CRL-15 2195TM) expressing human DLL3, by Flow Cytometry.

- Immunoglobulin (Ig) VH and VL genes were then amplified from identified positive clones. To isolate RNA from hybridomas, about 2×10^6 cells from single clones were pelleted and used as source material. For single B cells, 100 to 500 cells expanded from singularly isolated B cells were used as source material. RNA was isolated using RNeasy Plus (Qiagen, Hilden, Germany). cDNA was then synthesized using Smarter cDNA 20 synthesis kit (Clontech, Mountain View, CA) according to manufacturer's instructions. To facilitate cDNA synthesis, oligodT was used to prime reverse transcription of all messenger RNAs followed by "5' capping" with a Smarter IIA oligonucleotide. Subsequent amplification of the VH and VL fragments was performed using a 2-step PCR 25 amplification using 5' primers targeting the Smarter IIA cap and 3' primers targeting consensus regions in CH1. Briefly, each 50µl PCR reaction consists of 20µM of forward and reverse primer mixes, 25µl of PrimeStar Max DNA polymerase premix (Clontech), 2µl of unpurified cDNA, and 21µl of double-distilled H₂O. The cycling program starts at 94 °C for 3 min, followed by 35 cycles (94°C for 30 Sec, 50°C for 1 min, 68 °C for 1 min), 30 and ends at 72 °C for 7 min. The second round PCR was performed with VL and VH 2nd round primers containing 15bp complementary extensions that "overlap" respective regions in their respective pTT5 mother vector (VH and VL). Second round PCR was

performed with the following program: 94 °C for 3 min; 35 cycles (94 °C for 30 Sec, 50°C for 1 min, 68 °C for 1 min), and ends at 72 °C for 7 min.

In-Fusion® HD Cloning Kit (Clontech, U.S.A.) was used for directional cloning of VL gene into a pTT5 huIgK vector and VH gene into a pTT5 huIgG1KO vector. To facilitate

5 In-Fusion® HD Cloning, PCR products were purified and treated with Cloning Enhancer before In-Fusion HD Cloning. Cloning and transformation were performed according to manufacturer's protocol (Clontech, U.S.A.). Mini-prep DNAs were subjected to Sanger sequencing to confirm that complete V-gene fragments were obtained.

10 Using this methodology, pairs of Ig VH and VL genes encoding binding domains with specificity for DLL3 were prepared. Recombinant antibodies were produced by transient transfection of CHO-E37 cells with the corresponding heavy and light chain-encoding plasmids.

Confirmatory screening of recombinant antibodies

15 Supernatants containing expressed recombinant antibodies were assayed by flow cytometry for binding to cell lines expressing human or cyno DLL3. Briefly, cells were incubated with recombinant supernatants, washed, and bound mAbs from the supernatants were detected with anti-human-IgG-APC (Jackson ImmunoResearch 109-136-098). Signal-to-background ratios (S/B) were calculated by dividing the median fluorescence
20 intensity (MFI) of the sample by that of isotype control (variable regions against an unrelated protein and different constant region backbones).

Surface Plasmon Resonance (SPR) on Biacore 400 was performed on recombinant supernatants. Briefly, the non-optimized IgGs in the HTP supernatants were captured via Protein A/G onto the sensor surface for 60 sec at 10 ul/min. Binding of 100 nM human
25 DLL3 to the captured IgGs was monitored for 180 sec of association at 30 ul/min, followed by 120 sec of dissociation in the HBS-EP buffer. Regeneration of the Protein A/G surface was performed with Glycine pH 2.1 in between each binding cycle. The following materials were used in this assay: Protein reagent: recombinantly expressed human DLL3. System running buffer: HBS-EP (10 mM HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA,
30 and 0.005% v/v polysorbate P20). Capturing reagent: Protein A/G G (ThermoFisher Scientific, Waltham, MA, USA), with specificity towards all human IgG isotypes.

Clones of interest (with $K_d < 300 \text{ pM}$) were selected for multispecific formatting. Multispecific binding proteins were generated and further evaluated in mechanistic and functional screening (such as cell binding, cytotoxicity and T cell activation as described below).

5

Humanization of DLL3 and CD3 binders

Sequences of DLL3 binders as described above as well as CD3 binders described in the literature (Pessano et al., EMBO J. 1985 Feb; 4(2): 337-44; Salmerón A et al., J Immunol. 1991 Nov 1; 147(9):3047-52) were humanized and/or optimized. Sequence optimization/humanization of antibodies is a methodology to engineer antibodies raised in non-human species (against a specific antigen/ epitope) for use as therapeutics that resemble antibodies produced in humans and thereby eliminating potential adverse effects such as immunogenicity while retaining the specificity. The sequence optimization/humanization approach utilized here was as described by Singh et al, 2015 (Singh S et al., mAbs 2015: 7(4):778-91). In brief, closely matching human germlines were identified *in silico*, and optimization/humanization variants were evaluated using a phage screening method. Final lead candidate sequences were selected based on binding, percent human score and Epivax (in silico predictive tool for potential immunogenicity) score.

Construction of bispecific proteins binding DLL3 and CD3

The variable regions of the DLL3 and CD3 binders were cloned into the expression vector pTT5 (National Research Council, Canada), using common molecular biology techniques to form bispecific binding proteins with one DLL3 specific binding unit comprising a single chain Fab binding to DLL3 and an Fc region (such binding unit also referred to herein as “DLL3 arm” or “DLL3 chain”) and a CD3 specific binding unit comprising a single chain Fab binding to CD3 and an Fc region (such binding unit also referred to herein as “CD3 arm” or “CD3 chain. The Fc regions of the DLL3 and CD3 arms include either “Knob” or “Hole” mutations (Atwell et al, JMB, 1997, 270, 26-35) and the respective chains are referred to as Knob or Hole chains. For multi-fragment DNA assembly, a Gibson-assembly and NEBuilder HiFi DNA assembly approaches were used, following manufacturer’s protocols (New England Biolabs, Ipswich, MA, USA). DNA mini-preps were sequenced.

Each expression vector contains eukaryotic promoter elements for the chain-encoding gene (DLL3 or CD3 arm/chain), i.e., the gene encoding the signal sequence and the light and heavy chain, an expression cassette for a prokaryotic selection marker gene such as ampicillin, and an origin of replication. These DNA plasmids were propagated in
5 ampicillin resistant *E. coli* colonies and cultures and were purified.

Example 2: Expression and purification of bispecific, bivalent binding proteins binding DLL3 and CD3

Bispecific molecules binding DLL3 and CD3 were produced by transient transfection of
10 CHO-E cells with the pTT5 vectors carrying the DLL3/CD3-chain-encoding genes. Briefly, transfected CHO-E cells growing in suspension in serum-free media were cultivated in shake flasks under agitation at 140 rpm, 37°C and 5% CO₂ and kept at conditions of exponential growth. On the day of transfection, cells were chemically transfected with Knob-chain plasmid and Hole-chain plasmid in 1:3 mass ratio. They were
15 then seeded at 1 to 2x10⁶ cells/ml in 1 L of Gibco® FreeStyle™ CHO expression medium (LifeTechnologies, NY, US). Cells were then incubated under orbital shaking for 10 days with one-time feeding of 200 ml commercial feed solution to allow expression of the proteins. Antibody titers in the cell culture supernatants were determined using an Octet® instrument (Pall ForteBio, CA, US) and protA biosensor tips according to
20 manufacturer's instructions.

Recombinant DLL3/CD3 binding proteins were purified from culture supernatant in a two-step process: first by Protein A affinity chromatography using MabSelect™ column (GE Healthcare); second, by Cation exchange chromatography using a Poros 50 HS column (Applied Biosystems, Carlsbad, CA, USA). The two-step purified material was stored in
25 final buffer of 50mM Sodium Acetate and 100mM NaCl, pH 5.0 Purity and degree of heterogeneity of the samples were assessed by analytical size-exclusion chromatography, mass spectrometry and analytical ultracentrifugation. Samples that were advanced for functional testing comprised two-step purified material, with about 99% monomer content.

30 Table 1: Amino acid sequences and SEQ ID NOs of CDRs, VH, VL, scFabs, DLL3-arm and CD3-arm sequences of the proteins/antibody constructs described herein:

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO: 1	DLL3#1 LCCDR1	RASQSVSSNFLV
SEQ ID NO:2	DLL3#1 LCCDR2	GASTRAS
SEQ ID NO:3	DLL3#1 LCCDR3	QQYGDSPYT
SEQ ID NO:4	DLL3#1 HCCDR1	GNTFTNYMH
SEQ ID NO:5	DLL3#1 HCCDR2	IIDPSVGSKSYAQKFLG
SEQ ID NO:6	DLL3#1 HCCDR3	AGKRFGESYFDY
SEQ ID NO:7	DLL3#2 LCCDR1	RASQGISNYLA
SEQ ID NO:8	DLL3#2 LCCDR2	AASSLQS
SEQ ID NO:9	DLL3#2 LCCDR3	LQHNSSPYT
SEQ ID NO:10	DLL3#2 HCCDR1	GYTFTSYMH
SEQ ID NO:11	DLL3#2 HCCDR2	IINPSGGSTSYAQKFQG
SEQ ID NO:12	DLL3#2 HCCDR3	GEAVGGNYYYYGMDV
SEQ ID NO:13	DLL3#3 LCCDR1	RASQGISNYLV
SEQ ID NO:14	DLL3#3 LCCDR2	AVSSLYS
SEQ ID NO:15	DLL3#3 LCCDR3	LQHDSYPYT
SEQ ID NO:16	DLL3#3 HCCDR1	GYTFTSYVH
SEQ ID	DLL3#3	IINPGGGTTSYAQKFLG

SEQ ID Number	Brief description of sequence	Sequence
NO:17	HCCDR2	
SEQ ID NO:18	DLL3#3 HCCDR3	GEAVTGNYFYYGMDV
SEQ ID NO:19	DLL3#4 LCCDR1	RASKSVSSFGYSFMH
SEQ ID NO:20	DLL3#4 LCCDR2	LASNLES
SEQ ID NO:21	DLL3#4 LCCDR3	QHSRELPWT
SEQ ID NO:22	DLL3#4 HCCDR1	VYTFTSYFMY
SEQ ID NO:23	DLL3#4 HCCDR2	EISPTNGNSNLNERFKN
SEQ ID NO:24	DLL3#4 HCCDR3	GGDGYLDY
SEQ ID NO:25	DLL3#5 LCCDR1	QASQDISNYLN
SEQ ID NO:26	DLL3#5 LCCDR2	DASNLET
SEQ ID NO:27	DLL3#5 LCCDR3	QQYDNLPTWT
SEQ ID NO:28	DLL3#5 HCCDR1	GFTFTGYIYH
SEQ ID NO:29	DLL3#5 HCCDR2	WINPNSGGTNYAQKFQG
SEQ ID NO:30	DLL3#5 HCCDR3	GWDY
SEQ ID NO:31	DLL3#6 LCCDR1	RASQDISNYFA
SEQ ID NO:32	DLL3#6 LCCDR2	AASTLQS
SEQ ID NO:33	DLL3#6 LCCDR3	QQLNSYPYT

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:34	DLL3#6 HCCDR1	GGSISSYFWS
SEQ ID NO:35	DLL3#6 HCCDR2	RIYTSGSTNYNPSLNS
SEQ ID NO:36	DLL3#6 HCCDR3	RGDWGGFDI
SEQ ID NO:37	DLL3#1 VL	EIVLTQSPGTLSSLSPGERATLSCRASQSVSSNFLVWYQQKPG QAPRPLIYGASTRASGIPDRFSGSGSGADFTLTISRLEPEDFAL YYCQQYGDSPTYFTGQGTLEIK
SEQ ID NO:38	DLL3#1 VH	QVQLVQSGAEVKKPGASVKVSKASGNTFTNYMHVWRQ APGPGLEWMGIIDPSVGSKSYAQKFLGRVTIARDTSTSTVFL DLYSLRSEDNAVYFCARAGKRFGEYFDYWGQGLTVTVSS
SEQ ID NO:39	DLL3#2 VL	DIQMTQSPSAMSASVGDRVTITCRASQGISNYLAWFQQKPG KVPEPLIYAASSLQSGVPSRFSGSGSVTEFTLTISLQPEDFAT YYCLQHNSSPYFTGQGTKLEIK
SEQ ID NO:40	DLL3#2 VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYMHVWRQ APGQGLEWMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVY MELSSLRSEDNAVYYCARGEAVGGNYYYYGMDVWGQGT VTVSS
SEQ ID NO:41	DLL3#3 VL	DIQMTQSPSAMSASVGDRVTITCRASQGISNYLVWFQQKPG KAPKRLIYAVSSLYSGVPSRFSGSGSGTEFTLTISLQPEDFAT YYCLQHDSYPYFTGQGTKLEIK
SEQ ID NO:42	DLL3#3 VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYVHVWRQ APGQGLEWMVIINPGGGTTSYAQKFLGRVTMTRDTSTNTVY MELKSLRSEDNAVYYCARGEAVTGNYFYGYMDVWGQGT VTVSS
SEQ ID NO:43	DLL3#4 VL	DIVLTQSPASLAVSLGQRATISCRASKSVSSFGYSFMHWYQQ KPGQPPKLLIYLASNLESGVPARFSGSGSGTDFTLNHPVEEE DAATYYCQHSRELPTWTFGGGTKLEIK
SEQ ID NO:44	DLL3#4 VH	QVQLQQSGTELVKPGASVKLSCKASVYFTSYFMYWVKQR PGHGLEWIGEISPTNGNSNLNERFKNKATLTVDKSSSTAYM QLSSLTSEDSAVYYCTRGGDGYLDYWGQGTTLTVSS
SEQ ID	DLL3#5	DIQMTQSPSSLSASVGDRVTVTCQASQDISNYLNWYQQKPG

SEQ ID Number	Brief description of sequence	Sequence
NO:45	VL	KAPKLLIYDASNLETGVPSRFSGSGSGTDFTFTISSLQPEDIAT YYCQQYDNLPTWTFGQGTKVEIK
SEQ ID NO:46	DLL3#5 VH	QVQLVQSGAEVKKPGASVKVSCASGFTFTGYIHWVRQA PGQGLEWMGWINPNSGGTNYAQKFQGRVTMTRDSSINTAF MELSRLTSDDTAVYYCAAGWDYWGQGTLLVTVSS
SEQ ID NO:47	DLL3#6 VL	DIQLTQSPSFLSTSVGDRVITITCRASQDISNYFAWYQQKPGK APKLLIYAASLTQSGVPSRFSGSGSGTEFTLTISSLQPEDFAT YYCQQLNSYPYTFGQGTKLEIK
SEQ ID NO:48	DLL3#6 VH	QVQLQESGPGLVKPSETLSLTCTVSGGSISSYFWSWIRQPAG KGLEWIGRIYTSNSTNYNPSLNSRLTMSVDTSKNQFSLKLSS VTAADTAVYYCARRGDWGGFDIWGQGTVVTVSS
SEQ ID NO:49	DLL3#1 scFab	EIVLTQSPGTLSPGERATLSCRASQSVSSNFLVWYQQKPG QAPRPLIYGASTRASGIPDRFSGSGSGADFTLTISRLEPEDFAL YYCQQYGDSPYTFGQGTTLLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVSCASGNTFTNYMHVVR QAPGPGLEWMGIIDPSVGSKSYAQKFLGRVTIARDTSTSTVF LDLYSLRSEDTAVYFCARAGKRFGESYFDYWGQGTLLVTVSS ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV NPKPSNTKVDKRVEPKSC
SEQ ID NO:50	DLL3#2 scFab	DIQMTQSPSAMSASVGDRTITCRASQGISNYLAWFQQKPG KVPEPLIYAASSLQSGVPSRFSGSGSVTEFTLTISSLQPEDFAT YYCLQHNSSPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSG TASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVSCASGYTFTSYMHVVR QAPGQGLEWMGIINPSGGSTSYAQKFQGRVTMTRDSTSTV YMELSSLRSEDTAVYYCARGEAVGGNYYYGMDVWGQGT TVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP

SEQ ID Number	Brief description of sequence	Sequence
		VTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC
SEQ ID NO:51	DLL3#3 scFab	DIQMTQSPSAMSASVGDRVTITCRASQGISNYLVWFQQKPGKAPKRLIYAVSSLYSGVPSRFSGSGSGTEFTLTISLQPEDFATYYCLQHDSYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQVQLVQSGAEVKKPGASVKVSCKASGYTFTSYVHWVRQAPGQGLEWMVIINPGGGTTSYAQKFLGRVTMTRDTSTNTVYMELKSLRSEDYAVYYCARGEAVTGNFYGYMDVWGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC
SEQ ID NO:52	DLL3#4 scFab	DIVLTQSPASLAIVSLGQRATISCRASKSVSSFGYSFMHWYQQKPGQPPKLLIYLASNLESGVPARFSGSGSGTDFTLNIHPVEEEDAATYYCQHSRELPTWTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQVQLQQSGTELVKPGASVKLSCKASVYFTSYFMYWVKQRPFGHLEWIGEISPTNGNSNLNERFKNKATLTVDKSSSTAYMQLSSLTSEDSAVYYCTRGGDGYLDYWGQGTTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC
SEQ ID NO:53	DLL3#5 scFab	DIQMTQSPSSLSASVGDRVTITCRASQDISNYLNWYQQKPGKAPKLLIYDASNLETGVPSRFSGSGSGTDFTFTISLQPEDFATYYCQYDNLPTWTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQVQLVQSGAEVKKPGASVKVSCKASGYTFTGYYIHWV

SEQ ID Number	Brief description of sequence	Sequence
		RQAPGQGLEWMGWINPNSGGTNYAQKFQGRVTMTRDSSIN TAFMELSRLTSDDTAVYYCAAGWDYWGQGLTVTVSSASTK GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHK PSNTKVDKRVEPKSC
SEQ ID NO:54	DLL3#6 scFab	DIQLTQSPSFLSTSVGDRVITITCRASQDISNYFAWYQQKPGK APKLLIYAASLTQSGVPSRFSGGSGTEFTLTITSSLPEDFAT YYCQQLNSYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLQESGPGLVKPSSETLSLTCTVSGGSISSYFWSWIRQPA GKGLEWIGRIYTSSTNYNPSLNSRLTMSVDTSKNQFSLKLS SVTAADTAVYYCARRGDWGGFDIWGQGTVVTVSSASTKGP SVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPS NTKVDKRVEPKSC
SEQ ID NO:55	CD3#1 LCCDR1	RSSTGAVTTSNYAN
SEQ ID NO:56	CD3#1 LCCDR2	GTNKRAP
SEQ ID NO:57	CD3#1 LCCDR3	ALWYSNLWV
SEQ ID NO:58	CD3#1 HCCDR1	GFTFNTYAMN
SEQ ID NO:59	CD3#1 HCCDR2	RIRSKYNNYATYYADSVKD
SEQ ID NO:60	CD3#1 HCCDR3	HGNFGNSYVSWFAY
SEQ ID NO:61	CD3#2 LCCDR1	RSSTGAVTTSNYAN
SEQ ID NO:62	CD3#2 LCCDR2	GTNKRAP

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:63	CD3#2 LCCDR3	ALWYSNLWV
SEQ ID NO:64	CD3#2 HCCDR1	GFTFNTYAMN
SEQ ID NO:65	CD3#2 HCCDR2	RIRSKYINYATYYADSVKD
SEQ ID NO:66	CD3#2 HCCDR3	HGNFGNSYVSWFAY
SEQ ID NO:67	CD3#1 VL	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVL
SEQ ID NO:68	CD3#1 VH	EVQLVESGGGLVQPGGSLKLSAASGFTFNTYAMNWVRQA PGKGLEWVARIRSKYNNTYATYYADSVKDRFTISRDDSKNTA YLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFAYWGQGTL VTVSA
SEQ ID NO:69	CD3#2 VL	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVL
SEQ ID NO:70	CD3#2 VH	EVQLVESGGGLVQPGGSLKLSAASGFTFNTYAMNWVRQA PGKGLEWVARIRSKYINYATYYADSVKDRFTISRDDSKNTA YLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFAYWGQGTL VTVSA
SEQ ID NO:71	CD3#1 scFab	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVKVAWKADGSPVNTGVET TTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTV EKTVAPECSGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNTYAM NWVRQAPGKGLEWVARIRSKYNNTYATYYADSVKDRFTISR DDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFA YWGQGTLVTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT

SEQ ID Number	Brief description of sequence	Sequence
		VPSSSLGTQTYICNVNHNKPSNTKVDKRVEPKSC
SEQ ID NO:72	CD3#2 scFab	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVKVAWKADGSPVNTGVET TTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTV EKTVAPECSGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAM NWVRQAPGKGLEWVARIRSKYINYATYYADSVKDRFTISR DSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFAY WGQGTLLTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVK DYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP SSSLGTQTYICNVNHNKPSNTKVDKRVEPKSC
SEQ ID NO:73	DLL3#1 chain	EIVLTQSPGTLSPGERATLSCRASQSVSSNFLVWYQQKPG QAPRPLIYGASTRASGIPDRFSGSGSGADFTLTISRLEPEDFAL YYCQQYGDSPYTFGQGTTLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVCKASGNTFTNYMHVVR QAPGPGLEWMGIIDPSVGSKSYAQKFLGRVTIARDTSTSTVF LDLYSLRSEDYAVYFCARAGKRFGEYFDYWGQGTLLTVSS ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNV NHNKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFP PKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKV NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLW CLVKGFIYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS KLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:74	DLL3#2 chain	DIQMTQSPSAMSASVGDRVTITCRASQGISNYLAWFQQKPG KVPEPLIYAASSLQSGVPSRFSGSGSVTEFTLTISLQPEDFAT YYCLQHNSSPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSG

SEQ ID Number	Brief description of sequence	Sequence
		TASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS KDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHVVR QAPGQGLEWMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTV YMELSSLRSEDNAVYYCARGEAVGGNYYYGMDVWGQGT TVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGT QTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAG GPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMT KNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS LSLSPG
SEQ ID NO:75	DLL3#3 chain	DIQMTQSPSAMSASVGDRVTITCRASQGISNYLVWFQQKPG KAPKRLIYAVSSLYSGVPSRFSGSGSGTEFTLTISLQPEDFAT YYCLQHDSYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVSCKASGYTFTSYVHVVRQ APGQGLEWMVIINPGGGTTSYAQKFLGRVTMTRDTSTNTVY MELKSLRSEDNAVYYCARGEAVTGNFYFYGMDVWGQGT VTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPV TVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQ TYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGG PSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYV DGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPG

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:76	DLL3#4 chain	DIVLTQSPASLAVSLGQRATISCRASKSVSSFGYSFMHWYQQ KPGQPPKLLIYLASNLESGVPARFSGSGSGTDFTLNIHPVEEE DAATYYCQHSRELPWTFGGGTKLEIKRTVAAPSVFIFPPSDE QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESV TEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSQVQLQQSGTELVKPGASVKLSCKASVYTFSTSYFMY WVKQRPGHGLEWIGEISPTNGNSNLNERFKNKATLTVDKSS STAYMQLSSLTSEDSAVYYCTRGGDGYLDYWGQGTTTLTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICN VNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFL FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE VHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVS LWCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFF LYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSP G
SEQ ID NO:77	DLL3#5 chain	DIQMTQSPSSLSASVGDRVTVCQASQDISNYLNWYQQKPG KAPKLLIYDASNLETGVPSRFSGSGSGTDFTFTISLQPEDIA YYCQYDNLPTWTFGGGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQ DSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTK SFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGG GGSQVQLVQSGAEVKKPGASVKVSCKASGFTFTGYIHWV RQAPGQGLEWMGWINPNSGGTNYAQKFQGRVTMTRDSSIN TAFMELSRLTSDDTAVYYCAAGWDYWGQGLTVTVSSASTK GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHK PSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPK KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA LPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLV

SEQ ID Number	Brief description of sequence	Sequence
		KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:78	DLL3#6 chain	DIQLTQSPSFLSTSVGDRVITITCRASQDISNYFAWYQQKPGK APKLLIYAASSTLQSGVPSRFSGGSGTEFTLTISLQPEDFAT YYCQQLNSYPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLQESGPGLVKPSSETLSLTCTVSGGSISSYFWSWIRQPA GKGLEWIGRIYTSNSTYNPSLNSRLTMSVDTSKNQFSLKLS SVTAADTAVYYCARRGDWGGFDIWGQGTVVTVSSASTKGP SVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPS NTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKD TLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLVKG FYPSDIAVEWESNGQPENNYKTTTPVLDSDGSSFLYSKLTVD KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:79	CD3#1 chain	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVKVAWKADGSPVNTGVET TTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTV EKTVAPAECSGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAM NWVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRTISR DDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFA YWGQGTLLTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDP EVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ

SEQ ID Number	Brief description of sequence	Sequence
		DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFCFSVMHEALHNRF
SEQ ID NO:80	CD3#2 chain	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEPGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQPEDEAEYFCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVKVAWKADGSPVNTGVETTPPSKQSNKYAASSYLSLTPEQWKSRSYSCQVTHEGSTVEKTVAPAECSSGGGSEKSSSGSGSESKSTEGKSSSGSGSESKSTGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNTYAMNWVRQAPGKGLEWVARIRSKYINYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLVTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPTSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAEEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFCFSVMHEALHNRF
SEQ ID NO:81	Fc domain* (IgG1)	<u>DKTHTCPPCP</u> APPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:82	Fc W domain (IgG1, LALA)	<u>DKTHTCPPCP</u> APAAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFCFS

SEQ ID Number	Brief description of sequence	Sequence
		CSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:83	Fc SAV domain (IgG1, RF/LALA)	<u>DKTHTCPPCPAPEA</u> AGGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQ PREPQVYTLPPSREEMTKNQVSLSCAVKGFYPSDIAVEWESN GQPENNYKTTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSC SVMHEALHNRFTQKSLSLSPG
SEQ ID NO:84	Fc domain (IgG4Pro)	<u>ESKYGPPCPPCPAPEFL</u> GGPSVFLFPPKPKDTLMISRTPEVTC VVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQ PREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESN GQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSC SVMHEALHNHYTQKSLSLSLG
SEQ ID NO:85	Fc W domain (IgG4Pro)	<u>ESKYGPPCPPCPAPEFL</u> GGPSVFLFPPKPKDTLMISRTPEVTC VVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQ PREPQVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWES NGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFS CSVMHEALHNHYTQKSLSLSLG
SEQ ID NO:86	Fc SAV domain (IgG4Pro, RF)	<u>ESKYGPPCPPCPAPEFL</u> GGPSVFLFPPKPKDTLMISRTPEVTC VVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQ PREPQVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWES NGQPENNYKTTPPVLDSDGSFFLVSRLLTVDKSRWQEGNVFS CSVMHEALHNRFTQKSLSLSLG
SEQ ID NO:87	constant region of a kappa light chain	RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQW KVDNALQSGNSQESVTEQDSKDSTYSLSSTLTLSKADYEKH KQYACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO:88	constant region of a lambda light chain	GQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVKVA WKADGSPVNTGVETTTTPSKQSNKYAASSYLSLTPEQWKS HRSYSCQVTHEGSTVEKTVAPAEC

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:89	Linker	GGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGS
SEQ ID NO:90	Linker	GGGSGGGGSGGSGGSGGGGS
SEQ ID NO:91	Linker	GGGSGGGGSGGGGSGGGGSGGGGS
SEQ ID NO:92	Linker	GGGSGGGGGSGGGGGSGGGGSGGGGS
SEQ ID NO:93	Linker	GGGSGGGGSGGGSGGGSGGGGSGGGGSGGGGS
SEQ ID NO:94	Linker	GGGSGGGGSGGGSGGGSGGGGSGGGGSGGGGSGGGGS
SEQ ID NO:95	Linker	GGGSGGGGSGGGSGGGSGGGGSGGGGSGGGGSGGGSGG GGS
SEQ ID NO:96	CD3#3 LCCDR1	RSSTGAVTTSNYAN
SEQ ID NO:97	CD3#3 LCCDR2	GTNKRAP
SEQ ID NO:98	CD3#3 LCCDR3	ALWYSNLWV
SEQ ID NO:99	CD3#3 HCCDR1	GFTFNTYAMN
SEQ ID NO:100	CD3#3 HCCDR2	RIRSKYANYATYYADSVKD
SEQ ID NO:101	CD3#3 HCCDR3	HGNFGNSYVSWFAY
SEQ ID NO:102	CD3#3 VL	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVL
SEQ ID NO:103	CD3#3 HL	EVQLVESGGGLVQPGGSLKLSAASGFTFNTYAMNWVRQA PGKGLEWVARIRSKYANYATYYADSVKDRFTISRDDSKNTA YLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFAYWGQGTL VTVSA

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:104	CD3#3 scFab	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVKVAWKADGSPVNTGVET TTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTV EKTVAECSGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNTYAM NWVRQAPGKGLEWVARIRSKYANYATYYADSVKDRFTISR DDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFA YWGQGTLLTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSC
SEQ ID NO:105	CD3#3 chain	EAVVTQEPSLTVSPGGTVTLTCRSSTGAVTTSNYANWVQEK PGQLPRGLIGGTNKRAPWVPARFSGSLLGGKAALTLGAQP EDEAEYFCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVKVAWKADGSPVNTGVET TTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTV EKTVAECSGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNTYAM NWVRQAPGKGLEWVARIRSKYANYATYYADSVKDRFTISR DDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYVSWFA YWGQGTLLTVSAASTKGPSVFPLAPSSKSTSGGTAALGCLV KDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP EVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPP PSREEMTKNQVSLTSCAVKGFPYPSDIAVEWESNGQPENNYKT TPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSCSVMHEALHN RFTQKSLSLSPG
SEQ ID NO:133	DLL3#7 LCCDR1	RASQSVNSNFLA
SEQ ID	DLL3#7	GTSSRAT

SEQ ID Number	Brief description of sequence	Sequence
NO:134	LCCDR2	
SEQ ID NO:135	DLL3#7 LCCDR3	QQYGSSPWT
SEQ ID NO:136	DLL3#7 HCCDR1	GFTFSSYGMF
SEQ ID NO:137	DLL3#7 HCCDR2	VIWLDGDDDEDYVDSVKG
SEQ ID NO:138	DLL3#7 HCCDR3	VLDY
SEQ ID NO:139	DLL3#8 LCCDR1	KSSQSVLDTSNNKNYLV
SEQ ID NO:140	DLL3#8 LCCDR2	WASTRES
SEQ ID NO:141	DLL3#8 LCCDR3	QHYYNSPYT
SEQ ID NO:142	DLL3#8 HCCDR1	GYTFTDYMH
SEQ ID NO:143	DLL3#8 HCCDR2	WINPNSGGTNYEQKFQG
SEQ ID NO:144	DLL3#8 HCCDR3	DAVVIPMDY
SEQ ID NO:145	DLL3#9 LCCDR1	RASQISRSYLA
SEQ ID NO:146	DLL3#9 LCCDR2	GASSRAT
SEQ ID NO:147	DLL3#9 LCCDR3	QQYGTSPIT
SEQ ID NO:148	DLL3#9 HCCDR1	GGSISSYYWS
SEQ ID NO:149	DLL3#9 HCCDR2	YRYYSNGNTNYPNPSLKS
SEQ ID NO:150	DLL3#9 HCCDR3	IGVAGFYFDY

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:151	DLL3#10 LCCDR1	RASQSLNSIFLA
SEQ ID NO:152	DLL3#10 LCCDR2	GASSRAT
SEQ ID NO:153	DLL3#10 LCCDR3	QQYGGSMNT
SEQ ID NO:154	DLL3#10 HCCDR1	GYTFTGYMH
SEQ ID NO:155	DLL3#10 HCCDR2	WINPNSGGTIFAQRFQG
SEQ ID NO:156	DLL3#10 HCCDR3	DFGDTVGNAFDI
SEQ ID NO:157	DLL3#11 LCCDR1	SASSSVTYIH
SEQ ID NO:158	DLL3#11 LCCDR2	RTSYLAS
SEQ ID NO:159	DLL3#11 LCCDR3	QQRSSYPRT
SEQ ID NO:160	DLL3#11 HCCDR1	GYAFSDYWIT
SEQ ID NO:161	DLL3#11 HCCDR2	DIYPGSGSTKSSEKFKN
SEQ ID NO:162	DLL3#11 HCCDR3	LYYYGSHYLDI
SEQ ID NO:163	DLL3#12 LCCDR1	SASSSVTYIH
SEQ ID NO:164	DLL3#12 LCCDR2	RTSYLAS
SEQ ID NO:165	DLL3#12 LCCDR3	QQRSSYPRT
SEQ ID NO:166	DLL3#12 HCCDR1	GYAFSDYWIT
SEQ ID	DLL3#12	DIYPGSGSTKSSEKFKN

SEQ ID Number	Brief description of sequence	Sequence
NO:167	HCCDR2	
SEQ ID NO:168	DLL3#12 HCCDR3	LYYYGSYYLDT
SEQ ID NO:169	DLL3#13 LCCDR1	RSSQSIVHSNGNTYLE
SEQ ID NO:170	DLL3#13 LCCDR2	KVSNRFS
SEQ ID NO:171	DLL3#13 LCCDR3	FQGSHVPYT
SEQ ID NO:172	DLL3#13 HCCDR1	GYTFTNYGVT
SEQ ID NO:173	DLL3#13 HCCDR2	WINTYSGAPTYADDFNG
SEQ ID NO:174	DLL3#13 HCCDR3	LDDYDLYYFDY
SEQ ID NO:175	DLL3#14 LCCDR1	KASQSVVDYDGDSYMN
SEQ ID NO:176	DLL3#14 LCCDR2	AASTLES
SEQ ID NO:177	DLL3#14 LCCDR3	QQSDEDPWT
SEQ ID NO:178	DLL3#14 HCCDR1	GYTFTDYIYH
SEQ ID NO:179	DLL3#14 HCCDR2	YIYPGNSYTAYNQKFKD
SEQ ID NO:180	DLL3#14 HCCDR3	SGGSAMDY
SEQ ID NO:181	DLL3#15 LCCDR1	KASQSVVDYDGDSYLN
SEQ ID NO:182	DLL3#15 LCCDR2	AASNLES
SEQ ID NO:183	DLL3#15 LCCDR3	QQSSEDPRT

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:184	DLL3#15 HCCDR1	GYTFTNYGMN
SEQ ID NO:185	DLL3#15 HCCDR2	WINTYTGEPTYADDFKG
SEQ ID NO:186	DLL3#15 HCCDR3	FHFSSNGDAMDN
SEQ ID NO:187	DLL3#16 LCCDR1	RASQSVSDWLA
SEQ ID NO:188	DLL3#16 LCCDR2	RASSLES
SEQ ID NO:189	DLL3#16 LCCDR3	QLYNSYSPT
SEQ ID NO:190	DLL3#16 HCCDR1	GFTFSSYWMT
SEQ ID NO:191	DLL3#16 HCCDR2	NIKEDGSEKYYVDSVKG
SEQ ID NO:192	DLL3#16 HCCDR3	DWGYFDY
SEQ ID NO:193	DLL3#17 LCCDR1	RASENIYYSLA
SEQ ID NO:194	DLL3#17 LCCDR2	NTNSLED
SEQ ID NO:195	DLL3#17 LCCDR3	KQAYDFPLT
SEQ ID NO:196	DLL3#17 HCCDR1	GYTFISYYIH
SEQ ID NO:197	DLL3#17 HCCDR2	WIYPGDGSTNNNEKFKG
SEQ ID NO:198	DLL3#17 HCCDR3	GEGNAMDD
SEQ ID NO:199	DLL3#18 LCCDR1	RASENIYYSLA
SEQ ID	DLL3#18	NANSLED

SEQ ID Number	Brief description of sequence	Sequence
NO:200	LCCDR2	
SEQ ID NO:201	DLL3#18 LCCDR3	KQAYDVPLT
SEQ ID NO:202	DLL3#18 HCCDR1	GYTFTAYFIH
SEQ ID NO:203	DLL3#18 HCCDR2	YIDPFNDDTNYNVKFKG
SEQ ID NO:204	DLL3#18 HCCDR3	GTSATLDY
SEQ ID NO:205	DLL3#7 VL	EIVLTQSPDTLSLSPGERATLSCRASQSVNSNFLAWYQQKPGQTPRLLIFGTSSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQYGSSPWTFGQGTKVEIR
SEQ ID NO:206	DLL3#7 VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMFWVRQAPGKGLEWVAWIWLDGDDDEDYVDSVKGRFTISRDDSKNTLYLQMNSLRVDDTAIYYCARVLDYWGQGLTVTVSS
SEQ ID NO:207	DLL3#8 VL	DIVMAQSPDSLAVSLGERATINCKSSQSVLDTSNNKNYLWVWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQHYYNSPYTFGQGTKLEIK
SEQ ID NO:208	DLL3#8 VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTDYYMHWVRQAPGQGLEWMGWINPNSGGTNYEQKFQGRVTMTRDTSISTAYMELNRLRSDDTAIYYCTRDAVVIPMDYWGQGLTVTVSS
SEQ ID NO:209	DLL3#9 VL	EIVLTQSPGTLSLSPGERATLSCRASQSISRSLAWYQQKPGQAPRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQYGTSPITFGQGTRLEIK
SEQ ID NO:210	DLL3#9 VH	QVHLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWIRQTPGKGLDWIGYRYYSNTNYPNPSLKSRTVSLDMSNNQFSLKLSVTAADTAIYYCASIGVAGFYFDYWGQGLTVTVSS
SEQ ID NO:211	DLL3#10 VL	EIVLTQSPGTLSLSPGERATLSCRASQSLNSIFLAWYQQKPGQAPWLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYFCQQYGGSMNTFGQGTKLEIK
SEQ ID NO:212	DLL3#10 VH	QVQLVQSGAEVKKPGASVKVSKASGYTFTGYMHWVRLAPGQGLEWMGWINPNSGGTIFAQRFQGRVTMTRDTSISTVYMDLNRLRSDDTAIYYCARDFGDTVGNAFDIWGQGTMTVT

SEQ ID Number	Brief description of sequence	Sequence
		SS
SEQ ID NO:213	DLL3#11 VL	QIVLTQSPAIMASASPGEKVTITCSASSSVTYIHWFQQNPGTSP KLWIYRTSYLASGVPARFSGSGSGTSYSLTISRMEAEDAATY YCQQRSSYPRTFGGGGTKLEIK
SEQ ID NO:214	DLL3#11 VH	QVQLQQPGAELVQPGSSVKMSCKASGYAFSDYWITWVKQR PGQGLEWIGDIYPGSGSTKSSEKFKNKATLTADTSSSKAYIQ FSSLTPEDSAVYYCVSLYYYGSHYLDTWGQGTTTLTVSS
SEQ ID NO:215	DLL3#12 VL	QVVLTQSPAIMASASPGEKVTITCSASSSVTYIHWFQQNPGTSP KLWIYRTSYLASGVPARFSGSGSGTSYSLTISRMEAEDAATY YCQQRSSYPRTFGGGGTKLEIK
SEQ ID NO:216	DLL3#12 VH	QVQLQQPGAELVQPGSSVKMSCKASGYAFSDYWITWVKQR PGQGLEWIGDIYPGSGSTKSSEKFKNRATLTADTSSSTAYIQF SSLTPEDSAVYYCVSLYYYGSYYLDTWGQGTTTLTVSS
SEQ ID NO:217	DLL3#13 VL	DVLMTQTPLSLPVS LGDQAAISCRSSQSIVHSNGNTYLEWYL QKPGQSPKVLIIYKVS NRFSGV PDRFSGSGSGTDFTLKISRVE AEDLGVYYCFQGSHPYTFGGGTKLEIK
SEQ ID NO:218	DLL3#13 VH	QIQLVQSGPELKKPGETVKISCKASGYTFTNYGVTWVKQAP GKGLKWMGWINTYSGAPTYADDFNGRFALSLETSASTAYL QINN LKNEDTATYFCARLDDYDLYYFDYWGQGTALT TVSS
SEQ ID NO:219	DLL3#14 VL	DIVLTQSPASLSVSLGQRATISCKASQSVDDYDGDSYMNWYQ QKPGQPPKLLIYAAS TLESGIPARFSGSGSGTDFTLNIHPVEEE DAATYYCQQSDEDPWTFGGGTKLEIK
SEQ ID NO:220	DLL3#14 VH	QIQLVQSGPELVKPGVKISCKASGYTFTDYIHWKQRPQG GLEWIGYIYPGNSYTAYNQKFKDKATLTADNPSSTAYMQLS SLTSEDSAVYFCARSGGSAMDYWGQGTSTVTVSS
SEQ ID NO:221	DLL3#15 VL	DIVLTQSPASLAVSLGQRATISCKASQSVDDYDGDSYLNWYQ QKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNIHPVEE EDAATYYCQQSSEDPRTFGGGGTKLEIK
SEQ ID NO:222	DLL3#15 VH	QIQLVQSGPELKKPGETVKISCKASGYTFTNYGMNWVKQAP GKGLKWMGWINTYTGEPTYADDFKGRFAFSLETSASTAYL QINN LKNEDMATYFCTKFHFSSNGDAMD NWGQGTSTVTVSS
SEQ ID NO:223	DLL3#16 VL	DIQMTQSPSTLSASVGDRTITCRASQSVSDWLAWYQQKPG KAPKFLIYRASSLES GVP SRFS GSGSGTEFTLTISSLQPADFAT

SEQ ID Number	Brief description of sequence	Sequence
		YYCQLYNSYSPTFGQGTKVEIK
SEQ ID NO:224	DLL3#16 VH	EVHLVESGGGLVQPGGSLRLSCAASGFTSSYWMTWVRQA PGKGLEWVANIKEDGSEKYYVDSVKGRFTISRDNAKNSLYL QMNSLRAEDTALYYCARDWGYFDYWGQGTLLTVSS
SEQ ID NO:225	DLL3#17 VL	DIQMTQSPASLAASVGETVTITCRASENIYYSLAWYQQKQG KSPQLLIYNTNSLEDGVPSRFSGSGSGTQYSMKINSMQPEDT ATYFCKQAYDFPLTFGAGTKLELK
SEQ ID NO:226	DLL3#17 VH	QIQLQQSGPEVVKPGASVKISCKASGYTFISYYIHWVKQRPG QGLEWIGWIYPGDGSTNNNEKFKGKTTLTADKSSSTAYMLL SSLTSEDSAVYFCARGEGNAMDDWGQGTSTVTVSS
SEQ ID NO:227	DLL3#18 VL	DIQMTQSPASLAASVGETVTITCRASENIYYSLAWYQQKQG KSPQLLIYNANSLEDGVPSRFSGSGSGTQYSMKINNMQPEDT ATYFCKQAYDVPLTFGAGTKLELK
SEQ ID NO:228	DLL3#18 VH	QVQLQQSGPDLVKPGASVKMSCEASGYTFTAYFIHWVKQK PGQGLEWIGYIDPFNDNTNYNVKFKGKATLTSDTSSSIAYME LSSLTSEDSSFYYCARGTSATLDYWGHGTTTLTVSS
SEQ ID NO:229	DLL3#7 scFab	EIVLTQSPDTLSLSPGERATLSCRASQSVNSNFLAWYQQKPG QTPRLIFGTSSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAV YYCQQYGSSPWTFTGQGTKVEIRRTVAAPSVFIFPPSDEQLKS GTASVVCLLNMFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVESGGGVVQPGRLRLSCAASGFTSSYGMFWVRQ APGKGLEWVAWIWLDGDEEDYVDSVKGRFTISRDDSKNTL YLQMNSLRVDDTAIYYCARVLDYWGQGTLLTVSSASTKGP SVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPS NTKVDKRVKPKSC
SEQ ID NO:230	DLL3#8 scFab	DIVMAQSPDSLAVSLGERATINCKSSQSVLDTSNNKNYLVW YQQKPGQPPKLLIYWASTRESGVDPDRFSGSGSGTDFTLTISL QAEDVAVYYCQHYYNSPYTFGQGTKLEIKRTVAAPSVFIFPP SDEQLKSGTASVVCLLNMFYPREAKVQWKVDNALQSGNSQ ESVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGL

SEQ ID Number	Brief description of sequence	Sequence
		SSPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSES KSTGGGGSQVQLVQSGAEVKKPGASVKVSCKASGYTFTDY YMHWVRQAPGQGLEWMGWINPNSGGTNYEQKFQGRVTM TRDTSISTAYMELNRLRSDDTAVYYCTRDAVIPMDYWGQ GTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSL GTQTYICNVNHKPSNTKVDKRVEPKSC
SEQ ID NO:231	DLL3#9 scFab	EIVLTQSPGTLSLSPGERATLSCRASQSISRSLAWYQQKPGQ APRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVY YCQQYGTSPITFGQGTRLEIKRTVAAPSVFIFPPSDEQLKSGT ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR GECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQ VHLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIRQTPGK GLDWIGYRYYSNTNYPNPSLKSRTISLDMSNNQFSLKLSS VTAADTAIYYCASIGVAGFYFDYWGQGTTLVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSG VHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSN TKVDKRVEPKSC
SEQ ID NO:232	DLL3#10 scFab	EIVLTQSPGTLSLSPGERATLSCRASQSLNSIFLAWYQQKPGQ APWLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAV YFCQQYGGSMNTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVSCKASGYTFTGYMHWVR LAPGQGLEWMGWINPNSGGTIFAQRFQGRVTMTRDTSISTV YMDLNRLRSDDTAVYYCARDVFGDTVGNAFDIWGQGMVT VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTV SWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTY ICNVNHKPSNTKVDKRVEPKSC
SEQ ID NO:233	DLL3#11 scFab	QIVLTQSPAISASPGKEKVTITCSASSSVTYIHWFQQNPGTSP KLWIYRTSYLASGVPARFSGSGSGTSYSLTISRMEAEDAATY

SEQ ID Number	Brief description of sequence	Sequence
		YCQQRSSYPRTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGT ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR GECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQ VQLQQPGAELVQPGSSVKMSCKASGYAFSDYWITWVKQRP GQGLEWIGDIYPGSGSTKSSEKFKNKATLTADTSSSKAYIQF SSLTPEDSAVYYCVSLYYYGSHYLDTWGQGTTTLTVSSASTK GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHK PSNTKVDKRVEPKSC
SEQ ID NO:234	DLL3#12 scFab	QVVLTSQPAIMASAPGEKVTITCSASSSVTYIHWFQQNPGTSP KLWIYRTSYLASGVPARFSGSGSGTSYSLTISRMEAEDAATY YCQQRSSYPRTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGT ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR GECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQ VQLQQPGAELVQPGSSVKMSCKASGYAFSDYWITWVKQRP GQGLEWIGDIYPGSGSTKSSEKFKNRATLTADTSSSTAYIQFS SLTPEDSAVYYCVSLYYYGSYYLDTWGQGTTTLTVSSASTKG PSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHK SNTKVDKRVEPKSC
SEQ ID NO:235	DLL3#13 scFab	DVLMTQTPLSLPVSLGDQAAISCRSSQSIVHSNGNTYLEWYL QKPGQSPKVLIIYKVSNRFSGVDPDRFSGSGSGTDFTLKISRVE AEDLGVIYCFQGSHVPYTFGGGTKLEIKRTVAAPSVFIFPPS DEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQE SVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLS SPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSES KSTGGGGSQIQLVQSGPELKKPGETVKISCKASGYTFTNYGV TWVKQAPGKGLKWMGWINTYSGPTYADDFNGRFALSLE TSASTAYLQINNLKNEDTATYFCARLDDYDLYYFDYWGQG TALTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLG

SEQ ID Number	Brief description of sequence	Sequence
		TQTYICNVNHHKPSNTKVDKRVEPKSC
SEQ ID NO:236	DLL3#14 scFab	DIVLTQSPASLSVSLGQRATISCKASQSVDDYDGDSYMNWYQ QKPGQPPKLLIYAASLTESGIPARFSGSGSGTDFTLNHPVEEE DAATYYCQQSDEDPWTFGGGTKLEIKRTVAAPSVFIFPPSDE QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESV TEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSQIQLQQSGPELVKPGVKISCKASGYTFTDYYIHWM KQRPQGGLWIGYIYPGNSYTAYNQKFKDKATLTADNPSST AYMQLSSLTSEDSAVYFCARSGGSAMDYWGQGTSTVTVSSA STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTWWSN GALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNV NHHKPSNTKVDKRVEPKSC
SEQ ID NO:237	DLL3#15 scFab	DIVLTQSPASLAVSLGQRATISCKASQSVDDYDGDSYLNWYQ QKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNHPVEE EDAATYYCQQSSEDPRTFGGGTKLEIKRTVAAPSVFIFPPSDE QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESV TEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSQIQLVQSGPELKKPGETVKISCKASGYTFTNYGMN WVKQAPGKGLKWMGWINTYTGEPTYADDFKGRFAFSLETS ASTAYLQINNLKNEIDMATYFCTKHFHSSNGDAMDNWGQGT SVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTWWSNGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGT QTYICNVNHHKPSNTKVDKRVEPKSC
SEQ ID NO:238	DLL3#16 scFab	DIQMTQSPSTLSASVGRVTITCRASQSVSDWLAWYQQKPG KAPKFLIYRASSLESGVPSRFSGSGSGTEFTLTISSLQPADFAT YYCQLYNSYSPTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDYSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SEVHLVESGGGLVQPGGSLRLSCAASGFTSSYWMWVRQ APGKGLEWVANIKEDGSEKYYVDSVKGRFTISRDNKNSLY

SEQ ID Number	Brief description of sequence	Sequence
		LQMNSLRAEDTALYYCARDWGYFDYWGQGTLLVTVSSAST KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVN HKPSNTKVDKRVEPKSC
SEQ ID NO:239	DLL3#17 scFab	DIQMTQSPASLAASVGETVTITCRASENIYYSLAWYQQKQG KSPQLLIYNTNSLEDGVPSRFSGSGSGTQYSMKINSMQPEDT ATYFCKQAYDFPLTFGAGTKLELKRTVAAPSVFIFPPSDEQL KSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTE QDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTG GGGSQIQLQQSGPEVVKPGASVKISCKASGYTFISYYIHWVK QRPQGQLEWIGWIYPGDGSTNNNEKFKGKTTLTADKSSSTA YMLLSSLTSEDSAVYFCARGEGNAMDDWGQGTSVTVSSAS TKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVN HKPSNTKVDKRVEPKSC
SEQ ID NO:240	DLL3#18 scFab	DIQMTQSPASLAASVGETVTITCRASENIYYSLAWYQQKQG KSPQLLIYNANSLEDGVPSRFSGSGSGTQYSMKINNMQPEDT ATYFCKQAYDVPLTFGAGTKLELKRTVAAPSVFIFPPSDEQL KSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTE QDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTG GGGSQVQLQQSGPDLVKPGASVKMSCEASGYTFTAYFIHW VKQKPGQGLEWIGYIDPFNDNTNINVKFKGKATLTSDTSSSI AYMELSSLTSEDSSFYYCARGTSATLDYWGHGTTTLTVSSAS TKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVN HKPSNTKVDKRVEPKSC
SEQ ID NO:241	DLL3#7 chain	EIVLTQSPDTLSLSPGERATLSCRASQSVNSNFLAWYQQKPG QTPRLIFGTSSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAV YYCQQYGSSPWTFTGQGTKEIRRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF

SEQ ID Number	Brief description of sequence	Sequence
		NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMFWVRQ APGKGLEWVAVIWLDGDDDEDYVDSVKGRFTISRDDSKNTL YLQMNSLRVDDTAIYYCARVLDYWGQGTLVTVSSASTKGP SVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPS NTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKD TLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLVKG FYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVD KSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:242	DLL3#8 chain	DIVMAQSPDSLAVSLGERATINCKSSQSVLDTSNNKNYLWV YQQKPGQPPKLLIYWASTRESGVDPDRFSGSGSGTDFTLTISL QAEDVAVYYCQHYYNSPYTFGQGTKLEIKRTVAAPSVFIFPP SDEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQ ESVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGL SSPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSES KSTGGGGSQVQLVQSGAEVKKPGASVKVSCKASGYTFTDY YMHWRQAPGQGLEWMGWINPNSGGTNYEQKFQGRVTM TRDTSISTAYMELNRLRSDDTAVYYCTRDVAVIPMDYWGQ GTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSL GTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEA AGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLN GKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREE MTKNQVSLWCLVKGFYPSPDIAVEWESNGQPENNYKTTTPV LDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPG
SEQ ID NO:243	DLL3#9 chain	EIVLTQSPGTLSPGERATLSCRASQISRSYLAWYQQKPGQ APRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVY YCQQYGTSPITFGQGRLEIKRTVAAPSVFIFPPSDEQLKSGT

SEQ ID Number	Brief description of sequence	Sequence
		ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR GECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQ VHLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIRQTPGK GLDWIGYRYYSNTNYPNPSLKSRTISLDMNNQFSLKLSS VTAADTAIYYCASIGVAGFYFDYWGQGLTVTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSG VHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSN TKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDT LMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTK PREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDK SRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:244	DLL3#10 chain	EIVLTQSPGTLSPGERATLSCRASQSLNSIFLAWYQQKPGQ APWLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAV YFCQQYGGSMNTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SQVQLVQSGAEVKKPGASVKVSCKASGYTFTGYMHVVR LAPGQGLEWMGWINPNSGGTIFAQRFQGRVTMTRDTSISTV YMDLNRRLRSDDTAVYYCARDFTVGNAFDIWGQGTMTV VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTY ICNVNHKPSNTKVDRVEPKSCDKTHTCPPCPAPEAAGGPS VFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVD GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSL SPG
SEQ ID	DLL3#11	QIVLTQSPAISASPGEKVTITCSASSSVTYIHWFQQNPGTSP

SEQ ID Number	Brief description of sequence	Sequence
NO:245	chain	KLWIYRTSYLASGVPARFSGSGSGTSYSLTISRMEAEDAATY YCQQRSSYPRTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGT ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR GECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQ VQLQQPGAELVQPGSSVKMSCKASGYAFSDYWITWVKQRP GQGLEWIGDIYPGSGSTKSSEKFKNKATLTADTSSSKAYIQF SSLTPEDSAVYYCVSLYYYGSHYLDTWGQGTTTLTVSSASTK GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHK PSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKP KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA LPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLV KGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:246	DLL3#12 chain	QVVLTSQPAIMASAPGEKVTITCSASSSVTYIHWFQQNPGTSP KLWIYRTSYLASGVPARFSGSGSGTSYSLTISRMEAEDAATY YCQQRSSYPRTFGGGTKLEIKRTVAAPSVFIFPPSDEQLKSGT ASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSK DSTYSLSSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNR GECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGGSQ VQLQQPGAELVQPGSSVKMSCKASGYAFSDYWITWVKQRP GQGLEWIGDIYPGSGSTKSSEKFKNRATLTADTSSSTAYIQFS SLTPEDSAVYYCVSLYYYGSYYLDTWGQGTTTLTVSSASTKG PSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKP SNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKP DTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWCLVK GFYPSPDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTV DKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG

SEQ ID Number	Brief description of sequence	Sequence
SEQ ID NO:247	DLL3#13 chain	DVLMTQTPLSLPVSLGDQAAISCRSSQSIVHSNGNTYLEWYL QKPGQSPKVLIIYKVSNRFSGVPDRFSGSGSGTDFTLKISRVE AEDLGVIYCFQGSHVPYTFGGGTKLEIKRTVAAPSVFIFPPS DEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQE SVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLS SPVTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSES KSTGGGGSQIQLVQSGPELKKPGETVKISCKASGYTFTNYGV TWVKQAPGKGLKWMGWINTYSGAPTYADDFNGRFALSLE TSASTAYLQINNLIKNETATYFCARLDDYDLYYFDYWGQG TALTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLG TQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAA GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNG KEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM TKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQK SLSLSPG
SEQ ID NO:248	DLL3#14 chain	DIVLTQSPASLSVSLGQRATISCKASQSVDDYDGD SYMNWYQ QKPGQPPKLLIYAASLTESGIPARFSGSGSGTDFTLNHPVEEE DAATYYCQQSDEDPWTFGGGTKLEIKRTVAAPSVFIFPPSDE QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESV TEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSQIQLQQSGPELVKPGVKISCKASGYTFTDYIHW KQRPQGQLEWIGYIYPGNSYTAYNQKFKDKATLTADNPSS AYMQLSSLTSEDSAVYFCARSGGSAMDYWGQGTSTVTVSSA STKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNV NHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFP PKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLW

SEQ ID Number	Brief description of sequence	Sequence
		CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS KLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:249	DLL3#15 chain	DIVLTQSPASLAVSLGQRATISCKASQSVDDYDGD SYLNWYQ QKPGQPPKLLIYAASNLESGIPARFSGSGSGTDFTLNHPVEE EDAATYYCQQSSEDPRTFGGGTKLEIKRTVAAPS VFIFPPSDE QLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESV TEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKS TGGGGSQIQLVQSGPELKKPGETVKISCKASGYTFTNYGMN WVKQAPGKGLKWMGWINTYTGEPTYADDFKGRFAFSLETS ASTAYLQINN LKNEDMATYFCTKHFHSSNGDAMD NWGQGT SVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV VTVPSSSLGT QTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAG GPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMT KNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD DGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS LSLSPG
SEQ ID NO:250	DLL3#16 chain	DIQMTQSPSTLSASVGDRVTITCRASQSVSDWLAWYQQKPG KAPKFLIYRASSLESGVPSRFSGSGSGTEFTLTISSLQPADFAT YYCQLYNSYSPTFGQGTKVEIKRTVAAPS VFIFPPSDEQLKS GTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSF NRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTGGGG SEVHLVESGGGLVQPGGSLRLSCAASGFTSSYWMTWVRQ APGKGLEWVANIKEDGSEKYYVDSVKGRFTISRDN AKNSLY LQMNSLRAEDTALYYCARDWGYFDYWGQGT LTVTVSSAST KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSV VTVPSSSLGTQTYICNVN HKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPP KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH

SEQ ID Number	Brief description of sequence	Sequence
		NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWC LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSK LTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:251	DLL3#17 chain	DIQMTQSPASLAASVGETVTITCRASENIYYSLAWYQQKQG KSPQLLIYNTNSLEDGVPSRFSGSGSGTQYSMKINSMQPEDT ATYFCKQAYDFPLTFGAGTKLELKRTVAAPSVFIFPPSDEQL KSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTE QDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTG GGGSQIQLQQSGPEVVKPGASVKISCKASGYTFISYYIHWVK QRPQGQLEWIGWIYPGDGSTNNNEKFKGKTTLTADKSSSTA YMLLSSLTSEDSAVYFCARGEGNAMDDWGQGTSVTVSSAS TKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVN HKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPP KPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVH NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWC LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSK LTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:252	DLL3#18 chain	DIQMTQSPASLAASVGETVTITCRASENIYYSLAWYQQKQG KSPQLLIYNANSLEDGVPSRFSGSGSGTQYSMKINNMQPEDT ATYFCKQAYDVPLTFGAGTKLELKRTVAAPSVFIFPPSDEQL KSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTE QDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGECGGGGSEGKSSGSGSESKSTEGKSSGSGSESKSTG GGGSQVQLQQSGPDLVKPGASVKMSCEASGYTFTAYFIHW VKQKPGQGLEWIGYIDPFNDDTNYNVKFKGKATLTSDTSSSI AYMELSSLTSEDSSFYYCARGTSATLDYWGHGTTTLTVSSAS TKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVN HKPSNTKVDKRVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPP

SEQ ID Number	Brief description of sequence	Sequence
		KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVH NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLWC LVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSK LTVDKSRWQQGNVFCSSVMHEALHNHYTQKSLSLSPG
SEQ ID NO:253	Constant region of heavy chain CH1	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNV NHKPSNTKVDKRVEPKSC
SEQ ID NO:254	Mouse IgG2a HC	AKTTAPSVYPLAPVCGDTTGSSVTLGCLVKGYFPEPVTLTW NSGSLSSGVHTFPAVLQSDLYTLSSSVTVTSSTWPSQSITCNV AHPASSTKVDDKIEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPP KIKDVLMSLSPIVTCVVVDVSEDDPDVQISWVFNNEVHTA QTQTHREDYNSTLRVVSALPIQHQDWMGKEFKCKVNNKD LPAPIERTISKPKGSVRAPQVYVLPPEEEMTKKQVTLTCMV TDFMPEDIYVEWTNNGKTELNYKNTEPVLDSDGSYFMYSK LRVEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK
SEQ ID NO:255	Mouse Kappa LC	RADAAPTVSIFPPSSEQLTSGGASVVCFLNNFYPKDINVKWK IDGSERQNGVLNSWTDQDSKDSTYSMSSTLTTLTKDEYERHN SYTCEATHKTSTSPIVKSFNRNEC

*Underlined sequence indicates hinge region

Example 3: Production of recombinant proteins

• Human DLL3-His

- 5 A cell line to produce Human DLL3-His was generated using HEK-293 cells (Thermo Fisher), the Lenti-X Lentiviral System (Clontech), and plasmid encoding Human DLL3-His (human DLL3 Accession No. Q9NYJ7 huDLL3-His: SEQ ID NO:106). For expression, cells were cultured and expanded at 37C, 5% CO₂, and shaking at 140 rpm. On Day 0 of expression, cell were pelleted and re-suspended in Expi 293 media. On day 3 of
- 10 expression the conditioned culture supernatant was harvested by pelleting the cells for 40 minutes at 4700 rpm. Protease inhibitors were added to the biomass before purification. Expression was confirmed by western blot. The conditioned culture supernatant was

adjusted with 0.5 mM TCEP, 0.02% CHAPS, 10 mM imidazole. Purification was carried out on a HisTrap Ni excel column and Buffer A: 50 mM MES, 50 mM NaCl, 0.5 mM TCEP, 0.02% CHAPS, pH 6.5. The protein of interest was eluted in Buffer A supplemented with 0.5M Imidazole, pH 8.5, using an elution gradient from 20 mM imidazole to 500 mM imidazole. The pooled fractions were dialyzed in buffer: 50 mM MES, 50 mM NaCl, 1 mM TCEP, 0.02% CHAPS, 0.2M Arginine, 3% glycerol, pH 6.5. The purified material was qualified by mass spectrometry and analytical ultra-centrifugation.

• **Cyno DLL3-His**

A cell line to produce Cyno DLL3-His was generated using HEK-293 cells (Thermo Fisher), the Lenti-X Lentiviral System (Clontech), and plasmid encoding Cyno DLL3-His (Cyno DLL3 Accession No: XM_005589196.1 (RefSeq); Cyno DLL3-His: SEQ ID NO:107). For expression, cells were cultured and expanded at 37°C, 5% CO₂, and shaking at 140 rpm. On Day 0 of expression, cells were pelleted and re-suspended in Expi 293 media. On day 3 of expression the conditioned culture supernatant was harvested by pelleting the cells for 40 minutes at 4700 rpm. Protease inhibitors were added to the biomass before purification. Expression was confirmed by western blot. The conditioned culture supernatant was adjusted with 0.5 mM TCEP, 0.02% CHAPS, 10 mM imidazole. Purification was carried out on a HisTrap Ni excel column and Buffer A: 50 mM MES, 50 mM NaCl, 0.5 mM TCEP, 0.02% CHAPS, pH 6.5. The protein of interest was eluted in Buffer A supplemented with 0.5M Imidazole, pH 8.5, using an elution gradient from 20 mM imidazole to 500 mM imidazole. The pooled fractions were dialyzed in buffer: 50 mM MES, 50 mM NaCl, 1 mM TCEP, 0.02% CHAPS, 0.2M Arginine, 3% glycerol, pH 6.5. The purified material was qualified by mass spectrometry and analytical ultra-centrifugation.

• **Human CD3 E+G HuFc-6xHis (E+G indicates $\epsilon\gamma$ subunits)**

A cell line to produce Human CD3 E+G HuFc-6xHis was generated using HEK-293 cells (Thermo Fisher), the Lenti-X Lentiviral System (Clontech), and plasmid encoding Human CD3 E+G HuFc-6xHis (human CD3E Accession No: P07766; human CD3E+G-HuFc-His: SEQ ID NO:108). For expression, cells were cultured and expanded in Freestyle 293 media, at 37°C, humidified 8% CO₂ environment, and shaking at 135 rpm. The conditioned culture supernatant was harvested at Day 6 by centrifugation for 30 minutes at 9300xg. Expression was monitored by SDS-PAGE and Western Blotting. The conditioned culture

supernatant was adjusted with 0.2M Sucrose, 5% glycerol, 0.01% CHAPS, and 10mM Imidazole. The pH was then adjusted to 7.2. Purification was carried out in a two-step process: affinity purification using Ni/NTA resin (overnight incubation at 4C, and elution with 250mM Imidazole); followed by size-exclusion chromatography on a Superdex 200 column in destination buffer PBS with 0.2M Sucrose, 5% glycerol, 0.01% CHAPS, 1mM TCEP, pH7.2. The pooled material was concentrated using a 10K MWCO PES membrane viva cell 100 centrifugation device prior to final analysis and storage. The purified material was qualified by mass spectrometry and analytical ultra-centrifugation.

• **Cyno CD3 E+G HuFc-6xHis (E+G indicates $\epsilon\gamma$ subunits)**

A cell line to produce Cyno CD3 E+G HuFc-6xHis was generated using HEK-293 cells (Thermo Fisher), the Lenti-X Lentiviral System (Clontech), and plasmid encoding Cyno CD3 E+G HuFc-6xHis (Cynomolgus CD3E Accession No: Q95LI5 <, cyno CD3 E+G huFc-His: SEQ ID NO:109). For expression, cells were cultured and expanded in Freestyle 293 media, at 37C, humidified 8% CO2 environment, and shaking at 135 rpm. The conditioned culture supernatant was harvested at Day 6 by centrifugation for 30 minutes at 9300xg. Expression was monitored by SDS-PAGE and Western Blotting. The conditioned culture supernatant was adjusted with 0.2M Sucrose, 5% glycerol, 0.01% CHAPS, and 10mM Imidazole. The pH was then adjusted to 7.2. Purification was carried out in a two-step process: affinity purification using Ni/NTA resin (overnight incubation at 4C, and elution with 250mM Imidazole); followed by size-exclusion chromatography on a Superdex 200 column in destination buffer PBS with 0.2M Sucrose, 5% glycerol, 0.01% CHAPS, 1mM TCEP, pH7.2. The pooled material was concentrated using a 10K MWCO PES membrane viva cell 100 centrifugation device prior to final analysis and storage. The purified material was qualified by mass spectrometry and analytical ultra-centrifugation.

Example 4A: SPR based determination of affinities to recombinant DLL3 and CD3 $\epsilon\gamma$ subunits and interspecies cross-reactivity

The binding affinity of purified DLL3/CD3 bispecific constructs for recombinant human and cyno DLL3-ECD and Fc-fusion proteins of human and cyno CD3 $\epsilon\gamma$ subunits was determined by Surface Plasmon Resonance (SPR), using a ProteOn XPR36 instrument (Bio-Rad). The running buffer and all dilutions (except where stated) were done in PBS-T-EDTA. The GLM sensor chip (Bio-Rad) was normalized and pre-conditioned as per the manufacturer's recommendations. The sensor chip was activated with equal mixture of EDC/s-NHS in the horizontal direction for 300 sec at a flow rate of 30 ul/min and

immobilized with protein A/G (20 ug/ml in 10 mM acetate pH 4.5) in the horizontal direction for 300 sec at a flowrate of 30 μ l/min resulting in $\sim$ 5000 RU of protein A/G on the surface. The sensor chip was deactivated with 1M ethanolamine HCl in the horizontal direction for 300 sec at a flowrate of 30 μ l/min. The sensor chip was stabilized with 18 sec
 5 of 0.85% phosphoric acid at a flowrate of 100 μ l/min 3 times horizontally and 3 times vertically.

For binding kinetic determination to DLL3, bispecific molecules were captured individually on the protein A/G surface vertically for 250 sec at a flowrate of 25 μ l/min. The baseline was stabilized by injecting PBS-T-EDTA for 60 sec at a flowrate of 40
 10 μ l/min horizontally. The analyte (hu DLL3 or cy DLL3) was injected horizontally over the captured antibody for 600 sec at a flowrate of 40 μ l/min and a dissociation for 2700 sec. The concentrations of the analyte were 20 nM, 10 nM, 5 nM, 2.5 nM, and 1.25 nM. The surface was regenerated by injecting 0.85% phosphoric acid solution for 18 sec at a flowrate of 100 μ l/min one time horizontally and one time vertically.

15 For binding kinetic determination to CD3, hu CD3 $\epsilon\gamma$ -Fc and cy CD3 $\epsilon\gamma$ -Fc (dissolved in pH 4.5 acetate) was immobilized directly onto the sensor surface through amine coupling for 140 sec at a flowrate of 30 μ l/min vertically. The baseline was stabilized by injecting PBS-T-EDTA for 60 sec at a flowrate of 40 μ l/min horizontally. The analyte (the bispecific molecule) was injected horizontally over the captured CD3 $\epsilon\gamma$ -Fc construct for
 20 600 sec at a flowrate of 40 μ l/min and a dissociation for 600 sec. The concentrations of the analyte were 250 nM, 125 nM, 62.5 nM, 31.3 nM, and 15.6 nM or 2 μ M, 1 μ M, 0.5 μ M, 0.25 μ M, and 0.125 μ M. The surface was regenerated by injecting 0.85% phosphoric acid solution for 18 sec at a flowrate of 100 μ l/min one time horizontally and one time vertically.

25 The inter spot (interactions with sensor surface) and blank (PBS-T-EDTA) were subtracted from the raw data. Sensorgrams were then fit to 1:1 kinetic model to provide rate constants (k_a , k_d) as well as affinity (KD) value for hu DLL3 and cy DLL3, and fit globally to equilibrium to provide affinity (KD) value for hu CD3 $\epsilon\gamma$ and cy CD3 $\epsilon\gamma$.

DLL3/CD3 binding proteins described herein showed affinities to DLL3 in the pM range.
 30 Affinities to the CD3 $\epsilon\gamma$ subunit were in the low nM range. The interspecies gap between human and cyno is balanced. Affinities of three exemplary DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:75 and a CD3

chain of SEQ ID NO:79, SEQ ID NO:80 or SEQ ID NO:105, respectively) are shown in Table 2A.

Table 2A: Affinities (KD) of DLL3/CD3 binding proteins to human and cynomolgus monkey DLL3 and CD3 $\epsilon\gamma$ subunit as determined by SPR analysis, and interspecies gaps

DLL3/CD3 binding protein	Recomb. human DLL3	Recomb. cyno DLL3	Affinity gap KD hu/cyno DLL3	Recomb. Human CD3 $\epsilon\gamma$	Recomb. Cyno CD3 $\epsilon\gamma$	Affinity gap KD hu/cyno CD3 $\epsilon\gamma$
DLL3#3/CD3#1	< 10 pM	< 10 pM	1	7 nM	8 nM	1.1
DLL3#3/CD3#2	< 10 pM	< 10 pM	1	90 nM	100 nM	1.1
DLL3#3/CD3#3	< 10 pM	< 10 pM	1	68 nM	71 nM	1.04

5

Example 4B: SPR based determination of affinities to recombinant DLL3 and CD3 $\epsilon\gamma$

The binding affinity of purified DLL3/CD3 bispecific constructs for recombinant human DLL3-ECD and Fc-fusion protein of human CD3 $\epsilon\gamma$ subunits was determined by Surface Plasmon Resonance (SPR), using a ProteOn XPR36 instrument (Bio-Rad). The method was similar as in Example 4A, with a few differences as outlined below.

For binding kinetic determination to DLL3, anti-DLL3/CD3 molecules were captured individually on the protein A/G surface vertically for 200 sec at a flowrate of 30 $\mu\text{l}/\text{min}$. The baseline was stabilized by injecting PBS-T-EDTA for 60 sec at a flowrate of 30 $\mu\text{l}/\text{min}$ horizontally. The analyte (hu DLL3) was injected horizontally over the captured antibody for 300 sec at a flowrate of 30 $\mu\text{l}/\text{min}$ and a dissociation for 1800 sec. The concentrations of the analyte were 20 nM, 10 nM, 5 nM, 2.5 nM, and 1.25 nM. The surface was regenerated by injecting 0.85% phosphoric acid solution for 18 sec at a flowrate of 100 $\mu\text{l}/\text{min}$ one time horizontally and one time vertically.

For binding kinetic determination to CD3, HuCD3E_HuCD3G-Fc (dissolved in pH 4.5 acetate) was immobilized directly onto the sensor surface through amine coupling for 360 sec at a flowrate of 30 $\mu\text{l}/\text{min}$ vertically. The baseline was stabilized by injecting PBS-T-EDTA for 60 sec at a flowrate of 30 $\mu\text{l}/\text{min}$ horizontally. The analytes (all anti-DLL3/CD3 molecules) were injected horizontally over the captured antibody for 600 sec at a flowrate of 30 $\mu\text{l}/\text{min}$ and a dissociation for 1800 sec. The concentrations of the analyte were 250 nM, 83.3 nM, 27.8 nM, 9.3 nM, and 3.1 nM. The surface was regenerated by injecting

0.85% phosphoric acid solution for 18 sec at a flowrate of 100 μ l/min one time horizontally and one time vertically.

Affinities of exemplary DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, or SEQ ID NO:252 and a CD3 chain of SEQ ID NO:79, a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:105) are shown in 2B.

Table 2B: Affinities (KD) of DLL3/CD3 binding proteins to human DLL3 and CD3 $\epsilon\gamma$ subunit as determined by SPR analysis

DLL3/CD3 binding protein	Recomb. human huDLL3	Recomb. Human huCD3 $\epsilon\gamma$
DLL3#1/CD3#1	52 pM	2 nM
DLL3#2/CD3#1	91 pM	3 nM
DLL3#3/CD3#1	75 pM	2 nM
DLL3#3/CD3#2	120 pM	7 nM
DLL3#3/CD3#3	116 pM	4 nM
DLL3#4/CD3#1	< 20 pM	2 nM
DLL3#5/CD3#1	< 20 pM	3 nM
DLL3#6/CD3#1	152 pM	2 nM
DLL3#7/CD3#1	637 pM	1 nM
DLL3#8/CD3#1	40 pM	2 nM
DLL3#9/CD3#1	46 pM	2 nM
DLL3#10/CD3#1	141 pM	na
DLL3#11/CD3#1	37 pM	2 nM
DLL3#12/CD3#1	< 20 pM	2 nM
DLL3#13/CD3#1	23 pM	3 nM

DLL3/CD3 binding protein	Recomb. human huDLL3	Recomb. Human huCD3 $\epsilon\gamma$
DLL3#14/CD3#1	60 pM	3 nM
DLL3#15/CD3#1	na	na
DLL3#16/CD3#1	228 pM	2 nM
DLL3#17/CD3#1	< 20pM	2 nM
DLL3#18/CD3#1	< 20 pM	3 nM

Example 5: Generation of HEK293 cells expressing the extracellular domain and its subdomains on the cell surface

For generation of stable HEK293 cells expressing full-length human DLL3 (Uniprot: Q9NYJ7), cynomolgus DLL3 (UPI0003AB95BD), human DLL1 (Q00548) and human

5 DLL4 (Q9NR61), respectively, the respective coding sequence was cloned into pcDNA3.1 (Thermo Fisher Scientific) and a FLAG-tag was inserted between the signal sequence and the extracellular domain. The expression on the cell surface was verified by using an anti-FLAG antibody (clone M2, Sigma) followed by a monoclonal anti mouse IgG1 (Acris).

The extracellular domain of DLL3 consists of different subdomains:

- 10
- Hu DLL3 Signal peptide: aa 1-26
 - Hu DLL3 N-terminal ECD domain: aa 27-175
 - Hu DLL3 DSL: aa 176-215
 - Hu DLL3 EGF1: aa 216-249
 - Hu DLL3 EGF2: aa 274-310
 - 15 • Hu DLL3 EGF3: aa 312-351
 - Hu DLL3 EGF4: aa 353-389
 - Hu DLL3 EGF5: aa 391-427
 - Hu DLL3 EGF6: aa 429-465
 - Membrane proximal peptide: aa 466-492

20 The following subdomains of DLL3 were expressed on the cell surface of HEK293 cells:

- Hu DLL3 EGF1+2: Uniprot: Q9NYJ7 aa 216-310
- Hu DLL3 EGF 2+3: Uniprot: Q9NYJ7 aa 274-351
- Hu DLL3 EGF3+4: Uniprot: Q9NYJ7 aa 312-389
- Hu DLL3 EGF4+5: Uniprot: Q9NYJ7 aa 353-427
- 25 • Hu DLL3 EGF5+6: Uniprot: Q9NYJ7 aa 391-465

- Hu DLL3 EGF6+membrane proximal peptide: Uniprot: Q9NYJ7 aa 429-492

For the generation of HEK293 cells expressing 6-His-myc tagged subdomains of human DLL3, the respective coding sequences were cloned into pcDNA 3.1 (Thermo Fisher Scientific). Constructs contain an N-terminal mouse IgG Vk leading sequence, followed by a 6-Histidin-myc tag and the respective human DLL3 domain. To ensure cell surface localization of the human DLL3 domains, all constructs were followed by a Ser/Gly-Linker, the transmembrane domain (aa266-288) and the intracellular domain (aa 289-314) of human EpCAM (P16422). The expression of the subdomains was verified using mouse monoclonal IgG2a antibodies against various domains as described in WO 2013126746. Bound monoclonal antibody was detected with an goat anti-mouse IgG (F(ab')₂)-RPE (Jackson Immuno Research). The samples were measured by flow cytometry. The expression of human DLL1 and DLL4 was confirmed by FACS analysis using anti-DLL1 (R&D Systems, MAB1818) and anti-DLL4 R&D Systems, MAB1506) antibodies followed by PE-labeled anti-mouse or anti-rat secondary antibodies (see Figure 5, right panel).

Table 3: Sequences of various DLL3 subdomains expressed on HEK293 cells

ID	Construct	Sequence
SEQ ID NO:110	FLAG-tagged human DLL3 full length	MVSPRMSGLLSQTIVILALIFLPQTRPDYKDDDDKAGVFELQIHSGPGPGPGAPRSPCSARLPCRLFFRVCLKPGLSEEAAESPCALGAALSARGPVYTEQPGAPAPDLPLPDGLLQVPFRDAWPGTFSFIETWREELGDQIGGPAWSLLARVAGRRRLAAGGPWARDIQRAGAWELRF SYRARCEPPAVGTACTRLCRPRSAPSRGPGRLPCAPLEDECEAP LVCRAGCSPEHGFCEQPGECRCLEGWTGPLCTVPVSTSSCLSPRG PSSATTGCLVPGPGCDGNPCANGGSCSETPRSFECTCPRGFYGL RCEVSGVTCADGPCFNGGLCVGGADPDSAYICHCPPGFQGSNCE KRVDRCSLQPCRNGGLCLDLGHALRCRCRAGFAGPRCEHDLDD CAGRACANGGTCVEGGGAHRCSCALGFGGRDCRERADPCAARP CAHGGRCYAHFSGLVACAPGYMGARCEFPVHPDGASALPAAP PGLRPGDPQRYLLPPALGLLVAAGVAGAALLLVHVRRRGHSQD AGSRLLAGTPEPSVHALPDALNNLRTQEGSGDGPSSSDWNRPEDVDPQGIYVISAPSIYAREVATPLFPPLHTGRAGQRQHLLFPYPSSI LSVK

ID	Construct	Sequence
SEQ ID NO:111	FLAG- tagged cyno DLL3 full length	MVSPRMSRLLSQTVILALIFIPQARPDYKDDDDKAGVFELQIHSFG PGPGPGAPRSPCSARGPCRLFFRVCLKPGLSEEAESPALGAALS ARGPVYTEQPEAPAPDLPLPNGLLQVPFRDAWPGTFSLIETWREE LGDQIGGPAWSLLARVTRRRRLAAGGPWARDIQRAGAWELRFS YRARCELPVAGTACTRLCRPRSAPSRCPGLRPCAPLEDECEAPP VCRAGCSLEHGFCEQPGECRCLEGTGTPLCMPVSTSSCLGLRG PSSTTTGCLVPGPGPCDGNPCANGGSCSETPGSFECTCPRGFYGLR CEVSGVTCADGPCFNGGLCVGGADPDSAYICHCPPGFQGSNCEK RVDRCSLQPCRNGGLCLDLGHALRCRCRAGFAGPRCEHDLDDC AGRACANGGTCVEGGGAHRCSCALGFGGRNCRERADPCAARPC AHGGRCYAHFSGLVACAPGYMGARCEFPVHPDGVVSALPAAPP GLRPGDPQRYLLPPALGGLVAAGVAGAALLLVHVRRRGHAQDA GSRLLAGTPEPSVHALPDALNNLRTQEGPGDVPSSVDWNRPED VDSRGIYVISAPSIYAREVAMPLFPPLHTGRAGQRQNLLFPFPSSIL SVK
SEQ ID NO:112	FLAG- tagged human DLL1 full length	MGSRCALALAVLSALLCDYKDDDDKQVWSSGVFELKLQEFVNK KGLLGNRNCCRGAGPPPCACRTFFRVCLKHYQASVSPEPPCTY GSAVTPVLGVDSFSLPDGGGADSAFSPNPIRFPFGFTWPGTFSLIIEA LHTDSPDDLATENPERLISRLATQRHLTVGEEWSQDLHSSGRTDL KYSYRFVCDHYHYGEGCSVFCRPRDDAFGHFTCGERGEKVCNPG WKGPYCTEPICLPGCDEQHGFCDKPGECKCRVGVWQGRYCDCEIR YPGCLHGTCCQPWQCNCQEGWGGLFCNQDLNYCTHHKPKNG ATCTNTGQGSYTCSCRPGYTGTATCELGIDCDPSPCKNGGSCDTL ENSYSCTCPPGFYKICELSAMTCADGPCFNGGRCSDDPGGYSC RCPVGYSGFNCEKKIDYCSSSPCSNGAKCVDLGDAYLCRCQAGF SGRHCDNDVDDCASSPCANGGTCRDGVNDFSTCPPGYTGRNCS APVSRCEHAPCHNGATCHERGHRYVCECARGYGGPNCQFLLPEL PPGPAVVDLLEKLEGQGGPFPWVAVCAGVILVLMMLLGCAAVV VCVRLRLQKHRPPADPCRGETETMNNLANCQREKDISVSIIGATQ IKNTNKKADFHGDHSADKNGFKARYPAVDYNLVQDLKGDDTAV RDAHSKRDTKCQPQGSSEEGKTPTTLRGGEASERKRPDSGCSTS KDTKYQSVYVISEEKDECVIATEV
SEQ ID NO:113	FLAG- tagged human DLL4 full length	MAAASRSASGWALLLLVALWQQRAAGDYKDDDDKSGVFQLQL QEFINERGVLASGRPCEPGCRTFFRVCLKHFQAVVSPGPCTFGTV STPVLGTNSFAVRDSSGGGRNPLQLPFNFTWPGTFSLIIEAWHAP GDDLRLPEALPPDALISKIAIQGSLAVGQNWLLDEQSTLTRLRYSY RVICSDNYYGDNCSRLCKKRNDHFGHYVCQPDGNLSCLPGWTG EYCQQPICLSGCHEQNGYCSKPAECLCRPGWQGRCLNECIPHNG CRHGTCTPWQCTCDEGWGGLFCDQDLNYCTHHSPCKNGATCS NSGQRSYTCTCRPGYTGVDCLELSECDSPCRNGGSKDQEDG YHCLCPPGYYGLHCEHSTLSCADSPCFNGGSCRERNQGANYACE CPPNFTGSNCEKKVDRCTSNPCANGGQCLNRGPSRMCRCPGFT GTYCELHVSDCARNPCAHHGTCHDLENGLMCTCPAGFSGRRC VRTSIDACASSPCFNATCYTDLSTDTFVCNCPYGFVGSRCFPV GLPPSFPWVAVSLGVGLAVLLVLLGMVAVAVRQLRLRRPDDGS REAMNNLSDFQKDNLPAAQLKNTNQKKELEVDCGLDKSNCGK QQNHTLDYNLAPGPLGRGTMPGKFPHSDKSLGEKAPLRHLHSEKP ECRISAICSPRDSMYQSVCLISEERNECVIATEV

ID	Construct	Sequence
SEQ ID NO:114	FLAG tag	DYKDDDDK
SEQ ID NO:115	Vk leader	METDTLLLWVLLLVPGSTGD
SEQ ID NO:116	6His-myc tag	HHHHHHEQKLISEEDL
SEQ ID NO:117	Ser/Gly linker	SGGGGS
SEQ ID NO:118	EpCAM transmembrane domain	AGVIAVIVVVVIAVVAGIVLVI
SEQ ID NO:119	EpCAM intracellular domain	SRKKRMAKYEKAEIKEMGEMHRELNA
SEQ ID NO:120	EGF1	APLVCRAGCSPEHGFCEQPGECRCLEGWTGPLCT
SEQ ID NO:121	Hu DLL3 EGF 1+2	APLVCRAGCSPEHGFCEQPGECRCLEGWTGPLCTVPVSTSSCLSP RGPSSATTGCLVPGPGPCDGNPCANGGSCSETPRSFECTCPRGFY GLRCE
SEQ ID NO:122	EGF2	GPGPCDGNPCANGGSCSETPRSFECTCPRGFYGLRCE
SEQ ID NO:123	Hu DLL3 EGF 2+3	GPGPCDGNPCANGGSCSETPRSFECTCPRGFYGLRCEVSGVTCAD GPCFNGGLCVGGADPDSAYICHCPPGFQGSNCE
SEQ ID NO:124	EGF3	SGVTCADGPCFNGGLCVGGADPDSAYICHCPPGFQGSNCE
SEQ ID NO:125	Hu DLL3 EGF 3+4	SGVTCADGPCFNGGLCVGGADPDSAYICHCPPGFQGSNCEKRVD RCSLQPCRNGGLCLDLGHALRCRCRAGFAGPRCE
SEQ ID NO:126	EGF4	RVDRCSLQPCRNGGLCLDLGHALRCRCRAGFAGPRCE
SEQ ID NO:127	Hu DLL3 EGF 4+5	RVDRCSLQPCRNGGLCLDLGHALRCRCRAGFAGPRCEHLDLDDC AGRACANGGTCVEGGGAHRCSCALGFGGRDCR
SEQ ID NO:128	EGF5	DLDDCAGRACANGGTCVEGGGAHRCSCALGFGGRDCR
SEQ ID NO:129	Hu DLL3 EGF 5+6	DDCAGRACANGGTCVEGGGAHRCSCALGFGGRDCRERADPCAA RPCAHGGRCYAHFSGLVACAPGYMGARCE

ID	Construct	Sequence
SEQ ID NO:130	EGF6	RADPCAARPCAHGGRCYAHFSGLVACAPGYMGARCE
SEQ ID NO:131	Hu DLL3 EGF6-membrane proximal peptide	RADPCAARPCAHGGRCYAHFSGLVACAPGYMGARCEFPVHPD GASALPAAPPGLRPGDPQRYL
SEQ ID NO:132	Membrane proximal peptide	FPVHPDGASALPAAPPGLRPGDPQRYL

5 Example 6: Epitope domain mapping of DLL3/CD3 binding proteins

Epitope domains were determined by binding of DLL3/CD3 binding proteins to recombinant cell lines expressing subdomains of DLL3 (see Figure 2). Generation of these cell lines is described under Example 5.

10 DLL3/CD3 binding proteins were produced by transient transfection of CHO-E cells with the pTT5 vectors carrying the chain-encoding genes, as described in Example 2.

Cells expressing subdomains of DLL3 were stained with two-step purified DLL3/CD3 binding proteins at 1.6 nM in FACS buffer (PBS/0.5%BSA/0.05% sodium azide). Bound molecules were detected with PE-conjugated anti-human secondary antibody (Sigma-Aldrich, #P8047). As negative control, cells were incubated with secondary antibody only.

15 The samples were measured by flow-cytometry. Figure 3A-C show exemplary binders (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID
20 NO:251 or SEQ ID NO:252 and a CD3 chain of SEQ ID NO:79, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80) binding to different regions of DLL3. The sequences of the epitope domains are listed in Example 5, Table 3.

Example 7: Interspecies cross-reactivity

Interspecies cross-reactivity was determined by binding of DLL3/CD3 binding proteins to
5 NCI-H82 (HTB-175TM) and SHP77 (ATCC®, CRL-2195TM) cells, two SCLC cell lines, as
well as to a recombinant cell line expressing cynomolgus DLL3 (generation of cell line
described in Example 5).

DLL3/CD3 binding proteins were produced by transient transfection of CHO-E cells with
the pTT5 vectors carrying the chain-encoding genes, as described in Example 2. Cells
10 expressing human or cynomolgus DLL3 were stained with 1.6 nM of two-step purified
DLL3/CD3 binding proteins with increasing concentrations in FACS buffer
(PBS/0.5%BSA/0.05% sodium azide). Bound molecules were detected with PE-conjugated
anti-human secondary antibody (Sigma-Aldrich, #P8047). As negative control, cells were
incubated with secondary antibody only. The samples were measured by flow-cytometry.
15 Figure 4 shows binding to human and cyno DLL3-expressing cells of seven exemplary
DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of
SEQ ID NO:73 and a CD3 chain of SEQ ID NO:79, a DLL3 chain of SEQ ID NO:74 and a
CD3 chain of SEQ ID NO:79, a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ
ID NO:79, a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, a DLL3
20 chain of SEQ ID NO:76 and a CD3 chain of SEQ ID NO:79, a DLL3 chain of SEQ ID
NO:77 and a CD3 chain of SEQ ID NO:79 or a DLL3 chain of SEQ ID NO:78 and a CD3
chain of SEQ ID NO:79).

Example 8: Confirmation of the absence of binding to human DLL1 and DLL4

Absence of cross-reactivity to human DLL1 and DLL4 was determined by binding of
25 DLL3/CD3 binding proteins to recombinant cells expressing human DLL1 or DLL4
(generation of cell lines described in Example 5).

DLL3/CD3 binding proteins were produced by transient transfection of CHO-E cells with
the pTT5 vectors carrying the chain-encoding genes, as described in Example 2.

Cells expressing DLL1 and DLL4 were stained with 1.6 nM of two-step purified
30 DLL3/CD3 binding proteins with increasing concentrations in FACS buffer
(PBS/0.5%BSA/0.05% sodium azide). Bound molecules were detected with PE-conjugated
anti-human secondary antibody (Sigma-Aldrich, #P8047). As negative control, cells were

incubated with secondary antibody only. The samples were measured by flow-cytometry. Expression of human DLL1 and DLL4 was confirmed by FACS analysis using anti-DLL1 (R&D Systems, MAB1818) and anti-DLL4 R&D Systems, MAB1506) antibodies followed by PE-labeled anti-mouse or anti-rat secondary antibodies. Figure 5 shows

5 binding of seven exemplary DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73 and a CD3 chain of SEQ ID NO:79, a DLL3 chain of SEQ ID NO:74 and a CD3 chain of SEQ ID NO:79, a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:79, a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, a DLL3 chain of SEQ ID NO:76 and a CD3 chain of SEQ ID

10 NO:79, a DLL3 chain of SEQ ID NO:77 and a CD3 chain of SEQ ID NO:79 or a DLL3 chain of SEQ ID NO:78 and a CD3 chain of SEQ ID NO:79) to human DLL1 and DLL4-expressing cells.

Example 9: Binding of DLL3/CD3 binding proteins to SCLC cell lines and T cells

15 Binding of DLL3/CD3 binding proteins to human SCLC cell lines was tested by flow cytometry of NCI-H82 (HTB-175TM) and SHP77 (ATCC®, CRL-2195TM). DLL3/CD3 binding proteins were produced, as described in Example 2.

Human peripheral blood mononuclear cells (PBMCs) were prepared by Ficoll density gradient centrifugation from enriched lymphocyte preparations (buffy coats), a side

20 product of blood banks collecting blood for transfusions. All buffy coats were obtained after informed consent in accordance with the Declaration of Helsinki and with approval of the cantonal ethical committee in Austria and PBMCs were prepared the same day of collection. Therefore, mononuclear cells were isolated by Ficoll density gradient centrifugation (35 min without brake at 1400 rpm) and extensive washes with PBS.

25 Remaining erythrocytes were removed by incubating for 3 minutes in ACK lysis buffer (Thermo Fisher Scientific, A1049201), followed by washing in PBS, before suspension in assay medium containing RPMI 1640 GlutaMAX (Gibco #61870-010), 5% human AB serum AB (Gemini, GemCell cat # 100-512 LOT#H56500I) + 1% MEM-NEAA (Gibco #11140-035), 10mM HEPES (Affymetrix #7365-49-9), 10 µM beta-Mercaptoethanol

30 (Gibco #21985-023) and sodium pyruvate (Gibco #11360-039).

T-cells were isolated by negative selection using the Pan T Cell Isolation Kit II (Miltenyi Biotec #130-091-156). In brief, cells were resuspend in 40 µl buffer PBS/0,5% BSA

(Gibco ref#041-94553 M)/2mM EDTA (Invitrogen ref# 15575-038) per 10 Mio cells and incubated with 10 µl of Biotin-Antibody cocktail per 10 Mio cells for 5 min at 4 °C. Subsequently, 30 µl buffer and 20 µl anti-biotin MicroBeds/10 million cells were added and incubated for 10 min at 4°C. Subsequently the mixture was placed in a pre-rinsed

5 25LS column (Miltenyi Biotec #130-042-401) in the magnetic field of suitable MACS separator (Miltenyi Biotec). Flow-through was collected and washed in assay medium.

Cells (T cells or human SCLC cells) were stained with increasing concentrations of two-step purified DLL3/CD3 binding proteins with increasing concentrations in FACS buffer (PBS/0.5%BSA/0.05% sodium azide). Bound molecules were detected with PE-conjugated

10 anti-human secondary antibody (Sigma-Aldrich, #P8047). DLL3/CD3 binding proteins targeting DSL (DLL3#6), EGF1 (DLL3#5) or EGF4 (DLL#4) domains show stronger binding to SCLC cell lines expressing DLL3, compared to DLL3/CD3 binding proteins which target the membrane proximal peptide (DLL3#1, DLL3#2, DLL3#3). Binding to SHP77, NCI-H82 cells and T cells of exemplary DLL3/CD3 binding proteins (DLL3/CD3

15 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251 or SEQ ID NO:252 and a CD3 chain of SEQ ID NO:79, and a DLL3 binding protein comprising a

20 DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80) against different epitope domains is shown in Figure 6A-V.

Example 10A: Potency of redirecting non-stimulated PBMCs against human SCLC cell lines

25 Potency of non-stimulated PBMCs against SCLC cell lines was determined using lactate-dehydrogenase (LDH) release as readout for cell lysis. In this assay, DLL3-positive SCLC cell lines, SHP77 and NCI-H82, were co-cultured with human PBMCs as effector cells and increasing concentrations of DLL3/CD3 binding proteins for 72 hours at an effector to target cell ratio of 10:1. DLL3/CD3 binding proteins (DLL3/CD3 binding proteins

30 comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, and a CD3 chain of SEQ ID NO:79, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID

NO:80) were produced by transient transfection of CHO-E cells with the pTT5 vectors carrying the chain-encoding genes, as described in Example 2.

Human peripheral blood mononuclear cells (PBMCs) were prepared by Ficoll density gradient centrifugation from enriched lymphocyte preparations (buffy coats), a side
 5 product of blood banks collecting blood for transfusions. All buffy coats were obtained after informed consent in accordance with the Declaration of Helsinki and with approval of the cantonal ethical committee in Austria and PBMCs were prepared the same day of collection. Therefore, mononuclear cells were isolated by Ficoll density gradient centrifugation (35 min without brake at 1400 rpm) and extensive washes with PBS.
 10 Remaining erythrocytes were removed by incubating for 3 minutes in ACK lysis buffer (Thermo Fisher Scientific, A1049201), followed by washing in PBS, before suspension in assay medium containing RPMI 1640 GlutaMAX (Gibco #61870-010), 5% human AB serum AB (Gemini, GemCell cat # 100-512 LOT#H56500I) + 1% MEM-NEAA (Gibco #11140-035), 10mM HEPES (Affymetrix #7365-49-9), 10 µM beta-Mercaptoethanol
 15 (Gibco #21985-023) and sodium pyruvate (Gibco #11360-039).

Subsequently, target cells, SHP77 and NCI-H82, and PBMCs at a ratio of 1:10 were incubated with DLL3/CD3 binding proteins at concentrations from 0.0001 nM to 100 nM for 72 hours.

Cytotoxic activity was determined using the Cytotoxicity Detection Kit^{PLUS} (Roche),
 20 following the manufacturer's instructions. In brief, this method is based on the usage of the release of LDH from dead or plasma –membrane damaged cells. Cell culture supernatant is incubated with the reaction mixture from the kit for 30 minutes and the formation of Formazan, as a result of LDH activity is measured in a spectrophotometer at 500nm as surrogate for cell lysis. Percentage of cytotoxicity relative to the maximal lysis control
 25 was calculated according to the following formula:

$$\text{Cytotoxicity (relative to control)} = \frac{\text{measured value} - \text{background}}{\text{maximal lysis} - \text{minimal lysis}}$$

Background: Target cells + Effector cells

Maximal lysis: Target cells + 5% Triton X-100

Minimal lysis: Target cells

Using GraphPad Prism 5 software, the percentage of cytotoxicity relative to the maximal lysis control was plotted against the corresponding DLL3/CD3 binding protein concentrations. Dose response curves were analysed with the four-parameter logistic equation model for evaluation of sigmoidal dose-response curve and EC₉₀ values were calculated (EC₉₀ values are shown in Table 4A).

Figure 7A shows an example of potency in cell lysis of seven exemplary DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, or SEQ ID NO:78, and a CD3 chain of SEQ ID NO:79, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80) binding to different epitope domains.

Table 4A: EC₉₀ values [nM] of DLL3/CD3 binding proteins as measured in a 72 hour cytotoxicity assay with non-stimulated PBMCs as effector cells and SCLC cell lines, NCI-H82 and SHP77 as target cells.

DLL3/CD3 binding proteins	NCI-H82	SHP77
DLL3#1/CD3#1	1.7	1.4
DLL3#2/CD3#1	2.9	3.4
DLL3#3/CD3#1	0.9	1.5
DLL3#3/CD3#2	3.9	16.5
DLL3#4/CD3#1	no saturation at 100 nM	no saturation at 100 nM
DLL3#5/CD3#1	no saturation at 100 nM	no saturation at 100 nM
DLL3#6/CD3#1	no saturation at 100 nM	no saturation at 100 nM

Example 10B: Potency of redirecting non-stimulated T cells against human SCLC cell lines

T cells were isolated as described in Example 9. Subsequently purified T cells were used as effector cells in a cytotoxicity assay with SHP77 and NCI-H82 cells as target cells as described in Example 10A. Dose response curves are shown in Figures 7B-H (SHP77 cells) and Figure 7I-O (NCI-H82 cells). Calculated EC₉₀ values are shown in Table 4B.

Figure 7B-O show examples of potency in cell lysis of 19 exemplary DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ

ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, SEQ ID NO:252

- 5 Table 4B: EC₉₀ values [nM] of DLL3/CD3 binding proteins as measured in a 72 hour cytotoxicity assay with purified T cells as effector cells and SCLC and NCI-H82 cell lines as target cells.

DLL3/CD3 binding proteins	NCI-H82	SHP77
DLL3#1/CD3#1	0.66	0.15
DLL3#2/CD3#1	1.51	0.16
DLL3#3/CD3#1	0.27	0.02
DLL3#3/CD3#2	2.50	3.00
DLL3#4/CD3#1	no activity	no activity
DLL3#5/CD3#1	no activity	no activity
DLL3#6/CD3#1	no activity	no activity
DLL3#7/CD3#1	21.84	8.37
DLL3#8/CD3#1	no activity	no activity
DLL3#9/CD3#1	no activity	no activity
DLL3#10/CD3#1	no activity	no activity
DLL3#11/CD3#1	no activity	no activity
DLL3#12/CD3#1	no activity	no activity
DLL3#13/CD3#1	no activity	no activity
DLL3#14/CD3#1	no activity	no activity
DLL3#15/CD3#1	no activity	no activity
DLL3#16/CD3#1	no activity	no activity
DLL3#17/CD3#1	no activity	no activity
DLL3#18/CD3#1	no activity	no activity

Example 11: *In vitro* internalization assay

- 10 Changes in the potency of the DLL3/CD3 binding proteins after pre-incubated with DLL3-positive SHP77 cells as target cells were measured. If the DLL3/CD3 binding protein is

internalized, the effective concentration should decrease with this and thus the apparent potency should decrease.

DLL3/CD3 binding proteins were produced as described in Example 2. Human peripheral blood mononuclear cells (PBMCs) were prepared as described in Example 10.

- 5 SHP77 target cells were plated and incubated with DLL3/CD3 binding proteins at concentrations from 0.0001 to 100 nM for 2 and 4 hours before adding non-stimulated PBMCs for 48 hours. The effector to target cell ratio was 10:1. Cytotoxic activity was determined using the Cytotoxicity Detection Kit^{PLUS} (Roche) as described in Example 10.

- Figure 8 shows no difference in potency and efficacy of lysing cells of two exemplary
10 DLL3/CD3 binding proteins (DLL3/CD3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:79, and DLL3/CD3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80). Potency and efficacy have been observed compared to non-pre-incubated samples (0h). Results suggest that no significant internalization occurs with DLL3/CD3 binding proteins.

15

Example 12: Mouse PK study

The PK of DLL3/CD3 binding proteins was evaluated in C57BL/6 mice following a single 1 mg/kg i.v. dose. Serum concentrations of DLL3/CD3 binding proteins were determined using a DLL3 capture/CD3 detection assay.

- 20 In brief, male C57BL/6 mice received a single 1 mg/kg intravenous (IV) dose (n = 3 per molecule). Blood samples were collected pre-dose and 0.15, 2, 8, 24, 72, 96, 168, 240 and 336 hours post-dose. Serum drug levels were measured with an MSD-based ligand binding assay, using biotinylated DLL3 as the capture reagent and sulfo-tagged CD3 as the detection reagent. Pharmacokinetic (PK) parameters were calculated from serum
25 concentration time-profiles using non-compartmental analysis. The following PK parameters were assessed: AUC_{last} (area under the serum concentration-time curve from time zero to the last quantifiable time-point), AUC_{inf} (area under the serum concentration-time curve extrapolated to infinity), CL (systemic clearance), V_{ss} (steady-state volume of distribution) and t_{1/2} (terminal half-life).

- 30 Mean (SD) serum concentration time-profiles for exemplary DLL3/CD3 binding proteins (DLL3/CD3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:79, and DLL3/CD3 binding protein comprising a DLL3 chain of SEQ ID

NO:75 and a CD3 chain of SEQ ID NO:80) are summarized in Figure 9. Mean (SD) PK parameters for these exemplary DLL3/CD3 binding proteins are summarized in Table 5.

Table 5: Mean (SD) PK parameters of exemplary DLL3/CD3 binding proteins in male
5 C57BL/6 mice following a single 1 mg/kg intravenous dose

DLL3/CD3 binding proteins	AUC _{0-last} (μg*d/mL)	AUC _{0-inf} (μg*d/mL)	CL (mL/day/kg)	V _{ss} (mL/kg)	t _{1/2} (days)
DLL3#3/CD3#1	1180 (127)	1760 (179)	13.7 (1.48)	178 (57.7)	10.6 (4.67)
DLL3#3/CD3#2	1260 (170)	1440 (154)	16.8 (1.74)	109 (19.5)	4.97 (0.78)

10 **Example 13: *In vivo* xenograft efficacy study**

Efficacy studies were performed using a human xenograft mouse model reconstituted with human T cells. In detail, human SHP77 small cell lung cancer cells (2×10^7) were injected subcutaneously (*s.c.*) into the right dorsal flank of sub-lethally irradiated (2 Gy, day -1) female NOD.Cg-*Prkdc*^{scid} *Il2rg*^{tm1Sug}/JicTac mice (Day 1). In parallel, human CD3 positive
15 T cells (isolated from healthy human blood donor) were expanded *in vitro*.

Human peripheral blood mononuclear cells (PBMCs) were prepared as described in Example 10.

T-cells were isolated by negative selection using the Pan T Cell Isolation Kit II (Miltenyi Biotec #130-091-156). In brief, cells were resuspend in 40 μl buffer PBS/0,5% BSA (gibco
20 ref#041-94553 M)/2mM EDTA (Invitrogen ref# 15575-038)per 10 Mio cells and incubated with 10 μl of Biotin-Antibody cocktail per 10 Mio cells for 5 min at 4 °C. Subsequently, 30 μl buffer and 20 μl anti-biotin MicroBeds/10 million cells were added and incubated for 10 min at 4°C. Subsequently the mixture was placed in a pre-rinsed 25LS column (Miltenyi Biotec #130-042-401) in the magnetic field of suitable MACS
25 separator (Miltenyi Biotec). Flow-through was collected and washed in assay medium.

Subsequently T cells were expanded using the T Cell Activation/Expansion Kit human (Miltenyi Biotec Cat#130-091-441, Lot#5170720843) for 20 days. In brief, anti-Biotin MACSiBead™ Particles are loaded with CD2-, CD3-, CD28 Biotin and are transferred to

the purified T cells in a ratio of 2 cells per particle and incubated in presence of 20 Units recombinant IL-2 (R&D#202-IL-050/CF) at a density of 0.5-1 10⁶ cells/ml for 20 days.

Cells were supplemented with 20 Units fresh IL-2 every three days. Three days before injection into the animals, T cells were were restimulated with anti-Biotin

5 MACSiBead™ Particles are loaded with CD2-, CD3-, CD28 Biotin at a ratio of 1 bead per 4 cells for additional three days. Finally, beads were removed with a MACSiMAG Separator (Miltenyi Biotec) and T cells were washed in PBMCs.

On day 14, animals were randomized into treatment groups based on tumor volume and 2 x 10⁷ human T cells were injected intra-peritoneally (*i.p.*). Treatment was started on day 17
 10 and DLL3/CD3 binding protein(DLL3/CD3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:79, and DLL3/CD3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80) or Vehicle buffer (50mM NaOAc, 100mM NaCl, pH 5.0) was administered in a q7d dosing regimen by intravenous (*i.v.*) bolus injections into the lateral tail vein. Tumor growth was
 15 monitored by external caliper measurements and tumor volumes were calculated using a standard hemi-ellipsoid formula. Human T cell engraftment was assessed in the spleen by immunohistochemistry (IHC) staining for human CD3 at the end of the study. Only those animals showing human T-cell engraftment at the end of the study were included in the statistical analysis. Animals reaching sacrifice criteria were euthanized early during the
 20 studies for ethical reasons. Treatment of tumor-bearing mice with DLL3/CD3 binding proteins once weekly *i.v.* at 0.25 mg/kg induced significant tumor regression (Figure 10).

Example 14: Percent monomer content of DLL3/CD3 binding proteins

Percent monomer was determined for exemplary DLL3/CD3 binding proteins (DLL3/CD3
 25 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251, or SEQ ID NO:252 and a CD3 chain of SEQ ID NO:79, a DLL3 binding protein comprising a
 30 DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:105) by Analytical Size Exclusion Chromatography (aSEC) (shown in Table 6). aSEC was run on a Waters (Milford, MA, USA) Acquity UPLC system using a Protein BEH SEC column

200Å, 1.7µm, 4.6x150mm (Cat# 186005225). Running conditions were as follows: Mobile phase: 50mM Sodium Phosphate, 200mM Arginine and 0.05% Sodium Azide; Flow rate: 0.5ml/min; Runtime: 5 minutes; Sample loading amount: 10µg; Peak detection: A280nm; Automated processing method of chromatograms.

5 Table 6: Percent monomer after first and second purification step

DLL3/CD3 binding proteins	Percent monomer after 1 st step of purification	Percent monomer after 2 nd step of purification
DLL3#1/CD3#1	76.4	99.4
DLL3#2/CD3#1	54.4	98.0
DLL3#3/CD3#1	69.5	99.8
DLL3#3/CD3#2	77.6	99.5
DLL3#3/CD3#3	70.8	98.3
DLL3#4/CD3#1	72.2	99.1
DLL3#5/CD3#1	70.5	99.1
DLL3#6/CD3#1	55.2	99.9
DLL3#7/CD3#1	78.1	99.0
DLL3#8/CD3#1	54.8	99.5
DLL3#9/CD3#1	72.0	99.0
DLL3#10/CD3#1	70.2	98.3
DLL3#11/CD3#1	68.2	94.7
DLL3#12/CD3#1	66.2	98.7
DLL3#13/CD3#1	52.8	99.9
DLL3#14/CD3#1	52.6	87.0
DLL3#15/CD3#1	61.5	na
DLL3#16/CD3#1	71.8	99.5
DLL3#17/CD3#1	60.2	94.7
DLL3#18/CD3#1	61.5	98.2

Example 15A: Thermostability

Thermostability was determined by Differential Scanning Calorimetry (DSC) and results of the first melting transitions (T_{m1}) of DLL3/CD3 binding proteins are shown in Table 7. DSC thermal melts provide information regarding the thermal stability of a protein in solution relative to a buffer control. Thermal unfolding and aggregation of a 1 mg/ml solution of proteins in 20 mM Citrate, 115 mM NaCl, pH 6 was monitored from 20 °C to 110 °C at a scan rate of 60 °C/hr via an automated capillary DSC.

Table 7A

DLL3/CD3 binding proteins	Tm1
DLL3#3/CD3#1	64
DLL3#3/CD3#2	60
DLL3#3/CD3#3	63
DLL3#4/CD3#1	63

Example 15B: Thermostability

- 5 Thermostability was determined by Thermal Shift Analysis (TSA) and results of the first melting transitions (Tm1) of DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID
- 10 NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251 or SEQ ID NO:252 and a CD3 chain of SEQ ID NO:79, a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:105) are shown in Table 7B. The fluorescence intensity profile as a function of temperature was acquired using a
- 15 QuantStudio 6 Flex real-time PCR system (Applied Biosystems, Waltham, MA) with SYPRO Orange (Invitrogen, Carlsbad, CA) as the extrinsic fluorophore. Sample was diluted to 0.4 mg/ml in 10 mM histidine, pH 6.0 with 40 mM sodium chloride and 0.02% sodium azide. The melt curve was generated with a thermal ramp from 25°C to 95°C at a rate of 2°C/min, with data collected approximately every 0.4°C through the 'ROX' filter
- 20 set (Ex: 580 ±10 nm, Em: 623 ±14 nm). Data were analyzed using Protein Thermal Shift Software Version 1.3 (ThermoFisher Scientific, Waltham, MA).

Table 7B

DLL3/CD3 binding proteins	Tm1 (°C)
DLL3#1/CD3#1	65.6
DLL3#2/CD3#1	65.5
DLL3#3/CD3#1	65.6
DLL3#3/CD3#2	63.0
DLL3#3/CD3#3	65.5

DLL3/CD3 binding proteins	Tm1 (°C)
DLL3#4/CD3#1	65.8
DLL3#5/CD3#1	65.6
DLL3#6/CD3#1	65.8
DLL3#7/CD3#1	65.6
DLL3#8/CD3#1	65.8
DLL3#9/CD3#1	65.9
DLL3#10/CD3#1	64.6
DLL3#11/CD3#1	66.0
DLL3#12/CD3#1	65.9
DLL3#13/CD3#1	65.8
DLL3#14/CD3#1	65.9
DLL3#15/CD3#1	na
DLL3#16/CD3#1	65.8
DLL3#17/CD3#1	65.8
DLL3#18/CD3#1	65.8

Example 16: Predicted Immunogenicity scores *in silico* by Epivax:

Immunogenicity of sequences was evaluated *in silico* with a mathematical algorithm.

Specifically, EpiMatrix Treg-adjusted Scores (EpiVax Inc., Providence RI)) as a measure
of immunogenicity scores, were determined for DLL3/CD3 binding proteins (DLL3/CD3
binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID
NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID
NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID
NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251 or SEQ ID
NO:252 and a CD3 chain of SEQ ID NO:79, a DLL3 binding protein comprising a DLL3
chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, and a DLL3 binding protein
comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:105) and
compared to the scores of various Fc sequences. These scores are taking T-cell epitopes
and Treg epitopes into consideration. The lower the immunogenicity score, the less likely a
sequence to be immunogenic. In general, a negative score is considered low risk of
immunogenicity, while a highly positive score is viewed as indication for potential
immunogenicity. As shown in the tables below, DLL3/CD3 binding proteins described
herein have very low immunogenicity scores, indicating that the risk of being
immunogenic is low for these binding proteins.

Table 8: Adjusted Epivax scores of DLL3/CD3 binding proteins

DLL3/CD3 binding proteins	VH	VL	Full polypeptide chain (VL-CL-linker-VH-CH1-hinge-CH2-CH3)
DLL3#1 DLL3 chain	-36.88	-59.85	-43.31
DLL3#2 DLL3 chain	-46.99	31.40	-33.53
DLL3#3 DLL3 chain	-34.44	2.84	-35.50
DLL3#4 DLL3 chain	25.97	28.92	-21.93
DLL3#5 DLL3 chain	-59.87	-46.45	-44.95
DLL3#6 DLL3 chain	29.70	-14.55	-27.26
CD3#1 CD3 chain	-9.68	-50.52	-35.25
CD3#2 CD3 chain	-15.54	-50.52	-36.23
CD3#3 CD3 chain	-13.49	-50.52	-35.88
DLL3#7 DLL3 chain	-48.55	-34.92	-41.60
DLL3#8 DLL3 chain	-54.39	-37.79	-42.89
DLL3#9 DLL3 chain	35.95	-60.37	-31.83
DLL3#10 DLL3 chain	-33.82	-34.60	-40.73
DLL3#11 DLL3 chain	-24.00	26.66	-30.50
DLL3#12 DLL3 chain	-30.43	25.68	-31.67
DLL3#13 DLL3 chain	8.64	18.63	-25.96
DLL3#14 DLL3 chain	1.58	-10.63	-31.72
DLL3#15 DLL3 chain	8.93	-8.31	-29.91
DLL3#16 DLL3 chain	-55.14	13.98	-35.59
DLL3#17 DLL3 chain	11.89	12.55	-26.52
DLL3#18 DLL3 chain	5.82	15.85	-27.01

Table 9: Adjusted Epivax scores of Fc domains

Fc Protein Chain	Adjusted Epivax score
Fc-IgG1-WT	-25.64
Fc-IgG1-LALA	-29.83
Fc-IgG1-LALA-KNOB	-31.76
Fc-IgG1-LALA-HOLE	-18.01

5

Example 17A: Non-specific binding to surfaces

The specificity of the DLL3/CD3 binding proteins of the invention was further tested in a SPR-based assay using highly charged proteins. A non-specific binding assay was

developed using biosensor technology to determine if binding proteins have significant binding to unrelated charged proteins. In this assay, DLL3/CD3 binding proteins were passed over two SPR surfaces, one coated with a negatively charged protein (Trypsin Inhibitor) and one coated with a positively charged protein (Lysozyme). When a protein displays significant non-specific binding to these surfaces, it is likely that the cause of binding is the presence of positive or negative charged surface patches on the candidate. Non-specific binding of proteins may translate to poor pharmacokinetics (PK) and biodistribution and may also have downstream manufacturability impacts.

The experiment was performed on a Biacore T200. The dilution, surface preparation, and binding experiments were performed at 25 °C in 1X HBS-EP buffer prepared from 10X HBS-EP. The flow rate for both the immobilization protocol and binding experiment was at 5 µL/min.

To prepare the surface for the non-specific binding experiment, chicken egg white lysozyme and trypsin inhibitor from glycine max soybean were coupled manually to a series S CM5 chip with the surface density of 3000-5000 RU using the amine coupling kit according to the manufacture instructions.

Samples were prepared at 1 µM in 1X HBS-EP buffer. The samples were injected over activated surfaces with a 10 min association and 10 min dissociation. The data was collected using Biacore T200 Control Software version 2.0.1 and analyzed using Biacore T200 Evaluation Software version 3.0.

The DLL3/CD3 binding proteins did not bind to these highly charged surfaces. Table 10A shows absence of binding to the two highly charged proteins, Trypsin Inhibitor and Lysozyme, with two exemplary DLL3/CD3 binding proteins.

Table 10A:

DLL3/CD3 Binding Protein	Non-specific binding	
	Lysozyme (positive)	Tryp. Inhibitor (negative)
DLL3#3/CD3#1	No	No
DLL3#3/CD3#2	No	No

Example 17B: Non-specific binding to surfaces

The specificity of the DLL3/CD3 binding proteins (DLL3/CD3 binding proteins comprising a DLL3 chain of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID

NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO:242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO:245, SEQ ID NO:246, SEQ ID NO:247, SEQ ID NO:248, SEQ ID NO:249, SEQ ID NO:250, SEQ ID NO:251 or SEQ ID NO:252, and a CD3 chain of SEQ ID NO:79, a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:80, and a DLL3 binding protein comprising a DLL3 chain of SEQ ID NO:75 and a CD3 chain of SEQ ID NO:105) was further tested in an SPR-based assay using highly charged proteins, as described in Example 17a, with a 10 min association and 15 min dissociation. The results are shown in Table 10B.

Table 10B: Low RU numbers indicate no significant binding to unrelated charged proteins

DLL3/CD3 Binding Protein	Non-specific binding (RU)	
	Lysozyme (positive)	Tryp. Inhibitor (negative)
DLL3#1/CD3#1	4	1
DLL3#2/CD3#1	7	2
DLL3#3/CD3#1	54	85
DLL3#3/CD3#2	44	36
DLL3#3/CD3#3	17	28
DLL3#4/CD3#1	16	37
DLL3#5/CD3#1	4	0
DLL3#6/CD3#1	6	0
DLL3#7/CD3#1	9	3
DLL3#8/CD3#1	39	18
DLL3#9/CD3#1	15	27
DLL3#10/CD3#1	25	12
DLL3#11/CD3#1	11	13
DLL3#12/CD3#1	8	9
DLL3#13/CD3#1	11	3
DLL3#14/CD3#1	18	6
DLL3#15/CD3#1	na	na
DLL3#16/CD3#1	6	0
DLL3#17/CD3#1	15	8
DLL3#18/CD3#1	23	81

Example 18: Induction of lysis, T cell activation, T cell degranulation, T cell proliferation and cytokine secretion in presence of DLL3-positive and DLL3-negative tumor cells

PBMCs were purified as described in Example 9. To determine T cell activation, T cell degranulation and cytokine secretion, a cytotoxicity assay with PBMCs and DLL3-positive SHP77 or DLL3-negative RKO-E6 cells as target cells was setup as described in Example 10B. Potency of cell lysis by redirecting T cells towards human SHP77 or RKO-E6 cells was determined as described in Example 10A. To determine T cell activation, and T cell degranulation cells were centrifuged and stained with antibodies against CD4 (BD #550630), CD8 (BD #557834), CD25 (BD#340907), subsequently the cells were permabilized using the Fixation/Permeabilization Solution (BD #554714) and stained with antibodies against Perforin (BioLegend #308120) and Granzyme B (BD #560221) and measured by flow-cytometry. Cytokine levels in supernatants were determined by U-
 PLEX Biomarker Group 1 (hu) Assays (MSD, #K15067L-2, Kit).
 To determine the proliferation of T cells, PBMCs were labeled with 5 μ M Cell Trace™ CFSE (Invitrogen, C34554) and T cell stained with an anti-CD3 antibody (BioLegend cat#: 317336). Subsequently the labeled PBMCs were incubated with SHP77 or RKO-E6 cells at a ratio of 10:1 and increasing concentrations of an DLL3/CD3 binding protein for 6 days.
 Figures 11-15 show induction of T cell redirected lysis of SHP77 or RKO-E6 cells (Figure 11) and induction of dose-dependent T cell activation (Figure 12), T cell degranulation (Figure 13), secretion of cytokines (Figure 14A: Interferon gamma; 14B: MCP-1), and proliferation of T cells (Figure 15) in presence of SHP77 or RO-E6 cells and an DLL3/CD3 binding protein.

Example 19: Redirection of T cell subsets to SHP77 cells

PBMCs were isolated as described in Example 10a. Subsequently T cell subsets were isolated using the following reagents and protocols:

Table 11: Reagents and protocols for isolation of T cell subsets

T cell subset	Used reagents and protocols
Naïve T cells	Miltenyi Biotech; #130-097-095
CD4 ⁺ T cells	Miltenyi Biotech; #130-096-533

T cell subset	Used reagents and protocols
CD4 ⁺ Effector Memory T cells	Miltenyi Biotech; #130-094-125
CD4 ⁺ Central Memory T cells	Miltenyi Biotech; #130-094-302
CD8 ⁺ T cells	Miltenyi Biotech; #130-096-495
CD8 ⁺ CD45RA ⁺ Effector T cells	Miltenyi Biotech; #130-094-485
CD8 ⁺ Memory T cells	Miltenyi Biotech; #130-094-412

Figures 16-20 shows potency of redirecting of Pan T cells (Figure 16), naïve T cells (Figure 16), CD4⁺ T cells (Figures 17, 18), CD4⁺ Effector Memory T cells (Figure 18), CD4⁺ Central Memory T cells (Figure 17), CD8⁺ T cells (Figures 19, 20), CD8⁺CD45RA⁺ Effector T cells (Figure 19), CD8⁺ Memory T cells (Figure 20) against human SHP77 cells of an DLL3/CD3 binding protein.

Example 20: T cell infiltration in SHP77 xenograft tumor tissue with an exemplary DLL3/CD3 binding protein.

Remaining tumor tissues from mice in the study described in Example 13 were prepared, fixed in formalin and embedded in paraffin. Subsequently tissue sections were prepared and stained for CD3 expression on T cells (2GV6 Ventana). T cell infiltration in SHP77 xenograft tumor tissue with an exemplary DLL3/CD3 binding protein is shown in Figure 21. The scoring in Table 12 was used to quantify CD3 expression in xenograft tumor tissues.

Table 12: Scoring for quantification of infiltrating CD3-positive T cells

Score	Description
0	no CD3 ⁺ cells in plain of section
1	rare scattered cells or small clusters, mostly in the stroma
2	rare scattered cells or small clusters, in the stroma and margin
3	low cell numbers or clusters in stroma, epithelium and margin
4	moderate cell numbers or clusters in stroma and margin; low numbers of intra-epithelial cells
5	high cell numbers or clusters in stroma and margin; low numbers of intra-epithelial cells

Example 21: Production of anti-DLL3 antibodies

To obtain anti-DLL3 binders, hybridomas or single B cells derived from DLL3 immunized wild-type and AlivaMab humanized mice (Ablexis, San Francisco, CA, USA: AlivaMab transgenic mouse platform with human immunoglobulin loci) were cultured *in vitro*.

- 5 Supernatants were screened for reactivity against recombinant human DLL3, by AlphaLISA (PerkinElmer, Waltham, MA, USA), and against SHP-77 cells (ATCC®, CRL-2195TM) expressing human DLL3, by Flow Cytometry.

- Immunoglobulin (Ig) VH and VL genes were then amplified from identified positive clones. To isolate RNA from hybridomas, about 2×10^6 cells from single clones were
10 pelleted and used as source material. For single B cells, 100 to 500 cells expanded from singularly isolated B cells were used as source material. RNA was isolated using RNeasy Plus (Qiagen, Hilden, Germany). cDNA was then synthesized using Smarter cDNA synthesis kit (Clontech, Mountain View, CA) according to manufacturer's instructions. To facilitate cDNA synthesis, oligodT was used to prime reverse transcription of all
15 messenger RNAs followed by "5' capping" with a Smarter IIA oligonucleotide. Subsequent amplification of the VH and VL fragments was performed using a 2-step PCR amplification using 5' primers targeting the Smarter IIA cap and 3' primers targeting consensus regions in CH1. Briefly, each 50µl PCR reaction consists of 20µM of forward and reverse primer mixes, 25µl of PrimeStar Max DNA polymerase premix (Clontech), 2µl
20 of unpurified cDNA, and 21µl of double-distilled H₂O. The cycling program starts at 94 °C for 3 min, followed by 35 cycles (94 °C for 30 Sec, 50 °C for 1 min, 68 °C for 1 min), and ends at 72 °C for 7 min. The second round PCR was performed with VL and VH 2nd round primers containing 15bp complementary extensions that "overlap" respective regions in their respective pTT5 mother vector (VH and VL). Second round PCR was
25 performed with the following program: 94 °C for 3 min; 35 cycles (94 °C for 30 Sec, 50 °C for 1 min, 68 °C for 1 min), and ends at 72 °C for 7 min.

- In-Fusion® HD Cloning Kit (Clontech, U.S.A.) was used for directional cloning of VL gene into pTT5 huIgK vector and VH gene into pTT5 huIgG1KO vector. To facilitate In-Fusion® HD Cloning, PCR products were purified and treated with Cloning Enhancer
30 before In-Fusion HD Cloning. Cloning and transformation were performed according to manufacturer's protocol (Clontech, U.S.A.). Mini-prep DNAs were subjected to Sanger sequencing to confirm that complete V-gene fragments were obtained.

Using this methodology, pairs of Ig VH and VL genes encoding binding domains with specificity for DLL3 were prepared. Recombinant antibodies were produced by transient transfection of CHO-E37 cells with the corresponding heavy and light chain-encoding plasmids.

5

Confirmatory screening of recombinant antibodies

Supernatants containing expressed recombinant antibodies were assayed by flow cytometry for binding to cell lines expressing human DLL3. Briefly, cells were incubated with recombinant supernatants, washed, and bound mAbs from the supernatants were
10 detected with anti-human-IgG-APC (Jackson ImmunoResearch 109-136-098). Signal-to-background ratios (S/B) were calculated by dividing the median fluorescence intensity (MFI) of the sample by that of isotype control.

Surface Plasmon Resonance (SPR) on Biacore 400 was performed on recombinant supernatants. Briefly, the non-optimized IgGs in the HTP supernatants were captured via
15 Protein A/G onto the sensor surface for 60 sec at 10 ul/min. Binding of 100 nM human DLL3 to the captured IgGs was monitored for 180 sec of association at 30 ul/min, followed by 120 sec of dissociation in the HBS-EP buffer. Regeneration of the Protein A/G surface was performed with Glycine pH 2.1 in between each binding cycle. The following materials were used in this assay: Protein reagent: recombinantly expressed human DLL3.
20 System running buffer: HBS-EP (10 mM HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA, and 0.005% v/v polysorbate P20). Capturing reagent: Protein A/G, with specificity towards all human IgG isotypes.

Clones of interest (with $K_d < 300$ pM) were selected and further evaluated in various detection assays as described below.

25

Example 22: Binding of anti-DLL3 antibodies to recombinant human DLL3 protein

Binding of antibodies was performed in an ELISA. In brief, anti-DLL3 antibodies were coated at a concentration of 2 µg/ml at 4°C over night. After washing the plates were blocked with blocking buffer (Gibco, Gibco#043-90309A) for one hour at room
30 temperature, followed by washing and incubation with recombinant DLL3 protein at increasing concentrations. For detection the bound DLL3 protein was incubated with a polyclonal anti-DLL3 antibody (R&D Systems, AB4315) followed by a SULFO-TAG labelled anti-goat antibody (R32AG-1) for one hour. The bound antibody was quantified

by using a Read Buffer (MSD) in a Sector imager 6000 (MSD). Figure 22 shows binding to three exemplary anti-DLL3 antibodies to recombinant protein. Anti-DLL3 antibody targeting the membrane proximal peptide domain (DLL3#3) shows stronger binding to recombinant DLL3 protein, compared to anti-DLL3 antibodies which target the EGF4(DLL3#4) or EGF1 domain (DLL3#5).

Example 23: Binding of anti-DLL3 antibodies to SCLC cell lines

Cells (T cells or human SCLC cells) were stained with increasing concentrations of anti-DLL3 antibody with increasing concentrations in FACS buffer (PBS/0.5%BSA/0.05% sodium azide). Bound molecules were detected with PE-conjugated anti-mouse secondary antibody (Jackson Immuno Research, 115-116-072) by FACS analysis. Anti-DLL3 antibodies targeting DSL (DLL3#6), EGF1 (DLL3#5) or EGF4 (DLL3#4) domains show stronger binding to SCLC cell lines expressing DLL3, compared to anti-DLL3 antibodies which target the C-terminal peptide (DLL3#1, DLL3#2, DLL3#3) as shown in Figure 23.

Example 24: (IHC)

Anti-DLL3 antibodies were tested on a DLL3 mRNA positive SCLC sample at 20µg/ml concentration using Ventana Discovery ultra (Ventana Medical Systems, Inc, Arizona) following manufacturer's instructions. In brief, IHC was performed on formalin fixed paraffin embedded tissue and sections (4 µm) where stained with anti-DLL3 antibodies. IHC assay was performed on Ventana Discovery Ultra platform using RUO Discovery Universal protocol. Anti-DLL3 antibody staining was optimised using Proteinase K as Epitope Retrieval for 12 minutes. Anti-DLL3 antibody was incubated for 60 minutes, followed by anti-mouse HQ detection system and anti HQ HRP both for 12 minutes. Negative controls (IgG antibody) were performed for each tissue section, treated identically to the test slides.

The human SCLC cell line SHP77 were used as positive control. Digital images of whole-tissue sections were acquired using a Leica SCN400 histology scanner (Leica Microsystems, Milton Keynes, UK).

Immunohistochemistry analysis of a human SCLC sample with exemplary anti-DLL3 antibodies is shown in Figure 24.

Example 25: Determination of expression levels on SCLC cell lines

Cell surface expression of anti-DLL3 antibodies was quantified using the QIFIKIT (K0078; Dako). In brief, SCLC cell lines (SHP77, NCI-H82 and NCI-H2286) were labeled with the anti-DLL3 antibody (DL309, WO 2011/093097) or irrelevant antibody (isotype control) at 5 µg/ml. In a separate vial. Subsequently cells and beads provided in the kit were labeled in parallel with fluorescein-conjugated anti-mouse secondary antibodies. Samples were measured in a BD CantoII Fluorocytometer and consequently, the fluorescence is correlated with the number of bound anti-DLL3 antibodies on the cells and on the beads. Table 12 shows DLL3 molecules expressed on the cell surface of three SCLC cell lines.

Table 13: DLL3 molecules expressed on the cell surface of SCLC cell lines

SCLC CELL LINE	Cell surface DLL3 molecules/cell
SHP77 (ATCC®, CRL-2195™)	2061
NCI-H82 (HTB-175™)	824
NCI-H2286	129

Example 26: Determination of sensitivity of IHC protocol on SCLC cell line blocks

To evaluate the sensitivity of the IHC protocol, SCLC cell lines (SHP77, NCI-H82 and NCI-H2286) were processed like tissues in hospitals and fixed in Formalin and subsequently embedded in paraffin.

SCLC cell lines were cultured according to ATCC instructions. Cells were scraped from the plate and fixed in Formalin and subsequently added to solubilized Histogel (Thermo Fisher scientific, HG-4000-012) and incubated over night at 4°C, followed by a standard paraffin embedding procedure performed in routine pathology laboratory. In brief, cell block is incubated in Ethanol and Isopropanol before embedding in Paraffin (roti-Plast, #6642.5). Sections of the cell pellet blocks were performed and sections were stained with the protocol as described in Example 24. The results shown in Figure 25 show that an IHC protocol with anti-DLL3 antibody DLL3#5 is able to detect cells with very low DLL3 expression.

CLAIMS

1. A protein comprising a first antigen binding unit specifically binding to DLL3 and a second antigen binding unit specifically binding to CD3, wherein said first antigen binding unit specifically binding to DLL3 is selected from the group consisting of i) to xviii):
- 5
- i) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3);
- 10
- ii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3);
- 15
- iii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3);
- 20
- iv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3);
- 25
- v) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1), SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3);
- 30
- vi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:31 (CDR1), SEQ ID NO:32 (CDR2) and SEQ

ID NO:33 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1), SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3);

- 5 vii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:133 (CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:136 (CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3);
- 10 viii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:139 (CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:142 (CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3);
- 15 ix) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:145 (CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:148 (CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3);
- 20 x) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:151 (CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:154 (CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3);
- 25 xi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:157 (CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:160 (CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3);
- 30 xii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:163 (CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy chain CDRs comprising the amino acid

sequences of SEQ ID NO:166 (CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3);

xiii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3);

xiv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3);

xv) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3);

xvi) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3);

xvii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194 (CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3); and

xviii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3)..

2. The protein of claim 1, wherein said first antigen binding unit specifically binding to DLL3 comprises a first light chain variable domain and a first heavy chain variable domain and is selected from the group consisting of i) to xviii):

- 5 i) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:37 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:38;
- ii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequences of SEQ ID NO:39 and heavy chain variable domain
10 comprising the amino acid sequences of SEQ ID NO:40;
- iii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:42;
- iv) an antigen binding unit comprising a light chain variable domain comprising
15 the amino acid sequence of SEQ ID NO:43 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:44;
- v) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:45 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:46;
- 20 vi) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:47 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:48;
- vii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:205 and heavy chain variable domain
25 comprising the amino acid sequence of SEQ ID NO:206;
- viii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:207 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:208;
- ix) an antigen binding unit comprising a light chain variable domain comprising
30 the amino acid sequence of SEQ ID NO:209 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:210;

- x) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:211 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:212;
- 5 xi) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:213 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:214;
- xii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:215 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:216;
- 10 xiii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:217 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:218;
- xiv) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:219 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:220;
- 15 xv) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:221 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:222;
- xvi) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:223 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:224;
- 20 xvii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:225 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:226; and
- 25 xviii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:227 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:228.
3. The protein of claims 1 or 2, wherein said second antigen binding unit specifically
- 30 binding to CD3 is selected from the group consisting of i)-iii):

- i) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3);
- 5 ii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3); and
- 10 iii) an antigen binding unit comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:96 (CDR1), SEQ ID NO:97 (CDR2) and SEQ ID NO:98 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:99 (CDR1), SEQ ID NO:100 (CDR2) and SEQ ID NO:101 (CDR3).
- 15 4. The protein of any one of claims 1 to 3, wherein said second antigen binding unit specifically binding to CD3 comprises a second light chain variable domain and a second heavy chain variable domain selected from the group consisting of i) to iii):
- i) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain
- 20 comprising the amino acid sequence of SEQ ID NO:68;
- ii) an antigen binding unit comprising a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70; and
- iii) an antigen binding unit comprising a light chain variable domain comprising the
- 25 amino acid sequence of SEQ ID NO:102 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:103.
5. The protein of any one of claims 1 to 4, wherein
- i) said first antigen binding unit specifically binding to DLL3 comprises from its N-
- 30 to C-terminus a first light chain variable domain, a first light chain constant domain, a first peptide linker, a first heavy chain variable domain and a first heavy chain constant CH1 domain; and

- ii) said second antigen binding unit specifically binding to CD3 comprises from its N- to C-terminus a second light chain variable domain, a second light chain constant domain, a second peptide linker, a second heavy chain variable domain and a second heavy chain constant CH1 domain.

5

6. The protein of claim 5, wherein said first and/or second peptide linker comprises 26 to 42 amino acids, preferably any one of 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, more preferably 38 amino acids.

- 10 7. The protein of claims 5 or 6, wherein said first linker and/or second linker is a Gly-Ser linker, preferably comprising the amino acid sequence of SEQ ID NO:89, more preferably said first and second peptide linker comprise the same sequence.

8. The protein of any one of claims 5 to 7 further comprising a first and a second Fc domain, said first Fc domain covalently linked to said first antigen binding unit, and a
15 said second Fc domain covalently linked to said second antigen binding unit.

9. The protein of claim 8, wherein

- i) said first Fc domain comprises a tyrosine (Y) at position 366 [T366Y], and said
20 second Fc domain comprises a threonine (T) at position 407 [Y407T], or
- ii) said first Fc domain comprises a tryptophan (W) at position 366 [T366W], and said second Fc domain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V], or
- iii) said second Fc domain comprises a tyrosine (Y) at position 366 [T366Y], and
25 said first Fc domain comprises a threonine (T) at position 407 [Y407T], or
- iv) said second Fc domain comprises a tryptophan (W) at position 366 [T366W], and said first Fc domain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V],

- preferably wherein said first or said second Fc domain further comprises an arginine at
30 position 435 [H435R] and a phenylalanine at position 436 [Y436F].

10. The protein of claim 9, wherein said first and/or said second Fc domain further comprises an alanine at position 234[L234A] and at position 235 [L235A].
11. The protein of any one of claims 5 to 10, wherein the first light chain constant domain and the second light chain constant domain comprise a human kappa or lambda domain.
12. A protein comprising a first polypeptide chain specifically binding to DLL3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:241, SEQ ID NO: 242, SEQ ID NO:243, SEQ ID NO:244, SEQ ID NO: 245, SEQ ID NO: 246, SEQ ID NO: 247, SEQ ID NO: 248, SEQ ID NO: 249, SEQ ID NO: 250, SEQ ID NO: 251, and SEQ ID NO: 252 and a second polypeptide chain specifically binding to CD3 comprising the amino acid sequence of SEQ ID NO:79, SEQ ID NO:80 or SEQ ID NO:105.
13. An isolated nucleic acid molecule encoding the first antigen binding unit and/or the second antigen binding unit of a protein of any one of claims 1 to 5, optionally further encoding a first and/or a second Fc domain.
14. An isolated nucleic acid molecule encoding the first and/or the second polypeptide chain of claim 12.
15. An expression vector comprising the nucleic acid molecule of claim 13 or 14.
16. A host cell transfected with the expression vector of claim 15.
17. A method of manufacturing a protein of any one of claims 1 to 12 comprising
- i) cultivating the host cell of claim 16 under conditions allowing expression of the protein of any one of claims 1 to 12; and,
 - ii) recovering the protein; and optionally
 - iii) further purifying and/or modifying and/or formulating the protein.

18. A protein comprising a first polypeptide chain specifically binding to DLL3 and a second polypeptide chain specifically binding to CD3, wherein said first polypeptide chain comprises a first light chain, a first linker, and a first heavy chain and said second polypeptide chain comprises a second light chain, a second linker, and a second heavy chain, preferably wherein the C-terminus of said first light chain is covalently linked to the N-terminus of said first heavy chain via said first peptide linker and wherein the C-terminus of said second light chain is covalently linked to the N-terminus of said second heavy chain via said second peptide linker.
19. The protein of claim 18, wherein said first polypeptide chain specifically binding to DLL3 comprises a light chain variable domain and heavy chain variable domain comprising CDR sequences selected from the group consisting of i) to xviii):
- i) light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and a heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3);
 - ii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3);
 - iii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3);
 - iv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3);
 - v) light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy

chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1),
SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3);

vi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:31
(CDR1), SEQ ID NO:32 (CDR2) and SEQ ID NO:33 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1),
SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3)

vii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:133
(CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:136
(CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3);

viii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:139
(CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:142
(CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3);

ix) light chain CDRs comprising the amino acid sequences of SEQ ID NO:145
(CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:148
(CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3);

x) light chain CDRs comprising the amino acid sequences of SEQ ID NO:151
(CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:154
(CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3);

xi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:157
(CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:160
(CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3);

xii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:163
(CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy
chain CDRs comprising the amino acid sequences of SEQ ID NO:166
(CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3);

- xiii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3);

5 light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3);
- xiv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3);

10
- xv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3);

15
- xvi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194 (CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3); and

20
- xvii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3).

25

20. The protein of claim 18 or 19, wherein said second polypeptide chain specifically binding to CD3 comprises a light chain variable domain and heavy chain variable domain comprising CDR sequences selected from the group consisting of i) to iii):

- i) light chain CDRs comprising the amino acid sequences of SEQ ID NO:55 (CDR1), SEQ ID NO:56 (CDR2) and SEQ ID NO:57 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:58 (CDR1), SEQ ID NO:59 (CDR2) and SEQ ID NO:60 (CDR3);

30

- ii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:61 (CDR1), SEQ ID NO:62 (CDR2) and SEQ ID NO:63 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:64 (CDR1), SEQ ID NO:65 (CDR2) and SEQ ID NO:66 (CDR3); and
- 5 iii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:96 (CDR1), SEQ ID NO:97 (CDR2) and SEQ ID NO:98 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:99 (CDR1), SEQ ID NO:100 (CDR2) and SEQ ID NO:101 (CDR3).
- 10 21. The protein of any one of claims 18 to 20, wherein said first polypeptide chain specifically binding to DLL3 comprises a light chain variable domain and a heavy chain variable domain selected from the group consisting of i) to xviii):
- 15 i) a light chain variable domain comprising the amino acid sequences of SEQ ID NO:37 and heavy chain variable domain comprising the amino acid sequences of SEQ ID NO:38;
- 20 ii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:39 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:40;
- 20 iii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:41 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:42;
- 25 iv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:43 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:44;
- 25 v) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:45 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:46;
- 30 vi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:47 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:48;

- vii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:205 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:206;
- 5 viii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:207 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:208 ;
- ix) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:209 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:210;
- 10 x) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:211 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:212;
- xi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:213 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:214 ;
- 15 xii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:215 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:216;
- xiii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:217 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:218;
- 20 xiv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:219 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:220;
- xv) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:221 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:222 ;
- 25 xvi) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:223 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:224;
- 30

- xvii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:225 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:226; and
- xviii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:227 and heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:228.
22. The protein of any one of claims 19 to 22, wherein said second polypeptide chain specifically binding to CD3 comprises a light chain variable domain and a heavy chain variable domain selected from the group consisting of i)-iii):
- i) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:67 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:68;
- ii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:69 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:70; and
- iii) a light chain variable domain comprising the amino acid sequence of SEQ ID NO:102 and a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO:103.
23. The protein of any one of claims 18 to 22, wherein said first and/or second peptide linker comprises 26 to 42 amino acids, preferably any one of 30 to 40 amino acids, 34 to 40 amino acids, or 36 to 39 amino acids, more preferably 38 amino acids.
24. The protein of claim 23, wherein said first linker and/or second linker is a Gly-Ser linker, preferably comprising the amino acid sequence of SEQ ID NO:89, more preferably said first and second peptide linker comprise the same sequence.
25. The protein of any one of claims 18 to 24, wherein
- i) said first heavy chain comprises a tyrosine (Y) at position 366 [T366Y], and said second heavy chain comprises a threonine (T) at position 407 [Y407T], or

- ii) said first heavy chain comprises a tryptophan (W) at position 366 [T366W], and said second heavy chain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V], or
- iii) said second heavy chain comprises a tyrosine (Y) at position 366 [T366Y], and said first heavy chain comprises a threonine (T) at position 407 [Y407T], or
- iv) said second heavy chain comprises a tryptophan (W) at position 366 [T366W], and said first heavy chain comprises a serine (S) at position 366 [T366S], an alanine (A) at position 368 [L368A] and a valine (V) at position 407 [Y407V], preferably wherein said first or said second heavy chain further comprises an arginine at position 435 [H435R] and a phenylalanine at position 436 [Y436F].

26. The protein of claim 25, wherein said first and/or said second heavy chain further comprises an alanine at position 234[L234A] and at position 235 [L235A].
27. The protein of any one of claims 18 to 26, wherein said first light chain and said second light chain comprise a human kappa or lambda domain.
28. The protein of any one of claims 1 to 12 or 18 to 27 for use in medicine.
29. A pharmaceutical composition comprising the protein of any one of claims 1 to 12 or 18 to 27 and a pharmaceutically acceptable carrier.
30. A method of treating cancer comprising administering to a patient in need thereof a therapeutically effective amount of the protein of any one of claims 1 to 12 or 18 to 27.
31. Use of the protein of any one of claims 1 to 12 or 18 to 27 for preparing a pharmaceutical composition for treating cancer.
32. The protein of any one of claims 1 to 12 or 18 to 27 for use in a method of treating cancer.


33. The method of claim 30, the use of claim 31 or the protein for use of claim 32, wherein the cancer is a DLL3 expressing tumor, preferably wherein the cancer is SCLC, glioblastoma or a DLL3 expressing neuroendocrine tumor.
- 5 34. The method of claim 30 or 33, the use of claim 31 or 33, or the protein for use of claim 32 or 33, wherein said protein is used in combination with a chemotherapeutic agent, a therapeutically active compound that inhibits angiogenesis, a signal transduction pathway inhibitor, an EGFR inhibitor, an immune modulator, an immune checkpoint inhibitor, or a hormonal therapy agent.
- 10 35. The method, use or protein for use of claim 34, wherein said protein is in combination with an immune checkpoint inhibitor, preferably an anti- PD-1 antibody or an anti-PD-L1 antibody.
- 15 36. The method, use or protein for use of claim 35, wherein the anti-PD-1 antibody is selected from the group consisting of PD1-1, PD1-2, PD1-3, PD1-4, and PD1-5.
37. An anti-DLL3 antibody molecule comprising
- 20 i) light chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3); or
- 25 ii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:7 (CDR1), SEQ ID NO:8 (CDR2) and SEQ ID NO:9 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:10 (CDR1), SEQ ID NO:11 (CDR2) and SEQ ID NO:12 (CDR3); or
- 30 iii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:13 (CDR1), SEQ ID NO:14 (CDR2) and SEQ ID NO:15 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:16 (CDR1), SEQ ID NO:17 (CDR2) and SEQ ID NO:18 (CDR3); or
- iv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:19 (CDR1), SEQ ID NO:20 (CDR2) and SEQ ID NO:21 (CDR3) and heavy chain

- CDRs comprising the amino acid sequences of SEQ ID NO:22 (CDR1), SEQ ID NO:23 (CDR2) and SEQ ID NO:24 (CDR3); or
- 5 v) light chain CDRs comprising the amino acid sequences of SEQ ID NO:25 (CDR1), SEQ ID NO:26 (CDR2) and SEQ ID NO:27 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:28 (CDR1), SEQ ID NO:29 (CDR2) and SEQ ID NO:30 (CDR3); or
- 10 vi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:31 (CDR1), SEQ ID NO:32 (CDR2) and SEQ ID NO:33 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:34 (CDR1), SEQ ID NO:35 (CDR2) and SEQ ID NO:36 (CDR3); or
- vii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:133 (CDR1), SEQ ID NO:134 (CDR2) and SEQ ID NO:135 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:136 (CDR1), SEQ ID NO:137 (CDR2) and SEQ ID NO:138 (CDR3); or
- 15 viii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:139 (CDR1), SEQ ID NO:140 (CDR2) and SEQ ID NO:141 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:142 (CDR1), SEQ ID NO:143 (CDR2) and SEQ ID NO:144 (CDR3); or
- 20 ix) light chain CDRs comprising the amino acid sequences of SEQ ID NO:145 (CDR1), SEQ ID NO:146 (CDR2) and SEQ ID NO:147 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:148 (CDR1), SEQ ID NO:149 (CDR2) and SEQ ID NO:150 (CDR3); or
- x) light chain CDRs comprising the amino acid sequences of SEQ ID NO:151 (CDR1), SEQ ID NO:152 (CDR2) and SEQ ID NO:153 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:154 (CDR1), SEQ ID NO:155 (CDR2) and SEQ ID NO:156 (CDR3); or
- 25 xi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:157 (CDR1), SEQ ID NO:158 (CDR2) and SEQ ID NO:159 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:160 (CDR1), SEQ ID NO:161 (CDR2) and SEQ ID NO:162 (CDR3); or
- 30 xii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:163 (CDR1), SEQ ID NO:164 (CDR2) and SEQ ID NO:165 (CDR3) and heavy chain

- CDRs comprising the amino acid sequences of SEQ ID NO:166 (CDR1), SEQ ID NO:167 (CDR2) and SEQ ID NO:168 (CDR3); or
- xiii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:169 (CDR1), SEQ ID NO:170 (CDR2) and SEQ ID NO:171 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:172 (CDR1), SEQ ID NO:173 (CDR2) and SEQ ID NO:174 (CDR3); or
- xiv) comprising light chain CDRs comprising the amino acid sequences of SEQ ID NO:175 (CDR1), SEQ ID NO:176 (CDR2) and SEQ ID NO:177 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:178 (CDR1), SEQ ID NO:179 (CDR2) and SEQ ID NO:180 (CDR3); or
- xv) light chain CDRs comprising the amino acid sequences of SEQ ID NO:181 (CDR1), SEQ ID NO:182 (CDR2) and SEQ ID NO:183 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:184 (CDR1), SEQ ID NO:185 (CDR2) and SEQ ID NO:186 (CDR3); or
- xvi) light chain CDRs comprising the amino acid sequences of SEQ ID NO:187 (CDR1), SEQ ID NO:188 (CDR2) and SEQ ID NO:189 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:190 (CDR1), SEQ ID NO:191 (CDR2) and SEQ ID NO:192 (CDR3); or
- xvii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:193 (CDR1), SEQ ID NO:194 (CDR2) and SEQ ID NO:195 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:196 (CDR1), SEQ ID NO:197 (CDR2) and SEQ ID NO:198 (CDR3); or
- xviii) light chain CDRs comprising the amino acid sequences of SEQ ID NO:199 (CDR1), SEQ ID NO:200 (CDR2) and SEQ ID NO:201 (CDR3) and heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:202 (CDR1), SEQ ID NO:203 (CDR2) and SEQ ID NO:204 (CDR3).
38. The anti-DLL3 antibody molecule of claim 37 wherein said antibody molecule is a monoclonal, a chimeric, a humanized or a human antibody molecule.
39. The anti-DLL3 antibody molecule of claim 37 or 38 wherein said antibody molecule is a Fab, F(ab')₂, Fab', Fv or scFv.

40. The anti-DLL3 antibody molecule of any one of claims 37 to 39, wherein the heavy chain constant region is selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgM, IgA and IgE constant regions.
- 5 41. The anti-DLL3 antibody molecule of any one of claims 37 to 40, wherein the light chain constant region is kappa or lambda.
42. A method of detecting DLL3 in a biological sample comprising
(a) contacting the sample with an anti-DLL3 antibody of any one of claims 37 to
10 41;
(b) permitting formation of antibody-antigen complexes in the sample; and
(c) detecting the anti-DLL3 antibody.
43. The method of claim 42, wherein the anti-DLL3 antibody is detected by a
15 detectable signal, preferably said signal is detected by immunocytochemistry (ICC), immunohistochemistry (IHC), Western Blotting, Flow cytometry and/or ELISA.
44. The method of any of claims 42 or 43, wherein the biological sample is a tissue
20 sample, preferably a fixed tissue sample, more preferably a formalin-fixed and paraffin-embedded tissue sample.
45. The method of claim 44, wherein the anti-DLL3 antibody is detected by IHC.
- 25 46. The method of any one of claims 42 to 45, wherein the tissue sample is a tumor tissue sample, more preferably a tumor tissue sample isolated from a tumor selected from the group consisting of SCLC, LCNEC, glioma, glioblastoma, melanoma, or other neuroendocrine tumors.
- 30 47. A kit for detecting DLL3 comprising an antibody of any of claims 37 to 41, and instructions for use.

Fig. 1



02/63

Fig. 2




Fig. 3A






Fig. 3B



DLL3/CD3 Binding Proteins	Epitope Domain
DLL3#7/CD3#1	EGF6
DLL3#8/CD3#1	EGF4
DLL3#9/CD3#1	EGF3
DLL3#10/CD3#1	EGF1
DLL3#11/CD3#1	EGF1
DLL3#12/CD3#1	EGF1

Fig. 3C



DLL3/CD3 Binding Proteins	Epitope Domain
DLL3#6/CD3#1	DSL
DLL3#13/CD3#1	DSL
DLL3#14/CD3#1	DSL
DLL3#15/CD3#1	DSL
DLL3#16/CD3#1	Non-DSL – non-EGF1-6—non-membrane proximal peptide
DLL3#17/CD3#1	Non-DSL – non-EGF1-6—non-membrane proximal peptide
DLL3#18/CD3#1	Non-DSL – non-EGF1-6—non-membrane proximal peptide

Fig. 4




Fig. 5





Fig. 6A



09/63

Fig. 6B

non-DSL – non-EGF1-6–non-membrane proximal peptide




Fig. 6C




Fig. 6D




Fig. 6E




Fig. 6F





Fig. 6G



15/63

Fig. 6H




Fig. 6I

non-DSL – non-EGF1-6 – non-membrane proximal peptide





Fig. 6J



18/63

Fig. 6K




Fig. 6L




Fig. 6M




Fig. 6N




Fig. 6O




Fig. 6P

non-DSL – non-EGF1-6–non-membrane proximal peptide




Fig. 6Q




Fig. 6R




Fig. 6S




Fig. 6T






Fig. 6U



29/63


Fig. 6V

membrane proximal peptide



30/63

Fig. 7A



31/63

Fig. 7B

non-DSL – non-EGF1-6–non-membrane proximal peptide




Fig. 7 C




Fig. 7D




Fig. 7E




Fig. 7F




Fig. 7G




Fig. 7H




Fig. 7I

non-DSL – non-EGF1-6–non-membrane proximal peptide




Fig. 7J




Fig. 7K




Fig. 7L




Fig. 7M




Fig. 7N




Fig. 7O




Fig. 8






Fig. 9



47/63

Fig. 10



48/63

Fig. 11




Fig. 12




Fig. 13




Fig. 14A






Fig. 14B



53/63

Fig. 15



54/63

Fig. 16




Fig. 17




Fig. 18




Fig. 19





Fig. 20






Fig. 22




Fig. 23






Fig. 24


DLL3#1




DLL3#2



DLL3#4



DLL3#5



DLL3#6





Fig. 25



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2019/064942

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K16/28
ADD. A61K39/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2017/021349 A1 (AMGEN RES GMBH [DE]) 9 February 2017 (2017-02-09) cited in the application See claims, examples -----	1-47
A	WO 2013/126746 A2 (STEM CENTRX INC [US]) 29 August 2013 (2013-08-29) cited in the application See page 34, lines 28-43, figure 2, claims 1-28 -----	1-47
A	WO 2014/125273 A1 (OXFORD BIOTHERAPEUTICS LTD [GB]) 21 August 2014 (2014-08-21) cited in the application See claims -----	1-47



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

30 July 2019

Date of mailing of the international search report

06/08/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Nauche, Stéphane

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2019/064942

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2017021349	A1	09-02-2017	
		AU 2016302569 A1	07-12-2017
		BR 112018000475 A2	18-09-2018
		CA 2986848 A1	09-02-2017
		CL 2018000267 A1	05-10-2018
		CN 108271376 A	10-07-2018
		CO 2018000877 A2	19-04-2018
		CR 201800066 A	29-05-2018
		EA 201890336 A1	29-06-2018
		EP 3328889 A1	06-06-2018
		JP 2018527908 A	27-09-2018
		KR 20180033501 A	03-04-2018
		PE 11522018 A1	17-07-2018
		PH 12018500241 A1	13-08-2018
		TW 201708261 A	01-03-2017
		US 2017037130 A1	09-02-2017
		WO 2017021349 A1	09-02-2017
WO 2013126746	A2	29-08-2013	
		AU 2013203459 B2	03-11-2016
		AU 2016244333 A1	03-11-2016
		AU 2018282328 A1	17-01-2019
		BR 112014020826 A2	04-07-2017
		CA 2865404 A1	29-08-2013
		CL 2014002240 A1	15-05-2015
		CL 2016001616 A1	03-02-2017
		CN 104520324 A	15-04-2015
		CO 7111274 A2	10-11-2014
		CY 1119590 T1	07-03-2018
		DK 2817338 T3	23-10-2017
		EP 2817338 A2	31-12-2014
		EP 3093293 A1	16-11-2016
		EP 3093294 A1	16-11-2016
		EP 3095797 A1	23-11-2016
		ES 2645272 T3	04-12-2017
		HK 1200174 A1	31-07-2015
		HK 1231485 A1	22-12-2017
		HK 1231486 A1	22-12-2017
		HK 1231488 A1	22-12-2017
		HR P20171549 T1	17-11-2017
		HU E037021 T2	28-08-2018
		IL 234155 A	31-10-2018
		JP 6010241 B2	19-10-2016
		JP 6089047 B2	01-03-2017
		JP 2015509947 A	02-04-2015
		JP 2016098232 A	30-05-2016
		JP 2016098233 A	30-05-2016
		JP 2018162257 A	18-10-2018
		KR 20150003169 A	08-01-2015
		LT 2817338 T	10-11-2017
		ME 02928 B	20-04-2018
		NZ 628804 A	25-08-2017
		PE 00902015 A1	16-02-2015
		PE 20190658 A1	08-05-2019
		PH 12014501876 A1	17-11-2014
		PL 2817338 T3	29-12-2017
		PT 2817338 T	13-11-2017
		RU 2014138474 A	10-04-2016
		SG 10201806867V A	27-09-2018
		SG 11201405131X A	30-10-2014

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2019/064942

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		SI 2817338 T1	30-11-2017
		US 9133271 B1	15-09-2015
		US 9173959 B1	03-11-2015
		US 9334318 B1	10-05-2016
		US 9345784 B1	24-05-2016
		US 9352051 B1	31-05-2016
		US 9764042 B1	19-09-2017
		US 2014363455 A1	11-12-2014
		US 2014363826 A1	11-12-2014
		US 2014363887 A1	11-12-2014
		US 2014364590 A1	11-12-2014
		US 2014364593 A1	11-12-2014
		US 2014370037 A1	18-12-2014
		US 2015265724 A1	24-09-2015
		US 2016075779 A1	17-03-2016
		US 2016136296 A1	19-05-2016
		US 2016228571 A1	11-08-2016
		US 2016250348 A1	01-09-2016
		US 2016251431 A1	01-09-2016
		US 2017000901 A1	05-01-2017
		US 2017266312 A1	21-09-2017
		US 2017266313 A1	21-09-2017
		US 2017266315 A1	21-09-2017
		US 2017274094 A1	28-09-2017
		US 2017307622 A1	26-10-2017
		US 2017368195 A1	28-12-2017
		US 2018015178 A1	18-01-2018
		US 2018036421 A1	08-02-2018
		US 2018036422 A1	08-02-2018
		US 2019046656 A1	14-02-2019
		WO 2013126746 A2	29-08-2013

WO 2014125273	A1	21-08-2014	
		AU 2014217611 A1	06-08-2015
		CA 2900134 A1	21-08-2014
		CL 2015002223 A1	29-04-2016
		CN 105143266 A	09-12-2015
		EA 201500835 A1	29-02-2016
		EP 2956483 A1	23-12-2015
		JP 2016513094 A	12-05-2016
		KR 20150119164 A	23-10-2015
		PH 12015501744 A1	07-12-2015
		US 2016032006 A1	04-02-2016
		US 2018201691 A1	19-07-2018
		WO 2014125273 A1	21-08-2014
