



US011357846B2

(12) United States Patent
Fritzer et al.**(10) Patent No.: US 11,357,846 B2**
(45) Date of Patent: *Jun. 14, 2022**(54) METHOD OF PRODUCING
PHARMACEUTICAL COMPOSITIONS
COMPRISING IMMUNOGENIC
CHIKUNGUNYA VIRUS CHIKV-DELTA5NP3****(71) Applicant: Valneva SE, Nantes (FR)****(72) Inventors: Andrea Fritzer, Vienna (AT); Andreas
Meinke, Pressbaum (AT); Urban
Lundberg, Pressbaum (AT); Mario
Nebenführ, Vienna (AT); Jürgen
Heindl-Wruss, Vienna (AT); Robert
Schiegl, Siegenfeld (AT); Arnaud
Leon, Nantes (FR)****(73) Assignee: Valneva SE, Nantes (FR)****(*) Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.This patent is subject to a terminal dis-
claimer.**(21) Appl. No.: 17/407,499****(22) Filed: Aug. 20, 2021****(65) Prior Publication Data**

US 2022/0016230 A1 Jan. 20, 2022

Related U.S. Application Data**(63)** Continuation of application No. 16/641,012, filed as
application No. PCT/EP2018/075392 on Sep. 19,
2018.**(30) Foreign Application Priority Data**

Sep. 21, 2017 (EP) 17192374

(51) Int. Cl.**A61K 39/12** (2006.01)**A61K 9/00** (2006.01)**C12N 7/00** (2006.01)**A61K 9/19** (2006.01)**A61P 31/14** (2006.01)**C07K 14/005** (2006.01)**A61K 39/00** (2006.01)**(52) U.S. Cl.**CPC **A61K 39/12** (2013.01); **A61K 9/0019**
(2013.01); **A61K 9/19** (2013.01); **A61P 31/14**
(2018.01); **C07K 14/005** (2013.01); **C12N 7/00**
(2013.01); **A61K 2039/5254** (2013.01); **A61K**
2039/54 (2013.01); **C12N 2770/36121**
(2013.01); **C12N 2770/36122** (2013.01); **C12N**
2770/36123 (2013.01); **C12N 2770/36134**
(2013.01)**(58) Field of Classification Search**

None

See application file for complete search history.

(56) References Cited

U.S. PATENT DOCUMENTS

6,309,650 B1 10/2001 Kim et al.
7,871,814 B2 1/2011 Andino-Pavlovsky et al.
8,765,148 B2 7/2014 Wize et al.
8,865,184 B2 10/2014 Ella et al.
9,353,353 B2 5/2016 Nabel et al.
9,499,588 B2 11/2016 Mason et al.
10,086,061 B2 10/2018 Thomas et al.
10,537,630 B2 1/2020 Barbero Calzado et al.
10,660,950 B2 5/2020 Barbero Calzado et al.
11,207,397 B2 12/2021 Barbero Calzado et al.
2011/0171249 A1 7/2011 Frolov et al.
2012/0003266 A1 1/2012 Nabel et al.
2018/0362936 A1 12/2018 Barbero Calzado et al.
2018/0362937 A1 12/2018 Barbero Calzado et al.
2018/0369359 A1 12/2018 Barbero Calzado et al.
2018/0371027 A1 12/2018 Barbero Calzado et al.
2019/0008945 A1 1/2019 Barbero Calzado et al.
2020/0197506 A1 6/2020 Barbero Calzado et al.
2020/0368342 A1 11/2020 Barbero Calzado et al.
2021/0322534 A1* 10/2021 Fritzer A61K 9/0019

FOREIGN PATENT DOCUMENTS

CN 105749268 A 7/2016
WO WO 1999/011762 A1 3/1999
WO WO 2001/092552 A2 12/2001
WO WO 2010/062396 A2 6/2010
WO WO 2013/083726 A1 6/2013
WO WO 2016/145149 A1 9/2016
WO WO 2017/109225 A1 6/2017

OTHER PUBLICATIONS

Gardner et al. (PLOS Neglected Tropical Diseases. Feb. 2014; 8 (2):
e2719).
Edelman et al. The American journal of tropical medicine and
hygiene. 2000; 62 (6): 6: 681-685).
PCT/EP2016/082663, Apr. 19, 2017, International Search Report
and Written Opinion.
PCT/EP2016/082663, Jul. 5, 2018, International Preliminary Report
on Patentability.
PCT/EP2016/082662, Apr. 18, 2017, International Search Report
and Written Opinion.
PCT/EP2016/082662, Jul. 5, 2018, International Preliminary Report
on Patentability.
PCT/EP2018/075392, Apr. 2, 2020, International Search Report and
Written Opinion.
PCT/EP2018/075392, Nov. 20, 2018, International Preliminary
Report on Patentability.
[No Author Listed], Centers for Disease Control and Prevention.
Ingredients of vaccines fact sheet; continuously updated; [https://
www.cdc.gov/vaccines/vac-gen/additives.htm](https://www.cdc.gov/vaccines/vac-gen/additives.htm).
[No Author Listed], Japanese Encephalitis Vaccine. Centers for
Disease Control and Prevention, 2016. Retrieved from [https://www.
cdc.gov/japaneseencephalitis/vaccine/](https://www.cdc.gov/japaneseencephalitis/vaccine/) on Jun. 16, 2016.

(Continued)

Primary Examiner — Shanon A. Foley

(74) Attorney, Agent, or Firm — Wolf, Greenfield &
Sacks, P.C.**(57) ABSTRACT**The present invention relates to a process for producing an
immunogenic live attenuated Chikungunya virus, as well as
pharmaceutical compositions comprising the same.**17 Claims, 23 Drawing Sheets****Specification includes a Sequence Listing.**

(56)

References Cited

OTHER PUBLICATIONS

[No Author Listed], Pan-American Health Organization. 2015. Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory 2013-2014, Cumulative Cases (Updated Oct. 23, 2015).

[No Author Listed], Protamine sulfate. Wikimedia Foundation, Inc., 2015. Retrieved from https://en.wikipedia.org/wiki/Protamine_sulfate; updated Sep. 30, 2015 on Nov. 26, 2015.

[No Author Listed], Valneva Announces Successful Generation of a Highly-purified Zika Vaccine Candidate Using its FDA-EMA Approved Japanese Encephalitis Platform. Press release Jul. 7, 2016.

[No Author Listed], Valneva Reports Excellent Final Phase 1 Results for its Chikungunya Vaccine Candidate, Confirms Plans. Press release. Nov. 18, 2019.

[No Author Listed], World Health Organization, 2016 Zika Virus Fact Sheet 2016. Retrieved from <http://www.who.int/mediacentre/factsheets/zika/en/> on Mar. 11, 2016.

[No Author Listed], World Health Organization, 2016. Zika Situation Report Feb. 5, 2016.

[No Author Listed], Zika virus, strain H/PF/2013. European virus archive, 2016.

[No Author Listed], Valneva Reports Further Positive Results for Its Chikungunya Vaccine Candidate. Saint Herblain, France. May 22, 2019. 4 pages.

[No Author Listed], Valneva Reports Positive Phase 1 Interim Results for Its Chikungunya Vaccine Candidate. Saint Herblain, France. Jan. 7, 2019. 4 pages.

Abbink et al., Durability and correlates of vaccine protection against Zika virus in rhesus monkeys. *Sci. Transl. Med.* 2017;9:eaao4163.

Altschul et al., Basic Local Alignment Search Tool. *J. Mol. Biol.* 1990;215:403-410.

Altschul et al., Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nuc. Acids Res.* 1997;25(17):3389-3402.

Anez et al., Passage of dengue virus type 4 vaccine candidates in fetal rhesus lung cells selects heparin-sensitive variants that result in loss of infectivity and immunogenicity in rhesus macaques. *J. Virol.* Oct. 2009;83(20):10384-94. doi: 10.1128/JVI.01083-09. Epub Aug. 5, 2009.

Athmaram et al., A two step purification strategy for Chikungunya virions purification using sucrose buoyant density gradient separation. *J. Virology Res.* 2013;2(1):18-21.

Aubry et al., Inactivation of Zika virus in plasma with amotosalen and ultraviolet A illumination. *Transfusion.* Jan. 2016;56(1):33-40. doi: 10.1111/trf.13271. Epub Aug. 18, 2015.

Baronti et al., Complete Coding Sequence of Zika Virus from a French Polynesia Outbreak in 2013. *Genome Announc.* May-Jun. 2014; 2(3):e00500-14. Abstract.

Bender et al., Zika Virus Vaccine Candidate VLA1601: Cooperation Valneva & Emergent, Presentation at World Vaccine Congress Apr. 4, 2018.

Bender, Chikungunya Virus Vaccine Candidate Valneva's VLA1553. World Vaccine Conference 2019. Washington, D.C., Apr. 16, 2019. 43 pages.

Chroboczek et al., Virus-like particles as vaccine. *Acta Biochim Pol.* 2014;61(3):531-9. Epub Sep. 18, 2014.

Cohen, Infectious Disease. The race for a Zika vaccine is on. *Science.* Feb. 5, 2016;351(6273):543-4. doi: 10.1126/science.351.6273.543.

Cox et al., Predicting Zika virus structural biology: Challenges and opportunities for intervention. *Antivir Chem Chemother.* Aug. 2015;24(3-4):118-26. doi: 10.1177/2040206616653873. Epub Jun. 13, 2016.

Dowall et al., A susceptible mouse model for Zika virus infection. *PLOS Neglected Tropical Diseases.* 10(5):e0004658. May 5, 2016. DOI:10.1371/journal.pntd.0004658.

Eckels et al., Chikungunya virus vaccine prepared by Tween-ether extraction. *Appl Microbiol.* Feb. 1970;19(2):321-5.

Fritsche et al., Vaccine hypersensitivity—update and overview. *Swiss Med Wkly.* 2010;140(17-18):238-246.

Geradin et al., Chikungunya virus-associated encephalitis: A cohort study on La Réunion Island. 2005-2009. *Neurology.* 86(1):94-102.

Gubler et al., *Fields Virology*. Knipe DM, Howley PM, editors. Lippincott-Raven Publishers; Philadelphia: 2007:1153-1252.

Haddow et al., Genetic Characterization of Zika Virus Strains: Geographic Expansion of the Asian Lineage. *PLoS Negl Trop Dis* 6(2): e1477. doi:10.1371/journal.pntd.0001477.

Hallengard et al., Novel Attenuated Chikungunya Vaccine Candidates Elicit Protective Immunity in C57BL/6 mice. *J. Virology* 88(5):2858-2866.

Hallengard et al., Prime-Boost Immunization Strategies against Chikungunya Virus. *J. Virology.* 88(22):13333-13343.

Hombach et al., Report on a WHO consultation on immunological endpoints for evaluation of new Japanese encephalitis vaccines, WHO, Geneva, Sep. 2-3, 2004. *Vaccine.* 2005; 23(45):5205-5211.

Hutornjo et al., Comparison of ultracentrifugation methods for concentration of recombinant alphaviruses: sucrose and iodixanol cushions. *Environmental Experimental Biology.* 2012;10:117-123.

Katoh et al., Recent developments in the MAFFT multiple sequence alignment program. *Briefings in Bioinformatics.* 2008;9(4):286-298.

Kim et al., Design of Chimeric Alphaviruses with a Programmed, Attenuated, Cell Type-Restricted Phenotype. *J Virol.* 2011;85(9):4363-4376.

Konishi et al., Studies on structural proteins of Chikungunya Virus. I. Separation of three species of proteins and their preliminary characterization. *Microbiol Immunol.* 1980;24(5):419-28. doi: 10.1111/j.1348-0421.1980.tb02846.x.

Larkin et al., Clustal W and Clustal X version 2.0. *Bioinformatics.* 2007;23(21):2947-2948.

Larocca et al., Vaccine protection against Zika virus from Brazil. *Nature.* 2016;536:474-478. doi:10.1038/nature18952. Methods.

Lindenbach et al., *Fields Virology*. Knipe DM, Howley PM, editors. Lippincott-Raven Publishers; Philadelphia: 2007:1101-1152.

Malone et al., Zika Virus: Medical Countermeasure Development Challenges. *PLoS Negl Trop Dis.* 2016;10(3):e0004530. doi:10.1371/journal.pntd.0004530.

Modjarrad et al., Preliminary aggregate safety and immunogenicity results from three trials of a purified inactivated Zika virus vaccine candidate: phase 1, randomised, double-blind, placebo-controlled clinical trials. Dec. 4, 2017.

Monath, Yellow fever: an update. *Lancet Infect Dis.* 2001;1(1):11-20.

Needleman et al., A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Mol. Biol.* 1970;48(3):443-453.

Patkar et al., Yellow Fever virus NS3 plays an essential role in virus assembly independent of its known enzymatic functions. *J. Virol.* Apr. 2008;82(7):3342-52. doi: 10.1128/JVI.02447-07. Epub Jan. 16, 2008.

Pearson et al., Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA.* 1988;85(8):2444-8.

Pellerin, Walter Reed Scientists Test Zika Vaccine Candidate. U.S. Department of Defense. Jun. 9, 2016.

Pinto et al., A Temporal Role of Type I Interferon Signaling in CD8+ T Cell Maturation during Acute West Nile Virus Infection. *PLoS Pathog.* Dec. 2011;7(12):e1002407.

Plevka et al., Maturation of flaviviruses starts from one or more icosahedrally independent nucleation centres EMBO reports. 2011;12(6):602-606.

Powers et al., Re-emergence of Chikungunya and O'nyong-nyong viruses: evidence for distinct geographical lineages and distant evolutionary relationships. *J Gen Virol.* Feb. 2000;81(Pt 2):471-9. doi: 10.1099/0022-1317-81-2-471.

Reed et al., A simple method of estimating fifty percent endpoints. *American J Hygiene.* May 1938;27:493-497.

Rocha et al., Microcephaly: normality parameters and its determinants in northeastern Brazil: a multicentre prospective cohort study. *Bull World Health Organ, E-pub:* Feb. 8, 2016. doi:<http://dx.doi.org/10.2471/BLT.16.171215>.

(56)

References Cited

OTHER PUBLICATIONS

Roques et al., Attenuated and vectored vaccines protect nonhuman primates against Chikungunya virus. *JCI Insight*. Mar. 23, 2017;2(6):e83527. doi: 10.1172/jci.insight.83527.

Rozen-Gagnon et al., Alphavirus Mutator Variants Present Host-Specific Defects and Attenuation in Mammalian and Insect Models, *PLOS Pathogens*, 10(1):e1003877.

Schlegl et al., Influence of elemental impurities in aluminum hydroxide adjuvant on the stability of inactivated Japanese Encephalitis vaccine, *Ixiaro®*. *Vaccine*. 2015;33(44):5989-5996.

Shustov et al., Efficient, trans-complementing packaging systems for chimeric, pseudoinfectious dengue 2/yellow fever viruses. *Virology*. Apr. 25, 2010;400(1):8-17. doi: 10.1016/j.virol.2009.12.015.

Simizu et al., Structural Proteins of Chikungunya Virus, *J Virol*. 1984;51(1): 254-258.

Smith et al., Comparison of Biosequences. *Adv Appl Math*. 1981;2:482-489.

Srivastava et al., A purified inactivated Japanese encephalitis virus vaccine made in vero cells. *Vaccine*. 2001;19:4557-4565.

Tiwari et al., Assessment of immunogenic potential of Vero adapted formalin inactivated vaccine derived from novel ECSA genotype of Chikungunya virus. *Vaccine*. Apr. 21, 2009;27(18):2513-22. doi: 10.1016/j.vaccine.2009.02.062. Epub Feb. 27, 2009.

Vega-Rua et al., Chikungunya Virus Transmission Potential by Local Aedes Mosquitoes in the Americas and Europe. *PLOS Neglected Tropical Diseases*. 2015;9(5): e0003780.

Waterhouse et al., Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics*. 2009;25(9):1189-1191.

Way et al., Comparative Studies of some African Arboviruses in Cell Culture and in Mice, *J Gen. Virol*. 1976;30:123-130.

Weaver, Arrival of Chikungunya Virus in the New World: Prospects for Spread and Impact on Public Health. *PLoS Negl Trop Dis*. 2014;8(6):e2921. doi:10.1371/journal.pntd.0002921.

Garcia-Arriaza et al., A novel poxvirus-based vaccine, MVA-CHIKV, is highly immunogenic and protects mice against chikungunya infection. *J Virol*. Mar. 2014;88(6):3527-47. doi: 10.1128/JVI.03418-13. Epub Jan. 8, 2014.

Liljestrom et al., A new generation of animal cell expression vectors based on the Semliki Forest virus replicon. *Biotechnology (N Y)*. Dec. 1991;9(12):1356-61. doi: 10.1038/nbt1291-1356.

Pohjala et al., Inhibitors of alphavirus entry and replication identified with a stable Chikungunya replicon cell line and virus-based assays. *PLoS One*. 2011;6(12):e28923. doi: 10.1371/journal.pone.0028923. Epub Dec. 19, 2011. 13 pages.

* cited by examiner

FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4A





FIG. 4B



Plaque size, Immunogenicity

Virus titer

FIG. 4C




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 7




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 10

	Groups for subcutaneous immunization of C57Bl/6 mice				
	Group 1	Group 3	Group 5	Group 6	Group 7
Formulation ratio of P3 and P5B-07 (E168K)	P3:E168K 1:0.1	P3:E168K 1:1	P3:E168K 1:10	P3	P5B-7
Reference sequence	G	G	G	G	G
Heterogeneity	G>A	G=A	G<A	G>A	A
Chromatogram					

FIG. 11A




FIG. 11B




FIG. 11C





FIG. 11D

P3:E168K	Antigen dose/mouse (TCID ₅₀)	Single mouse sera: ++positive +low positive -negative
1:0.1	1.12E+04	7 ++, 2 +, 1 -
1:1	2.03E+04	7 ++, 3 -
1:10	1.12E+04	2 ++, 3 +, 5 -

FIG. 12



35°C

Log₁₀ (TCID₅₀)

Days post-infection

Legend:

- 0.1D2
- 0.1D4
- △ 0.1D5
- 0.01D2
- 0.01D4
- ▲ 0.01D5
- 0.001D2
- 0.001D4
- ▲ 0.001D5

Days post-infection	0.1D2	0.1D4	0.1D5	0.01D2	0.01D4	0.01D5	0.001D2	0.001D4	0.001D5
1	8.8	9.1	8.5	8.7	8.9	8.6	8.6	8.8	8.4
2	8.4	8.4	7.7	7.7	7.8	7.4	7.7	7.8	6.4
3	6.8	5.8	4.8	6.5	6.2	4.8	6.5	6.2	4.8

FIG. 13C





FIG. 14



Term	P-Value	P-value Prob.>F
A-Time of infection	4.75	0.0436
B-MOI	0.049	0.8269
C-Temperature	14.93	0.0012
AB	0.2	0.6583
BC	0.17	0.6815
AC	2.8	0.1126
A ²	3.33	0.0856
B ²	0.5	0.4895
C ²	20.27	0.0003

FIG. 15A




FIG. 15B

Term	F-value	p-value Prob>F
A-Time of infection	12.82	0.0023
B-MOI	3.72	0.0708
C-Temperature	67.85	<0.0001
AB	0.025	0.8752
BC	1.68	0.2118
AC	0.56	0.4653
A ²	5.29	0.0344
B ²	0.03	0.8653
C ²	10.94	0.0042

FIG. 16A




FIG. 16B

Pos. 9649 (Silent)

Term	F-Value	p-value Prob>F
A-MOI	32.19	0.0024
B-Temperature	10.16	0.0243
AB	8.74	0.0317

Pos. 9119 (E247K)

Term	F-Value	p-value Prob>F
A-MOI	2.42	0.1712
B-Temperature	51.72	0.0002

Pos. 8882 (E168K)

Term	F-Value	p-value Prob>F
A-MOI	17.65	0.0057
B-Temperature	2.5	0.1651

FIG. 17A






FIG. 17B

Pos. 9649 (Silent)

Term	F-Value	p-value Prob.>F
A-MOI	17.31	0.0253
B-Harvest Day	0.46	0.5456




Pos. 9119 (E247K)

Term	F-Value	p-value Prob.>F
A-MOI	27.0	0.0351
B-Harvest Day	32.0	0.0298
AB	27.0	0.0351

Pos. 8882 (E168K)

Term	F-Value	p-value Prob.>F
A-MOI	16.2	0.0276
B-Harvest Day	7.5	0.0714

FIG. 18A



○ Design Points

FIG. 18B

1

**METHOD OF PRODUCING
PHARMACEUTICAL COMPOSITIONS
COMPRISING IMMUNOGENIC
CHIKUNGUNYA VIRUS CHIKV-DELTA5NP3**

RELATED APPLICATIONS

This Application is a Continuation of U.S. application Ser. No. 16/641,012, filed Feb. 21, 2020, entitled "METHOD OF PRODUCING PHARMACEUTICAL COMPOSITIONS COMPRISING IMMUNOGENIC CHIKUNGUNYA VIRUS CHIKV-DELTA5NP3", which is a national stage filing under 35 U.S.C. 371 of International Patent Application Serial No. PCT/EP2018/075392, filed Sep. 19, 2018. The entire contents of these applications are incorporated herein by reference in their entirety.

**REFERENCE TO A SEQUENCE LISTING
SUBMITTED AS A TEXT FILE VIA EFS-WEB**

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Aug. 18, 2021, is named I042270131US01-SEQ-JRV, and is 45,568 bytes in size.

FIELD OF THE INVENTION

The invention relates to a process for producing an immunogenic live attenuated Chikungunya virus, as well as pharmaceutical compositions comprising the same.

BACKGROUND OF THE INVENTION

Chikungunya virus (CHIKV) is a positive-sense, single-stranded RNA virus from the genus Alphavirus, family Togaviridae. Chikungunya virus disease is mainly an outbreak disease and is associated with high attack rates. The virus is transmitted to humans via a mosquito vector and causes fever, rash, fatigue and severe polyarthralgia. Infections with CHIKV generally resolve spontaneously and are not usually fatal, except in rare cases involving CNS infection, where the death rate is between 10 to 30 percent. Particularly at risk for CHIKV CNS disease are infants under one year and adults over 65 years, with an infection rate 25-fold and 6-fold higher than the general population, respectively. The rate of persistent disabilities in children following CHIKV encephalitis is estimated at between 30 and 45 percent (Gerardin P, et al. Chikungunya virus-associated encephalitis A cohort study on La Réunion Island, 2005-2009 (2016) *Neurology* 86:1-9). Furthermore, about 30 percent of all Chikungunya patients experience arthralgia for months to years after recovery. In some cases, neurological, renal, cardiac, respiratory or hepatic complications can occur.

Currently no vaccines or medications are available for the prevention or treatment of Chikungunya virus disease. Outbreaks in the past have occurred mainly in Africa, but the East-Central South African (ECSA) genotype has recently expanded its geographical range, resulting in outbreaks in India, Asia, and even temperate Europe (Weaver, S., Arrival of Chikungunya Virus in the New World: Prospects for Spread and Impact on Public Health (2014) *PLOS Neglected Tropical Diseases* 8(6): e2921). Although CHIKV has been repeatedly imported into the Americas since 1995, autochthonous transmission had not been reported until 2013 in the Caribbean. By 2015, the epidemic had spread to the main-

2

land and caused upwards of one million suspected cases in 27 countries in the Americas (Pan-American Health Organization (2015) Number of Cumulative Cases of Chikungunya Fever in the Americas). Further epidemics may be aided in part by the spread of the CHIKV mosquito vector into non-endemic regions, as well as the ability of CHIKV to adapt to local mosquito species (Vega-Rua A, et al., Chikungunya Virus Transmission Potential by Local Aedes Mosquitoes in the Americas and Europe (2015) *PLOS Neglected Tropical Diseases* DOI:10.1371/journal.pntd.0003780). The high rate of contagion of Chikungunya virus, its geographical spread, and its potential for long-lasting complications underscore the need for developing preventative measures, such as vaccines.

Vaccines against Chikungunya virus may comprise live attenuated CHIKV particles; i.e., live CHIKV particles which have been altered to reduce virulence, but still maintain immunogenicity. One example of an attenuated CHIKV contains a deletion mutation in the non-structural protein 3 (CHIKV-Δ5nsP3; see FIG. 2). CHIKV-Δ5nsP3 has been shown to confer protective immunity in mice (Hallengård D, et al. Novel attenuated Chikungunya vaccine candidates elicit protective immunity in C57BL/6 mice (2014) *J. Virol.* 88:2858-2866) and non-human primates (Rogues P, et al. Attenuated and vectored vaccines protect non-human primates against Chikungunya virus (2017) *J. Clin. Invest.* Insight 2(6):e83527). These preliminary in vivo studies in mice and non-human primates were done with CHIKV-Δ5nsP3 virus produced on BHK-21 cells—a cell type generally not favored for the production of human vaccines. It would be necessary, therefore, as disclosed herein, to adapt CHIKV-Δ5nsP3 virus production to a more suitable cell culture platform. Such adaptation is not a trivial process; for example, it is known in the art that the adaptation of viruses to a particular host cell can lead to mutations that change the surface charge of the virus particles. Such acquired mutations can serve to attenuate viruses; in fact, serial passaging has been used to develop such attenuated virus particles that are used in many vaccines against viruses. With regard to CHIKV, it has been shown that repeated in vitro passaging of virulent wild-type Chikungunya virus can lead to certain point mutations, resulting in partial or complete attenuation of the virus (Gardner C L, et al., Deliberate Attenuation of Chikungunya Virus by Adaptation to Heparan Sulfate-Dependent Infectivity: A Model for Rational Arboviral Vaccine Design (2014) *PLOS Neglected Tropical Diseases* 8(2): e2719).

It has now surprisingly been found that certain point mutations resulting in loss of immunogenicity can occur early during the cell substrate adaptation process of the attenuated CHIKV-Δ5nsP3, making control or reduction of said point mutations an essential consideration for the production of a successful vaccine candidate. Thus, the current invention provides a process with well-defined parameters for the propagation of the CHIKV-Δ5nsP3 vaccine candidate, allowing production of highly immunogenic virus particles, while simultaneously achieving high production titers in cell culture suitable for industrial application.

SUMMARY OF THE INVENTION

The present invention relates to a pharmaceutical composition comprising a sufficient amount of immunogenic Chikungunya virus to elicit a neutralizing immune response in a subject; i.e., an immune response that is protective against infection with and/or disease caused by Chikungunya virus. In particular, the invention provides a pharma-

ceutical composition comprising live attenuated CHIKV-Δ5nsP3 particles wherein the percentage of said viral particles with immunogenicity-reducing mutations, particularly immunogenicity-reducing mutations in the E2 protein, are minimized. The disclosure further provides a process for producing a pharmaceutical composition comprising a live attenuated CHIKV-Δ5nsP3, wherein the process minimizes the presence of immunogenicity-reducing mutations in the viral genome, particularly mutations at E168 of viral E2 protein and/or other E2 residues and/or residues in other structural or non-structural CHIKV proteins. The current disclosure further provides pharmaceutical compositions comprising an immunogenic live attenuated Chikungunya virus obtainable by the process of the invention.

Efforts to develop a vaccine against Chikungunya virus are currently underway. One of the most advanced vaccine candidates provides a chimeric construct in a measles virus platform (see themisbio.com). The vaccine, currently in Phase 2 trials, is delivered in two doses (clinicaltrials.gov ID: NCT02861586). A one-shot vaccine would represent a distinct advantage in the field.

Accordingly, in one embodiment, it is an object of the current invention to provide a stable, well-defined, safe and effective pharmaceutical composition such as, e.g. a vaccine, against Chikungunya virus, preferably an improved vaccine conferring protection with only one vaccination; i.e., a so called "one-shot" vaccine at industrial scale using common cell substrates, to provide processes to generate such a stable, well-defined, safe and effective vaccine and methods and uses for said stable, well-defined safe and effective vaccine.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings are not intended to be drawn to scale. The Figures are illustrative only and are not required for enablement of the disclosure. For purposes of clarity, not every component may be labeled in every drawing. In the drawings:

FIG. 1 Map of pMX plasmid used for full assembly of the CHIKV-Δ5nsP3 genome from synthesized fragments. For the full assembly of the CHIKV-Δ5nsP3 sequence, the pMX plasmid with an Ampicillin resistance cassette (pMA) was used. The pMX vector series is based on pUC-like cloning vectors, but leaves out unnecessary promoters (biosafety level 1).

FIG. 2 Schematic illustration of the CHIKV-Δ5nsP3 genome structure. The Chikungunya virus genome encodes two polyproteins: non-structural proteins 1-4 (nsP1-4) and structural proteins (C, E3, E2, 6K, E1). Compared with the wild-type genomic sequence, the CHIKV-Δ5nsP3 sequence contains a 183-bp deletion in the 3' part of the sequence encoding nsP3 (amino acids 1656 to 1717 in the nsP1-4 polyprotein), which results in a 60 amino acid deletion in the nsP3 replicase protein (indicated by Δ60aa). SP, subgenomic promoter; UTR, untranslated region. (Figure adapted from Hallengård D, et al., 2014, supra.)

FIGS. 3A and 3B Cloning strategy for assembly of the CHIKV-Δ5nsP3 genome in pMA. (FIG. 3A) Design schematic of synthesized polynucleotide fragments covering the full CHIKV-Δ5nsP3 genome. (FIG. 3B) Cloning strategy for assembly of CHIKV-Δ5nsP3 genome in pMA plasmid. 1. Cloning of CHIKV-Δ5nsP3 fragment 2 into pMA containing fragment 1 (pMA fragment 1) via EcoRI and PacI. 2. Assembly of fragment 4 and fragment 3 via ClaI and PacI. 3. Preparation of fragment 5 digested with XhoI and PacI in pMA for final full assembly. 4. Full assembly of CHIKV-

Δ5nsP3 genome in pMA by fusion of AgeI/XhoI-digested fragments 3 and 4 and XhoI/PacI-digested fragment 5 with AgeI/PacI-linearized pMA fragments 1 and 2. Correct CHIKV-Δ5nsP3 genome assembly was verified via Sanger sequencing.

FIGS. 4A-4C Yield, plaque size and immunogenicity of CHIKV-Δ5nsP3 following passaging in Vero cells. Virus was passaged in three parallel replicates (A, B and C) starting from a common P0 (rescue) as detailed in Table 2. (FIG. 4A) Virus titers 24 h after infection of Vero cells from passage 0 to passage 16. The average titer of the three replicates (A, B and C) is shown. (FIG. 4B) Relative titers of P0, P5 and P15 CHIKV-Δ5nsP3 as assessed by plaque assay. Vero cells were seeded at a density of 4×10^5 cells per well in 6-well plates in MEM supplemented with 5% FBS, 2 mM L-Glutamine and 1% Antibiotic-Antimycotic (Anti-Anti) and were incubated overnight at 35° C. and 5% CO₂. On the next day, culture supernatant was removed from Vero cells and serial dilutions of CHIKV-Δ5nsP3 were added onto the cells. Following incubation for 1 hour at 35° C./5% CO₂, a methylcellulose overlay with a final concentration of 2% was added and cells were further incubated 3 days at 35° C./5% CO₂. Finally, plaques were counted after crystal violet staining (0.5% crystal violet in 5% formaldehyde) to assess virus titer (pfu/ml) and plaque morphologies. (FIG. 4C) Immunogenicity of P0, P5B, P8B and P15C CHIKV-Δ5nsP3 as assessed by neutralization of CHIKV-Δ5nsP3 (P0) in PRNT on Vero cells. Vero cells were seeded in 12-well plates at a density of 3×10^5 and incubated overnight at 35° C./5% CO₂. Groups of five C57Bl/6 mice were immunized once subcutaneously with a dose of 10^5 TCID₅₀ of the respective CHIKV-Δ5nsP3 passages. CHIKV-Δ5nsP3 at P0 (virus rescue), also at 10^5 TCID₅₀, was used as a positive control. Day 21 serum pools at 4-fold serial dilutions ranging from 1:20 to 1:327,680 were mixed with 560 pfu/ml CHIKV-Δ5nsP3 (at P0) and incubated for one hour. The CHIKV-Δ5nsP3/neutralization mixes were then added to the Vero cells and the plates were incubated for 2 hours at 35° C./5% CO₂. This step was followed by a 2% methylcellulose overlay and plates were incubated for ~60 hours at 35° C./5% CO₂. After removal of the overlay, cells were stained with crystal violet/5% formaldehyde and plaques were counted.

FIGS. 5A-5B Effect of controlled MOI (0.01) during passaging of CHIKV-Δ5nsP3 on immunogenicity. (FIG. 5A) Schematic illustration of CHIKV-Δ5nsP3 passaging on Vero cells under uncontrolled and controlled conditions. CHIKV-Δ5nsP3 P0 was passaged on Vero cells to P3B under uncontrolled conditions with varying MOI (as outlined in Table 2; Replicate B). Using P3B as a starting material, a controlled infection process was carried out with all subsequent infections at the defined MOI of 0.01 to generate one P4 passage, two P5 passages and one P6 passage for analysis of immunogenicity in mice. (FIG. 5B) Immunogenicity of P0 (○), P2B (□), P5 #1 (●), P5 #2 (◆), P6 (■) and P15 (Δ) CHIKV-Δ5nsP3 as assessed by neutralization of CHIKV-Δ5nsP3 (P2) in PRNT on Vero cells. Groups of ten C57Bl/6 mice were subcutaneously immunized with a single dose of the respective CHIKV-Δ5nsP3 preparations at an intended dose of 10^5 TCID₅₀ and at day 21 following immunization, pooled sera were assessed for CHIKV-Δ5nsP3 (560 pfu/ml) neutralization capacity at 4-fold serial dilutions ranging from 1:20 to 1:327,680 as described for FIGS. 4A-4C.

FIGS. 6A-6D Immunogenicity and observed genomic heterogeneities of single plaque isolates of CHIKV-Δ5nsP3 from P5B and P8B. Two CHIKV-Δ5nsP3 virus rescue harvests (4399gr1 P0 (#1) and 4415gr1 P0 (#2)) and a P15

5

CHIKV-Δ5nsP3 harvest (P15C-DS) served as immunogenic and non-immunogenic controls, respectively. (FIG. 6A) Immunogenicity of CHIKV-Δ5nsP3 single plaque isolates P5B-11, P5B-03 and P8B-05 as assessed by neutralization of CHIKV-Δ5nsP3 (P2) in PRNT on Vero cells. Groups of ten C57Bl/6 mice were subcutaneously immunized with a single dose of the respective CHIKV-Δ5nsP3 isolates at the indicated TCID₅₀ doses and at day 19 following immunization, pooled sera were assessed for CHIKV-Δ5nsP3 (560 pfu/ml) neutralization capacity at 4-fold serial dilutions ranging from 1:20 to 1:327,680 as described for FIGS. 4A-4C. (FIG. 6B) Schematic genomes of CHIKV-Δ5nsP3 single plaque isolates P5B-11, P5B-03 and P8B-05 derived from Sanger sequencing covering the full CHIKV-Δ5nsP3 genome with identified point mutations indicated. (FIG. 6C) Immunogenicity of further CHIKV-Δ5nsP3 single plaque isolates P5B-02, P5B-04, P5B-07 and P8B-01 was assessed as in (A). (FIG. 6D) Schematic genomes of CHIKV-Δ5nsP3 single plaque isolates P5B-02, P8B-01, P5B-04 and P5B-07 derived from Sanger sequencing covering the full CHIKV-Δ5nsP3 genome with identified point mutations indicated.

FIG. 7 Chromatograms from Sanger sequencing of CHIKV-Δ5nsP3 at passage 0 (P0) and four independently generated CHIKV-Δ5nsP3 samples at passage 3 (P3 Examples 1-4), revealing E168K and E247K heterogeneities in the E2 protein arising by passage 3. The four independently generated CHIKV-Δ5nsP3 samples at P3 (P3 Examples 1-4) were all derived from the same MVSB (P1). The regions shown cover genomic regions encoding amino acids 168 and 247 of the E2 protein and were sequenced with primer pairs 16 and 17 as indicated. (For primer sequences, see Table 1). For comparison, sequencing chromatograms of CHIKV-Δ5nsP3 at passage 0 (P0) are shown. Sites of sequence heterogeneities are indicated by boxes.

FIGS. 8A-8C Next generation sequencing (NGS) of a Master Virus Seed Bank (P1) and two independently generated passage three (P3) CHIKV-Δ5nsP3 preparations: (FIG. 8A) P1-MVSB; (FIG. 8B) P3-Example 3; (FIG. 8C) P3-Example 4. A background level of genomic heterogeneities is set at 15% as indicated by the dotted line. More frequent heterogeneities arising by passage 3 included point mutations at genomic nucleic acid positions 8882 and 9119, which correspond to E168K and E247K mutations, respectively, in the E2 protein as indicated in (FIG. 8B) and (FIG. 8C).

FIGS. 9A-9C Location of amino acid positions with frequently observed heterogeneities (G55R, G82R, E168K and E247K) within the E2 glycoprotein marked on equivalent positions on a cryo-EM structure of CHIKV VLPs of strain Senegal 37997 (PDB 3J2W; Sun S, et al., Structural analyses at pseudo atomic resolution of Chikungunya virus and antibodies show mechanisms of neutralization (2013) eLife 2:e00435. DOI: 10.7554/eLife.00435). G55R is indicated by a square, G82R is indicated by a circle, E168K is indicated by a triangle and E247K is indicated by a star. The figures show the CHIKV proteins of the unit-cell as they would be assembled in the viral membrane, composed of four copies of E1/E2 dimers with the transmembrane portion (TM) indicated by brackets, and the capsid protein (C) at the inner (bottom) side of the membrane. Three E1/E2/capsid-assemblies compose the trimeric q3-spike near the 5-fold axis and the fourth assembly is one of three elements of the i3-spike at the 3-fold axis, all in surface representation and viewed from three different angles: (FIG. 9A) CHIKV unit-cell side view; viewpoint above membrane plane. (FIG. 9B) CHIKV unit-cell side view; viewpoint below membrane plane (FIG. 9C) CHIKV unit-cell, top view.

6

FIG. 10 Groups for subcutaneous immunization of C57Bl/6 mice with virus preparations of different P3:P5B-07 (E168K) ratios to determine the threshold at which the presence of the E168K point mutation results in loss of CHIKV-Δ5nsP3 immunogenicity at a dose of approximately 3×10^4 TCID₅₀. The formulation ratios of P3 and P5B-07 (E168K mutant) for groups 1, 3 and 5 were 1:0.1, 1:1 and 1:10, respectively, with an intended dose of 3×10^4 TCID₅₀. Additionally, group 6 and group 7 represent P3 alone and P5B-07 (E168K) alone, respectively. The reference sequence (wild-type) in the heterogenic position (corresponding to nucleotide 8882) was G and depending on the ratio of either 1:0.1, 1:1 or 1:10, a shift towards the nucleotide A, i.e., G>A to G=A or G<A, was in accordance with sequencing chromatograms. The P3 viral population displayed ~20% E168K mutants. The P5B-07 (E168K) viral population did not show heterogeneity (i.e., ~100% contained the E2 protein E168K mutation).

FIGS. 11A-11D Effect on immunogenicity of different P3 to E168K-mutant CHIKV-Δ5nsP3 ratios assessed by VRP-based neutralization assay. Single mouse sera were generated by subcutaneous immunization of C57Bl/6 mice with an intended dose of 3×10^4 TCID₅₀ of CHIKV-Δ5nsP3 P3:E168K at ratios 1:0.1, 1:1 and 1:10 as outlined in FIG. 10. Day 21 single sera were analyzed for neutralization of LR-CHIKV virus replicon particles (VRPs) using a BHK-21 cell-based Luciferase assay. VRPs resemble a replication-deficient wild-type CHIKV displaying capsid and envelope proteins of the LR2006-OPY1 wild-type virus. Briefly, 2×10^4 BHK-21 cells were seeded in 96-well plates and incubated overnight at 35° C./5% CO₂. CHIKV-Δ5nsP3 VRPs were incubated with serial dilutions (starting at 1:20 to 1:312,500) of mouse sera for 1 hour at 35° C. CHIKV VRP/serum mixes were added to BHK-21 cells at an MOI of 5 and incubated for 1 hour at 35° C./5% CO₂. Cells were washed and fresh medium was added. Luciferase activity was measured in the supernatant 24 hours post-infection using the *Renilla* Luciferase Assay System (Promega). P-MVSB positive control (○), P5B-07 (E168K) negative control (□) and individual mouse sera immunized with respective P3:E168K ratio (filled circles). (FIG. 11A) Immunogenicity of CHIKV-Δ5nsP3 P3:E168K at a ratio of 1:0.1 (Group 1). (FIG. 11B) Immunogenicity of CHIKV-Δ5nsP3 P3:E168K at a ratio of 1:1 (Group 3). (FIG. 11C) Immunogenicity of CHIKV-Δ5nsP3 P3:E168K at a ratio of 1:10 (Group 5). (FIG. 11D) Summary of immunogenicity results shown in A, B, and C: strong immunogenic (++) positive), low immunogenic (+low positive) and negative (–) in comparison to P3.

FIG. 12 Vero cell culture train from thawing to infection with CHIKV-Δ5nsP3 in 850 cm² roller bottles. P: Passage, M: Million, T75: T-Flask 75 cm², T175: T-Flask 175 cm², RB850: Roller Bottle CellBIND 850 cm².

FIGS. 13A-13C Effect on virus titer of cell growth temperature, multiplicity of infection and post-seeding infection time during production of CHIKV-Δ5nsP3 in Vero cells. Vero cells were expanded at 35° C. in MEM, 2 mM glutamine and 10% FBS and seeded in 850 cm² roller bottles for cell infection. For virus production, different temperatures (37° C., 35° C., 28° C.), MOIs (0.1; 0.01; 0.001 TCID₅₀/cell) and times of cell infection post Vero cell seeding in days (D2, D4, D5) were tested in culture medium deprived of FBS. Shown are viral productivities measured on Vero cells according to the Reed & Muench method and expressed in TCID₅₀/mL. (FIG. 13A) 37° C.; (FIG. 13B) 35° C.; (FIG. 13C) 28° C.

FIG. 14 Total CHIKV-Δ5nsP3 virus productivity from each of the conditions described in FIG. 13.

FIGS. 15A-15B Maximum CHIKV-Δ5nsP3 virus titers: Response surface quadratic model. (FIG. 15A) ANOVA analysis of the model. Grey line: significant model term (Prob(F)<0.05; values greater than 0.1 indicate the model term is not significant). (FIG. 15B) Contour Plot of the model.

FIGS. 16A-16B CHIKV-Δ5nsP3 virus stability: Response surface quadratic model. (FIG. 16A) ANOVA analysis of the model. Grey line: significant model term (Prob(F)<0.05; values greater than 0.1 indicate the model term is not significant). (FIG. 16B) Contour Plot of the model.

FIGS. 17A-17B Heterogeneity of the CHIKV-Δ5nsP3 E2 viral protein at genomic nucleic acid positions 8882, 9112 and 9649, corresponding to E2 amino acids 168, 247 and 423, respectively: Model analysis from Day 2 CHIKV-Δ5nsP3 sample harvests. (FIG. 17A) ANOVA analysis of the model. Grey line: significant model term (Prob(F)<0.05). (FIG. 17B) Contour Plot of the model.

FIGS. 18A-18B Heterogeneity of the CHIKV-Δ5nsP3 E2 viral protein at genomic nucleic acid positions 8882, 9112 and 9649; corresponding to E2 amino acids 168, 247 and 423, respectively: Model analysis from Day 2 and Day 5 CHIKV-Δ5nsP3 sample harvests (28° C.). (FIG. 18A) ANOVA analysis of the model. Grey line: significant model term (Prob(F)<0.05). (FIG. 18B) Contour Plot of the model.

DETAILED DESCRIPTION OF THE INVENTION

During the course of industrialization of the CHIKV-Δ5nsP3 attenuated virus vaccine candidate, it was observed that passing of the virus on Vero cells resulted in higher virus titers with increasing passages; however, a concomitant increase in sequence heterogeneity of the CHIKV-Δ5nsP3 viral genome was also observed. Certain point mutations arising during passaging on Vero cells were found to be reproducible from batch to batch and appeared already in early passages on the new cell substrate. It was surprisingly observed that some of these mutations correlated with a significant loss of or decrease in neutralizing immunogenicity conferred by the CHIKV-Δ5nsP3 virus. Other reproducible mutations did not reduce immunogenicity and/or acted as “rescuing” mutations for the immunogenicity-reducing mutations. A correlation between a low multiplicity of infection (“MOI”) and generation of increased sequence heterogeneity in CHIKV-Δ5nsP3 was identified; however, because of the need to have a single source of virus over years of manufacturing, high MOIs are generally not feasible for industrial use. It was therefore not clear at the outset whether culturing conditions allowing the generation of immunogenic CHIKV-Δ5nsP3 particles with a production yield sufficient for reproducible and reliable manufacturing could be found (problem of the invention).

Provided herein are methods to control and minimize the herein observed immunogenicity-reducing mutations while still enabling high production yields. Also provided herein are pharmaceutical compositions comprising an effective amount of an immunogenic Chikungunya virus with a residual amount of a non-immunogenic variant of Chikungunya virus. In a preferred embodiment, the pharmaceutical composition is produced using a low MOI such as an MOI of less than 0.1, e.g. 0.01 or 0.001, but produced under such controlled conditions (e.g., reduced passage numbers following rescue, optimized temperature and host cell conflu-

ency of Chikungunya virus as described herein. In some embodiments, the virus particle is a live virus, a chimeric virus, an attenuated live virus, a modified live virus, or a recombinant live virus. In one embodiment, the virus particles of the invention may be optionally inactivated. In some embodiments, the virus particle is an attenuated form of the virus particle. For example, the virus may have reduced infectivity, virulence, and/or replication in a host, as compared to a wild-type virus. In some embodiments, the virus is a mutated or modified virus, for example the nucleic acid of the virus may contain at least one mutation relative to the wild-type virus, such as a substitution or deletion. In some embodiments, the virus is a recombinant live virus, meaning a virus that is generated recombinantly and may contain nucleic acid sequences from different sources. In some aspects, the wild-type Chikungunya virus is inactivated. In a preferred embodiment, the virus is inactivated with formaldehyde.

In one embodiment, the immunogenic Chikungunya virus is a live attenuated virus. In a preferred embodiment, the live attenuated Chikungunya virus is the protective CHIKV-Δ5nsP3 as described by Hallengård D, et al. (supra), referred to herein as CHIKV-Δ5nsP3 and defined by the nucleic acid sequence of SEQ ID NO: 1. Briefly, the wild-type CHIKV genome carries a positive-sense single-stranded RNA genome of 11 kb containing two open reading frames encoding nonstructural proteins (nsP1 to nsP4) and structural proteins (C, E3, E2, 6K, and E1), respectively. An attenuated CHIK virus, CHIKV-Δ5nsP3, based on the La Reunion CHIKV strain LR2006-OPY1, was constructed by substituting amino acid residues 1656 to 1717 of the P1234 polyprotein with a small linker (AA sequence AYRAAAG) in the hypervariable region of the nsP3 protein (see FIG. 2). CHIKV-Δ5nsP3 has been shown to be infectious, highly immunogenic and protective against challenge with wild-type CHIKV (Hallengård D, et al., supra and Hallengård D, et al., Prime-Boost Immunization Strategies against Chikungunya Virus (2014) J. Virology, 88(22):13333-13343, and Rogues P, et al. 2017, supra). In one embodiment, the live attenuated Chikungunya virus may be a variant of the CHIKV-Δ5nsP3 attenuated mutant virus. In a preferred embodiment, the live attenuated Chikungunya virus as provided herein comprises the CHIKV-Δ5nsP3 as encoded by the nucleic acid sequence defined by SEQ ID NO: 1. As used herein, the term “CHIKV-Δ5nsP3” is used interchangeably with “CHIKV-Δ5nsP3 virus”, “CHIKV-Δ5nsP3 particle”, “CHIKV-Δ5nsP3 virus particle” or plural versions thereof.

Provided herein is a pharmaceutical composition comprising an effective amount of a CHIKV-Δ5nsP3. In one aspect, an effective amount of an immunogenic CHIKV-Δ5nsP3 virus is defined as an amount sufficient to elicit neutralizing antibodies to Chikungunya virus. In a further aspect, an effective amount of an immunogenic CHIKV-Δ5nsP3 virus is defined as an amount to elicit an immune response in a vaccinated subject which confers protective immunity against Chikungunya virus infection. In a preferred aspect, an effective amount of CHIKV-Δ5nsP3 is defined as at least 10², at least 10³, at least 10⁴, at least 10⁵, at least 10⁶, preferably at least 10³ immunogenic CHIKV-Δ5nsP3 particles. In one aspect, immunogenic CHIKV-Δ5nsP3 particles are defined as CHIKV-Δ5nsP3 particles which express an E2 structural protein as defined by the polypeptide sequence of SEQ ID NO: 2. In one aspect, the E2 structural protein contains one or more point mutations that do not affect the immunogenicity of the virus, i.e., are not immunogenicity reducing. In one embodiment, the point mutations that do not affect the immunogenicity of the virus

may be at amino acids 232 and/or 247 of the E2 protein, such as H232Y and/or E247K. In one embodiment, the E2 structural protein of the CHIKV-Δ5nsP3 contains no more than about ten point mutations. In one embodiment, the E2 structural protein of the CHIKV-Δ5nsP3 contains no more than 9, 8, 7, 6, 5 or 4 point mutations. In a preferred embodiment, the E2 structural protein of the CHIKV-Δ5nsP3 contains three or less point mutations, most preferably only one or two point mutations.

As defined herein, an immunogenic CHIKV-Δ5nsP3 is a CHIKV-Δ5nsP3 which is capable of stimulating an effective immune response in vivo when delivered e.g. at a dose of about 3×10^4 TCID₅₀, i.e., an immune response in which neutralizing antibodies are produced which are sufficient for reducing or preventing signs or symptoms of Chikungunya virus disease. In a preferred embodiment, the immunogenic CHIKV-Δ5nsP3 as defined herein is a CHIKV-Δ5nsP3 which expresses an E2 structural protein according to the amino acid sequence provided by SEQ ID NO: 2. In a further preferred embodiment, the immunogenic CHIKV-Δ5nsP3 as defined herein is defined by the polynucleotide sequence according to SEQ ID NO: 1. As an alternative or additional definition, the immunogenic CHIKV-Δ5nsP3 of the current invention stimulates the production of antibodies with neutralizing capacity in an immunized subject, i.e., neutralization of Chikungunya virus in an in vitro assay of at least 50%, preferably at least 60%, preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, most preferably at least 95% at a 1:80 or higher serum dilution.

As defined herein, a non-immunogenic CHIKV-Δ5nsP3 is a CHIKV-Δ5nsP3 which elicits levels of neutralizing antibodies in a vaccinated subject inadequate to prevent signs or symptoms of Chikungunya virus disease. In a preferred embodiment, a non-immunogenic CHIKV-Δ5nsP3 is a CHIKV-Δ5nsP3 which expresses an E2 structural protein with at least one amino acid substitution, especially amino acid substitutions in the E2 structural protein, especially E168K and/or G55R substitutions, particularly an E168K substitution. A non-immunogenic CHIKV-Δ5nsP3 is further defined as eliciting antibodies in an immunized subject which show poor capacity to neutralize infection of cells with Chikungunya virus (wild-type or attenuated) in an in vitro assay. In particular, a non-immunogenic CHIKV-Δ5nsP3 is defined as eliciting levels of neutralizing antibodies in an immunized subject which provide less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, especially less than 10%, neutralization of Chikungunya virus in an in vitro neutralization assay at a 1:80 or higher serum dilution.

In a further aspect, the effective amount of CHIKV-Δ5nsP3 is defined as an amount sufficient to elicit neutralizing antibodies against wild-type Chikungunya virus. In one aspect, the pharmaceutical composition is a two-shot pharmaceutical composition. In a preferred aspect, the pharmaceutical composition is a one-shot pharmaceutical composition. In a preferred aspect, the pharmaceutical composition comprises at least 10^2 , at least 10^3 , at least 10^4 , at least 10^5 , at least 10^6 , preferably between about 10^3 to 10^5 total CHIKV-Δ5nsP3 viral particles, especially about 10^3 or 10^4 CHIKV-Δ5nsP3 comprised in a total pool of particles with and without point mutations, especially immunogenicity-reducing point mutations. In a preferred aspect, the pharmaceutical composition comprises a detectable amount of non-immunogenic CHIKV-Δ5nsP3 as defined herein; pref-

erably a non-immunogenic CHIKV-Δ5nsP3 with at least one point mutation compared with the wild-type E2 protein as defined by SEQ ID NO: 2.

In a preferred embodiment, the pharmaceutical composition comprises CHIKV-Δ5nsP3 and comprises an increased amount of a non-immunogenic variant(s) of CHIKV-Δ5nsP3, e.g. compared to a vaccine composition comprising CHIKV-Δ5nsP3 produced in BHK-21 cells as used in mouse studies described in Hallengård D, et al. 2014, supra, but still comprises sufficient immunogenic particles of CHIKV-Δ5nsP3 to produce protective immunity in a vaccinated subject. For instance, the pharmaceutical composition may comprise (i) CHIKV-Δ5nsP3 which expresses an E2 structural protein as defined by the polypeptide sequence of SEQ ID NO: 2 in an amount sufficient to produce protective immunity in a vaccinated subject; (ii) an increased amount of CHIKV-Δ5nsP3 having at least one mutation in said E2 structural protein, e.g. compared to a vaccine composition comprising CHIKV-Δ5nsP3 produced in BHK-21 as used in mouse studies described in Hallengård D, et al. 2014, supra; and (iii) optionally a pharmaceutically acceptable excipient.

It is demonstrated herein that production of CHIKV-Δ5nsP3 by serial passaging five or more times in Vero cells results in high levels of sequence heterogeneity, particularly in the E2 structural protein (see e.g. Example 2 below). For instance, E168K and/or G55R mutations of the E2 protein often appeared by passage 5 (see e.g. Table 3 below), and both correlated with a drop in immunogenicity. Accordingly, production of CHIKV-Δ5nsP3 using five or more passages in Vero cells as described in Hallengård D, et al. 2014 supra can unfavorably result in high levels of non-immunogenic mutants of CHIKV-Δ5nsP3 (such as E168K[E2]) in the vaccine composition. In contrast, it is demonstrated below that sequence heterogeneity in the E2 structural protein after fewer than five passages was much lower (see e.g. Example 3—although the E168K mutation was present after three passages, its frequency was only 18%).

Accordingly, in one aspect the pharmaceutical composition comprises (i) CHIKV-Δ5nsP3; and (ii) optionally a pharmaceutically acceptable excipient; wherein at least 30% of the CHIKV-Δ5nsP3 particles present in the composition express an E2 structural protein as defined by the polypeptide sequence of SEQ ID NO: 2. In this embodiment, at least 30% of the CHIKV-Δ5nsP3 particles are non-mutants with respect to the E2 structural protein, i.e. express the E2 structural protein of SEQ ID NO: 2. In other words, the frequency of sequence heterogeneity (i.e. mutant CHIKV-Δ5nsP3 particles expressing at least one mutation in the E2 structural protein of SEQ ID NO: 2) is 70% or less. Unless specified otherwise, when referring to “CHIKV-Δ5nsP3” or “CHIKV-Δ5nsP3 particles” in general it is intended to encompass both non-mutant and mutant forms of CHIKV-Δ5nsP3, i.e. CHIKV-Δ5nsP3 which express an E2 structural protein of SEQ ID NO: 2 and CHIKV-Δ5nsP3 which express an E2 structural protein having one or more mutations in SEQ ID NO: 2. In one embodiment, the E2 structural protein of the CHIKV-Δ5nsP3 contains no more than about ten point mutations. In one embodiment, the E2 structural protein of the CHIKV-Δ5nsP3 contains no more than 9, 8, 7, 6, 5 or 4 point mutations. In a preferred embodiment, the E2 structural protein of the CHIKV-Δ5nsP3 contains three or less point mutations, most preferably only one or two point mutations.

In preferred embodiments, at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% of the CHIKV-Δ5nsP3 particles present

in the composition are non-mutants, i.e. express an E2 structural protein as defined by the polypeptide sequence of SEQ ID NO: 2.

In one aspect, the pharmaceutical composition comprises (i) CHIKV-Δ5nsP3; and (ii) optionally a pharmaceutically acceptable excipient; wherein less than 70% of the CHIKV-Δ5nsP3 particles present in the composition express an E2 structural protein having one or more mutations with respect to the polypeptide sequence of SEQ ID NO: 2.

In one aspect, the pharmaceutical composition comprises (i) CHIKV-Δ5nsP3; and (ii) optionally a pharmaceutically acceptable excipient; wherein less than 70% of the CHIKV-Δ5nsP3 particles present in the composition express an E2 structural protein having the mutation E168K in the polypeptide sequence of SEQ ID NO: 2.

In a preferred embodiment, the mutations (e.g. the mutation E168K) in the E2 structural protein are present at a frequency of 70% or less, e.g., less than 70% of the total CHIKV-Δ5nsP3 particles comprise one or more mutations (or the mutation E168K) and 30% or more of the total CHIKV-Δ5nsP3 particles express a non-mutated E2 structural protein or an E2 structural protein that does not comprise the mutation E168K.

In a preferred embodiment, less than 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2% or 1% of the CHIKV-Δ5nsP3 particles present in the composition express an E2 structural protein having one or more mutations (such as, e.g., E168K) with respect to the polypeptide sequence of SEQ ID NO: 2. For instance, the composition may comprise 1-70%, 1-50%, 1-30%, 1-20%, 5-70%, 5-50%, 5-30%, 5-20%, 10-70%, 10-50%, 10-30% or 10-20% of mutant particles (i.e. CHIKV-Δ5nsP3 particles expressing an E2 structural protein having one or mutations (such as, e.g., E168K) with respect to the polypeptide sequence of SEQ ID NO: 2), compared to the total number of CHIKV-Δ5nsP3 particles (mutant and non-mutant) present in the composition. In one embodiment, the CHIKV-Δ5nsP3 particles expressing an E2 structural protein having an E168K mutation further comprise a mutation which mitigates the loss of immunogenicity conferred by the E168K mutation. In one embodiment, the mutation is in the nsP1 protein, especially at residue A38. In a preferred embodiment, the CHIKV-Δ5nsP3 particles expressing an E2 structural protein with an E168K mutation also express an nsP1 with an A38S mutation.

Also provided herein is a process for producing a pharmaceutical composition of the invention, comprising the steps of 1) growing a CHIKV-Δ5nsP3 virus on a cell line, and 2) minimizing the presence of immunogenicity-reducing mutations of the CHIKV-Δ5nsP3 virus. In one embodiment, the immunogenic CHIKV-Δ5nsP3 virus is propagated in a cell line selected from the group consisting of an EB66 cell line, a Vero cell line, a Vero-αHis cell line, a HeLa cell line, a HeLa-S3 cell line, a 293 cell line, a PC12 cell line, a CHO cell line, a 3T3 cell line, a PerC6 cell line, a MDSK cell line, a chicken embryonic fibroblast cell line, a duck cell line and a diploid avian cell line. In some embodiments, said cell line is a duck cell line. In some embodiments, said cell line is a diploid avian cell line. In some embodiments, said cell line is EB66 cell line. In a preferred embodiment, said cell line is a Vero cell line.

In one embodiment, the presence of immunogenicity-reducing mutations is minimized by passaging CHIKV-Δ5nsP3 less than 5 times, preferably less than 4 times, preferably less than 3 times, preferably less than 2 times, more preferably only one time, most preferably at most 3 times. As used herein, the passage numbers refer to the

number of in vitro passages following virus rescue (P0). In a preferred embodiment, the virus is passaged on Vero cells. In one aspect, the virus is grown at an optimal temperature. In a preferred embodiment, said optimal temperature is between about 28° C. and 37° C., preferably about 35° C.

In one embodiment, the host cell culture is infected with CHIKV-Δ5nsP3 at an optimal MOI. In one aspect, an optimal MOI is defined as an MOI low enough as to not require excessive amounts of working virus seed bank culture, but high enough to minimize immunogenicity-reducing mutations as described herein. In a preferred aspect, the optimized MOI is an MOI of less than 0.1, preferably an MOI of between about 0.1 and 0.001, more preferably an MOI of between about 0.09 to 0.0011, even more preferably an MOI of about 0.05 to 0.005, most preferably an MOI of about 0.01. In one aspect, the host cell confluency is assessed before infection. In one aspect, the host cell confluency is between about 20 and 90%, preferably between about 30 and 75%, more preferably between about 40 and 60%, especially about 50 to 60%. In one aspect, the cell culture is infected at an optimal timepoint post-host cell seeding; i.e., at between day 2 and day 5 after host cell seeding, preferably at about 4 days after host cell seeding. In one aspect, the virus particles are harvested between day one and day 6 after host cell infection, preferably between day one and day 4, preferably on day one or day 2 after host cell infection, preferably on both day one and day 2 after host cell infection.

In one aspect, immunogenicity-reducing mutations are point mutations at any location in the genome of CHIKV-Δ5nsP3 as defined by the polynucleotide sequence of SEQ ID NO: 1. In one embodiment, the immunogenicity-reducing mutations are present in the genome at a location other than the E2 protein. In a preferred embodiment, the immunogenicity-reducing mutations are located in the E2 protein, preferably at amino acid residues 55 and/or 168, e.g., G55R and/or E168K mutations, especially E168K. In some embodiments, the immunogenicity-reducing mutations as described herein are mitigated or “rescued” by other mutations in the genome. In one embodiment, a mitigating mutation of E168K is an A38S mutation of nonstructural protein 1 (nsP1).

In one aspect, the frequency of the E168K mutation of the E2 protein of the CHIKV-Δ5nsP3 is less than 90%, less than 80%, less than 70%, less than 60%, less than 50%, less than 40%, less than 30%, less than 20%, less than 10%, preferably less than 50% in the total pool of harvested CHIKV-Δ5nsP3.

In one aspect, the current invention provides an immunogenic CHIKV-Δ5nsP3 obtainable by the process provided herein. In another aspect, the current invention provides a pharmaceutical composition comprising an immunogenic CHIKV-Δ5nsP3 obtainable by the process provided herein.

Aspects of the invention provide a use of the process described herein for manufacturing a composition for immunization against a Chikungunya virus infection. In a preferred embodiment, the composition is a vaccine. In one embodiment, the vaccine is administered to the subject once, twice or three or more times. In one aspect, CHIKV-Δ5nsP3 viral particles isolated from immunized subjects have a similar point mutation profile to the vaccine composition administered, particularly with regard to point mutations in the E2 structural protein. In one embodiment, the vaccine is administered once or twice. In a preferred embodiment, the vaccine is administered only once; e.g., a one-shot vaccine. In one aspect, a booster vaccination is optionally adminis-

tered. In certain preferred aspects, the pharmaceutical composition is provided in lyophilized form.

Other aspects provide compositions comprising the virus particles obtainable by the process described herein for treating and/or preventing a Chikungunya virus infection. In one aspect, the compositions are for use in a method of stimulating an immune response in a subject and/or in a method of treating or preventing a Chikungunya virus infection. As used herein, the term “preventing” also means “protecting from”. The Chikungunya virus infection in one aspect may be caused by West African, East/Central/South African (ECSA) and/or Asian genotypes of Chikungunya virus.

Virus preparations produced using any of the processes described herein may be further subjected to additional processing steps, including additional filtration steps and/or lyophilization. The virus preparation may be subjected to analysis for purity of the preparation. For example, the virus preparations may be assessed for the presence of impurities and contaminants, such as, e.g., host cell genomic DNA, and/or host cell proteins. The purity of a virus preparation may be assessed using any method known in the art, such as size exclusion chromatography (SEC), optical density at different wavelengths, protein gel electrophoresis (e.g., SDS-PAGE), Western Blotting, ELISA, PCR, and/or qPCR.

In some embodiments, the virus preparation is assessed for residual impurities or contaminants. In some embodiments, the amount of residual impurities or contaminants is compared to the amount of impurities or contaminants at an earlier stage in the purification process, such as, e.g., directly after viral harvest. In some embodiments, the relative reduction of impurities in the final virus preparation is between 60-95% relative to the presence of impurities at an earlier stage in the purification process. In some embodiments, the relative reduction of impurities in the final virus preparation is approximately 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, or 95%. In some embodiments, the final virus preparation contains less than 5% impurities or contaminants. In some embodiments, the final virus preparation contains less than 5, 4, 3, 2, 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, or less than 0.1% impurities. In a preferred embodiment, the final virus preparation contains less than 1% impurities.

Any of the processes described herein may be used in the manufacture of a composition comprising purified virus for administration to a subject. In some embodiments, the subject is a mammalian subject, such as a human or a non-human animal, including livestock, pets or companion animals. In some embodiments, the composition is administered to a subject in need of immunization against the virus or similar virus as that of the virus preparation. In some embodiments, the virus preparations or compositions comprising viruses purified using the processes described herein are for treating or preventing infection with the virus or a similar virus as that of the virus preparation. In a preferred embodiment, the virus preparations or compositions comprising viruses purified using the processes described herein are for treating or preventing a Chikungunya virus infection, particularly a Chikungunya virus infection caused by West African, East/Central/South African (ECSA) and/or Asian genotypes of Chikungunya virus.

The CHIKV-Δ5nsP3 pharmaceutical compositions or CHIKV-Δ5nsP3 viruses purified using the processes described herein may be administered to a subject by any route known in the art. In some embodiments, the preparations or compositions may be administered via conventional

routes, such as parenterally or orally. As used herein, “parenteral” administration includes, without limitation, subcutaneous, intracutaneous, intradermal, intravenous, intramuscular, intraarticular, intraperitoneal, intrathecal or by infusion.

Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. The methods and techniques of the present disclosure are generally performed according to conventional methods well known in the art. Generally, nomenclatures used in connection with, and techniques of biochemistry, enzymology, molecular and cellular biology, microbiology, virology, cell or tissue culture, genetics and protein and nucleic chemistry described herein are those well-known and commonly used in the art. The methods and techniques of the present disclosure are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated.

TABLE A-1

Abbreviations	
Abbreviation	Definition
CHIKV	Chikungunya virus
CHIKV-Δ5nsP3	CHIKV with a defined deletion mutation in nsP3 (SEQ ID NO: 1)
ECSA	East-central south African
E (1, 2, 3) proteins	Envelope proteins
C protein	Capsid protein
6K protein	6 kilodalton protein
nsP	Non-structural protein
MEM	Minimum essential medium
EMEM	Eagle's MEM
FBS	Fetal bovine serum
PBS	Phosphate buffered saline
Pfu	Plaque forming unit
PRNT	Plaque reduction neutralization test
NGS	Next generation sequencing
h	hour
d	Day
AA	Amino acid
PCR	Polymerase chain reaction
qPCR	Quantitative PCR
s.c.	Subcutaneous
TSE	Transmissible Spongiform Encephalopathy
p.i.	Post-infection
PP	Polypropylene
μF	Micro-Farad
μm	micrometer
kDa	kilodalton
mg	milligram
RNA	Ribonucleic acid
Anti-anti	Antibiotic-antimycotic
SINV	Sindbis virus
SFV	Semliki Forest virus
WHO	World Health Organization
MVSB or P1-MVSB	Master virus seed bank (Passage 1 after rescue)
WVSB or P2-WVSB	Working virus seed bank (Passage 2 after rescue)
VRP	Virus replicon particle
TCID ₅₀	Tissue culture infective dose
LR-VRP	VRP based on the La Reunion CHIKV isolate
BHK(-21)	Baby hamster kidney cells
MOI	Multiplicity of infection
M	Million

TABLE A-1-continued

Abbreviations	
Abbreviation	Definition
T75, T150, T175	T flask 75, 150, 175 cm ²
RB850	Roller bottle 850 cm ²
Prob.F	Probability as determined by the F-test in one-way analysis of variance
ANOVA	Analysis of variance
DS	Drug substance
GLuc	Gaussia luciferase
GMP	Good manufacturing practice
R&D	Research and development
SEC	Size exclusion chromatography
SDS-PAGE	Sodium dodecyl sulfate-polyacrylamide gel electrophoresis
ELISA	Enzyme-linked Immunosorbant Assay
w/w	Weight/weight
mL	Milliliter
CPE	Cytopathic effect
V	Volts
mM	millimolar
TBS	Tris-buffered saline
NaCl	Sodium chloride
LR2006-OPY1	La Reunion CHIKV isolate
D-PBS	Dulbecco's phosphate buffered saline
RPM	Rotations per minute
RRV	Ross River virus
MCB	Master cell bank
ATCC	American Type Culture Collection

EXAMPLES

Example 1. Initial Trials for CHIKV-Δ5nsP3 Drug Substance (DS) Production

Assembly of Synthesized CHIKV-Δ5nsP3 Genome

The CHIKV-Δ5nsP3 virus genome was synthesized in five fragments at MWG Eurofins (Germany) and was fully assembled in the pMA plasmid (pMX vector with ampicillin resistance), a standard cloning vector. The pMX vector backbone is shown in FIG. 1. All cloning and plasmid preparation procedures were carried out under TSE-free conditions using electro-competent NEB10β *E. coli* cells. The cloning strategy used for the full assembly of the CHIKV-Δ5nsP3 genome in pMA is outlined in FIG. 3B. Briefly, the pMA plasmid containing fragment 1, which covers nsP1 and part of nsP2 (as shown in FIG. 3A), was

linearized via EcoRI/PacI restriction digestion and fragment 2 covering parts of nsP2 and nsP3 was fused to fragment 1. In parallel, fragment 3 (covering nsP4 and C) was fused to fragment 4 (covering C and E2) via ClaI/PacI cloning. In a third cloning step, fragments 3 and 4 were cloned via AgeI/XhoI and fragment 5 via XhoI/PacI into the AgeI/PacI-linearized pMA already containing fragments 1 and 2. The cloning resulted in the pMA_CHIKV-Δ5nsP3 vector encoding CHIKV-Δ5nsP3, as verified by sequencing.

CHIKV-Δ5nsP3 Rescue from Vero Cells ("Virus Rescue")

For the production of CHIKV-Δ5nsP3 virus particles from the engineered pMA_CHIKV-Δ5nsP3 vector on Vero cells (virus rescue), the pMA_CHIKV-Δ5nsP3 plasmid was linearized by NotI restriction digestion and subjected to in vitro transcription using Ambion's mMessage mMachine SP6 Kit (AM130). RNA integrity was confirmed via gel electrophoresis (not shown). In parallel, Vero cells were prepared for electroporation with viral RNA. Briefly, Vero cells were detached from cell culture flasks using TrypLESelect (Gibco) and washed twice with PBS. All centrifugation steps were performed at 300 g at room temperature. Viral RNA was mixed with 8×10⁶ Vero cells in 800 μl PBS and the Vero cell/RNA mix was transferred into 0.4 cm electroporation cuvettes. Two pulses were performed at 850 V, 25 μF, 200 Ohm. After electroporation, Vero cells were kept at room temperature for 10 minutes and finally resuspended in MEM/5% FCS/1% Antibiotic-Antimycotic (Anti-Anti)/2 mM L-Glutamine and incubated in T75 flasks for 48 hours at 35° C./5% CO₂. Cell culture supernatant containing rescued CHIKV-Δ5nsP3 (passage 0; P0) was harvested and centrifuged at 3,000 g for 10 minutes at 4° C. The virus titer was determined by plaque and TCID₅₀ assay on Vero cells. The rescued CHIKV-Δ5nsP3 (P0) was stored at -80° C. The genomic structure of the thus obtained CHIKV-Δ5nsP3 virus vaccine candidate, also referred to as VLA1553, is shown in FIG. 2.

Verification of CHIKV-Δ5nsP3 Sequence

In order to verify the viral genome sequence, viral nucleic acids were extracted from harvested cell culture supernatant using the QIAamp MinElute Virus Spin Kit (QIAGEN #57704) and cDNA-synthesis was done using the SuperScript III First-Strand Synthesis System (Life Technologies, Catalog #18080-051) using random hexamers. PCR with Phusion High Fidelity Polymerase was performed with primers amplifying overlapping regions of the CHIKV-Δ5nsP3 genome, and the PCR products were subjected to Sanger sequencing at MWG Eurofins, Germany. The sequences of primer pairs used for PCR and sequencing are shown in Table 1.

TABLE 1

Primer pairs used for CHIKV-Δ5nsP3 genome sequencing.			
Primer Pair	Forward primer sequence (5'-3') restriction sites (lower case)	SEQ ID NO: Reverse primer sequence (5'-3') restriction sites (lower case)	SEQ ID NO:
1	ttaggatccGATGGCTGCGTGAGACAC	8 taactcgagCCGTCAGGTCTGTTGAACAT	9
2	ttaggatccTACCACCAGGCGATTAAAG	10 taactcgagCTTTGCCCACTTACTGAAGG	11
3	ttaggatccTGCTACAGAAGTCACGCC	12 taactcgagGCCAAGCGTGAATCAG	13
4	ttaggatccAACAGCTTGAGGACAGAGCG	14 taactcgagCTCTGTCTCATCAGCTCGG	15
5	ttaggatccAAATTGCAGTCATAGGAGTCTTC	16 taactcgagAGTACGTTGACGTGCTCTGA	17
6	ttaggatccGTGGGTTAAACAACTGCAAA	18 taactcgagGGTTAAAGTCTTCTATCCTCTGG	19

TABLE 1-continued

Primer pairs used for CHIKV-Δ5nsP3 genome sequencing.					
Primer Pair	Forward primer sequence (5'-3') restriction sites (lower case)	SEQ ID NO:	Reverse primer sequence (5'-3') restriction sites (lower case)	SEQ ID NO:	
7	ttaggatccGGATAACCACTGGGATAATAGG	20	taactcgagAGTTGTGAAATTCCTTCTGCC	21	
8	ttaggatccCCAGATAGAACCAGTGAAC	22	taactcgagCAGCAGCTCTACTTGGGTC	23	
9	ttaggatccAGGAGGGAAAGACAGGCT	24	taactcgagCCCTCGCCTTCTTCTG	25	
10	ttaggatccCAAAATAGAAGGAGTGCAAAAAG	26	taactcgagCCTGGAGTTTCTTAAGTAATAGTTGC	27	
11	ttaggatccACCGGTCCAGGTCATTTA	28	taactcgagGCAGCAAATTCCTCCAG	29	
12	ttaggatccCCATTCCAGAACACTACAG	30	taactcgagATACCTGATTTTCATCATGGC	31	
13	ttaggatccCCTTTGATAAGAGCCAAAGATG	32	taactcgagTACAAAGTTATGACGGGTCTT	33	
14	ttaggatccCAACGAACAGGGCTAATTG	34	taactcgagGACCGCTTAAAGGCCAG	35	
15	ttaggatccGTGCATGAAATCGAAAATG	36	taactcgagTGGTCTTGTGGCTTTATAGACA	37	
16	ttaggatccAACCGGAGGAAACCTAC	38	taactcgagGTACCGCACCGTCTGG	39	
17	ttaggatccAGTACCTTGCAGCACGT	40	taactcgagCCCACATCGACAGG	41	
18	ttaggatccCGAGCCGTATAAGTATTGGC	42	taactcgagCGCCGTGAAGACCTTAC	43	
19	ttaggatccACTACTGTCACTACTTTGGAGC	44	taactcgagTACCGGTTTGTGTATTT	45	
20	ttaggatccCACAACTGGTACTGCAGAGAC	46	taactcgagGCGTAGCCCTTTGATCTATAG	47	
21	ttaggatccGGTGCTATGCGTGTCTGT	48	taactcgagATCTCTACGTCCCTGTGGG	49	

Passaging of CHIKV-Δ5nsP3 on Vero Cells

Following rescue of CHIKV-Δ5nsP3, Vero cells were infected and the virus was serially passaged in three replicates (see Table 2). For passaging, Vero cells seeded in T150 flasks and grown to confluency (1-3 days) were washed twice with 1×DPBS before the addition of 20 mL infection medium (EMEM w/o serum). Inoculum was added directly to the flask at the indicated volume and the cells were incubated for 24 h at 35° C., 5% CO₂. Passaging was done in three replicates (A, B and C) with infection of Vero cells at an MOI of 0.01 for the first passage from virus rescue (P0). The 20 mL harvests were transferred to a 50 mL PP tube 24 h p.i., cell debris was removed by centrifugation (3000 g, 10 min) and the supernatant transferred to a fresh 50 mL PP tube. A 49% (w/w) sucrose solution was added to a final concentration of 10% (w/w) and 1 mL aliquots of stabilized harvest were stored at ≤-70° C.

Subsequent infections were carried out with harvest without sucrose. The infections were carried out using different volumes of harvest which were roughly calculated based on the observed cytopathic effect in the previous passage and in parallel replicates. Volumes of harvest used for infection varied between 5 μL and 1 mL. Infections were followed by a single harvest after 24 h. Note that the MOI used for production of passages 2-16 was determined retrospectively after TCID₅₀ results were available, resulting in a wide (“uncontrolled”) range of MOIs throughout the experiment (See Table 2). This procedure was performed up to 16 passages in the three parallel replicates (replicates A, B and C) allowing systematic observation of various parameters during adaptation of CHIKV-Δ5nsP3 to Vero cell passaging. During this experiment, yield (TCID₅₀/mL), volume of infection, number of Vero cells per flask, Vero cell passage number and cytopathic effect (CPE) were recorded. The multiplicity of infection (MOI) was determined retrospec-

tively based on the measured TCID₅₀ and also recorded. The CHIKV-Δ5nsP3 passages were simultaneously assessed for plaque size in all three replicates.

TABLE 2

Serial passaging of CHIKV-Δ5nsP3 on Vero cells following virus rescue, performed in triplicate. Data shown include harvest yield (24 h; TCID ₅₀), infection volume (mL), number of cells/flask, Vero cell passage number, multiplicity of infection (MOI) and observed cytopathic effect (CPE) at 24 h post-infection.						
Replicate A						
Passage	TCID ₅₀ /mL	Infection (mL)	Cells @ T150	Vero passage	MOI (TCID ₅₀)	CPE @ 24 h
P0	9.77E+06					n.a.
P1A	2.59E+07	0.02	2.00E+07	149	0.010	50
P2A	1.90E+07	1	2.90E+07	149	0.893	50
P3A	2.92E+07	0.1	2.80E+07	149	0.068	20
P4A	2.14E+08	0.37	2.00E+07	150	0.540	90
P5A	1.01E+08	0.1	2.70E+07	150	0.792	60
P6A	3.50E+08	0.1	2.60E+07	150	0.390	60
P7A	7.43E+08	0.037	1.30E+07	151	0.997	50
P8A	8.02E+08	0.05	1.60E+07	151	2.323	90
P9A	1.51E+09	0.01	2.40E+07	151	0.334	60
P10A	3.60E+09	0.005	1.98E+07	152	0.381	80
P11A	1.88E+09	0.005	1.84E+07	152	0.982	90
P12A	6.94E+08	0.005	1.30E+07	152	0.724	90
P13A	4.22E+09	0.01	1.44E+07	153	0.482	60
P14A	3.05E+09	0.01	1.90E+07	153	2.219	90
P15A	3.03E+09	0.01	2.20E+07	153	1.384	90
P16A	2.91E+09	0.01	1.70E+07	154	1.781	90
Replicate B						
Passage	TCID ₅₀ /mL	Infection (mL)	cells @ T150	Vero passage	MOI (TCID ₅₀)	CPE @ 24 h
P0	9.77E+06					n.a.
P1b	6.47E+07	0.02	2.00E+07	149	0.010	50

TABLE 2-continued

Serial passaging of CHIKV-Δ5nsP3 on Vero cells following virus rescue, performed in triplicate. Data shown include harvest yield (24 h; TCID ₅₀), infection volume (mL), number of cells/flask, Vero cell passage number, multiplicity of infection (MOI) and observed cytopathic effect (CPE) at 24 h post-infection.						
P2B	5.32E+07	1	2.90E+07	149	2.232	50
P3B	2.27E+07	0.1	2.80E+07	149	0.190	20
P4B	1.65E+08	0.08	2.00E+07	150	0.091	80
P5B	1.65E+08	0.1	2.70E+07	150	0.613	60
P6B	4.06E+08	0.1	2.60E+07	150	0.636	60
P7B	9.19E+08	0.092	1.30E+07	151	2.875	60
P8B	7.43E+08	0.1	1.60E+07	151	5.744	90
P9B	2.14E+09	0.025	2.40E+07	151	0.774	75
P10B	4.06E+09	0.01	1.98E+07	152	1.080	90
P11B	2.09E+09	0.01	1.84E+07	152	2.214	90
P12B	8.76E+08	0.01	1.30E+07	152	1.608	90
P13B	2.91E+09	0.01	1.44E+07	153	0.609	60
P14B	4.22E+09	0.01	1.90E+07	153	1.530	90
P15B	3.79E+09	0.01	2.20E+07	153	1.917	95
P16B	3.79E+09	0.01	1.70E+07	154	2.232	90

Replicate C						
Pas-sage	TCID ₅₀ /mL	Infection (mL)	cells @ T150	Vero passage	MOI (TCID ₅₀)	CPE @ 24 h
P0	9.77E+06					n.a.
P1C	3.24E+07	0.02	2.00E+07	149	0.010	50
P2C	3.29E+07	1	2.90E+07	149	1.117	50
P3C	2.54E+07	0.1	2.80E+07	149	0.118	20
P4C	1.29E+08	0.15	2.00E+07	150	0.191	90
P5C	4.13E+08	0.1	2.70E+07	150	0.477	60
P6C	5.91E+08	0.1	2.60E+07	150	1.587	60
P7C	1.85E+09	0.185	1.30E+07	151	8.415	70
P8C	4.83E+08	0.2	1.60E+07	151	23.073	90
P9C	1.03E+09	0.1	2.40E+07	151	2.013	90
P10C	5.14E+09	0.015	1.98E+07	152	0.783	90
P11C	1.51E+09	0.015	1.84E+07	152	4.205	90
P12C	1.15E+09	0.015	1.30E+07	152	1.738	90
P13C	4.22E+09	0.01	1.44E+07	153	0.802	65
P14C	3.81E+09	0.01	1.90E+07	153	2.219	90
P15C	1.07E+09	0.01	2.20E+07	153	1.730	90
P16C	2.48E+09	0.01	1.70E+07	154	0.631	90

Trends Observed During Serial Passaging Under “Uncontrolled” MOI Conditions

During CHIKV-Δ5nsP3 adaptation to Vero cell passaging, it was observed that total CHIKV-Δ5nsP3 virus yield increased substantially with increased passage number on Vero cells as shown in Table 2. As shown in FIG. 4A, passages 6 and above yielded an approximately 100-fold increase in titer compared with virus rescue (P0) and early passages. Also observed was a concomitant decrease in CHIKV-Δ5nsP3 plaque size (FIG. 4B). The effect of in vitro passaging of wild-type Chikungunya virus on plaque size in other cell lines has been previously described by Gardner C L, et al. (Gardner C L, et al., 2014, supra). A high virus yield would be highly desirable for industrial production of the inactivated virus; however, it was also herein observed that CHIKV-Δ5nsP3 became less immunogenic with increased passaging as shown in FIG. 4C. Briefly, to determine the immunogenicity of different passages of CHIKV-Δ5nsP3 as generated in Table 2, above, groups of five C57Bl/6 mice were immunized once subcutaneously with a dose of 10⁵ TCID₅₀ CHIKV-Δ5nsP3 passage 5 (P5B), passage 8 (P8B) or P15 (P15C). CHIKV-Δ5nsP3 at P0 (virus rescue), also at 10⁵ TCID₅₀, was used as a positive control. Day 21 serum pools were assessed for their capacity to neutralize CHIKV-Δ5nsP3 (P0) in a PRNT assay on Vero cells, by testing 4-fold serial serum dilutions ranging from 1:20 to 1:327,680. As

shown in FIG. 4C, the immunogenicity in mice of the P5B CHIKV-Δ5nsP3 showed slightly shifted immunogenicity compared with the unpassaged CHIKV-Δ5nsP3 (P0), whereas a P8B virus showed substantially reduced immunogenicity comparable to the P15C virus. (As the P15C virus was non-immunogenic, it served as a negative control in subsequent PRNT assays.)

Finally, selected passages of the CHIKV-Δ5nsP3 were tested for genetic stability by Sanger sequencing. Upon passaging of CHIKV-Δ5nsP3 on Vero cells up to 16 times, it was verified that the 60 amino acid deletion in the nsP3 gene responsible for the attenuation of the virus was genetically stable, indicating that the virus does not revert back to wild-type, an important safety consideration for live attenuated vaccines.

Trends Observed During Serial Passaging Under Controlled MOI Conditions

The use of high MOIs (e.g., higher than 0.1) is not conducive to an industrial scale process as too much starting material is needed. In this regard, the use of a lower MOI (0.01) over three passages was tested and the immunogenicity of the resulting passages was determined. As shown in FIG. 5A, three passages from Replicate B in Table 2, were passaged under uncontrolled conditions, with MOIs ranging from 0.01 to 2.23. From the thus-obtained passage 3 (P3B), further passages were done up to P6 using an MOI of 0.01 for each passage and shown as “controlled conditions at MOI 0.01” in FIG. 5A. Under conditions using an MOI of 0.01, the immunogenicity of the resulting virus was lost very quickly as shown in FIG. 5B. By passage 5, the CHIKV-Δ5nsP3 was rendered non-immunogenic. This result was in contrast to the results seen with P5B in FIG. 4C, which was produced with a much higher MOI (see Table 2, Replicate B) and was still immunogenic. This observation seemed to indicate that a lower MOI leads to faster selection during Vero cell passaging for mutations that affect the immunogenicity of CHIKV-Δ5nsP3.

Example 2. Defining Sequence Heterogeneities of CHIKV-Δ5nsP3 which Affect Immunogenicity

Due to the observed reduction/loss of immunogenicity (neutralizing antibody titer) and decreased plaque size at higher CHIKV-Δ5nsP3 passages, it was of interest to analyze possible sequence heterogeneities within the viral populations at different passage numbers. In addition, it was of interest to analyze the sequence of individual plaques of the viral population. Unpassaged CHIKV-Δ5nsP3 (P0) did not show sequence heterogeneities based on Sanger sequencing. In general, with increased passage numbers an increase in sequence heterogeneities for all 3 replicates was observed (Replicates A, B and C; Table 3). In the case of passaging replicate C at passage 8 (P8C), the virus population was still heterogeneous (sequence heterogeneities shown in Table 3), whereas the P15C passage showed a more homogenous virus population with defined point mutations (indicated by *). The immunogenicity data shown in FIG. 4C (Example 1, above) focused on P5B and P8B, which showed sequence heterogeneities in the CHIKV-Δ5nsP3 non-structural proteins (nsP) and envelope protein E2 as shown in Table 3 below.

TABLE 3

Sequence heterogeneities in CHIKV-Δ5nsP3 at passages P5, P8 and P15 (Replicates A, B and C as produced in Table 2). Sequence heterogeneities were determined by Sanger sequencing. For that purpose, viral nucleic acids were extracted from harvested cell culture supernatant using the QIAamp MinElute Virus Spin Kit (Qiagen). The cDNA was synthesized using the SuperScript III First-Strand Synthesis System (ThermoFischer) using random hexamers. PCR with Phusion High Fidelity Polymerase was performed with primers amplifying overlapping regions of the CHIKV-Δ5nsP3 genome (Table 1), which were sequenced by Sanger sequencing at MWG Eurofins, Germany. The readout shows results of automated base calling (>20%) as well as heterogeneities detected by visual analyses of sequencing chromatograms. *Full mutations (i.e., 100%) are indicated by an asterisk.

Gene	Replicate A			Replicate B			Replicate C		
	P5A	P8A	P15A	P5B	P8B	P15B	P5C	P8C	P15C
nsP2		G577W	G577W		G577W	L494I G577W G621R		G577W	G577W*
nsP3					R470S	R470S			
nsP4					V113I				
E2	E168K	G55R	G55R	E168K	G55R	G55R	G55R	G55R	G55R*
	E247K	T230I	M171V	A423A	H99Y	H99Y	E247K	T230I	T230I*
	A423A	H232Y	T230I		E168K	E168K		E247D	
		E247D	H232Y		T230I	T230I			
					E247D				

Expansion and Sequencing of Single CHIKV-Δ5nsP3 Plaques

To understand the effect of individual mutations on immunogenicity and consequently develop a controlled and reproducible production process for a highly immunogenic CHIKV-Δ5nsP3 vaccine, individual plaques from CHIKV-Δ5nsP3 isolates P5B and P8B were picked. Briefly, serial dilutions of P5B and P8B CHIKV-Δ5nsP3 were used for infection of Vero cells in a plaque assay (described under FIG. 4) and after an incubation of 72 hours, single plaques of different morphologies were picked (small plaques were preferentially picked, since they began to appear after several passages on Vero cells) and expanded via re-infection of Vero cells (5×10^5) in 6-well plates. The different CHIKV-Δ5nsP3 samples that derived from single plaques were selected based on mutations in the E2 gene sequence prior to one further expansion on Vero cells. Clones P5B-02, P5B-03, P5B-04, P5B-07, P5B-11, P8B-01 and P8B-05 were expanded and purified for in vivo immunogenicity experiments. Individual plaques were expanded on Vero cells grown in Roller Bottles using a single 850 cm² Roller Bottle with CellBIND surface for each isolate. Upon reaching confluency, cells were washed twice with 100 mL D-PBS+Ca+Mg before 100 mL infection medium (EMEM w/o serum) containing the inoculum at an MOI of 0.01 was added and the cells were incubated for 24 h at 35° C., 5% CO₂. The 100 mL harvests were transferred to 50 mL PP tubes 24 h p.i. and the cell debris was removed by centrifugation, followed by a 0.2 μm filtration using Steriflip® vacuum filtration devices (Merck). Individual clarified virus harvests were first concentrated using Amicon® Centrifugal Filter devices, diafiltrated to TBS buffer and purified by

protamine sulphate and Capto™ Core700 treatment. Purified CHIKV-Δ5nsP3 clones were then used for further studies.

The full genome sequences of the expanded CHIKV-Δ5nsP3 samples, P5B+1 and P8B+1; namely P5B-02, P5B-03, P5B-04, P5B-07, P5B-11, P8B-01 and P8B-05 as described above, derived from single plaques, P5B and P8B, respectively, were assessed by Sanger sequencing. The observed point mutations of the individual plaques are summarized in Table 4 and schematic genomic sequences are shown in FIGS. 6B and 6D.

In order to assess the effect of specific point mutations on the immunogenicity of CHIKV-Δ5nsP3, day 19 mouse sera from mice immunized with the individual plaque-derived viruses were generated and analyzed in PRNT. Briefly, a single dose of CHIKV-Δ5nsP3 at an intended TCID₅₀ dose of 10^5 was administered subcutaneously to C57Bl/6 mice (10 per treatment group) and pools of day 19 sera were analyzed in a PRNT assay at 4-fold serial dilutions ranging from 1:20 to 1:327,680 for their virus neutralization capacity. The virus that was neutralized in the PRNT corresponded to a passage 2 CHIKV-Δ5nsP3 (P2, 560 pfu/ml) which did not show sequence heterogeneities and therefore was identical in sequence to the unpassaged CHIKV-Δ5nsP3 (P0). The neutralization mix (560 pfu/ml CHIKV-Δ5nsP3 P2 and serial serum dilutions) was incubated for 1 hour at room temperature and added onto Vero cells, followed by incubation for 2 hours. Finally, a methylcellulose overlay (0.8%) was added followed by incubation for 72 hours. The plaque readout was done following crystal violet staining (0.5% crystal violet in 5% Formaldehyde).

TABLE 4

Mutations in single picked plaques of P5 and P8 CHIKV-Δ5nsP3 passages. Sequencing of the full CHIKV-Δ5nsP3 genome was performed at MWG Eurofins, Germany, using the primers in Table 1. All P5B and P8B clones corresponded to a P6 (P5B + 1) and P9 (P8 + 1) CHIKV-Δ5nsP3, respectively, expanded on Vero cells. The PRNT₅₀ titers were calculated in GraphPad Prism using non-linear – 3 parameter calculations. The viral protein in which the respective point mutations were identified are shown in brackets. PRNT₅₀ values for non-immunogenic isolates were not measurable as indicated.

Experiment #	Isolate	Passage	Point mutation(s)	Immunogenicity	PRNT ₅₀
4399gr1	P0 #1	0	N/A	Positive control	386
4415gr1	P0 #2	0	N/A	Positive control	4215
4415gr2	P5B-02	5 + 1	E168K[E2] + A38S [nsP1]	retained	373
4415gr3	P5B-03	5 + 1	H232Y[E2]	retained	112
4415gr4	P5B-04	5 + 1	E168K[E2]	lost	—
4415gr5	P5B-07	5 + 1	E168K[E2]	lost	—
4415gr6	P5B-11	5 + 1	E247K[E2]	retained	219
4415gr7	P8B-01	8 + 1	H99Y/E168K[E2] + R470S[nsP3]	lost	—
4415gr8	P8B-05	8 + 1	G55R/H232Y[E2] + G577W[nsP2]	lost	—
4399gr2	P15C-DS	15	G55R/T230I[E2] + G577W[nsP2]	Negative control	—

— = not measurable

As can be seen in FIG. 6A, the immunogenicity of CHIKV-Δ5nsP3 P5B-11 and P5B-03, both with single point mutations in the E2 protein, E247K and H232Y (FIG. 6B), respectively, is not affected when compared to the immunogenicity of P0 CHIKV-Δ5nsP3. The neutralization capacity of P0 serum is shown from two independent mouse experiments (4415; P0 #2 and 4399; P0 #1) delineating the acceptable range of immunogenicity. On the other hand, P8B-05, characterized by 3 point mutations, G577W, G55R and H232Y in nsP2 and E2, respectively, is non-immunogenic in mice. The G55R mutation was already described in literature by Gardner C L, et al., as being a result of passaging CHIKV in vitro. The viral particle is affected by an increased dependence on heparan sulfate binding in vitro which leads to an attenuation in vivo (Gardner C L, et al., 2014, supra).

As can be seen in FIG. 6C, only one of the four CHIKV-Δ5nsP3 samples derived from single plaques (P5B-02) was still immunogenic and was characterized by two point mutations, A38S and E168K in nsP1 and E2 (FIG. 6D), respectively. Also, P8B-01 with three point mutations, R470S in the nsP3 and H99Y and E168K in the E2 protein, was non-immunogenic in mice and was comparable to the negative control P15C. P5B-04 and P5B-07 each had single point mutations in E2, i.e., Glutamic acid 168 was mutated to a Lysine (E168K), which was also shown to have a direct effect on immunogenicity, since CHIKV-Δ5nsP3 neutralization capacity was lost.

In summary, it was observed that many of the mutations arising during passaging on Vero cells were located in the E2 protein. Some of the identified point mutations in the E2 protein and/or other parts of the genome did not substantially affect immunogenicity of the virus; particularly the H232Y[E2] and E247K[E2] mutations. However, some of the other identified mutations in the CHIKV-Δ5nsP3 resulted in loss of immunogenicity; particularly the frequently-occurring E168K[E2] mutation, whether alone or in combination with other mutations. An interesting exception was the mutant with both E168K[E2] and A38S[nsP1] mutations, which maintained immunogenicity. This observation suggests that the A38S[nsP1] mutation has a mitigating effect on the reduced immunogenicity conferred by the E168K[E2] mutation. Furthermore, an isolate with G55R/H232Y[E2] and G577W[nsP2] mutations also demonstrated

poor immunogenicity, perhaps mainly due to the G55R mutation in E2, as the H232Y mutation alone had little effect (see P5B-03).

The E168K and G55R mutations in Chikungunya virus E2 protein were previously described as conferring increased positive surface charge, leading to increased interaction with heparan sulfate and/or other Glycosaminoglycans (GAGs), ultimately resulting in increased specific infectivity. On the background of wild-type CHIKV, the mutations were shown to cause a smaller plaque size, due to lower spread on plates mediated by binding to heparan sulfate. Furthermore, the mutations resulted in attenuation of CHIKV in a mouse model of musculoskeletal disease (MSD), with decreased spread in mice to organs and thus lower levels of viremia (Gardner C L, et al., 2014, supra; Silva L A, et al., A single-amino-acid polymorphism in Chikungunya virus E2 glycoprotein influences glycosaminoglycan utilization (2014) J. Virol.; 88(5):2385-97). The fact that the presence of E168K and G55R mutations in an otherwise wild-type CHIKV resulted in intermediate attenuation is consistent with the present disclosure with regard to reduced plaque size or reduced immunogenicity in vivo. However, it was unexpected that the said two mutations on the background of the attenuated CHIKV-Δ5nsP3 would result in loss of immunogenicity in mice as reported herein.

It has also been reported as a common phenomenon for other cell culture passaged alphaviruses such as Sindbis virus (SINV; Klimstra W B, et al., Infection of neonatal mice with Sindbis virus results in a systemic inflammatory response syndrome (1999) J. Virol.; 73(12):10387-98; Klimstra W B, et al., The furin protease cleavage recognition sequence of Sindbis virus PE2 can mediate virion attachment to cell surface heparan sulfate (1999) J. Virol.; 73(8): 6299-306; Byrnes and Griffin, Binding of Sindbis virus to cell surface heparan sulfate (1998) J. Virol.; 72(9):7349-56), Ross River virus (RRV; Heil M L, et al., An amino acid substitution in the coding region of the E2 glycoprotein adapts Ross River virus to utilize heparan sulfate as an attachment moiety (2001) J. Virol.; 75(14):6303-9) and Semliki Forest virus (SFV; Smit J M, et al., Adaptation of alphaviruses to heparan sulfate: interaction of Sindbis and Semliki forest viruses with liposomes containing lipid-conjugated heparin (2002) J. Virol.; 76(20):10128-37) that substitutions for positively-charged residues in E2 confer enhanced heparan-sulfate dependent infectivity in vitro and

that these mutations can be selected within a few serial in vitro passages. Further, it was shown that such mutations led to attenuation of the viruses in vivo (Byrnes A P and D E Griffin, Large-plaque mutants of Sindbis virus show reduced binding to heparan sulfate, heightened viremia, and slower clearance from the circulation (2000) *J. Virol.*; 74(2):644-51; Klimstra W B, et al. 1999, supra).

Because sequence heterogeneities, with a concomitant drop in immunogenicity, were already apparent at passages P5 and P8 of CHIKV-Δ5nsP3, sequence heterogeneities at earlier passages, as well as their effects on immunogenicity, were examined more closely as outlined below.

Example 3. Defining Sequence Heterogeneities and Immunogenicity of CHIKV-Δ5nsP3 at Passage P3

The occurrence at later passages of sequence heterogeneities with adverse effects on the immunogenicity of CHIKV-Δ5nsP3 as measured by neutralizing antibody titers warranted finding the optimal passage which was characterized by both high immunogenicity as well as a viral titer sufficient for production of an effective vaccine.

To determine genetic stability of the CHIKV-Δ5nsP3 during MVSB (P1), WWSB (P2) and CHIKV-Δ5nsP3 drug substance ("VLA1553") (P3) production, independently generated passages 1, 2 and 3 were sequenced. As determined by Sanger sequencing, P0 (virus rescue), P1 (MVSB) and P2 (WWSB) did not show any obvious sequence heterogeneities. The next step was to demonstrate reproducibility of genetic stability of P3 derived purified drug substance (DS) using P2 (WWSB) for infection. In total, four independent P3 harvests, consisting of combined day 1 and day 2 harvests, were produced in two T150 T-flasks using P2 (WWSB) for infection (MOI 0.01). For each replicate, the individual harvests at day 1 and day 2 were pooled (total volume ~50 mL) and concentrated approximately 10-fold (Amicon 100 kDa ultrafiltration device). Diafiltration was done against 25 mM Tris/150 mM NaCl, pH 7.4, followed by protamine sulfate treatment (2 mg/mL final concentration) to precipitate host cell DNA. The clear supernatant was then further purified by batch adsorption chromatography using CaptoCore 700 resin (addition of ~1 mL of 50% slurry in Tris/NaCl buffer). The resin was removed by centrifugation and sucrose was added to a final concentration of 10% to allow freezing and thawing of CHIKV-Δ5nsP3. The final formulation was then 0.2 μm sterile filtered and stored frozen (<-65° C.) until further processing.

At passage 3 (P3), no heterogeneities by automatic base calling were detected (Eurofins –all <20%). However, by visual inspection, a small fraction of the viral population showed a consistent increase in the E168K and E247K sequence heterogeneities in the gene for the CHIKV glycoprotein E2, which was absent in the rescued CHIKV-Δ5nsP3 (P0) as well as the MVSB and WWSB samples. FIG. 7 shows sequencing chromatograms for the four independently generated P3 DS samples (Examples 1-4) compared with the virus at passage 0 (P0). As indicated by the box outlines, a G/A heterogeneity at the codon for amino acid 168 (genomic nucleic acid position 8882) was detectable in all four replicates and verified by the reverse sequencing reaction, which revealed the same heterogeneity. The same position in the P0 sample showed a sharp peak for G in the forward sequencing reaction (and C in the reverse sequencing reaction). The E247K heterogeneity (genomic nucleic acid position 9119), on the other hand, was present but barely detectable in the sequencing chromatograms.

Additionally, next generation sequencing of P3 was carried out and compared with sequencing of passage 1 (P1-MVSB) in order to quantify the amount of E168K and E247K within the viral population. As can be seen in FIG. 8, at the stage of P1-MVSB, only a very low level of sequence heterogeneities was detectable (all below background; 15% cut-off indicated by the dotted line). The two representative P3 sample, however, showed an overall increase in sequence heterogeneities, with two mutations in the E2 gene that reached or rose above background (E168K in 18% and 15% of the viral population—genomic nucleic acid position 8882 and, to a lower extent, E247K—genomic nucleic acid position 9119).

In summary, the presence of an E168K mutation in the E2 protein of CHIKV-Δ5nsP3 was identified by Sanger sequencing and NGS in eight independently-generated P3 samples, demonstrating the reproducibility of this result. Representative sequencing examples are shown in FIGS. 7 and 8. These data demonstrate that the emergence of an E168K mutation in the E2 protein of CHIKV-Δ5nsP3 virus was highly reproducible upon passaging in Vero cells and that it diminished the immunogenicity of the attenuated CHIKV-Δ5nsP3 virus as shown in Example 2 where P5B plaque-derived viruses (P5B-04 and P5B-07 in FIG. 6) were non-immunogenic. Two additional mutations, G55R and G82R, which were observed during passaging of CHIKV-Δ5nsP3 on Vero cells, like E168K have been reported to affect the virulence of CHIKV (Gardner C L, et al., 2014, supra, Silva L A, et al., 2014, supra and Gorchakov R, et al., 2012, supra). Attenuation of Chikungunya virus vaccine strain 181/clone 25 is determined by two amino acid substitutions in the E2 envelope glycoprotein (2012) *J. Virol.*; 86(11):6084-96; Epub 2012/03/30, leading to attenuation of the virus. Since these two mutations also negatively impacted the immunogenicity of CHIKV-Δ5nsP3, their emergence should be avoided for the production of a highly immunogenic CHIKV-Δ5nsP3 vaccine candidate.

The locations of amino acids prone to mutation within the E1/E2 dimer are shown in FIG. 9. The three E2 amino acids G55, G82 and E168 are all accessible on the surface and therefore their mutation may potentially affect interactions of CHIKV with its cellular receptors. In contrast, E247[E2] is more buried within the protein structure and thus may not be surface exposed to the same extent. The position of E247 may be one reason why the E247K mutation did not noticeably affect the immunogenicity of CHIKV-Δ5nsP3 as shown above.

Example 4. Determining the Threshold of the E168K Mutation for Loss of Immunogenicity of a Heterogeneous CHIKV-Δ5nsP3 Virus Population

The above observations indicated that the E168K[E2] mutation appears early and frequently during passaging of CHIKV-Δ5nsP3 on Vero cells and is associated with lost immunogenicity. In order to develop a process for the reliable manufacture of an effective, immunogenic live-attenuated Chikungunya virus vaccine, the tolerance for this mutation in a sample of the CHIKV-Δ5nsP3 vaccine was tested by preparing different ratios of the P3 drug substance and the virus P5B-07 (E168K single mutant; see Table 3).

Passage 3 (P3) drug substance, which displayed about 20% E168K heterogeneity (data not shown), was mixed with a preparation of CHIKV-Δ5nsP3 from the P5B-07 isolate (E168K mutant) at ratios of 1:0.1, 1:1 and 1:10. The mixtures were sequenced to verify the approximate frequency of the E168K mutation in each virus preparation. As

shown in FIG. 10 (see row labeled heterogeneity), the relative amount of nucleotide A, compared with the wild-type nucleotide G, was shown to increase with increasing levels of added P5B-07 isolate (up to 100% A in the E168K control group 7).

In order to determine the effect of the E168K mutation on immunogenicity, C57Bl/6 mice were immunized s.c. with the different CHIKV-Δ5nsP3 samples specified in FIG. 10 with an intended dose of 3×10^4 TCID₅₀ (actual dose shown in FIG. 11D). Day 21 mouse sera were collected and individual sera from the 1:0.1, 1:1 and 1:10 groups were tested and compared with pooled sera from the two control groups, P3 (Group 6) and P5B-07 (E168K; Group 7) using a virus replicon particle (VRP)—based neutralization assay (FIG. 11). VRPs resemble a replication deficient wild-type CHIKV displaying capsid and envelope proteins of the LR2006-OPY1 virus. The method was carried out essentially as described by Gläsker S, et al. (Virus replicon particle based Chikungunya virus neutralization assay using *Gaussia* luciferase as readout (2013) Virol. J.; 10:235). The VRPs were produced by co-transfecting BHK-21 cells with a CHIKV replicon expressing *Gaussia* luciferase (GLuc) and two helper RNAs expressing wild-type CHIKV capsid protein and the remaining structural proteins (E3, E2, 6K and E1), respectively. The resulting single round infectious particles were used in the CHIKV neutralization assay using secreted GLuc as a readout. Upon neutralization of VRPs in the Luciferase assay, a reduction of GLuc expression by BHK-21 cells can be measured. It was observed that the capacity for eliciting neutralizing immunity dropped with higher amounts of the E168K mutant (FIGS. 11A, B and C, respectively). The number of individual mice showing either positive, low positive or negative immune responses to Chikungunya virus is shown in FIG. 11D.

This finding confirms that, as the ratio of E168K mutant to wild-type viral particles in a virus population increases, the immunogenicity of CHIKV-Δ5nsP3 in mice is diminished. It is therefore crucial to closely monitor position E168 in E2 to ensure high immunogenicity of the CHIKV-Δ5nsP3 vaccine. Based on previous passaging processes and quantification of E168K within the viral population at passage 8, it was observed that at a rate of about 70% of the E168K mutation within the CHIKV-Δ5nsP3 population, the immunogenicity was lost when analyzing mouse serum pools in PRNT.

Example 5. Upstream Process for Reducing E168K Mutations in CHIKV-Δ5nsP3

The aim of this example was to characterize an optimized Vero cell culture based process for the production of CHIKV-Δ5nsP3 in roller bottles. The impact of several upstream process parameters (MOI, day of Vero cell infection following plating and incubation temperature) on viral productivity and sequence heterogeneity of the E2 protein were tested using the GMP Working Virus Seed Bank (GMP WWSB B3005044; passage 2, also referred to herein as “P2 CHIKV-Δ5nsP3”) and the R&D Vero working cell bank to produce drug substance (DS; passage 3, i.e. also referred to herein as “P3 CHIKV-Δ5nsP3”).

Preparation of GMP WWSB B3005044

A characterized Pre-Master Virus Seed Bank (PMVSB, Pre-Master Virus Seed Bank AFR886/197579 from virus rescue from Vero cells) was established under R&D conditions and a Pre-Master Virus Seed Bank was used to generate the Master Seed Banks of the CHIKV-Δ5nsP3 under GMP conditions. The GMP Working Virus Seed Bank,

VLA78-1553-WWSB-2016, batch B3005044 was produced at Halix B.V. under the same production method and GMP conditions as described for the VLA78-1553-MVSB-2016, batch #B3005567. Briefly, the VERO Working Cell Bank (internal designation: ICB 2014/002) was expanded in four stages in a seed train using T75 cm² flasks (1×), then T175 cm² flasks (3×) and in the last stage 6×850 cm² roller bottles as shown in FIG. 12. Four of the six roller bottles were used for production of the CHIKV-Δ5nsP3 seed banks. The total amount of cells in the four roller bottles was determined. The Pre-Master Virus Seed Bank was used to generate the CHIKV-Δ5nsP3 Master Virus Seed Bank. Subsequently, the CHIKV-Δ5nsP3 Master Virus Seed Bank was used to generate the CHIKV-Δ5nsP3 Working Virus Seed Bank. Infection was done at all stages at an MOI of 0.01. After 24 hours of infection (35° C.; 5% CO₂; 0.5 RPM) the CHIKV-Δ5nsP3 was harvested. Tris and sucrose stock solution was added to a final concentration of 5% Sucrose and 25 mM Tris in the harvest pool. After formulation, the pool was filtered through a steam sterilized 0.22 μm filter into a sterile 500 mL bio process container by using a peristaltic pump. Filling of vials was done one by one with 0.7 mL formulated and filtered harvest using a peristaltic filling pump. A ThermoFisher capper/decapper device was used to open and close the 2 D Matrix Cryo vials. Filled vials of the GMP seed banks were stored at <−65° C.

Culture of Vero Cells

Culturing of Vero cells was performed at 35° C. and 5% CO₂ in T75 cm² (T75), T175 cm² (T175) T-Flasks and 850 cm² roller bottles (850RB). Vero cells used in the different experiments were derived from the GMP master cell bank MCB ICB/2014/001. The internal designation of this research working cell bank was Bk5685. The GMP master cell bank was derived from the WHO Vero cell bank 10-87 P134 which originated from the Institut Merieux (Aventis Pasteur) P129 bank and ultimately from the original ATCC CCL 81 P113 bank. More detail regarding the cell culture train is shown in FIG. 12. Cells were maintained in MEM medium supplemented with 10% FBS and 2 mM L-glutamine.

Virus Production in 850 cm² Roller Bottles

Following two, four or five days of cell expansion at 35° C. in 850RB, cells were washed with PBS and infected with the CHIKV-Δ5nsP3 (WWSB B3005044) at MOIs of 0.1, 0.01 or 0.001 TCID₅₀/cell. For virus production, infected cells were incubated at 37° C., 35° C. or 28° C. in 100 mL of MEM medium supplemented with 2 mM glutamine.

Virus Titration

Virus titers were determined on Vero cells using the TCID₅₀ assay. Cells were seeded in microplates and infected with 10-fold serially diluted virus samples in EMEM supplemented with 0.5% FBS and 2 mM glutamine. After a one week incubation at 35° C./5% CO₂, virus-induced cytopathic effects were monitored and viral titers were calculated according to the Reed and Muench method (Reed, L. J.; Muench, H. A simple method of estimating fifty percent endpoints (1938) The American Journal of Hygiene 27:493-497).

Virus Genome Extraction and Sequencing

Viral nucleic acid was extracted and purified from Vero cell culture supernatant at the indicated timepoints using QIAamp MinElute Virus Spin Kit (Qiagen) and cDNA synthesis was performed using SuperScript III First-Strand Synthesis System (ThermoFischer) using random hexamers. For sequencing of the E2 gene region, first, PCRs with Phusion High Fidelity Polymerase (ThermoFischer) were done using primers 16F, 16R, 17F, 17R, 18F and 18R (for

primer sequences see Table 1) to amplify overlapping regions of the CHIKV E2 gene. After purification of PCR amplicons, Sanger sequencing was performed at MWG Eurofins, Germany. In addition to analyses of sequence heterogeneities that were detected by automatic base calling (>20%), all sequencing chromatograms were manually read to detect also heterogeneities below the detection limit (<20%).

Optimization of a Process for Producing an Immunogenic P3 CHIKV-Δ5nsP3 Drug Substance

To optimize the process for producing passage 3 CHIKV-Δ5nsP3 on Vero cells, different MOIs, times of Vero cell infection post-seeding and temperatures of incubation were tested in all combinations as shown in Table 5. Additionally, yields were analyzed at different days following infection. Three aspects of the harvested virus were monitored: viral productivity, stability of the titer as well as the level of sequence heterogeneity of the E2 structural protein.

TABLE 5

Parameters tested for production of CHIKV-Δ5nsP3 on Vero cells in 850 cm ² roller bottles and the parameters of the identified optimized process.			
	Tested*	Optimized Process	
MOI (TCID ₅₀ /cell)	0,1, 0,01, 0,001	0,01	
Time of cell infection (post cell seeding)	D2, D4, D5	D4	
Temperature (° C.)	37, 35, 28	35	

*all combinations were performed.

Virus Production

CHIKV-Δ5nsP3 production kinetics achieved for all the conditions tested are shown in FIG. 13. The optimal process currently used to produce CHIKV-Δ5nsP3 (35° C., MOI 0.01, infection on day 4 after Vero cell seeding) was also included in the experiment, showing virus productivity at the expected level of 10^{8.0-8.5}/mL 24 h after infection (FIG. 13B).

Compared to MOI and time of infection, temperature had the most impact on viral production kinetics. At 37° C. and 35° C. (FIGS. 13A and 13B, respectively), maximum productivity was achieved at day one post-infection. The CHIKV-Δ5nsP3 infectivity dropped substantially during the subsequent 24 hours, with a slightly less pronounced viral titer loss at 35° C. Temperature reduction to 28° C. resulted in a striking change of the viral kinetics. Maximal viral productivity was delayed one to four days depending on the

MOI used; and significantly improved virus titer stability was observed (FIG. 13C). The total virus yield produced from all of the harvests is shown in FIG. 14. For the 28° C. condition as shown in FIG. 13A, the yield of two harvests were combined (day 1 and day 2 post-infection); for the 35° C. condition (FIG. 13B), the yield of all three harvests were combined (days 1-3) and for the 37° C. condition (FIG. 13C), the yield of five harvests (days 1-5) were combined.

To complete initial observations, viral productivity and titer stability data were analyzed using a response surface quadratic model (FIGS. 15 and 16). For viral productivity, the maximum virus titer values were used. For virus titer stability, the delta value resulting from the subtraction of the maximum titer from the titer measured one day later in the production process was calculated and used. For total virus productivity, the individual titers at each time point were added.

With ANOVA analysis of both models, it was possible to indicate the statistically significant influencing factors (FIGS. 15A and 16A). Confirming previous observations, temperature was the strongest factor affecting viral productivity and stability. In particular, low temperatures (28° C.) enabled higher viral titers while keeping the virus titer reasonably stable over time. However, there was not a significantly higher overall total virus productivity at this temperature compared with 35° C. (see esp. FIG. 14).

Time of infection after Vero cell seeding also influenced the response, but to a lower extent. MOI did not have a significant impact. For both models, infection at 72 h post cell seeding was an adequate time for cell infection. Conversely, a single temperature did not allow combining optimal virus production and titer stability since the highest viral yields were found at 35° C. and the most stabilized titers were observed at 28° C. (FIGS. 15B and 16B). Maximal total virus productivity within the shortest time post infection was achieved at 35° C. (see FIG. 14).

Analysis of E2 Protein Gene Sequence

Virus samples collected at either day 2 or day 5 after infection of Vero cells (infected at day 4 post-seeding) were selected to conduct an analysis of genomic RNA sequence of the viral E2 structural protein. These samples are most representative for Vero cell confluence on roller bottles. Tables 6 and 7 below summarize the percentage of heterogeneities estimated for four amino acid (AA) positions based on the nucleic acid sequence determined by Sanger sequencing. Table 6 shows data for CHIKV-Δ5nsP3 grown at three different temperatures and harvested two days post-infection and Table 7 shows data for CHIKV-Δ5nsP3 grown at 28° C. and harvested 5 days post-infection.

TABLE 6

Analysis of RNA genome sequence for E2 viral protein from D2 post-infection sample harvests. Shown are the estimated percentages of nucleic acid heterogeneities corresponding to four AA positions (indicated in parentheses), as determined by Sanger sequencing. The heterogeneity at nucleic acid position 9649 is a silent mutation.									
Time post seeding in roller bottles	D4								
	28° C.			35° C.			37° C.		
MOI (TCID ₅₀ /cell)	0.1	0.01	0.001	0.1	0.01	0.001	0.1	0.01	0.001
Pos. 8543 (G55R)	0	0	0	0-5	0-5	0-5	0	0	0

TABLE 6-continued

Analysis of RNA genome sequence for E2 viral protein from D2 post-infection sample harvests. Shown are the estimated percentages of nucleic acid heterogeneities corresponding to four AA positions (indicated in parentheses), as determined by Sanger sequencing. The heterogeneity at nucleic acid position 9649 is a silent mutation.									
Time post seeding in roller bottles		D4							
Temperature	28° C.			35° C.			37° C.		
MOI (TCID ₅₀ /cell)	0.1	0.01	0.001	0.1	0.01	0.001	0.1	0.01	0.001
Pos. 8882 (E168K)	30	30	40-50	25	25	30-50	25	25	30-50
Pos. 9119 (E247K)	0-5	0-5	0-5	5-10	10	10-20	10-20	10-20	10-20
Pos. 9649 (A423, Silent)	0-5	5-10	10-20	5	5	5-10	0-5	0-5	5-10

TABLE 7

Analysis of RNA genome sequence for E2 viral protein from D5 post-infection sample harvests. Shown are the estimated percentages of nucleic acid heterogeneities corresponding to four AA positions (indicated in parentheses), as determined by Sanger sequencing. The heterogeneity at nucleic acid position 9649 is a silent mutation.

Time post seeding in roller bottles		D4		
Temperature		28° C.		
MOI (TCID ₅₀ /cell)		0.1	0.01	0.001
Pos. 8543 (G55R)		0	0	0
Pos. 8882 (E168K)		30-40	40-50	50
Pos. 9119 (E247K)		0-5	0-10	10
Pos. 9649 (A423, Silent)		0-5	0-5	10-20

MOI, temperature, day of infection post-Vero cell seeding and day of sample harvest all influenced the productivity and the quality of CHIKV-Δ5nsP3 when produced in Vero cells. The strength of each parameter, however, was of different importance. For example, the results suggested a correlation between MOI and heterogeneity levels; i.e., the lower the viral input at infection, the higher the observed level of heterogeneity at harvest. The incubation temperature did not appear to impact the stability of the nucleotide sequence, with the exception of Pos. 9119 (E247K) where a higher level of heterogeneity was observed at 37° C. (Table 6).

Also, the sample harvest collected later in the viral kinetic triggered a slightly higher level of heterogeneity for the same AA position.

To complete this first analysis, mathematical modelling of the raw data was also performed (FIGS. 17 and 18). Only three nucleic acid positions (Pos. 8882, 9119, 9649) could be analyzed as no significant model was calculated for the position 8543. As indicated in FIG. 17, temperature and MOI had different impacts depending on the nucleic acid position considered. The MOI significantly influenced the heterogeneities observed for nucleic acid positions 8882 and 9649, but not for 9112. The temperature affected positions 9112 and 9649, but not 8882 (FIG. 17). MOI was shown to consistently influence the level of heterogeneity when D2 and D5 post-infection harvest samples from 28° C.-produced virus were compared—minimized levels of AA heterogeneity were observed when cells were inoculated with a high quantity of virus (MOI 0.1 TCID₅₀/cell). This observation might prompt the establishment of new process parameter settings with a higher MOI as a means to achieve maximal virus production/titer stability while ensuring a low level of heterogeneity within the E2 protein. However, an MOI of 0.1 results in significantly higher consumption of GMP working virus seed bank, thus limiting its practical industrial applicability.

Post-infection harvest day only impacted the variation of nucleic acid position 9119 (FIG. 18).

SEQUENCES

Nucleotide sequence of the CHIKV-Δ5nsP3

SEQ ID NO: 1

GATGGCTGCGTGAGACACGCTAGCCTACAGTTTCTTACTGCTCTACTCTGCAAAGCAAGAGATTAATAACCCATCATGGATC

CTGTGTACGTGGACATAGACGCTGACAGCGCTTTTGAAGGCCCTGCAACGTGCGTACCCCATGTTTGAGGTGGAACCAAGG

CAGGTCACACCGAATGACCATGCTAATGCTAGAGCGTTCTCGCATCTAGCTATAAACTAATAGAGCAGGAAATTGACCCCGA

CTCAACCATCCTGGATATCGGCAGTGCGCCAGCAAGGAGGATGATGTGCGACAGGAAGTACCACTGCGTCTGCCCAGTGC

AGTGCGGAAGATCCCGAGAGACTCGCCAATTATGCGAGAAAGCTAGCATCTGCCGAGGAAAGTCTTGGACAGAAACATCT

CTGGAAGATCGGGGACTTACAAGCAGTAATGGCCGTGCCAGACACGAGACGCCAACATTCTGCTTACACACAGACGTCTCA

TGTAGACAGAGACGACGCTGCTATATACCAAGACGCTCTATGCTGTACACGCACCCACGTCGCTATACCACCAGGCGATTAA

AGGGGTCCGAGTGGCGTACTGGGTGGGTTTGACACACCCCGTTTCATGTACAATGCCATGGCGGGTGCCTACCCCTCATAT

CGACAAACTGGGCAGATGAGCAGGTACTGAAGGCTAAGAACATAGGATTATGTTCAACAGACCTGACGGAAGGTAGACGAG

-continued

GCAAGTTGTCTATTATGAGAGGGAAAAAGCTAAAACCGTGCACCGTGTGCTGTTCTCAGTAGGGTCAACGCTCTACCCGGAA
AGCCGCAAGCTACTTAAGAGCTGGCACCTGCCATCGGTGTCCATTAAAGGGCAAACCTCAGCTTCACATGCCGCTGTGATACA
GTGGTTTCGTGTGAGGGCTACGTCGTTAAGAGAATAACGATGAGCCCAGGCCCTTATGGAAAAACACAGGGTATGCGGTAA
CCCACCACGCAGACGGATTCTGTATGTGCAAGACTACCGACACGGTTGACGGCGAAAGAATGTCATTCTCGGTGTGCACATAC
GTGCCGGCGACCATTTGTGATCAAATGACCGGCATCCTTGCTACAGAAGTCACGCCGAGGATGCACAGAAGCTGTTGGTGG
GGCTGAACCAGAGAATAGTGGTTAACGGCAGAACGCAACGGAATACGAACACCATGAAAAATTATCTGCTTCCCGTGGTCGC
CCAAGCCTTCAGTAAGTGGGCAAAGGAGTGCCGGAAAGACATGGAAGATGAAAACTCCTGGGGGTGAGAGAAAGAACT
GACCTGTCTGTCTATGGGCATTCAAGAAGCAGAAAAACACACCGGTCTACAAGAGGCCGTGATACCCAGTCAATTGAGAAGG
TTCAGGCCGAGTTTGACAGCTTTGTGGTACCGAGTCTGTGGTCTGTCGGGTGTCAATCCCTTTGAGGACTAGAATCAAATGGT
TGTTAAGCAAGGTGCCAAAAACCGACCTGATCCCATACAGCGGAGACGCCGAGAAGCCCGGGACGCAGAAAAAGAAGCAG
AGGAAGAACGAGAAGCAGAACTGACTCGCGAAGCCCTACCACTCTACAGGCAGCACAGGAAGATGTTGAGTGAATCGA
CGTGAACAGCTTGAGGACAGAGCGGCGCAGGAATAATAGAGACTCCGAGAGGAGCTATCAAAGTTACTGCCCAACCAAC
AGACCACGTCTGGGAGAGTACCTGGTACTCTCCCGCAGACCGTACTACGTAGCCAGAAGCTCAGTCTGATTACGCTTTGG
CGGAGCAAGTGAAGACGTGCACGCACAACGGACGAGCAGGGAGGTATGCGGTGGAAGCGTACGACGGCCGAGTCTTAGTGC
CCTCAGGCTATGCAATCTCGCCTGAAGACTTCAGAGTCTAAGCGAAAGCGCAACGATGGTGTATAACGAAAGAGAGTTCTGTA
AACAGAAAGCTACACCATATTGCGATGCACGGACCGCCCTGAACACCGACGAAGAGTCTGATGAGCTGGTGAAGGCAGAGA
GGACAGAACACGAGTACGTCTACGACGTGGATCAGAGAAGATGCTGTAAGAAGGAAGAAGCCGAGGACTGGTACTGGTGG
GCGACTTGACTAATCCGCCCTACCAAGAATTGCGATATGAAGGGCTAAAAATCCGCCCTGCCTGCCATACAAAATTGCACTCA
TAGGAGTCTTCGGAGTACCGGGATCTGGCAAGTCAGCTATTATCAAGAACCCTAGTTACCAGGCAGGACCTGGTGACTAGCGG
AAAGAAAGAAAAGTCCCAAGAAATCACACCGACGTGATGAGACAGAGAGGTCTAGAGATATCTGCACGTACGGTTGACTCG
CTGCTCTTGAATGGATGCAACAGACCAGTCGACGTGTTGTACGTAGACGAGGCGTTTGCGTGCCACTCTGGAACGCTACTTGC
TTTGATCGCCTTGGTGAGACCAAGGCAGAAAGTTGTACTTTGTGGTGACCCGAAGCAGTGCGGCTTCTTCAATATGATGCAGA
TGAAAGTCAACTATAATCACAACATCTGCACCAAGTGTACCAACAAAGTATCTCCAGGCGGTGTACACTGCCTGTGACCGCCA
TTGTGTATCGTTGCAATTACGAAGGCAAAATGCGCACTACGAATGAGTACAACAAGCCGATTGTAGTGGACACTACAGGCTCA
ACAAAACCTGACCTGGAGACCTCGTGTAACTGCTTCAAGGGTGGGTAAACAACGTCAAATTGACTATCGTGATACGA
GGTCATGACAGCAGCCGATCCCAAGGGTTAACGAAAAAGGAGTTTACGCAAGTTAGACAAAAAGTTAATGAAAAACCGCTCT
ATGCATCAACGTGAGACAGTCAACGTACTCCTAACGCGTACGGAAGGTAACCTGGTATGGAAGACACTTCCGGCGACCCG
TGGATAAAGACGCTGCAGAACCACCGAAAGGAACTTCAAAGCAACTATTAAGGAGTGGGAGGTGGAGCATGCATCAATAA
TGGCGGGCATCTGCAGTACCAAATGACCTTCGATACATTCCAAAATAAAGCCAACGTTTGTGGGCTAAGAGCTTGGTCCCTA
TCCTCGAAACAGCGGGGATAAACTAAATGATAGGCAGTGGTCTCAGATAATTCAAGCCTTCAAAGAAGACAAAGCATACTCA
CCTGAAGTAGCCCTGAATGAAATATGTACGCGCATGTATGGGGTGGATCTAGACAGCGGGCTATTTCTAAACCGTTGGTGT
TGTGTATTACGCGGATAACCACTGGGATAATAGGCCTGGAGGGAAAAATGTTGCGATTAAACCCGAGGCAGCATCCATTCTAG
AAAGAAAGTATCCATTACAAAAGGGAAGTGAACATCAACAAGCAGATCTGCGTGACTACCAGGAGGATAGAAGACTTTAA
CCCTACCACCAACATCATACCGCCAACAGGAGACTACCACACTCATTAGTGGCCGAACACCGCCAGTAAAAGGGGAAAGAA
TGGAATGGCTGGTTAAACAAGATAAACGGGCCACACGTGCTCCTGGTCAGTGGCTATAACCTTGCACTGCCTACTAAGAGAGTC
ACTTGGGTAGCGCGTTAGGTGTCCGCGAGCGGACTACACATACAACCTAGAGTTGGGTCTGCCAGCAACGCTTGGTAGGT
ATGACCTAGTGGTCATAAACATCCACACACCTTTTCGCATACACCATTACCAACAGTGCCTCGACCACGCAATGAACTGCAAA
TGCTCGGGGGTGACTCATTGAGACTGCTCAAACCGGGCGGCTCTCTATTGATCAGAGCATATGGTTACGCAGATAGAACCAGT
GAACGAGTCATCTGCGTATTGGGACGCAAGTTTAGATCGTCTAGAGCGTTGAAACCACCATGTGTACCAGCAACACTGAGAT
GTTTTTCTATTGACCACTTTGACAATGGCAGAAGGAATTCACAACCTCATGTGATGAACAATCACTGAATGCAGCCTTCGTA

- continued

GGACAGGTCACCCGAGCAGGATGTGCACCGTCGTACCGGGTAAAACGCATGGACATCGCGAAGAACGATGAAGAGTGCCTA
GTCAACGCGCTAACCTCGCGGGTTACCGGGTGGCGGTGTTTGAAGGCAGTATACAAAAATGGCCGGAGTCTTTAAGA
ACAGTGCAACACCAGTGGGAACCGCAAAAACAGTTATGTGCGGTACGTATCCAGTAATCCACGCTGTTGGACCAACTTCTCT
AATTATTGCGAGTCTGAAGGGGACCGGAATTGGCAGCTGCCTATCGAGAAGTCGCAAGGAAGTAAGTAGGCTGGGAGTA
AATAGTGTAGCTATACCTCTCCTCTCCACAGGTGTATACTCAGGAGGGAAAGACAGGCTGACCCAGTCACTGAACCACTCTTT
ACAGCCATGGACTCGACGGATGCAGACGTGGTCATCTACTGCCGCGCAAAAGATGGGAGAAGAAAATATCTGAGGCCATAC
AGATGCGGACCCAAGTAGAGCTGCTGGATGAGCACATCTCCATAGACTGCGATATTGTTGCGGTGCACCCCTGACAGCAGCTTG
GCAGGCAGAAAAGGATACAGCACACGGAAGGCGCACTGTACTCATATCTAGAAGGGACCCGTTTTCATCAGACGGCTGTGG
ATATGGCGGAGATACATACTATGTGGCCAAAGCAAACAGAGGCCAATGAGCAAGTCTGCCTATATGCCCTGGGGGAAAGTAT
TGAATCGATCAGGCAGAAATGCCCGGTGGATGATGCAGACGCATCATCTCCCCCAAACCTGTCCTGTCCTTTGCGGTACGC
TATGACTCCAGAACGCGTCACCCGGCTTCGCATGAACACGTCACAAGCATAATTGTGTGTTCTTCGTTTCCCTCCCAAAGTAC
AAAAAGAGGAGTGCAAAAAGTCAAATGCTCTAAGGTAATGCTATTTGACCACAACCTGCCATCGCGCTAAGTCCAAGGG
CTTATAGAGGTGCCGTGCCGGTAACCTTGCGGCCGTGTCTGATTGGGTAATGAGCACCGTACCTGTCGCGCCGCCAGAAGA
AGGCGAGGGAGAAACCTGACTGTGACATGTGACGAGAGAGAAGGGAATATAACACCCATGGCTAGCGTCCGATTCTTTAGG
GCAGAGCTGTGTCCGGTCGTACAAGAAACAGCGGAGACGCGTGACACAGCAATGTCTCTTCAGGCACCAACCGAGTACCGCA
CGGAACCGAATCATCCGCCGATCTCCTTCGGAGCATCAAGCGAGACGTTCCCCATTACATTTGGGGACTTCAACGAAGGAGAA
ATCGAAAGCTTGCTTCTGAGCTACTAATTTTCGGAGACTTCTTACCAGGAGAAGTGGATGACTTGACAGACAGCGACTGGTC
CACGTGCTCAGACACGGACGACGAGTTAAGACTAGACAGGGCAGGTGGGTATATATTCTCGTCGGACACCGGTCCAGGTCTAT
TTACAACAGAAGTCAGTACGCCAGTCAGTGCTGCCGCTGAACACCCCTGGAGGAATCCACCAGGAGAAGTGTACCCACCTA
AGCTGGATGAAGCAAAGGAGCAACTATTACTTAAGAACTCCAGGAGAGTGCATCCATGGCCAAACAGAAGCAGGTATCAGTC
GCGCAAAGTAGAAAACATGAAAGCAGCAATCATCCAGAGACTAAAGAGAGGCTGTAGACTATACTTAATGTGAGAGCCCCA
AAAGTCCCTACTTACCGGACTACATATCCGGCGCCTGTGTACTCGCCTCCGATCAACGTCCGATTGTCCAATCCCGAGTCCGCA
GTGCGAGCATGCAATGAGTCTTAGCTAGAACTATCCAACTGTCTCATATACCAAATTACCGACGAGTATGATGCATATCTA
GACATGGTGGACGGGTCGGAGAGTTGCCTGGACCGAGCGACATTCAATCCGTCAAAACCTCAGGAGCTACCCGAAACAGCACG
CTTACCACGCGCCCTCCATCAGAAGCGCTGTACCGTCCCCATTCCAGAACACACTACAGAATGTACTGGCAGCAGCCACGAAAA
GAACTGCAACGTACACAGATGAGGAATTACCCACTTTGGACTCAGCAGTATTCAACCTGGAGTGTTCAAAAAATTGCA
TGCAACCAAGAATACTGGGAAGAATTGCTGCGAGCCCTATTAGGATAACAACCTGAGAATTTAGCAACCTATGTTACTAAACTA
AAAGGGCCAAAAGCAGCAGCGCTATTCGCAAAAACCCATAATCTACTGCCACTACAGGAAGTACCAATGGATAGGTTACAGT
AGATATGAAAAGGACGCTAAAGGTGACTCCTGGTACAAAGCATACAGAGGAAAGACCTAAGGTGCAGTTATACAGGCGGC
TGAACCTTGCGCAGCATACTATGTGGGATTACAGAGAGCTGGTTAGGAGGCTGAACGCCGTCTCTTACCCAATGTAC
ATACACTATTTGACATGTCTGCCGAGGATTTGATGCCATCATAGCCGCACACTTTAAGCCAGGAGACACTGTTTTGGAACGG
ACATAGCCTCCTTTGATAAGAGCCAAGATGATTCACTTTCGCTTACTGCTTTGATGCTGTTAGAGGATTTAGGGGTGGATCACT
CCCTGCTCGACTTGATAGAGGCTGCTTCCGAGAGATTTCCAGCTGTACCTACCGACAGGTACGCGCTTCAAGTTCGGCGCC
ATGATGAAATCAGTATGTTCTTAATCTGTTTCGTCAACACATGTTAAACATCACCATCGCCAGCCGAGTGTGGAAGATCGT
CTGACAAAATCCGCGTGCAGCGCTTTCATCGGCGACGACAACATAATACATGGAGTCGTCTCCGATGAATTGATGGCAGCCAG
ATGTGCCACTTTGGATGAACATGGAAGTGAAGATCATAGATGCAGTTGTATCCTTGAAAGCCCTTACTTTTGTGGAGGGTTTAT
ACTGCACGATACTGTGACAGGAACAGCTTCAGAGTGGCAGACCCGCTAAAAAGGCTTTTAACTGGGCAACCCGCTAGCG
GCAGGTGACGAACAAGATGAAGATAGAAGACGAGCGCTGGCTGACGAAGTGATCAGATGGCAACGAACAGGGCTAATTGAT
GAGCTGGAGAAAGCGGTATCTCTAGGTACGAAGTGCAGGGTATATCAGTTGTGGTAATGTCCATGGCCACCTTTGCAAGCTC
CAGATCCAACCTTCGAGAAGCTCAGAGGACCCGTCATAACTTTGTACGGCGGTCTAAATAGGTACGCACTACAGCTACCTATTT
TGCAGAAGCCGACAGCAAGTATCTAAACACTAATCAGCTACAATGGAGTTCATCCCAACCAAACCTTTTACAATAGGAGGTAC

-continued

CAGCCTCGACCCTGGACTCCGCGCCCTACTATCCAAGTCATCAGGCCAGACCGCGCCCTCAGAGGCAAGCTGGGCAACTTGC
CCAGCTGATCTCAGCAGTTAATAAACTGACAATGCGCGCGGTACCACAACAGAAGCCACGCAGGAATCGGAAGAATAAGAAG
CAAAAGCAAAACAACAGGCGCCACAAAACAACACAAATCAAAAGAAGCAGCCACCTAAAAAGAAACCGGCTCAAAAGAAAA
AGAAGCCGGGCGCAGAGAGAGGATGTGCATGAAAAATCGAAAAATGATTGTATTTTCGAAGTCAAGCACGAAGGTAAGGTAA
CAGGTTACGCGTGCCTGGTGGGGACAAAGTAATGAAACAGCACACGTAAAGGGGACCATCGATAACGCGGACCTGGCCA
AACTGGCCTTTAAGCGGTCTAAGTATGACCTTGAATGCGCCAGATACCCGTGCACATGAAGTCCGACGCTTCGAAGTTC
ACCCATGAGAAACCGGAGGGGTACTACAACCTGGCACACGAGCAGTACAGTACTCAGGAGGCCGGTTCACCATCCCTACAG
GTGCTGGCAAAACAGGGGACAGCGGAGACCGGATCTTCGACAACAAGGGACGCGTGGTGGCCATAGTCTTAGGAGGAGCTA
ATGAAGGAGCCGTACAGCCCTCTCGGTGGTGACCTGGAATAAAGACATTTGTACTAAAAACACCCCGAGGGGGCCGAAGA
GTGGAGTCTTGCCATCCAGTTATGTGCTGTTTGGCAAAACACCAGTTCCCTGCTCCAGCCCCCTTGACGCCCCGTGCTGCTAC
GAAAAGGAACCGGAGGAAACCTTACGCATGCTTGAGGACAACGTCTAGACCTGGGTACTATCAGCTGCTACAAGCATCCTT
AACATGTTCTCCCCACCGCCAGCGAGCAGCAGCACCAAGGACAACCTTCAATGTCTATAAAGCCACAAGACCATACTTAGCTCACTG
TCCCGACTGTGGAGAAGGGCACTCGTGCCATAGTCCCGTAGCACTAGAACGCATCAGAAATGAAGCGACAGACGGGACGCTG
AAAATCCAGGTCTCCTTGCAAAATCGGAATAAAGACGGATGACAGCCACGATTGGACCAAGCTGCGTTATATGGACAACCACAT
GCCAGCAGACGCAGAGAGGGCGGGGCTATTTGTAAGAACATCAGCACCGGTGTACGATTACTGGAACAATGGGACACTTCATC
CTGGCCCGATGTCCAAAGGGGAACTCTGACGGTGGGATTCACTGACAGTAGGAAGATTAGTCACTCATGTACGCACCCATT
TCACCAGACCCCTCCTGTGATAGGTGCGGAAAAATTCATTCCCGACCGCAGCAGGTAAAGAGCTACCTTGACGACGTACG
TGCAGAGCACCGCGCAACTACCGAGGAGATAGAGGTACACATGCCCCAGACACCCCTGATCGCACATTAATGTCACAACAG
TCCGGCAACGTAAGATCAGTCAATGGCCAGACGGTGCAGTACAAGTGAATGCGGTGGCTCAAATGAAGGACTAACAA
CTACAGACAAAGTGATTAATAACTGCAAGTTGATCAATGTCTATGCCGCGGTACCAATCACAAAAAGTGGCAGTATAACTCC
CCTCTGGTCCCGCTAATGTGAACCTGGGGACCGAAAAGGAAAAATTCACATCCCGTTTCCGCTGGCAATGTAACATGCAG
GGTGCCATAAGCAAGGAACCCACCGTGACGTACGGGAAAAACCAAGTCATCATGCTACTGTATCCTGACCACCAACACTCC
TGTCTACCGGAATATGGGAGAAGAACCAACTATCAAGAAGAGTGGGTGATGCATAAGAAGGAAGTCGTGCTAACCGTGCC
GACTGAAGGGCTCGAGGTACAGTGGGGCAACACGAGCCGTATAAGTATTGGCCGCGAGTTATCTACAAACGGTACAGCCCAT
GGCCACCCGCATGAGATAATTCTGTATTATTATGAGCTGTACCCACTATGACTGTAGTAGTTGTGTCAGTGGCCACGTTTCATA
CTCTGTGTCGATGGTGGGTATGGCAGCGGGGATGTGCATGTGTGCACGACGAGATGCATCACACCGTATGAAGTACACCCAG
GAGCTACCGTCCCTTTCTGCTTAGCCTAATATGCTGCATCAGAACAGCTAAAGCGGCCACATACCAAGAGGCTGCGATATACC
TGTGGAACGAGCAGCAACCTTTGTTTGGCTACAAGCCCTTATTCGCTGGCAGCCCTGATTGTTCTATGCAACTGTCTGAGAC
TCTTACCATGCTGCTGTAAACGTTGGCTTTTGTAGCCGTAATGAGCGTCGGTGCCCACTGTGAGCGGTACGAACAGTAA
CAGTGATCCCGAACACGGTGGGAGTACCGTATAAGACTCTAGTCAATAGACCTGGCTACAGCCCATGGTATTGGAGATGGA
ACTACTGTCACTGCTTTGGAGCCAACACTATCGCTTGATTACATCAGTGCAGTACAAAACCGTCATCCCGTCTCCGTACGT
GAAGTGCTGCGGTACAGCAGAGTGCAAGGACAAAAACCTACCTGACTACAGCTGTAAGGTCTTCACCGCGCTTACCCATTTA
TGTGGGGCGCGCTACTGCTTCTGCAGCGTGAAAACACGCAAGTTGAGCGAAGCACACGTGGAGAAGTCCGAATCATGCAA
AACAGAATTTGCATCAGCATACAGGCTCATACCGCATCTGCATCAGCTAAGCTCCGCGTCCCTTACCAAGGAAATAACATCAC
TGTAACATGCCTATGCAAACGGCGACCATGCCGTACAGTTAAGGACGCCAAATTCATTGTGGGGCCAATGTCTTCAGCCTGGA
CACCTTTCGACAACAAAATGTGGTGTACAAAGGTGACGTCTATAACATGGACTACCGCCCTTTGGCGCAGGAAGACCAGGA
CAATTTGGCGATATCCAAAGTCGCACACCTGAGAGTAAAGACGTCTATGCTAATACACAACCTGGTACTGCAGAGACCGGCTGT
GGGTACCGTACAGTGCCATACTCTCAGGCACCATCTGGCTTTAAGTATTGGCTAAAAGAACCGGGGCGTGGTGCAGCACA
CAGCACCATTTGGCTGCCAAATAGCAACAAACCCGGTAAGAGCGGTGAACTGCGCCGTAGGGAACATGCCCATCTCCATCGAC
ATACCGGAAGCGGCTTCACTAGGTCGTGACGCGCCCTCTTTAACGAGACATGTCGTGCGAGGTACCAGCCTGCACCCATTTC

-continued

CTCAGACTTTGGGGGCGTCGCCATTATTAATATGCAGCCAGCAAGAAAGGCAAGTGTGCGGTGCATTTCGATGACTAACGCCG
TCACTATTTCGGGAAGCTGAGATAGAAGTTGAAGGGAATTCCTCAGCTGCAATCTCTTTCTCGACGGCCTTAGCCAGCGCCGAA
TTCGCGCTACAAGTCTGTTCTACACAAGTACACTGTGCAGCCGAGTGCACCCCCGAAGGACCACATAGTCAACTACCCGGC
GTCACATACCACCCTCGGGGTCCAGGACATCTCCGCTACGCGATGTTCATGGGTGCAGAAGATCACGGGAGGTGTGGGACTG
GTTGTTGCTGTTGCCGCACTGATTCTAATCGTGGTGTATGCGTGTCTTCAGCAGGCCTAACTTGACAATTAAGTATGAAGG
TATATGTGTCCCTAAGAGACACACTGTACATAGCAATAATCTATAGATCAAAGGGCTACGCAACCCCTGAATAGTAACAAAA
TACAAAATCACTAAAAATTATAAAAAACAGAAAAATACATAAATAGGTATACGTGTCCCTAAGAGACACATTGTATGTAGGTG
ATAAGTATAGATCAAAGGGCCGAATAACCCCTGAATAGTAACAAAAATGAAAAATCAATAAAAAATCATAAATAGAAAAACCA
TAAACAGAAGTAGTTCAAAGGGCTATAAAACCCCTGAATAGTAACAAAAATATAAATAAATAAATAAATAAATAAATAAATAA
TTGGCAAACGAAGAGATGTAGGTACTTAAGCTTCCTAAAGCAGCCGAACCTCACTTTGAGAAGTAGGCATAGCATACCGAAC
TCTTCCACGATTCTCCGAACCCACAGGGACGTAGGAGATGTTATTTTGTGTTTAAATATTTCAAAAAAAAAAAAAAAAAAAAAA

A

Amino acid sequence of E2 protein from LR22006_OPY1 Chikungunya virus strain - amino acids 339-742 from structural polyprotein GenBank Accession: ABD95938.1 (1-1248 aa)

SEQ ID NO: 2

STKDNFNVYKATRPYLAHCPDCGEGHSCCHSPVALERIRNEATDGLTKIQVSLQIGIKTDDSHDWTCLRYMDNHMPADAERAGLFVR
TSAPCTITGTMGHFILARCPKGETLTVGFTDSRKISHSCTHPFHDPPIVIGREKFHSRPQHGKELPCSTYVQSTAATTEEIEVHMPPDT
PDHTLMSQQSGNVKITVNGQTVRYKCNCGGSNEGLTTDDKVINNCKVDQCHAAVTNHKKWQYNSPLVPRNAELGDRKGKIHIPF
PLANVTCTRVKARNPTVTYGNQVIMLLYPDHTLLSYRNMGEENPYQEEWVMHKKEVVLTVPTGLEVTWGNNEPYKYWPQLS
TNGTAHGHPHEIILYYEYPTMTVVVSVATFILLSMVGMAAGMCMCARRRCITPYELTPGATVPFLLSLICCIRTAKA

Some E2 variants identified herein

E168K variant of E2 protein from Chikungunya virus

SEQ ID NO: 3

STKDNFNVYKATRPYLAHCPDCGEGHSCCHSPVALERIRNEATDGLTKIQVSLQIGIKTDDSHDWTCLRYMDNHMPADAERAGLFVR
TSAPCTITGTMGHFILARCPKGETLTVGFTDSRKISHSCTHPFHDPPIVIGREKFHSRPQHGKELPCSTYVQSTAATTEEIKVHMPPDT
PDHTLMSQQSGNVKITVNGQTVRYKCNCGGSNEGLTTDDKVINNCKVDQCHAAVTNHKKWQYNSPLVPRNAELGDRKGKIHIPF
PLANVTCTRVKARNPTVTYGNQVIMLLYPDHTLLSYRNMGEENPYQEEWVMHKKEVVLTVPTGLEVTWGNNEPYKYWPQLS
TNGTAHGHPHEIILYYEYPTMTVVVSVATFILLSMVGMAAGMCMCARRRCITPYELTPGATVPFLLSLICCIRTAKA

G55R variant of E2 protein from Chikungunya virus

SEQ ID NO: 4

STKDNFNVYKATRPYLAHCPDCGEGHSCCHSPVALERIRNEATDGLTKIQVSLQIRIKTDDSHDWTCLRYMDNHMPADAERAGLFVR
TSAPCTITGTMGHFILARCPKGETLTVGFTDSRKISHSCTHPFHDPPIVIGREKFHSRPQHGKELPCSTYVQSTAATTEEIEVHMPPDT
PDHTLMSQQSGNVKITVNGQTVRYKCNCGGSNEGLTTDDKVINNCKVDQCHAAVTNHKKWQYNSPLVPRNAELGDRKGKIHIPF
PLANVTCTRVKARNPTVTYGNQVIMLLYPDHTLLSYRNMGEENPYQEEWVMHKKEVVLTVPTGLEVTWGNNEPYKYWPQLS
TNGTAHGHPHEIILYYEYPTMTVVVSVATFILLSMVGMAAGMCMCARRRCITPYELTPGATVPFLLSLICCIRTAKA

E247K variant of E2 protein from Chikungunya virus

SEQ ID NO: 5

STKDNFNVYKATRPYLAHCPDCGEGHSCCHSPVALERIRNEATDGLTKIQVSLQIGIKTDDSHDWTCLRYMDNHMPADAERAGLFVR
TSAPCTITGTMGHFILARCPKGETLTVGFTDSRKISHSCTHPFHDPPIVIGREKFHSRPQHGKELPCSTYVQSTAATTEEIEVHMPPDT
PDHTLMSQQSGNVKITVNGQTVRYKCNCGGSNEGLTTDDKVINNCKVDQCHAAVTNHKKWQYNSPLVPRNAELGDRKGKIHIPF
PLANVTCTRVKARNPTVTYGNQVIMLLYPDHTLLSYRNMGEENPYQEEWVMHKKEVVLTVPTGLEVTWGNNEPYKYWPQLS
TNGTAHGHPHEIILYYEYPTMTVVVSVATFILLSMVGMAAGMCMCARRRCITPYELTPGATVPFLLSLICCIRTAKA

G82R variant of E2 protein from Chikungunya virus

SEQ ID NO: 6

STKDNFNVYKATRPYLAHCPDCGEGHSCCHSPVALERIRNEATDGLTKIQVSLQIGIKTDDSHDWTCLRYMDNHMPADAERARLFVR
TSAPCTITGTMGHFILARCPKGETLTVGFTDSRKISHSCTHPFHDPPIVIGREKFFISRPQHGKELPCSTYVQSTAATTEEIEVHMPPDT
PDHTLMSQQSGNVKITVNGQTVRYKCNCGGSNEGLTTDDKVINNCKVDQCHAAVTNHKKWQYNSPLVPRNAELGDRKGKIHIPF

-continued

PLANVTCTRVKARNPTVTYGNQVIMLLYPDHPTLLSYRNMGEENPYQEEWVMHKKEWLTVPTEGLEVTWGNNEPYKYWPQLS
 TNGTAHGHPHEIILYLYEYPTMTVVVSVATFILLSMVGMAAGMCMCARRRCITPYELTPGATVPFLLSLICCIRTAKA
 H232Y variant of E2 protein from Chikungunya virus
 SEQ ID NO: 7
 STKDNFNVYKATRPYLAHCPDCGEGHSCSPVALERIRNEATDGLKIQVSLQIGIKTDDSHDWTCLRYMDNHPADAERAGLFVR
 TSAPCTITGTMGHFILARCPKGETLTVGFTDSRKISHSCTHPFHDDPPVIGREKFSRPQHGKELPCSTYVQSTAATTEEVHMPDPT
 PDHTLMSQQSGNVKITVNGQTVRYKCNCGSGNEGLTTTDDKVINNCKVDQCHAAVTN~~Y~~KKWQYNSPLVPRNAELGDRKGKIHIPFP
 LANVTCTRVKARNPTVTYGNQVIMLLYPDHPTLLSYRNMGEENPYQEEWVMHKKEVLTVPTEGLEVTWGNNEPYKYWPQLST
 NNGTAHGHPHEIILYLYEYPTMTVVVSVATFILLSMVGMAAGMCMCARRRCITPYELTPGATVPFLLSLICCIRTAKA
 Primer 1F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 8
 TTAGGATCCGATGGCTGCGTGAGACAC
 Primer 1R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 9
 TAACTCGAGCCGTCAGGTCTGTTGAACAT
 Primer 2F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 10
 TTAGGATCCTACCACCAGCGATTAAAG
 Primer 2R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 11
 TAACTCGAGCTTTGCCCACTTACTGAAGG
 Primer 3F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 12
 TTAGGATCCTGCTACAGAAAGTCACGCC
 Primer 3R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 13
 TAACTCGAGGCCAAAGCGTGAATCAG
 Primer 4F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 14
 TTAGGATCCAACAGCTTGAGGACAGAGCG
 Primer 4R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 15
 TAACTCGAGCTCTGTCTCATCAGTCGG
 Primer 5F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 16
 TTAGGATCCAAATTGCAGTCATAGGAGTCTTC
 Primer 5R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 17
 TAACTCGAGAGTACGTTGACGTGCTCTGA
 Primer 6F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 18
 TTAGGATCCGTGGGTAAACAAGTCAAA
 Primer 6R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 19
 TAACTCGAGGGTTAAAGTCTTCTATCCTCCTGG
 Primer 7F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 20
 TTAGGATCCGATAACCACTGGGATAATAGG
 Primer 7R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 21
 TAACTCGAGAGTTGTGAAATTCCTTCTGCC
 Primer 8F for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 22
 TTAGGATCCCGCAGATAGAACCAGTGAAC
 Primer 8R for CHIKV-Δ5nsP3 sequencing
 SEQ ID NO: 23
 TAACTCGAGCAGCAGCTCTACTTGGGTC

-continued

Primer 9F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCAGGAGGAAAGACAGGCT

Primer 9R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGCCCTCGCCTTCTTCTG

Primer 10F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCCAAATAGAAGGAGTGCAAAAAG

Primer 10R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGCCTGAGTTTCTTAAGTAATAGTTGC

Primer 11F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCACCGGTCCAGGTCATTTA

Primer 11R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGGCAGCAAATTCTTCCCAG

Primer 12F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCCCATTCCAGAACACTACAG

Primer 12R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGATACCTGATTTCATCATGGC

Primer 13F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCCCTTTGATAAGAGCCAAGATG

Primer 13R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGTACAAAGTTATGACGGGTCCT

Primer 14F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCCAACGAACAGGGCTAATTG

Primer 14R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGGACCGCTTAAAGGCCAG

Primer 15F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCGTGCATGAAATCGAAANTG

Primer 15R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGTGGTCTTGCGCTTTATAGACA

Primer 16F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCAACCGGAGGAAACCTTAC

Primer 16R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGGTACCGCACCGTCTGG

Primer 17F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCAGCTACCTTGACGACGT

Primer 17R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGCCCACCATCGACAGG

Primer 18F for CHIKV-Δ5nsP3 sequencing

TTAGGATCCCGAGCCGTATAAGTATTGGC

Primer 18R for CHIKV-Δ5nsP3 sequencing

TAACTCGAGCGCCGGTGAAGACCTTAC

SEQ ID NO: 24

SEQ ID NO: 25

SEQ ID NO: 26

SEQ ID NO: 27

SEQ ID NO: 28

SEQ ID NO: 29

SEQ ID NO: 30

SEQ ID NO: 31

SEQ ID NO: 32

SEQ ID NO: 33

SEQ ID NO: 34

SEQ ID NO: 35

SEQ ID NO: 36

SEQ ID NO: 37

SEQ ID NO: 38

SEQ ID NO: 39

SEQ ID NO: 40

SEQ ID NO: 41

SEQ ID NO: 42

SEQ ID NO: 43

-continued

Primer 19F for CHIKV-Δ5nsP3 sequencing

SEQ ID NO: 44

TTAGGATCCACTACTGTCAGTCACTTTGGAGC

Primer 19R for CHIKV-Δ5nsP3 sequencing

SEQ ID NO: 45

TAACTCGAGTACCGGGTTTGTGCTATTT

Primer 20F for CHIKV-Δ5nsP3 sequencing

SEQ ID NO: 46

TTAGGATCCCACAACCTGGTACTGCAGAGAC

Primer 20R for CHIKV-Δ5nsP3 sequencing

SEQ ID NO: 47

TAACTCGAGGCGTAGCCCTTTGATCTATAG

Primer 21F for CHIKV-Δ5nsP3 sequencing

SEQ ID NO: 48

TTAGGATCCGGTGCTATGCGTGTGCGT

Primer 21R for CHIKV-Δ5nsP3 sequencing

SEQ ID NO: 49

TAACTCGAGATCTCCTACGTCCCTGTGGG

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 49

<210> SEQ ID NO 1

<211> LENGTH: 11674

<212> TYPE: DNA

<213> ORGANISM: Chikungunya virus

<400> SEQUENCE: 1

gatggctgcg tgagacacac gtagcctacc agtttcttac tgctctactc tgcaaagcaa	60
gagattaata acccatcatg gatcctgtgt acgtggacat agacgctgac agcgcctttt	120
tgaaggccct gcaacgtgcg taccocatgt ttgaggtgga accaaggcag gtcacaccga	180
atgaccatgc taatgctaga gcgttctcgc atctagctat aaaactaata gacgaggaaa	240
ttgacccoga ctcaaccatc ctggatatcg gcagtgcgcc agcaaggagg atgatgtcgg	300
acaggaagta ccactgcgtc tgcccgatgc gcagtgcgga agatcccag agactcgcca	360
attatgcgag aaagctagca tctgccgag gaaaagtcct ggacagaaac atctctggaa	420
agatcgggga cttacaagca gtaatggccg tgccagacac ggagacgcca acattctgct	480
tacacacaga cgtctcatgt agacagagag cagacgtcgc tatatacaca gacgtctatg	540
ctgtacacgc acccacgtcg ctataccacc aggcgattaa aggggtccga gtggcgtact	600
gggttggtt cgacacaacc ccgttcatgt acaatgccat ggcgggtgcc taccctcat	660
actcgacaaa ctgggcagat gacgaggtac tgaaggctaa gaacatagga ttatgttcaa	720
cagacctgac ggaaggtaga cgaggcaagt tgtctattat gagagggaaa aagctaaaac	780
cgtgcgaccg tgtgtgttcc tcagttaggt caacgtctca cccggaaaag cgcaagctac	840
ttaagagctg gcacctgcca tcggtgttcc atttaaaggg caaactcagc ttcacatgcc	900
gctgtgatac agtggtttcg tgtgagggct acgtcgttaa gagaataacg atgagcccag	960
gcctttatgg aaaaaccaca gggatgcgg taaccaccca cgcagacgga ttctgatgt	1020
gcaagactac cgacacggtt gacggcgaaa gaatgtcatt ctcggtgtgc acatacgtgc	1080
cggcgaccat ttgtgatcaa atgaccggca tccttctac agaagtcacg ccggaggatg	1140
cacagaagct gttggtgggg ctgaaccaga gaatagtgtg taacggcaga acgcaacgga	1200
atcgaacac catgaaaaat tatctgcttc ccgtggtcgc ccaagccttc agtaagtggg	1260

-continued

caaaggagtg	ccggaagac	atggaagatg	aaaaactcct	gggggtcaga	gaaagaacac	1320
tgacctgtg	ctgtctatgg	gcattcaaga	agcagaaaac	acacacggtc	tacaagaggc	1380
ctgataccca	gtcaattcag	aagggttcagg	ccgagtttga	cagctttgtg	gtaccgagtc	1440
tgtggtcgtc	cgggttgtca	atccctttga	ggactagaat	caaaggttg	ttaagcaagg	1500
tgccaaaaac	cgacctgatc	ccatacagcg	gagacgcccg	agaagcccg	gacgcagaaa	1560
aagaagcaga	ggaagaacga	gaagcagaac	tgactcgcg	agccctacca	cctctacagg	1620
cagcacagga	agatgttcag	gtcgaaatcg	acgtggaaca	gcttgaggac	agagcgggcg	1680
caggaataat	agagactccg	agaggagcta	tcaaagttac	tgcccaacca	acagaccacg	1740
tcgtgggaga	gtacctggta	ctctcccgc	agaccgtact	acgtagccag	aagctcagtc	1800
tgattcacgc	tttgccggag	caagtgaaga	cgtgcacgca	caacggacga	gcagggaggt	1860
atgcggtcga	agcgtacgac	ggccgagtc	tagtgcctc	aggctatgca	atctcgctg	1920
aagacttcca	gagtctaagc	gaaagcgcaa	cgatggtgta	taacgaaaga	gagttcgtaa	1980
acagaaagct	acaccatatt	gcgatgcacg	gaccagccct	gaacaccgac	gaagagtcgt	2040
atgagctggt	gagggcagag	aggacagaac	acgagtacgt	ctacgacgtg	gatcagagaa	2100
gatgctgtaa	gaaggaagaa	cccgcaggac	tggtactggt	ggcgacttg	actaatccgc	2160
cctaccacga	attcgcatat	gaagggctaa	aaatccgcc	tgctgcca	tacaaaattg	2220
cagtcatagg	agtcctcgga	gtaccgggat	ctggcaagtc	agctattatc	aagaacctag	2280
ttaccaggca	ggacctggtg	actagcggaa	agaaagaaaa	ctgccaagaa	atcaccaccg	2340
acgtgatgag	acagagaggt	ctagagatat	ctgcacgtac	ggttgactcg	ctgctcttga	2400
atggatgcaa	cagaccagtc	gacgtgtgt	acgtagacga	ggcgtttcg	tgccactctg	2460
gaacgctact	tgttttgatc	gccttggtga	gaccaaggca	gaaagttgta	ctttgtggtg	2520
acccgaagca	gtgcggcttc	ttcaatatga	tgcatagaa	agtcactat	aatcacacaa	2580
tctgcaccca	agtgtaccac	aaaagtatct	ccaggcggtg	tacactgcct	gtgaccgcca	2640
ttgtgtcatc	gttgcatcac	gaaggcaaaa	tgcgactac	gaatgagtac	aacaagccga	2700
ttgtagtgga	cactacagcg	tcaacaaaa	ctgaccctgg	agacctcgtg	ttaactgtct	2760
tcagaggggtg	ggttaaacaa	ctgcaaattg	actatcgtgg	atacgaggtc	atgacagcag	2820
ccgcatccca	agggtaaac	agaaaaggag	tttacgcagt	tagacaaaa	gttaaatgaa	2880
acccgtctta	tgcataacg	tcagagcacg	tcaacgtact	cctaacgcgt	acggaaggta	2940
aactggtatg	gaagacactt	tccggcgacc	cgtggataaa	gacgctgcag	aaccaccga	3000
aaggaaactt	caaagcaact	attaaggagt	gggaggtgga	gcctgcacga	ataatggcgg	3060
gcctctgcag	tcaccaaattg	accttcgata	cattccaaaa	taaagccaac	gtttgttggg	3120
ctaagagctt	ggctccctatc	ctcgaaacag	cggggataaa	actaatgat	aggcagtggt	3180
ctcagataat	tcaagccttc	aaagaagaca	aagcatactc	acctgaagta	gcctgaatg	3240
aaatatgtac	gcgcgtgtat	gggtggatc	tagacagcgg	gctatcttct	aaaccgttgg	3300
tgtctgtgta	ttacgcggat	aaccactggg	ataataggcc	tggagggaaa	atgttcggat	3360
ttaaccccca	ggcagcatcc	attctagaaa	gaaagtatcc	attcacaaaa	gggaagtgga	3420
acatcaacaa	gcagatctgc	gtgactacca	ggaggataga	agactttaac	cctaccacca	3480
acatcatacc	ggccaacagg	agactaccac	actcattagt	ggccgaacac	cgccagtaa	3540
aaggggaaag	aatggaatgg	ctggttaaca	agataaacgg	ccaccacgtg	ctcctggtca	3600
gtggctataa	ccttgcactg	cctactaaga	gagtcacttg	ggtagcgccg	ttaggtgtcc	3660

-continued

gcggagcgga ctacacatac aacctagagt tgggtctgcc agcaacgctt ggtaggtatg	3720
acctagtggc cataaacatc cacacacctt ttgcataca ccattaccaa cagtgcgtcg	3780
accacgcaat gaaactgcaa atgctcgggg gtgactcatt gagactgctc aaaccgggcg	3840
gctctctatt gatcagagca tatggttacg cagatagaac cagtgaacga gtcactgctg	3900
tattgggacg caagtttaga tcgtctagag cgttgaaacc accatgtgtc accagcaaca	3960
ctgagatggt tttcctatto agcaactttg acaatggcag aaggaatttc acaactcatg	4020
tcatgaacaa tcaactgaat gcagccttcg taggacaggt caccgcagca ggatgtgcac	4080
cgctgtaccg ggtaaaacgc atggacatcg cgaagaacga tgaagagtgc gtagtcaacg	4140
ccgctaacc tcgcgggtta ccgggtggcg gtgtttgcaa ggcagtatac aaaaaatggc	4200
cggagtcctt taagaacagt gcaacaccag tgggaaccgc aaaaacagtt atgtgcggta	4260
cgtatccagt aatccacgct gttggacca acttctctaa ttattcggag tctgaagggg	4320
accgggaatt ggcagctgcc tatcgagaag tcgcaaagga agtaactagg ctgggagtaa	4380
atagtgtagc tatacctctc ctctccacag gtgtatactc aggagggaaa gacaggctga	4440
cccagtcact gaaccacctc ttacagcca tggactcgac ggatgcagac gtggctcatc	4500
actgcccgca caaagaatgg gagaagaaaa tatctgaggc catacagatg cggacccaag	4560
tagagctgct ggatgagcac atctccatag actgcgatat tgttcgctg caccctgaca	4620
gcagcttggc aggcagaaaa ggatacagca ccacggaagg cgcactgtac tcatatctag	4680
aagggaaccg ttttcatcag acggctgtgg atatggcgga gatacatact atgtggccaa	4740
agcaaacaga ggccaatgag caagtctgcc tatatgcctc gggggaaagt attgaatcga	4800
tcaggcgaaa atgcccgtg gatgatgcag acgcatcatc tcccccaaa actgtcccgt	4860
gcctttgcg ttacgctatg actccagaac gcgtcaccgc gcttcgcatg aaccacgtca	4920
caagcataat tgtgtgttct tcgtttcccc tcccaaagta caaaatagaa ggagtgcaa	4980
aagtcaaatg ctctaaggta atgctatttg accacaacgt gccatcgcgc gtaagtccaa	5040
gggcttatag aggtgccgct gccggtaacc ttgcggccgt gtctgattgg gtaatgagca	5100
ccgtacctgt cgcgcgcgcc agaagaaggc gagggagaaa cctgactgtg acatgtgacg	5160
agagagaagg gaatataaca cccatggcta gcgtccgatt ctttagggca gagctgtgtc	5220
cggctgtaca agaacacgag gagacgcgtg acacagcaat gtctcttcag gcaccaccga	5280
gtaccgcac ggaaccgaat catccgcga tctccttcgg agcatcaagc gagacgttcc	5340
ccattacatt tggggacttc aacgaaggag aaatcgaaag cttgtcttct gagctactaa	5400
ctttcggaga cttcttacca ggagaagtgg atgacttgac agacagcgac tggtecacgt	5460
gctcagacac ggacgacgag ttaagactag acagggcagg tgggtatata ttctcgtcgg	5520
acaccggtcc aggtcattta caacagaagt cagtacgcca gtcagtgtg cgggtgaaca	5580
ccctggagga agtccacgag gagaagtgtt acccacctaa gotggatgaa gcaaaggagc	5640
aactattact taagaaactc caggagagtg catccatggc caacagaagc aggtatcagt	5700
cgcgcaaagt agaaaacatg aaagcagcaa tcatccagag actaaagaga ggctgtagac	5760
tatacttaat gtcagagacc caaaaagtcc ctacttaccg gactacatat ccggcgctg	5820
tgtactcgcc tccgatcaac gtccgattgt ccaatcccga gtccgcagtg gcagcatgca	5880
atgagttctt agctagaaac tatccaactg tctcatcata ccaaattacc gacgagtatg	5940
atgcatactc agacatggcg gacgggtcgg agagttgcct ggaccgagcg acattcaatc	6000

-continued

cgctaaaact caggagctac ccgaacacgc acgcttacca cgcgcctcc atcagaagcg	6060
ctgtaccgtc cccattccag aacacactac agaattgtact ggcagcagcc acgaaaagaa	6120
actgcaacgt cacacagatg aggggaattac ccactttgga ctcagcagta ttcaacgtgg	6180
agtgtttcaa aaaattcgca tgcaaccaag aatactggga agaatttgct gccagcccta	6240
ttaggataac aactgagaat ttagcaacct atgttactaa actaaaaggg ccaaaagcag	6300
cagcgctatt cgcaaaaacc cataatctac tgccactaca ggaagtacca atggataggt	6360
tcacagtaga tatgaaaagg gacgtaaagg tgactcctgg tacaagcat acagaggaaa	6420
gacctaaagt gcaggttata caggcggtg aacccttggc gacagcatc ctatgtggga	6480
ttcacagaga gctggttagg aggtgaacg ccgtcctcct acccaatgta catacactat	6540
ttgacatgtc tgccgaggat ttgatgccca tcatagccgc acactttaag ccaggagaca	6600
ctgttttggg aacggacata gcctccttg ataagagcca agatgattca cttgcgctta	6660
ctgctttgat gctgttagag gatttagggg tggatcactc cctgctggac ttgatagagg	6720
ctgctttcgg agagatttcc agctgtcacc taccgacagg tacgcgcttc aagtctggcg	6780
ccatgatgaa atcaggtatg ttctaactc tgttcgtcaa cacattgtta aacatcacca	6840
tcgccagccg agtgctggaa gatcgtctga caaaatccgc gtgcgcggcc ttcacggcg	6900
acgacaacat aatacatgga gtgctctccg atgaattgat ggcagccaga tgtgccactt	6960
ggatgaacat ggaagtgaag atcatagatg cagttgtatc cttgaaagcc ccttactttt	7020
gtggagggtt tatactgcac gatactgtga caggaacagc ttgcagagtg gcagaccgc	7080
taaaaaggct ttttaactg ggcaaacgc tagcggcagg tgacgaacaa gatgaagata	7140
gaagacgagc gctggtgac gaagtgatca gatggcaacg aacagggcta attgatgagc	7200
tggagaaagc ggtatactct aggtacgaag tgcagggtat atcagttgtg gtaatgtcca	7260
tggccacctt tgcaagctcc agatccaact tcgagaagct cagaggaccc gtcataactt	7320
tgtacggcgg tcctaaatag gtacgcacta cagctaccta ttttcagaa gccgacagca	7380
agtatctaaa cactaatcag ctacaatgga gttcatccca acccaaactt ttacaatag	7440
gaggtaccag cctcgacct ggactccgcg ccctactatc caagtcacga gccccagacc	7500
cgccctcag aggcgaagctg ggcaacttgc ccagctgac tcagcagta ataaactgac	7560
aatgcgcgcg gtaccacaac agaagccagc caggaatcgg aagaataaga agcaaaagca	7620
aaaacaacag gcgccacaaa acaacacaaa tcaaaagaag cagccaccta aaaagaaacc	7680
ggctcaaaag aaaaagaagc cgggccgcag agagaggatg tgcataaaaa tcgaaaatga	7740
ttgtatttcc gaagtcaagc acgaaggtaa ggtaacaggt tacgcgtgcc tgggtgggga	7800
caaagtaatg aaaccagcac acgtaaaggg gaccatcgat aacgcggacc tggccaaact	7860
ggcctttaag cggatcatct agtatgacct tgaatgcgcg cagatacccg tgcacatgaa	7920
gtccgacgct tcgaagttca cccatgagaa accggagggg tactacaact ggcaccacgg	7980
agcagtacag tactcaggag gccggttcac catcctaca ggtgctggca aaccagggga	8040
cagcggcaga ccgactctcg acaacaaggg acgcgtgggt gccatagtct taggaggagc	8100
taatgaagga gcccgtagc ccctctcggt ggtgacctgg aataaagaca ttgtcactaa	8160
aatcaccccc gagggggcgg aagagtggag tcttgccatc ccagttatgt gctgttggc	8220
aaacaccacg ttccctgct cccagcccc ttgcacgccc tgctgctacg aaaaggaacc	8280
ggaggaaacc ctacgcatgc ttgaggacaa cgtcatgaga cctgggtact atcagctgct	8340
acaagcatcc ttaacatgtt ctccccaccg ccagcgacgc agcaccaagg acaacttcaa	8400

-continued

tgtctataaa gccacaagac catacttagc tcaactgtccc gactgtggag aagggcactc	8460
gtgccatagt ccgtagcac tagaacgcat cagaaatgaa gcgacagacg ggacgctgaa	8520
aatccaggtc tccttgcaaa tcggaataaa gacggatgac agccacgatt ggaccaagct	8580
gcgttatatg gacaaccaca tgccagcaga cgcagagagg gcggggctat ttgtaagaac	8640
atcagcaccg tgtacgatta ctggaacaat gggacacttc atcctggccc gatgtccaaa	8700
aggggaaact ctgacgggtg gattcactga cagtaggaag attagtcaact catgtacgca	8760
cccatttcac cagcaccctc ctgtgatagg tcgggaaaaa ttccattccc gaccgcagca	8820
cggtaaagag ctaccttgca gcacgtacgt gcagagcacc gccgcaacta ccgaggagat	8880
agaggtagac atgccccag acacccctga tcgcacatta atgtcacaac agtccggcaa	8940
cgtaaagatc acagtcaatg gccagacggt gcggtacaag tgtaattgctg gtggctcaaa	9000
tgaaggacta acaactacag acaaagtgat taataactgc aaggttgatc aatgtcatgc	9060
cgcggtcacc aatcacaaaa agtggcagta taactccctc ctggtccgc gtaatgctga	9120
acttggggac cgaaaaggaa aaattcacat ccggtttccg ctggcaaatg taacatgcag	9180
ggtgcctaaa gcaaggaacc ccaccgtgac gtacgggaaa aaccaagtca tcatgctact	9240
gtatcctgac caccacacac tcctgtccta ccggaatatg ggagaagaac caaactatca	9300
agaagagtgg gtgatgcata agaaggaagt cgtgctaacc gtgcccactg aagggctcga	9360
ggtcacgtgg ggcaacaacg agccgtataa gtattggccg cagttatcta caaacggtac	9420
agcccatggc caccgcgatg agataattct gtattattat gagctgtacc ccactatgac	9480
tgtagtagtt gtgtcagtgg ccacgttcac actcctgtcg atggtgggta tggcagcggg	9540
gatgtgcatg tgtgcacgac gcagatgcat cacaccgtat gaactgacac caggagctac	9600
cgtccctttc ctgcttagcc taatatgtcg catcagaaca gctaaagcgg ccacatacca	9660
agaggctgcg atatacctgt ggaacgagca gcaacctttg ttttggtac aagcccttat	9720
tccgctggca gccctgattg ttctatgcaa ctgtctgaga ctcttaccat gctgctgtaa	9780
aacgttggct tttttagccg taatgagcgt cggtgccac actgtgagcg cgtacgaaca	9840
cgtaacagtg atcccgaaca cgggtggagt accgtataag actctagtca atagacctgg	9900
ctacagcccc atggtattgg agatggaact actgtcagtc actttggagc caaactatc	9960
gcttgattac atcacgtgcg agtacaaaac cgtcatcccg tctccgtacg tgaagtgtg	10020
cggtagagca gagtgaagc acaaaaacct acctgactac agctgtaagg tcttcaccg	10080
cgtctaccca tttatgtggg gcggcgcccta ctgcttctgc gacgtgaaa acacgcagtt	10140
gagcgaagca cagctggaga agtccgaatc atgcaaaaca gaatttgcac cagcatagag	10200
ggctcatacc gcatctgcat cagctaagct ccgcgtcctt taccaaggaa ataacatcac	10260
tgtaactgcc tatgcaaacg gcgaccatgc cgtcacagtt aaggacgcca aattcattgt	10320
ggggccaatg tcttcagcct ggacaccttt cgacaacaaa atttgtgtgt acaaagggtga	10380
cgtctataac atggactacc cgccctttgg cgcaggaaga ccaggacaat ttggcgatat	10440
ccaaagtgc acacctgaga gtaaagacgt ctatgctaac acacaactgg tactgcagag	10500
accggctgtg ggtacggtac acgtgccata ctctcaggca ccatctggct ttaagtattg	10560
gctaaaagaa cgcggggcgt cgctgcagca cacagcacca tttggctgcc aaatagcaac	10620
aaacccggtg agagcggtga actgcgcgt agggaacatg cccatctcca tcgacatacc	10680
ggaagcggcc ttcactaggg tcgtcgacgc gccctcttta acggacatgt cgtgcgaggt	10740

-continued

```

accagcctgc acccattcct cagactttgg gggcgctgcc attattaaat atgcagccag 10800
caagaaaggc aagtgtgctg tgcattcgat gactaacgcc gtcactattc gggaagctga 10860
gatagaagtt gaagggaatt ctcagctgca aatctctttc tcgacggcct tagccagcgc 10920
cgaattccgc gtacaagtct gttctacaca agtacactgt gcagccgagt gccaccccc 10980
gaaggaccac atagtcaact acccgggctc acataccacc ctcggggtcc aggacatctc 11040
cgctacggcg atgtcatggg tgcagaagat cacgggaggt gtgggactgg ttgttgctgt 11100
tgccgcactg attctaactg tgggtgctatg cgtgtcgttc agcaggcact aacttgacaa 11160
ttaagtatga aggtatatgt gtcccctaag agacacactg tacatagcaa ataatctata 11220
gatcaaaggg ctacgcaacc cctgaatagt aacaaaatac aaaatcacta aaaattataa 11280
aaacagaaaa atacataaat aggtatacgt gtcccctaag agacacattg tatgtagggtg 11340
ataagtatag atcaaagggc cgaataaccc ctgaatagta acaaaatatg aaaatcaata 11400
aaaatcataa aatagaaaaa ccataaacag aagtagttca aagggtataa aaaccctga 11460
atagtaacaa aacataaaat taataaaaat caaatgaata ccataattgg caaacggaag 11520
agatgtagggt acttaagctt cctaaaagca gccgaactca ctttgagaag taggcatagc 11580
ataccgaact cttccacgat tctccgaacc cacagggacg taggagatgt tattttgttt 11640
ttaatatattc aaaaaaaaaa aaaaaaaaaa aaaa 11674

```

```

<210> SEQ ID NO 2
<211> LENGTH: 423
<212> TYPE: PRT
<213> ORGANISM: Chikungunya virus

```

```

<400> SEQUENCE: 2

```

```

Ser Thr Lys Asp Asn Phe Asn Val Tyr Lys Ala Thr Arg Pro Tyr Leu
1           5           10           15
Ala His Cys Pro Asp Cys Gly Glu Gly His Ser Cys His Ser Pro Val
20           25           30
Ala Leu Glu Arg Ile Arg Asn Glu Ala Thr Asp Gly Thr Leu Lys Ile
35           40           45
Gln Val Ser Leu Gln Ile Gly Ile Lys Thr Asp Asp Ser His Asp Trp
50           55           60
Thr Lys Leu Arg Tyr Met Asp Asn His Met Pro Ala Asp Ala Glu Arg
65           70           75           80
Ala Gly Leu Phe Val Arg Thr Ser Ala Pro Cys Thr Ile Thr Gly Thr
85           90           95
Met Gly His Phe Ile Leu Ala Arg Cys Pro Lys Gly Glu Thr Leu Thr
100          105          110
Val Gly Phe Thr Asp Ser Arg Lys Ile Ser His Ser Cys Thr His Pro
115          120          125
Phe His His Asp Pro Pro Val Ile Gly Arg Glu Lys Phe His Ser Arg
130          135          140
Pro Gln His Gly Lys Glu Leu Pro Cys Ser Thr Tyr Val Gln Ser Thr
145          150          155          160
Ala Ala Thr Thr Glu Glu Ile Glu Val His Met Pro Pro Asp Thr Pro
165          170          175
Asp His Thr Leu Met Ser Gln Gln Ser Gly Asn Val Lys Ile Thr Val
180          185          190
Asn Gly Gln Thr Val Arg Tyr Lys Cys Asn Cys Gly Gly Ser Asn Glu
195          200          205

```

-continued

Gly Leu Thr Thr Thr Asp Lys Val Ile Asn Asn Cys Lys Val Asp Gln
 210 215 220
 Cys His Ala Ala Val Thr Asn His Lys Lys Trp Gln Tyr Asn Ser Pro
 225 230 235 240
 Leu Val Pro Arg Asn Ala Glu Leu Gly Asp Arg Lys Gly Lys Ile His
 245 250 255
 Ile Pro Phe Pro Leu Ala Asn Val Thr Cys Arg Val Pro Lys Ala Arg
 260 265 270
 Asn Pro Thr Val Thr Tyr Gly Lys Asn Gln Val Ile Met Leu Leu Tyr
 275 280 285
 Pro Asp His Pro Thr Leu Leu Ser Tyr Arg Asn Met Gly Glu Glu Pro
 290 295 300
 Asn Tyr Gln Glu Glu Trp Val Met His Lys Lys Glu Val Val Leu Thr
 305 310 315 320
 Val Pro Thr Glu Gly Leu Glu Val Thr Trp Gly Asn Asn Glu Pro Tyr
 325 330 335
 Lys Tyr Trp Pro Gln Leu Ser Thr Asn Gly Thr Ala His Gly His Pro
 340 345 350
 His Glu Ile Ile Leu Tyr Tyr Tyr Glu Leu Tyr Pro Thr Met Thr Val
 355 360 365
 Val Val Val Ser Val Ala Thr Phe Ile Leu Leu Ser Met Val Gly Met
 370 375 380
 Ala Ala Gly Met Cys Met Cys Ala Arg Arg Arg Cys Ile Thr Pro Tyr
 385 390 395 400
 Glu Leu Thr Pro Gly Ala Thr Val Pro Phe Leu Leu Ser Leu Ile Cys
 405 410 415
 Cys Ile Arg Thr Ala Lys Ala
 420

<210> SEQ ID NO 3
 <211> LENGTH: 423
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: mutation arising from Vero passaging

<400> SEQUENCE: 3

Ser Thr Lys Asp Asn Phe Asn Val Tyr Lys Ala Thr Arg Pro Tyr Leu
 1 5 10 15
 Ala His Cys Pro Asp Cys Gly Glu Gly His Ser Cys His Ser Pro Val
 20 25 30
 Ala Leu Glu Arg Ile Arg Asn Glu Ala Thr Asp Gly Thr Leu Lys Ile
 35 40 45
 Gln Val Ser Leu Gln Ile Gly Ile Lys Thr Asp Asp Ser His Asp Trp
 50 55 60
 Thr Lys Leu Arg Tyr Met Asp Asn His Met Pro Ala Asp Ala Glu Arg
 65 70 75 80
 Ala Gly Leu Phe Val Arg Thr Ser Ala Pro Cys Thr Ile Thr Gly Thr
 85 90 95
 Met Gly His Phe Ile Leu Ala Arg Cys Pro Lys Gly Glu Thr Leu Thr
 100 105 110
 Val Gly Phe Thr Asp Ser Arg Lys Ile Ser His Ser Cys Thr His Pro
 115 120 125
 Phe His His Asp Pro Pro Val Ile Gly Arg Glu Lys Phe His Ser Arg
 130 135 140

-continued

```

Pro Gln His Gly Lys Glu Leu Pro Cys Ser Thr Tyr Val Gln Ser Thr
145                      150                      155                      160

Ala Ala Thr Thr Glu Glu Ile Lys Val His Met Pro Pro Asp Thr Pro
                      165                      170                      175

Asp His Thr Leu Met Ser Gln Gln Ser Gly Asn Val Lys Ile Thr Val
                      180                      185                      190

Asn Gly Gln Thr Val Arg Tyr Lys Cys Asn Cys Gly Gly Ser Asn Glu
                      195                      200                      205

Gly Leu Thr Thr Thr Asp Lys Val Ile Asn Asn Cys Lys Val Asp Gln
210                      215                      220

Cys His Ala Ala Val Thr Asn His Lys Lys Trp Gln Tyr Asn Ser Pro
225                      230                      235                      240

Leu Val Pro Arg Asn Ala Glu Leu Gly Asp Arg Lys Gly Lys Ile His
                      245                      250                      255

Ile Pro Phe Pro Leu Ala Asn Val Thr Cys Arg Val Pro Lys Ala Arg
                      260                      265                      270

Asn Pro Thr Val Thr Tyr Gly Lys Asn Gln Val Ile Met Leu Leu Tyr
                      275                      280                      285

Pro Asp His Pro Thr Leu Leu Ser Tyr Arg Asn Met Gly Glu Glu Pro
290                      295                      300

Asn Tyr Gln Glu Glu Trp Val Met His Lys Lys Glu Val Val Leu Thr
305                      310                      315                      320

Val Pro Thr Glu Gly Leu Glu Val Thr Trp Gly Asn Asn Glu Pro Tyr
                      325                      330                      335

Lys Tyr Trp Pro Gln Leu Ser Thr Asn Gly Thr Ala His Gly His Pro
                      340                      345                      350

His Glu Ile Ile Leu Tyr Tyr Tyr Glu Leu Tyr Pro Thr Met Thr Val
                      355                      360                      365

Val Val Val Ser Val Ala Thr Phe Ile Leu Leu Ser Met Val Gly Met
370                      375                      380

Ala Ala Gly Met Cys Met Cys Ala Arg Arg Arg Cys Ile Thr Pro Tyr
385                      390                      395                      400

Glu Leu Thr Pro Gly Ala Thr Val Pro Phe Leu Leu Ser Leu Ile Cys
                      405                      410                      415

Cys Ile Arg Thr Ala Lys Ala
                      420

```

```

<210> SEQ ID NO 4
<211> LENGTH: 423
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mutation arising from Vero passaging

```

```

<400> SEQUENCE: 4

```

```

Ser Thr Lys Asp Asn Phe Asn Val Tyr Lys Ala Thr Arg Pro Tyr Leu
1                      5                      10                      15

Ala His Cys Pro Asp Cys Gly Glu Gly His Ser Cys His Ser Pro Val
                      20                      25                      30

Ala Leu Glu Arg Ile Arg Asn Glu Ala Thr Asp Gly Thr Leu Lys Ile
35                      40                      45

Gln Val Ser Leu Gln Ile Arg Ile Lys Thr Asp Asp Ser His Asp Trp
50                      55                      60

Thr Lys Leu Arg Tyr Met Asp Asn His Met Pro Ala Asp Ala Glu Arg
65                      70                      75                      80

```


Ala	His	Cys	Pro	Asp	Cys	Gly	Glu	Gly	His	Ser	Cys	His	Ser	Pro	Val
			20				25						30		
Ala	Leu	Glu	Arg	Ile	Arg	Asn	Glu	Ala	Thr	Asp	Gly	Thr	Leu	Lys	Ile
			35				40						45		
Gln	Val	Ser	Leu	Gln	Ile	Gly	Ile	Lys	Thr	Asp	Asp	Ser	His	Asp	Trp
			50	55						60					
Thr	Lys	Leu	Arg	Tyr	Met	Asp	Asn	His	Met	Pro	Ala	Asp	Ala	Glu	Arg
			65	70						75			80		
Ala	Gly	Leu	Phe	Val	Arg	Thr	Ser	Ala	Pro	Cys	Thr	Ile	Thr	Gly	Thr
			85						90			95			
Met	Gly	His	Phe	Ile	Leu	Ala	Arg	Cys	Pro	Lys	Gly	Glu	Thr	Leu	Thr
			100			105						110			
Val	Gly	Phe	Thr	Asp	Ser	Arg	Lys	Ile	Ser	His	Ser	Cys	Thr	His	Pro
			115			120						125			
Phe	His	His	Asp	Pro	Pro	Val	Ile	Gly	Arg	Glu	Lys	Phe	His	Ser	Arg
			130			135						140			
Pro	Gln	His	Gly	Lys	Glu	Leu	Pro	Cys	Ser	Thr	Tyr	Val	Gln	Ser	Thr
			145			150						155			
Ala	Ala	Thr	Thr	Glu	Glu	Ile	Glu	Val	His	Met	Pro	Pro	Asp	Thr	Pro
			165						170			175			
Asp	His	Thr	Leu	Met	Ser	Gln	Gln	Ser	Gly	Asn	Val	Lys	Ile	Thr	Val
			180			185						190			
Asn	Gly	Gln	Thr	Val	Arg	Tyr	Lys	Cys	Asn	Cys	Gly	Gly	Ser	Asn	Glu
			195			200						205			
Gly	Leu	Thr	Thr	Thr	Asp	Lys	Val	Ile	Asn	Asn	Cys	Lys	Val	Asp	Gln
			210			215						220			
Cys	His	Ala	Ala	Val	Thr	Asn	His	Lys	Lys	Trp	Gln	Tyr	Asn	Ser	Pro
			225			230						235			
Leu	Val	Pro	Arg	Asn	Ala	Lys	Leu	Gly	Asp	Arg	Lys	Gly	Lys	Ile	His
			245						250			255			
Ile	Pro	Phe	Pro	Leu	Ala	Asn	Val	Thr	Cys	Arg	Val	Pro	Lys	Ala	Arg
			260			265						270			
Asn	Pro	Thr	Val	Thr	Tyr	Gly	Lys	Asn	Gln	Val	Ile	Met	Leu	Leu	Tyr
			275			280						285			
Pro	Asp	His	Pro	Thr	Leu	Leu	Ser	Tyr	Arg	Asn	Met	Gly	Glu	Glu	Pro
			290			295						300			
Asn	Tyr	Gln	Glu	Glu	Trp	Val	Met	His	Lys	Lys	Glu	Val	Val	Leu	Thr
			305			310						315			
Val	Pro	Thr	Glu	Gly	Leu	Glu	Val	Thr	Trp	Gly	Asn	Asn	Glu	Pro	Tyr
			325			330						335			
Lys	Tyr	Trp	Pro	Gln	Leu	Ser	Thr	Asn	Gly	Thr	Ala	His	Gly	His	Pro
			340			345						350			
His	Glu	Ile	Ile	Leu	Tyr	Tyr	Tyr	Glu	Leu	Tyr	Pro	Thr	Met	Thr	Val
			355			360						365			
Val	Val	Val	Ser	Val	Ala	Thr	Phe	Ile	Leu	Leu	Ser	Met	Val	Gly	Met
			370			375						380			
Ala	Ala	Gly	Met	Cys	Met	Cys	Ala	Arg	Arg	Arg	Cys	Ile	Thr	Pro	Tyr
			385			390						395			
Glu	Leu	Thr	Pro	Gly	Ala	Thr	Val	Pro	Phe	Leu	Leu	Ser	Leu	Ile	Cys
			405			410						415			
Cys	Ile	Arg	Thr	Ala	Lys	Ala									
			420												

-continued

```

<210> SEQ ID NO 6
<211> LENGTH: 423
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mutation arising from Vero passaging

<400> SEQUENCE: 6

Ser Thr Lys Asp Asn Phe Asn Val Tyr Lys Ala Thr Arg Pro Tyr Leu
1          5          10          15

Ala His Cys Pro Asp Cys Gly Glu Gly His Ser Cys His Ser Pro Val
20          25          30

Ala Leu Glu Arg Ile Arg Asn Glu Ala Thr Asp Gly Thr Leu Lys Ile
35          40          45

Gln Val Ser Leu Gln Ile Gly Ile Lys Thr Asp Asp Ser His Asp Trp
50          55          60

Thr Lys Leu Arg Tyr Met Asp Asn His Met Pro Ala Asp Ala Glu Arg
65          70          75          80

Ala Arg Leu Phe Val Arg Thr Ser Ala Pro Cys Thr Ile Thr Gly Thr
85          90          95

Met Gly His Phe Ile Leu Ala Arg Cys Pro Lys Gly Glu Thr Leu Thr
100         105         110

Val Gly Phe Thr Asp Ser Arg Lys Ile Ser His Ser Cys Thr His Pro
115         120         125

Phe His His Asp Pro Pro Val Ile Gly Arg Glu Lys Phe His Ser Arg
130         135         140

Pro Gln His Gly Lys Glu Leu Pro Cys Ser Thr Tyr Val Gln Ser Thr
145         150         155         160

Ala Ala Thr Thr Glu Glu Ile Glu Val His Met Pro Pro Asp Thr Pro
165         170         175

Asp His Thr Leu Met Ser Gln Gln Ser Gly Asn Val Lys Ile Thr Val
180         185         190

Asn Gly Gln Thr Val Arg Tyr Lys Cys Asn Cys Gly Gly Ser Asn Glu
195         200         205

Gly Leu Thr Thr Thr Asp Lys Val Ile Asn Asn Cys Lys Val Asp Gln
210         215         220

Cys His Ala Ala Val Thr Asn His Lys Lys Trp Gln Tyr Asn Ser Pro
225         230         235         240

Leu Val Pro Arg Asn Ala Glu Leu Gly Asp Arg Lys Gly Lys Ile His
245         250         255

Ile Pro Phe Pro Leu Ala Asn Val Thr Cys Arg Val Pro Lys Ala Arg
260         265         270

Asn Pro Thr Val Thr Tyr Gly Lys Asn Gln Val Ile Met Leu Leu Tyr
275         280         285

Pro Asp His Pro Thr Leu Leu Ser Tyr Arg Asn Met Gly Glu Glu Pro
290         295         300

Asn Tyr Gln Glu Glu Trp Val Met His Lys Lys Glu Val Val Leu Thr
305         310         315         320

Val Pro Thr Glu Gly Leu Glu Val Thr Trp Gly Asn Asn Glu Pro Tyr
325         330         335

Lys Tyr Trp Pro Gln Leu Ser Thr Asn Gly Thr Ala His Gly His Pro
340         345         350

His Glu Ile Ile Leu Tyr Tyr Tyr Glu Leu Tyr Pro Thr Met Thr Val
355         360         365

Val Val Val Ser Val Ala Thr Phe Ile Leu Leu Ser Met Val Gly Met

```

-continued

370	375	380	
Ala Ala Gly Met Cys Met Cys Ala Arg Arg Arg Cys Ile Thr Pro Tyr			
385	390	395	400
Glu Leu Thr Pro Gly Ala Thr Val Pro Phe Leu Leu Ser Leu Ile Cys			
	405	410	415
Cys Ile Arg Thr Ala Lys Ala			
	420		
<210> SEQ ID NO 7			
<211> LENGTH: 423			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: mutation arising from Vero passaging			
<400> SEQUENCE: 7			
Ser Thr Lys Asp Asn Phe Asn Val Tyr Lys Ala Thr Arg Pro Tyr Leu			
1	5	10	15
Ala His Cys Pro Asp Cys Gly Glu Gly His Ser Cys His Ser Pro Val			
	20	25	30
Ala Leu Glu Arg Ile Arg Asn Glu Ala Thr Asp Gly Thr Leu Lys Ile			
	35	40	45
Gln Val Ser Leu Gln Ile Gly Ile Lys Thr Asp Asp Ser His Asp Trp			
	50	55	60
Thr Lys Leu Arg Tyr Met Asp Asn His Met Pro Ala Asp Ala Glu Arg			
	65	70	75
Ala Gly Leu Phe Val Arg Thr Ser Ala Pro Cys Thr Ile Thr Gly Thr			
	85	90	95
Met Gly His Phe Ile Leu Ala Arg Cys Pro Lys Gly Glu Thr Leu Thr			
	100	105	110
Val Gly Phe Thr Asp Ser Arg Lys Ile Ser His Ser Cys Thr His Pro			
	115	120	125
Phe His His Asp Pro Pro Val Ile Gly Arg Glu Lys Phe His Ser Arg			
	130	135	140
Pro Gln His Gly Lys Glu Leu Pro Cys Ser Thr Tyr Val Gln Ser Thr			
	145	150	155
Ala Ala Thr Thr Glu Glu Ile Glu Val His Met Pro Pro Asp Thr Pro			
	165	170	175
Asp His Thr Leu Met Ser Gln Gln Ser Gly Asn Val Lys Ile Thr Val			
	180	185	190
Asn Gly Gln Thr Val Arg Tyr Lys Cys Asn Cys Gly Gly Ser Asn Glu			
	195	200	205
Gly Leu Thr Thr Thr Asp Lys Val Ile Asn Asn Cys Lys Val Asp Gln			
	210	215	220
Cys His Ala Ala Val Thr Asn Tyr Lys Lys Trp Gln Tyr Asn Ser Pro			
	225	230	235
Leu Val Pro Arg Asn Ala Glu Leu Gly Asp Arg Lys Gly Lys Ile His			
	245	250	255
Ile Pro Phe Pro Leu Ala Asn Val Thr Cys Arg Val Pro Lys Ala Arg			
	260	265	270
Asn Pro Thr Val Thr Tyr Gly Lys Asn Gln Val Ile Met Leu Leu Tyr			
	275	280	285
Pro Asp His Pro Thr Leu Leu Ser Tyr Arg Asn Met Gly Glu Glu Pro			
	290	295	300
Asn Tyr Gln Glu Glu Trp Val Met His Lys Lys Glu Val Val Leu Thr			

-continued

305	310	315	320
Val Pro Thr Glu Gly Leu Glu Val Thr Trp Gly Asn Asn Glu Pro Tyr			
	325	330	335
Lys Tyr Trp Pro Gln Leu Ser Thr Asn Gly Thr Ala His Gly His Pro			
	340	345	350
His Glu Ile Ile Leu Tyr Tyr Tyr Glu Leu Tyr Pro Thr Met Thr Val			
	355	360	365
Val Val Val Ser Val Ala Thr Phe Ile Leu Leu Ser Met Val Gly Met			
	370	375	380
Ala Ala Gly Met Cys Met Cys Ala Arg Arg Arg Cys Ile Thr Pro Tyr			
385	390	395	400
Glu Leu Thr Pro Gly Ala Thr Val Pro Phe Leu Leu Ser Leu Ile Cys			
	405	410	415
Cys Ile Arg Thr Ala Lys Ala			
	420		

<210> SEQ ID NO 8
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 8

ttaggatccg atggctgcgt gagacac

27

<210> SEQ ID NO 9
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 9

taactcgagc cgtaggtct gttgaacat

29

<210> SEQ ID NO 10
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 10

ttaggatcct accaccaggc gattaaag

28

<210> SEQ ID NO 11
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 11

taactcgagc ttgcccact tactgaagg

29

<210> SEQ ID NO 12
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

-continued

<400> SEQUENCE: 12

ttaggatcct gctacagaag tcacgcc

27

<210> SEQ ID NO 13

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 13

taactcgagg ccaaagcgtg aatcag

26

<210> SEQ ID NO 14

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 14

ttaggatcca acagcttgag gacagagcg

29

<210> SEQ ID NO 15

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 15

taactcgagc tctgtctcat cacgtcgg

28

<210> SEQ ID NO 16

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 16

ttaggatcca aattgcagtc ataggagtct tc

32

<210> SEQ ID NO 17

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 17

taactcgaga gtacgttgac gtgctctga

29

<210> SEQ ID NO 18

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 18

ttaggatccg tgggttaaac aactgcaaa

29

<210> SEQ ID NO 19

<211> LENGTH: 33

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 19

taactcgagg gttaaagtct tctatcctcc tgg

33

<210> SEQ ID NO 20
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 20

ttaggatccg gataaccact gggataatag g

31

<210> SEQ ID NO 21
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 21

taactcgaga gttgtgaaat tccttctgcc

30

<210> SEQ ID NO 22
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 22

ttaggatccc gcagatagaa ccagtgaac

29

<210> SEQ ID NO 23
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 23

taactcgagc agcagctcta cttgggtc

28

<210> SEQ ID NO 24
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 24

ttaggatcca ggagggaag acaggct

27

<210> SEQ ID NO 25
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 25

-continued

taactcgagc cctcgcttc ttctg 25

<210> SEQ ID NO 26
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 26

ttaggatccc aaaatagaag gaggcaaaa ag 32

<210> SEQ ID NO 27
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 27

taactcgagc ctggagtttc ttaagtaata gttgc 35

<210> SEQ ID NO 28
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 28

ttaggatcca ccggtccagg tcattta 27

<210> SEQ ID NO 29
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 29

taactcgagg cagcaaattc ttcccag 27

<210> SEQ ID NO 30
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 30

ttaggatccc cattccagaa cacactacag 30

<210> SEQ ID NO 31
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 31

taactcgaga tacctgattt catcatggc 29

<210> SEQ ID NO 32
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 32

ttaggatccc ctttgataag agccaagatg 30

<210> SEQ ID NO 33
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 33

taactcgagt acaaagttat gacgggtcct 30

<210> SEQ ID NO 34
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 34

ttaggatccc aacgaacagg gctaattg 28

<210> SEQ ID NO 35
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 35

taactcgagg accgcttaaa ggccag 26

<210> SEQ ID NO 36
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 36

ttaggatccg tgcataaaaa tcgaaaatg 29

<210> SEQ ID NO 37
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 37

taactcgagt ggtcttgtgg ctttatagac a 31

<210> SEQ ID NO 38
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 38

ttaggatcca accggaggaa accctac 27

-continued

<210> SEQ ID NO 39
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 39

taactcgagg taccgcaccg tctgg

25

<210> SEQ ID NO 40
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 40

ttaggatcca gctaccttgc agcacgt

27

<210> SEQ ID NO 41
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 41

taactcgagc ccaccatcga cagg

24

<210> SEQ ID NO 42
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 42

ttaggatccc gagccgtata agtattggc

29

<210> SEQ ID NO 43
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 43

taactcgagc gccggtgaag accttac

27

<210> SEQ ID NO 44
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 44

ttaggatcca ctactgtcag tcactttgga gc

32

<210> SEQ ID NO 45
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

-continued

<400> SEQUENCE: 45

taactcgagt accgggtttg ttgctattt

29

<210> SEQ ID NO 46

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 46

ttaggatccc acaactggta ctgcagagac

30

<210> SEQ ID NO 47

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 47

taactcgagg cgtagccctt tgatctatag

30

<210> SEQ ID NO 48

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 48

ttaggatccg gtgctatgcg tgcctg

26

<210> SEQ ID NO 49

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer

<400> SEQUENCE: 49

taactcgaga tctcctacgt cctctgtggg

29

The invention claimed is:

1. A pharmaceutical composition comprising:

- (i) CHIKV-Δ5nsP3 particles expressing an E2 structural protein as defined by the amino acid sequence of SEQ ID NO: 2;
- (ii) CHIKV-Δ5nsP3 particles expressing an E2 structural protein having an E168K and/or G55R mutation in the amino acid sequence of SEQ ID NO: 2; and
- (iii) optionally a pharmaceutically acceptable excipient, wherein 1-50% of the CHIKV-Δ5nsP3 particles present in the composition express an E2 structural protein having said E168K and/or G55R mutation.

2. The pharmaceutical composition according to claim 1, wherein said composition induces neutralizing antibodies against CHIKV-Δ5nsP3 in a mouse immunized with said pharmaceutical composition resulting in a serum comprising said neutralizing antibodies, and wherein said serum neutralizes Chikungunya virus (CHIKV) infection of Vero cells at least 50% in an in vitro neutralization assay at a 1:80 serum dilution.

3. The pharmaceutical composition according to claim 1, wherein the composition induces neutralizing antibodies

against CHIKV-Δ5nsP3 in a mouse immunized with said pharmaceutical composition resulting in a serum comprising said neutralizing antibodies, and wherein said serum neutralizes Chikungunya virus (CHIKV) infection of Vero cells at least 80% in an in vitro neutralization assay at a 1:80 serum dilution.

4. The pharmaceutical composition according to claim 1, wherein the composition induces neutralizing antibodies against CHIKV-Δ5nsP3 in a mouse immunized with said pharmaceutical composition resulting in a serum comprising said neutralizing antibodies, and wherein said serum neutralizes Chikungunya virus (CHIKV) infection of Vero cells at least 90% in an in vitro neutralization assay at a 1:80 serum dilution.

5. The pharmaceutical composition according to claim 1, wherein the composition induces neutralizing antibodies against CHIKV-Δ5nsP3 in a mouse immunized with said pharmaceutical composition resulting in a serum comprising said neutralizing antibodies, and wherein said serum neutralizes Chikungunya virus (CHIKV) infection of Vero cells at least 95% in an in vitro neutralization assay at a 1:80 serum dilution.

83

6. The pharmaceutical composition according to claim 1, wherein said CHIKV-Δ5nsP3 particles in (i) are defined by the polynucleotide sequence of SEQ ID NO: 1.

7. The pharmaceutical composition according to claim 1, wherein said pharmaceutical composition comprises at least 10^3 , at least 10^4 CHIKV-Δ5nsP3 particles that express an E2 structural protein as defined by the amino acid sequence of SEQ ID NO: 2.

8. The pharmaceutical composition according to claim 1, wherein said pharmaceutical composition comprises about 10^4 CHIKV-Δ5nsP3 particles that express an E2 structural protein as defined by the amino acid sequence of SEQ ID NO: 2.

9. The pharmaceutical composition according to claim 1, wherein said pharmaceutical composition is a one-shot pharmaceutical composition.

10. The pharmaceutical composition according to claim 1, wherein said pharmaceutical composition is a vaccine.

11. The pharmaceutical composition according to claim 10, wherein said pharmaceutical composition is provided in a lyophilized form.

12. A process for producing the pharmaceutical composition of claim 1 comprising the step of:

84

growing CHIKV-Δ5nsP3 particles on host cells in such a way as to minimize the presence of immunogenicity-reducing mutations of the virus.

13. The process according to claim 12, wherein said CHIKV-Δ5nsP3 particles are defined by a polynucleotide sequence of SEQ ID NO: 1 and are passaged on host cells in culture less than five times.

14. The process according to claim 13, wherein said CHIKV-Δ5nsP3 particles are defined by a polynucleotide sequence of SEQ ID NO: 1 and are passaged on host cells in culture at most three times.

15. A method of protection against disease caused by Chikungunya virus in a subject in need thereof, comprising administering to the subject an effective amount of the pharmaceutical composition according to claim 1.

16. The method according to claim 15, wherein said pharmaceutical composition comprises an effective amount of CHIKV-Δ5nsP3 particles which express an E2 structural protein as defined by the amino acid sequence of SEQ ID NO: 2, wherein said effective amount is defined as an amount sufficient to prevent Chikungunya virus viremia in a vaccinated subject.

17. The method according to claim 15, wherein the subject is a human subject.

* * * * *