



(51) International Patent Classification:

C12N 1/16 (2006.01) A01N 63/32 (2020.01)

(21) International Application Number:

PCT/IB2022/053387

(22) International Filing Date:

11 April 2022 (11.04.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

102021000009128 12 April 2021 (12.04.2021) IT

(71) Applicant: AGROVENTURES S.R.L. [IT/IT]; Piazza B. Buozzi, 9, 04100 Latina (LT) (IT).

(72) Inventors: CASTORIA, Raffaello; Via della Canzone del Piave, 24, 00143 Roma (RM) (IT). DE CICCIO, Vincenzo; Via Suppa, 32, 70121 Bari (BA) (IT). DE CURTIS, Filippo; C/da Cappuccini, 1, 86035 Larino (CB) (IT). FOLCHI, Bruno; 712 Main St. North, Southbury, Connecticut 06488 (US). IANIRI, Giuseppe; Via San Giovanni, 79, 86100 Campobasso (CB) (IT). LIMA, Giuseppe; C/da Vazzieri Poggio Verde, 1, 86010 Ferrazzano (CB) (IT). PALMIERI, Davide; C/da Loie, 469, 86016 Riccia (CB) (IT).

(74) Agent: MARTURANO, Pasqualino et al.; c/o Barzano' & Zanardo Milano S.p.A., Via Borgonuovo, 10, 20121 Milan (IT).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))
- with sequence listing part of description (Rule 5.2(a))
- in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(54) Title: NEW YEAST STRAIN AND ITS USES FOR THE CONTROL OF PHYTOPATHOGENS




Fig. 1

(57) Abstract: The present invention relates to a new yeast strain for the control of the main phytopathogens, which is effective in the prevention, suppression, treatment or control of a wide range of phytopathogens and related diseases affecting the whole plant (both the aerial part and the radical part).



NEW YEAST STRAIN AND ITS USES FOR THE CONTROL OF PHYTOPATHOGENS

5 The present invention regards a new yeast strain and its uses for
the control of phytopathogens. Importantly, the present invention concerns
a new yeast strain for the control of the main phytopathogens, which is
effective in the prevention, treatment and control of a wide range of
phytopathogens and the related diseases affecting the whole plant (both
the aerial and root parts).

10 It is well known that crop plants can be affected by various fungal,
bacterial and viral diseases, as well as by numerous harmful insects that
can also be vectors of phytopathogens.

 Currently, plant protection is mainly based on the use of chemical
pesticides. However, concerns about the emergence of resistance to
15 chemical active ingredients in pathogen populations and the impact of
these chemicals on human and animal health and on the environment,
require the development of alternative plant protection approaches, such
as biological and integrated control of plant diseases. Integrated pest
management (IPM) consists of the combined use of both chemical and
20 biological plant protection products along with agronomic measures useful
in controlling plant pathogens, i.e. that are not conducive to pathogens
attack. Biological management (i.e. biological control or biocontrol), on the
other hand, mainly relies on the use of products formulated with microbial
antagonists, plant-derived compounds or semiochemicals, which generally
25 are naturally occurring and pose low risks to human health and the
environment. In the last decades, the use of microorganisms for plant
protection has resulted in a number of well-established commercial
products, mainly based on soil-borne bacteria and fungi, which mainly act
via mycoparasitism and the production of natural compounds, such as
30 antibiotics or secondary metabolites with antimicrobial activity. The use of
microbial antagonistic agents aims to increase the range of biocontrol
formulates with low or no environmental impact, which can replace or
integrate chemical active ingredients for which risk mitigation measures

are being enforced for reducing their use or even withdrawing these chemicals from the market.

Some ascomycetous and basidiomycetous yeast isolates are of particular interest for various agricultural applications, including as
5 biocontrol agents (BCAs) to counteract plant pathogens. Epiphytic fungal microflora abundantly populates the aerial parts of plant organs, being adapted to a wide range of environmental stresses. These natural features make yeasts and yeast-like fungi excellent candidates for the development of plant protection products for the control of plant pathogens, both in the
10 field and during storage. Several yeast species with antagonistic properties against different pathogens have been selected and characterized for commercial use as BCAs e.g. *Candida oleophila* (Aspire®, Ecogen, Langhorne, PA, USA; Nexy®, Lesaffre-Bionext, France), *Cryptococcus albidus* (YieldPlus®, Lallemand, Montreal,
15 Canada), *Candida sake* (Candifruit®, IRTA, Lleida, Spain), *Metschnikowia fructicola* (Shemer®, Bayer, Leverkusen, Germany) and *Aureobasidium pullulans* (Boniprotect®, Biofa, Musingen, Germany), the latter an yeast-like fungus.

For the commercial registration of BCA-based products, biological,
20 ecological and toxicological studies are required to demonstrate that the selected microorganisms are effective against plant pathogens and harmless to humans, animals and the environment.

However, the range of efficacy for many biocontrol yeasts is too narrow both in terms of target pathogens and in terms of cultures. In fact,
25 most of microbial-based formulates have extremely reduced scope of application, thus leading to the necessity of an integrated use with other active ingredients. Furthermore, the use of combination of biocontrol microorganisms with synthetic chemical active ingredients has a major constraint in the limited ability of BCAs to tolerate chemical active
30 ingredients. Therefore, there are numerous technical restrictions for the

combinations of most microbial species used as BCAs with synthetic chemicals, which further limits the use of these products.

In the light of this situation, it appears evident the need for using new BCAs that can solve the above-mentioned issues and represent alternatives to the ones presently marketed.

In this context, the present invention represents a possible solution, since it consists of the formulate of a new microorganism to be used as a biocontrol agent of the main plant pathogens.

According to the present invention, it has now been demonstrated that the yeast species *Papiliotrema terrestris* and its extracellularly secreted substances can be advantageously used against plant pathogens.

In particular, the Applicant has now isolated a new yeast strain (PT22AV), belonging to the basidiomycete yeast species *Papiliotrema terrestris*. The aforementioned yeast strain was filed for patent purposes in according to the procedures laid down in the Budapest Treaty, on 26 October 2020, at the Westerdijk Fungal Biodiversity Institute (CBS), depositor AgroVentures srl, with access number provided by the International patent filing Authority CBS147138 and with identification reference provided by the depositor of PT22AV.

In line with the invention, Table 1 shows the taxonomic classification of the PT22AV strain of *Papiliotrema terrestris*.

Table 1

Taxonomical classification of PT22AV

Scientific classification	
Kingdom	Fungi
Phylum	Basidiomycota
Subphylum	Agaricomycotina
Class	Tremellomycetes
Subclass	Tremellomycetidae

Order	Tremellales
Family	Rhynchogastremataceae
Genus	<i>Papiliotrema</i>
Species	<i>Papiliotrema terrestris</i>
Synonyms	
<i>Cryptococcus terrestris</i>	
Strain	
<i>Papiliotrema terrestris</i> PT22AV	

The genus *Papiliotrema* (Class Tremellomycetes, Order Tremellales, Family Rhynchogastremataceae) was first described in 2002 with the aim of classifying a new microbial species, *Papiliotrema bandonii*.

5 Recently, the taxonomic reclassification, involving many species of basidiomycetous yeasts, has transferred numerous species from genus *Cryptococcus* to genus *Papiliotrema*. At present, this new fungal genus includes about 24 species and among these many are described as biological control agents, such as *P. laurentii* (formerly *C. laurentii*).
 10 *Papiliotrema terrestris* is a relatively new species, in fact it was first described in 2015.

In particular, as shown in the examples reported below, the PT22AV strain of *Papiliotrema terrestris*, isolated according to the present invention, has shown better or comparable performances as a biocontrol
 15 agent as compared to already known chemical and biological products. In particular, fresh, dried or dehydrated PT22AV cells, applied in biological or integrated protocols for seed coating, in seedbed, in the field or in the post-harvest stage, showed a reduction in the disease incidence that was comparable to that obtained with synthetic chemical products.

20 As shown below, the PT22AV strain of *Papiliotrema terrestris* according to the present invention was tested in efficacy and selectivity assays conducted for a period of 3 years (2018-2019-2020) in 4 EU countries on 6 crops against a wide range of biotrophic and necrotrophic

fungus pathogens, in the field and in the post-harvest. Different formulations based on strain PT22AV were tested, which were different from each other for composition and concentration of yeast cells, and were obtained through different approaches and technologies.

5 On the basis of the obtained results, strain PT22AV of *Papiliotrema terrestris* according to the present invention has proved to be a better biocontrol agent against plant pathogens as compared to:

- a) commercial microbial formulations based on bacteria (e.g. *Bacillus subtilis* and *B. amyloliquefaciens*), yeasts (e.g. *Aureobasidium pullulans*)
10 and fungi (e.g. *Trichoderma spp.*);
- b) numerous commercially established chemical plant protection products.

 As regards the comparison with formulations based on bacteria and yeasts (point a), *Papiliotrema terrestris* PT22AV, according to the present
15 invention, has proved to yield better performances since it can be used at much lower cell concentrations with the same efficacy. Overall, strain PT22AV according to the present invention, used alone or in combination with other agrochemicals (fungicides, adjuvants, fertilizers, biostimulants, etc.) was demonstrated to be much more effective than other commonly
20 used microbe-based formulations (e.g. *Bacillus spp.*) in the control of phytopathogenic fungi (including mycotoxigenic ones) and other plant pathogens, when applied both in the field and post-harvest, both in the aerial and in the root part. Furthermore, the strain according to the present invention, unlike chemical formulations indicated above (b), does not pose
25 any risk of selecting populations of resistant pathogens, is safer for humans and the environment, acts on a wider range of crops, diseases, phenological stages of plants, has a better adaptability to environmental conditions, can be better used in synergistic integration with various chemical or biological fungicides, adjuvants and other agrochemicals.

30 On the other hand, as regards the comparison with chemical products (point b), the strain according to the present invention has shown

comparable or even higher performances as compared to a wide range of chemical fungicides. Furthermore, according to the present invention, *P. terrestris* PT22AV does not produce any toxic chemical residues unlike chemical products. In addition, the strain can be applied both before and after harvest, does not allow the onset of resistance in pathogens' populations, is compatible with many agrochemicals when used in combination when used in combination with these products, and can be used both in organic farming and in integrated pest management.

According to the present invention, the efficacy of the treatment based on strain PT22AV has been clearly demonstrated for both fresh and dehydrated cells.

According to the present invention, formulates including the strain PT22AV are particularly advantageous since, if applied to crops, they actively protect the plant, promote its growth, stimulate systemic resistance against pathogens, reduce the production and accumulation of mycotoxins, mitigate the harmful effects of nematodes and other biotic and abiotic stressors. Furthermore, according to the present invention, strain PT22AV of *P. terrestris* is highly active in preventing, suppressing and controlling a wide range of diseases/pathogens affecting the whole plant (aerial parts and roots), with the possibility of application in the field, in post-harvest, in the soil and for seed coating. In particular, strain PT22AV according to the present invention can be advantageously used for root and seed treatment and for the entire aerial parts (including flowers and fruits) of plants, and can be applied alone or in synergistic combinations with other products for agriculture, in particular with pesticides (acaricides, bactericides, fungicides, insecticides, herbicides, algaecides, molluscicides, nematocides, rodenticides, plant growth regulators, repellents, semiochemicals, virucides), bio-pesticides, adjuvants and other agrochemicals (additives, fertilizers, biostimulants).

Therefore, according to the present invention, *P. terrestris* PT22AV, compared to other known biocontrol yeasts, can be advantageously used

against a wide range of pathogens in different types of crops and is tolerant towards synthetic chemical active principles, which makes it possible its effective use in combinations with these chemicals, as shown in the examples below.

5 In particular, the PT22AV strain of *P. terrestris*, according to the present invention, can be advantageously used as a biocontrol agent (BCA) for the management of plant pathogens both in organic agriculture and in integrated disease management (IDM).

10 Furthermore, according to the present invention, it has been found that strain PT22AV has a very active secondary metabolism, and secretes a series of secondary metabolites in the extracellular environment. Among these, extracellular polymers are of particular importance, a complex group of substances largely made up of polysaccharides. The composition and production of these polymers are very complex and it has been
15 observed that it is conditioned by several factors, including the composition of the growth medium, pH, temperature and oxygenation during biomass production. According to the present invention, it has been found that the extracellular polymers produced by *P. terrestris* strain PT22AV are mainly composed of carbohydrates (50-90%); the most
20 represented monomers are: mannose, xylose, glucose, galactose, arabinose. The structure of these polymers is very similar to starch and glycogen, it is indeed organized in a main chain consisting of monomeric units linked through α - (1→3) bonds and branches anchored through β - (1→2) bonds. These compounds are mainly produced under conditions of
25 trophic stress and in all conditions limiting cell growth. These polymers are active in the stimulation of plant metabolism, in inducing resistance against pathogens, to improve the chemical, physical and biological characteristics of the soil, to promote the healing of wounds in plants, and if they are distributed on the aerial parts they constitute a physical obstacle to
30 penetration of pathogens.

Therefore, according to the present invention, the aforementioned extracellular substances produced by the PT22AV yeast strain can be advantageously used against plant pathogens.

Scientific papers on molecular taxonomy have highlighted a high genetic homogeneity within the species *P. terrestris*; this indicates that in this species, the rate of genetic recombination linked to sexual reproduction is low, and that asexual reproduction is prevalent, which is a guarantee of genetic and phenotypic stability, and therefore of constant high performance in the field, in other application and also over time.

No technical and/or scientific publications have so far documented evidence of both biocontrol and biostimulant activities of *P. terrestris* species in the pre-harvest stage - in the ground or in the open field, in soil applications, on seeds and on aerial plants parts. Therefore, according to the present invention, the biostimulant activity of the yeast species *P. terrestris*, in particular of the strain PT22AV, in pre-harvest stage, is shown for the first time (see Example 5).

Moreover, according to the present invention, the biocontrol activity by the yeast species *P. terrestris*, in particular of strain PT22AV, is shown for the first time in pre-harvest stage. According to the international scientific literature, the only reported biocontrol activity by the species *P. terrestris* regards strain LS28 (previously classified as *Cryptococcus laurentii*), which was not applied in the open field but it was only applied on the harvested products (post-harvest) against post-harvest fungal pathogens. Nevertheless, biocontrol activity by PT22AV strain is significantly higher than the biocontrol activity by LS28, as compared by the Applicant. As shown in the comparative experimental data reported below (see Example 6), the efficacy of PT22AV is significantly higher than the efficacy of LS28. In addition, PT22AV showed a significantly greater ability to resist a wide range of abiotic stresses by PT22AV as compared to LS28.

The PT22AV strain, according to the present invention, is safe for humans and the environment, its manufacturing process has no environmental impact and satisfies green and circular economy principles, with manufacturing by-products to be used as soil organic conditioners and with possibility to use food industry by-products such as molasse for its fermentation process. The efficacy, selectivity, toxicological and ecotoxicological properties of the formulations based on the PT22AV strain have been extensively tested under GAP - GLP conditions (as required by current legislation regarding the European registration of microbial-based pesticides).

As far as human toxicology is concerned, *Papiliotrema terrestris* is a naturally occurring and non-pathogenic yeast species. In this regard, Ke et al. (2018) report that *Papiliotrema terrestris* was non-pathogenic in Sprague-Dawley rats after intravenous (IV) administration of 2.4×10^8 CFU and oral administration of 1.3×10^9 CFU. The genotoxic results for the galactosidase enzyme concentrate are negative, both in a bacterial reverse mutation test (Ames's test) and in a chromosome aberration test in cultured Chinese hamster lung fibroblast (CHL/IU) cells. In addition, in the 13-week gavage study in Sprague-Dawley rats, no adverse effects were observed in any of the tested groups and a No Observed Adverse Effects Level (NOAEL) of 2000 mg/kg bw/day [total organic solids (TOS) 1800 mg/kg bw/day] was established, which was the highest dose tested. Allergenicity sequence analysis revealed no evidence that galactosidase enzyme is an allergen. The data presented in this study support the conclusion that galactosidase produced by *P. terrestris* is safe for use in food production. The Applicant has carried out additional toxicological studies (data not shown), which confirm that strain PT22AV according to the present invention has a favorable toxicological profile based on values obtained from oral, pulmonary, intraperitoneal, dermal, eye irritation and other relevant toxicological tests.

As regards environmental toxicology, the Applicant has carried out various studies (data not shown), necessary for the commercial registration process in Europe and in the United States, which confirm that the formulations based on the PT22AV strain are safe for the environment.

5 Studies include evaluations of birds, mammals, non-target arthropods such as bees, predatory mites, soil organisms such as earthworms and others, aquatic organisms such as fish, daphnia and algae, as well as other non-target organisms, for which selectivity studies are required by regulatory bodies in Europe and USA.

10 According to the present invention, the *P. terrestris* strain PT22AV can be effectively used as a broad-spectrum fungicide (in terms both of crop plants and of plant pathogens) in organic farming, as well as for integrated pest management programs.

Furthermore, the ability to quickly colonize the plant surfaces,
15 natural openings and wounds advantageously enables the PT22AV strain to effectively counteract even the phytopathogenic bacteria that penetrate their hosts through these openings (e.g. *Pseudomonas* spp., *Xanthomonas* spp., *Clavibacter* spp., *Ralstonia* spp., *Erwinia* spp). In particular, the strain PT22AV produces a large quantity of polysaccharides
20 in extracellular environments during growth both *in vitro* and *in vivo*. This very complex matrix protects the strain from abiotic stressors, improves adhesion to surfaces and triggers plant systemic resistance, improving tolerance to a wide range of pathogens, including viruses.

In addition, the *P. terrestris* species, in particular the PT22AV strain,
25 according to the present invention, can be advantageously applied also to the soil, in particular for the improvement of the soil structure itself and for nematodes control. In fact, the application of the PT22AV strain to the soil produces a series of beneficial effects, including a positive action on the soil and rhizosphere microbiome. The polysaccharide matrix, which
30 facilitates the formation of aggregates and therefore improves the soil structure, is a source of nutrients for useful bacteria, microfauna and

predatory protists, and contributes to the ecological biogeochemical processes that are essential for the preservation of the soil ecosystem. Furthermore, formulations including the PT22AV strain according to the present invention are particularly advantageous since, if applied to crops, they actively protect the plant and promote its growth and resistance to biotic stressors, such as damage from nematodes. In particular, when applied as a root treatment, the PT22AV strain while partially reducing the proliferation of nematodes it is also especially effective in limiting their harmful effects, as shown by measuring plant height (see Example 7). Possible nematodes species are *Meloidogyne* spp., *Heterodera* spp., *Globodera* spp., *Belonolaimus* spp., *Pratylenchus* spp., *Rotylenchulus* spp., *Trichodorus* spp., *Paratylenchus* spp. etc.

The PT22AV strain is effective at a broad spectrum of concentrations by improving the overall health of plants.

The mechanisms underlying the antagonistic action of the PT22AV strain according to the present invention advantageously allow its application on a broad spectrum of host plants and pathogens (fungi, bacteria and viruses), and a high flexibility in its application both in terms of time (all the phenological phases) and in terms of the treated organs (seed, root, epigeous plant parts, flower, fruit). Furthermore, the ability to adapt to very heterogeneous environmental conditions (humidity, temperature, UV irradiation etc.) allows its application in different climatic areas.

Underlying the efficacy of the PT22AV strain in controlling plant diseases is the ability to survive and persist on plant surface or on wounded tissues for long periods. In particular, the tolerance to environmental stresses such as dehydration, oxidative stress, UV radiation, osmotic stress and tolerance to fungicides allow the adaptation of the PT22AV strain to the application environment.

According to the present invention, the plant protection against many fungal pathogens exerted by *P. terrestris* PT22AV is mainly

preventive. In particular, the application of the live cells of the strain prevents the establishment of the infection process in the early stages of plant-pathogen interaction (e.g. conidia germination, germ tube elongation, infection site colonization). Furthermore, bio-protection is the result of
5 several mechanisms with different relevance, which regulate the interactions both with pathogens and with other microorganisms that populate the same ecological niche.

The modes of action that characterize the biocontrol activity of the PT22AV yeast strain according to the present invention are listed below.

10 - Wound competence: for most necrotrophic fungi, wounded plant tissues represent the main penetration site to establish the infectious process. As already widely described in the literature, plant tissues respond to a wide range of stresses by triggering oxidative nature processes. Among these, a robust oxidative response is triggered by
15 wounded plant tissues, which brings along the accumulation of reactive oxygen species (mainly superoxide anion and hydrogen peroxide) and other oxidizing substances. Studies carried out by the Applicant demonstrate how the biological control agent PT22AV effectively colonizes injured tissues by resisting the oxidative stress exerted by reactive oxygen
20 species (ROS: superoxide anion O_2^- , and hydrogen peroxide, H_2O_2) generated by plant tissues as a consequence of wounding. The selected yeast strain, according to the invention, tolerates oxidative stressors through two enzymes (superoxide dismutase and catalase). Resistance to oxidative stress can be considered the first mechanism implemented by
25 PT22AV against post-harvest wound pathogens. In particular, the Applicant has demonstrated the PT22AV strain strong ability to resist oxidative phenomena by the assessment of the population dynamics in artificial wounds on apple, which showed how this microorganism is able to effectively colonize this environment. The strain population was
30 assessed by plating on nutrient agar medium yeast samples withdrawn from the wounded tissue inoculated with the PT22AV strain. From this

analysis it emerged that the PT22AV strain colonizes the wound very quickly, doubling in concentration within 6 hours from inoculation. Furthermore, strain resistance to oxidative stress was further investigated by the Applicant by evaluating the growth of the microorganism in the presence of hydrogen peroxide. For this purpose, increasing concentrations of hydrogen peroxide were added to the NYDA agar plates, subsequently inoculated with a suspension of PT22AV strain cells. From this analysis it emerged that this microorganism effectively withstands hydrogen peroxide concentrations up to 30 mM, therefore to a level of oxidative stress much higher than the one that is produced by wounding of a plant tissue.

- Competition for space and nutrients: PT22AV acts by efficiently competing for space and nutrients with fungal pathogens. *Papiliotrema terrestris* isolate effectively inhibit the development of the pathogen in its early growth stages by preventing the infection process. *In vivo* experiments, carried out by the Applicant, have highlighted how the addition of exogenous nutrients in artificial wounds drastically reduces the biocontrol efficacy of the yeast. Furthermore, in space competition, the PT22AV yeast strain is aided by the formation of an extracellular polysaccharide capsule which can promote adhesion to the fruit surface, thus preventing contact between host fruit and pathogen (Fig. 2). Therefore, competition for space and nutrients is a key mechanism of *P. terrestris* PT22AV in its antagonism against necrotrophic fungi and is closely related to oxidative stress resistance, hence to the ability to colonize plant tissues. Effective and timely space competition is a major trait of PT22AV making it effective against any disease and any plant culture (i.e. it confers this yeast a wide spectrum biocontrol activity). More specifically, efficacious competition of PT22AV for space (and consequently for nutrients) is a highly complex phenotypic trait relying on many genes, which enables this BCA to exert an action that is analogous to multi-site fungicides, but with a much wider range of mechanisms of

action and number of involved genes; in the analogy with these fungicides, in fact, competition makes it very unlikely, if not impossible in the case of PT22AV, the selection of pathogen populations that are resistant to this BCA.

5 - β -1,3-glucanase activity: *P. terrestris* strain PT22AV, in addition to competition for space and nutrients, can directly interact with fungal pathogens through the degradation of their cell wall mediated by the production of the extracellular β -1,3-glucanase enzyme. *In vitro* studies carried out by the Applicant have shown that *P. terrestris* PT22AV is able
10 to produce significantly higher levels of extracellular β -1,3-glucanase when cultured in the presence of hyphal cell walls of the pathogens *P. expansum* and *B. cinerea* as sole sources of carbon. This mechanism is to be considered second level in terms of relevance compared to the aforementioned mechanisms.

15 - No antibiosis: metabolomic studies carried out by the Applicant have shown that, unlike other microorganisms active against plant pathogens and registered as the active ingredient in microbial-based formulates (in particular bacteria) *P. terrestris* PT22AV does not produce molecules with antifungal activity. Under experimental conditions, the
20 antibiosis appears not to be involved in the antagonist activity exerted by strain PT22AV, thus ruling out the possibility that fungal pathogens develop resistance. This makes PT22AV a much safer and more sustainable BCA than bacteria-based formulates, the number of which is currently higher than other microbial species (fungi and yeasts).

25 - Plant resistance induction: plants have an innate "immune system" to recognize and respond to pathogens (fungi, bacteria and viruses); this plant "immune response" can be triggered by various beneficial microorganisms inducing resistance both locally and systemically. Biocontrol yeasts can elicit plant systemic resistance against a broad
30 range of pathogens and this action is implicit in their biocontrol activity. The induction of resistance can be triggered by microbial cells, parts of

them or even by the molecules secreted in the extracellular environment once recognized by plants. *Papiliotrema terrestris* PT22AV produces many extracellular substances that can trigger a systemic defense response in the host plant thus improving tolerance to pathogens. This extracellular
5 substance production activity is unmatched by any other commercially available biocontrol microorganism.

- Adhesion to the plant surface: during biomass production and after application in the field, the PT22AV strain produces an abundant matrix of polysaccharides that allows yeast cells to adhere to solid surfaces. This
10 extracellular matrix promotes the PT22AV adhesion to plants and wounded tissues, improves resistance to washout caused by rain and minimizes the negative effects of biotic and abiotic stressors. This activity is much higher than that of any other BCA commercially available or described in the literature.

15 It is therefore a specific object of the present invention a yeast strain belonging to *Papiliotrema terrestris* species, said strain being deposited at the Westerdijk Fungal Biodiversity Institute with deposit number CBS147138, hereinafter also referred to as PT22AV strain.

According to the present invention, said yeast strain can be in a form
20 selected from fresh cells, dried cells, dehydrated cells, devitalized cells, inactivated cells, frozen cells or cells in aqueous suspension, including derivatives (mainly extracellular polysaccharides) of the cell biomass production process.

The present invention also relates to a phytosanitary composition
25 for agricultural use, in particular for controlling phytopathogens in plants or crops, said composition comprising the PT22av yeast strain of the *Papiliotrema terrestris* species defined above, as the active principle, together with one or more excipients or phytopharmacologically acceptable adjuvants.

30 According to the present invention, said composition can be in liquid or solid form.

Furthermore, according to the present invention, the concentration of said yeast strain within the composition can range from 10^3 CFU to 10^{13} CFU per gram of solid composition or from 10^2 CFU to 10^{12} CFU per mL of liquid composition. Solid and liquid formulates at a concentration higher than 10^7 CFU per gram or mL are mainly suitable for foliar and post-harvest applications, while for formulates with a concentration lower than 10^7 CFU per gram or mL they can be used for soil treatments, enrichment of compost or other substrates for sowing, transplanting or growing plants and seed coating.

The concentration of the active ingredient in the formulates according to the invention is expressed as colony forming units (CFU) per gram or mL. The quantification can also be performed by other techniques such as real-time PCR, flow-cytometry, etc.

The PT22AV strain concentration in solutions for fertigation or in hydroponic cultivation etc can range between 10^2 and 10^{12} CFU per mL of solution. In distribution systems that involve the use of pollinating insects, the concentration of yeast in the trays dispenser can range between 10^2 and 10^{12} .

The composition according to the present invention can be in the form of tablets, capsules, granules, pellets, powder, such as dry powder or wettable powder, wettable granules, fluid, dry fluid, emulsion, suspension, solution, dispersion.

In particular, a composition according to the present invention can be in the form of: Aerosol (A); bait for pollinators dispersion (B); Dust (D); Dry flowable (DF); Emulsifiable (E); Emulsifiable concentrate (EC); Flowable (FL); Granule (G); Microencapsulated (M); Pellet (P); Soluble powder (SP); Wettable powder (WP); Water-dispersible granule (WG/WDG).

More specifically, composition can be in the form of: grain bait (AB); bait concentrate (CB); capsule suspension for seed treatment (CF); encapsulated granule (CG); capsule suspension (CS); dispersible

concentrate (DC); dispersible powder (DP); powder for dry seed treatment (DS); emulsifiable concentrate (EC); emulsion, water-in-oil (EO); emulsion for seed treatment (ES); emulsion, oil in water (EW); fine granule (FG); flowable concentrate for seed treatment (FS); granular bait (GB);
5 microgranule (GG); flo-dust (GP); Granule (GR); solution for seed treatment (LS); micro-emulsion (ME); microgranule (MG); oil dispersion (OD); oil miscible flowable concentrate (OF); oil miscible liquid (OL); oil dispersible powder (OP); paste (PA); seed coated with pesticide (PS); suspension concentrate (SC); suspo-emulsion (SE); water soluble
10 granules (SG); soluble concentrate (SL); water soluble powder (SP); water soluble powder for seed treatment (SS); ultra-low volume suspension (SU); tablet (TB); technical material (TC); technical concentrate (TK); water-dispersible granular (WG/WDG); wettable powder (WP); wettable dispersible powder for slurry seed treatment (WS).

15 According to an embodiment of the present invention, said composition can further comprise one or more chemical compounds, for example synthetic chemical compounds, and/or one or more biological agents, for example microorganisms such as bacteria, yeasts, fungi, viruses, algae, said one or more chemical compounds and said one or
20 more biological agents being selected from among fungicides, insecticides, fertilizers, biostimulants (e.g. humic and fulvic acids, protein hydrolysates, seaweed and plant extracts, chitosan and other biopolymers, inorganic compounds, beneficial microbes), macronutrients(e.g. nitrogen (N), phosphorous (P) and potassium (K)), micronutrients, elicitors, phyto-
25 regulators, microbial biocontrol agents, plant growth regulators, foliar nutrients, antibiotics, herbicides, acaricides, food additives(e.g. all compounds included in following categories: antioxidants, colors, preservatives and flour treatment agents), adjuvants (e.g. emulsifiers, wetting agents, dispersants, defoarmers, chelating agents, neutralizers,
30 solvents), microorganisms of agri-food interest.

According to the present invention, the excipients that can be used in the composition are for example co-formulants based on minerals (mineral powders) or organic (simple, complex or cyclic sugars), dispersants, emulsifiers, wetting agents, stabilizers, synergists, corroborants.

5 The aforementioned composition, further comprising one or more chemical compounds and/or one or more biological agents and/or one or more co-formulants, can be a tank-mix or a co-formulation.

In particular:

10 said foliar nutrients can be selected from nitrogen, phosphorus, potassium, calcium, sulfur, magnesium;

 said micronutrients can be selected from iron, boron, chlorine, manganese, zinc, copper, molybdenum and nickel;

15 said insecticides can be selected from inorganic compounds, oils, botanical extracts, organochlorines, organophosphates, carbamates, neonicotinoids, avermectins, organosulfur, organotins, synthetic pyrethroids;

20 said elicitors can be chosen from chemical synthesis elicitors, such as salicylic acid, methyl salicylate, benzothiadiazole, benzoic acid and chitosan, or from plant extracts elicitors, such as tannins, citrus oils, bioflavonoids, such as dihydroquercetin, polysaccharics, arabino-galactans, polysaccharides and other plant extracts;

25 said plant growth regulators can be chosen from (international nomenclature) auxins, gibberellins, cytokinins, ethylene, abscissic acid, ancymidol; chlormequat; chloro IPC; daminozide; flurprimidol; hydrogen cyhanamide, cyanamide (H_2CN_2); mefluidide; mepiquat chlorure; paclobutrozo; proexadione calcium; succinic acid (SADH), forchlorfenuron.

30 According to an embodiment of the present invention, said composition is a fungicidal composition. For example, the fungicidal action is carried out by the specific strain and/or by further fungicides added to the composition.

According to the present invention, the fungicides can belong for example to strobirulins, benzimidazoles, sterol inhibitors, dicarboxamides, dithiocarbamates, inorganic fungicides, phenylpyrroles or their combinations, and to other chemical groups.

5 In particular, said fungicides (chemical and/or biological) can be selected from among (international nomenclature) 2-methoxyethylmercury chloride, 2-phenylphenol, 3-ethoxypropyl mercury bromide, 8-hydroxyquinoline sulfate, 8-phenylmercurioxyquinoline, acibenzolar, acibenzolar-S-methyl, acylamino acid fungicides, acypetacs, Adavelt, 10 *Agrobacterium radiobacter* K84, aldimorph, allyl alcohol, ametoctradin, aminopyrifin, amisulbrom, ampropylfos, anilazine, aureofungin, azaconazole, azithiram, azoxystrobin, *Bacillus amyloliquefaciens* (formerly *subtilis*) strain QST713, *Bacillus amyloliquefaciens* strain AH2, strain IT-45, strain FZB24, strain MBI600, strain D747, *Bacillus mycoides* isolate J, 15 *Bacillus nakamurai* strain F727, *Bacillus pumilus* strain QST 2808, *Bacillus subtilis* var. *amyloliquefaciens* strain FZB24, *Bacillus subtilis* strain AFS032321, strain GB03, strain IAB/BS03, barium polysulfide, benalaxyl-M (=kiralaxyl), benodanil, benomyl, benquinox, bentluron, benthiavalicarb, Benthiavalicarb-isopropyl, benzalkonium chloride, 20 benzamacril, benzamide fungicides, benzamorf, benzohydroxamic acid, benzovindiflupyr, bethoxazin, binapacryl, biphenyl, bitertanol, bithionol, bixafen, blasticidin-S, Bordeaux mixture, boric acid, boscalid, bromuconazole, bupirimate, Burgundy mixture, buthiobate, calcium polysulfide, captafol, captan, carbamorph, carbendazim, carboxin, 25 carpropamid, carvone, cell walls of *Saccharomyces cerevisiae* strain LAS117, Cheshunt mixture, chinomethionat, chlobenthiazone, chloraniformethan, chloranil, chlorfenazole, chlorodinitronaphthalene, chloroneb, chloropicrin, Chlorothalonil, chlorquinox, chlozolate, ciclopirox, climbazole, *Clonostachys rosea* strain CR-7, *Coniothyrium minitans*, copper hydroxide, copper naphthenate, copper oleate, copper 30 oxychloride, copper soap, copper sulfate, basic, copper zinc chromate,

copper(II) acetate, copper(II) carbonate, basic, copper(II) sulfate, coumoxystrobin, cufraneb, cuprobam, cuprous oxide, cyazofamid, cyclafuramid, cyclobutrifluram, cycloheximide, cyflufenamid, cymoxanil, cypendazole, cyproconazole, cyprodinil, Cyprofuram, dazomet, DBCP, debacarb, decafentin, dehydroacetic acid, dichlobentiazox, dichlofluanid, dichlone, dichlorophen, dichlorophenyl, dichlozoline, diclobutrazol, diclocymet, diclomezine, dicloran, diethofencarb, diethyl pyrocarbonate, difenoconazole, diflumetorim, dimethachlone, dimethirimol, dimethomorph, dimoxystrobin, diniconazole, diniconazole-M, dinobuton, dinocap, dinocap-4, dinocap-6, dinoceton, dinopenton, dinosulfon, dinoterbon, diphenylamine, Dipymetitrone, dipyrithione, disulfiram, ditalimfos, dithianon, DNOC, dodemorph, dodicin, dodine, donatodine, drazoxolon, edifenphos, enestrobin, enestroburin, enoxastrobin, epoxiconazole, etaconazole, etem, ethaboxam, ethirimol, ethoxyquin, ethylene oxide, ethylmercury 2,3-dihydroxypropyl mercaptide, ethylmercury acetate, ethylmercury bromide, ethylmercury chloride, ethylmercury phosphate, etridiazole, extract from *Melaleuca alternifolia* (tea tree), extract from *Reynoutria sachalinensis* (giant knotweed), extract from the cotyledons of lupine plantlets ("BLAD"), extract of *Swinglea glutinosa*, F500, famoxadone, fenamidone, fenaminosulf, fenaminstrobin, fenamistrobin, fenapanil, fenarimol, fenazaquin, fenbuconazole, fenfuram, fenhexamid, fenitropan, fenoxanil, fempiclonil, fempicoxamid, fenpropidin, fenpropimorph, fenpyrazamine, fentin, fentin acetate, fentin chloride, fentin hydroxide, ferbam, ferimzone, Florylpicoxamid, fluazaindolizine, fluazinam, fludioxonil, flufenoxystrobin, fluindapyr, flumorph, fluopicolide, fluopimomide, fluopyram, fluoroimide, fluotrimazole, fluoxapiprolin, fluoxastrobin, fluquinconazole, flurozolamide, flusilazole, flusulfamide, Flutianil, flutolanil, flutriafol, fluxapyroxad, folpet, fosetyl, fosetyl-Al, fthalide (phthalide), fuberidazole, furalaxyl, furametpyr, furcarbanil, furfural, furmecyclox, furophanate, *Gliocladium catenulatum* J1446, *Gliocladium virens* GL-21, glyodin, griseofulvin, guazatine, halacrinat, hexachlorobenzene, hexachlorophene, hexaconazole,

hexylthiofos, hymexazol, imazalil, imibenconazole, iminoctadine, Inatreq
 (fenpicoxamid), inorganic oils, inpyrfluxam, iodocarb, ipconazole,
 Ipflufenquin, iprobenfos, iprodione, iprovalicarb, isofetamid, isoflucypram,
 isopropanol azole, isoprothiolane, isopyrazam, isotianil, isovaledione, Jun
 5 Si Qi, kasugamycin, kresoxim-methyl, laminarin, Lime sulfur (lime sulphur),
 mancopper, mancozeb, mandestrobin, mandipropamid, maneb, mebenil,
 mecarbinzid, mefenoxam, mefentrifluconazole, mepanipirim, mepronil,
 meptyldinocap, metalaxyl, metalaxyl-M (=mefenoxam), metam,
 metazoxolon, metconazole, methasulfocarb, methfuroxam, methyl
 10 bromide, methyl isothiocyanate, methylmercury benzoate, methylmercury
 dicyandiamide, methylmercury pentachlorophenoxide, metiram,
 metominostrobin, metrafenone, metsulfovax, metyltetraprole, milneb,
 myclobutanil, myclozolin, N-(ethylmercury)-p-toluenesulfonanilide, nabam,
 naftifine, natamycin, neem oil, nitrothal-isopropyl, nuarimol, octhilinone,
 15 ofurace, organic oils, orthophenyl phenol, orysastrobin, oxadixyl,
 oxathiapiprolin, oxazosulfyl, oxine copper, oxolinic acid, oxpoconazole,
 oxycarboxin, oxytetracycline, PCNB, pefurazoate, penconazole,
 pencycuron, penflufen, pentachlorophenol, penthiopyrad, phenamacril,
 phenylmercuriurea, phenylmercury acetate, phenylmercury chloride,
 20 phenylmercury nitrate, phosdiphen, Phosphite, phosphorous acid and
 salts, phosphorus acid, phthalide, picarbutrazox, picolinamides,
 picoxystrobin, piperalin, plant oils (mixtures): eugenol, geraniol, thymol,
 polycarbamate, polyoxin-D, potassium azide, potassium bicarbonate,
 potassium polysulfide, potassium thiocyanate, probenazole, prochloraz,
 25 procymidone, propamocarb, propiconazole, propineb, proquinazid,
 prothiocarb, prothioconazole, *Pseudomonas chlororaphis* strain AFS009,
Pseudomonas syringae ESC-10, pydiflumetofen, pyracarbolid,
 pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyrapropoyne,
 pyraziflumid, pyrazophos, pyribencarb, pyributicarb, pyridachlometyl,
 30 pyridinitril, pyrifenox, pyrimethanil, pyrimorph, pyrioifenone, pyrisoxazole,
 pyroquilon, pyroxychlor, pyroxyfur, quinacetol, quinazamid, quinconazole,

quinofumelin, quinomethionate, quinoxifen, quintozone, rabenzazole,
 Reynoutria sachalinensis, salicylanilide, sec-butylamine, sedaxane,
 sesame oil, silthiofam, silver, simeconazole, SJC17, sodium azide, sodium
 bicarbonate, sodium hypochlorite, sodium orthophenylphenoxide, sodium
 5 pentachlorophenoxide, sodium polysulfide, spiroxamine, *Streptomyces*
griseovirides strain K61, *Streptomyces lydicus* strain WYEC108,
 streptomycin, sulfur, sulfuryl fluoride, sultropen, tea tree oil, tebuconazole,
 tebufloquin, tecloftalam, tecnazene, tecoram, tetraconazole,
 thiabendazole, thiadifluor, thicyofen, thifluzamide, thiochlorfenphim,
 10 thiophanate, thiophanate-methyl, thioquinox, thiram, THQ25, tiadinil,
 tioxyimid, tolclofos-methyl, tolfenpyrad, tolprocarb, tolylfluanid, tolylmercury
 acetate, triadimefon, triadimenol, triamiphos, triarimol, triazbutil, triazoxide,
 trichlamide, *Trichoderma afroharzianum* (formerly *harzianum*) strain T-22,
Trichoderma asperellum (formerly *harzianum*) strain ICC012, strain T25,
 15 strain TV1, *Trichoderma asperellum* strain T34, *Trichoderma*
atrobrunneum (formerly *harzianum*) strain ITEM908, *Trichoderma*
atroviride (formerly *harzianum*) strain IMI206040, strain T11, *Trichoderma*
atroviride strain I-1237, *Trichoderma atroviride* strain LU132, *Trichoderma*
atroviride strain SC1, *Trichoderma gamsii* (formerly *viride*) strain ICC080,
 20 triclopyricarb, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine,
 triticonazole, UCQ09, *Ulocladium oudemansii* strain U3, uniconazole-P,
 urea, validamycin, valifenalate, vinclozolin, voriconazole, WLR08,
 zarilamid, zinc naphthenate, zinc thiazole, zineb, ziram, zoxamide,
 preferably azoxystrobin, trifloxystrobin, benomyl, carbendazim,
 25 thiabendazole, tebuconazole, imazalil, penconazole, procymidone,
 vinclozolin, mancozeb, ziram, copper oxychloride, sulfur, fludioxonil and/or
 iprovalicarb.

The phytosanitary composition according to the present invention
 can be prepared according to various processes known to a person skilled
 30 in the art, for example by a method chosen among spray dry, extrusion,
 fluidized-bed granulation with dehydration, freeze-drying, emulsion

process, suspension in mixture with one or more chemical compounds and/or biological agents and/or with one or more co-formulants, described before.

It is a further object of the present invention a kit comprising:

5 a) the PT22AV yeast strain of *Papiliotrema terrestris* species as defined above, or a composition as defined above, and

b) one or more chemical compounds, for example synthetic chemical compounds, and/or one or more biological agents, for example microorganisms such as bacteria, yeasts, fungi, viruses, algae, or
10 compositions of them. Both chemical compounds and biological agents can be selected among fungicides, insecticides, fertilizers, biostimulants, macronutrients, micronutrients, elicitors, phyto-regulators, microbial biological control agents, plant growth regulators, foliar nutrients, antibiotics, herbicides, acaricides, food additives, antioxidants, adjuvants,
15 microorganisms of agri-food interest.

In particular:

said foliar nutrients can be selected from nitrogen, phosphorus, potassium, calcium, sulfur, magnesium;
aforementioned micronutrients can be selected from iron, boron, chlorine,
20 manganese, zinc, copper, molybdenum and nickel;

said insecticides can be selected from inorganic compounds, oils, botanical extracts, organochlorines, organophosphates, carbamates, neonicotinoids, avermectins, organosulfurs, organotins, synthetic pyrethroids;

25 said elicitors can be chosen from chemical synthesis elicitors, such as salicylic acid, methyl salicylate, benzothiadiazole, benzoic acid and chitosan, or elicitors consisting of plant extracts, such as tannins, citrus oils, bioflavonoids, such as dihydroquercetin, arabinba galactane, polysaccharides and others plant extracts;

30 said plant growth regulators can be chosen from (international nomenclature): auxins, gibberellins, cytokinins, ethilene, abscissic acid,

ancymidol; chlormequat; chloro IPC; daminozide; flurprimidol; hydrogen cyanamide (H_2CN_2); mefluidide; mepiquat chlorure; paclobutrozo; proexadione calcium; succinic acid (SADH), forchlorfenuron.

As mentioned above, fungicides can belong, for example, to
 5 strobirulins, benzimidazoles, sterol inhibitors, dicarboxamides, dithiocarbamates, inorganic fungicides, phenylpyrroles or their combinations, and to other chemical groups.

According to the kit of the present invention, said fungicides (chemical and/or biological) can be selected from (international
 10 nomenclature) 2-methoxyethylmercury chloride, 2-phenylphenol, 3-ethoxypropyl mercury bromide, 8-hydroxyquinoline sulfate, 8-phenylmercurioxyquinoline, acibenzolar, acibenzolar-S-methyl, acylamino acid fungicides, acypetacs, Adavelt, *Agrobacterium radiobacter* K84, aldimorph, allyl alcohol, ametocetradin, aminopyrifin, amisulbrom,
 15 ampropylfos, anilazine, aureofungin, azaconazole, azithiram, azoxystrobin, *Bacillus amyloliquefaciens* (formerly *subtilis*) strain QST713, *Bacillus amyloliquefaciens* strain AH2, strain IT-45, strain FZB24, strain MBI600, strain D747, *Bacillus mycoides* isolate J, *Bacillus nakamurai* strain F727, *Bacillus pumilus* strain QST 2808, *Bacillus subtilis* var. *amyloliquefaciens*
 20 strain FZB24, *Bacillus subtilis* strain AFS032321, strain GB03, strain IAB/BS03, barium polysulfide, benalaxyl-M (=kiralaxyl), benodanil, benomyl, benquinox, bentazone, benthiavalicarb, Benthiavalicarb-isopropyl, benzalkonium chloride, benzamacril, benzamide fungicides, benzamorf, benzohydroxamic acid, benzovindiflupyr, bethoxazin,
 25 binapacryl, biphenyl, bitertanol, bithionol, bixafen, blastocidin-S, Bordeaux mixture, boric acid, boscalid, bromuconazole, bupirimate, Burgundy mixture, buthiobate, calcium polysulfide, captan, carbamorph, carbendazim, carboxin, carpropamid, carvone, cell walls of *Saccharomyces cerevisiae* strain LAS117, Cheshunt mixture,
 30 chinomethionat, chlobenthiazole, chloranilformetan, chloranil, chlorfenazole, chlorodinitronaphthalene, chloroneb, chloropicrin,

Chlorothalonil, chlorquinox, chlozolate, ciclopirox, climbazole, *Clonostachys rosea* strain CR-7, *Coniothyrium minitans*, copper hydroxide, copper naphthenate, copper oleate, copper oxychloride, copper soap, copper sulfate, basic, copper zinc chromate, copper(II) acetate, copper(II) carbonate, basic, copper(II) sulfate, coumoxystrobin, cufraneb, cuprobam, cuprous oxide, cyazofamid, cyclafuramid, cyclobutrifluram, cycloheximide, cyflufenamid, cymoxanil, cypendazole, cyproconazole, cyprodinil, Cyprofuram, dazomet, DBCP, debacarb, decafentin, dehydroacetic acid, dichlobentiazox, dichlofluanid, dichlone, dichlorophen, dichlorophenyl, dichlozoline, diclobutrazol, diclocymet, diclomezine, dicloran, diethofencarb, diethyl pyrocarbonate, difenoconazole, diflumetorim, dimethachlone, dimethirimol, dimethomorph, dimoxystrobin, diniconazole, diniconazole-M, dinobuton, dinocap, dinocap-4, dinocap-6, dinoceton, dinopenton, dinosulfon, dinoterbon, diphenylamine, Dipymetitrone, dipyrithione, disulfiram, ditalimfos, dithianon, DNOC, dodemorph, dodicin, dodine, donatodine, drazoxolon, edifenphos, enestrobin, enestroburin, enoxastrobin, epoxiconazole, etaconazole, etem, ethaboxam, ethirimol, ethoxyquin, ethylene oxide, ethylmercury 2,3-dihydroxypropyl mercaptide, ethylmercury acetate, ethylmercury bromide, ethylmercury chloride, ethylmercury phosphate, etridiazole, extract from *Melaleuca alternifolia* (tea tree), extract from *Reynoutria sachalinensis* (giant knotweed), extract from the cotyledons of lupine plantlets ("BLAD"), extract of *Swinglea glutinosa*, F500, famoxadone, fenamidone, fenaminosulf, fenaminstrobin, fenamistrobin, fenapanil, fenarimol, fenazaquin, fenbuconazole, fenfuram, fenhexamid, fenitropan, fenoxanil, fenciclonil, fencicoxamid, fenpropidin, fenpropimorph, fenpyrazamine, fentin, fentin acetate, fentin chloride, fentin hydroxide, ferbam, ferimzone, Florylpicoxamid, fluazaindolizine, fluazinam, fludioxonil, flufenoxystrobin, fluindapyr, flumorph, fluopicolide, fluopimomide, fluopyram, fluoroimide, fluotrimazole, fluoxapiprolin, fluoxastrobin, fluquinconazole, flurozalamide, flusilazole, flusulfamide, Flutianil, flutolanil, flutriafol, fluxapyroxad, folpet, fosetyl, fosetyl-Al, fthalide

(phthalide), fuberidazole, furalaxyl, furametpyr, furcarbanil, furfural, furmecyclox, furophanate, *Gliocladium catenulatum* J1446, *Gliocladium virens* GL-21, glyodin, griseofulvin, guazatine, halacrinat, hexachlorobenzene, hexachlorophene, hexaconazole, hexylthiofos, hymexazol, imazalil, imibenconazole, iminoctadine, Inatreq (fenpicoxamid), inorganic oils, inpyrfluxam, iodocarb, ipconazole, Ipflufenquin, iprobenfos, iprodione, iprovalicarb, isofetamid, isoflucypram, isopropanol azole, isoprothiolane, isopyrazam, isotianil, isovaledione, Jun Si Qi, kasugamycin, kresoxim-methyl, laminarin, Lime sulfur (lime sulphur), mancopper, mancozeb, mandestrobin, mandipropamid, maneb, mebenil, mecarbinzid, mefenoxam, mefentrifluconazole, mepanipyrim, mepronil, meptyldinocap, metalaxyl, metalaxyl-M (=mefenoxam), metam, metazoxolon, metconazole, methasulfocarb, methfuroxam, methyl bromide, methyl isothiocyanate, methylmercury benzoate, methylmercury dicyandiamide, methylmercury pentachlorophenoxide, metiram, metominostrobin, metrafenone, metsulfovax, metyltetraprole, milneb, myclobutanil, myclozolin, N-(ethylmercury)-p-toluenesulfonanilide, nabam, naftifine, natamycin, neem oil, nitrothal-isopropyl, nuarimol, octhilinone, ofurace, organic oils, orthophenyl phenol, orysastrobin, oxadixyl, oxathiapiprolin, oxazosulfyl, oxine copper, oxolinic acid, oxpoconazole, oxycarboxin, oxytetracycline, PCNB, pefurazoate, penconazole, pencycuron, penflufen, pentachlorophenol, penthiopyrad, phenamacril, phenylmercuriurea, phenylmercury acetate, phenylmercury chloride, phenylmercury nitrate, phosdiphen, Phosphite, phosphorous acid and salts, phosphorus acid, phthalide, picarbutrazox, picolinamides, picoxystrobin, piperalin, plant oils (mixtures): eugenol, geraniol, thymol, polycarbamate, polyoxin-D, potassium azide, potassium bicarbonate, potassium polysulfide, potassium thiocyanate, probenazole, prochloraz, procymidone, propamocarb, propiconazole, propineb, proquinazid, prothiocarb, prothioconazole, *Pseudomonas chlororaphis* strain AFS009, *Pseudomonas syringae* ESC-10, pydiflumetofen, pyracarbolid,

pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyrapropoyne,
 pyraziflumid, pyrazophos, pyribencarb, pyributicarb, pyridachlometyl,
 pyridinitril, pyrifenox, pyrimethanil, pyrimorph, pyrioifenone, pyrisoxazole,
 pyroquilon, pyroxychlor, pyroxyfur, quinacetol, quinazamid, quinconazole,
 5 quinofumelin, quinomethionate, quinoxifen, quintozone, rabenzazole,
 Reynoutria sachalinensis, salicylanilide, sec-butylamine, sedaxane,
 sesame oil, silthiofam, silver, simeconazole, SJC17, sodium azide, sodium
 bicarbonate, sodium hypochlorite, sodium orthophenylphenoxide, sodium
 pentachlorophenoxide, sodium polysulfide, spiroxamine, *Streptomyces*
 10 *griseovirides* strain K61, *Streptomyces lydicus* strain WYEC108,
 streptomycin, sulfur, sulfuryl fluoride, sultropen, tea tree oil, tebuconazole,
 tebufloquin, tecloftalam, tecnazene, tecoram, tetraconazole,
 thiabendazole, thiadifluor, thicyofen, thifluzamide, thiochlorfenphim,
 thiophanate, thiophanate-methyl, thioquinox, thiram, THQ25, tiadinil,
 15 tioxyimid, tolclofos-methyl, tolfenpyrad, tolprocarb, tolylfluanid, tolylmercury
 acetate, triadimefon, triadimenol, triamiphos, triarimol, triazbutyl, triazoxide,
 trichlamide, *Trichoderma afroharzianum* (formerly *harzianum*) strain T-22,
Trichoderma asperellum (formerly *harzianum*) strain ICC012, strain T25,
 strain TV1, *Trichoderma asperellum* strain T34, *Trichoderma*
 20 *atrobrunneum* (formerly *harzianum*) strain ITEM908, *Trichoderma*
atroviride (formerly *harzianum*) strain IMI206040, strain T11, *Trichoderma*
atroviride strain I-1237, *Trichoderma atroviride* strain LU132, *Trichoderma*
atroviride strain SC1, *Trichoderma gamsii* (formerly *viride*) strain ICC080,
 triclopyricarb, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine,
 25 triticonazole, UCQ09, *Ulocladium oudemansii* strain U3, uniconazole-P,
 urea, validamycin, valifenalate, vinclozolin, voriconazole, WLR08,
 zarilamid, zinc naphthenate, zinc thiazole, zineb, ziram, zoxamide,
 preferably azoxystrobin, trifloxystrobin, benomyl, carbendazim,
 thiabendazole, tebuconazole, imazalil, penconazole, procymidone,
 30 vinclozolin, mancozeb, ziram, copper oxychloride, sulfur, fludioxonil and/or
 iprovalicarb.

The present invention also concerns the use of the PT22AV yeast strain as defined above, of the extracellular substances secreted by said strain, of a composition or of a kit as defined above against phytopathogens of one or more plants or crops, for example agricultural crops, ornamental crops, forest crops, wild species, industrial crops, plants in parks and gardens. According to the present invention, said yeast strain, extracellular substances, composition or kit can be used before sowing, pre-harvest or post-harvest.

In particular, according to the present invention, the yeast strain PT22AV belonging to *Papiliotrema terrestris* species is effective in the control of phytopathogens, since it acts by controlling their development and so limiting the establishment of infections in plant, without directly killing the phytopathogen. Therefore, the term "against phytopathogens" mentioned above is meant for the control of phytopathogens.

The extracellular substances (or secondary metabolites) secreted by the PT22AV yeast strain, belonging to *Papiliotrema terrestris* species, during the biomass production phase, are essentially polymers of a glucidic nature, the most frequent monomer units of which are: mannose; xylose; glucose, galactose, arabinose. The structure of these polymers is organized in main chains of monomers linked through linear and branches bonds. Therefore, the present invention includes the use of wastewater, (i.e. cultural growth broth applied as it is (crude) or concentrate, with high content of EPSs (extracellular polysaccharides) compounds) from the *Papiliotrema terrestris* PT22AV biomass production process against phytopathogens, i.e. as plant protection agents or biocontrol agents against plant pathogens.

In particular, according to the present invention, the yeast strain can be used in the direct and indirect control of all harmful organisms for plants. For example, the strain can be used for the induction of resistance and biostimulation of plants, improving the tolerance of plants to pathogens, including viruses, thanks to the production of polysaccharides

and other metabolites. This highly complex matrix protects the strain from abiotic stressors, improves adhesion to surfaces and triggers an induction of plant resistance thus improving tolerance to a wide range of pathogens, including viruses.

5 Liquid or dehydrated preparations based on extracellular polymers, of protein and carbohydrate nature, or secondary metabolites, which accumulate during the fermentation and subsequent industrial transformation of the PT22AV strain, can be used as phytostimulants, inducers of resistance against pathogens, abiotic stress relievers (osmotic, 10 water, thermal, UV, etc.), healing films, adjuvants in the distribution of foliar applications or seed coating, growth medium for microorganisms, including Bio Packaging (integration or replacement of Packaging to preserve food products, etc).

 According to the use of the present invention, said one or more 15 plants or crops can be selected from cereals, such as wheat, barley, rye, oats, rice, corn, sorghum; fruit trees, such as olive, apple, pear, apricot, nashi, plum, peach, almond, cherry, persimmon, banana, vine, strawberry, raspberry, blackberry trees; citrus fruits, such as oranges, lemons, mandarins, clementine trees, grapefruits; legumes, such as beans, peas, 20 lentils, soy; vegetable crops, such as spinach, lettuce, asparagus, artichokes, cabbage, carrots, onions, garlic, tomatoes, potatoes, aubergines, peppers, fennel; cucurbits, such as squashes, zucchini, watermelons, melons; oil plants, such as soy, sunflower, canola, peanut, castor, coconut; tobacco; coffee; tea trees; cocoa; sugar beet; sugar cane; 25 cotton.

 According to the present invention, said phytopathogens can be fungi, such as for example Basidiomycetes, Ascomycetes, Deuteromycetes, Oomycetes, fungi with sexual or asexual reproduction, fungi with biotrophic or necrotrophic activity, protists or Chromista. For 30 example, the strain can be used for the control of mycotoxigenic fungi that

affect plant production both in the field and in the post-harvest stage, preventing the accumulation of mycotoxins on any agricultural crop.

In particular, according to the present invention, said fungi can be selected among the species *Albugo* spp.; *Alternaria* spp.; *Anthraco*se;
 5 *Armillaria* spp.; *Ascochyta* spp.; *Aspergillus* spp.; *Blumeria graminis*; *Botrytis cinerea*; *Botrytis* spp.; *Bremia lactucae*; *Cercospora kikuchii*; *Cercospora soja*na; *Cercospora* spp.; *Cercospora* *herpotrichoides*; *Cladosporium* spp.; *Claviceps purpurea*; *Colletotrichum* spp.; *Corynespora cassicola*; *Diaporthe* spp.; *Erysiphe* spp.; *Fusarium graminearum*;
 10 *Fusarium oxysporum*; *Fusarium* spp.; *Helminthosporium* spp.; *Leveillula Taurica*; *Macrophomina phaseolina*; *Magnaporthe oryzae*; *Magnaporthe* spp.; *Melampsora lini*; *Monilinia* spp.; *Mucor* spp.; *Mycosphaerella graminicola*; *Mycosphaerella* spp.; *Oidium* spp.; *Penicillium* spp.; *Peronospora manshurica*; *Peronospora* spp.; *Phaeosphaeria* spp.;
 15 *Phakopsora pachyrhizi*; *Phakopsora* spp.; *Phoma* spp.; *Phytophthora* spp.; *Plasmopara viticola*; *Podosphaera* spp.; *Pseudoperonospora cubensis*; *Puccinia* spp.; *Pyrenochaeta lycopersici*; *Pyrenophora* spp.; *Pyricularia oryzae*; *Pythium* spp.; *Ramularia* spp.; *Rhizoctonia solani*; *Rhizoctonia* spp.; *Rhizopus* spp.; *Rhynchosporium* spp.; *Sclerotinia sclerotiorum*;
 20 *Sclerotinia* spp.; *Sclerotium cepivorum*; *Sclerotium rolfsii*; *Sclerotium* spp.; *Septoria glycines*; *Septoria* spp.; *Sphaerotheca* spp.; *Stemphylium* spp.; *Stenocarpella maydis*; *Thielaviopsis basicola*; *Thielaviopsis* spp.; *Tilletia* spp.; *Ucinula* spp.; *Uromyces* spp.; *Ustilago maydis*; *Ustilago* spp.; *Venturia* spp.; *Verticillium* spp.

25 Furthermore, according to the use of the present invention, said phytopathogens can be bacteria, viruses or parasitic nematodes of crops. In particular, the bacteria can be selected from among the species *Pseudomonas* spp., *Xanthomonas* spp., *Clavibacter* spp., *Ralstonia* spp., *Erwinia* spp., while nematodes can be chosen among the species
 30 *Meloidogyne* spp., *Heterodera* spp., *Globodera* spp., *Belonolaimus* spp.,

Pratylenchus spp., *Rotylenchulus* spp., *Trichodorus* spp., *Paratylenchus* spp. etc.

According to the use of the present invention, said yeast strain or said extracellular substances or said composition or said kit can be applied
5 on one or more parts, intact or wounded, of the plant selected from seeds, aerial parts, leaves, stems, trunks, buds, gems, branches, stems, flowers, fruits, roots.

In particular, for field treatments on plants aerial part, the PT22AV strain, or the extracellular substances, can be applied using the distribution
10 equipment currently in use or the aerial irrigation system, for example by means of spraying equipment or overhead irrigation systems. Treatments on fruit, flowers and fresh-cut produce during the post-harvest stage can be applied using distribution equipment or added to the washing water.
According to the present invention, the quantity of living cells of said yeast
15 strain distributed per unit of culture surface can range from 10^8 to 10^{15} CFU per hectare.

According to the present invention the dosage of PT22AV-based compositions can vary according to the type of formulate and the field of application, generally from a few grams to tens of kilos per hectare. The
20 dosage can also depend on the crop, the pathogen to be countered, the pressure of the disease, the weather and other agronomic factors.

According to the use of the present invention, aforementioned yeast strain or its extracellular substances or the composition or the kit, as described above, can be applied to solid plant growth substrates, such as soil,
25 preferably at a live cell concentration of the yeast strain ranging from 10^2 to 10^{12} CFU per gram of solid growth substrate (soil or another matrix).

According to the present invention, soil treatment can be carried out by applying the formulate based on the PT22AV strain of *Papiliotrema terrestris*, by dilution in the substrate, sprayed or distributed with a
30 fertigation system. Soil treatment can be aimed at improving the chemical-

physical and microbiological characteristics of the soil and/or for the control of phytopathogenic nematodes.

Moreover, according to the present invention, PT22AV strain can also be used for the treatment of a hydroponic or floating crop system, for example against pathogens and root parasites; in this case the product
5 can be applied directly in the nutrient solution.

Furthermore, said yeast strain or said composition can be used as seed coating agents; in this case the strain concentration can range from 10^2 to 10^{10} CFU per gram of seed. According to the invention, PT22AV
10 strain used as seed treatment can be applied in combination with filming, surfactant or tackifying agents.

In addition, said yeast strain or said composition can be used by means of pollinating insects to counteract plant pathogens capable of penetrating the host through the flower (*Erwinia* spp., *Pseudomonas syringae*, *Botrytis cinerea*, etc.).
15

According to the use of the present invention, said phytopathogen can be chosen from *Botrytis cinerea* and *Plasmopara viticola* of vine; *Botrytis cinerea* of strawberry, *Botrytis cinerea* and *Phytophthora infestans* of tomato, *Venturia inequalis* and *Stemphylium* spp. of apple and/or pear
20 tree, *Monilia* spp. of stone fruit and/or *Phytophthora* spp. of potato.

According to the present invention, PT22AV yeast strain, its extracellular substances, its composition or kit, as described above, can also be applied to the seed and their use can take place in green-house, in the field or in post-harvest stage, in a biological or integrated regime.
25 Furthermore, when the kit as defined above is used, said components a) and b) can be used separately or sequentially.

"Separate use" is intended as the simultaneous application of components a) and b) of the kit in distinct forms. "Sequential use" is intended as the subsequent application, in any order, of components a)
30 and b) of the kit, each in a distinct form.

The present invention also relates to a method for the control of phytopathogens in agricultural crops comprising or consisting in applying an effective dose of the PT22AV strain, of the extracellular substances secreted by it, of a composition or a kit as defined above on said crops or
5 on the soil where the crop grows, before sowing, pre-harvest or post-harvest. The phytopathogens and the crops according to the method of the present invention can be those described above.

According to the method of the present invention, said application can take place on one or more plant parts, intact or wounded, selected
10 from seeds, aerial parts, leaves, stems, trunks, buds, shoots, branches, stems, flowers, fruits, roots.

Furthermore, according to the method of the invention, the dose of aforementioned composition varies from 10^8 CFU to 10^{15} CFU per hectare. Some examples of application protocols, with reference to the dosage, the
15 volume of application, the method of application and the site of application of *Papiliotrema terrestris* PT22AV, in the main crops of agricultural interest and against the main crops' pathogens, are shown in Table 2.

20

25

30

Table 2
Application details of formulation based on PT22AV

Crop	Pests or Group of pests controlled	Growth stage & season	Application rate per treatment			Application	
			Active substance/ha, mL	Kg, min	Water L/ha / max	Method	Site
Tomato	<i>Botrytis cinerea</i> <i>Phytophthora infestans</i>	Till BBCH 63 every 7-10 days depending on the pressure	0.5 - 1.5 x 10 ¹² per ha		500 - 1000 Taking into account the vegetation	Spray	Foliar
Grape (Wine Table)	<i>Botrytis cinerea</i> <i>Oidium tuckeri</i>	Till BBCH 67 every 7-10 days depending on the pressure	0.5 - 1.5 x 10 ¹² per ha		500 - 800 Taking into account the vegetation	Spray	Foliar
	Soil diseases & nematodes	Seed coating	0.5 - 1.5 x 10 ¹² per Kg seed			Spray	Seed coating
		Seedling (radical pre-transplant application)	1 - 5 x 10 ⁷ per mL			Suspension	Root soaking
		Transplant (fertigation)	1 - 5 x 10 ⁶ per mL				Fertigation
		Fertigation (applications following transplantation)	1 - 5 x 10 ⁶ per mL				

Strawberry	<i>Botrytis cinerea</i>	Till BBCH 65 every 5-10 days depending on the pressure	0.5 - 1.5 x 10 ¹² per ha	400 - 600 Taking into account the vegetation	Spray	Foliar
Stone fruit	<i>Monilinia</i> spp.	Till BBCH 69 every 10-15 days depending on the pressure	0.5 - 1.5 x 10 ¹² per ha	800 - 1000 Taking into account the vegetation	Spray	Foliar
Pome fruit	<i>Stemphylium</i> spp.	Till BBCH 79 every 10-15 days depending on the pressure	0.5 - 1.5 x 10 ¹² per ha	800 - 1000 Taking into account the vegetation	Spray	Foliar
	Post-harvest diseases	Post-harvest	5 x 10 ⁶ - 5 x 10 ⁷ per mL	-	Suspension in washing water	Fruit
Wheat	<i>Blumeria graminis</i> <i>Septoria tritici</i> <i>Fusarium</i> spp. <i>Puccinia</i> spp.	First application BBCH 39-42 Second application BBCH 51-53	0.5 - 1.5 x 10 ¹² per ha	400 - 600 Taking into account the vegetation	Spray	Foliar
Potato	<i>Phytophthora infestans</i> .	Till BBCH 66 every 5-10 days depending on the pressure	0.5 - 1.5 x 10 ¹² per ha	500 - 800 Taking into account the vegetation	Spray	Foliar

Ornamentals	Foliar pathogens	All phenological stages with intervals of 5-15 days depending on the pressure of the disease	5×10^6 - 5×10^7 per mL	-	Spray	Foliar
Turf	Fungal pathogens	All phenological stages with intervals of 5-15 days depending on the pressure of the disease	$0.5 - 1.5 \times 10^{12}$ per ha	500 - 1000 Taking into account the vegetation	Spray	Foliar
Plant Biostimulant	-	Application to soil/root during all phenological stages with intervals of 5-10 days	$0.5 - 1.5 \times 10^{12}$ per ha	In fertigation	Drip irrigation	Foliar

The dosages and volumes of distribution are exemplary and can be varied according to the disease pressure and the type of vegetation.

The present invention will now be described, for illustrative but not limitative purposes, with particular reference to the examples and attached figures, in which:

- **Figure 1** shows the colonization of *P. expansum* hyphae by the biocontrol agent PT22AV. Image of the *in vivo* interaction between *P. expansum* and *P. terrestris* strain PT22AV observed under a scanning electron microscope. The yeast colonizes the fungal hyphae and adheres to them by means of an extracellular matrix of a polysaccharide nature;

- **Figure 2** shows foliar colonization by the PT22AV strain 5 days after the application. The image shows the yeast-leaf interaction 5 days after application observed under a scanning electron microscope, in particular the yeast on the upper and lower page of tomato leaves, where colonization of a stoma, natural opening, is also shown;

- **Figure 3** shows the PT22AV yeast grown on an agarized growth medium after 3 days of incubation at 28 °C;

- **Figure 4** shows the phylogenetic tree of the PT22AV strain obtained by analyzing the nucleotide sequence of the large subunit ribosomal RNA gene and comparing it with that of other related species;

- **Figure 5** shows the phylogenetic tree generated by comparing the entire proteome of the PT22AV strain with that of other species of basidiomycete's yeast.

EXAMPLE 1. *Isolation and characterization of the PT22AV strain of Papiliotrema terrestris.*

The PT22AV strain was selected from a yeast collection isolated from the epiphytic microbial community of various fruits and vegetables in central Italy.

The PT22AV strain was identified by sequencing the large subunit ribosomal RNA gene, clearly highlighting belonging to the *Papiliotrema terrestris* species (Figure 4).

In particular, the nucleotide sequence of the large subunit ribosomal RNA gene region of the PT22AV strain is shown below:

```

      CCGGGAAGAGCTCAAATTTGAAATCTGGCGTGCTCAGTGCGTC
      CGAGTTGTAATCTATAGAGGCGTTTTCCGTGCCGGACTGTGTCTAAG
5      TCCCTTGGAACAGGGTATCAAAGAGGGTGATAATCCCGCACTTGACA
      CAATGACCGGTGCTCTGTGATACGTCTTCTACGAGTCGAGTTGTTTG
      GGAATGCAGCTCAAATGGGTGGTGAGTTCCATCTAAAGCTAAATATT
      GGCGAGAGACCGATAGCGAACAAGTACCGTGAGGGAAAGATGAAAA
      GCACTTTGGAAAGAGAGTTAAACAGTACGTGAAATTGTTGAAAGGGA
10     AACGATTGAAGTCAGTCGTGACTGAGAGGCTCAGCCGGTTCTGCCG
      GTGTATTCCCTCAGTCGGGTCAACATCAGTTTTGTTCCGTGGATAAG
      GGCAGCTGGAAGGTGGCACCTCCGGGTGTGTTATAGCCAGCTGTCTG
      CATACATCGAATGAGACTGAGGAATGCAGCTCGCCTTTATGGCGGGG
      TCGCCCACGTCGAG (SEQ ID NO:1).

```

15 The Applicant obtained the entire genomic sequence of the PT22AV BCA strain by Nanopore sequencing technology. The bioinformatic analysis of the sequence of the entire PT22AV genome allowed the prediction of the entire proteome, also used for the taxonomic analysis of the strain, hence precisely defining the differences with other strains of related species (Figure 5).

20 The entire mitochondrial genome of the PT22AV strain consists of the following sequence:

```

      CTATGAGTATAGCACTAGGAAGCTAACCATTAGGGCGAATAGCCCTG
      TAGCTTCGGCTAGAGCGAATCCTAGAATAGCGTATGTGAATAGTTGT
25     CCTCGAACAGCAGGGTTTTCGAGCTGTACCAGCGATTAGAGCGCTGAA
      TACTGTACCGATCCCAACACCGGCTCCTGAGAGACCGATAGCGGCTA
      GACCAGCACCAATGAACTTTGCAGCAGCCATTGTGTATTGTGATTGTC
      TGAACNNNNNNNNNTCAATGAATTGAGCTTTAGCCTTTATAATCTA
      AAGGCTAGTGATGACATCCATGAACGGCTATCACCTGTATAAAGACA
30     GTGATAGAATATATGTGATCATGGAGGGGACAGACGTGGATCTCTAT
      TATGAACTCTATAGTGCGGTAGCTGGATCTCAGACTATATTACCGTAT

```

CTACCCTTTTAGGACCTATAATCATGCTTATAATATCTATAGATGGTAG
ATTACCTAACACTCGGTGCTATAGTAGGTTATAGGAGTATTATACTCC
TATAACCATAAACTATGAATGGCTACTATCAGCATGAGCAATGACTCT
ATTTACCGCTATAGTAATGGTGTACTCTACCATTTTTCTATAGGATCC
5 TACCTATCCCTATAAGCTATAGGGTTTTCTATGAAAATGCTCTACAATA
CATGTATGCTAACGTGAGTCAATGAGTATAGCCTCGGAACTTTATTT
ATGCACTTCTGACGCCATGCTGAATATAATAGCGCCACATGTATGCAT
AGCTTTTATCTCTCTCTATAACAAGTTATTATTTATTAGTAAATTTATTAC
TAAATTTATTTTTAATAAATAAATTATTATTATTATTATATTGGTATGGAC
10 GGACAGAGCAAGATTCGAACTTGCGGTGGGCGGGACCCACTTTGAG
GTTCAAGCTCAATGCCTTAGACCACTAGGCTATCTGTCCTGTTATCTC
ATCGACTCACCTGCTACCCTTAAAGAGGGGTAGGAAAAATGGTGAAA
TATAGTATATTGTCTATCTATAAACTTTTATAAGAATAGTTACGGTCAT
AGATTTTTATCGTCTCGTCTCTACACAATGGCTATAACCCCTTACTATT
15 AAGGCCAGCCTATATTTTTTGCTGCTATAGGTGACACAGAATATTTAT
GTTAAAGAACATAAGGGGGCTAGTATAAAGTCCAAGCATAATAGCTTT
GTAACGTATACACGGCTCACAATCAGTTGAAGGATCGGAGGTAGTAC
GAACACTGATACAACGTATACCAGTGCAGCCAGAGTAAGGAGAACGA
ATGCTCCTTGGTTAAGGAAGTAAAATGGGAGAGCTTGAGGCATAGGC
20 TGATATGCAACAGGCAATGTGATAGGTGTTTCCATCATCACTATGAAT
ACTATAATTATGAGTGACTATATAGCATGCTCATAATAACAATAGACTC
TTACATACTTTCCTTTCNNNNNNNNNCTTCTGACGCCATGCTGAATA
TAATAGCGCCACATGTATGCATAGCTTTTATCTCTCTCTATAACAAGTTA
TTATTTATTAGTAAATTTATTACTAAATTTATTTTTAATAAATAAATTATT
25 ATTATTATTATATTGGTATGGACGGACAGAGCAAGATTTCGAACTTGCG
GTGGGCGGGACCCACTTTGAGGTTCAAGCTCAATGCCTTAGACCACT
AGGCTATCTGTCCTGTTATCTCATCGACTCACCTGCTACCCTTAAAGA
GGGGTAGGAAAAATGGTGAAATATAGTATATTGTCTATCTATAAACTT
TTATAAGAATAGTTACGGTCATAGATTTTTATCGTCTCGTCTCTACACA
30 ATGGCTATAACCCCTTACTATTAAGGCCAGCCTATATTTTTGCTGCTA
TAGGTGACACAGAATATTTATGTTAAAGAACATAAGGGGGCTAGTATA

AAGTCCAAGCATAATAGCTTTGTAACGTATACACGGCTCACAATCAGT
TGAAGGATCGGAGGTAGTACGAACACTGATACAACGTATACCAGTGC
AGCCAGAGTAAGGAGAACGAATGCTCCTTGGTTAAGGAAGTAAATG
GGAGAGCTTGAGGCATAGGCTGATATGCAACAGGCAATGTGATAGGT
5 GTTTCCATCATCACTATGAATACTATAATTATGAGTGA CTATATAGCAT
GCTCATAATAACAATAGACTCTTACATACTTTCTTTCTATGAGAGGTT
GGTTAGTTAGCAAGGGCAGAAGGGGACTATAGTTATTGGTAACCATA
TAAGGTCGTTATAACGTTTCAACAAGCCTATGATTGGATAGGTAGCAT
GTCGAATGCGTGGTGTGGAGTAGGTGATGGTAGTACCCACTCGATTG
10 TGTAGGCGAACCGTGTGCTTCGTTTCGTACGTTAGGTTGTCCGTGAAG
TATAGCGGTTCTCCCCATGGGTTGTCACCAACGTA CTGTGCGTTAGT
GAACATGTCGTAAACAGTGTACAGGAACAGAAGAGTAGCAGAGACGC
TAACGAATGAACCCCATGATGAGATTAGGTTCCATCCCGCGTAAGCG
TCAGGGTAGTCAGCGAATCGTCGTGGCATTCTTGTAGTCCTAGGAA
15 GTGTTGTGGCATAAATGTCATGTTAACACCGATGAAGAATGTCCAGAA
GTGCGCTTGAGCTAGGTGTTCTTCGTACATCTTACCGAACATCTTTGG
AGCCCAGTAGTAGAACCCGGCAAAGATACTGAATACGGCACCCATTG
AAAGCACGTAGTGGAAGTGAGCCACAACGTAGTATGTGTCATGCATC
GCTACGTCAACGCTCGCGTTCTGCTAGAACTACACCTGTCAGTCCACC
20 AATAGTGAACAGACCGACGAATCCTAGAGCGAATAGCATAGGTGCTA
GCAGTCGGATAGAACCACCGTAAGCAGTCGCTAGCCAACTGAAGAC
CTTGATACCTGTAGGTACCGCAATAATCATAGATGCAGCTGTGAAGTA
AGCTCGTGTGTCTACGTCCATCCCTACGGCGTACATGTGGTGGCTCC
ATACAATGAATCCAAGCACACCGATAGAGGCAATGGCGTACACCATA
25 CCAAGGTAACCGAAAACAGGCTTTCCACTGAATGTTGAAACAACGTG
GCTAACCATACCAAATCCAGGGATAATCATTAGGTATACTTCAGGGTG
TCCGAAGAACCAGAATAGGTGTTGGTATAGGACAGGGTCCCCTCCTC
CAGCAGGGTCTGAAGAATGATGTGTTGAAGTTACGGTCAGTAAGGATC
ATAGTAAGAGCTCCGGCTAGAACTGGAATACATAGAATAATAA TAACT
30 GACTGTGATAGCATCGCCCATACGAATAGAGGTGCCTTGTGTAGGCG
TAGCCCTGGAGCTCGCATGTTAAGGATTGTAGTGATAATGTTAATGG

CACCAAGTAGAGATGAAATCCCTGAGAGGTGCAGGCTGAATACAGCT
AGATCAACTGATCCACCACTGTGTGATTGTACACCTGTTAGAGGCAT
GTACATAGTCCAACCGCTACCCATACCTTGTTCCACGAAGACGCTTG
AAAGGATAAGAACGATTGCAGGAGGTAGTAGCCAGAAGCTGATGTTG
5 TTCAGCCGTGGGAACGCCATGTCCGGCGCACCGATAAGAACGGGAG
CCATGTAGTTAGCGAACCCAGCCATAGCAGGTACCACCATGAAGTAA
ATCATCATGATACCGTGAGTTGTAGCAATTACGTTGTAGAGGTGGTG
GTTACCAGAAAGGAACTGGTTACCAGGAGCTGATAGCTCCATCCGGA
TAAGTACACTGAACGCAGTACCCAGAAGACCGGTAAAGACAGCAAAA
10 AGGATATAAAGGGTACCAATATCCTTGGCGTTTGTAGAGAAGAGCCA
TCGGCTAGCCATGTTAATAAGAGGGTTATGCTGTTATGGGGAGGGGT
AGGTAGGTTGACAGCTGCTGGTGAGAGTAACTATAGTGCATAGCAAT
ATTCATGGTAGAGTATTCTATCACTATAGGCTCTAGTTATCTACCATAA
AAATGGACATAGCGGATAGCCATGAGAACTAAACAGTAGAACTGA
15 TAGATGCCTTAATAGGCAGAAAGTGTCACAAATGTGAGGTCTTTAATC
CACAATAAGACTTAGAGTAAATAGAAGGGAGGAGGTTACCGGTATT
AGTAAATCATAGAGGAGAGAGAGAGATGCATCACTAGTAACCCATAG
GGTACACAGCAGTGCTAGTAGCAAGTCTTTAATAATCATATGTGATAG
ATGGCGGAATAGCCTAATCTACACACTATTAGATGTAGGTACTTTAGA
20 TATGATATAATTTAAACATTTATAGCTGTAATTATAAAAACCTCGCTATA
GGCTTATAATATTATAATATGACCATATATTTTATATAAAGCATAACAA
CTATGTATGGCAATATAAGTCCTATAGCCATCTAGCACTACAATGCAG
AGTGTCTAGGTATGACAGTTTAAGTAACAGCCTATATACATAACAAC
AAACCTTATAACTGTACAGGGTCTTAGTCCACCACATGTCACACACGA
25 GTATGGACTTCTGCTTAGGCAAAGCCCCTTTATACTAACCTTCCCATC
AGAAACCTTTATACTCAACCAAAGGGTTAATGTTAGGTTTATAGGGTA
GTTGTTATTTAAAAAATAAGGAGGGGTATGGAGGCATGCTTTGCGA
GTATACCATAAAAAACAAATAACATATAAATATGACTGCACAACAACGAT
CATTTGTGCGAGGCGCTACTACGTCTACGGCCGCTATTCCAATCACAC
30 CCTCACCACCTTACTCTGGAATCACCATGGCCAATCCTTACAAGTGG
AGCTGTGCTAGCTATGCTTTCAAGCGCTGCCCTTTGGTTCAACGGTC

TACAATACTCAGGTACATTTCCTTATCCTTGGGATGCTTAGCACACTAA
CAGCTATGATCCTATGGTGGGCTGACTGTGTTAAGGAAGGTACATAC
CTTGGACACCACACTAAGGTAGTTCAACAAAACCTTGCTATGGGTGTT
GCGCTTTTCATTGTAACAGAAGCCGCACTATTTGTGAGCATCTTCTGG
5 GCCTACTTCCACTCATCACTAGCACCTACAGTGGAAGTACTAGGTACTCAA
TGGCCACCCGCTGGTATCACAGCCCTATCACCTATGGCTATCCCCCT
ACTCAATACCGTTCTACTACTTAGCAGCGGTGCTACAGTTACATGGG
GTCACCACGCATTCTTCTGCCGAAACCGATCAGCTGCTCTTCAAGGA
CTTATNNNNNNNNNCAAGGACTTATGTTTACAGTAGCTCTAGCTATT
10 ATCTTCACTGCCCTACAAGGTGTTGAATACAACATGGCTGGATTACT
ATTGCTGACGGTGCTTACGGTAGCTGTTTCTATATGGCCACAGGTGC
TCACGGACTACACGTGCTGGTAGGTACACTAGCTATCACTGTTGGTC
TTGCACGACTTATTAACCTACCAACTTACATCAACTCACCACATTGGTC
TAGAAGGAGCAATCCTATACTGGCACTTTGTGGACGTCGTTTGGGTC
15 ATCCTATACGTTACAGTTTACTGGTGGGGAGCTCCATTATCCCTATCT
AAATTAGTCCCCTATTTATATATTTTCCCCTTCCAGAATACTACTTACC
TTGTTATACTTAACCTTTTAATTATAAGTTAATTGGCTGAGTAGAATAG
GGTTACAATAGAGGTAGATGAGGAACAAGAACACTATAGTATGCTTA
GTTACCATAGAAAATTTAAATTTTTATAAGCTATGAAGTGACTATTTCT
20 TAAATGGCTATCAGTTATTTAAGACAACCTTGTTAGAAGTCAATAGTCAT
AATATCTAATAGCATGGCTCATTTTTTATATTAATAAATATTAACAGTATTTA
AAATTTAAAGTCATTTAATGGTATACTGCGTTTAGTATAAAAATAGAGG
ATTATATTGACACGTTTACCAGCCGAATTAATACAATTACTAAATTTAG
AGTGAGTCTAATTCAGGTGGGCTGGGCTCGAACCAGCGGCCTCATG
25 CTCCCAAAGCATACGTCTTAACCAAACCTCGACTACCACCTGTGTACTC
TTAGCTACCGGCCCCCATACAAACCATCATTAAAGTTGATAATTATTGA
TACTATTAACCTATTGTAGTATCTTTAATGAGGGGTCTAAACTGGTACA
ACGTCAAACGCAACAAGTACGCTTGGTACCAGTAGAAGTAGAGCGAT
AGCTGGTGGTAGCATTCCAGTCCAACAGAAAGAAAGTAGTTGATCGT
30 ACCGAAGTCGTGGTAGTGTAGCCCGGAACCACACAAAACCAAACAA
CCGGCACAAGTCTTAAGCGCCAGCACTAGTGATGATAGTGACAGTGG

TGTACCATTGGCAATAACTTCAGGGAATCCCCATCCACCTAGGAATA
GTGTAGCTGATAGCGTGCTCATGAGAACGATAGAACAGTATTCTCCT
AGGAAGAAGAGTACGAAAGGCATAGCACTGTGCTCTGTCATGAACCC
AGCAACAAGTTCTGATTTCAGCTTCAGGAAGGTCAAAGGGTGTTCGGT
5 TTGTTTCGGCTAGTACAGATACAATGAATAGCATGAATAGAGGCGCTA
GAGGAACACAGTACCAGATAGGTGCTTGTGATTCTACAATAGCCGTG
TGGTTAAGGCTACCGGCCATGATCAGGACTGCTAGAACAGCAGAACC
TAGGATCAGTTCATAGCTAATCATTTGAGCTGTTGATCGTAGCCCACC
AAGAAGAGCGTATGATTGTTGGCAGACCATCCAGCTAGTAGTACCC
10 CATAAACACCAACACTAGAAACAGCTAGAGAGTATAGAACCCCTAGTT
CTAGGTCAGCAATAGCGGCACCAGGACCGAATGGCATAACAGCCCA
ACCAAGAAGGGCGAACACCAGTGTAATAACAGGTGCTAGGTAGAAGA
TGACGGTGTTAGAGGCCCCCGGAACAACAGTTTCCTTTACCACCAGC
TTCAGGGCGTCAGCGAACGGTTGTAGTACTCCATAGACACCTACTGC
15 GTTAGGCCCTACTCGTCGTTGCATTGCAGCCAGAACCTTACGTTCGA
TAATGGTCATGAACGCAACAGCTAGAAGAATAGGTACCAGTACAGCT
AGCACAGACAGAATGGAGAAGAGAGTAGAAAGCATATTACGTAGTTT
GTAGACATAAAACCCGTGATACTTAAAAGACTAAGACCAGTCGGGAG
GTATGGGAGAGGTACACTATAAATAATAATAATGAATGAATATATATA
20 GACAGACAGTACATAACTGTGCCGTATGGTTATATTTATCAATCTTAC
CTAATTGTTAAAACATGGTTATAATAACCACCCTTATAAGTATAACTAA
GGTAGTCTATAATGGTTATAGGAACCAGAATAGTCATGAATATAAGAG
CACAATGGCTCCTAGCGATACGTTAAGATAAATAGTGTGGTGATAGTT
AAATAAAGGGGTCTATAGCAATGGCAGAAATAACCATAGTCAATAAGA
25 CCTAGGCTTGTGACTTTAGCCATGCTAGGTATTCTGGCAGGCTAACC
GCTTCAATACAAATAGGCATGAAGGCGTGGTAAGCTCCACAAAGTTC
GCTACATTGTCCGTAGTAAACACCTTGTCGTTGGATCAGAGTAGATAC
TTGGTTTtagTCGACCTGGACAAGCATCGACCTTGATACCTAGTGAAG
GAATAGCCCAGTCATGCAGAACGTCAGTAGCTGTTACGATGGCTCGG
30 ATGTGCGTGTCAACTGGAAGAACTAGCCGGTTGTCCACTTCTAGTAG
TCGTAGTTGCCCAGCTTCTAGTTTCGTCAGCAGGAACCATGTATGAAT

CAAATGCAACTGTTTCGTTGTCGTCACCAGCAAAGTCTGATAGTTCGT
ATGACCAGTACCATTGGTGTCCAATAACCTTAACAGTCATAGCTGGGT
TAACCACCTCCATGTGGTTCCCCTTGTAACCTATAGAAAAGGAATAG
GGTTGATTATGGACTATGCCTTGCTTCCCACCACGATAGTTCATAACT
5 ATGTTAACAATCCTGAAAGGACTTACCCCGACCCTCCTAACACCTAGT
CTTTTTTATTACCCTACCCCCCAAGGGGGGGTTAAAAGGTGGGTTA
GGGCGTAAGGAGGGAAGTAGGGGGTTAATGTGGGATTATAGCACCG
TTGTAGTCTCTGAGGCCTACGGCCTACCTGTACGGGATATAACCCAG
CCAGGTATACAGTGGTAAATGCATGGACGAGAGTAACTGTGTTGATT
10 TTTCCAATTTTTTTTACCTATACTGAGCCTAGTATAGGGGAGGTTTAATT
AAAATTATCAAACACAGCTCATAGCTATGATTTCTATGTCCTATAGGG
AGTCCATGAGGTATAGCAGCTTGAATGAAGGGAAGGCAATAGCCACA
AGGATAAGTGCTGGCACAACAGTCCATACAACCTTCAATAAGAGTACC
GTGGTTGTTGTACCGGTGAGCGATAGGCGCTTGTGTGTCCTTGAAAG
15 AGATAACGATAGATCCGAGCATCCATCCTACTGCCACACCAATAACG
ATGAGGTAGAAGAAGATAGTGTCGTGTAGTTCAGTGATCCCCTCGAA
TGATGGAGAAGCACCGTCTTGAATCCTAGAAGCCAAGGTTCTGGAG
CGTCACAAGTAGCTGTGAATAGTGATGTAAATAGTGAAAGCATGCTAT
ATAATGAATATTAAATATAAACTAGTCGTGTCAATCGCTACCAATGATA
20 AAACACTGATAGTGGTACTAGACCAATGTGACGTCATAGTCACTTTAT
TAATCGCATAATACAACCTTGTTGAGAGGTGAAAATGAGATAATAGGT
NNNNNNNNNGCGGACGTGTCAGGTAGTAAGGGGTACTGATAACCA
TAAATGGTTACTAGCTAATAACGGTTAGCTGAATCGAACAGCCTTTAC
TAATGAGATCAACCCATTCTTTAGTAATCCAATACCGTGTATATTAATA
25 AAAGTCAGTCTAAATAAAAGATATTGTATTGTTAATAGACAACTCTATT
ATAACTATTTGGTAGGTAAGAAGAGAAATATATTACACACCATAGCTA
TAGGTCTCTCTAAGTACAATGAGAGTGGTAAATAACCTATATCTATTA
GGGGAATATTAGTCCTTATGACCTAACAGCCCCAACCCAATAACTAAT
GGGTATGTAAGAAGGTTAATAAGAGGTTGGTAGAATTTCTCTATTGCT
30 ACTAGTGTATAGGCCTATACTTATAACTATAGGTCTCTACTTATATATA
TATATCAGTTAGGTGTACTTATAACCTATTACTAGTAGTGTATTATTAA

TACCAAAAGGTGAGTAGGGGGCTAATTTACCATTCATCTATCAGCCG
CAGGTTCCCCTACGGCTACCATGGTTCGACTTCAGACCAGTCAGCTG
GCCAGGCTCTGTAGCAAGCTCCGCCATTATTACTGTCTTCCAGCACC
CATTGATATCAATGGGAGGTCTCTCCATCATTATGACTATAGGTTGAT
5 GTATAATTATAATGTGGATATATATGAGACTATAGTTAAATACAAGCAG
CGCTCACCCACGTTACCCTACCAGCATACTAGCCCGTGACGAGCGGT
TAGTACCCAGAGAGATTAACTTCACCGCTCCATACTTATGAGCGAT
TACTCGGGACTCCAGCTTCACGCGAACGAGTATCAGCTCGCGATCTG
TATGCTTGTGGTAGTTTTGGATGCTTTCCCATTTTATAAGGCTAGCTC
10 TCATCTGACACACACACTTGTGGCAGGCCTGTAGCCCTTCCCATAAG
GGCCATGACGGCCTGACATAGGCCCGTACTCGTATCCCTTATAAGGC
TACTACTGCCTATACACTAATCAAATTAATTAGCATCATAGACAGACG
GGTCTGCGTTCGTTTCATTGGACTCAACCGTAGGGCTCACGACCACGA
ACTGACGACGGCCATGCAACGCCTGTAAATTACCAGACTCGCTCTAA
15 ATTAATAAACTATTAAATAATGAGCGAGCCAACAGTAAAATATGCAAG
GGAAGGTAAGGTTTTAAGCGTATCGTCTGATTAAACAGCATGCCCCG
CGCCCGTGTGCTCTCCGGCGGTCTCATCGTTAGTGTGTGCGGTCTTG
GACTATACCCTACAGGCGAGTTGCTCAGGCGTTAGCCTTCCAGGGGT
TAAAATTGTGACTATGCTTGACGTCAAGTATTACACTCTATATTCAAGT
20 CGCACTCCCCACTGCGGCAACTATGGTTCAGTCTGGGGCTAAACAG
GCTCTACTCTGTGTCGGTATCCCAGCCTTCGTACCTTACCGTCAGTCT
GTCACCAGCACCAATTGCATTGCCACCCCTAGGGATCATAGGATCT
CAGCCCTCCCCCAGTGTTGGTGGTACCAAATGTAGACTCGTATCCT
ATTAGAGATGGCTGCGTACCCTATACGCCCATTC AACACCATAAGAG
25 CTGACCCGTCTGTGTTACCGCGACTGCTGGCACAGATGGTTGGACG
GGTCTTGACATAATAAGCATCAATATGACTTATTATATCTGGATATTA
AGATATCCAATCTAGGTCACCTGGTCACACCTTTTCCCTAACCCCCCT
ACGGGGGGTTAGGGACGCGGTGCATTGCCAAGGATTCTTCACTGCT
GCGCACATGGGTGCTCTGGCTGGCCAGTGTCCGATGCATAGCTATC
30 ACTATCCCCTTCCCGTCATAGCCTAAGTAACCTCATAGCTGGTAGGG
GCTGAATCTACGACAATATGGTGGATTACCTTTGCACCATGTATAAGC

ATTAAGTATGATATCACACAACTCTTAAAAAATAAAAGTAGTGATCAAT
TTAGAATACTATTCATGATGATTATCCCTGCGCTATAGGGGCCATTGC
GTATGATTCCAGCACTGCTGACCATTTGGCTCCTATTCACCATAATGA
ATCGGAAAACTTGCATGTGTTATACGCCTAGATACCCTTCATGGCTCG
5 CCACGATGAAACGATCGCGATTATCCCTTCTATCAATACCTATATTATT
ACTTATTATCATCAATATAGGTACTACGCCCCACTAACCCCTTACTAA
CCCTTACCCCTCCAATGGAGAGGTTAAAAGGTGGGTAAGGGCTTGG
GCGGAAAGGGTCTAGATCATCCCAGTTACCACATTCAATNNNNNNNN
NNATTGTGACTATGCTTGACGTCAAGTATTACACTCTATATTCAAGTC
10 GCACTCCCCACTGCGGCAACTATGGTTCCTGCTGGGGCTAAACAG
GCTCTACTCTGTGTCGGTATCCCAGCCTTCGTACCTTACCGTCAGTCT
GTCACCAGCACCAATTGCATTGCCACCCCTAGGGATCATAGGATCT
CAGCCCTCCCCCAGTGTTGGTGGTACCAAATGTAGACTCGTATCCT
ATTAGAGATGGCTGCGTACCCTATACGCCCATTCAACACCATAAGAG
15 CTGACCCGTCTGTGTTACCGCGACTGCTGGCACAGATGGTTGGACG
GGTCTTGACATAATAAGCATCAATATGACTTATTATATCTGGATATTA
AGATATCCAATCTAGGTACCTGGTCACACCTTTTCCCTAACCCCCCT
ACGGGGGGTTAGGGACGCGGTGCATTGCCAAGGATTCTTCACTGCT
GCGCACATGGGTGCTCTGGCTGGCCAGTGTCGGATGCATAGCTATC
20 ACTATCCCCTTCCCGTCATAGCCTAAGTAACCTCATAGCTGGTAGGG
GCTGAATCTACGACAATATGGTGGATTACCTTTGCACCATGTATAAGC
ATTAAGTATGATATCACACAACTCTTAAAAAATAAAAGTAGTGATCAATT
TAGAATACTATTCATGATGATTATCCCTGCGCTATAGGGGCCATTGCG
TATGATTCCAGCACTGCTGACCATTTGGCTCCTATTCACCATAATGAA
25 TCGGAAAACTTGCATGTGTTATACGCCTAGATACCCTTCATGGCTCGC
CACGATGAAACGATCGCGATTATCCCTTCTATCAATACCTATATTATTA
CTTATTATCATCAATATAGGTACTACGCCCCACTAACCCCTTACTAAC
CCTTACCCCTCCAATGGAGAGGTTAAAAGGTGGGTAAGGGCTTGGG
CGGAAAGGGTCTAGATCATCCANNNNNNNNNNNATCATCACAGTTAC
30 CACCTTCAATAAACTTATTGGTAGGTTGAAGGCTATAGTCTGGATCTA
ATGATCGTCTTATGCGTAATTAAACGCGGTTCGTGTTGTACTATTAGG

TAAAAATTTTGTAGCTAAACGCTCTAAGTTGATATAGCTATGATAAGAG
CGATGAATTTATTGACCAACTTCTGAGTCGTCAATGGGGACGAATACT
TTACCTAATCTGTATGATACACCATTTCATAGGCATAGGGTAAATTAGA
ATCTATATTACTAGCAAAATTACCCCTATCAAAGTATAATCTTTCGTAC
5 TAGGTAATAACATGGTGGTACTCTTCACTAAGTTTTACCATTTATCAAC
TCTAGAAGAGGGTGTAGTAGGAGGGGTGCGAAACCATAAGGACCAC
AGTGGACATTATTGATAGTTGGTGGGAATAATTAGTCATCATCATCAT
CATAACAATAGGCCATGATAAACAGGGTATTTTTTTGTAATCAATAATTA
TAAATATAACTATTACCAGTAGTGATGAGTTTATAATTATGCCCTATAA
10 CTGTCTATGTTTTCCATTCCATGTTGATTATAAGGGGTCTATGCTAT
GATGATAGGTTGTGTAATGCTTATCGTGATAGCTTTGATGGAGCGTTG
TCACCAGGTAGTAGTGAAACAGGTGTAATCGCTGAACGTTGGTCTGT
GATGGCCAGAACTCCAGAGGCTACTTCTACTACGAATCCTAGTGTCA
GTACTAGGAAGAAGATAACGGCAACCCAGAAACCGTATAGACCAACG
15 TGTCCAAGGCTAGGAGCAAGAGGGATCAGTGTAGCCACTTCTAGGTC
AAAGGCTAGGAATAGCACAGCAACAAGGTAGAAGCTGATACTGAATG
GAGCTCGTGTTTGTCCGCGAGAGGCGTCATAGCCACATTCGTAAGTA
GTGACCTTTTCAGTGTCAGGACGAGCTACAGAAAGTAGTGCGTTTAC
AAGCAGTAGAATCACGGCCAGTACAGGCACAATGATAACAAGGCTGT
20 ATAGAGTAGTACTCATAGTCATGGTTATACGCTAAATAGTGAAAGAGC
CAGGAGACGAGCGCTATCTAGTAGTAGTGATGGTGATGCCATGTAGA
AGACTAGGATGGCAGTAAGTAGAGAAATAGTGTATGATTGTACAGCT
GTAATGTCTACAGGCTTTTTCTGAGTTACGCATAACAGCCCAGTAATAG
TCCCATCGAGAGTTAGCAAAGTGGTCATGAGGGAATGTTTGTGGGCT
25 GACTGGTGCTGATGGGTCAAACCAGGCTACTCGAATAACGTTAAGGT
AGTATGATGCACTGATAACGCTAGAGAGGATAGCAACTAGTGATACG
AAGTAGTACCCACCACTGATAGCGCTGTAAAGCACTTCTAGCTTACC
GAAGAACCCTAGTAGTGGTGGAAACACCAGCTAGTGAGAATAGTAGTA
CTAGTAGGCTAAAGGCTAGAGGTAGGTTTCGTACACGTAGACCAGCT
30 AGTTGTGAGATCAGTGTTACGTCATGGGTAAACGTCACCTCCACGAGA
TGAGCTATGGCTAATAGCTCCTAGCAGGCTGAATGATAGGATAGCTG

TGAGTGTGTATTGTACTAGGTAGAACACAAAAGCGCTAGTAGCTTCA
GGGCTGTTTACACTCATAGCTAGTAGCATGAACCCTACGTGTGAGAC
TGTAGAGTATGCTAGTAGTCGCTTGATACGTCGTTGTGCTAGACCAA
CTACAGCACCTACAACAAGAGACAGTGTGCTACTCATAAGTAGTAGTA
5 GTGCCCATACGTCGTAAGAGGGCTGTATCAACAACAACAAGCATAGCA
CTGATAAGACCACTGCTTAGGGTGTATAGTAGTGTGCTAGGCTGAT
CTTAGGTAGAACTGCAATCCATGTAGTCACAATTGTTGGTACCCCGTC
ATAAACGTCAGGACCCCAGTTGTGGAAAGGAGCAGCTGTAACCTTAA
ATAGAAGACCAGTCATAACAGCTGTAAAGCCAAGAGCAAGAGCCCAG
10 TTGATGCTGATAGAGTCTGATCCACCAACACTCAGAAGTGTGACTAG
AGATTCAAGGTTAGTAATACCTGTTTGGTTGTAACTACGGCAAAACC
TAGTAGGATAAGACATGATGATAGACCACCTAGTAGGAAGTATAGTA
GACCACTATGAGTAGCAGACTCTGAATCACGGTAAAGAGCTGCTAGT
ACATATACAGCAAATGACTGTAGTTCTAGAGCCAGGTATAGAGTAACC
15 ATGTCACCGCTAGCGACAAGGAGACATCCACCCATGCTAGAAAGAAG
TGCAAATAGAGGATAAGTGGCAATACGAGGAGTAGCCATAGGTGATA
GATGAGCGTGAGGAGCCCATTGGACTAGAGCAATAGCACCTACAGAA
AGTAGGAATACTTGTGCAGCTAGTGAAAGACCAGTCAGATCAATAAG
TCCACTGTATAGACCAAGACCTGATCCTACCCCCTCCCAGCTATAGG
20 CATTGGCGGCCAGTGCAGCAGCTGAAAGGAATGTGAGGACTGTAAC
TCGTGTGAGTAGAGCTGGCGTAGCATAGACCGAAGGAAGCGCAACG
GCTGTCAGTAGAATAAATAGAGCTAGTGTAAGCATTACTGTTGGCGG
TAATAATAATAGTTGTTATCTCAATAATCATAAAGAGCTATAGTACTAT
ACCTACTCCCTATTATAGACCCTATTCATGCCTAATAACTATAATATCC
25 AAGAAATCTTTGATTACTAAATAACAACCTCTATCATTATATATGCCTAT
ATTTTCTGGGTATATTAATTAGACGACAATCCATAGATATGAATAATAC
ATATGGCTTATACCTACCTATAGGCTTTAAGAATTAGATCCTAATACCA
CCCCTCAGTGTTAGGCATGCCATGTCTTTAATAACATTACTATTCTCTA
TCCCTGTGAATCTCGTAATACAATAAAAAGTATGTTAAGAGTATGTTTG
30 TTTTAAATTTAAATCTACTCACTAACCCTATGGCCATAGGACTCAAGCT
GTACCGTTAATTAGCATTTAACCTATAATGATACAAATTAGAAATCATAT

CCTAAGGAATATCTGTCGTATAGGTTTTAATACAAGTACCGTGATTTG
CACACGGACCATCTGCTTGGAAGGCAGAGACGCTACTCTTACGCCAT
ACTTGTCAGGGAAAGGGTCATTTAATCTATTGTGATAAGTATATTTAC
CATTATCGGTTGGGCTACTTCACTGTACTATTTAAATGGTGAGTCTTG
5 TACTATTAGAAGAATAAATCAGATATTCTTAGTAACACTTCGCTTGAAT
AGGATTAGGTATAGTAAACAGTGGTACTACTTCTTGTTATACCCTCTA
CCACCCCTACTAACCCACCAAAGGTGTGTTAGTAGGCAGGTAAAGTG
GGTAGAGGGAAGGAAGAGAAGGGTTAGTATCGGAGGGATAATCTTAA
TGATTGTAGACGTATTATTACTTATACCTCATAGGTGGTGTATAGAGTT
10 TATAATTTTGATCCTTTAAAGAGGCTATATAAATGGAGTGACACGGAC
TCGAACCGTGCTCTCAGCGATGCAAACGCAGTATACTACCAACTGTA
CTATCACCCCTCCCNNNNNNNNNNCACCCCTCCCTTAGCACTAGTTA
TTTAAGTTTTTTCTATTCTAAGGGGTCTATAGATAATATCTATTGTCGT
GCTCGTAGACATAGAGCAATAGGACCCAGCATCGCTAGTAGCAGTAC
15 GAAACTGATAAGAACGAGCCAGAGGAATCCATAGGTGTAAAGGCTAA
GACCAAGAGCTTGCACTTGTGATAGGTATGCGTATGAGGCATCAGCT
GCTTGTGAACCGTGTACAACGAATACATCAGTCATTGATGATGTAATA
ACTGGTGCTCCATCACCAAGAAGCAGAGTGTTAAGGCTGTATAGTGC
TCCAGTGATAGCGCTGACGGTTGTTGATGTTACGCTAGGCATTACGC
20 TAAAGGCTTCCAGTAGGAATACTCCAGCAATGATGAAAAGTAGTGGTA
GTCCTTGAGTAAATTCTTGCCCGCTAGCAGTAATTTAGAGAGACGTA
CGTTAAGCATCATCACCAAGCAATAGGAATAGCACAGCAACAGCTCCC
ACGTATACGACTAGGTATGTGAGACCTACGTACGTGATACCTAGACA
TACTAGGTAACAAGCAGCTAGGACGAATACACCAATTAGATATAGGAT
25 AGCTACAATAGGGCTGCTAGCTGTGATCGTCATTACAGCGCTTACAA
CAGTGGCTAGTGCCAGTAGTTCTAGTACAAAGCTAGTCANNNNNNNN
NNGTGACAGTCATGCTGATACTAAGCGATAAAGTAAAAAGCATTTATT
CACTGTCATAGCATCCAGGTCGAGCTTGCTGCCATAGATGTCCCACT
GGGAATACACATCGTCACCACTGTGGATCGAACACAGGTCGGAGCAT
30 TAACAGTGCCCTGCTCTACCACTGAGCTATGGTAACCATCGTCTATCA
CCCCTCTTATAGACCTAAACTATATGCTTGCTTATTAAATGGTAAAT

ACTATCACTAGTTACACACTTAACCATTTATTAACCCCTCTACCCCTTT
AGTTACACTCTTATACCCTCTCCCCTGCCTATCCTCTTACCCCCCCCC
TTGGGGGGGTTAAAAGGTGGGTAAAAGGGGATAGGGGGTCAAGATG
TGATCCCATTAATTTTCAATTTATTTAAATTAATTTAACTAGTGAAGTT
5 CAATAGCGTCCTTCAGGTATGAACATGTTAGAACACAGAATACGTATG
CTTGAATCAGAGATACAGCTAGTTCTAGACCAATAAGAGCTACAAACA
CAGCAAATGGAATTAGTGTGATAACTGCTACAATAACTCCTGCGCTGA
ATAGCTTAGCTAGGAAAGTAGCAAGAATCTTCAGTAGTGTGTGACCA
GCAGTCATGTTACTGAATAGCCGAACACCTAGGCTAGCAGCTCGTGC
10 AAGGTATGAACTGTTTTCGATCAGTACCAGCATAGGAACCAGAGCTA
GAGGAGTACCGCTTGGTACGAAGTATGAGAAGAAGTGTACTCCGTGT
AGACGTAGTCCTAGGATAGTTACAGCAAGGAATACAAATACTGATAGT
CCAAGAGCAGCAATACCGCTAGTAGTTACTGTGTATCCGTATGGAAT
GTTACCGTTTTAGGTTAGCTACAAGAATAAACATGAATAGCGCGTAGAT
15 GAACGGTGTGTAGATTTCTGTGACGGCTACCTACTTGGTCTCGAACCA
TACCTGCAAGTGTTGCGTAGGCAGCTTCTAGGGCTAGACTCCATCGA
CTTGGTACTAGGCTGTACCCGTTGTTAGCTACAACGTGAAGTGATACT
AGAAGTCCTACAGTAAGGAGTGTGTAAAGACCTAGGTTAGTAAGAGC
TAGAACAAAATCACCAACCACAGGTACGTGGAGGCTAATGAGAGGAA
20 CAACTTCAAATTGTTCAAGAGGAGAAAGGATTACTGCTTGTGTTGTCA
TTTATTGAGTTTTTAAGATTAATTAACAGATGTTAAGAGGTGTTGACAAC
AATTATTAGCAGTTTTTAATAGGCATAATAATATATTAACCCTAAAAG
AATTTAATAGTTCGAAAAAGGCTTAATAAGCAAGAAAAAGGAGGCAGG
TTTAAGACCCCTAATAGGTTTGGTTTAATAGTTGTTAGAAAAGAGCTTA
25 ATAAGTAACTGTTTAAAACATTAAAAGGTTGTTATTACTACTCTTACT
GAAGATTTATTAATCTAGGCTAGGGAGCTTCCTAGCCTCTTATTTTCA
ACAAGGAAACAAGAAAGGNNNNNNNNNNATAGTTTGTAAATGATGAG
CTTGTGAGCGGCCTCACGCGTGGACTGCAACTCTGTGCGAAGGGGT
TCAACTCCCCTCAAGCTCTATTCCTTGTCTTGGCCTAGCCCATAACTA
30 TTATGATAGTTATGTCTCATAGCTATTTAATGAAAAACCATTCAAAGCT
ATTGATTATAGCCTACTTATAGTTATTAATAATAAATCCCCTCTAGACC

CTTTTAGCAATAGGGATGGTAATAGTGGGTTGTAGGCGTACAGTGCT
GAACGGTATCGCACGCCTTTTTATTTTTTATTAAAAAAAATCAGACC
GTAGGAAGCTCAGAGAGCCATATATAGTATGCCTTGCCTTATAAGCTA
CTCACTATACAAATCCTTCACTAACATTATCATTCCCTTACTAACCCAT
5 ATATATATATATATATATTTTAATTGAGAATGAGGGGAAAGTGTTAGTA
ACATGTGAGGCAGTTATTACATGATAAGGATAGCACAAAGCTATGGCA
ATAGGCATAGAAACCTATGGTTATTACTATCTGGGATTTACCCATAAC
TAGTGATTATCCATGAGAAGAGCCACCCATACTAGGCCCATGTATATA
GACCATATCAGCATCAACAGGACTCTGGCTATGAATGTCTATAGCCC
10 CCTCCTCCCCACCATTTCTAACGCTACAACTTAATTTGTGTACTCTCC
TAAGCATAAAGATAATCTTATTAGCATAAAAAGTAACCAACCTAATAGT
ATAGGTCTAGTATAGATATTTGTATTTATCACTATAAGTATGGTTAATG
TATCACTTTTATTAAAATATTAATAGGGGTCTATTTATCCACTAATAGT
CGGTGTGTAAGCATAACCTTGACTGTATACACGTTCTTCTTGTGATG
15 GCTGTGGTAGAGGCTGTGTCTAGCACTTGTTGGTCGTTAGCCGTTAG
TGAACCTACAACGATTTCGTTGTTCTGTCATACGTGGTCGTGCTGGTTC
GGCAGATAGACGACCATGAACTGATACTTGTACCCCTACCAGACTAG
CTGGTAGTTGTA CTCCATTGACACTAGGGTAGGTAGTGATAGGGGTA
GCATTAGATACTGTACCAATAGTACCAAGCATAACGGTTGATAACCCGC
20 CCAAACCTTCTTAGTCTTAAGTTCCCGTGCTAGGTAGTGACCAAGAACG
GTAGCGTCCATATATGGGCCTGATAGTCGTGTAACGATTAGCATTACT
GGTCGTTGGCATACTTCGGACAGTGCCGTTTCGTAGTTGCATAATAGT
TAGTGACGATAGGGCATGGTTTGATGGTGCGTAGTAACCGACCTTCA
CAGTAATACCTCCTACAGGAAGTCCAGCTGCAGGACGTGTAATGATT
25 GTAGGTCGTCCCCAGGCGACTTGATGTCCAGCCTCAAAATGGTTGAA
TACCTTTAGGGTATTGTTGATGGTGTGTAGTACACGTTTCGTCTACGTA
AGAGTCTGATACGTATGTGTGTAGTGACCTGGTGCTAGTTGTAGAT
GTGCCATATTAATAATGATGAAAAAAGTGTGCGACCAATGGATAATGA
TCATAAGGATGTATTGAGATGGAACAGTTAATTGCTATCGAGCATATA
30 TTATCATTCATACTGATTCCCGAGAGGAAGGATTGAACTTCCGTATTA
TCGTTATGAGTGATACGCTCTAACCAACTGAGCTATCTCGAGTATTAC

TCTCTGTTACTAATTTTATCTATATAGTTAGGAACTAATAATCGTAAAC
TTTTTTAGGTAGGTATCTAAACCATGCACTGATAGGTTCTATAGGACG
TATTGGTCATAACTATGATCATTTATAAGCAGTACGAGATTCGAACTC
ATGACACACTGGTTAAGAGCCAGGCGCTCTACCGCTGAGCTAACCGC
5 TTAACACCGTACTTTTTATTAGGATAGGGTCCTATAGGAATGTTATCCA
TCTCATGCTGACCAACTACGAGTCGAACGTAGACAGACTCCACCAAA
AAAGAGCAGGGCTACCATTACCCCATTTGGTCATACACTGTGCACCAT
TACCTACCCTCCTCTACCCATCTTTTCTACCCTTTACTCCCCCCCCCTTT
TTAAGGGGGTTAGTAANNNNNNNNNNGACAGAGGTAACAAGATGAAA
10 TACTGCTTATAGTGATGGACAGCTGACAGGAGCAGACTCGAACGTG
CAGTGGACAATAGCCCGACTGGTTTACAGCCAGCTCCCTTACCAATT
AGGGGTACCCTGCCTTTGTTTATACGATACTGCCCCAACGTGGAGTC
GAACCACGGCCTGCTGAACTTCACTCAGCTGCTCTGACCAACTAAGC
CATCGGGGCTATCAATGCTCTATTAGAGATTGCAGATATCTATAACGA
15 CAGGTGAAAAGCAAATGCTGATAGTTAATAGGCAATGAACTTGCTAG
GCACAGAATGATAGAAAAATATTGTTACAAAAATGGTAAGAGATAATA
ATACATTGTAAGGAGGCATAGTACAATCCTACTCTGTGCCTAAACACC
TAGAACGATTCGAACGTCCATCTGTCTGGGCCGAAACCGAAGGCTCTA
CCATTGAGCTATAGGTGCAGGTTCCCTAGCTATCGACTATGAAAACCTT
20 CTCGATTAGGGGCATTGACTAGCACTATCATCAAGCATAGGAATATAT
AGCCTGATGGGGTAATTACTGTGGTAATACATATTTTCTATAGTATTTA
GGTAAACACGCAGCCTTATATAGTATTTATACGTTTGATAGGACTCGA
ACCTACACANNNNNNNNNNGCTACACATCTCTCGATACACGCTCCTA
AGGCGCGCCTGGCTACCAATTACAGCACAAACGTCATAGTGTTCTAT
25 ACACTATTAGGGCATAGAATACCTAGAATAAAATAGTAGATGCATTGC
TTATTACCACAACTATGTTACTATTCATAATACTATCACCATGCCTATT
AATTATGGAACCACTATGATGAGTTCTATTATCTATACTCTAATGAATG
AAAGGGGTCTATAGTATATCTGTTAGCTATGATTGTAGTGATGGGGCT
GTAGCAGCTGGAACATCAGCAGCTGGCACGTCCAGGTCAGATAGTG
30 TGTTTTCCACAATACCAACTACAGGTACAATAACAAGGTACCATCCGA
AGTAGAAGATACTTGCAATAGCACCGATAGTAATGAACGGTTCTTCAG

CGTGTTGGCTACCAAGCCATCCTAGTAGTACTAGGTCAACAGCTAGG
AACCAGAACGCTACTCGCCATAGTGGTCGGAATTGTCCACCACGTAC
ACGGCTAGTGTCCAGAATAGGCATAGCTAGCAGGATAAGTAGTGATC
CAAGCATGGCTACCACACCTAGCAGCTTGTTAGGTACGGCCCGTAGG
5 ATAGCGTAGTAAGGAAGTAGGTACCACTCAGGCACGATTGAGGCCG
GTGTACTCATAGGGTTGGCTGGGATGTAGTTGTCGCTGTGACCTAGC
AGGTTAGGTGCGTAGCACACAATACCTGTGATGGCAGCAAGGAATAC
CCACATAGTCACCAGGTCCTTGAATAGATAGTATGGAGCCATTGGCA
GCTTATCGCTGTTAGATGAGAGTCCAAGAGGGTTGCTTGACCCGTGT
10 GTGTGCAGCGTCATCATGTGCACTAGAGCTAGTGCAGATAGAACGAA
TGGTAGCACAAAGTGAAGAGAGAAGAATCGGTAAAGCGTAGCGTTGT
TTACACTAAATCCACCCCAAACAATTGTACGAAGTCTTGTCCGATCC
ATGGAATAGCTGAAAGCATGTTTGTAAATCACAGTAGCTCCCATAGTG
ACATTTGACCGTATGGTAGAACGTATCCAAGGAAAGCAATAGCCATC
15 ATAAGGACTAGGATAATTACACCTACTGACCATGGCATTACCCGTGGT
GCCTTGTATGAAGAGTAGTAAAGACCACGTCCGATGTGGGCGTATAC
TAGGATGAAGAAGAATGAGGCACCGTTAGCGTGTAGGTACCGTAGTA
GCCAACCGTAGTTAACGTCACGCATGATATGCTCAACACTCACAAAA
GCCAGGTCTACGTTAGGTGTGTAGTGCATCGCTAGAATGACACCAGT
20 CACAATTTGTAGACCTAGACATGTGGCTAGTAGTGATCCAAAGTTCCA
CATGTATGATAGGTTTCGAGGTTGTGGACTGTCGACGACGTAAGAGT
TTACTAGTCGTAGTAGTGGGTACTGCTTTAGCAGACGCATAGTACACT
AGTAATGTCCACCACATGGATTACAATGAAGCATGATGGACAGGTAT
CCCCGCAGGGACATTAAGAAGAGAGATGTCATGTTAATAGTGATAAA
25 TCTAACATGTTGTTAATGATAGTGAGGTTATTTAGCAGACTATTTAGT
ATGATACGTTACCAACGCTAGACGTATGAGGAGTCGAACCTCAATCA
CCACTATGAGTGGTGGACTACCAGCCTAATCTGATACACATCTTGTAC
AGTTTTAGACGTAATTAGATATCCATCACCTTTTAACTTTTCGTATACC
TCCAGAGACTCATGTATAATATAAGAACAGTGATACGACTCTATAGTC
30 ATAAGTACAGGAATCCCACCCCTTGCTATAAAAAGATATAATATTACC
ACATTCACACAGACGTTATCAGTTTTATCATGTTTAAGAATATCCTTTA

CCCGGATCATACTTAAGAGCATTATAATTTAGATGTGTTGAGGGGCTA
GGTCCACGTAGGGGGAGTGTAGTGCATCGCCAGAATCACTCCTGTAA
CGATTTGTAGACCTAGGCATGTAGTAGTGACCCAAAGTTCCACATGTA
TGATAAGTTCGCAGGTTGTGGACTGTCGACGACGTAAGAGTTTACCA
5 GTCGTAGTAGTGGGTACTGCTTTAGCAGACGCATCGTTCCTACTTAC
CCCCCTGTAAGGGGGGGTAAGGTGGCCATAGAATGGATTACATTAA
TGTCCATGACCGGGTACCCTATAGGGGCCTTACCTTAAAAAATAAACT
TCCAGAATAGGCATGGTGCTCTGATTAGAGGGTGAGAAAAGAAAAAT
ATAATTTCAATAAATAGGGCTATTACAAGCTTTAAAGCCTTAACGTTGC
10 CATCGCTAATAGACATACGAGGAGTCGAACCTCGCCCGCCGCTACAT
CAAAGCGGTGCACTAACCAGATATGCTATACGTCTTGTCTTGTCTGT
TATGCTTAGTTAGTTAATTATAACGTTTCAGTATTCAATGTTATTAATG
CCTGTAATTTATTACGGTTAACTATGCTTATGAGCAGGAATATAATGAA
TATATTTATTAAAAGGCTGGCTACCTTATCCATATTAGAGCAAGTTGG
15 CTACTTTTATGGATATTTAGAAAAAGGGTCTATAGTGATGGTACAACA
TATGCTAATACACTATGACTTGCTACCTTGGCTTACTGATGGTAGTAG
CCCTAGCCCTGCTACGATTAGTAGCACTAGAGACACGTCAGTAACAC
TGGCCGATGGTACCCAGGTAGTACTAGAGGAGCGATCACCATAAG
GGCAAGTGATAGGAAGCCTAGAACAAATGTAAAGGGCGTAGTTCGTTA
20 CGACTGATGTGTCGTAAGAAGCAATAGTACGTCCCATGTTAGGCAGT
ACTGTTGATAGTCCATAAGGACCAACAGTTTCGATCAGACCACGGTC
AAGAACCTTAGATGCAACGTATCCCAGGTTCATGAGAGGCGTGATTA
CTAGACCTGTTACAGCAGCGTTCCATCGCCATTGTCCGTTGATGAAC
CCGTATAGGGCTCGTGCTAGGGCACTGTCGGTTAGGCTTACAGCCA
25 GACGCATACCAGGACCGTATAGGAATACAGCTCCCATTGCTCCGAGT
AGTGTCATAACCAGTGGCAGTTGCTTGTATAGTACTGGTAGGGCAAA
TTCAGCTTCTACCAGTGATACGTGAGAAGGGTGTTGGGGCAGTACCG
CGCTGAGGTAGTCAGTACCCATTCCCTAGCCATAGGTCCTTAGCCATG
TATCCAAACAGTACTGAAAGCACTGCTAGGAATACAAGTGGAGCACC
30 TAGTAGTACTGGAGCTTCGTGCACGTTGTCGTAAGCTTGTCGTGTTC
CATTGACAGTACCAAAGAACGTTAGGGCGATCAGACGGATGCTATAG

AACGCTGTAAGACCAGCAGCCAGTGTACCGGCCAGTAGATTACTGT
TCCGTTAAGGCTGTATGTTCCCATAGCTAGTTCTAGTAGATGGTCCTT
ACTGTAGTACCCTGTCAGTCCTGGCAGGGCCATAAGTGACAGTGAAC
CAATCAGCATAGCCGCGTATGTGAACGGTAGTAGTGGAGCTAGTCCA
5 CCTAGTCGTCGTAGGTCTTGTTGGTCAGCCATAGCGTGGATAACACT
ACCAGCAGCCATAAACAGTAGAGCCTTGAAACAGGCGTGTGTAACAA
GGTGGAAGAGCCCTACGCTGAATTGTGATAGCCCTACAGCCATCATC
ATGTATCCCAGCTGTGAACATGTAGAGAAAGCGATCACTCGCTTTAG
GTCGTTCACTAGTAGACCACATGTAGCAGCGTAAGTAGCTGTTGCCG
10 CACCGAGCCAAGCAGTTACTACAAGGGCGGTAGGACCGTACTCTAGT
AGTGGTGATGTACGAAGCATAAGGTATACCCAGCAGTCACCAGCGT
AGCAGCGTGAATTAGAGCAGACACAGGAGTTGGACCCTCCATGGCC
AATAATTGTTAGTAGCAATGACAATCCCTTTCAGGATTGAGCAGGCCA
TGTC AACCATCAGGAAGTACTTATATGCAAAAAAATATCTTGAATCAT
15 GTTTAAGCATAACAACCTATGGGTACTAGTATGGTCTTTGGGCATGAA
TCACA ACTAGTGATTCTATGCTGCCAGTATTGAATAGTTACAAATACT
GGCAAATCATAGTACAGCAGACGCTTTAAAACCAAGACAGAAAAAGG
GGACTATCGTACAAGATGCCGAGCCCATACGGGTAGTTGAGCACCAA
TAAGCTCGTCTAGTACATGAAGGTGTT CAGTAGGTACCCAAACGTAG
20 TAGAGAGTGGTACGTGACCATGTGGGGCTGGGCGTTGGGAAGACA
GGGCTGGTACACCTAGGGATCGTAGTCGTGCAGCGAATAGGTCGTA
TTCGTAATCTTCAGCACCACTCCAAAGGGGTAGTACCAGTCCGCCTG
CTGGTTGTGAGAAGTACTCACGCCAAGTGTCTCGATAAGCGCTCAGA
ATAATATGAACAACAAGAGAATGAAGATTGTAGCAACCCAGTGCTTGG
25 GCTGTTAGACGTCGTACACCAGGGCCTGCAGGCGTCAGGTAGAAGA
GTGAATACAGAGGCCATAGTGCTGGTAGTGTACTTGTGAGCATCAGA
GTGAAGCTGACCGGTGGGAGGTGAGGACGACGGTTATCTGTCACCT
CAGTAGTAACAGGCTTATTGACCATAAACATGGTCATTAGTGATTGAA
CATATTCCCACTCAGTTTGGCTCCACATAGAAGAAATACGCTTGATTG
30 ATAGCCGTGCTGTGGGTAGACCACTAGTCAGGTACTTGCTGAACATT
TGGAATGGCCCCGCCACGAGCATAGCTACGAAAGCTGCAAGTTGATC

AGGGCTCATAGAGGTGGGCATATCCTTCTTAGCACTAGCACCCCTTGT
TACCATACGAACCAGGATTAGTACCAGGACCTGTCTTTTCACCTGAAC
TTGATGGAGTAGTACCCGTACCCACCTTTGTGGGATCTGTCCCGGTA
ATAAGATTTAGTTCATCAGCTAGCCACGTATGTAGTGGTAGTTGAGCA
5 CTCTTACTCATAGCCCCACCTAGGAAGAGTAGACCAATAACAGTTAGT
ACTGTTTTCGTTTAGAAGAGGTGCTGTGCTCAGAATAGTTGCGTAGTC
CAGGCTACCAAAGACAGCGAGAGCCAGGAAGAAACCAATGCTGAGT
GAGGTGTCACCAACACGGTTAACCATCATGGCCTTGTTAGCGGCCTT
AGACGCTTGTACCCGTGTGAGCCAGAACCCAATAAGGAGGAACGAT
10 GCCACCCCAATAAGTTCCCAACCAATGAACATCAGTGGGTATGAATC
CGCTGTTACCAGCAGAGCCATGCTCCCAGTGAACAGCGAAAGATAGC
TGAAGAATCGTGGTGTGTGTGGGTCTTCAGCCATGTATGACATGCTA
TAGACGTGTACAGCGCTAGATACGCATAGGACAGGTAGAAGCATCGC
AACTGTAAGGTCATCGAAGCTAAATGCCCATTGAGCTTGTAGCACAC
15 CAACGGTAAGCCAGTCACCTACAACGATAGTAACAGGTGAACGGCAA
AGAGCTACTTCGTAGAAGGCTACTAGTGATAGACATGCCGATAGTAC
TAGGCATACAGTGGCAACTAGTTGTGAACCTGTGCTACCAAGTGCTC
GCCCTCGTAGACCAGCGGCTGCTGAACCTAGGAAGGGTAGAAGAAT
AAGAGCTAGGTACATAGACTATAGTCGTGGAGATGATAGGCTCAGTG
20 TCCCCCGTAGGCGGTAATAGGCAACTAGTACTCCAAGACCAATAGCG
CTTTCAGCTCCCGCCGTAGCGATGATAATGATACTGAATGTCATTCCA
GCAGCATCGCTATAGCTGTTAGCTGTTGTGAGAACGAGTAGTGTCAC
CGCGAGCAGCATCAGTTCAATGCTCACAAGTAGCATGAGGAGGCTAC
GCCGGTTAAGGCTAAACCCCAGCACACCAACAAGGAACAGTACGAGT
25 GATAGAGTCATGGGACAGTCATTCGTGATAGGCGGAAGGCCATGTTT
AAACATCGTACAGTAGGAGTACTTGGCATCCATGATAGTGACCATAT
CACTCACTTCCCCCCCCGAAAGGGGGTAGTGGTAAGTAGCAGACTGTT
GCCAAACAGACTACCTATATCAGTAGTATAGTATCAGTACGTGGACCA
GCAACATTGAGTTACGGGCGACCCACTAACTCTCTCCACCCTTTACC
30 CCCCTATGAGGGGGGGTTGAAGGGATAGGTAAAAGGATTTAGTAAGA
AGGGATGTTTAAAAGATGGGTAAATATATTATTATAAGGTAGGGTTTA

GATTATGTGGGAGATTATTATAAGAATCCCGCGGCAGATGGGACTCG
AACCCACACTCACAACGTTGACAACGTTGGACACTGCCATTATGCTA
CTACCGCTAGGCACTGTCCTATTACTCTAATTACTATCTTTGTGATAAT
ACAGTTGTATTAGTATAGTAACAATAAAAGGTATCATCTGACNNNNNN
5 NNNNCCTTTTGGTGAAAGGAGAAAAGAAAAATCTATACCGCTGGTAC
GTTGTAGAGCAGTGCAGAAGCTTCAACTTCTAGACCAGACAGTAGTA
CGCCAGGGATCACACCAAGGAATAGAGTCATGACGATCAGAGGTAG
AACTGTCATAGCTTCTGTTTCGTGTCAGGTCACCTGCGATTGTTAGGTA
TGGACTTAGAGTTCCTGCGCTGATACGGGCCCATAGCCATGTAGAGT
10 AGCAAGCGCTAAGCACGATACCTGTAGCTCCAAGAGCACAAGCTAGG
AATGATTGTTCCATCACCTGCTAGAGCCATGAATTCACCCACCCAGT
TGA CTGAGAGAGGAACACCCATGTTAGCACAGATAGCAAAGAAGAAC
ACTACTGTGAATACTGGCATAACGCTAGCCATACCTCGGTAGTAGCT
GATAGACCGTGTGTGGTATCGGTTGTATAGAACACCACCCACCATCA
15 CGAATAGCGCTGGTGAGATTAGACCGTGCGCTAGTGATAGTAGTAGA
GCACCACTGATACCAATTACGCTGTTACTGAAGAGCCCTAGGACGAC
CACAGCCATGTGCCCTACTGATGAGTATGCAATAAGCTGCTTAGTGT
CTACTGACCGTAGTGTGGCTAGACTAGAGTAGATCAGACTTACAACA
GCAAGTGTTTGTACTAGAGGTGCGTAATAGTCGCAAGCATCAGGTAG
20 TACTGGTAGTACTAGTCGGTATACTCCATACGTGGCAAGCTTCAGCA
CTGTACCAGCTAGAAGCATAGATCCAGGCAGTGGAGCGTCAGCGTG
AGCACGAGGTAGCCAGATATGGAAAGGCACGATTGGAGTCTTTACCA
TAAGGGCTAGACCAAAGCACACAAACAGGTATCGTTGTACAGCAGGG
TCTAGTGATACGTGTTGTAGCACTGTCATGTCTGTTGTACCTGTGTGG
25 CTATATAGACCTAGAATACCTAGTAGCATGAATAGAGATCCTGCTAGT
GTATATAGGAATAGAAGTAGGGCTGACCGTACTCGTGTTGGTGAGCC
ACCATAGAGACCAACCATCACGTAAAGAGGCATCAGTACAGCTTCGA
ATGCTACGTAGAAGACGATGTAGTCAGTAGTAGTGAATGCAGCGTTA
ATAAGAGCTGTAGCAAGGAACACAAGAGCCAGTTGGGCCTTTTGTGA
30 AGCGTGAGGAGTGTGAGCTAGAGCTAGGACCATAGGTTCTACTAGGA
AGCATGAAAGTGTTACTAGTAGCATGCTAGTGTAGTCCACACCAAAG

AGAAGAGGAGCACCAATACCAATAAATTCGTGGTATTCCATAGCTCC
CCAGGGGTCAGTAGCCAGTAGCATCAGACAGACCATGTAACGAAGT
GTTTGTGACGACGCCATGGCCACAGTCATAGTAACGCCACCTGGTTG
TTCTGGATTAGCTCCACCTTCAGTAGGTTGCCATGCCAGGTAGATGG
5 CGGTGATAATAGGATATGCAATTAGATATGTAATCATAGATTTTATTAG
GGGGGTAGTAGCCTTAGTAGAATGGCCATTTGTCCCTCTACCCTGTT
AGTTAATAAATAAAGCTCTGCTTGAATATTATAGAAGTACTTATTATTA
TGATGATAGAACCGTTTATGGCGATAAACCTATATTATACGTATAGCA
GTAGAGTTCTACTAAACCTTTAACGATAGAGCACAATTTATTTATACCT
10 ATAACCATAGGGCCCATGTTTATGAAGATAGGTTCTTGTATTATGTTTA
TGGTGATAGAACCTATATATATATATATATACGGTTACTCACCATTGTT
TATAAGAGGTAGCCGGATCGAACGGCTGCTAGCGCTGTGTAAAAGC
GCTGACCTAACCCTAGTCGAACCTCCTTGTACTGTTAACATTATCTG
AGCCTATTAATGATTAGAAAATATACTATTTAGTACCATTAGCATAGCT
15 AGGAGAATACAGGGATACATGGGATCGAACCACGCCTAAGAGCTTTG
GAGACTCTTGTACTACCACTATACGATATCCCTATCAGGCTTAGGACC
GGACATAATATCTATCATCTTAATAGATAGGGGGGCTATCATGCCTAT
TCACAACAGATCATGATAGTATGATAAGCTATGGGTCAAGCCGCGGG
CTCGAACCGCGCACCTCTCCGACCACAACGGAACGCTCTGCCAAAT
20 GAGCTAGAATGACCTGTGTTATGGTCATACCAATACACCAGAGGAAT
AATAATGCTTAGAGGCAATAATATATTATTCAATACTAATAGTTTCCTC
TGTTTGCCTCCGCGTGGGATCGAACCACGATTTTAGGTTTACAAGAC
CCACGTTGCACCATTCAACTACAGAGGCTCATAGTGATCGATAAAGA
CCCCATATCTATGACCATTGTAACGCTACTGGGACTTGAACCCAGTCT
25 ACCTGTGTGAAGGACAGGTGTGCTACCCCTACACCATAGCGTCTACT
ACTGTTATATATGACATACTATGTTTCATGAGCACGCGCACATAAAAAT
ATCCCTAATAATATAACACTTGAGCTATGGTAAATAGTGGGGGTCTAT
TATTATGCTACCCATTACTACCCTTTTTAGGTTAGTTAAGGAAAAGTAA
AGCGGANNNNNNNNNNTAGCTAGGAGAATACAGGGATACATGGGAT
30 CGAACCCACGCCTAAGAGCTTTGGAGACTCTTGTACTACCACTATACG
ATATCCCTATCAGGCTTAGGACCGGACATAATATCTATCATCTTAATA

GATAGGGGGGCTATCATGCCTATTCAACAACAGATCATGATAGTATGAT
AAGCTATGGGTCAAGCCGCGGGCTCGAACCGCGCACCTCTCCGACC
ACAACGGAACGCTCTGCCAAATGAGCTAGAATGACCTGTGTTATGGT
CATACCAATACACCAGAGGAATAATAATGCTTAGAGGCAATAATATAT
5 TATTCAATACTAATAGTTTCCTCTGTTTGCCTCCGCGTGGGATCGAAC
CACGATTTTAGGTTTACAAGACCCACGTTGCACCATTCAACTACAGAG
GCTCATAGTGATCGATAAAGACCCCATATCTATGACCATTGTAACGCT
ACTGGGACTTGAACCCAGTCTACCTGTGTGAAGGACAGGTGTGCTAC
CCCTACACCATAGCGTCTACTACTGTTATATATGACATACTATGTTTCAT
10 GAGCACGCGCACATAAAAATATCCCTAATAATATAACACTTGAGCTAT
GGTAAATAGTGGGGGTCTATTATTATGCTACCCATTACTACCCTTTTT
AGGTTAGTTAAGGAAAAGTAAAGCGGANNNNNNNNNNNGTAAAGCGG
AGCCAGGACTCGAACCATGGACCTACAGCTCATGAGGCTGCAATGCT
GCCAAATTACACTACCCCGCGATTGATTATGACCATATTATATATAAA
15 GACAGGTTTATCTCCGTACTCGGACGCGATAGGAGTCGAACCTACAA
CTGGTCATCCCAGATACAGCTAGCAACCGTATACACCTCACCAATAG
TTGTGCACGTCCCGTCCTGTTACACATAGGTGAGTAAGCCTATAATAA
TGGGTCATACTATTAACGATAGGTAAATTAATCCTATAACAACGGG
ACATGTTATTAGCGATGTTGTTGTTTTCTTGTTATTCATGATTAGGTTT
20 GCCAACAATAGTGTGTGGTAAAAAGGGTGAGGGGTCATTAGGTAGTT
GGTATGGGTATTGGACTIONTGGGTGTGACTACGCTCTGGACGCATGC
CCATTACACGTGTCACCCCTTTACTATCTTTGATAGTTAATCTTGCGC
AGTAGTCTACTGCGTGTCCATAGCGTTTGAACATCGTTATTAGCAATG
TTCAAACGCGTTGATGCATGTCCATAGCTGGATTGCGGCTTTATATGC
25 TGGCAGCCCTTATCCATTGGGTGCTTAGCTGCCCTGCGGTGGCTCAT
TAATAGGCACAACAGGTACACCAGAGGTACCCTGACACTGGTCCTTT
CGTACTAGGTGCCATCCTCGAACATGCTAATAAAGCACAGCGGGTAT
AAGCTATACTGCTTCACAGCGTATTGACAATAGTGTTATTCATAATAAA
ACAGTGGCTTTACCACTGGTCAGACTCTATCATAGGCAGCAAGCTT
30 GCTACCGTGCAGGATAGTTATAGGCTCTGAGCCTTATAGTCGTTGGA
CGTGATACTTTAGAATAATTCACGCTCCTAGTTGTAAGGTATATACTTC

CAGGATGGTTTCTGCTTTTGGTGATCAATCACCGTTGGGCTCAATTTA
CATTATTCTCAACAAAACCTATGACTTAAACAAAGGGGTCTAGTTTAG
GTCAACTCGAGGACCTAGGCGGTACAGCATAGATTCATGCATATGAG
GTGTAAGCATGGCTTGTAAGTGCCTTGATGGGCTTTGTAATCGCA
5 AGCTTGTATTGCTTTTGTCCATCCTTTCCATAACGGTCAGCACGGACA
GTAGTGGGGATACCCATGGCAGTAATGGCCGCTGCTAGGCGTACGC
AATCTGCATGAGTATAGCTATTTACGTAGATGAATACGGTACTGTTTG
CGATGTCATATGAGCCGTCACTCATGATGAGATGGGCCAGTACAACA
GGCGTCACAAGATCCTTAATCATAGCCGGAACCTTGATCCACTC
10 ACCAGCAGCCCCCATCACATAAAACATATTGTGATATTGGTTGAAAAC
GCCGAGAGAAAGAGTCTTCAGCCGGAATTGTCCAGAGGGTAGTGTA
GTAACCCCCGTTTGGCAGAACTGGCTAAATAGTCCATATACCCAATA
GGCATAATCACGATAAGGGGGCCCCAAAAGACCATTCCAGGCGTGTAT
TAGATGTAGCACTTGAACGCGATGCATGGCTATCGCCCAGAAGTTGT
15 CCAATAACCACTTGGTGTAGGTAATATGGGAGCATTACAATGGATCGT
AGACCCCGAATAATTTTCGCTTGTAAGCGTTGAAGAAGTTGTAGTGTT
GATGTCATAGTTGTTTTTTGTTGATTGAATTAAATATATGTCGCTTAAG
CGAACCCAACTCACGTAGCTCCTTAGACTGCGAACAGCAGTACCCTT
CCAAGCTTGTGCACCTGGAGGATGAGCTGAGTCGACATCGAGGTCG
20 CAAACAGCACGGGCGATATGAACTCTCGCGTGCTATAACGCTGTTAT
CCCTAGCGTAGCGTTAATCGCCTAGCCCAGAGTGTACCCTGACACGT
CTGGGGTTCACTTGGCCATGCTTGCGCACTTTAAGCACTGCTCAGTG
CTATCAATCAGGCACCCATTTCATGTCCATGAACGATAGTAGCCCCTAC
ATGGGCAATACCTTAAAAAAGGCTATTCGGGTTTCCATCCCGGGCAA
25 GGGTGCCATACGTGGAGGCCTCCGGTACTGTATAGGAGGCGACCGT
CCCAGTCAAACCTAGCTGATGACCTATCCTAGCCCATTGCCCGTAAGA
GCGGGGCTTCACTACCACGAATCCTCATGGCGGTGTAATAGTCGAAC
GACATACTTCCCCAGATTCATGTGGCGACCGGCCATCACTAGTAAAG
CTGCATAGGGTCTTCTTGCCCTGCTGTGCTAAAACCGCATCTTCACG
30 GCTTATTCAGTTTTCGCTGAGGGGGCCCCCGGAGACAGCGGGTGCGTC
GTGACCCTCTTCATGCGGGACCATAATTAGTGGCCAAGGAATTTTCG

TACCTTAGCACCCCTTATGGTTAAGGCGGTCGTTTACCTGAGTTAGGTA
CACGTTGAAGCCCTTAAGCTCTACGCAGTACATTCACTCTAGGCACC
GGACAAGGGACGGCCCCCTATACGAGCCACAGCTGGTGCTGGGCATA
GCAGGGACCTGTGTTCTTGGTGACAGTCGCACACCCCGTTCCACTG
5 CGTCCAGCCAGGATGTTTAATTTCTAAGTTAAACTAAGCTTAAAAAAT
AACTATCCCCATAAGCTGGACACCCTTATAGTAGGACGTACGGGTT
TCTGTTGCCGAGTTCCTTTGGGAGCCCTATCTCTCACTCTGGTGTACT
CCACCACGGGACCTGTGTCGGTCATAGGCACGGTAAACGCACGCAG
CTTCAAGGCATATTTATTATATAGTATGCTCAGCTTGTGTTAACTCATC
10 GCTTTATACTTAGAGACCGGTTTAACTCAGGATGACTGCCATTGTCTA
TCTGAGACCCACCCCGTTCAAGTGCAAGGTGTCTAGACCTTGTATCG
CTACTCGTGTGCGCCATGCTCTATCCACGTTGTCCAGACAATGCTACA
CTGCCCTTCTTTCAACATGGGTGCTCATCTACCCTATAATCTTACTAC
ATCATTATTATTATAGCCCGCACCCGCCCATGTTACAAACCCCTTTGG
15 GGGTAAGAGCAGGTGAGGCATGTGAACTTTAGACATAAATACCCCTA
TCTTTCATGAAAAAGATAGGTAAATAATATTATATGTTTACAGTTTTAG
TAAGTATTTATAGGCGTTGTGTGGGACGATAGCTGAGACCCGTTTATC
TTCGGCGCACTGGTACTCGGCCACTGGGTTGTTACACCATCGTTAAC
AGGTGGCTACCTCCAGGCACACTGCATGACTGTCATTGTAGCAGCAC
20 CTCCTTACGTGTCATTTAGCTATATCTGTTTGGGTCCTCGCTCAACGC
TCATGACTGTCTTCATCTAGGCTGCTGACCTTATCGCCTACAGCCTGT
GTACAGGACGTAGTAGTCCCCCTTCCATGGTAAGCTACCCCTGATAG
AGCACGGATGCCCCCAGAACTGCACTCATTTCCCTATAAAACCTCCT
AGATGGTGATCTTATATTAATAAATACATGGCGGCCTGAACAGTACA
25 GGGTCCTACCACTCGATGCCTGTCTGACGCCTTAACGTCTTTGATG
AGAGCCGGCTAGCTCGCTGTTAGATAGGCCTATCACCCCTAGCCTGA
GCTCGTCAGAGTACAATAACGGACACCTGTTCGGCCTCTGTCATG
GCCATCACACGTCAATCAAGTACTTAAATACTTGACGTAAGAGGAC
ACAACAGGCCTGGCCCAAGCTAGATCACAGCGGTTCAAGTCTGGGC
30 AGGAGGACTAATTCCCCTATTGCACCATAATAAGCACCAATATAGCCT
ATTGACTATAACTGACACATATCACATCATTACCTTTATCAATCATTTT

TATTGTAATGATCCATACCTGATTTACCAATTTATAGTGTAGCTATAT
CTGACGGTTTATCTGATATAACAACAACTAAACTGATGGTCAATGG
CTTATATTGGTAGGAGATGTATGGATAGCAATGGGCGATTTTGTCTGT
CGTTTAGGCTATTGGCCTCGCCCTCCTTCCCAACTCGGCGACCCATT
5 ATGCAAAAGGTACGCTGTCTGGGCGTTACTATGGCGATCCTAGAAACC
CGACCATAGCATTGCAGCCCTTCAGCTGCTTATGGACCATAGCTGCT
TCAGGCTCTATTTCACTGCGGTACAACCGCTCTCTTTCATCGTTCACT
CGCGTTACTATTTTCTATCACTCGGACCACCTATTAGCCGTTGGGAGA
GATCCCAATCACTCTACAAAATGCAGAATGCTCTTTAACACATCAGCA
10 GTCACACGGGACTTTTACCCTCTATGGTTATCTGGCGTTCCATCCAGT
TCAAAGTGTGATGACATGTTACAGCTACACCTACTTTTCGCTCGCCG
CTACTAAGGAGGTACGGTTGGTTGCCTGTAGGCCCTACTATGATGAT
TCACTTTGGGCCATGGCTATCATTGATTTACCGTGAGGTTGTCATTAA
TTGACCTTTGGACTAGGGCNNNNNNNNNNCAACAAAAGATAGTGTGT
15 TTCCTACTACTCATTACTAACCCTTCTAATTAGAATTACCCTAACCCC
ACCAAAGGGGGGTAAGTGGTGGGTAAAAGGTAGGTAAAAGTAATAG
TACTCCACTAGTCATAATGGTTATAGGTGTAGGTACTCACTCATGAA
ATTATGACCATGAGTGTTTATAAATAACTATGACCATGAGTATAACTA
GTAGTATTGGTGGTTAAAAAGTTTAATGGTAATAGAATATGGCTTATG
20 CTTATGATGGGTGGGGTCTATAATACATGAATGTACTATGGACACCC
GGGTTTGGTGGTCATAAT (SEQ ID NO:2).

The entire coding region for the ribosomal genes of the PT22AV strain consists of the following sequence:

ACCTGGTTGATCCTGCCAGTAGTCATATGCTTGTCTCAAAGATTAAGC
25 CATGCATGTCTAAGTATAACAAATTCATACTGTGAACTGCGAATGG
CTCATTAATCAGTTATAGTTTATTTGATGGTATCTTGCTACATGGATA
ACTGTGGTAATTCTAGAGCTAATACATGCTGAAAAGCCCCGACTTCTG
GAAGGGGTGTATTTATTAGATAAAAAACCAATGGGTGCAAGCCCTCTA
TGGTGATTCATAATAACTTCTCGAATCGCATGGCCTTGCGCCGGCGA
30 TGCTTCATTCAAATATCTGCCCTATCAACTTTTCGATGGTAGGATAGGG
GCCTACCATGGTATCAACGGGTAACGGGGGAATTAGGGTTTCGATTCC

GGAGAGGGAGCCTGAGAAACGGCTACCACATCCAAGGAAGGCAGCA
GGNNNNNNNNNNATCCCGACACGGGGAGGTAGTGACAATAAATAAC
AATATAGGGCCCTATTGGGTCTTATAATTGGAATGAGTACAATTTAAA
TCCCTTAACGAGGAACAACCTGGAGGGCAAGTCTGGTGCCAGCAGCC
5 GCGGTAATTCCAGCTCCAGTAGCGTATATTAAAGTTGTTGCAGTTAAA
AAGCTCGTAGTCGAACTTCGGGCCTGGCTGGGCGGTCCGCCTTACG
GTGTGTACTGTCCGGCCGGGCCTTACCTCATGGTGAGCCCGTATGC
CCTTTACTGGGTGTGCGGTGGAACCATGAATTTTACCTTGAGAAAATT
AGAGTGTTCAAAGCAGGCATAAGCCCGAATACATTAGCATGGAATAA
10 TAGAATAGGACGTGCGGTTCTATTTTGGTTTCTAGGATCGCCGTA
ATGATTAATAGGGACGGTCGGGGGCATTAGTATTCAGTTGCTAGAGG
TGAAATTCTTAGATTTACTGAAGACTAACTTCTGCGAAAGCATTTGCC
AAGGACGTTTTTCATTGATCAAGAACGAAGGTTAGGGGATCAAAAACG
ATTAGATACCGTTGTAGTCTTAACAGTAACTATGCCGACTAGGGATC
15 GGGCCACGTTAATTTCTGACTGGCTCGGCACCTTACGAGNNNNNNNN
NNTTCTGGGGGGAGTATGGTCGCAAGGCTGAACTTAAAGGAATTGA
CGGAAGGGCACCAACAGGTGTGGAGCCTGCGGCTTAATTTGACTCA
ACACGGGGAACTCACCAAGGTCCAGACATAGTAAGGATTGACAGATT
GATAGCTCTTTCTTGATTCTATGGGTGGTGGTGCATGGCCGTTCTTAG
20 TTGGTGGAGTGATTTGTCTGGTTAATTCCGATAACGAACGAGACCTTA
ACCTGCTAAATAGTCAGGCCGGCTTCGGCTGGTCGTCGACTTCTTAG
AGGGACTGTCGGCGTTTAGCCGACGGAAGTTTGAGGCAATAACAGG
TCTGTGATGCCCTTAGATGTTCTGGGCCGCACGCGCTACACTGACTG
AGCCAGCGAGTTTATCACCTTAGCCGAGAGGCTTGGGTAATCTTGTG
25 AAACCTCAGTCGTGCTGGGGATAGAGCATTGCAATTATTGCTCTTCAAC
GAGGAATACCTAGTAAGCGTAAGTCATCAACTTGCGTTGATTACGTCC
CTGCCCTTTGTACACACCGCCCGTCGCTACTACCGATTGAATGGCTT
GGTGAGATTTCCGGATTGGCGTTGGGGAGCCGGCAACGGCACCCCTT
GGCTGAGAAGCTACTCAAACCTGGTCATTTAGAGGAAGTAAAAGTCG
30 TAACAAGGTTTCCGTAGGTGAACCTGCGGAAGGATCATTATTGATTG
GTCGAAAGACCTTATCAGATTCTACCACCTCTGTGAACCGTTGACCTC

CGGGTTAATAATCAAACATCAGTGTAACGAACGTAAGAGTATCTTAAC
GAAACAAAACCTTTCAACAACGGATCTCTTGGCTCTCGCATCGATGAAG
AACGCAGCGAAATGCGATAAGTAATGTGAATTGCAGAATTGAGTGAAT
CATCGAATCTTTGAACGCACCTTGCGCCTTTTGGTATTCCGAAAGGCA
5 TGCCTGTTTCAGTGTCATGAAATCTCAATCTAATATGTTTTCTGAACAT
GTTAGACTTGGACTTGGGCGTCTGCCAGTGATGGCTCGCCTCAAATG
ACTTAGTGGAACATCCACATCAGTGTTAGACGTAATAAGTTTCGTCT
CTCCTTGTGGTGATGACTGCTCAGAACCTGCCATCGCGCACTTTTGA
CTTTGACCTGAAATCAGGTAGGGCTACCCGCTGAACTTAAGCATATC
10 AATAAGCGGAGGAAAAGAACTAACAAGGATTCCCTAGTAACGGCGA
GTGAACCGGGAAGAGCTCAAATTTGAAATCTGGCGTGCTCAGTGCGT
CCGAGTTGTAATCTATAGAGGCGTTTTCCGTGCCGGACTGTGTCTAA
GTCCCTTGGAACAGGGTATCAAAGAGGGTGATAATCCCGCACTTGAC
ACAATGACCGGTGCTCTGTGATACGTCTTCTACGAGTCGAGTTGTTT
15 GGGAATGCAGCTCAAAATGGGTGGTGAGTTCCATCTAAAGCTAAATA
TTGGCGAGAGACCGATAGCGAACAAGTACCGTGAGGGAAAGATGAA
AAGCACTTTGGAAAGAGAGTTAAACAGTACGTGAAATTGTTGAAAGG
GAAACGATTGAAGTCAGTCGTGACTGAGAGGGCTCAGCCGGTTCTGCC
GGTGTATTCCCTCAGTCGGGTCAACATCAGTTTTGTTCCGGTGGATAA
20 GGGCAGCTGGAAGGTGGCACCTCCGGGTGTGTTATAGCCAGCTGTC
GCATACATCGAATGAGACTGAGGAATGCAGCTCGCCTTTATGGCGGG
GTTGCGCCACGTCCGAGCTTAGGATGTTGACATAATGGCTTTAAACG
ACCCGTCTTGAAACACGGACCAAGGAGTCTAACATATCTGCGAGTGT
TTGGGTGTCAAACCCAAGCGCGTAATGAAAGTAAACGTAGGAGGGAT
25 CCGCAAGGAGCACCTTCGACCGATCTGGATCTTCTGTGATGGATTTG
AGTAAGAGCATATATGCTGGGACCCGAAAGATGGTGAACATATGCCTG
AATAGGGCGAAGCCAGGGAACTCTGGTGGAGGCTCGTAGCGATTCT
TGACGTGCAAATCGATCGTCAAATTTGGGTATAGGGGCGAAAGACTA
ATCGAACCATCTAGTAGCTGGTTCCTGCCGAAGTTTCCTCAGGATAG
30 CAGAACTCGCATCAGTTTTATGAGGTAAAGCGAATGATTAGAGGAAT
TGGGGACGAAACGTCCTTAACCTATTCTCAAACCTTTAAATGTGTAAGA

AGGACTTGTCACTTAATTGGACGAGTCCATGCGAATGAGAGTTTCTA
GTGGGCCATTTTTGGTAAGCAGAACTGGCGATGCGGGATGAACCGAT
CGTGAGGTAAAGGTGCCGAATACACGCTCATCAGACACCACAAAGG
TGTTAGTTCATCTAGACAGCAGGACGGTGGCCATGGAAGTCGGAATC
5 CGCTAAGGAGTGTGTAACAACTCACCTGCCGAATGAACTAGCCCTGA
AAATGGATGGCGCTCAAGCGTGTTACCCATACCTCACCGTCGGCGTT
GAAGTGACGCGCCGACGAGTAGGCAGGCGTGGAGGTTTGTGAAGAA
GCCTTGGCAGTGATGCTGGGTGAAACAGCCTCTNNNNNNNNNNNAGA
TAGGGAAGCTCCGTTTCAAAGTGCACGCTTATCCGTGCCGCCTATCG
10 AAAGGGAATCCGGTTAAGATTCCGGAACCAGGATGTGGATCATTGAC
GGCAACGTAAATGAAGTTGGAGACGCTGGCAAGGGCCCCGGGAAGA
GTTCTCTTTTCTCCTTGACCGCTTACGACCTCGAAATCGGATTATCCG
GAGATGAGGTTATATGGCGGGCAGAGCACGACACCTCTGTCGTGTC
CGGTGCGTCCTTGACAGTCCTTGAAAATCCGACGGAACGTATAAGTC
15 TCACGCCTGGTCGTACTCATAACCGCAGCAGGTCTCCAAGGTGAACA
GCCTCTAGTTGATAGAACAATGTAGATAAGGGAAGTCGGCAAATAG
ATCCGTAACTTCGGGAAAAGGATTGGCTCTAAGGGTTGGGTACGTGC
GGCCATTGGCGGAACAGAGCTGGACCAGGTCCGACTTGCTGGGGCA
ACCCGGCTGGACTGGCTCGGACCGGCGATGGGATGNNNNNNNNNNN
20 AACTATGACTCTCTTAAGGTAGCCAAATGCCTCGTCATCTAATTAGTG
ACGCGCATGAATGGATTAACGAGATTCCCACTGTCCCTATCTACTATC
TAGCGAAACCACAGCCAAGGGAACGGGCTTGGCAGAATCAGCGGGG
AAAGAAGACCCTGTTGAGCTTGACTCTAGTTTGACATTGTGAAAAGAC
ATGGAGGGTGTAGAATAAGTGGGAGCTTCGGCGCCGGTGAAATACC
25 ACTACCTCCATCGTTTTTTTTACTTATTCAATGAGGCGGAGCTGGGATT
AACGTCCACCTTTTTGTCTTAAGGTCCTTTACGGGCTGATCCGGGTT
GAAGACATTGTCAGGTGGGGAGTTTGGCTGGGGCGGCACATCTGTT
AAAAGATAACGCAGGTGTCCTAAGGGGGACTCATGGAGAACAGAAAT
CTCCAGTAGAACA AAAAGGTAAAAGTCCCCTTGATTTTGATTTTCAGT
30 GTGAATACAAACCATGAAAGTGTGGCCTATCGATCCTTTAGTCCCTCG
GAACTCGAGGCTAGAGGTGCCAGAAAAGTTACCACAGGGATAACTG

GCTTGTGGCAGCCAAGCGTTCATAGCGACGTTGCTTTTTGATCCTTC
GATGTCGGCTCTTCCTNNNNNNNNNNGGAGTAAGTACTGACTCTCTTA
AGGTAGCCAAATGCCTCGTCATCTAATTAGTGACGCGCATGAATGGA
TTAACGAGATTCCCCTGTCCCTATCTACTATCTAGCGAAACACAGC
5 CAAGGGAACGGGCTTGGCAGAATCAGCGGGGAAAGAAGACCCTGTT
GAGCTTGACTCTAGTTTGACATTGTGAAAAGACATGGAGGGTGTAGA
ATAAGTGGGAGCTTCGGCGCCGGTCAAATACCACTACCTCCATCGTT
TTTTTTTACTTATTCAATGAGGCGGAGCTGGGATTAACGTCCACCTT
TTTGTCTTAAGGTCCTTTACGGGCTGATCCGGGTTGAAGACATTGTCA
10 GGTGGGGAGTTTGGCTGGGGCGGCACATCTGTTAAAAGATAACGCA
GGTGTCTTAAGGGGGACTCATGGAGAACAGAAATCTCCAGTAGAACA
AAAGGGTAAAAGTCCCCTTGATTTTGATTTTCAGTGTGAATACAAACC
ATGAAAGTGTGGCCTATCGATCCTTTAGTCCCTCGGAACTCGAGGCT
AGAGGTGCCAGAAAAGTTACCACAGGGATAACTGGCTTGTGGCAGC
15 CAAGCGTTCATAGCGACGTTGCTTTTTGATCCTTCGATGTCGGCTCTT
CCTGTCNNNNNNNNNNNAGAATTCGGTAAGCGTTGGATTGTTACCCA
CTAATAGGGAACGTGAGCTGGGTTTAGACCGTCGTGAGACAGGTTAG
TTTTACCCTACTGATGGAGGGTTATCGTAACAGTAATTGAGGGTAGTA
CGAGAGGAACTGCTCATTGAGATAATTGGTATTTGCGCCTGTCCGAT
20 CGGGCAATGGCGCGAAGCTATCATCTGCTAGATTATGGCTGAACGCC
TCTAAGTCAGAATCTGTACTGGAAACGATGATGTTGGTCCCGCATGT
GTTAGTTGTGTCAAATAGGCTTCGGCTGTGAACCATATCTGGGCTG
GGTTGTTTGGACGGAAAGGTCCTTGCAGCTTGCTCTTGATTGAAATG
GAATATACGCGGGGGGTGAATCCTTTGCAGACGACTTGAATTGGAAC
25 NNNNNNNNNNTAGTAGAGTAGCCTTGTTGCTACGATCTACTGAGGCT
AAGCCCTTGT (SEQ ID NO:3).

The COI1 gene of the PT22AV strain consists of the following nucleotide sequence:

ATGGCTAGCCGATGGCTCTTCTCTACAAACGCCAAGGATATTGGTAC
30 CCTTTATATCCTTTTTGCTGTCTTTACCGGTCTTCTGGGTACTGCGTT
CAGTGTACTTATCCGGATGGAGCTATCAGCTCCTGGTAACCAGTTCC

TTTCTGGTAACCACCACCTCTACAACGTAATTGCTACAACCTCACGGTA
TCATGATGATTTACTTCATGGTGGTACCTGCTATGGCTGGGTTCGCTA
ACTACATGGCTCCCGTTCTTATCGGTGCGCCGGACATGGCGTTCCCA
CGGCTGAACAACATCAGCTTCTGGCTACTACCTCCTGCAATCGTTCTT
5 ATCCTTTCAAGCGTCTTCGTGGAACAAGGTATGGGTAGCGGTTGGAC
TATGTACATGCCTCTAACAGGTGTACAATCACACAGTGGTGGATCAGT
TGATCTAGCTGTATTCAGCCTGCACCTCTCAGGGATTTTCATCTCTACT
TGGTGCCATTAACATTATCACTACAATCCTTAACATGCGAGCTCCAGG
GCTACGCCTACACAAGGCACCTCTATTTCGTATGGGCGATGCTATCAC
10 AGTCAGTTATTATTATTCTATGTATTCCAGTTCTAGCCGGAGCTCTTAC
TATGATCCTTACTGACCGTAACTTCAACACATCATTCTTCGACCCTGC
TGGAGGAGGGGACCCTGTCCTATAACCAACACCTATTCTGGTTCTTCG
GACACCCTGAAGTATACCTAATGATTATCCCTGGATTTGGTATGGTTA
GCCACGTTGTTTCAACATTTCAGTGGAAAGCCTGTTTTCGGTTACCTTG
15 GTATGGTGTACGCCATTGCCTCTATCGGTGTGCTTGGATTCATTGTAT
GGAGCCACCACATGTACGCCGTAGGGATGGACGTAGACACACGAGC
TTACTTCACAGCTGCATCTATGATTATTGCGGTACCTACAGGTATCAA
GGTCTTCAGTTGGCTAGCGACTGCTTACGGTGGTTCTATCCGACTGC
TAGCACCTATGCTATTCGCTCTAGGATTCGTCGGTCTGTTCACTATTG
20 GTGGACTGACAGGTGTAGTTCTAGCGAACGCGAGCGTTGACGTAGC
GATGCATGACACATACTACGTTGTGGCTCACTTCCACTACGTGCTTTTC
AATGGGTGCCGTATTCAGTATCTTTGCCGGGTTCTACTACTGGGCTC
CAAAGATGTTTCGGTAAGATGTACGAAGAACACCTAGCTCAAGCGCAC
TTCTGGACATTCTTCATCGGTGTTAACATGACATTTATGCCACAACAC
25 TTCCTAGGACTACAAGGAATGCCACGACGATTCGCTGACTACCCTGA
CGCTTACGCGGGATGGAACCTAATCTCATCATGGGGTTCATTGTTA
GCGTCTCTGCTACTCTTCTGTTCCCTGTACACTGTTTACGACATGTTCA
CTAACGACAAGTACGTTGGTGACAACCCATGGGGAGAACCGCTATAC
TTCACGGACAACCTAACGTACGAACGAAGCACACGGTTCGCCTACAC
30 AATCGAGTGGGTACTACCATCACCTACTCCACACCACGCATTCGACA
TGCTACCTATCCAATCATAG (SEQ ID NO:4).

The taxonomic classification of the PT22AV strain of *Papiliotrema terrestris* is shown in Table 1.

EXAMPLE 2. *Analysis of the antagonistic capacity and mechanisms of action of Papiliotrema terrestris PT22AV.*

5 The PT22AV strain was first characterized *in vitro* by analyzing its antagonistic capacity and mechanisms of action. The assay consisted of a pure culture in which the radial growth of fungal pathogens on an agar growth medium was evaluated and compared to the growth of the same plant pathogens in dual culture with *Papiliotrema terrestris* PT22AV as
10 biocontrol agent. By evaluating the differences in radial growth of the fungal pathogens in the different conditions, it was possible to quantify the antifungal effect produced by the antagonist yeast. The average inhibition produced by the PT22AV strain against the various fungal pathogens tested is approximately 18%, specifically: *Penicillium expansum* 25%;
15 *Monilinia fructigena* and *M. laxa* 20%; *Botrytis cinerea* 15%; *Fusarium* spp. 10%; *Rhizoctonia solani* 20%; *Alternaria solani* 15%; *Aspergillus* spp. 25%; *Colletotrichum* spp. 20%.

EXAMPLE 3. *Evidence of efficacy of Papiliotrema terrestris PT22AV in comparison with known products.*

20 The PT22AV strain of *Papiliotrema terrestris* species was tested in the context of efficacy trials with GAP certification, according to the guidelines: PP 1/135(4) Phytotoxicity assessment, PP 1/152(4) Design and analysis of efficacy evaluation trials and PP 1/181(5), Conduct and reporting of efficacy evaluation trials, including good experimental practice.
25 In particular, 32 efficacy and selectivity tests conducted over a period of 3 years (2018-2019-2020) in 4 EU countries, on 6 crops, against a wide range of biotrophic and necrotrophic fungal pathogens, in the field and in the post-harvest.

30 Tables 3a and 3b shows the efficacy of *Papiliotrema terrestris* PT22AV applied in the field or post-harvest against areal part fungal pathogens (Table 3a), and as soil inoculant and/or in fertigation against

soil-borne fungal pathogens (Table 3b), in different formulations (WG: wettable granule - WP: wettable powder - SC: suspension concentrated) and at different dosages.

The WG and WP formulations have the following composition: dehydrated yeast cells, cryoprotectants (mono and disaccharides), inert substances of mineral or organic nature, disintegrants. The quantitative ratios of the different components that make up the formulate vary according to the microbial concentration of the formulate. The SC formulate has the following composition: yeast cells, water, stabilizers.

The effectiveness of the PT22AV strain was compared with that of biological and chemical competitors using the following formula:

$$\{[(\text{Competitor efficacy} \times 100) / \text{PT22AV efficacy}] - 100\} \times -1.$$

In this formula, the efficacy of PT22AV is considered as the 100% and the percentage of fungicide competitor's efficacy is calculated in the first step. Subsequently, the percentage difference between the two treatments is calculated (-100). Finally, by multiplying for -1 the result will acquire a positive sign in the comparisons showing that PT22AV has a higher efficacy and a negative one in the comparisons where PT22AV displays a lower efficacy.

Strawberry	<i>Botrytis cinerea</i>	Grey mold (in field)	WP	9.0 x 10 ¹¹	20			+8			-40
			SC	2.0 x 10 ¹²	81			+13			+65
			WP	2.5 x 10 ¹²	55			+18			-29
			WG	2.0 x 10 ¹²	68			+33			-4
			WG	2.0 x 10 ¹²	58			+22			-22
			WP	2.5 x 10 ¹²	78			+42			+8
			WP	1.2 x 10 ¹²	57			+21			-24
			WP	2.0 x 10 ¹²	85			+47			+16
			WP	9.0 x 10 ¹¹	42			-7			-69
			SC	2.0 x 10 ¹²	88			+48		Cyprodinil	+19
			WP	2.5 x 10 ¹²	47			+14		Fludioxonil	-42
			WG	2.0 x 10 ¹²	56			+28			-19
			WG	2.0 x 10 ¹²	49			+22			-36
			WP	2.5 x 10 ¹²	80			+50			+16
			WP	1.2 x 10 ¹²	61			+34			-9
			WP	2.0 x 10 ¹²	88			+54			+23
Potato	<i>Phytophthora infestans</i>	Potato blight	WP	9.0 x 10 ¹¹	45			+11			-48
			SC	2.0 x 10 ¹²	78			+48			+14
			WP	2.5 x 10 ¹²	45		Copper oxychloride	+51			
			WG	2.0 x 10 ¹²	61			+63			

Apple tree				WG	2.0 x 10 ¹²	46			+52		
				WP	2.5 x 10 ¹²	76			+71		
				WP	1.2 x 10 ¹²	43			+48		
				WP	2.0 x 10 ¹²	80			+72		
	<i>Stemphylium</i> <i>sp.</i>	Brown spot		WG	2.0 x 10 ¹²	87	<i>Aerobasidium</i> <i>pullulans</i>		+64	<i>Fludioxonil</i>	+25
				WG	2.0 x 10 ¹²	80			+70		+18
				WG	3.0 x 10 ¹²	88			+63		+26
				WP	2.5 x 10 ¹²	80			+70		+18
				WP	1.2 x 10 ¹²	77			+72		+15
				WP	3.0 x 10 ¹²	90			+62		+27
				WP	9.0 x 10 ¹¹	68			+54		+5
				SC	2.0 x 10 ¹²	79			+60		+17
				WG	2.0 x 10 ¹²	79			+36		+25
				WG	2.0 x 10 ¹²	54			-1		-9
				WG	3.0 x 10 ¹²	87			+36		+32
				WP	2.5 x 10 ¹²	72			+23		+18
				WP	1.2 x 10 ¹²	67			+17		+12
				WP	2.0 x 10 ¹²	96			+42		+40
				WG	100gr/100l	75			+66		+17
				WP	50gr/100l	80			+68		+22

		{post-harvest treatment}											
Peach tree	<i>Monilia spp.</i>	Brown rot				WG	2.0 × 10 ¹²	82	<i>Bacillus subtilis</i>		+18	-8	
							WG	2.0 × 10 ¹²			70	+5	-27
							WG	3.0 × 10 ¹²			91	+26	+3
							WP	2.5 × 10 ¹²			86	+22	-2
							WP	1.2 × 10 ¹²			73	+9	-21
							WP	2.0 × 10 ¹²			88	+23	0
						WP	9.0 ×10 ¹¹	60			-11	-46	
						SC	4.0 × 10 ¹²	85			+21	-3	
Tomato	<i>Botrytis cinerea</i>	Grey mold				WP	2.5 × 10 ¹²	40	<i>Bacillus subtilis</i>		-10	-37	
							WG	2.0 × 10 ¹²			62	+30	+11
							WG	2.0 × 10 ¹²			57	+22	+3
							WP	2.5 × 10 ¹²			79	+44	+30
							WP	1.2 × 10 ¹²			63	+30	+12
							WP	2.0 × 10 ¹²			82	+46	+32
						WP	9.0 ×10 ¹¹	51			+13	-7	
						SC	2.0 × 10 ¹²	88			+50	+37	

Table 3b
Efficacy comparison between PT22AV formulations and fungicide competitors (soil-borne pathogens).

Soil single application				
Host crop	Pest	Application rate PT22AV (CFU / g soil)	Efficacy PT22AV (%)	Efficacy compared to <i>Trichoderma viride</i> (%)
Tomato	<i>Rhizoctonia solani</i>	1x10 ⁷	78	+17
	<i>Fusarium oxysporum</i>	1x10 ⁷	74	-5
	<i>Pythium ultimum</i>	1x10 ⁸	75	+7
Lettuce	<i>Sclerotinia sclerotiorum</i>	1x10 ⁷	71	-13
	<i>Fusarium oxysporum</i>	1x10 ⁷	73	+4
	<i>Pythium ultimum</i>	1x10 ⁸	79	+6
Melon	<i>Fusarium oxysporum</i>	1x10 ⁷	87	+16
	<i>Pythium ultimum</i>	1x10 ⁸	81	-4
	<i>Rhizoctonia solani</i>	1x10 ⁸	82	+10
Aubergine	<i>Phytophthora capsici</i>	1x10 ⁷	72	+13
	<i>Rhizoctonia solani</i>	1x10 ⁸	85	+18
	<i>Pythium ultimum</i>	1x10 ⁸	78	+6
Soil application + Irrigation				
Host crop	Pest	Application rate PT22AV (CFU / ml)	Efficacy PT22AV (%)	Efficacy compared to <i>Trichoderma viride</i> (%)
Tomato	<i>Rhizoctonia solani</i>	5x10 ⁶ CFU	84	+23
	<i>Fusarium oxysporum</i>		87	+10
	<i>Pythium ultimum</i>		88	+20
Lettuce	<i>Sclerotinia sclerotiorum</i>		91	+12
	<i>Fusarium oxysporum</i>		85	+18
	<i>Pythium ultimum</i>		85	+13
Melon	<i>Fusarium</i>		92	+21

	<i>oxysporum</i>			
	<i>Pythium ultimum</i>		93	+10
	<i>Rhizoctonia solani</i>		90	+18
Aubergine	Phytophthora capsici		82	+23
	<i>Rhizoctonia solani</i>		88	+20
	<i>Pythium ultimum</i>		87	+16

EXAMPLE 4. *Compatibility of Papiliotrema terrestris PT22AV with fungicides and efficacy in protocols of integrated disease management.*

5 The microbial formulates based on *Papiliotrema terrestris* strain PT22AV can be effectively applied in integrated pest management (IPM) in combination with synthetic chemical active ingredients. This possibility arises from the ability of the PT22AV yeast to tolerate the active ingredients with fungicidal action, allowing the two formulations to be used
10 simultaneously or in succession in time (Table 4).

The resistance or tolerance of the yeast object of the present invention was established following two main approaches:

i) The synthetic chemical active principles were incorporated, at increasing concentrations, in an agarized growth medium (NYDA: Nutrient
15 broth 8 gr l⁻¹; Yeast extract 5 gr l⁻¹; Glucose 10 gr l⁻¹; Agar 20 gr l⁻¹). Cell suspensions of the PT22AV strain, suitably diluted, were distributed on the Petri dishes containing the growth medium with the antifungal molecules, so the microorganism growth in the presence of fungicides was compared to that in the negative control (NYDA without fungicides).

20 ii) The active ingredients tested were dissolved in sterile distilled water at the concentrations suggested by the manufacturer. The PT22AV yeast-based formulates were suspended in these solutions and incubated at 20° C with stirring at 150 rpm for 1 hour. The yeast suspension in distilled water was used as the negative control. The suitably diluted cell
25 suspensions were inoculated on growth agar medium (NYDA) to verify their viability.

The first test method highlighted a good level of resistance of the PT22AV yeast to most of the fungicides tested, although with different tolerance levels among the various active ingredients tested (Table 4).

The second test method allowed to establish tolerance to synthetic chemical active ingredients in tank mix conditions, assuming a residence time in the spraying machine tanks for one hour. In this case, no tested fungicide caused a reduction in the viability of the PT22AV yeast as compared to the negative control. The broad spectrum of resistance that emerged is due to the fact that the yeast is placed in a state of quiescence determined by the formulation process and the cell activation time, following rehydration, is approximately 2 hours. In this period of time, the yeast cells resist better to abiotic stress than those in active growth; this allows the simultaneous use of the yeast with all the antifungal active ingredients.

Table 4

Minimum inhibitory concentration of different fungicides against PT22AV

Active Ingredient (AI)	AI concentration in the commercial formulation	Minimum inhibitory concentration (mg / l)	Resistance level
Strobilurins			
Azoxystrobin	22,9%	1000	Resistant
Trifloxystrobin	50%	1000	Resistant
Benzimidazoles			
Benomyl	50%	1000	Resistant
Carbendazim	41,7%	1000	Resistant
Thiabendazole	41,8%	1000	Resistant
Sterol inhibitors			

Tebuconazole	25%	100	Sensitive
Imazalil	10%	300	Sensitive
Penconazol	10,2%	200	Sensitive
Dicarboximide			
Procymidone	50%	1000	Resistant
Vinclozolin	50%	1000	Resistant
Dithiocarbamates			
Mancozeb	50%	300	Sensitive
Ziram	81%	500	Resistant
Inorganic fungicides			
Copper oxychloride	40%	500	Resistant
Sulfur	80%	2000	Resistant
Phenylpyrrole			
Fludioxonil	50%	1000	Resistant
Fungicides assembled			
Iprovalicarb + Copper oxychloride	4,2% + 35,6%	500	Resistant

EXAMPLE 5. *Biostimulant activity by the yeast species P. terrestris, in particular strain PT22AV, on crop plants.*

According to the present invention, for the first time biostimulant activity by the yeast species *P. terrestris*, in particular strain PT22AV, on crop plants has been shown.

The PT22AV strain has been applied via root application on melon, strawberry and lettuce crops as indicated in Table 2 and the biostimulant activity of the strain was assessed.

The results, reported in Table 5, show that the radical application of microbial formulations based on *P. terrestris* strain PT22AV increases the biomass of all the three crops assayed (melon, strawberry and lettuce), compared to the untreated check. The stimulation produced by the roots application of the yeast PT22AV is also evident in terms of leaf surface and the number of leaves/flowers.

Table 5

Crop: MELON						
Treatments		Leaf surface (cm²)	Number of leaves	Number of flowers	Dry weight of the root system (gr)	Dry weight of the aerial part (gr)
1	Untreated control	179 (a)	8.25 (a)	4.5 (a)	2.9 (a)	1.5 (a)
2	PT22AV WP	451.75 (c)	14 (b)	7.25 (b)	4.45 (b)	2.3 (b)
3	PT22AV WG	421.75 (c)	14 (b)	7.30 (b)	4.2 (b)	2.35 (b)
4	Radifarm	299.75 (b)	12.75 (b)	7.75 (b)	3.77 (ab)	2.12 (b)
Crop: STRAWBERRY						
Treatments		Leaf surface (cm²)	Number of leaves	Number of flowers	Dry weight of the root system (gr)	Dry weight of the aerial part (gr)
1	Untreated control	344 (a)	9 (a)	2.25 (a)	2.42 (a)	-
2	PT22AV WP	868 (c)	17.75 (c)	7.75 (c)	4.3 (b)	-
3	PT22AV WG	811.75 (c)	17.25 (c)	7.5 (c)	4.17 (b)	-
4	Radifarm	508.5 (b)	14 (b)	4.75 (b)	3.85 (b)	-

Crop: <i>LETTUCE</i>						
Treatments		Leaf surface (cm ²)	Number of leaves	Number of flowers	Dry weight of the root system (gr)	Dry weight of the aerial part (gr)
1	Untreated control	-	12.6 (a)	-	3.32 (a)	1.47 (a)
2	PT22AV WP	-	18.75 (b)	-	4.45 (b)	2.25 (b)
3	PT22AV WG	-	19.7 (b)	-	4.62 (b)	2.2 (b)
4	Radifarm	-	17.72 (b)	-	4.2 (b)	2.07 (b)

WP = Wettable powder, WG = Water dispersible granules

The values marked with the same letters are not statistically different (at $P \leq 0,05$) according to Tuckey's test.

5 **EXAMPLE 6.** *Efficacy comparison between PT22AV and LS28 isolates in three representative crops.*

The efficacy of PT22AV and LS28 strains was compared in in-field and post-harvest trials that were carried out for a period 5 years (from 2016 to 2020) in Italy, at the Molise University (Plant Pathology laboratory,
10 Department of Agricultural, Environmental and Food Sciences), on 7 crops in field trials and on 4 crops in post-harvest trials, against a wide range of biotrophic and necrotrophic fungal pathogens.

PT22AV and LS28 were applied following the protocol described in Table 2. For the field trials the efficacy was evaluated at harvest, while in
15 the post-harvest the measurements were performed when in the untreated control at least 10% of infection was obtained. At the same level of CFU (Colony Forming Units) per hectare, significantly higher efficacy of strain PT22AV than strain LS28 was observed. The comparison of application protocols between strains PT22AV and LS28, with reference to dosage,
20 crop plants and relevant plant pathogens are shown in Table 6.

Table 6
Efficacy comparison between PT22AV and LS28 isolates in three
representative crops

Field trials				
Culture	Disease	Pathogen	CFU per ha	Efficacy PT22AV vs LS28 (%)
Stone-fruit	Brown rot	<i>Monilia</i> spp.	1x10 ¹²	+17%
	Gray mold	<i>Botrytis cinerea</i>		+22%
Grape	Powdery mildew	<i>Erysiphe necator</i>	2x10 ¹²	+29%
	Gray mold	<i>Botrytis cinerea</i>	1x10 ¹²	+32%
	Downy mildew	<i>Plasmopara viticola</i>	2x10 ¹²	+41%
	Bunch rot	<i>Aspergillus</i> spp.		+17%
	Acid rot	yeasts and bacteria		+15%
Strawberry	Gray mold	<i>Botrytis cinerea</i>	1x10 ¹²	19%
Post-harvest trials				
Culture	Disease	Pathogen	CFU per ha	Efficacy PT22AV vs LS28 (%)
Stone-fruit	Brown rot	<i>Monilia</i> spp	5x10 ⁷	+18%
	Gray mold	<i>Botrytis cinerea</i>		+10%
Grape	Gray mold	<i>Botrytis cinerea</i>		+32
	Bunch rot	<i>Aspergillus</i> spp.		+12%

	Acid rot	Yeast and bacteria		+45%
Strawberry	Gray mold	<i>Botrytis cinerea</i>		+31%

Analogous comparisons of the Applicant have shown a significantly greater ability to resist a wide range of abiotic stresses by PT22AV as compared to LS28 (Table 7). This greater resistance to stressors is associated with higher adaptability, survival and viability in the environment, and with the colonization of plant tissues; these traits are in turn intuitively linked to the biocontrol activity against plant pathogens. Furthermore, greater resistance to stressors is associated with greater resistance and survival during the stage of yeast biomass production and to the stresses due to dehydration and freeze-drying methods, common steps of the formulation process, thus ensuring the production of a greater quantity of vital and active yeast biomass for biocontrol. In particular, the biomass production process results in a CFU concentration of strain PT22AV that is on average 70% higher than strain LS28. In addition, the successive formulation step further reduces the CFU concentration of LS28 of about 40% with respect to PT22AV strain with freeze dry process and over 90% with fluid bed formulation.

Table 7

Effect of stress on PT22AV and LS28 growth on agarized media.

Stress type	Compound	Tolerance level	
		PT22AV	LS28
Oxidative	H ₂ O ₂	+++	+
	Menadione	++	-
	Patulin	+++	+
	Sodium hypochlorite	+++	++
	Cumene hydroperoxide	+	-
Genotoxic	UV radiation	+++	++

	Hydroxyurea	+	-
Cell wall stress	Red congo	+++	++
	SDS	+++	+++
	Calcofluor white	++	-
	Caffeine	++	+
Osmotic stress	NaCl	+++	++
	KCl	+++	+
	Lithium chlorite	+++	++

The symbol (+++) stands for highly tolerant, (++) moderately tolerant, (+) low tolerant and (-) sensitive

EXAMPLE 7. *PT22AV efficacy against nematodes (*Meloidogyne* spp.) on tomato plants.*

The PT22AV strain has been applied on tomato plants as described in Table 2 and the efficacy of the strain against *Meloidogyne* spp nematodes was assessed. The results, reported in Table 8, show that when PT22AV strain is applied as a root treatment, while partially reducing the proliferation of nematodes it is also especially effective in limiting their harmful effects, as shown by measuring plant height.

Table 8
PT22AV efficacy against nematodes (*Meloidogyne* spp.) on tomato plants

Application timing	A: drenching before transplantation; BCD: through irrigation, with 7 days-time interval from each other			
Treatment	Active substance	Application rate (g or mL/ha)	% efficacy (Abbot transformation)	Average Height ©
Untreated Control	-	-	0.00 b	70.68 ab
PT22AV	<i>Papiliotrema terrestris</i> 3.5x10 ⁹ CFU/g	A: 1000 g/100 L	14.89 b	76.10 a
		BCD: 1000 g/ha		
PT22AV	<i>Papiliotrema</i>	A: 500	10.11 b	74.60 ab

	<i>terrestris</i> 3.5x10 ⁹ CFU/g	g/100 L BCD: 500 g/ha		
AV-YD-20-01	Yeast Derivatives 20%	10 l/ha	1.18 b	67.30 ab
BIOACT	<i>Paecilomyces lilacinus</i>	A: 750 ml/ha BCD: 750 ml/ha	3.81 b	65.08 ab
ERGOFERT NEMACONTROL	Yeast fluid extract 42.29 %	A: 6000 ml/ha BCD: 3000 ml/ha	3.72 b	63.83 b
VELUM PRIME	Fluopyram 400 g/L	625 ml/ha	47.06 a	74.45 ab

The values marked with the same letters are not statistically different (at $P \leq 0,05$) according to Tuckey's test.

5 The present invention has been described for illustrative, but not
limitative purposes, according to its preferred embodiments, but it is to be
understood that variations and/or modifications may be made by those
skilled in the art without thereby departing from the relative scope of
protection, as defined from the attached claims.

CLAIMS

1) Yeast strain belonging to *Papiliotrema terrestris* species, said strain being deposited at the Westerdijk Fungal Biodiversity Institute with deposit number CBS147138.

5 2) Yeast strain according to claim 1, wherein said yeast strain is in a form selected from fresh cells, dried cells, dehydrated cells, devitalized cells, inactivated cells, frozen cells or cells in aqueous suspension.

 3) Composition comprising the yeast strain of the *Papiliotrema terrestris* species, as defined in any one of claims 1-2, as active principle,
10 together with one or more excipients or adjuvants.

 4) Composition according to claim 3, wherein said composition is in liquid or solid form.

 5) Composition according to any one of claims 3-4, wherein the concentration of said yeast strain within the composition ranges from 10^3
15 CFU to 10^{13} CFU per gram of solid composition or from 10^2 CFU to 10^{12} CFU per mL of liquid composition.

 6) Composition according to any one of claims 3-5, wherein said composition is in the form of tablets, capsules, granules, pellets, powder, such as dry powder or wettable powder, fluid, dry fluid, emulsion,
20 suspension, solution, dispersion.

 7) Composition according to any one of claims 3-6, said composition further comprising one or more chemical compounds and/or one or more biological agents selected from fungicides, insecticides, fertilizers, biostimulants, macronutrients, micronutrients, elicitors, plant
25 regulators, microbial biocontrol agents, plant growth regulators, foliar nutrients, antibiotics, herbicides, acaricides, food additives, adjuvants, microorganisms of agri-food interest.

 8) Composition according to claim 7, wherein said fungicides are selected from 2-methoxyethylmercury chloride, 2-phenylphenol, 3-ethoxypropyl mercury bromide, 8-hydroxyquinoline sulfate, 8-phenylmercurioxyquinoline, acibenzolar, acibenzolar-S-methyl, acylamino
30

acid fungicides, acypetacs, Adavelt, *Agrobacterium radiobacter* K84, aldimorph, allyl alcohol, ametoctradin, aminopyrifen, amisulbrom, ampropylfos, anilazine, aureofungin, azaconazole, azithiram, azoxystrobin, *Bacillus amyloliquefaciens* (formerly *subtilis*) strain QST713, *Bacillus amyloliquefaciens* strain AH2, strain IT-45, strain FZB24, strain MBI600, strain D747, *Bacillus mycoides* isolate J, *Bacillus nakamurai* strain F727, *Bacillus pumilus* strain QST 2808, *Bacillus subtilis* var. *amyloliquefaciens* strain FZB24, *Bacillus subtilis* strain AFS032321, strain GB03, strain IAB/BS03, barium polysulfide, benalaxyl-M (=kiralaxyl), benodanil, benomyl, benquinox, bentaluron, benthiavalicarb, Benthiavalicarb-isopropyl, benzalkonium chloride, benzamacril, benzamide fungicides, benzamorf, benzohydroxamic acid, benzovindiflupyr, bethoxazin, binapacryl, biphenyl, bitertanol, bithionol, bixafen, blasticidin-S, Bordeaux mixture, boric acid, boscalid, bromuconazole, bupirimate, Burgundy mixture, buthiobate, calcium polysulfide, captafol, captan, carbamorph, carbendazim, carboxin, carpropamid, carvone, cell walls of *Saccharomyces cerevisiae* strain LAS117, Cheshunt mixture, chinomethionat, chlobenthiazole, chloraniformethan, chloranil, chlorfenazole, chlorodinitronaphthalene, chloroneb, chloropicrin, Chlorothalonil, chlorquinox, chlozolate, ciclopirox, climbazole, *Clonostachys rosea* strain CR-7, *Coniothyrium minitans*, copper hydroxide, copper naphthenate, copper oleate, copper oxychloride, copper soap, copper sulfate, basic, copper zinc chromate, copper(II) acetate, copper(II) carbonate, basic, copper(II) sulfate, coumoxystrobin, cufraneb, cuprobam, cuprous oxide, cyazofamid, cyclafuramid, cyclobutrifluram, cycloheximide, cyflufenamid, cymoxanil, cypendazole, cyproconazole, cyprodinil, Cyprofuram, dazomet, DBCP, debacarb, decafentin, dehydroacetic acid, dichlobentiazox, dichlofluanid, dichlone, dichlorophen, dichlorophenyl, dichlozoline, diclobutrazol, diclocymet, diclomezine, dicloran, diethofencarb, diethyl pyrocarbonate, difenoconazole, diflumetorim, dimethachlone, dimethirimol, dimethomorph, dimoxystrobin, diniconazole,

diniconazole-M, dinobuton, dinocap, dinocap-4, dinocap-6, dinoceton,
 dinopenton, dinosulfon, dinoterbon, diphenylamine, Dipymetitrone,
 dipyrithione, disulfiram, ditalimfos, dithianon, DNOC, dodemorph, dodicin,
 dodine, donatodine, drazoxolon, edifenphos, enestrobin, enestroburin,
 5 enoxastrobin, epoxiconazole, etaconazole, etem, ethaboxam, ethirimol,
 ethoxyquin, ethylene oxide, ethylmercury 2,3-dihydroxypropyl mercaptide,
 ethylmercury acetate, ethylmercury bromide, ethylmercury chloride,
 ethylmercury phosphate, etridiazole, extract from *Melaleuca alternifolia*
 (tea tree), extract from *Reynoutria sachalinensis* (giant knotweed), extract
 10 from the cotyledons of lupine plantlets ("BLAD"), extract of *Swinglea*
glutinosa, F500, famoxadone, fenamidone, fenaminosulf, fenaminstrobin,
 fenamistrobin, fenapanil, fenarimol, fenazaquin, fenbuconazole, fenfuram,
 fenhexamid, fenitropan, fenoxanil, fempiclonil, fempicoxamid, fenpropidin,
 fenpropimorph, fenpyrazamine, fentin, fentin acetate, fentin chloride, fentin
 15 hydroxide, ferbam, ferimzone, Florylpicoxamid, fluazaindolizine, fluazinam,
 fludioxonil, flufenoxystrobin, fluindapyr, flumorph, fluopicolide,
 fluopimomide, fluopyram, fluoroimide, fluotrimazole, fluoxapiprolin,
 fluoxastrobin, fluquinconazole, flurozolamide, flusilazole, flusulfamide,
 Flutianil, flutolanil, flutriafol, fluxapyroxad, folpet, fosetyl, fosetyl-Al, fthalide
 20 (phthalide), fuberidazole, furalaxyl, furametpyr, furcarbanil, furfural,
 furmecyclox, furophanate, *Gliocladium catenulatum* J1446, *Gliocladium*
virens GL-21, glyodin, griseofulvin, guazatine, halacrinat, halacrinat,
 hexachlorobenzene, hexachlorophene, hexaconazole, hexylthiofos,
 hymexazol, imazalil, imibenconazole, iminoctadine, Inatreq
 25 (fempicoxamid), inorganic oils, inpyrfluxam, iodocarb, ipconazole,
 Ipflufenquin, iprobenfos, iprodione, iprovalicarb, isofetamid, isoflucypram,
 isopropanol azole, isoprothiolane, isopyrazam, isotianil, isovalledione, Jun
 Si Qi, kasugamycin, kresoxim-methyl, laminarin, Lime sulfur (lime sulphur),
 mancopper, mancozeb, mandestrobin, mandipropamid, maneb, mebenil,
 30 mecarbinzid, mefenoxam, mefentrifluconazole, mepanipyrim, mepronil,
 meptyldinocap, metalaxyl, metalaxyl-M (=mefenoxam), metam,

metazoxolon, metconazole, methasulfocarb, methfuroxam, methyl
 bromide, methyl isothiocyanate, methylmercury benzoate, methylmercury
 dicyandiamide, methylmercury pentachlorophenoxide, metiram,
 metominostrobin, metrafenone, metsulfovax, metyltetraprole, milneb,
 5 myclobutanil, myclozolin, N-(ethylmercury)-p-toluenesulfonanilide, nabam,
 naftifine, natamycin, neem oil, nitrothal-isopropyl, nuarimol, octhilinone,
 ofurace, organic oils, orthophenyl phenol, orysastrobin, oxadixyl,
 oxathiapiprolin, oxazosulfyl, oxine copper, oxolinic acid, oxpoconazole,
 oxycarboxin, oxytetracycline, PCNB, pefurazoate, penconazole,
 10 pencycuron, penflufen, pentachlorophenol, penthiopyrad, phenamacril,
 phenylmercuriurea, phenylmercury acetate, phenylmercury chloride,
 phenylmercury nitrate, phosdiphen, Phosphite, phosphorous acid and
 salts, phosphorus acid, phthalide, picarbutrazox, picolinamides,
 picoxystrobin, piperalin, plant oils (mixtures): eugenol, geraniol, thymol,
 15 polycarbamate, polyoxin-D, potassium azide, potassium bicarbonate,
 potassium polysulfide, potassium thiocyanate, probenazole, prochloraz,
 procymidone, propamocarb, propiconazole, propineb, proquinazid,
 prothiocarb, prothioconazole, *Pseudomonas chlororaphis* strain AFS009,
Pseudomonas syringae ESC-10, pydiflumetofen, pyracarbolid,
 20 pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyrapropoyne,
 pyraziflumid, pyrazophos, pyribencarb, pyributicarb, pyridachlometyl,
 pyridinitril, pyrifenox, pyrimethanil, pyrimorph, pyriofenone, pyrisoxazole,
 pyroquilon, pyroxychlor, pyroxyfur, quinacetol, quinazamid, quinconazole,
 quinofumelin, quinomethionate, quinoxifen, quintozone, rabenzazole,
 25 Reynoutria sachalinensis, salicylanilide, sec-butylamine, sedaxane,
 sesame oil, silthiofam, silver, simeconazole, SJC17, sodium azide, sodium
 bicarbonate, sodium hypochlorite, sodium orthophenylphenoxide, sodium
 pentachlorophenoxide, sodium polysulfide, spiroxamine, *Streptomyces*
griseovirides strain K61, *Streptomyces lydicus* strain WYEC108,
 30 streptomycin, sulfur, sulfuryl fluoride, sultropen, tea tree oil, tebuconazole,
 tebufloquin, tecloftalam, tecnazene, tecoram, tetraconazole,

thiabendazole, thiadifluor, thicyofen, thifluzamide, thiochlorfenphim, thiophanate, thiophanate-methyl, thioquinox, thiram, THQ25, tiadinil, tioxyimid, tolclofos-methyl, tolfenpyrad, tolprocarb, tolylfluand, tolylmercury acetate, triadimefon, triadimenol, triamiphos, triarimol, triazbutil, triazoxide, 5 trichlamide, *Trichoderma afroharzianum* (formerly *harzianum*) strain T-22, *Trichoderma asperellum* (formerly *harzianum*) strain ICC012, strain T25, strain TV1, *Trichoderma asperellum* strain T34, *Trichoderma atrobrunneum* (formerly *harzianum*) strain ITEM908, *Trichoderma atroviride* (formerly *harzianum*) strain IMI206040, strain T11, *Trichoderma atroviride* strain I-1237, *Trichoderma atroviride* strain LU132, *Trichoderma atroviride* strain SC1, *Trichoderma gamsii* (formerly *viride*) strain ICC080, 10 triclopyricarb, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, UCQ09, *Ulocladium oudemansii* strain U3 , uniconazole-P, urea, validamycin, valifenalate, vinclozolin, voriconazole, WLR08, 15 zarilamid, zinc naphthenate, zinc thiazole, zineb, ziram, zoxamide, preferably azoxystrobin, trifloxystrobin, benomyl, carbendazim, thiabendazole, tebuconazole, imazalil, penconazole, procymidone, vinclozolin, mancozeb, ziram, copper oxychloride, sulfur, fludioxonil and/or iprovalicarb.

20 9) Kit comprising

a) the yeast strain of *Papiliotrema terrestris* species as defined in any one of claims 1-2, or a composition as defined in any one of claims 3-8, and

25 b) one or more chemical compounds and/or one or more biological agents or compositions including them, selected from fungicides, insecticides, fertilizers, biostimulants, macronutrients, micronutrients, elicitors, plant growth regulators, microbial biological control agents, plant growth regulators, foliar nutrients, antibiotics, herbicides, acaricides, food additives, antioxidants, adjuvants, microorganisms of agri-food interest.

30 10) Kit , according to claim 9, wherein said fungicides are selected from 2-methoxyethylmercury chloride, 2-phenylphenol, 3-ethoxypropyl

mercury bromide, 8-hydroxyquinoline sulfate, 8-phenylmercurioxyquinoline, acibenzolar, acibenzolar-S-methyl, acylamino acid fungicides, acypetacs, Adavelt, *Agrobacterium radiobacter* K84, aldimorph, allyl alcohol, ametocradin, aminopyrifen, amisulbrom, 5 ampropylfos, anilazine, aureofungin, azaconazole, azithiram, azoxystrobin, *Bacillus amyloliquefaciens* (formerly *subtilis*) strain QST713, *Bacillus amyloliquefaciens* strain AH2, strain IT-45, strain FZB24, strain MBI600, strain D747, *Bacillus mycoides* isolate J, *Bacillus nakamurai* strain F727, *Bacillus pumilus* strain QST 2808, *Bacillus subtilis* var. *amyloliquefaciens* 10 strain FZB24, *Bacillus subtilis* strain AFS032321, strain GB03, strain IAB/BS03, barium polysulfide, benalaxyl-M (=kiralaxyl), benodanil, benomyl, benquinox, bentaluron, benthiavalicarb, Benthiavalicarb-isopropyl, benzalkonium chloride, benzamacril, benzamide fungicides, benzamorf, benzohydroxamic acid, benzovindiflupyr, bethoxazin, 15 binapacryl, biphenyl, bitertanol, bithionol, bixafen, blasticidin-S, Bordeaux mixture, boric acid, boscalid, bromuconazole, bupirimate, Burgundy mixture, buthiobate, calcium polysulfide, captafol, captan, carbamorph, carbendazim, carboxin, carpropamid, carvone, cell walls of *Saccharomyces cerevisiae* strain LAS117, Cheshunt mixture, 20 chinomethionat, chlobenthiazole, chloraniformethan, chloranil, chlorfenazole, chlorodinitronaphthalene, chloroneb, chloropicrin, Chlorothalonil, chlorquinox, chlozolate, ciclopirox, climbazole, *Clonostachys rosea* strain CR-7, *Coniothyrium minitans*, copper hydroxide, copper naphthenate, copper oleate, copper oxychloride, copper soap, 25 copper sulfate, basic, copper zinc chromate, copper(II) acetate, copper(II) carbonate, basic, copper(II) sulfate, coumoxystrobin, cufraneb, cuprobam, cuprous oxide, cyazofamid, cyclafuramid, cyclobutrifluram, cycloheximide, cyflufenamid, cymoxanil, cypendazole, cyproconazole, cyprodinil, Cyprofuram, dazomet, DBCP, debacarb, decafentin, dehydroacetic acid, 30 dichlobentiazox, dichlofluanid, dichlone, dichlorophen, dichlorophenyl, dichlozoline, diclobutrazol, diclocymet, diclomezine, dicloran,

diethofencarb, diethyl pyrocarbonate, difenoconazole, diflumetorim, dimethachlone, dimethirimol, dimethomorph, dimoxystrobin, diniconazole, diniconazole-M, dinobuton, dinocap, dinocap-4, dinocap-6, dinoceton, dinopenton, dinosulfon, dinoterbon, diphenylamine, Dipymetitrone, 5 dipyrithione, disulfiram, ditalimfos, dithianon, DNOC, dodemorph, dodicin, dodine, donatodine, drazoxolon, edifenphos, enestrobin, enestroburin, enoxastrobin, epoxiconazole, etaconazole, etem, ethaboxam, ethirimol, ethoxyquin, ethylene oxide, ethylmercury 2,3-dihydroxypropyl mercaptide, ethylmercury acetate, ethylmercury bromide, ethylmercury chloride, 10 ethylmercury phosphate, etridiazole, extract from *Melaleuca alternifolia* (tea tree), extract from *Reynoutria sachalinensis* (giant knotweed), extract from the cotyledons of lupine plantlets ("BLAD"), extract of *Swinglea glutinosa*, F500, famoxadone, fenamidone, fenaminosulf, fenaminstrobin, fenamistrobin, fenapanil, fenarimol, fenazaquin, fenbuconazole, fenfuram, 15 fenhexamid, fenitropan, fenoxanil, fencpiclonil, fencpicoxamid, fenpropidin, fenpropimorph, fenpyrazamine, fentin, fentin acetate, fentin chloride, fentin hydroxide, ferbam, ferimzone, Florylpicoxamid, fluazaindolizine, fluazinam, fludioxonil, flufenoxystrobin, fluindapyr, flumorph, fluopicolide, fluopimomide, fluopyram, fluoroimide, fluotrimazole, fluoxapiprolin, 20 fluoxastrobin, fluquinconazole, flurozolamide, flusilazole, flusulfamide, Flutianil, flutolanil, flutriafol, fluxapyroxad, folpet, fosetyl, fosetyl-Al, fthalide (phthalide), fuberidazole, furalaxyl, furametpyr, furcarbanil, furfural, furmecyclox, furophanate, *Gliocladium catenulatum* J1446, *Gliocladium virens* GL-21, glyodin, griseofulvin, guazatine, halacrinat, 25 hexachlorobenzene, hexachlorophene, hexaconazole, hexylthiofos, hymexazol, imazalil, imibenconazole, iminoctadine, Inatreq (fencpicoxamid), inorganic oils, inpyrfluxam, iodocarb, ipconazole, Ipflufenquin, iprobenfos, iprodione, iprovalicarb, isofetamid, isoflucypram, isopropanol azole, isoprothiolane, isopyrazam, isotianil, isovaledione, Jun 30 Si Qi, kasugamycin, kresoxim-methyl, laminarin, Lime sulfur (lime sulphur), mancopper, mancozeb, mandestrobin, mandipropamid, maneb, mebenil,

mecarbinzid, mefenoxam, mefentrifluconazole, mepanipirim, mepronil,
 meptyldinocap, metalaxyl, metalaxyl-M (=mefenoxam), metam,
 metazoxolon, metconazole, methasulfocarb, methfuroxam, methyl
 bromide, methyl isothiocyanate, methylmercury benzoate, methylmercury
 5 dicyandiamide, methylmercury pentachlorophenoxide, metiram,
 metominostrobin, metrafenone, metsulfovax, metyltetraprole, milneb,
 myclobutanil, myclozolin, N-(ethylmercury)-p-toluenesulfonanilide, nabam,
 naftifine, natamycin, neem oil, nitrothal-isopropyl, nuarimol, octhilinone,
 ofurace, organic oils, orthophenyl phenol, orysastrobin, oxadixyl,
 10 oxathiapiprolin, oxazosulfyl, oxine copper, oxolinic acid, oxpoconazole,
 oxycarboxin, oxytetracycline, PCNB, pefurazoate, penconazole,
 pencycuron, penflufen, pentachlorophenol, penthiopyrad, phenamacril,
 phenylmercuriurea, phenylmercury acetate, phenylmercury chloride,
 phenylmercury nitrate, phosdiphen, Phosphite, phosphorous acid and
 15 salts, phosphorus acid, phthalide, picarbutrazox, picolinamides,
 picoxystrobin, piperalin, plant oils (mixtures): eugenol, geraniol, thymol,
 polycarbamate, polyoxin-D, potassium azide, potassium bicarbonate,
 potassium polysulfide, potassium thiocyanate, probenazole, prochloraz,
 procymidone, propamocarb, propiconazole, propineb, proquinazid,
 20 prothiocarb, prothioconazole, *Pseudomonas chlororaphis* strain AFS009,
Pseudomonas syringae ESC-10, pydiflumetofen, pyracarbolid,
 pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyrapropoyne,
 pyraziflumid, pyrazophos, pyribencarb, pyributicarb, pyridachlometyl,
 pyridinitril, pyrifenox, pyrimethanil, pyrimorph, pyrioferone, pyrisoxazole,
 25 pyroquilon, pyroxychlor, pyroxyfur, quinacetol, quinazamid, quinconazole,
 quinofumelin, quinomethionate, quinoxyfen, quintozone, rabenzazole,
Reynoutria sachalinensis, salicylanilide, sec-butylamine, sedaxane,
 sesame oil, silthiofam, silver, simeconazole, SJC17, sodium azide, sodium
 bicarbonate, sodium hypochlorite, sodium orthophenylphenoxide, sodium
 30 pentachlorophenoxide, sodium polysulfide, spiroxamine, *Streptomyces*
griseovirides strain K61, *Streptomyces lydicus* strain WYEC108,

streptomycin, sulfur, sulfuryl fluoride, sultropen, tea tree oil, tebuconazole, tebufloquin, tecloftalam, tecnazene, tecoram, tetraconazole, thiabendazole, thiadiflur, thicyofen, thifluzamide, thiochlorfenphim, thiophanate, thiophanate-methyl, thioquinox, thiram, THQ25, tiadinil, 5 tioxymid, tolclofos-methyl, tolfenpyrad, tolprocarb, tolylfluanid, tolylmercury acetate, triadimefon, triadimenol, triamiphos, triarimol, triazbutil, triazoxide, trichlamide, *Trichoderma afroharzianum* (formerly *harzianum*) strain T-22, *Trichoderma asperellum* (formerly *harzianum*) strain ICC012, strain T25, strain TV1, *Trichoderma asperellum* strain T34, *Trichoderma atrobrunneum* (formerly *harzianum*) strain ITEM908, *Trichoderma atroviride* (formerly *harzianum*) strain IMI206040, strain T11, *Trichoderma atroviride* strain I-1237, *Trichoderma atroviride* strain LU132, *Trichoderma atroviride* strain SC1, *Trichoderma gamsii* (formerly *viride*) strain ICC080, 10 triclopyricarb, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, UCQ09, *Ulocladium oudemansii* strain U3, uniconazole-P, urea, validamycin, valifenalate, vinclozolin, voriconazole, WLR08, zarilamid, zinc naphthenate, zinc thiazole, zineb, ziram, zoxamide, preferably azoxystrobin, trifloxystrobin, benomyl, carbendazim, thiabendazole, tebuconazole, imazalil, penconazole, procymidone, 15 vinclozolin, mancozeb, ziram, copper oxychloride, sulfur, fludioxonil and/or iprovalicarb.

11) Use of the yeast strain belonging to the yeast species *Papiliotrema terrestris* as defined in any one of claims 1-2, of extracellular substances secreted by said strain, of a composition as defined in any of 25 the claims 3-8, or of a kit as defined in any one of claims 9-10, against phytopathogens of one or more plants or crops, in particular before sowing, pre-harvest or post-harvest.

12) Use according to claim 11, wherein said one or more plants or crops are selected from cereals, such as wheat, barley, rye, oats, rice, 30 corn, sorghum; fruit trees, such as olive, apple, pear trees, apricot, nashi, plum, peach, almond, cherry, persimmon, banana, grape, strawberry,

raspberry, blackberry; citrus fruits, such as oranges, lemons, mandarins, clementines, grapefruits; legumes, such as beans, peas, lentils, soy; horticultural crops, such as spinach, lettuce, asparagus, artichokes, cabbage, carrots, onions, garlic, tomatoes, potatoes, aubergines, peppers, fennel; cucurbits, such as squashes, zucchini, watermelons, melons; oil plants, such as soy, sunflower, canola, peanut, castor, coconut; tobacco; coffee; you; cocoa; sugar beet; sugar cane; cotton.

13) Use according to any one of claims 11-12, wherein said phytopathogens are fungi, such as for example Basidiomycetes, Ascomycetes, Deuteromycetes, Oomycetes, fungi with sexual or asexual reproduction, fungi with biotrophic or necrotrophic activity, protists or Chromista.

14) Use according to claim 13, wherein said fungi are selected from the *Albugo* spp.; *Alternaria* spp.; Anthracnose; *Armillaria* spp.; *Ascochyta* spp.; *Aspergillus* spp.; *Blumeria graminis*; *Botrytis cinerea*; *Botrytis* spp.; *Bremia lactucae*; *Cercospora kikuchii*; *Cercospora soja*; *Cercospora* spp.; *Cercospora herpotrichoides*; *Cladosporium* spp.; *Claviceps purpurea*; *Colletotrichum* spp.; *Corynespora cassiicola*; *Diaporthe* spp.; *Erysiphe* spp.; *Fusarium graminearum*; *Fusarium oxysporum*; *Fusarium* spp.; *Helminthosporium* spp.; *Leveillula Taurica*; *Macrophomina phaseolina*; *Magnaporthe oryzae*; *Magnaporthe* spp.; *Melampsora lini*; *Monilinia* spp.; *Mucor* spp.; *Mycosphaerella graminicola*; *Mycosphaerella* spp.; *Oidium* spp.; *Penicillium* spp.; Downy mildew manshurica; Downy mildew spp.; *Phaeosphaeria* spp.; *Phakopsora pachyrhizi*; *Phakopsora* spp.; *Phoma* spp.; *Phytophthora* spp.; *Plasmopara viticola*; *Podosphaera* spp.; *Pseudoperonospora cubensis*; *Puccinia* spp.; *Pyrenochaeta lycopersici*; *Pyrenophora* spp.; *Pyricularia oryzae*; *Pythium* spp.; *Ramularia* spp.; *Rhizoctonia solani*; *Rhizoctonia* spp.; *Rhizopus* spp.; *Rhynchosporium* spp.; *Sclerotinia sclerotiorum*; *Sclerotinia* spp.; *Sclerotium cepivorum*; *Sclerotium rolfsii*; *Sclerotium* spp.; *Septoria glycines*; *Septoria* spp.; *Sphaerotheca* spp.; *Stemphylium* spp.; *Stenocarpella maydis*;

Thielaviopsis basicola; *Thielaviopsis* spp.; *Tilletia* spp.; *Uncinula* spp.; *Uromyces* spp.; *Ustilago maydis*; *Ustilago* spp.; *Venturia* spp.; *Verticillium* spp.

5 15) Use according to any one of claims 11-12, wherein said phytopathogens are bacteria, viruses or nematodes.

 16) Use according to claim 15, wherein the bacteria are selected from the species *Pseudomonas* spp., *Xanthomonas* spp., *Clavibacter* spp., *Ralstonia* spp., *Erwinia* spp.

10 17) Use according to claim 15, wherein the nematodes are selected from the species *Meloidogyne* spp., *Heterodera* spp., *Globodera* spp., *Belonolaimus* spp., *Pratylenchus* spp., *Rotylenchulus* spp., *Trichodorus* spp., *Paratylenchus* spp.

15 18) Use according to any one of claims 11-17, wherein said yeast strain or said extracellular substances or said composition or said kit are applied on one or more parts, intact or wounded, of the plant selected from seeds, aerial parts, leaves, stems, trunks, buds, gems, branches, stems, flowers, fruits, roots.

20 19) Use according to any one of claims 11-18, wherein the quantity of live cells of said yeast strain distributed per unit of culture surface ranges from 10^8 to 10^{15} CFU per hectare.

25 20) Use according to any one of claims 11-18, wherein said yeast strain or said extracellular substances or said composition or said kit are applied to a solid growth substrate, such as soil, preferably at a concentration of live cells of yeast strain ranging from 10^2 to 10^{12} CFU per gram of solid growth substrate.

30 21) Use according to any one of claims 11-20, wherein said phytopathogen is selected between *Botrytis cinerea* and *Plasmopara viticola* of vine; *Botrytis cinerea* of strawberry, *Botrytis cinerea* and *Phytophthora infestans* of tomato, *Venturia inequalis* and *Stemphylium* spp. of apple and/or pear tree, *Monilia* spp. of stone fruit and/or *Phytophthora* spp of potato.

22) Use according to any one of claims 11-21, wherein when the kit as defined in any one of claims 9-10 is used, said a) and b) are used separately or sequentially.

5 23) Method for the control of phytopathogens in agricultural crops comprising or consisting in applying the yeast strain belonging to *Papiliotrema terrestris* species as defined in any one of claims 1-2, extracellular substances secreted by said strain, a composition as defined in any of the claims 3-8 or a kit as defined in any one of claims 9-10 on said crops or on the soil in which the crop grows, before sowing, pre-
10 harvest or post-harvest.

 24) Method, according to claim 23, characterized in that said application takes place on one or more parts, intact or wounded, of the plant selected from seeds, aerial parts, leaves, stems, trunks, buds, shoots, branches, stems, flowers, fruits, roots.

15 25) Method according to any one of claims 22-24, wherein the dose of said composition ranges from 10^8 CFU to 10^{15} CFU per hectare.




Fig. 1




Fig. 2



Fig. 3

Tree scale: 0.01




Fig. 4

Tree scale: 0.1




Fig. 5

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2022/053387

A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N1/16 A01N63/32

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12N A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LIMA GIUSEPPE ET AL: "Activity of the Yeasts Cryptococcus laurentii and Rhodotorula glutinis Against Post-harvest Rots on Different Fruits", BIOCONTROL SCIENCE AND TECHNOLOGY., vol. 8, no. 2, 1 June 1998 (1998-06-01), pages 257-267, XP055869872, GB ISSN: 0958-3157, DOI: 10.1080/09583159830324 Whole doc., in partic. Table 1 (p.60) ----- -/--	1-25



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 June 2022

Date of mailing of the international search report

22/06/2022

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2

NL - 2280 HV Rijswijk

Tel. (+31-70) 340-2040,

Fax: (+31-70) 340-3016

Authorized officer

Roscoe, Richard

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2022/053387

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TOLAINI V ET AL: "Lentinula edodes enhances the biocontrol activity of Cryptococcus laurentii against Penicillium expansum contamination and patulin production in apple fruits", INTERNATIONAL JOURNAL OF FOOD MICROBIOLOGY, ELSEVIER BV, NL, vol. 138, no. 3, 15 April 2010 (2010-04-15), pages 243-249, XP026979159, ISSN: 0168-1605 [retrieved on 2010-02-14] Whole doc., in partic. abstract; col.2, p.243 -----	1-25
X	GIUSEPPE LIMA ET AL: "Integrated control of blue mould using new fungicides and biocontrol yeasts lowers levels of fungicide residues and patulin contamination in apples", POSTHARVEST BIOLOGY AND TECHNOLOGY, ELSEVIER, AMSTERDAM, NL, vol. 60, no. 2, 20 December 2010 (2010-12-20), pages 164-172, XP028171762, ISSN: 0925-5214, DOI: 10.1016/J.POSTHARVBIO.2010.12.010 [retrieved on 2010-12-24] Abstract, sections 2.6, 2.7, Fig.1, Table 1 -----	1-25
X	WO 2008/114304 A2 (UNI DEGLI STUDI DEL MOLISE [IT]; CASTORIA RAFFAOLLO [IT] ET AL.) 25 September 2008 (2008-09-25) Whole doc., in partic. p.7, 9-11, claims -----	1-25
A	MICCOLI CECILIA ET AL: "The necessity for molecular classification of basidiomycetous biocontrol yeasts", BIOCONTROL, KLUWER ACADEMIC PUBLISHERS, DORDRECHT, NL, vol. 65, no. 4, 23 March 2020 (2020-03-23), pages 489-500, XP037194352, ISSN: 1386-6141, DOI: 10.1007/S10526-020-10008-Z [retrieved on 2020-03-23] the whole document -----	1-25

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2022/053387

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☒ forming part of the international application as filed:
 - ☒ in the form of an Annex C/ST.25 text file.
 - ☐ on paper or in the form of an image file.
 - b. ☐ furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 - ☐ in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).
 - ☐ on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2022/053387

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2008114304	A2	25-09-2008	NONE